



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:



- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search


Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>

C

522,807



The Map Accompanying This
Text Is In
THE UNIV. OF MICH.
MAP COLLECTION







DEPARTMENT OF THE INTERIOR,
CENSUS OFFICE.

FRANCIS A. WALKER, Superintendent,
Appointed April 1, 1879; resigned November 3, 1881.

CHAS. W. SEATON, Superi
Appointed November 4, 1881.


REPORT

ON THE

FORESTS OF NORTH AMERI
(EXCLUSIVE OF MEXICO),

BY

CHARLES S. SARGENT,
ARNOLD PROFESSOR OF ARBORICULTURE IN HARVARD COLLEGE,
SPECIAL AGENT TENTH CENSUS.



WASHINGTON:
GOVERNMENT PRINTING OFFICE.
1884.

Class 317

Book Um 3-9

Accession 6534

LETTER OF TRANSMITTAL.

DEPARTMENT OF THE INTERIOR,
CENSUS OFFICE,
Washington, D. C., September 1, 1884.

Hon. H. M. TELLER,
Secretary of the Interior.

SIR: I have the honor to transmit herewith the Report on the Forests of North America (exclusive of Mexico),
by Charles S. Sargent, Arnold Professor of Arboriculture in Harvard College.

This report constitutes the ninth volume of the series forming the final report on the Tenth Census.

I have the honor to be, most respectfully, your obedient servant,

CHAS. W. SEATON,
Superintendent of Census.



For 1898
Library




TABLE OF CONTENTS.

LETTER OF TRANSMITTAL	Page ix
-----------------------------	------------

PART I.

THE FOREST TREES OF NORTH AMERICA, EXCLUSIVE OF MEXICO.

THE FORESTS OF NORTH AMERICA—GENERAL REMARKS	3-16
THE ATLANTIC REGION	3-6
THE PACIFIC REGION	6-10
DISTRIBUTION OF GENERA	10-12
DISTRIBUTION OF SPECIES	12-16
A CATALOGUE OF THE FOREST TREES OF NORTH AMERICA, EXCLUSIVE OF MEXICO, WITH REMARKS UPON THEIR SYNONYMY, BIBLIOGRAPHICAL HISTORY, DISTRIBUTION, ECONOMIC VALUES, AND USES	17-219
INDEX TO CATALOGUE	220-243

PART II.

THE WOODS OF THE UNITED STATES.

PRELIMINARY REMARKS	247
SPECIFIC GRAVITY AND ASH	248-251
FUEL VALUE	251, 252
THE STRENGTH OF WOOD	252
COMPARATIVE VALUES	252
TABLE OF RELATIVE VALUES	253-255
TABLE OF AVERAGES	256-259
TABLE ILLUSTRATING THE RELATION BETWEEN TRANSVERSE STRENGTH AND SPECIFIC GRAVITY IN THE WOOD OF CERTAIN SPECIES	259-264
GENERAL REMARKS	264, 265
TANNIN VALUES	265
TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES	266-349
TABLE II.—ACTUAL FUEL VALUE OF SOME OF THE MORE IMPORTANT WOODS OF THE UNITED STATES	350-353
TABLE III.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE UNITED STATES UNDER TRANSVERSE STRAIN	354-415
TABLE IV.—BEHAVIOR OF SOME OF THE WOODS OF THE UNITED STATES UNDER TRANSVERSE STRAIN: SPECIMENS EIGHT CENTIMETERS SQUARE	414-417
TABLE V.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE UNITED STATES UNDER COMPRESSION	418-481

PART III.

THE FORESTS OF THE UNITED STATES IN THEIR ECONOMIC ASPECTS.

GENERAL REMARKS	485
THE LUMBER INDUSTRY	485-489
FUEL	489
WOOD USED AS FUEL FOR VARIOUS PURPOSES	489
ESTIMATED CONSUMPTION OF WOOD FOR DOMESTIC PURPOSES	489
CONSUMPTION OF CHARCOAL	489
FOREST FIRES	491-493
TABLE OF FOREST FIRES OCCURRING DURING THE CENSUS YEAR	491, 492
NORTH ATLANTIC DIVISION	494-510
MAINE	494-496
NEW HAMPSHIRE	496-498
VERMONT	498-500
MASSACHUSETTS, RHODE ISLAND, AND CONNECTICUT	500, 501
NEW YORK	501-506
NEW JERSEY	506
PENNSYLVANIA	506-510

	Page.
SOUTH ATLANTIC DIVISION	511-523
DELAWARE	511
MARYLAND	511
DISTRICT OF COLUMBIA	511
VIRGINIA	511, 512
WEST VIRGINIA	512-515
NORTH CAROLINA	515-518
Naval stores	516, 517
SOUTH CAROLINA	518, 519
Burning off dead herbage	518
GEORGIA	519, 520
FLORIDA	520-523
Pencil cedar	522
Cypress	522
SOUTHERN CENTRAL DIVISION	524-546
ALABAMA	524-530
The Maritime pine region	525-527
Cypress swamps of the Tensas river	525-527
The forests of the Chattahoochee in eastern Alabama, mixed forest growth, etc	527, 528
Forests of the Tennessee valley	528, 529
General remarks	529
The pine belt of central Alabama	529
The pine region of the Coosa	529
Naval stores	529, 530
MISSISSIPPI	530-536
The pine forests of southern Mississippi	531, 532
The northeastern counties	532-534
Central pine hills	534
Western Mississippi	534, 535
The Yazoo delta	535, 536
LOUISIANA	536-540
Moss ginning	536, 537
TEXAS	540-543
INDIAN TERRITORY	543
ARKANSAS	543, 544
TENNESSEE	544, 545
Effect of fires upon the forest	545
KENTUCKY	545, 546
Pasturage of woodlands	546
NORTHERN CENTRAL DIVISION	547-563
OHIO	547
INDIANA	547
ILLINOIS	547-550
MICHIGAN	550-554
Forest fires	550, 551
Statistics of growing timber	551
WISCONSIN	554-558
MINNESOTA	558-560
Forests on Indian reservations	559, 560
IOWA	560
MISSOURI	560, 561
DAKOTA	561, 562
NEBRASKA	562
KANSAS	562, 563
WESTERN DIVISION	564-580
MONTANA	564-566
WYOMING	566, 567
COLORADO	567, 568
NEW MEXICO	568
ARIZONA	568, 569
UTAH	569-571
Lake range, west of Utah lake	570
Sanpete Valley range	570
Sevier River mountains	570
NEVADA	571
IDAHO	571-573
WASHINGTON	573-576
OREGON	576-578
CALIFORNIA	578-580
Pasturage of mountain forests	579, 580
ALASKA	580

TABLE OF CONTENTS.

vii

LIST OF ILLUSTRATIONS.

	Page.
MAP OF THE UNITED STATES, SHOWING THE CHARACTER OF THE FUEL USED IN THE DIFFERENT SECTIONS OF THE SETTLED PORTION OF THE COUNTRY.....	489
MAP OF THE UNITED STATES, SHOWING THE PROPORTION OF WOODLAND WITHIN THE SETTLED AREA BURNED OVER DURING THE CENSUS YEAR.....	491
MAP SHOWING DENSITY OF FORESTS IN MAINE, NEW HAMPSHIRE, VERMONT, MASSACHUSETTS, RHODE ISLAND, CONNECTICUT, NEW YORK, NEW JERSEY, AND PENNSYLVANIA.....	495
MAP OF MAINE, SHOWING THE DISTRIBUTION OF PINE AND SPRUCE FORESTS.....	496
MAP OF NEW HAMPSHIRE AND VERMONT, SHOWING THE DISTRIBUTION OF THE PINE AND SPRUCE FORESTS.....	497
MAP OF PENNSYLVANIA, SHOWING THE DISTRIBUTION OF THE PINE AND HEMLOCK FORESTS.....	506
MAP SHOWING DENSITY OF FORESTS IN DELAWARE, MARYLAND, WEST VIRGINIA, VIRGINIA, NORTH CAROLINA, OHIO, KENTUCKY, TENNESSEE, INDIANA, AND ILLINOIS.....	511
MAP OF WEST VIRGINIA, SHOWING THE DISTRIBUTION OF THE HARDWOOD, SPRUCE, AND PINE FORESTS.....	512
MAP OF NORTH CAROLINA, SHOWING THE DISTRIBUTION OF THE PINE FORESTS.....	515
MAP SHOWING DENSITY OF FORESTS IN SOUTH CAROLINA, GEORGIA, FLORIDA, ALABAMA, MISSISSIPPI, AND LOUISIANA.....	518
MAP OF SOUTH CAROLINA, SHOWING THE DISTRIBUTION OF THE PINE FORESTS.....	519
MAP OF GEORGIA, SHOWING THE DISTRIBUTION OF THE PINE FORESTS.....	520
MAP OF FLORIDA, SHOWING THE DISTRIBUTION OF THE PINE FORESTS.....	522
MAP OF ALABAMA, SHOWING THE DISTRIBUTION OF THE PINE FORESTS.....	524
MAP OF MISSISSIPPI, SHOWING THE DISTRIBUTION OF THE PINE FORESTS.....	530
MAP OF LOUISIANA, SHOWING THE DISTRIBUTION OF THE PINE FORESTS.....	536
MAP SHOWING DENSITY OF FORESTS IN TEXAS.....	540
MAP OF TEXAS, SHOWING THE DISTRIBUTION OF THE PINE FORESTS.....	541
MAP SHOWING DENSITY OF FORESTS IN MISSOURI, ARKANSAS, KANSAS, AND INDIAN TERRITORY.....	543
MAP OF ARKANSAS, SHOWING THE DISTRIBUTION OF THE PINE AND HARDWOOD FORESTS.....	544
MAP SHOWING DENSITY OF FORESTS IN MICHIGAN, WISCONSIN, MINNESOTA, AND IOWA.....	550
MAP OF THE LOWER PENINSULA OF MICHIGAN, SHOWING THE DISTRIBUTION OF THE HARDWOOD AND PINE FORESTS.....	551
MAP OF THE UPPER PENINSULA OF MICHIGAN, SHOWING THE DISTRIBUTION OF THE HARDWOOD AND PINE FORESTS.....	551
MAP OF WISCONSIN, SHOWING THE DISTRIBUTION OF THE HARDWOOD AND PINE FORESTS.....	554
MAP OF MINNESOTA, SHOWING THE DISTRIBUTION OF THE HARDWOOD AND PINE FORESTS.....	558
MAP SHOWING DENSITY OF FORESTS IN DAKOTA.....	561
MAP SHOWING DENSITY OF FORESTS IN NEBRASKA.....	562
MAP SHOWING DENSITY OF FORESTS IN MONTANA.....	564
MAP SHOWING DENSITY OF FORESTS IN WYOMING.....	566
MAP SHOWING DENSITY OF FORESTS IN COLORADO.....	567
MAP SHOWING DENSITY OF FORESTS IN NEW MEXICO.....	568
MAP SHOWING DENSITY OF FORESTS IN ARIZONA.....	569
MAP SHOWING DENSITY OF FORESTS IN UTAH.....	570
MAP SHOWING DENSITY OF FORESTS IN NEVADA.....	571
MAP SHOWING DENSITY OF FORESTS IN IDAHO.....	572
MAP SHOWING DENSITY OF FORESTS IN WASHINGTON.....	574
MAP SHOWING DENSITY OF FORESTS IN OREGON.....	576
MAP SHOWING DENSITY OF FORESTS IN CALIFORNIA.....	578
MAP OF A PORTION OF CALIFORNIA, SHOWING THE DISTRIBUTION OF THE REDWOOD FORESTS.....	580

MAPS CONTAINED IN PORTFOLIO ACCOMPANYING THIS VOLUME.

- No. 1.—MAP SHOWING THE POSITION OF THE FOREST, PRAIRIE, AND TREELESS REGIONS OF NORTH AMERICA, EXCLUSIVE OF MEXICO.
- No. 2.—MAP SHOWING THE NATURAL DIVISIONS OF THE NORTH AMERICAN FORESTS, EXCLUSIVE OF MEXICO.
- No. 3.—MAP SHOWING THE DISTRIBUTION OF THE GENUS FRAXINUS (THE ASHES) IN NORTH AMERICA, EXCLUSIVE OF MEXICO.
- No. 4.—MAP OF THE UNITED STATES, SHOWING THE DISTRIBUTION OF THE GENERA CARYA AND UMBELLULARIA (THE HICKORIES AND CALIFORNIA LAUREL).
- No. 5.—MAP OF THE UNITED STATES, SHOWING THE DISTRIBUTION OF THE GENUS JUGLANS (THE WALNUTS).
- No. 6.—MAP SHOWING THE DISTRIBUTION OF THE GENUS QUERCUS (THE OAKS) IN NORTH AMERICA, EXCLUSIVE OF MEXICO.
- No. 7.—MAP OF THE UNITED STATES, SHOWING THE DISTRIBUTION OF THE GENERA CASTANEA AND CASTANOPSIS (THE CHESTNUTS AND CHINQUAPINS).
- No. 8.—MAP SHOWING THE DISTRIBUTION OF THE GENUS PINUS (THE PINES) IN NORTH AMERICA, EXCLUSIVE OF MEXICO.
- No. 9.—MAP SHOWING THE DISTRIBUTION OF THE GENERA ABIES AND PICEA (THE FIRS AND SPRUCES) IN NORTH AMERICA, EXCLUSIVE OF MEXICO.
- No. 10.—MAP OF THE UNITED STATES, SHOWING THE DISTRIBUTION OF LIRIODENDRON TULIPIFERA AND PINUS LAMBERTIANA.
- No. 11.—MAP OF THE UNITED STATES, SHOWING THE DISTRIBUTION OF PROSOPIS JULIFLORA, QUERCUS ALBA, AND QUERCUS DENSIFLORA.
- No. 12.—MAP SHOWING THE DISTRIBUTION OF FRAXINUS AMERICANA AND PINUS PONDEROSA IN NORTH AMERICA, EXCLUSIVE OF MEXICO.
- No. 13.—MAP SHOWING THE DISTRIBUTION OF THE GENERA CHAMÆCYPARIS AND CUPRESSUS IN NORTH AMERICA, EXCLUSIVE OF MEXICO.
- No. 14.—MAP SHOWING THE DISTRIBUTION OF THE GENERA THUJA, TAXODIUM, AND SEQUOIA IN NORTH AMERICA, EXCLUSIVE OF MEXICO.
- No. 15.—MAP SHOWING THE DISTRIBUTION OF PINUS STROBUS, PINUS PALUSTRIS, AND PSEUDOTSUGA DOUGLASHII IN NORTH AMERICA, EXCLUSIVE OF MEXICO.
- No. 16.—MAP OF THE UNITED STATES, SHOWING THE RELATIVE AVERAGE DENSITY OF EXISTING FORESTS.

The Map Accompanying This
Text is in
THE UNIV. OF MICH.
LIB. COLLECTION

1

LETTER OF TRANSMITTAL

BROOKLINE, MASSACHUSETTS, *July 1, 1883.*

TO THE SUPERINTENDENT OF CENSUS.

SIR: I have the honor to submit the following report upon the nature and condition of the forests of the United States, to which are added statistics of the lumber and other industries directly dependent upon the forest for their support.

Mr. Andrew Robeson, of Brookline, Massachusetts, has prepared the maps which accompany this report; he has supervised the entire statistical work of this division and has conducted its correspondence.

Mr. Stephen P. Sharples, of Cambridge, Massachusetts, has conducted the various experiments undertaken with the view of determining the value of the different woods produced in the forests of the United States.

Mr. C. G. Pringle, of East Charlotte, Vermont, has examined the forests of northern New England and New York, Pennsylvania, and West Virginia; and subsequently, as an agent for the American Museum of Natural History, has greatly increased our knowledge of the trees of Arizona and southern California.

Mr. A. H. Curtiss, of Jacksonville, Florida, has studied the forests of Georgia and Florida, and subsequently, as an agent of the American Museum of Natural History, has added to our knowledge of the semi-tropical forests of southern Florida.

Dr. Charles Mohr, of Mobile, Alabama, has explored the forests of the Gulf states.

Mr. H. C. Putnam, of Eau Claire, Wisconsin, has gathered the forest statistics of Pennsylvania, Michigan, Wisconsin, and Minnesota.

Mr. George W. Letterman, of Allenton, Missouri, has examined the forests extending west of the Lower Mississippi River, and Professor F. L. Harvey, of Fayetteville, Arkansas, has gathered the forest statistics of that state.

Mr. Sereno Watson, of Cambridge, Massachusetts, has studied, during a long and arduous journey, the forests of the northern Rocky Mountain region, and Mr. Robert Douglas, of Waukegan, Illinois, those of the Black hills of Dakota.

I take this opportunity to call your attention to the faithful and admirable manner in which my associates have performed the difficult duties to which they were assigned; their zeal and intelligence have made possible the preparation of this report.

It is my pleasant duty also to call your attention to the fact that this investigation has been greatly aided from the first by the experience and knowledge of Messrs. G. M. Dawson, John Macoun, and Robert Bell, members of the Geological Survey of Canada; the information in regard to the distribution northward of the trees of the eastern United States is entirely derived from the latter's paper upon the Canadian forests, published in the Report of the Geological Survey of Canada for the years 1879-'80.

I am under special obligation to Dr. George Engelmann, of Saint Louis, Missouri, my companion in a long journey through the forests of the Pacific region, for valuable assistance and advice; his unrivaled knowledge of our oaks, pines, firs, and other trees has been lavishly placed at my disposal.

Mr. M. S. Bebb, of Rockford, Illinois, the highest American authority upon the willow, has given me the benefit of his critical advice in the study of this difficult genus. I desire to express to him and to Dr. Laurence Johnson, of New York, who has furnished me with a full series of notes upon the medical properties of the trees of the United States, the deep sense of my obligation. My thanks are also due to Mr. Henry Gannett, Geographer of the Tenth Census, for cordial co-operation in the work of this division; to Colonel T. T. S. Laidley, of the United States army, in command of the arsenal at Watertown, Massachusetts, and to Mr. James E. Howard, in charge of the testing machine there, for advice and assistance afforded Mr. Sharples while conducting the experiments upon the strength of woods, as well as to a large number of correspondents in all parts of the United States who have favored me with their cordial co-operation.

I am, sir, your obedient servant,

CHARLES S. SARGENT,
Special Agent.

PART I.

THE FOREST TREES OF NORTH AMERICA,
EXCLUSIVE OF MEXICO.

THE FORESTS OF NORTH AMERICA.

GENERAL REMARKS.

The North American continent, or that part of it situated north of Mexico, which will alone be considered here, may be conveniently divided, with reference to its forest geography, into the Atlantic and the Pacific regions, by a line following the eastern base of the Rocky mountains and its outlying eastern ranges from the Arctic circle to the Rio Grande. The forests which cover these two divisions of the continent differ as widely, in natural features, composition, and distribution, as the climate and topography of eastern America differ from the climate and topography of the Pacific slope. The causes which have produced the dissimilar composition of these two forests must be sought in the climatic conditions of a geological era earlier than our own and in the actual topographical formation of the continent; they need not be discussed here.

The forests of the Atlantic and the Pacific regions, dissimilar in composition in the central part of the continent, are united at the north by a broad belt of subarctic forests extending across the continent north of the fiftieth degree of latitude. One-half of the species of which this northern forest is composed extends from the Atlantic to the Pacific; and its general features, although differing east and west of the continental divide, in conformity with the climatic conditions peculiar to the Atlantic and the Pacific sides of the continent, still possess considerable uniformity. The forests of the Atlantic and the Pacific regions are also united at the south by a narrow strip of the flora peculiar to the plateau of northern Mexico, here extending northward into the United States. Certain characteristic species of this flora extend from the gulf of Mexico to the shores of the Pacific, and while the peculiar features of the eastern and the western slopes of the interior mountain system of the continent are still maintained here, the Atlantic and the Pacific regions of the Mexican forest belt possess many general features in common. Typical North American species, moreover, peculiar to the forests of the Atlantic or of the Pacific, mingle upon the Black hills of Dakota, and upon the Guadalupe and other mountains of western Texas, the extreme eastern ridges of the Rocky Mountain range, and the outposts between the Atlantic and the Pacific regions.

THE ATLANTIC REGION.

The forests of the Atlantic region may be considered under six natural divisions: the Northern Forest, the Northern Pine Belt, the Southern Maritime Pine Belt, the Deciduous Forest of the Mississippi Basin and the Atlantic Plain, the Semi-tropical Forest of Florida, and the Mexican Forest of Southern Texas (Map No. 2, portfolio):

These natural divisions, although composed in part of species found in other divisions and possessing many general features in common, are still for the most part well characterized by predominant species or groups of species, making such a separation natural and convenient.

The *Northern Forest* stretches along the northern shores of Labrador nearly to the sixtieth degree of north latitude, sweeps to the south of Hudson bay, and then northwestward to within the Arctic circle. This Northern Forest extends southward to the fiftieth degree of north latitude on the Atlantic coast, and nearly to the fifty-fourth degree at the 100th meridian. It occupies 10 degrees of latitude upon the Atlantic sea-board and nearly 20 degrees in its greatest extension north and south along the eastern base of the Rocky mountains. The region occupied by this Northern Forest, except toward its southwestern limits, enjoys a copious rainfall; it is divided by innumerable streams and lakes, and abounds in swampy areas often of great extent. The nature of the surface and the low annual mean temperature check the spread of forest growth and reduce the number of arborescent species, of which this forest is composed, to eight; of these, four cross to the Pacific coast, while the remainder, with a single exception, are replaced west of the continental divide by closely allied forms of the Pacific forest. The white and the black spruces are characteristic trees of this region; they form an open, stunted forest upon the low divides of the

water-sheds, and reach a higher latitude than any other arborescent species of the continent; the valleys and wide bottoms are clothed with broad sheets of poplars, dwarf birches, and willows. The forest of this entire region is scattered, open, stunted, and of no great economic value. It embraces, south of the sixtieth degree of north latitude, the northern extension of the great midcontinental plateau, which will be considered hereafter.

South of the Northern Forest the *Northern Pine Belt* extends from the Atlantic coast to the ninety-sixth meridian of longitude; east of the Apalachian Mountain system it extends south over nearly 6 degrees of latitude, with a long, narrow spur following the higher Alleghany ridges for nearly 3 degrees farther south; west of the Alleghany mountains, in the region of the great lakes, the pine forest is replaced south of the forty-third degree of latitude by the deciduous growth of the Mississippi basin. This second division of the Atlantic forest may be characterized by the white pine (*Pinus Strobus*), its most important, if not its most generally-distributed, species. East of the Apalachian system this tree often forms extensive forests upon the gravelly drift plain of the Saint Lawrence basin, or farther south and west appears in isolated groves, often of considerable extent, scattered through the deciduous forest. Forests of black spruce are still an important feature of this region, especially at the north, and within its boundaries the hemlock, the yellow cedar, the basswood, the black and the white ash, the sugar maple, and several species of birch and elm find their northern limits and the center of their most important distribution. The hickories and the oaks, characteristic features of the deciduous forests of all the central portion of the Atlantic region, reach here the northern limits of their distribution, as do the chestnut, the sassafras, the tulip tree, the magnolia, here represented by a single species, the red cedar, the tupelo, the sycamore, the beech, and other important genera.

The *Southern Maritime Pine Belt* extends from the thirty-sixth degree of north latitude along the coast in a narrow belt, varying from one hundred to two hundred miles in width, as far south as cape Malabar and Tampa bay; it stretches across the Florida peninsula and along the coast of the gulf of Mexico until the alluvial deposits of the Mississippi are encountered; it reappears west of that river in Louisiana, north and south of the Red river, and here gradually mingles with the deciduous forests of the Mississippi basin in Arkansas and eastern Texas. This belt is well characterized by the almost continuous growth, outside of the broad river bottoms and the immediate neighborhood of the coast, by the open forest of the long-leaved pine (*P. palustris*). The live oak, the palmetto, and various species of pine characterize the coast forest of this region; through the river bottoms and along the borders of the shallow ponds, scattered through the pine forest, different gums, water oaks, hickories, and ashes attain noble dimensions. The southern cypress (*Taxodium*), although extending far beyond the limits of this natural division, here attains its greatest development and value, and, next to the long-leaved pine, may be considered the characteristic species of the maritime pine belt.

The *Deciduous Forest of the Mississippi Basin and the Atlantic Plain* occupies, with two unimportant exceptions to be considered hereafter, the remainder of the Atlantic region. Through this deciduous forest, where peculiar geological features have favored the growth of *Coniferae*, belts of pine, growing gregariously or mixed with oaks and other broad-leaved trees, occur, especially upon some portions of the Atlantic plain and toward the limits of the Southern Maritime Pine Belt, west of the Mississippi river. The characteristic features of the forest of this whole region are found, however, in the broad-leaved species of which it is largely composed. Oaks, hickories, walnuts, magnolias, and ashes give variety and value to this forest, and here, with the exception of a few species peculiar to a more northern latitude, the deciduous trees of the Atlantic region attain their greatest development and value. Upon the slopes of the southern Alleghany mountains and in the valley of the lower Red river, regions of copious rainfall and rich soil, the deciduous forest of the continent attains unsurpassed variety and richness. Upon the Alleghany mountains northern and southern species are mingled, or are only separated by the altitude of these mountains; rhododendrons, laurels, and magnolias, here attaining their maximum development, enliven the forests of northern pines and hemlocks which clothe the flanks of these mountains or are scattered through forests of other broad-leaved species. The cherry, the tulip tree, and the chestnut here reach a size unknown in other parts of the country. The forest of the Red River valley is hardly less varied. The northern species which the elevation of the Alleghany mountains has carried south are wanting, but other species peculiar to the southern Atlantic and Gulf coasts are here mingled with plants of the southern deciduous forest. The seven species of *Carya* (the hickories) are nowhere else closely associated. A great variety of the most important oaks grow here side by side; here is the center of distribution of the North American hawthorns, which do not elsewhere attain such size and beauty. The osage orange is peculiar to this region; the red cedar, the most widely distributed of American *Coniferae*, the southern and the yellow pine (*Pinus palustris* and *mitis*) here reach their best development. Just outside of this region, upon the "bluff" formation of the lower Mississippi valley and of western Louisiana, the stately southern magnolia, perhaps the most beautiful of the North American trees, and the beech assume their greatest beauty, and give a peculiar charm to this southern forest.

The western third of the Atlantic region is subjected to very different climatic conditions from those prevailing in the eastern portion of the continent; it consists of an elevated plateau which falls away from the eastern base of the Rocky mountains, forming what is known as the Great Plains. This great interior region, on account of its remoteness from natural reservoirs of moisture, receives a meager and uncertain rainfall, sufficient to insure a growth of herbage, but not sufficient to support, outside the narrow bottoms of the infrequent streams, the scantiest

forests. This treeless plateau extends north to the fifty-second degree of north latitude; it follows southward the trend of the Rocky mountains far into Mexico, extending eastward at the point of its greatest width, in about latitude 40° N., nearly to the ninety-seventh meridian. This whole region is generally destitute of forest. The narrow bottoms of the large streams are lined, however, with willows, poplars, elms, and hackberries, trees adapted to flourish under such unfavorable conditions. These diminish in size and number with the rainfall, and often disappear entirely from the banks of even the largest streams toward the western limits of the plateau, south of the forty-fifth degree of latitude. North and east of these central treeless plains a belt of prairie extends from the sixtieth degree of north latitude to southern Texas. The average width east and west of this prairie region, through much of its extent, is not far from 150 miles. Its eastern extension, between the fortieth and forty-fifth degrees of latitude, is much greater, however, here reaching the western shores of lake Michigan, and forming a great recess in the western line of the heavy forest of the Atlantic region with a depth of nearly 600 miles. The transition from the heavy forest of the eastern and central portions of the Atlantic region to the treeless plateau is gradual. The change occurs within the prairie region. Here is the strip of debatable ground where a continuous struggle between the forest and the plain takes place. There is here sufficient precipitation of moisture to cause, under normal conditions, a growth of open forest, but so nicely balanced is the struggle that any interference quickly turns the scale. Trees planted within this prairie belt thrive if protected from fire and the encroachment of the tough prairie sod, and so extend the forest line westward; if the forest which fringes the eastern edge of the prairie is destroyed it does not soon regain possession of the soil, and the prairie is gradually pushed eastward.

The eastern line of the plain where arborescent vegetation is confined to the river bottoms, and which divides it from the prairie where trees grow naturally, to some extent, outside of the bottoms, and where they may be made to grow under favorable conditions everywhere, is determined by the rainfall enjoyed by this part of the continent. The extreme eastern point reached by this line is found, upon the fortieth degree of north latitude, near the northern boundary of the state of Kansas. North of the fortieth degree it gradually trends to the west, reaching the eastern base of the Rocky mountains in about latitude 52°. This northwestern trend of the eastern plain line may be ascribed to the comparatively small evaporation which takes place during the shorter summer of the north and to a slight local increase of spring and summer rainfall. South of the fortieth degree the plain line gradually trends to the southwest under the influence of the gulf of Mexico, reaching its extreme western point in Texas upon the one hundredth meridian.

Other causes, however, than insufficient rainfall and a nicely balanced struggle between the forest and the plain have prevented the general growth of trees in the prairie region east of the ninety-fifth meridian. The rainfall of this region is sufficient to insure the growth of a heavy forest. The rain falling upon the prairies of Minnesota, Wisconsin, Iowa, Illinois, and Missouri equals in amount that enjoyed by the Michigan peninsula and the whole region south of lakes Ontario and Erie, while prairies exist within the region of the heaviest forest growth. It is not want of sufficient heat, or of sufficient or equally distributed moisture, which has checked the general spread of forest over these prairies. The soil of which the prairies are composed, as is shown by the fact that trees planted upon them grow with vigor and rapidity, is not unsuited to tree growth. It is not perhaps improbable that the forests of the Atlantic region once extended continuously as far west at least as the ninety-fifth meridian, although circumstantial evidence of such a theory does not exist; and the causes which first led to the destruction of the forests in this region, supposing that they ever existed, cannot with the present knowledge of the subject be even guessed at. It is, however, fair to assume that forests once existed in a region adapted, by climate, rainfall, and soil, to produce forests, and that their absence under such conditions must be traced to accidental causes. It is not difficult to understand that the forest once destroyed over such a vast area could not easily regain possession of the soil protected by an impenetrable covering of sod and subjected to the annual burnings which have occurred down to the present time; while the force of the wind, unchecked by any forest barrier, over such an area would, even without the aid of fires, have made the spread of forest growth slow and difficult. The assumption that these eastern prairies may have once been covered with forests is strengthened by the fact that since they have been devoted to agriculture, and the annual burning has been stopped, trees which were formerly confined to the river bottoms have gradually spread to the uplands. Small prairies situated just within the western edge of the forest have entirely disappeared within the memory of persons still living; the oak openings—open forests of large oaks through which the annual fires played without greatly injuring the full-grown trees—once the characteristic feature of these prairies, have disappeared. They are replaced by dense forests of oak, which only require protection from fire to spring into existence. In western Texas, the mesquit, forced by annual burning to grow almost entirely below the surface of the ground, is, now that prairie fires are less common and destructive, spreading over what a few years ago was treeless prairie. The prairies, then, or the eastern portions of them situated in the region of abundant rainfall, are fast losing their treeless character, and the forest protected from fire is gradually gaining in every direction; regions which fifty years ago were treeless outside the river bottoms now contain forests covering 10 or even 20 per cent. of their area. These eastern, well-watered prairies must not, however, be confounded with their dry western rim adjoining the plains—the debatable ground between forest and plain—or with the plains themselves. There is now no gradual, constant spread of forest growth upon the plains. They are treeless, on account of insufficient moisture to develop forest growth; and while trees may, perhaps, if planted, survive during a few years

FOREST TREES OF NORTH AMERICA.

at the western limits of the prairie as here laid down, the permanent establishment of forests there does not seem practicable, and, sooner or later, a period of unusual drought must put an end to all attempts at forest cultivation in a region of such insufficient and uncertain rainfall (Map No. 1, portfolio).

It remains to consider the *Semi-tropical Forest of Florida* and the *Mexican Forest of Southern Texas*.

A group of arborescent species of West Indian origin occupies the narrow strip of coast and islands of southern Florida. This belt of semi-tropical vegetation is confined to the immediate neighborhood of the coast and to occasional hummocks or islands of high ground situated in the savannas which cover a great portion of southern Florida, checking, by the nature of the soil and want of drainage, the spread of forest growth across the peninsula. This semi-tropical forest belt reaches cape Malabar on the east and the shores of Tampa bay on the west coast, while some of its representatives extend fully 2 degrees farther north. It is rich in composition; nearly a quarter of all the arborescent species of the Atlantic forest are found within this insignificant region. The semi-tropical forest, in spite of its variety, is of little economic importance. The species of which it is composed here reach the extreme northern limit of their distribution; they are generally small, stunted, and of comparatively little value. Certain species, however, attain respectable proportions; the mahogany, the mastic, the royal palm, the mangrove, the sea-grape, the Jamaica dogwood, the manchineel, and other species here become considerable and important trees.

In western and southern Texas the trees of the Mississippi basin, checked by insufficient moisture from farther extension southward outside the river bottoms, are replaced by species of the plateau of northern Mexico. The streams flowing into the gulf of Mexico are still lined, however, east of the one-hundredth meridian, with the species of the Atlantic basin, which thus reach southward to beyond the Rio Grande. The Mexican forest belt of Texas extends from the valley of the Colorado river, near the ninety-eighth meridian, to the Rio Grande. It touches the coast not far from the Nueces river and extends to the eastern base of the mountain ranges west of the Pecos; here the species of which it is composed mingle with those peculiar to the Pacific-Mexican forest. The forest of this region, like that of all countries of insufficient moisture, is open, stunted, and comparatively of little value. It is characterized by enormous areas covered with chaparral (dense and often impenetrable thickets of thorny shrubs and small trees), by a stunted and occasional arborescent growth upon the hills and plains, and by fringes of heavier timber along the river bottoms. The most valuable and perhaps the most characteristic species of this whole region, the mesquit, extends to the Pacific coast. With this exception, none of the arborescent species peculiar to this region attain any considerable size or importance, although the forest of small junipers which covers the low limestone hills of the Colorado valley are locally valuable in a country so generally destitute of trees. The region immediately adjoining the Rio Grande abounds in different species of *Acacia*, *Leucæna*, and other Mexican *Leguminosæ*; and farther west, upon the dry plains of the Presidio, the Spanish bayonet (*Yucca baccata*) covers wide areas with a low, open, and characteristic forest growth.

THE PACIFIC REGION.

The Pacific forest region is coextensive with the great Cordilleran Mountain system of the continent. The causes which have influenced the present position and density of these forests must be sought in the peculiar distribution of the rainfall of the region. The precipitation of moisture upon the northwest coast is unequaled by that of any other part of the continent. It gradually decreases with the latitude until, in southern California, the temperature of the land so far exceeds that of the ocean that precipitation is impossible through a large part of the year. The interior of all this great region, shut off by the high mountain ranges which face the ocean along its entire extent, is very imperfectly supplied with moisture. It is a region of light, uncertain, and unequally distributed rainfall, heavier at the north, as upon the coast, and decreasing gradually with the latitude in nearly the same proportion. This entire region is composed of a mass of mountain ranges with a general north and south trend, separating long and generally narrow valleys. The precipitation of moisture within the interior region is largely regulated by the position of the mountain chains. Warm currents ascending their sides become cold and are forced to deposit the moisture they contain. It follows that, while the interior valleys are rainless or nearly so, the mountain ranges, and especially the high ones, receive during the year a considerable precipitation of both rain and snow. If the distribution of the forests of any region is dependent upon the distribution and amount of moisture it receives, forests exceeding in density those of any other part of the continent would be found upon the northwest coast; they would gradually diminish toward the south, and entirely disappear near the southern boundary of the United States, while the forests of all the interior region, from the summit of the principal Coast Ranges to the eastern base of the Rocky mountains, would be confined to the flanks and summits of the mountains. These forests would be heavy upon the high ranges, especially toward the north; they would disappear entirely from the valleys and low mountain ranges. An examination of the forests of the Pacific region will show that in general distribution and density they actually follow the distribution of the rainfall of the region. These forests well illustrate the influence of moisture upon forest growth. Within the Pacific region the heaviest and the lightest forests of the continent coexist with its heaviest and lightest rainfall.

The forests of the Pacific region may be considered under four divisions: the Northern Forest, the Coast Forest, the Interior Forest, and the Mexican Forest (Map No. 2, portfolio).

The *Northern Forest* of the Pacific region extends from nearly the seventieth to about the fifty-eighth degree of north latitude, or, immediately upon the coast, is replaced by the Coast Forest nearly 2 degrees farther north; it extends from the continental divide, here mingled with the Northern Forest of the Atlantic region, to the shores of the Pacific. The southern limit of this open, scanty Northern Forest, composed of species which extend across the continent, or of species closely allied to those of the Northern Forest of the Atlantic region, is still imperfectly known, especially in the interior. The determination of the southern range in Alaska and British Columbia of several species, as well as the northern range here of a few others, must still be left to further exploration. The white spruce, the most important and the most northern species of the forest of the North Atlantic region, is here also the most important species. It attains a considerable size as far north as the sixty-fifth degree, forming, in the valley of the Yukon, forests of no little local importance. The canoe-birch, the balsam poplar, and the aspen, familiar trees of the North Atlantic region, also occur here. The gray pine and the balsam fir of the Atlantic region are replaced by allied forms of the same genera. The larch alone, of the denizens of the extreme Northern Forest of the Atlantic coast, finds no congener here in the northern Pacific forest.

The Pacific *Coast Forest*, the heaviest, although far from the most varied, forest of the continent, extends south along the coast in a narrow strip from the sixtieth to the fiftieth parallel; here it widens, embracing the shores of Puget sound and extending eastward over the high mountain ranges north and south of the boundary of the United States. This interior development of the Coast Forest, following the abundant rainfall of the region, is carried northward over the Gold, Selkirk, and other interior ranges of British Columbia in a narrow spur extending north nearly to the fifty-fourth parallel. It reaches southward along the Cœur d'Alène, Bitter-Root, and the western ranges of the Rocky Mountain system to about latitude 47° 30', covering northern Washington territory, Idaho, and portions of western Montana.

The Coast Forest south of the fiftieth degree of latitude occupies the region between the ocean and the eastern slopes of the Cascade Range; in California the summits of the principal southern prolongation of these mountains, the Sierra Nevada, marks the eastern limits of the Coast Forest, which gradually disappears south of the thirty-fifth parallel, although still carried by the high ridges of the southern Coast Range nearly to the southern boundary of the United States. The Coast Forest, like the forests of the whole Pacific region, is largely composed of a few coniferous species, generally of wide distribution. The absence of broad-leaved trees in the Pacific region is striking; they nowhere form great forests as in the Atlantic region; when they occur they are confined to the valleys of the coast and to the banks of mountain streams, and, economically, are of comparatively little value or importance. The characteristic and most valuable species of the northern Coast Forest are the Alaska cedar (*Chamaecyparis*), the tide-land spruce, and the hemlock. These form the principal forest growth which covers the ranges and islands of the coast between the sixty-first and the fiftieth parallels. Other species of the Coast Forest reach here the northern limits of their distribution, although the center of their greatest development is found farther south.

The red fir (*Pseudotsuga*), the most important and widely-distributed timber tree of the Pacific region, reaches the coast archipelago in latitude 51°; farther inland it extends fully 4 degrees farther north, and in the region of Puget sound and through the Coast Forest of Washington territory and Oregon it is the prevailing forest tree. The characteristic forest of the northwest coast, although represented by several species extending south as far as cape Mendicino, near the fortieth parallel, is replaced south of the Rogue River valley by a forest in which forms peculiar to the south rather than to the north gradually predominate. The forest of the northwest coast reaches its greatest density and variety in the narrow region between the summits of the Cascade Range and the ocean. North of the fifty-first parallel it gradually decreases in density, and south of the forty-third parallel it changes in composition and character. This belt of Coast Forest is only surpassed in density by that of some portions of the redwood forest of the California coast. The red fir, the great tide-land spruce, the hemlock, and the red cedar (*Thuja*) reach here enormous dimensions. The wide river bottoms are lined with a heavy growth of maple, cottonwood, ash, and alder, the narrow interior valley with an open growth of oak. In this great coniferous forest the trunks of trees two or three hundred feet in height are often only separated by the space of a few feet. The ground, shaded throughout the year by the impenetrable canopy of the forest, never becomes dry; it is densely covered by a thick carpet of mosses and ferns, often of enormous size. The more open portions of this forest are choked by an impenetrable growth of various *Vaccinææ* of almost arborescent proportions, of hazel, the vine-maple, and other shrubs. The soil which has produced the maximum growth of forest in this region is, outside the river bottoms, a thin, porous gravel of glacial origin, rarely more than a few inches in depth; the luxuriance of vegetable growth, therefore, illustrates the influence of a heavy rainfall and temperate climate upon the forest.

The general character of this forest in the interior, although composed largely of the species peculiar to the coast, differs somewhat from the Coast Forest proper in composition and largely in natural features. The dense, impenetrable forest of the coast is replaced, east of the summit of the Cascade Range, by a more open growth, generally largely destitute of undergrowth. The red fir, the hemlock, and the red cedar (*Thuja*) are still important elements of the forest. Less valuable species of the Coast Forest—the white fir (*Abies grandis*), the yew, the alders, the mountain hemlock (*Tsuga Pattoniana*), the hawthorn, the buckthorn, and the white pine (*Pinus monticola*)—are still represented. The latter, a local species upon the coast, only reaches its greatest development toward the eastern limit of this region, here forming considerable and important forests. Other species peculiar to the Coast Forest, the maples, the ash, the oak, the arbutus, and the Alaska cedar, do not extend east of the Cascades. The tide-

land spruce is replaced by an allied species of the interior region. The widely-distributed yellow pine (*Pinus ponderosa*), barely represented in the northern portions of the immediate Coast Forest, becomes east of the mountains one of the most important and characteristic elements of the forest. The Coast Forest south of the forty-third degree of latitude changes in composition. The tide-land spruce, the hemlock, and the *Thuya* are gradually replaced by more southern species. The sugar pine (*P. Lambertiana*) here first appears. The California laurel (*Umbellularia*) covers with magnificent growth the broad river bottoms. The *Libocedrus*, several oaks, and the chinquapin here reach the northern limits of their distribution. The change from the northern to the southern forest is marked by the appearance of the Port Orford cedar (*Chamaecyparis Lawsoniana*), adding variety and value to the forests of the southern Oregon coast. Farther south, near the northern boundary of California, the redwood forests (*Sequoia*) appear.

The Coast Forest of California will be most conveniently discussed under three subdivisions: the forest of the Coast Range, the forest of the western slope of the Sierra Nevada, which, toward the northern boundary of the state, extends to the coast, covering the mass of mountains which here unite the Sierra Nevada and the Coast Range; and, third, the open forest of the long, narrow valleys lying between the Coast Range and the Sierra Nevada, south of this northern connection. The important feature of the Coast Range, as far south as the thirty-seventh degree of latitude, is the belt of redwood occupying an irregular, interrupted strip of territory facing the ocean, and hardly exceeding thirty miles in width at the points of its greatest development. The heaviest growth of the redwood forest occurs north of the bay of San Francisco, and here, along the slopes and bottom of the narrow cañons of the western slope of the Coast Range, the maximum productive capacity of the forest is reached. No other forest of similar extent equals in the amount of material which they contain the groups of redwood scattered along the coast of northern California. The red fir reaches, in the California Coast Range, a size and value only surpassed in the more northern forests of the coast; the yellow pine is an important tree in the northern portions of this region, and here flourish other species of the genus endemic to this region. The forest of the Coast Range is marked by the presence within its limits of several species of singularly restricted distribution. *Ocupressus macrocarpa* and *Pinus insignis* are confined to a few isolated groves upon the shores of the bay of Monterey; *Abies bracteata* occupies three or four cañons high up in the Santa Lucia mountains; it is found nowhere else; and *Pinus Torreyana*, the most local arborescent species of North America, has been detected only in one or two small groups upon the sand-dunes just north of the bay of San Diego. The characteristic forest of the Coast Range is checked from farther southern development, a little below the thirty-fifth parallel, by insufficient moisture; the scanty forests which clothe the high declivities of the Coast Range farther south belong in composition to the Sierra forests.

The heavy forest which covers the western slopes of the Sierra Nevada, a forest only surpassed in density by the redwood belt of the coast and the fir forest of Puget sound, occupies, in its greatest development, a belt situated between 4,000 and 8,000 feet elevation. This forest belt extends from about the base of mount Shasta at the north to the thirty-fifth parallel; farther south it diminishes in density and disappears upon the southern ridges of the Coast Range just north of the southern boundary of California. Its greatest width occurs in northern California, where to the south of mount Shasta the Sierra system is broken down into a broad mass of low ridges and peaks. The characteristic species of this forest is the great sugar pine (*P. Lambertiana*), which here reaches its greatest development and value, and gives unsurpassed beauty to this mountain forest. With the sugar pine are associated the red fir, the yellow pine, two noble *Abies*, the *Libocedrus*; and, toward the central part of the state, the great *Sequoia*, appearing first in small isolated groups, and then, farther south, near the headwaters of Kern river, in a narrow belt extending more or less continuously for several miles. This heavy forest of the Sierras, unlike the forest which farther north covers the western flanks of the Cascade Range, is almost destitute of undergrowth and young trees. It shows the influence of a warm climate and unevenly distributed rainfall upon forest growth. The trees, often remote from one another, have attained an enormous size, but they have grown slowly. Above this belt the Sierra forest stretches upward to the limits of tree growth. It is here subalpine and alpine in character and of little economic value. Different pines and firs, the mountain hemlock, and the western juniper are scattered in open stretches of forest upon the high ridges of the Sierras. The forest below the belt of heavy growth gradually becomes more open. Individual trees are smaller, while the number of species increases. The small pines of the upper foot-hills are mingled with oaks in considerable variety. These gradually increase in number. Pines are less frequent and finally disappear.

The forest of the valleys is composed of oaks, the individuals often widely scattered and of great size, but nowhere forming a continuous, compact growth. The Coast Forest of the Pacific region, unsurpassed in density, is composed of a comparatively small number of species, often attaining enormous size. It presents the same general features throughout its entire extent, except as modified by the climatic conditions of the regions which it covers. The species which compose this forest range through nearly 26 degrees of latitude, or northern species, are replaced in the south by closely allied forms; and, as in the Atlantic region, the southern species far exceed in number those peculiar to the north.

The *Interior Forest* extends from the southern limits of the northern subarctic forest to the plateau of northern Mexico; it occupies the entire region between the eastern limits of the Pacific Coast Forest and the extreme western limits of the Atlantic region. The forests of this entire region, as compared with the forests east and west of it, are stunted and remarkable in their poverty of composition. They are confined to the high slopes

and cañons of the numerous mountain ranges composing the interior region, while the valleys are treeless, or, outside of the narrow river bottoms, nearly treeless. The interior forest attains its greatest development and considerable importance upon the western slope of the California Sierras and upon the flanks of the high peaks of the southern Rocky Mountain system, from Colorado, where the timber line reaches an extreme elevation of 13,500 feet, to southern New Mexico and western Arizona. The minimum in North American forest development, outside the absolutely treeless regions, both in the number of species and in the proportion of forest to entire area, is found south of the Blue mountains of Oregon, in the arid region between the Wahsatch mountains and the Sierra Nevada, known as the Great Basin. Here the open, stunted forest is confined to the highest ridges and slopes of the infrequent cañons of the low mountain ranges which occupy, with a general north and south trend, this entire region. The individuals which compose this forest are small, although often of immense age, and everywhere show the marks of a severe struggle for existence. Seven arborescent species only have been detected in the forests of the northern and central portions of this region. The mountain mahogany (*Cercocarpus*), the only broad-leaved species of the region, with the exception of the aspen, which throughout the entire interior region borders, above an elevation of 8,000 feet, all mountain streams, reaches here its greatest development. This tree, with the nut pine (*Pinus monophylla*), characterizes this region. Stunted junipers are scattered over the lowest slopes of the mountains, or farther south often cross the high valleys, and cover with open growth the *mesas*, as the lower foot-hills are locally known. An open forest of arborescent yuccas (*Yucca brevifolia*) upon the high Mojave plateau is a characteristic and peculiar feature of the flora of this interior region. The red fir and the yellow pine, widely distributed throughout the Pacific region, do not occur upon the mountain ranges of the Great Basin.

The heavy forests of the interior region, found along the western slopes of the California Sierras and upon the Rocky Mountain system, are, for the most part, situated south of the forty-second degree of latitude. The forests of the whole northern interior portion of the continent, outside the region occupied in the northern Rocky mountains by the eastern development of the Coast Forest, feel the influence of insufficient moisture; the number of species of which they are composed is not large; the individuals are often small and stunted, while the forests are open, scattered, without undergrowth, and confined to the cañons and high slopes of the mountains. The most generally distributed species of this northern region, a scrub pine (*Pinus Murrayana*), occupies vast areas, almost to the exclusion of other species, and is gradually taking possession of ground cleared by fire of more valuable trees. South of the fifty-second parallel the red fir (*Pseudotsuga*) and the yellow pine (*Pinus ponderosa*) appear; with them is associated, in the Blue mountains and in some of the ranges of the northern Rocky mountains, the western larch (*Larix occidentalis*), the largest and most valuable tree of the Columbian basin.

The forest covering the eastern slope of the Sierra Nevada consists almost exclusively of various species of pine, often of great size and value. The characteristic species of this region are the yellow pine and the closely-allied *Pinus Jeffreyi*, here reaching its greatest development. The red fir is absent from this forest, while the oaks, multiplied in many forms on the western slopes of these mountains, have here no representative.

The forests of the southern Rocky Mountain region, less heavy and less generally distributed than those of the western slope of the Sierras, are, as compared with those of the Great Basin, heavy, dense, and valuable. They owe their existence to the comparatively large precipitation of moisture distributed over this elevated region. The characteristic species of the Colorado mountains is a spruce (*Picea Engelmanni*); it forms, at between 8,000 and 10,000 feet elevation, extensive and valuable forests of considerable density and great beauty; with it are associated a balsam fir of wide northern distribution, and various alpine and subalpine species of pine; at lower elevations forests of yellow pine and red fir cover the mountain slopes, while the bottoms of the streams are lined with cottonwood, alder, and maple, or with an open growth of the white fir (*Abies concolor*), a species of the Coast Forest, here reaching the eastern limits of its distribution; the foot-hills above the treeless plain are covered with scant groves of the nut-pine (*Pinus edulis*), stunted junipers, and a small oak, which in many forms extends through a large area of the southern interior region. A forest similar in general features to that of Colorado, and largely composed of the same species, extends over the high mountains of New Mexico to those of western Texas and western and northwestern Arizona, where a heavier forest of pine covers the elevated region lying along the thirty-fifth parallel, culminating in the high forest-clad San Francisco mountains of northern Arizona.

The species of the interior Pacific region mingle along its southern borders with the species peculiar to the plateau of northern Mexico. The Pacific-Mexican Forest, although differing widely in natural features from the Atlantic-Mexican Forest, possesses several species peculiar to the two. The forests of this region are confined to the high mountains and their foot-hills, and to the banks of the rare water-courses. They disappear entirely from the Colorado desert and from the valleys and low mountain ranges of southwestern Arizona. The most important and generally distributed species peculiar to the valleys of this region is the mesquit, the characteristic species of the Atlantic-Mexican region. The suwarrow, however, the great tree cactus, is perhaps the most remarkable species of the region, giving an unusual and striking appearance to the dry *mesas* of central and southern Arizona. The high mountain ranges, extending across the boundary of the United States, between the one hundred and fifth and the one hundred and eleventh meridians, enjoy a larger and more regularly distributed rainfall than the regions east, and especially west, of these meridians. The forests which cover these southern mountain ranges are often dense and varied. Upon their summits and almost inaccessible upper slopes the firs and pines of

the Pacific region are mingled with pines, a juniper, an arbutus, and various other species peculiar to the Mexican plateau. Extensive forests of a cypress of Mexican origin also characterize this mountain vegetation. The bottoms of the cañons are lined with a dense growth of cottonwood, hackberry, a noble sycamore, an ash, a cherry, and other deciduous trees. The high foot-hills and *mesas* are covered with open groves of various oaks peculiar to the Mexican-Pacific region, here reaching, within the United States at least, their greatest development.

Such are some of the prominent forest features of North America; a dense forest, largely composed, except at the north, of a great variety of broad-leaved species, and extending from the Atlantic sea-board in one nearly unbroken sheet until checked by insufficient moisture from further western development—the forest of the Atlantic region; a forest of conifers, occupying the ranges of the great Cordilleran mountain system, unsurpassed in density in the humid climate of the coast, open and stunted in the arid interior—the forest of the Pacific region.

A more detailed examination of the distribution of North American arborescent genera and species will serve to illustrate the wealth of the forests of the Atlantic and the comparative poverty of those of the Pacific region. It will show, too, more clearly how widely the forests of these two great regions differ in composition.

DISTRIBUTION OF GENERA.

The forests of North America contain arborescent representatives of 158 genera; 142 genera occur in the Atlantic and 59 genera in the Pacific region. Of the Atlantic genera, 48 are not represented in the United States outside the semi-tropical region of Florida.

The following table illustrates the distribution of these genera; the genera of semi-tropical Florida are designated by a *.

	Genera represented by arborescent species in the Atlantic region.	Genera represented by arborescent species in the Pacific region.		Genera represented by arborescent species in the Atlantic region.	Genera represented by arborescent species in the Pacific region.
Magnolia	✓		Eysenhardtia	✓	✓
Liriodendron	✓		Dalea		✓
Asimina	✓		Robinia	✓	✓
*Anona	✓		Olneya		✓
*Capparis	✓		*Piscidia	✓	
*Canella	✓		Cladrastis	✓	
*Clusia	✓		Sophora	✓	
Gordonia	✓		Gymnocladus	✓	
Fremontia		✓	Gleditschia	✓	
Tilia	✓		Parkinsonia	✓	✓
*Byrsonima	✓		Cercis	✓	
*Guaiacum	✓		Prosopis	✓	✓
Porlieria	✓		Leucæna	✓	
Xanthoxylum	✓		Acacia	✓	✓
Ptelia	✓	✓	*Lysiloma	✓	
Canotia		✓	*Pithecolobium	✓	
*Simaruba	✓		*Chrysobalanus	✓	
*Bursera	✓		Prunus	✓	✓
*Amyris	✓		Vauquellinia		✓
*Swietenia	✓		Cercocarpus		✓
*Ximenia	✓		Pyrus	✓	✓
Ilex	✓		Cratægus	✓	✓
Cyrilla	✓		Heteromeles		✓
Cliftonia	✓		Amelanchier	✓	
Euonymus	✓		Hamamelis	✓	
*Myginda	✓		Liquidambar	✓	
*Schæfferia	✓		Rhizophora	✓	
*Reynosa	✓		*Conocarpus	✓	
Condalia	✓	✓	*Laguncularia	✓	
Rhamnus	✓	✓	*Calyptranthes	✓	
Ceanothus		✓	*Eugenia	✓	
*Colubrina	✓		Cereus		✓
Æsculus	✓	✓	Cornus	✓	✓
Unguadia	✓	✓	Nyssa	✓	
Sapindus	✓	✓	Sambucus	✓	✓
*Hypelate	✓		Viburnum	✓	
Acer	✓	✓	*Exostemma	✓	
Negundo	✓	✓	Pinckneya	✓	
Rhus	✓		*Genipa	✓	
Pistacia	✓		*Guettarda	✓	

	Genera represented by arbo- rescent species in the Atlantic region.	Genera represented by arbo- rescent species in the Pacific region.		Genera represented by arbo- rescent species in the Atlantic region.	Genera represented by arbo- rescent species in the Pacific region.
Vaccinium	✓		Planera	✓	
Andromeda	✓		Celtis	✓	✓
Arbutus	✓	✓	*Ficus	✓	
Oxydendrum	✓		Morus	✓	✓
Kalmia	✓		Maclura	✓	
Rhododendron	✓		Platanus	✓	✓
*Myrsine	✓		Juglans	✓	✓
*Ardisia	✓		Carya	✓	
*Jacquinia	✓		Myrica	✓	✓
*Chrysophyllum	✓		Quercus	✓	✓
*Sideroxylon	✓		Castanopsis		✓
*Dipholis	✓		Castanea	✓	
Bumelia	✓	✓	Fagus	✓	
*Mimusops	✓		Ostrya	✓	
Diospyros	✓		Carpinus	✓	
Symplocos	✓		Betula	✓	✓
Halesia	✓		Alnus	✓	✓
Fraxinus	✓	✓	Salix	✓	✓
Forestiera	✓		Populus	✓	✓
Chionanthus	✓		Libocedrus		✓
Osmanthus	✓		Thuja	✓	✓
Cordia	✓		Chamaecyparis	✓	✓
*Bourreria	✓		Cupressus		✓
*Ehretia	✓		Juniperus	✓	✓
Catalpa	✓	✓	Taxodium	✓	
Chilopsis	✓	✓	Sequoia		✓
*Crescentia	✓		Taxus	✓	✓
*Citharexylum	✓		Torreya	✓	✓
*Avicennia	✓		Pinus	✓	✓
*Pisonia	✓		Picea	✓	✓
*Coccoloba	✓		Tsuga	✓	✓
Persea	✓		Pseudotsuga		✓
*Nectandra	✓		Abies	✓	✓
Sassafras	✓		Larix	✓	✓
Umbellularia		✓	Sabal	✓	
*Drypetes	✓		Washingtonia		✓
*Sebastiania	✓		*Thrinax	✓	
*Hippomane	✓		*Oreodoxa	✓	
Ulmus	✓		Yucca	✓	✓

Arborescent species of 43 genera occur within the limits of the two regions. They are :

Ptelia.	Robinia.	Arbutus.	Quercus.	Taxus.
Condalia.	Parkinsonia.	Bumelia.	Betula.	Torreya.
Rhamnus.	Prosopis.	Fraxinus.	Alnus.	Pinus.
Æsculus.	Acacia.	Chilopsis.	Salix.	Picea.
Ungnadia.	Prunus.	Celtis.	Populus.	Tsuga.
Sapindus.	Pyrus.	Morus.	Thuja.	Abies.
Acer.	Cratægnus.	Platanus.	Chamaecyparis.	Larix.
Negundo.	Cornus.	Juglans.	Juniperus.	Yucca.
Eysenhardtia.	Sambucus.	Myrica.		

The following genera, 44 in number, of the Atlantic region, exclusive of those of semi-tropical Florida, are not represented in the Pacific forest :

Magnolia.	Cliftonia.	Rhizophora.	Forestiera.	Maclura.
Liriodendron.	Pistacia.	Nyssa.	Chionanthus.	Carya.
Asimina.	Cladrastis.	Viburnum.	Osmanthus.	Castanea.
Gordonia.	Sophora.	Pinckneya.	Cordia.	Fagus.
Tilia.	Gymnocladus.	Andromeda.	Catalpa.	Ostrya.
Porlieria.	Gleditschia.	Oxydendrum.	Persea.	Carpinus.
Xanthoxylum.	Leucæna.	Diospyros.	Sassafras.	Taxodium.
Ilex.	Hamamelis.	Symplocos.	Ulmus.	Sabal.
Cyrilla.	Liquidambar.	Halesia.	Planera.	

The following genera of the Atlantic region, 9 in number, are represented in the Pacific flora by one or more frutescent, but by no arborescent, species:

Euonymus.	Amelanchier.	Vaccinium.	Rhododendron.
Rhus.	Viburnum.	Kalmia.	Forestiera.
Cercis.			

Ptelia, *Condalia*, *Sapindus*, *Robinia*, *Bumelia*, *Celtis*, *Morus*, and *Juglans*, genera reaching their greatest development in North America in the Atlantic region, extend with a single arborescent representative into the Pacific region. *Rhamnus*, *Aesculus*, *Acer*, *Negundo*, *Prunus*, *Pyrus*, *Crataegus*, *Cornus*, *Sambucus*, *Fraxinus*, *Platanus*, *Myrica*, *Quercus*, *Betula*, *Alnus*, *Salix*, *Populus*, *Thuja*, *Chamaecyparis*, *Juniperus*, *Taxus*, *Torreya*, *Pinus*, *Picea*, *Tsuga*, *Abies*, and *Larix*, characteristic North American genera, are widely represented in the two regions.

Ungnadia, *Eysenhardtia*, *Parkinsonia*, *Prosopis*, *Acacia*, *Chilopsis*, and *Yucca*, genera of the Mexican flora, are common to the two regions.

Arbutus, a genus of the Pacific region, just reaches, with a doubtful species, the Atlantic region through western Texas.

The following genera of the Pacific region, 13 in number, have no representatives in the Atlantic region:

Fremontia.	Cercocarpus.	Castanopsis.	Sequoia.
Canotia.	Heteromeles.	Libocedrus.	Pseudotsuga.
Olneya.	Umbellularia.	Cupressus.	Washingtonia.
Vauquelinia.			

The following genera of the Pacific, 3 in number, are represented in the Atlantic region by frutescent species:

Ceanothus.	Dalea.	Cereus.
------------	--------	---------

The Atlantic forest, exclusive of semi-tropical Florida, contains 45 genera entirely unrepresented in the Pacific region and 7 genera without Pacific arborescent representatives. The Pacific forest contains 13 genera unrepresented in the Atlantic region and 3 genera without Atlantic arborescent representatives.

The following genera of the Mexican region, 14 in number, are not elsewhere represented in North America. Genera with arborescent representatives in both the Atlantic and Pacific-Mexican regions are designated by a star (*):

Portiera.	Pistacia.	Olneya.	Acacia.	*Chilopsis.
Canotia.	*Eysenhardtia.	*Parkinsonia.	Vauquelinia.	Washingtonia.
*Ungnadia.	Dalea.	Leucæna.	Cereus.	

Portiera and *Leucæna* belong to the Atlantic; *Canotia*, *Dalea*, *Olneya*, *Vauquelinia*, *Cereus*, and *Washingtonia* to the Pacific region.

DISTRIBUTION OF SPECIES.

In the forests of North America 412 arborescent species have been detected; of these, 292 species belong to the Atlantic region, and 153 occur within the limits of the Pacific region. Species common to the two regions are rare; they are principally confined to the subarctic Northern Forest and to the narrow belt along the southern boundary of the United States.

The following species, 10 in number, cross the continent:

<i>Prosopis juliflora</i> .	<i>Sambucus Mexicana</i> .	<i>Salix longifolia</i> .	<i>Populus balsamifera</i> .	<i>Picea alba</i> .
<i>Pyrus sambucifolia</i> .	<i>Betula papyrifera</i> .	<i>Populus tremuloides</i> .	<i>Juniperus Virginiana</i> .	<i>Yucca baccata</i> .

Prosopis juliflora, *Sambucus Mexicana*, and *Yucca baccata* belong to the Mexican flora of the south; *Salix longifolia* also belongs here, although extending northward into the Atlantic and through the Pacific Coast region of the United States. *Populus balsamifera*, *Betula papyrifera*, and *Picea alba* belong to the Northern Forest. *Pyrus sambucifolia*, *Populus tremuloides* and *Juniperus Virginiana* are widely distributed through the central portions of the Atlantic and Pacific regions; they are the only really continental arborescent species.

The following species of the Atlantic region, 15 in number, extend from the Atlantic into the Pacific region:

<i>Ptelia trifoliata</i> .	<i>Negundo aceroides</i> .	<i>Crataegus tomentosa</i> .	<i>Quercus Emoryi</i> .
<i>Condalia obovata</i> .	<i>Parkinsonia aculeata</i> .	<i>Fraxinus viridis</i> .	<i>Alnus incana</i> .
<i>Sapindus marginatus</i> .	<i>Prunus Americana</i> .	<i>Celtis occidentalis</i> .	<i>Salix nigra</i> .
<i>Ungnadia speciosa</i> .	<i>Prunus Pennsylvanica</i> .	<i>Morus microphylla</i> .	

Ptelia trifoliata, a widely distributed species of the Atlantic region, extends through western Texas into the extreme southeastern portion of the Pacific region. *Condalia obovata*, *Ungnadia speciosa*, *Parkinsonia aculeata*, *Morus microphylla*, and *Quercus Emoryi*, of the Atlantic-Mexican forest, extend into the Pacific-Mexican region. *Sapindus marginatus*, of the southern Atlantic region, extends through western Texas to the Pacific-Mexican region. *Prunus Americana*, *Prunus Pennsylvanica*, and *Alnus incana*, widely distributed through the northern portions of the Atlantic region, just reach the eastern limits of the central Pacific region.

Negundo aceroides, *Crataegus tomentosa*, *Fraxinus viridis*, and *Celtis occidentalis* are widely distributed through the interior Pacific region, although nowhere reaching the coast.

The following species of the Pacific region, 8 in number, extend through the Mexican into the Atlantic region :

<i>Eysenhardtia orthocarpa.</i>	<i>Acacia Greggii.</i>	<i>Chilopsis saligna.</i>	<i>Juniperus occidentalis.</i>
<i>Prosopis pubescens.</i>	<i>Fraxinus pistaciæfolia.</i>	<i>Juglans rupestris.</i>	<i>Juniperus pachyphloea.</i>

Juglans rupestris and *Juniperus occidentalis* reach their greatest development in the Pacific Coast region, and extend through the Pacific-Mexican region into western Texas; no other species are common to the Pacific Coast forest and the Atlantic-Mexican region. The 6 remaining Pacific-Atlantic species belong to the Pacific-Mexican region, just reaching western Texas.

The following species of the Southern Pacific region extends into the Atlantic region :

Salix amygdaloides.

The following species of the Pacific forest, 12 in number, endemic to the interior arid region, do not extend beyond its limits :

<i>Acer grandidentatum.</i>	<i>Cratægus rivularis.</i>	<i>Populus angustifolia.</i>	<i>Pinus monophylla.</i>
<i>Robinia Neo-Mexicana.</i>	<i>Fraxinus anomala.</i>	<i>Pinus flexilis.</i>	<i>Picea pungens.</i>
<i>Cercocarpus ledifolius.</i>	<i>Quercus undulata.</i>	<i>Pinus edulis.</i>	<i>Yucca brevifolia.</i>

A detailed examination of the distribution of the arborescent species composing the North American forests shows that—

Magnolia is represented by seven Atlantic species, with the center of its distribution in the southern Alleghany region.

Liriodendron is represented by a single species, widely-distributed through the eastern and central portions of the Atlantic region.

Asimina is represented by a single widely-distributed arborescent species and by three frutescent species of the Atlantic region.

Anona, *Capparis*, *Oanella*, and *Olusia* are represented each by a single semi-tropical species.

Gordonia is represented by two species of the southern Atlantic region, one of wide distribution, the other rare and local.

Fremontia, a genus endemic to the Pacific region, is represented by a single species of the southern Pacific Coast region.

Tilia is represented by two Atlantic species, with its center of distribution in the southern Alleghany region.

Byrsonima is represented by a single semi-tropical species.

Guaiacum is represented by a single semi-tropical species.

Portiera is represented by a single species of the Atlantic-Mexican region.

Xanthoxylum is represented by two species of the Atlantic region, by a semi-tropical species, and by a second semi-tropical species which reaches the Atlantic-Mexican region.

Ptelia is represented by a single arborescent species of wide distribution in the Atlantic, reaching also the Pacific region, where a frutescent species occurs, and by a second frutescent species of the south Atlantic region.

Canotia, a genus endemic to the Pacific-Mexican region, is represented by a single species.

Simaruba, *Amyris*, *Swietenia*, *Ximenia*, are each represented by a single semi-tropical species.

Bursera is represented by a single semi-tropical species and by a second frutescent species of the Pacific-Mexican region.

Ilex, an Atlantic genus, is represented by four arborescent and several frutescent species, with its center of distribution in the southern Atlantic region.

Cyrilla and *Cliftonia* are each represented by a single species of the southern Atlantic region.

Euonymus is represented by a widely-distributed arborescent species in the Atlantic, and by a frutescent species in both the Atlantic and the Pacific regions.

Myginda, *Schæfferia*, and *Reynosia* are each represented by a single semi-tropical species.

Condalia is represented by one semi-tropical and by one species of the Atlantic-Mexican reaching the Pacific-Mexican region.

Rhamnus is represented by one arborescent and by one frutescent species in the Atlantic, by two arborescent and one frutescent species in the Pacific region, and by one frutescent species common to the two regions.

Ceanothus is represented by a single arborescent species in the Pacific Coast region and by several frutescent species widely distributed through the Atlantic and the Pacific regions.

Colubrina is represented by a single semi-tropical species.

Æsculus is represented by two arborescent and by three frutescent species in the Atlantic, and by an arborescent species in the Pacific region.

Ungnadia, an endemic genus of the Atlantic-Mexican region, and just reaching the Pacific-Mexican region, is represented by a single species.

Sapindus is represented by one species widely distributed through the southern Atlantic, and reaching the Pacific region, and by one semi-tropical species.

Acer is represented by five Atlantic and four Pacific species.

Negundo is represented by one species widely distributed through the Atlantic and the Pacific regions and by a second species in the Pacific region.

Rhus is represented by five arborescent species in the Atlantic and by several frutescent species in both the Atlantic and the Pacific regions.

Pistacia is represented by a single species in the Atlantic-Mexican region.

Eysenhardtia is represented by a single arborescent species in the Pacific-Mexican, extending into the Atlantic-Mexican region, where a second frutescent species occurs.

Dalea is represented by a single arborescent species in the Pacific-Mexican and by numerous frutescent and herbaceous species in the Atlantic and the Pacific regions.

Robinia, with its center of distribution in the southern Alleghany region, is represented by two arborescent and one frutescent species in the Atlantic and by one arborescent species in the Pacific region.

Olneya, an endemic genus of the Pacific-Mexican region, is there represented by a single species.

Piscidia is represented by a single semi-tropical species.

Ciadrastis is represented by a single local species in the southern Atlantic region.

Sophora is represented by a species in the southern Atlantic and by a second species in the Atlantic-Mexican region, and by four frutescent or suffrutescent species.

Gymnocladus is represented by a single species in the central Atlantic region.

Gleditschia is represented by two widely-distributed species in the Atlantic region.

Parkinsonia is represented by an arborescent species common to the Atlantic and the Pacific-Mexican regions, by two arborescent and one frutescent species in the Pacific-Mexican, and by a frutescent species in the Atlantic-Mexican region.

Cercis is represented by a widely-distributed species in the Atlantic, by a second species in the Atlantic-Mexican, and a frutescent species of the California Coast region.

Prosopis is represented by two arborescent species common to the Atlantic and the Pacific-Mexican regions, and by two frutescent species.

Leucæna is represented by two species in the Atlantic-Mexican region.

Acacia is represented by two arborescent species in the Atlantic-Mexican, by one arborescent species of the Pacific-Mexican extending into the Atlantic-Mexican region, and by several frutescent species widely distributed through the two regions.

Lysiloma is represented by a single semi-tropical species.

Pithecolobium is represented by a single polymorphous arborescent species of semi-tropical Florida, and by a shrubby species of the Mexican Boundary region.

Chrysobalanus is represented by one arborescent and one frutescent semi-tropical species.

Prunus is represented by seven arborescent species in the Atlantic region; of these, one is semi-tropical and two extend into the Pacific region. This genus is represented in the Pacific region by four species, of which one belongs to the Mexican region, and by several frutescent species.

Vauquelinia, an endemic genus of the Pacific-Mexican region, is there represented by a single species.

Cercocarpus is represented by two widely-distributed species in the Pacific region.

Pyrus is represented by one species common to both Atlantic and Pacific, by three arborescent and one frutescent species in the Atlantic, and by one arborescent species in the Pacific region.

Cratægus is represented by twelve arborescent and frutescent species in the Atlantic, of which one extends into the Pacific region, and by two species in the Pacific region.

Heteromeles is represented by a single species in the Pacific Coast region.

Amelanchier is represented by one arborescent species in the Atlantic and by one frutescent species in the Pacific region.

Hamamelis and *Liquidambar* are each represented by one widely-distributed species in the Atlantic region.

Rhizophora is represented by a single species in the southern Atlantic region.

Conocarpus, *Laguncularia*, and *Calyptranthes* are each represented by a single semi-tropical species.

Eugenia is represented by five semi-tropical species.

Cereus is represented by a single arborescent species in the Pacific and by several frutescent species in the Atlantic and Pacific regions.

Cornus is represented by two arborescent species in the Atlantic, by a single arborescent species in the Pacific region, and by several frutescent and herbaceous species in the two regions.

Nyssa is represented by three species in the Atlantic region.

Sambucus is represented by one arborescent species of wide distribution in the Pacific, by one species in the Pacific-Mexican extending into the Atlantic-Mexican, by a frutescent species in the Atlantic, by a second frutescent species in the Pacific, and by a frutescent species common to the Atlantic and Pacific regions.

Viburnum is represented by two arborescent species in the Atlantic and by several frutescent species in the Atlantic and the Pacific regions.

Exostemma is represented by a single semi-tropical species.

Pinckneya, an endemic genus of the southern Atlantic region, is there represented by a single species.

Genipa is represented by a single semi-tropical species.

Guettarda is represented by one arborescent and by one frutescent semi-tropical species.

Vaccinium is represented by one arborescent species in the Atlantic and by several frutescent species in the Atlantic and the Pacific regions.

Andromeda is represented by an arborescent and several frutescent species in the Atlantic region.

Arbutus is represented by one species in the Pacific Coast, by a second species in the Pacific-Mexican, and by one species in the Atlantic-Mexican region.

Oxydendrum, an endemic genus of the Atlantic region, is there represented by a single species.

Kalmia is represented by one arborescent species and by three frutescent species in the Atlantic region, of which one extends to the Pacific region.

Rhododendron is represented by one arborescent and by several frutescent species in the Atlantic and by several frutescent species in the Pacific region.

Myrsine, *Ardisia*, *Jacquinia*, *Chrysophyllum*, *Sideroxylon*, and *Dipholis* are each represented by a single semi-tropical species.

Bumelia is represented by four species in the Atlantic and by one species in the Pacific-Mexican region.

Mimusops is represented by one semi-tropical species.

Diospyros is represented by one species in the Atlantic and by one in the Atlantic-Mexican region.

Symplocos is represented by one species in the southern Atlantic region.

Halesia is represented by two arborescent and by one frutescent species in the southern Atlantic region.

Fraxinus, with its center of distribution in the southern Atlantic region, is represented by seven species in the Atlantic, of which one extends into the Pacific region, and one belongs to the Mexican region, and by three arborescent and one frutescent species in the Pacific, of which one belongs to the Mexican region.

Forestiera is represented by one arborescent and seven frutescent species in the Atlantic region, of which one reaches the Mexican-Pacific region.

Chionanthus and *Osmanthus* are each represented by a single species in the southern Atlantic region.

Cordia is represented by one arborescent and by one frutescent semi-tropical species and by one arborescent and one frutescent species in the Atlantic-Mexican region.

Bourreria and *Ehretia* are each represented by a single semi-tropical species.

Catalpa is represented by two species in the southern Atlantic region.

Chilopsis is represented by a single species in the Pacific-Mexican region, extending into the Atlantic-Mexican region.

Crescentia, *Oitharexylum*, and *Avicennia* are each represented by a single semi-tropical species.

Pisonia is represented by one arborescent and by two frutescent semi-tropical species.

Coccoloba is represented by two semi-tropical species.

Persea is represented by one species in the southern Atlantic region.

Nectandra is represented by one semi-tropical species.

Sassafras is represented by one widely-distributed species in the Atlantic region.

Umbellularia is represented by a single species in the Pacific Coast region.

Drypetes, *Sebastiania*, and *Hippomane* are each represented by a single semi-tropical species.

Ulmus, with its center of distribution in the Mississippi basin, is represented in the Atlantic region by five species.

Planera is represented by a single species in the southern Atlantic region.

Celtis is represented by a single polymorphous species of wide distribution in the Atlantic region, extending into the Pacific region, and by a frutescent species common to the Atlantic-Mexican and the Pacific-Mexican regions.

Ficus is represented by three semi-tropical species.

Morus is represented by one widely-distributed species in the Atlantic region, and by one species in the Atlantic-Mexican, extending into the Pacific-Mexican region.

Maclura is represented by a single local species in the southern Atlantic region.

Platanus is represented by one widely-distributed species in the Atlantic region, by a species in the Pacific coast, and by a species in the Pacific-Mexican region.

Juglans is represented by two widely-distributed species in the Atlantic region and by a species in the Pacific coast, extending through the Pacific-Mexican into the Atlantic-Mexican region.

Carya, an endemic genus of the Atlantic region, with its center of distribution west of the Mississippi river, is represented by seven species.

Myrica is represented by one arborescent and two frutescent species in the Atlantic region and by one arborescent species in the Pacific Coast region.

Quercus, with its center of most important distribution in the basin of the lower Ohio river, is represented in the Atlantic region by twenty-four arborescent species, of which one, belonging to the Mexican region, extends into the Pacific-Mexican region; and in the Pacific region by twelve arborescent species, of which one belongs to the interior and four to the Mexican region, and by two frutescent species.

Castanopsis is represented by a single species in the Pacific Coast region.

Castanea is represented by two species in the Atlantic region.

Fagus, *Ostrya*, and *Carpinus* are each represented by a single widely-distributed species in the Atlantic region.

Betula, with its center of distribution in the northern Atlantic region, is represented by one arborescent and by one frutescent species common to the Atlantic and the Pacific regions, by four arborescent and one frutescent species in the Atlantic region, and by one arborescent species in the Pacific region.

Alnus is represented by three arborescent species in the Atlantic, of which one extends to the Pacific region, by three arborescent species in the Pacific region, and by two frutescent species common to the Atlantic and the Pacific regions.

Salix is represented in the Atlantic region by five arborescent species, of which three are found in the Pacific region, and by many frutescent species. This genus is represented in the Pacific region by ten arborescent and by many frutescent species.

Populus is represented by two species common to the Atlantic and the Pacific regions, by three species in the Atlantic region, and by three species in the Pacific region.

Libocedrus is represented by a single species in the Pacific Coast region.

Thuja is represented by one species in the Atlantic and by one species in the Pacific region.

Chamaecyparis is represented by one species in the Atlantic and by two species in the Pacific Coast region.

Cupressus is represented by four species in the Pacific region, of which three occur in the coast and one in the Mexican region.

Juniperus is represented by one arborescent species in the Atlantic region, by three arborescent species in the Pacific, of which one belongs to the Pacific-Mexican and one extends to the Atlantic-Mexican region, and by two frutescent species common to both regions.

Taxodium is represented by a single species in the southern Atlantic region.

Sequoia, an endemic genus of the Pacific Coast region, is there represented by two species.

Taxus is represented by an exceedingly local arborescent species in the southern Atlantic region, by a frutescent species in the northern Atlantic region, and by an arborescent species in the Pacific Coast region.

Torreya is represented by a single exceedingly local arborescent species in the southern Atlantic region and by a single species in the Pacific Coast region.

Pinus, with its center of distribution in the southern Pacific Coast region, is represented by thirteen species in the Atlantic and by twenty-two species in the Pacific region, of which three belong to the interior and four to the Mexican region.

Picea is represented by one species common to the Atlantic and the Pacific regions, by one species in the Atlantic, and by three species in the Pacific region, of which one belongs to the interior region.

Tsuga is represented by two species in the Atlantic and by two species in the Pacific region.

Pseudotsuga, an endemic genus of the Pacific region, is there represented by a single widely-distributed species.

Abies is represented by one widely-distributed and by one exceedingly local species in the Atlantic region and by seven species in the Pacific region, of which one is exceedingly local.

Larix is represented by one species in the Atlantic and by two species in the Pacific region.

Sabal is represented by a single species in the southern Atlantic region.

Washingtonia is represented by a single species in the Pacific Mexican region.

Thrinax is represented by two semi-tropical species, and *Oreodoxa* by one.

Yucca is represented by one arborescent and one frutescent species common to the Atlantic and the Pacific regions, by one arborescent and by two frutescent species in the Atlantic, and by two arborescent and by one frutescent species in the Pacific region.

A CATALOGUE
OF THE
FOREST TREES OF NORTH AMERICA, EXCLUSIVE OF MEXICO,
WITH
REMARKS UPON THEIR SYNONYMY, BIBLIOGRAPHICAL HISTORY,
DISTRIBUTION, ECONOMIC VALUE, AND USES.

FOREST TREES OF NORTH AMERICA.

Species which grow from the ground with a single stem, either wholly or over a large portion of the area of their distribution, are admitted as trees into the following catalogue, without reference to the height or size they may attain.

The line which divides trees from shrubs is entirely arbitrary, and is often unsatisfactory in application. A separation of this nature, however, based upon habit rather than upon size, is perhaps less objectionable, all things considered, than any other, and serves at least to keep this catalogue within reasonable limits.

The word "compact", used in the description of various woods mentioned in the catalogue, indicates that they show no tendency to check or open in drying, and does not refer to their structure.

CATALOGUE OF FOREST TREES.

MAGNOLIACEÆ.

1.—*Magnolia grandiflora*, Linnæus,

Spec. 2 ed. 755.—Marshall, Arbustum, 84.—Am. Gewach. t. 185, 186.—Walter, Fl. Caroliniana, 158.—Gærtner, Fruct. i, 343, t. 70.—B. S. Barton, Coll. i, 13; ii, 20.—Aiton, Hort. Kew. ii, 251; 2 ed. iii, 329.—Bartram, Travels, 2 ed. 82.—Lamarck, Dict. iii, 672; Ill. iii, 35, t. 490.—Mönch, Meth. 274.—Willdenow, Spec. ii, 1255; Enum. i, 579.—Michaux, Fl. Bor.-Am. i, 327.—Nouveau Duhamel, ii, 219, t. 65.—Desfontaines, Hist. Arb. ii, 5.—Robin, Voyages, iii, 265.—Andrews, Bot. Rep. viii, t. 518.—Titford, Hort. Bot. Am. 76.—Michaux f. Hist. Arb. Am. iii, 71, t. 1; N. American Sylva, 3 ed. ii, 8, t. 51.—Pursh, Fl. Am. Sept. ii, 380.—Nuttall, Genera, ii, 18; Sylva, i, 81; 2 ed. i, 96.—De Candolle, Syst. i, 450; Prodr. i, 80.—Hayne, Dend. Fl. 116.—Elliott, Sk. ii, 36.—Loddiges, Bot. Cab. t. 814.—Sprengel, Syst. ii, 642.—Audubon, Birds, t. 5, 32.—Rafinesque, Med. Bot. ii, 32.—Don, Miller's Dict. i, 82.—Eaton, Manual, 6 ed. 218.—Croom in Am. Jour. Sci. 1 ser. xxvi, 314.—London, Arboretum, i, 261 & t.—Hooker, Jour. Bot. i, 188.—Eaton & Wright, Bot. 312.—Torrey & Gray, Fl. N. America, i, 42.—Spach, Hist. Veg. vii, 470.—Dietrich, Syn. iii, 308.—Seringe, Fl. Jard. iii, 225.—Darby, Bot. S. States, 210.—Cooper in Smithsonian Rep. 1858, 250.—Chapman, Fl. S. States, 13.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 66.—Wood, Cl. Book, 214; Bot. & Fl. 24.—Porcher, Resources S. Forests, 38.—Baillon, Hist. Pl. i, 133, f. 165-169.—Koch, Dendrologie, i, 367.—Young, Bot. Texas, 148.—Vasey, Cat. Forest Trees, 6.

M. Virginiana, var. *β. foetida*, Linnæus, Spec. 1 ed. 536, in part.

M. grandiflora, var. *elliptica* and *obovata*, Pursh, Fl. Am. Sept. ii, 380.

M. grandiflora, var. *lanceolata*, Pursh, Fl. Am. Sept. ii, 380.—Bot. Mag. t. 1952.—Eaton, Manual, 6 ed. 218.

BIG LAUREL. BULL BAY.

Cape Fear river, North Carolina, south near the coast to Mosquito inlet, and Tampa bay, Florida; basin of the Mississippi river south of latitude 32° 30', extending westward to southwestern Arkansas, and along the Texas coast to the valley of the Brazos river.

One of the most magnificent trees of the Atlantic forest, evergreen, 18 to 27 meters in height, with a trunk 0.60 to 1.20 meter in diameter; reaching its greatest development on the "bluff" formations along the eastern bank of the Mississippi river from Vicksburg to Natchez, and of western Louisiana.

Wood heavy, hard, not strong, close-grained, compact, easily worked, satiny; medullary rays very numerous, thin; color, creamy white or often light brown, the heavier sap-wood nearly white; specific gravity, 0.6360; ash, 0.53; little used except as fuel; suitable for interior finish, fine cabinet work, etc.

2.—*Magnolia glauca*, Linnæus,

Spec. 2 ed. 755.—Kalm, Travels, English ed. i, 204.—Schœpf, Mat. Med. Am. 91.—Marshall, Arbustum, 83.—Wangenheim, Amer. 60, t. 19, f. 46.—Walter, Fl. Caroliniana, 158.—B. S. Barton, Coll. i, 13; ii, 20.—Lamarck, Dict. iii, 674.—Aiton, Hort. Kew. ii, 251; 2 ed. iii, 329.—Mönch, Meth. 274.—Willdenow, Spec. ii, 1256; Enum. i, 579.—Schkuhr, Handb. ii, 1441, t. 148.—Michaux, Fl. Bor.-Am. i, 327.—Nouveau Duhamel, ii, 223, t. 66.—Desfontaines, Hist. Arb. ii, 5.—Titford, Hort. Bot. Am. 76.—Bonpland, Pl. Malm. 103, t. 42.—Michaux f. Hist. Arb. Am. iii, 77, t. 2; N. American Sylva, 3 ed. ii, 12, t. 52.—Pursh, Fl. Am. Sept. ii, 381.—Eaton, Manual, 6 ed. 218.—Bigelow, Med. Bot. ii, 67, t. 27; Fl. Boston. 3 ed. 244.—Nuttall, Genera, ii, 18.—Barton, Prodr. Fl. Philadelph. 59; Med. Bot. i, 77, t. 7; Compend. Fl. Philadelph. ii, 17.—Loddiges, Bot. Cab. t. 215.—De Candolle, Syst. i, 452; Prodr. i, 80.—Hayne, Dend. Fl. 116.—Elliott, Sk. ii, 37.—Bot. Mag. t. 2164.—Sprengel, Syst. 642.—Torrey, Compend. Fl. N. States, 221; Fl. N. York, i, 17, t. 5.—Audubon, Birds, t. 118.—Rafinesque, Med. Bot. ii, 34.—Don, Miller's Dict. i, 82.—Eaton, Manual, 6 ed. 218.—Hooker, Jour. Bot. i, 188.—Beck, Bot. 15.—Sertum Botanicum, v & t.—Eichenbach, Fl. Exot. v, 37, t. 342.—Lindley, Fl. Med. 23.—Eaton & Wright, Bot. 312.—Torrey & Gray, Fl. N. America, i, 42.—Spach, Hist. Veg. vii, 473.—Dietrich, Syn. iii, 308.—Griffith, Med. Bot. 96, f. 56.—London, Arboretum, i, 267 & t.—Emerson, Trees Massachusetts, 527; 2 ed. ii, 603 & t.—Seringe, Fl. Jard. iii, 226.—Gray, Genera, i, 61, t. 23; Manual N. States, 5 ed. 49.—Schnizlein, Icon. t. 176.—Darlington, Fl. Cestrica, 3 ed. 8.—Darby, Bot. S. States, 211.—Cooper in Smithsonian Rep. 1858, 250.—Chapman, Fl. S. States, 13.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 66.—Lesquereux in Owen's 2d Rep. Arkansas, 374.—Wood, Cl. Book, 214; Bot. & Fl. 24.—Porcher, Resources S. Forests, 36.—Koch, Dendrologie, i, 369.—Young, Bot. Texas, 148.—Vasey, Cat. Forest Trees, 6.

M. Virginiana, var. *a. glauca*, Linnæus, Spec. 1 ed. 535.

M. fragrans, Salisbury, Prodr. 379.—Rafinesque, Fl. Ludoviciana, 91; Med. Bot. ii, 32.

M. longifolia, Sweet, Hort. Brit. 11.—Don, Miller's Dict. i, 83.—Dietrich, Syn. iii, 308.

M. glauca, var. *latifolia*, Aiton, Hort. Kew. 2 ed. iii, 350.—Pursh, Fl. Am. Sept. ii, 381.—Eaton, Manual, 6 ed. 218.

M. glauca, var. *longifolia*, Aiton, Hort. Kew. 2 ed. iii, 330.—Pursh, Fl. Am. Sept. ii, 381.—Rafinesque, Fl. Ludoviciana, 91.—Hayne, Dend. Fl. 116.—Eaton, Manual, 6 ed. 218.

SWEET BAY. WHITE BAY. BEAVER TREE. WHITE LAUREL. SWAMP LAUREL.

Cape Ann, Massachusetts; New Jersey southward, generally near the coast, to bay Biscayne and Tampa bay, Florida; basin of the Mississippi river south of latitude 35°, extending west to southwestern Arkansas and the valley of the Trinity river, Texas.

A tree 15 to 22 meters in height, with a trunk sometimes 1.20 meter in diameter, or toward its northern limits reduced to a low shrub; swamps or low wet woods, reaching its greatest development on the rich hummocks of the interior of the Florida peninsula and along the low sandy banks of pine-barren streams of the Gulf states.

Wood light, soft, not strong, close-grained, compact; medullary rays very numerous, thin; color, light brown tinged with red, the sap-wood nearly white; specific gravity, 0.5035; ash, 0.47; in the Gulf states sometimes used in the manufacture of broom handles and small woodenware.

The dried bark, especially of the root, of this species and of *M. acuminata* and *M. Umbrella* is included in the American *Materia Medica*, furnishing an aromatic tonic and stimulant used in intermittent and remittent fevers; a tincture made by macerating the fresh fruit or bark in brandy is a popular remedy for rheumatism (*U. S. Dispensatory*, 14 ed. 567.—*Nat. Dispensatory*, 2 ed. 891).

3.—*Magnolia acuminata*, Linnæus,

Spec. 2. ed. 756.—Marshall, Arbustum, 83.—Walter, Fl. Caroliniana, 159.—B. S. Barton, Coll. i, 13.—Aiton, Hort. Kew. ii, 251; 2 ed. iii, 331.—Lamarck, Dict. iii, 674.—Willdenow, Spec. ii, 1257; Enum. i, 579.—Michaux, Fl. Bor.-Am. i, 329.—Nouveau Duhamel, ii, 222.—Desfontaines, Hist. Arb. ii, 5.—Michaux f. Hist. Arb. Am. iii, 82, t. 3; N. American Sylva, 3 ed. ii, 15, t. 53.—Pursh, Fl. Am. Sept. ii, 381.—De Candolle, Syst. i, 453; Prodr. i, 80.—Loddiges, Bot. Cab. t. 418.—Nuttall, Genera, ii, 18.—Bot. Mag. t. 2427.—Hayne, Dend. Fl. 117.—Elliott, Sk. ii, 37.—Rafinesque, Med. Bot. ii, 32.—Guimpel, Otto & Hayne, Abb. Holz. 18, t. 17.—Sprengel, Syst. ii, 642.—Torrey, Compend. Fl. N. States, 221; Fl. N. York, i, 28.—Rafinesque, Med. Bot. ii, 34.—Beck, Bot. 15.—Sertum Botanicum, v. & t.—Don, Miller's Dict. i, 83.—Reichenbach, Fl. Exot. t. 251.—Eaton, Manual, 6 ed. 218.—London, Arboretum, i, 273 & t.—Eaton & Wright, Bot. 312.—Torrey & Gray, Fl. N. America, i, 43.—Dietrich, Syn. iii, 308.—Griffith, Med. Bot. 98.—Darlington, Fl. Cestrica, 3. ed. 9.—Darby, Bot. S. States, 211.—Cooper in Smithsonian Rep. 1858, 250.—Chapman, Fl. S. States, 14.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 67.—Wood, Cl. Book, 214; Bot. & Fl. 24.—Porcher, Resources S. Forests, 38.—Baillon, Hist. Pl. i, 140.—Gray, Manual N. States, 5. ed. 49.—Koch, Dendrologie, i, 371.—Young, Bot. Texas, 149.—Vasey, Cat. Forest Trees, 6.—Nat. Dispensatory, 2 ed. 891.—Ridgway in Proc. U. S. Nat. Mus. 1882, 58.

M. Virginiana, var. *e.* Linnæus, Spec. 1 ed. 536.

M. DeCandollii, Savi, Bibl. Ital. i, 224 & t.

Tulipastrum Americanum, Spach, Hist. Veg. vii, 483.

CUCUMBER TREE. MOUNTAIN MAGNOLIA.

Western New York to southern Illinois, southward along the Alleghany mountains, and scattered through eastern and middle Kentucky and Tennessee, usually on Carboniferous deposits, to southern Alabama (Stockton, *Mohr*) and northeastern Mississippi; Arkansas, Crowley's ridge, and in the southern and southwestern part of the state (Texarkana, *Harvey*, and in Polk, Howard, Cross, and Pike counties).

A large tree, 20 to 30 meters in height, with a trunk 0.60 to 1.20 meter in diameter; rich woods, reaching its greatest development on the slopes of the southern Alleghany mountains.

Wood durable, light, soft, not strong, close-grained, compact, satiny; medullary rays numerous, thin; color, yellow-brown, the sap-wood lighter, often nearly white; specific gravity, 0.4690; ash, 0.29; used for pump-logs, water-troughs, flooring, cabinet-making, etc.

4.—*Magnolia cordata*, Michaux,

Fl. Bor.-Am. i, 328.—Aiton, Hort. Kew. 2 ed. iii, 331.—Poiret, Suppl. iii, 547.—Michaux f. Hist. Arb. Am. iii, 87, t. 4; N. American Sylva, 3 ed. ii, 18, t. 54.—Pursh, Fl. Am. Sept. ii, 382.—Lindley, Bot. Reg. iv, t. 325.—Nuttall, Genera, ii, 18.—De Candolle, Syst. i, 455; Prodr. i, 80.—Hayne, Dend. Fl. 118.—Elliott, Sk. ii, 38.—Loddiges, Bot. Cab. t. 474.—Sprengel, Syst. ii, 642.—Rafinesque, Med. Bot. ii, 32.—Eaton, Manual, 6 ed. 218.—Sertum Botanicum, v & t.—Don, Miller's Dict. i, 83.—Reichenbach, Fl. Exot. t. 250.—London, Arboretum, i, 275 & t.—Eaton & Wright, Bot. 312.—Torrey & Gray, Fl. N. America, i, 43.—Dietrich, Syn. iii, 308.—Darby, Bot. S. States, 211.—Cooper in Smithsonian Rep. 1858, 250.—Chapman, Fl. S. States, 14.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 68.—Wood, Cl. Book, 214; Bot. & Fl. 25.—Koch, Dendrologie, i, 371.—Vasey, Cat. Forest Trees, 6.

Tulipastrum Americanum, var. *subcordatum*, Spach, Hist. Veg. vii, 483.

CUCUMBER TREE.

Southern Alleghany Mountain region, near Augusta, Georgia (*Michaux, Elliott*), head of Sipsey creek, "valley of Davidson creek", Winston county, Alabama (*Mohr*).

A tree 22 to 24 meters in height, with a trunk sometimes 0.60 meter in diameter; low, rich woods; very rare and local.

Wood light, soft, not strong, close-grained, compact; medullary rays very numerous, thin; color, light brown streaked with yellow, the sap-wood light yellow; specific gravity, 0.4139; ash, 0.32.

5.—*Magnolia macrophylla*, Michaux,

Fl. Bor.-Am. i, 327.—Nouveau Duhamel, ii, 221.—Desfontaines, Hist. Arb. ii, 5.—Aiton, Hort. Kew. 2 ed. iii, 331.—Poirot, Suppl. iii, 573.—Michaux f. Hist. Arb. Am. iii, 99, t. 7; N. American Sylva, ii, 26, t. 57.—Bonpland, Pl. Malm. 84, t. 33.—Pursh, Fl. Am. Sept. ii, 381.—Nuttall, Genera, ii, 18; Sylva, i, 83; 2 ed. i, 99.—De Candolle, Syst. i, 454; Prodr. i, 80.—Bot. Mag. t. 2189.—Hayne, Dend. Fl. 117.—Elliott, Sk. ii, 40.—Sprengel, Syst. ii, 642.—Rafinesque, Med. Bot. ii, 31, t. 62.—Eaton, Manual, 6 ed. 218.—Sertum Botanicum, v & t.—Don, Miller's Dict. i, 83.—Croom in Am. Jour. Sci. 1 ser. xxv, 76.—Reichenbach, Fl. Exot. ii, 44, t. 139.—Loudon, Arboretum, i, 271 & t.—Eaton & Wright, Bot. 312.—Torrey & Gray, Fl. N. America, i, 43.—Spach, Hist. Veg. vii, 479.—Dietrich, Syn. iii, 308.—Griffith, Med. Bot. 98, f. 57.—Darby, Bot. S. States, 211.—Cooper in Smithsonian Rep. 1858, 250.—Seringe, Fl. Jard. iii, 230.—Chapman, Fl. S. States, 14.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 67.—Wood, Cl. Book, 214; Bot. & Fl. 25.—Gray, Manual N. States, 5 ed. 49.—Koch, Dendrologie, i, 374.—Vasey, Cat. Forest Trees, 6.

LARGE-LEAVED CUCUMBER TREE.

North Carolina, eastern base of the Alleghany mountains (Iredell and Lincoln counties); southeastern Kentucky southward to middle and western Florida and southern Alabama, extending west to the valley of Pearl river, Louisiana; central Arkansas (Garland, Montgomery, Hot Springs, and Sebastian counties).

A tree 6 to 18 meters in height, with a trunk rarely 0.60 meter in diameter; rich woods, reaching its greatest development in the limestone valleys of northern Alabama; rare and local.

Wood light, hard, not strong, close-grained, compact, satiny; medullary rays numerous, thin; color, brown, the sap-wood light yellow; specific gravity, 0.5309; ash, 0.35.

6.—*Magnolia Umbrella*, Lamarck,

Dict. iii, 673.—Nouveau Duhamel, ii, 221.—De Candolle, Prodr. i, 80.—Loiseleur, Herb. Amat. iii, t. 198.—Sprengel, Syst. ii, 642.—Don, Miller's Dict. i, 83.—Torrey & Gray, Fl. N. America, i, 43.—Spach, Hist. Veg. vii, 475.—Dietrich, Syn. iii, 308.—Seringe, Fl. Jard. iii, 227.—Gray, Genera, i, 62, t. 24; Proc. Linnæan Soc. ii, 106, f. 1-18; Manual N. States, 5 ed. 49.—Cooper in Smithsonian Rep. 1858, 250.—Chapman, Fl. S. States, 13.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 67.—Wood, Cl. Book, 214; Bot. & Fl. 25.—Porcher, Resources S. Forests, 38.—Vasey, Cat. Forest Trees, 6.

M. Virginiana, var. *tripetala*, Linnæus, Spec. 1 ed. 536.

M. tripetala, Linnæus, Spec. 2 ed. 756.—Marshall, Arbustum, 84.—Walter, Fl. Caroliniana, 159.—B. S. Barton, Coll. i, 14.—Aiton, Hort. Kew. ii, 252; 2 ed. iii, 331.—Willdenow, Spec. ii, 1258; Enum. i, 579.—Michaux, Fl. Bor.-Am. i, 327.—Desfontaines, Hist. Arb. ii, 5.—De Candolle, Syst. i, 452.—Michaux f. Hist. Arb. Am. iii, 90, t. 5; N. American Sylva, 3 ed. ii, 20, t. 5.—Pursh, Fl. Am. Sept. ii, 381.—Nuttall, Genera, ii, 18; Sylva, i, 84; 2 ed. i, 100.—Guimpel, Otto & Hayne, Abb. Holz. 20, t. 18.—Hayne, Dend. Fl. 116.—Elliott, Sk. ii, 38.—Torrey, Compend. Fl. N. States, 221.—Rafinesque, Med. Bot. ii, 32.—Eaton, Manual, 6 ed. 218.—Eaton & Wright, Bot. 312.—Griffith, Med. Bot. 98.—Loudon, Arboretum, i, 269, t. 5.—Darby, Bot. S. States, 211.—Koch, Dendrologie, i, 370.—Nat. Dispensatory, 2 ed. 891.

UMBRELLA TREE. ELK WOOD.

Southeastern Pennsylvania, southward along the Alleghany mountains to central Alabama (Prattville, *Mohr*) and northeastern Mississippi, westward through Kentucky and Tennessee; in central (Hot Springs) and southwestern Arkansas (Fulton, valley of the Red river, *Harvey*).

A small tree, rarely exceeding 12 meters in height, with a trunk 0.10 to 0.40 meter in diameter; rich, shady hillsides; most common and reaching its greatest development along the western slope of the southern Alleghany mountains.

Wood light, soft, not strong, close-grained, compact; medullary rays very numerous, thin; color, brown, the heavier sap-wood nearly white; specific gravity, 0.4487; ash, 0.20.

FOREST TREES OF NORTH AMERICA.

7.—*Magnolia Fraseri*, Walter,

Fl. Caroliniana, i, 59 & t.—Torrey & Gray, *Fl. N. America*, i, 43.—Walpers, *Rep.* i, 70.—Dietrich, *Syn.* iii, 308.—Chapman, *Fl. S. States*, 14.—Curtis in *Rep. Geological Surv. N. Carolina*, 1860, iii, 68.—Wood, *Cl. Book*, 214; *Bot. & Fl.* 25.—Gray, *Manual N. States*, 5 ed. 49.—Koch, *Dendrologie*, i, 372.—Vasey, *Cat. Forest Trees*, 6.

M. auriculata, Lamarck, *Dict.* iij, 673.—Bartram, *Travels*, 2 ed. 337.—Willdenow, *Spec.* ii, 1258; *Enum.* i, 579.—Michaux, *Fl. Bor.-Am.* i, 328.—Nouveau Duhamel, ii, 222.—Desfontaines, *Hist. Arb.* ii, 5.—Michaux f. *Hist. Arb. Am.* iii, 94, t. 6; *N. American Sylva*, 3 ed. ii, 23, t. 56.—Andrews, *Bot. Rep.* ix, t. 573.—*Bot. Mag.* t. 1206.—Cubières, *Mem. Mag. & t.*—Aiton, *Hort. Kew.* 2 ed. iii, 332.—Pursh, *Fl. Am. Sept.* ii, 382.—Nuttall, *Genera*, ii, 18; *Sylva*, i, 84; 2 ed. i, 98.—De Candolle, *Syst.* i, 454; *Prodr.* i, 80.—Hayne, *Dend. Fl.* 117.—Elliott, *Sk.* ii, 39.—Sprengel, *Syst.* ii, 642.—Audubon, *Birds*, t. 38.—Don, *Miller's Dict.* i, 83.—Eaton, *Manual*, 6 ed. 218.—Hooker, *Jour. Bot.* i, 188.—Spach, *Hist. Veg.* vii, 477.—London, *Arboretum*, i, 276 & t.—Seringe, *Fl. Jard.* iii, 229.

M. pyramidata, Bartram, *Travels*, 2 ed. 338.—Pursh, *Fl. Am. Sept.* ii, 382.—De Candolle, *Syst.* i, 454; *Prodr.* i, 80.—Hayne, *Dend. Fl.* 117.—Lindley, *Bot. Reg.* v, t. 407.—Loddiges, *Bot. Cab.* t. 1092.—Rafinesque, *Med. Bot.* ii, 32.—Don, *Miller's Dict.* i, 83.—Eaton, *Manual*, 6 ed. 221.—London, *Arboretum*, i, 277 & t.—Seringe, *Fl. Jard.* iii, 230.—Darby, *Bot. S. States*, 211.

M. auricularis, Salisbury, *Parad. Lond.* i, t. 43.—Kerner, *Hort.* t. 360.

LONG-LEAVED CUCUMBER TREE.

Alleghany mountains, from Virginia southward to the Chattahoochee region of western Florida, and southern Alabama (Clark county, *Mohr*), extending west to the valley of Pearl river, Mississippi.

A small tree, 8 to 12 meters in height, with a trunk 0.15 to 0.20 meter in diameter; rich woods.

Wood light, soft, not strong, close-grained, compact; medullary rays very numerous, thin; color, brown, the sap-wood nearly white; specific gravity, 0.5003; ash, 0.28.

8.—*Liriodendron Tulipifera*, Linnaeus,

Spec. 1 ed. i, 535.—Kalm, *Travels*, English ed. i, 202.—Marshall, *Arbustum*, 78.—Wangenheim, *Amer.* 32, t. 13, f. 32.—Walter, *Fl. Caroliniana*, 158.—Schmidt, *Arb.* i, 48.—B. S. Barton, *Coll.* i, 14, 45.—Aiton, *Hort. Kew.* ii, 250; 2 ed. iii, 329.—Gärtner, *Fruct.* ii, t. 178.—*Bot. Mag.* t. 275.—Möench, *Meth.* 222.—Abbot, *Insects Georgia*, ii, t. 102.—Schkuhr, *Handb.* ii, 93, t. 147.—Trew, *Icon.* t. 10.—Willdenow, *Spec.* ii, 1254; *Enum.* i, 579.—Michaux, *Fl. Bor.-Am.* i, 326.—Nouveau Duhamel, iii, 62, t. 18.—Desfontaines, *Hist. Arb.* ii, 15.—Poirot in Lamarck, *Dict.* viii, 137; *Ill.* iii, 36, t. 491.—St. Hilaire, *Pl. France*, iii, t. 377.—Titford, *Hort. Bot. Am.* 76.—Michaux f. *Hist. Arb. Am.* iii, 202, t. 5; *N. American Sylva*, 3 ed. ii, 35, t. 61.—Eaton, *Manual*, 63; 6 ed. 208.—Nuttall, *Genera*, ii, 18; *Sylva*, i, 84; 2 ed. i, 100.—Barton, *Prodr. Fl. Philadelph.* 59; *Med. Bot.* i, 91, t. 8; *Compend. Fl. Philadelph.* ii, 18.—De Candolle, *Syst.* i, 462; *Prodr.* i, 82.—Bigelow, *Med. Bot.* ii, 107, t. 31.—Hayne, *Dend. Fl.* 115.—Elliott, *Sk.* ii, 40.—Torrey, *Compend. Fl. N. States*, 221; *Fl. N. York*, i, 28.—Rafinesque, *Med. Bot.* ii, 239.—Guimpel, Otto & Hayne, *Abb. Holz.* 34, t. 29.—Cobbett, *Woodlands*, No. 516.—Sprengel, *Syst.* ii, 642.—Audubon, *Birds*, t. 12.—Don, *Miller's Dict.* i, 86.—Beck, *Bot.* 15.—Lindley, *Fl. Med.* 23.—Spach, *Hist. Veg.* vi, 488.—London, *Arboretum*, i, 284 & t.—Eaton & Wright, *Bot.* 302.—Penn. *Cycl.* xxv, 341.—Torrey & Gray, *Fl. N. America*, i, 44.—Dietrich, *Syn.* iii, 309.—Griffith, *Med. Bot.* 98, f. 58.—Emerson, *Trees Massachusetts*, 529; 2 ed. ii, 605 & t.—Seringe, *Fl. Jard.* iii, 240.—Gray, *Genera*, i, 64, t. 25; *Manual N. States*, 5 ed. 50.—Darlington, *Fl. Cestrica*, 3 ed. 9.—Darby, *Bot. S. States*, 212.—Agardh, *Theor. & Syst. Pl.* t. 11, f. 2.—Cooper in *Smithsonian Rep.* 1858, 250.—Chapman, *Fl. S. States*, 14.—Curtis in *Rep. Geological Surv. N. Carolina*, 1860, iii, 77.—Lemaire, *Ill. Hort.* 15, t. 571.—Wood, *Cl. Book*, 215; *Bot. & Fl.* 25.—Porcher, *Resources S. Forests*, 39.—Engelmann in *Trans. Am. Phil. Soc. new ser.* xii, 183.—Baillon, *Hist. Pl.* i, 143, f. 175-178.—Koch, *Dendrologie*, i, 380.—Guibourt, *Hist. Drogues*, 7 ed. iii, 748.—Ridgway in *Am. Nat.* vi, 663; *Proc. U. S. Nat. Mus.* 1892, 59.—Vasey, *Cat. Forest Trees*, 6.—Eichler, *Sit. Bot. Brand.* xxii, 83, f. 1-3.—Bell in *Geological Rep. Canada*, 1879-'80, 53°.

Tulipifera Liriodendron, Miller, *Dict.* No. 1.

L. procera, Salisbury, *Prodr.* 379.

TULIP TREE. YELLOW POPLAR. WHITE WOOD.

Southwestern Vermont, through western New England, southward to northern Florida (latitude 30°); west through New York, Ontario, and Michigan to lake Michigan, south of latitude 43° 30', thence south to latitude 31° in the Gulf states east of the Mississippi river; through southern Illinois and southeastern Missouri to Crowley's ridge, northeastern Arkansas.

One of the largest and most valuable trees of the Atlantic forests, 30 to 60 meters in height, with a trunk 2 to 4 meters in diameter (*Ridgway*); rich woods and intervale lands, reaching its greatest development in the valley of the lower Wabash river and along the western slopes of the Alleghany mountains in Tennessee and North Carolina.

Wood light, soft, not strong, brittle, very close straight-grained, compact, easily worked; medullary rays numerous, not prominent; color, light yellow or brown, the thin sap-wood nearly white; specific gravity, 0.4230; ash, 0.23; largely manufactured into lumber and used for construction, interior finish, shingles, in boat-building, and especially in the manufacture of wooden pumps, woodenware, etc.; varieties varying slightly in color and density are recognized by lumbermen.

Liriodendrin, a stimulant tonic, with diaphoretic properties, is obtained by macerating the inner bark, especially of the root (*Jour. Philadelphia Col. Phar.* iii, 5.—*U. S. Dispensatory*, 14 ed. 556.—*Nat. Dispensatory*, 2 ed. 871).

ANONACEÆ.

9.—*Asimina triloba*, Dunal,

Mon. Anon. 83.—De Candolle, Syst. i, 479; Prodr. i, 87.—Elliott, Sk. ii, 42.—Guimpel, Otto & Hayne, Abb. Holz. 66, t. 53.—Hayne, Dend. Fl. 118.—Sprengel, Syst. ii, 639.—Torrey, Compend. Fl. N. States, 222; Ann. Lye. N. York, ii, 165.—Beck, Bot. 16.—Don, Miller's Dict. i, 91.—Nuttall in Jour. Philadelphia Acad. vii, 11.—Dietrich, Syn. iii, 304.—London, Arboretum, i, 293, f. 39.—Gray, Genera, i, 69, t. 26, 27; Manual N. States, 5 ed. 50.—Parry in Owen's Rep. 609.—Darlington, Fl. Cestrica, 3 ed. 9.—Darby, Bot. S. States, 212.—Cooper in Smithsonian Rep. 1858, 250.—Chapman, Fl. S. States, 15.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 94.—Lesquereux in Owen's 2d Rep. Arkansas, 247.—Maout & Decaisne, Bot. English ed. 199 & figs.—Bot. Mag. t. 5854.—Wood, Cl. Book, 215; Bot. & Fl. 26.—Porcher, Resources S. Forests, 41.—Engelmann in Trans. Am. Phil. Soc. new ser. xii, 183.—Koch, Dendrologie, ii, 383.—Young, Bot. Texas, 149.—Vasey, Cat. Forest Trees, 6.—Ridgway in Proc. U. S. Nat. Mus. 1882, 60.—Burgess in Coulter's Bot. Gazette, vii, 95.

Anona triloba, Linnaeus, Spec. 1 ed. 537.—Marshall, Arbustum, 10.—Lamarck, Dict. ii, 125.—Walter, Fl. Caroliniana, 158.—B. S. Barton, Coll. i, 29.—Aiton, Hort. Kew. ii, 254; 2 ed. iii, 335.—Willdenow, Spec. ii, 1267; Enum. i, 580.—Nouveau Duhamel, ii, 83, t. 25.—Desfontaines, Hist. Arb. ii, 21.—Michaux f. Hist. Arb. Am. iii, 161, t. 9; N. American Sylva, 3 ed. ii, 33, t. 60.—Barton, Prodr. Fl. Philadelph. 59.—Schkuhr, Handb. ii, 95, t. 149.

Anona pendula, Salisbury, Prodr. 380.

Orchidocarpum arietinum, Michaux, Fl. Bor.-Am. i, 329.

Porcelia triloba, Persoon, Syn. ii, 95.—Pursh, Fl. Am. Sept. ii, 383.—Rafinesque, Fl. Ludoviciana, 92.—Barton, Compend. Fl. Philadelph. ii, 18.—Nuttall, Genera, ii, 19.—Poirot, Suppl. iv, 529.—Eaton, Manual, 6 ed. 278.—Audubon, Birds, t. 2, 162.—Eaton & Wright, Bot. 371.

Uvaria triloba, Torrey & Gray, Fl. N. America, i, 45.—Torrey, Fl. N. York, i, 30.—Caruel in Ann. Mus. Firenze, 1864, 9, t. 1, f. 1-7.—Baillon, Adansonia, viii, 333; Hist. Pl. i, 193, f. 220-228.

A. campaniflora, Spach, Hist. Veg. vii, 529.

PAPAW. CUSTARD APPLE.

Western New York (Lockport and in Monroe county); Ontario (Queenstown heights); eastern and central Pennsylvania, west to southern Michigan, southern Iowa, and eastern Kansas (Manhattan), south to middle Florida and the valley of the Sabine river, Texas.

A small tree, sometimes 12 meters in height, with a trunk rarely exceeding 0.30 meter in diameter, or often reduced to a slender shrub; rich, rather low woods, reaching its greatest development in the lower Wabash valley and in the valley of the White river, Arkansas.

Wood very light, very soft and weak, coarse-grained, spongy; layers of annual growth clearly marked by several rows of large open ducts; color, light yellow shaded with green, the sap-wood lighter; specific gravity, 0.3969; ash, 0.21.

10.—*Anona laurifolia*, Dunal,

Mon. Anon. 65.—De Candolle, Syst. i, 468; Prodr. i, 84.—Sprengel, Syst. ii, 641.—Lindley, Bot. Reg. xvi, t. 1328.—Schnitzlein, Icon. t. 174, f. 9.—Grisebach, Fl. British West Indies, 4.—Cooper in Smithsonian Rep. 1860, 439.—Chapman, Fl. S. States, Suppl. 603.

A. glabra, Chapman in Coulter's Bot. Gazette, iii, 2 [not Linnaeus].

A. species, Vasey, Cat. Forest Trees, 6.

POND APPLE.

Semi-tropical Florida, cape Malabar to bay Biscayne, on the west coast, Pease creek to the Caloosa river, and through the West Indies.

A small tree, sometimes 9 meters in height, with a trunk 0.30 meter in diameter, or toward its northern limit and on the west coast often reduced to a stout, wide-spreading shrub; common and reaching its greatest development within the United States on the low islands and shores of the Everglades in the neighborhood of bay Biscayne.

Wood light, soft, not strong, rather close-grained, compact, containing many scattered open ducts; color, light brown streaked with yellow, sap-wood lighter; specific gravity, 0.5053; ash, 4.86.

The large fruit (0.14 to 0.28 meter long) scarcely edible.

CAPPARIDACEÆ.

11.—Capparis Jamaicensis, Jacquin,

Stirp. Am. 160, t. 101.—Aiton, Hort. Kew. 2 ed. iii, 285.—De Candolle, Prodr. i, 252.—Descourtilz, Fl. Med. Antilles, v. t. 273.—Macfadyen, Fl. Jamaica, 39.—Grisebach, Fl. British West Indies, 18.—Chapman, Fl. S. States, 32.—Porcher, Resources S. Forests, 75.—Eichler in Martius, Fl. Brasil. xiii, 270, t. 64, f. 11.—Vasey, Cat. Forest Trees, 6.

C. Breynia, Linnæus, Spec. 2 ed. 721, in part.—Aiton, Hort. Kew. 2 ed. iii, 285.—De Candolle, Prodr. i, 252, in part.—Swartz, Obs. 210 [not Jacquin].—Macfadyen, Fl. Jamaica, 39.

C. cynophyllophora, Linnæus, Spec. 1 ed. 504 [not subsequent ed. *vide* Eichler, l. c.].—Aiton, Hort. Kew. 2 ed. iii, 285.—Macfadyen, Fl. Jamaica, 39.

C. siliquosa, Linnæus, Spec. 2 ed. 721.

C. torulosa, Swartz, Prodr. 81.—De Candolle, Prodr. i, 252.—Grisebach, Fl. British West Indies, 18.

C. uncinata, Loddiges, Cat. [not Wallich].

C. emarginata, Richard, Fl. Cuba, 78, t. 9.—Walpers, Rep. i, 201.

Semi-tropical Florida, cape Canaveral to the southern keys; in the West Indies and southward to Brazil.

A small tree, sometimes 6 meters in height, with a trunk 0.15 meter in diameter, or reduced to a low shrub; common and reaching its greatest development within the United States on Upper Metacombe and Umbrella Keys.

Wood heavy, hard, close-grained, compact, satiny, containing many evenly-distributed large open ducts; medullary rays numerous, obscure; color, yellow tinged with red, the sap-wood lighter; specific gravity, 0.6971; ash, 4.76.

CANELLACEÆ.

12.—Canella alba, Murray;

Linnæus, Syst. 14 ed. iv, 443.—Swartz, Obs. 190; Trans. Linnæan Soc. i, 96, t. 8.—Willdenow, Spec. ii, 851; Enum. i, 496.—Aiton, Hort. Kew. 2 ed. iii, 144.—Titford, Hort. Bot. Am. Suppl. 3, t. 10, f. 4.—De Candolle, Prodr. i, 563.—Hayne, Arzn. 9, t. 5.—Stevenson & Churchill, Med. Bot. ii, t. 66.—Woodville, Med. Bot. 3 ed. iv, 694, t. 237.—Lindley, Med. Bot. 116.—Carson, Med. Bot. i, 24, t. 16.—Griffith, Med. Bot. 181, f. 98.—Miers in Ann. Nat. Hist. 3 ser. i, 348; Contrib. i, 116.—Grisebach, Fl. British West Indies, 109.—Chapman, Fl. S. States, 93.—Guibourt, Hist. Drogues, 7 ed. iii, 621, f. 767.—Vasey, Cat. Forest Trees, 7.—Bentley & Trimen, Med. Pl. i, 26, t. 26.

C. Winterana, Gärtner, Fruct. i, 377, t. 77.

Wintera Canella, Linnæus, Spec. 2 ed. 636.—Poiret in Lamarck, Dict. viii, 799, t. 399.

C. laurifolia, Loddiges, Cat.—Sweet, Hort. Brit. 65.—Don, Miller's Diet. i, 680.

WHITE WOOD. CINNAMON BARK. WILD CINNAMON.

Semi-tropical Florida, on the southern keys (Elliott's Key, Key Largo to Jew Fish Key); through the West Indies.

A small tree, often 10 meters in height, with a trunk 0.22 meter in diameter; not rare.

Wood very heavy, exceedingly hard, strong, close-grained, compact; medullary rays numerous, thin; color, dark reddish-brown, the sap-wood light brown or yellow; specific gravity, 0.9893; ash, 1.75.

The pale inner bark appears in the *Pharmacopœa* under the name of *Cortex canellæ albæ*, furnishing an aromatic stimulant and tonic, occasionally employed in cases of debility of the digestive organs, or as an adjunct to more active remedies (*Miers, l. c.—Flückiger & Hanbury, Pharmacographia, 68.—U. S. Dispensatory, 14 ed. 210.—Nat. Dispensatory, 2 ed. 337*).

GUTTIFERÆ.

13.—*Clusia flava*, Linnæus,

Spec. 2 ed. 1495.—Willdenow, Spec. iv, 977; Enum. ii, 1043.—Aiton, Hort. Kew. 2 ed. v, 444.—Titford, Hort. Bot. Am. 105.—De Candolle, Prodr. i, 559.—Macfadyen, Fl. Jamaica, 134.—Nuttall, Sylva, ii, 111, t. 77; 2 ed. ii, 58, t. 77.—Grisebach, Fl. British West Indies, 407.—Cooper in Smithsonian Rep. 1858, 264.—Chapman, Fl. S. States, 43.—Planchon & Triana in Ann. Sci. Nat. 4 ser. xiii, 352.—Walpers, Ann. vii, 340.—Vasey, Cat. Forest Trees, 7.

C. rosea, Torrey & Gray, Fl. N. America, i, 168.

Jamaica and other West Indian islands; Key West (*Blodgett*) prior to 1840. Not detected by later explorers (*Palmer, Garber, Chapman, Ourtiss*) of the botany of semi-tropical Florida, and probably not now growing spontaneously within the limits of the United States.

Wood not examined.

TERNSTROMIACEÆ.

14.—*Gordonia Lasianthus*, Linnæus,

Mant. i, 570.—Ellis, Phil. Trans. 60, 518, t. 11; Letters, t. 2.—L'Heritier, Stirp. Nov. 156.—Cavanilles, Diss. ii, 307, t. 161.—Walter, Fl. Caroliniana, 177.—Aiton, Hort. Kew. ii, 231; 2 ed. iv, 234.—Lamarck, Dict. ii, 770; Ill. iii, 146, t. 594, f. 1.—Swartz, Obs. 271.—Willdenow, Spec. iii, 840.—Michaux, Fl. Bor.-Am. ii, 43.—Bot. Mag. t. 668.—Nouveau Duhamel, ii, 236, t. 68.—Desfontaines, Hist. Arb. i, 484.—Persoon, Syn. ii, 259.—Michaux f. Hist. Arb. Am. iii, 131, t. 1; N. American Sylva, 3 ed. ii, 29, t. 58.—Pursh, Fl. Am. Sept. i, 451.—Nuttall, Genera, ii, 84.—De Candolle, Prodr. i, 528.—Elliott, Sk. ii, 171.—Sprengel, Syst. iii, 125.—Don, Miller's Dict. i, 573, f. 99.—Audubon, Birds, t. 168.—Reichenbach, Fl. Exot. t. 151.—Spach, Hist. Veg. iv, 79.—London, Arboretum, i, 379, f. 93.—Torrey & Gray, Fl. N. America, i, 223.—Eaton, Manual, 6 ed. 161.—Eaton & Wright, Bot. 258.—Browne, Trees of America, 52.—Dietrich, Syn. iv, 862.—Gray, Genera, ii, 103, t. 140, 141; Manual N. States, 5 ed. 104.—Choisy, Mem. Ternst. & Camel. 51.—Darby, Bot. S. States, 256.—Cooper in Smithsonian Rep. 1858, 250.—Chapman, Fl. S. States, 60.—Curtis in Rep. Geological Surv. N. Carolina 1860, iii, 80.—Maout & Decaisne, English ed. 274 & figs.—Wood, Cl. Book, 274; Bot. & Fl. 65.—Baillon, Hist. Pl. iv, 230, f. 254, 255.—Vasey, Cat. Forest Trees, 7.

Hypericum Lasianthus, Linnæus, Spec. 1 ed. 783.—Hill, Veg. Syst. xv, t. 1, f. 3.

G. pyramidalis, Salisbury, Prodr. Stirp. 386.

LOBLOLLY BAY. TAN BAY.

Southern Virginia, south near the coast to cape Malabar, and cape Romano, Florida, west along the Gulf coast to the valley of the Mississippi river.

A tree 15 to 24 meters in height, with a trunk often 0.45 to 0.50 meter in diameter; low, sandy swamps.

Wood light, soft, not strong, close-grained, compact, not durable; medullary rays numerous, thin; color, light red, the sap-wood lighter; specific gravity, 0.4728; ash, 0.76; somewhat employed in cabinet-making.

The bark, rich in tannin, was once occasionally used, locally, in tanning leather (*Bartram, Travels*, 2 ed. 160).

15.—*Gordonia pubescens*, L'Heritier,

Stirp. Nov. 156.—Lamarck, Dict. ii, 770.—Cavanilles, Diss. ii, 308, t. 162.—Aiton, Hort. Kew. ii, 231; 2 ed. iv, 234.—Willdenow, Spec. iii, 841.—Michaux, Fl. Bor.-Am. ii, 43.—Venant, Jard. Malm. t. 1 (Schrader, Nenes Jour. Bot. 1806, 121).—Nouveau Duhamel, ii, 237.—Koenig & Sims, Ann. Bot. i, 171.—Desfontaines, Hist. Arb. i, 484.—Persoon, Syn. ii, 259.—Michaux f. Hist. Arb. Am. iii, 135, t. 2; N. American Sylva, 3 ed. ii, 31, t. 59.—Pursh, Fl. Am. Sept. ii, 451.—Nuttall, Genera, ii, 84.—Loiseleur, Herb. Amat. iv, t. 236.—Elliott, Sk. ii, 171.—De Candolle, Prodr. i, 528.—Sprengel, Syst. iii, 125.—Don, Miller's Dict. i, 573.—Eaton, Manual, 6 ed. 161.—Audubon, Birds, t. 185.—Spach, Hist. Veg. iv, 80.—London, Arboretum, i, 380, f. 94.—Torrey & Gray, Fl. N. America, i, 223.—Eaton & Wright, Bot. 258.—Browne, Trees of America, 54.—Dietrich, Syn. iv, 862.—Gray, Genera, ii, 102, t. 141, f. 11–14, t. 142.—Choisy, Mem. Ternst. & Camel. 51.—Darby, Bot. S. States, 257.—Cooper in Smithsonian Rep. 1858, 250.—Chapman, Fl. S. States, 60.—Wood, Cl. Book, 274; Bot. & Fl. 65.—Vasey, Cat. Forest Trees, 7.—Goodale & Sprague, Wild Flowers, 193, t. 47.

Franklinia Altamaha, Marshall, Arbustum, 49.—Bartram, Travels, 2 ed. 16, 465.—Rafinesque, Atlant. Jour. 79 & f.

G. Franklini, L'Heritier, Stirp. Nov. 156.—Willdenow, Spec. iii, 841.—Nouveau Duhamel, ii, 237.—Desfontaines, Hist. Arb. i, 484.—Persoon, Syn. ii, 259.—Poiret, Suppl. ii, 816.

Michauxia sessilis, Salisbury, Prodr. Stirp. 386.

Lacathea florida, Salisbury, Parad. Lond. t. 56.—Colla, Hort. Ripul. Appx. i, 134.

FRANKLINIA.

Near Fort Barrington, on the Altamaha river, Georgia (*J. & W. Bartram, Dr. Moses Marshall*).

Careful explorations of Bartram's original locality by later botanists, especially by Mr. H. W. Ravenel, have failed to rediscover this species, which is, however, still preserved in cultivation through the original plants introduced by the Bartrams. "Florida" given as a locality by Torrey & Gray, *l. c.*, on the authority of *Herb. Schweinitz*, and followed by Chapman, *l. c.*, is probably an error (*Ravenel in Am. Naturalist*, xvi, 235).

STERCULIACEÆ.

16.—*Fremontia Californica*, Torrey,

Smithsonian Contrib. vi, 5, t. 2, f. 2; Proc. Am. Assoc. iv, 191; Pacific R. R. Rep. iv, 15, 71.—Newberry in Pacific R. R. Rep. vi, 68.—Walpers, Ann. iv, 319.—Gray in Jour. Boston Soc. Nat. Hist. vii, 146.—Bentham & Hooker, Genera, i, 212, 962.—Bot. Mag. t. 5691.—Lemaire, Ill. Hort. xiii, t. 496.—Belge Hort. xvii, 236, t. 13.—Carrière in Rev. Hort. 1867, 91 & t.—Koch, Dendrologie, i, 483.—Masters in London Gard. Chronicle, 1869, 610.—Seemann, Jour. Bot. vii, 297.—London Garden, 1873, 54 & t.—Planchon in Fl. des Serres, xxii, 175.—Brewer & Watson, Bot. California, i, 88; ii, 437.—Rothrock in Wheeler's Rep. vi, 41, 357.

Cheiranthodendron Californicum, Baillon, Hist. Pl. iv, 70.

SLIPPERY ELM.

California, valley of Pitt river, southward along the western foot-hills of the Sierra Nevada, and in the Santa Lucia mountains southward through the Coast ranges to the San Jacinto mountains; rare at the north, most common and reaching its greatest development on the southern sierras and the San Gabriel and San Bernardino ranges.

A small tree, 6 to 10 meters in height, the short trunk often 0.30 to 0.45 meter in diameter, or more often a tall, much branched shrub; dry, gravelly soil.

Wood heavy, hard, very close-grained, compact, satiny, containing many groups of small ducts parallel to the thin, conspicuous medullary rays, layers of annual growth obscure; color, dark brown tinged with red, the thick sap-wood lighter; specific gravity, 0.7142; ash, 1.69.

The mucilaginous inner bark used locally in poultices.

TILIACEÆ.

17.—*Tilia Americana*, Linnaeus,

Spec. 1 ed. 514.—Marshall, Arbustum, 153.—Wangenheim, Amer. 55.—Aiton, Hort. Kew. ii, 229; 2 ed. iii, 299.—Willdenow, Spec. ii, 1162; Enum. i, 565.—Desfontaines, Hist. Arb. ii, 37.—Persoon, Syn. ii, 66.—Michaux f. Hist. Arb. Am. iii, 311, t. 1; N. American Sylva, 3 ed. iii, 81, t. 131.—Barton, Prodr. Fl. Philadelph. 58; Compend. Fl. Philadelph. ii, 6.—Eaton, Manual, 59.—James in Long's Exped. i, 69.—Watson, Dend. Brit. ii, 134, t. 134.—Torrey, Compend. Fl. N. States, 214; Fl. N. York, i, 116.—London, Arboretum i, 373 & t.—Torrey & Gray, Fl. N. America, i, 239.—Bigelow, Fl. Boston. 3 ed. 227.—Emerson, Trees Massachusetts, 511; 2 ed. ii, 584 & t.—Browne, Trees of America, 47.—Gray, Genera, ii, 96, t. 136; Manual N. States, 5 ed. 103; Hall's Pl. Texas, 5.—Darlington, Fl. Cestrica, 3 ed. 38.—Darby, Bot. S. States, 262.—Cooper in Smithsonian Rep. 1868, 250.—Chapman, Fl. S. States, 59.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 79.—Lesquereux in Owen's 2d Rep. Arkansas, 352.—Wood, Cl. Book, 272; Bot. & Fl. 64.—Porcher, Resources S. Forests, 103.—Engelmann in Trans. Am. Phil. Soc. new ser. xii, 186.—Walpers, Ann. vii, 449.—Koch, Dendrologie, i, 480.—Young, Bot. Texas, 188.—Vasey, Cat. Forest Trees, 7.—Macoun in Geological Rep. Canada, 1875-76, 191.—Sears in Bull. Essex Inst. xiii, 174.—Bell in Geological Rep. Canada, 1879-80, 51c.—Ridgway in Proc. U. S. Nat. Mus. 1882, 61.

T. nigra, Borkhausen, Handb. d. Forstbot. ii, 1219.

T. glabra, Ventenat in Mem. Acad. Sci. iv, 9, t. 2.—Nouveau Duhamel, i, 228.—Poiret in Lamarck, Dict. vii, 681.—Pursh, Fl. Am. Sept. ii, 362.—Nuttall, Genera, ii, 3.—De Candolle, Prodr. i, 513.—Hayne, Dend. Fl. 112.—Elliott, Sk. ii, 2.—Guimpel, Otto & Hayne, Abb. Holz. 55, t. 45.—Hooker, Fl. Bor.-Am. i, 108.—Don, Miller's Dict. i, 553.—Eaton, Manual, 6 ed. 365.—Beck, Bot. 59.—Darlington, Fl. Cestrica, 2 ed. 312.—Eaton & Wright, Bot. 452.—Dietrich, Syn. iii, 237.—Richardson, Arctic Exped. 422.

T. latifolia, Salisbury, Prodr. 367.

T. Canadensis, Michaux, Fl. Bor.-Am. 306.—Persoon, Syn. ii, 66.—Poiret in Lamarck, Dict. vii, 683.

T. neglecta, Spach, Ann. Sci. Nat. 2 ser. ii, 346, s. 15; Hist. Veg. iv, 27, 29.—Walpers, Rep. i, 359.

LIME TREE. BASS WOOD. AMERICAN LINDEN. LIN. BEE TREE.

Northern New Brunswick, westward in British America to about the one hundred and second meridian, southward to Virginia and along the Alleghany mountains to Georgia and southern Alabama; extending west in the United States to eastern Dakota, eastern Nebraska, eastern Kansas, the Indian territory, and southwest to the valley of the San Antonio river, Texas.

A large tree, 20 to 24 meters in height, with a trunk 0.90 to 1.20 meter in diameter, or, exceptionally, 30 to 45 meters in height, with a trunk 0.92 to 1.84 meter in diameter (valley of the lower Wabash river, *Ridgway*); common in all northern forests, and always an indication of rich soil; toward its western and southwestern limits only along river bottoms.

Wood light, soft, not strong, very close-grained, compact, easily worked; medullary rays numerous, rather obscure; color, light brown, or often slightly tinged with red, the sap-wood hardly distinguishable; specific gravity, 0.4525; ash, 0.55; largely used in the manufacture of woodenware and cheap furniture, for the panels and bodies of carriages, the inner soles of shoes, in turnery, and the manufacture of paper-pulp (the quickly-discolored sap renders it unfit for making white paper).

The inner bark, macerated, is sometimes manufactured into coarse cordage and matting; the flowers, rich in honey, highly prized by apiarists.

Aqua tilia, an infusion of the flowers, buds, and leaves of the different species of *Tilia*, is used in Europe as a domestic remedy in cases of indigestion, nervousness, etc. (*Nat. Dispensatory*, 2 ed. 1429).

Var. *pubescens*, London,

Arboretum, i, 374 & t.—Browne, *Trees of America*, 48.—Gray, *Manual N. States*, 5 ed. 103; *Hall's Pl. Texas*, 5.

T. Caroliniana, Miller, *Dict.* No. 4.—Wangenheim, *Amer.* 56.—Marshall, *Arbustum*, 154.

T. Americana, Walter, *Fl. Caroliniana*, 153 [not Linnaeus].

T. pubescens, Aiton, *Hort. Kew.* ii, 229; 2 ed. iii, 299.—Willdenow, *Spec.* ii, 1162; *Enum.* i, 566.—Ventenat in *Mem. Acad. Sci.* iv, 10, t. 3.—Nouveau Duhamel, i, 228, t. 51.—Persoon, *Syn.* ii, 66.—Desfontaines, *Hist. Arb.* ii, 37.—Michaux f. *Hist. Arb. Am.* iii, 317, t. 3; *N. American Sylva*, 3 ed. iii, 85, t. 133.—Pursh, *Fl. Am. Sept.* ii, 363.—De Candolle, *Prodr.* i, 513.—Hayne, *Dend. Fl.* 112.—Elliott, *Sk.* ii, 3.—Watson, *Dend. Brit.* ii, t. 135.—Torrey, *Comp. Fl. N. States*, 215.—Don, *Miller's Dict.* i, 553.—Eaton, *Manual*, 6 ed. 365.—Beck, *Bot.* 59.—Eaton & Wright, *Bot.* 452.—Penn. *Cycl.* xxiv, 447.—Dietrich, *Syn.* iii, 237.—Darby, *Bot. S. States*, 262.—Chapman, *Fl. S. States*, 59.—Curtis in *Rep. Geological Surv. N. Carolina*, 1860, iii, 79.—Walpers, *Ann.* vii, 449.—Koch, *Dendrologie*, i, 479.—Vasey, *Cat. Forest Trees*, 7.

T. laxiflora, Michaux, *Fl. Bor.-Am.* i, 306.—Poiret in *Lamarek, Dict.* vii, 683.—Persoon, *Syn.* ii, 66.—Willdenow, *Enum. Suppl.* 38.—De Candolle, *Prodr.* i, 513.—Hayne, *Dend. Fl.* 113.—Torrey, *Compend. Fl. N. States*, 215.—Don, *Miller's Dict.* i, 553.—Eaton, *Manual*, 6 ed. 365.—Beck, *Bot.* 59.—Spach, *Ann. Sci. Nat.* 2 ser. ii, 343, t. 15; *Hist. Veg.* iv, 32.—Browne, *Trees of America*, 48.—Dietrich, *Syn.* iii, 237.

T. grata, Salisbury, *Prodr.* 367.

T. pubescens, var. *leptophylla*, Pursh, *Fl. Am. Sept.* ii, 63.

? *T. stenopetala*, Rafinesque, *Fl. Ludoviciana*, 92.—Robin, *Voyages*, iii, 484.

T. truncata, Spach, *Ann. Sci. Nat.* 2 ser. ii, 342; *Hist. Veg.* iv, 30.—Dietrich, *Syn.* iii, 237.

T. Americana, var. *Walteri*, Wood, *Cl. Book*, 272; *Bot. & Fl.* 64.

North Carolina to the Chattahoochee region of western Florida, usually near the coast; Houston, Texas (*H. Hall*).

A small tree, rarely exceeding 15 meters in height, with a trunk 0.30 meter in diameter; swamps or low ground; rare, or often confounded with the typical *T. Americana*.

Wood lighter, but not otherwise distinguishable from that of *T. Americana*; specific gravity 0.4074; ash, 0.65.

18.—*Tilia heterophylla*, Ventenat,

Mem. Acad. Sci. iv, 16, t. 5.—Nouveau Duhamel, i, 229.—Poiret in *Lamarek, Dict.* vii, 683.—Pursh, *Fl. Am. Sept.* ii, 363.—Nuttall, *Genera*, ii, 3; *Sylva*, i, 90, t. 23; 2 ed. i, 107, t. 23.—De Candolle, *Prodr.* i, 513.—Don, *Miller's Dict.* i, 553.—Eaton, *Manual*, 6 ed. 365.—Spach in *Ann. Sci. Nat.* 2 ser. ii, 345; *Hist. Veg.* iv, 34.—Torrey & Gray, *Fl. N. America*, i, 239.—Eaton & Wright, *Bot.* 452.—Penn. *Cycl.* xxiv, 447.—Walpers, *Rep.* i, 359.—Dietrich, *Syn.* iii, 237.—Cooper in *Smithsonian Rep.* 1858, 250.—Chapman, *Fl. S. States*, 60.—Curtis in *Rep. Geological Surv. N. Carolina*, 1860, iii, 79.—Wood, *Cl. Book*, 272; *Bot. & Fl.* 64.—Gray, *Manual N. States*, 5 ed. 103.—Vasey, *Cat. Forest Trees*, 7.—*Nat. Dispensatory*, 2 ed. 1429.—*Ridgway in Proc. U. S. Nat. Mus.* 1882, 61.

T. alba, Michaux f. *Hist. Arb. Am.* iii, 315, t. 2; *N. American Sylva*, 3 ed. iii, 84, t. 132 [not Waldstein & Kitaibel].—Eaton & Wright, *Bot.* 452.—Darby, *Bot. S. States*, 262.

T. laxiflora, Pursh, *Fl. Am. Sept.* ii, 363 [not Michaux].—Elliott, *Sk.* ii, 2.

T. Americana, var. *heterophylla*, Loudon, *Arboretum*, i, 375 & t.

T. heterophylla, var. *alba*, Wood, *Cl. Book*, 272; *Bot. & Fl.* 64.

WHITE BASS WOOD. WAHOO.

Mountains of Pennsylvania, southward along the Alleghany mountains to northern Alabama and Florida (valley of the Apalachicola river, opposite Chattahoochee, *Mohr*), west to middle Tennessee and Kentucky, southern Indiana, and southern and central Illinois (valley of the Illinois river).

A tree 15 to 20 meters in height, with a trunk 0.60 to 1.20 meter in diameter; rich woods and river bottoms, often on limestone; most common and reaching its greatest development along the western slopes of the southern Alleghany mountains and in middle Tennessee.

Wood light, soft, not strong, close-grained, compact, easily worked; medullary rays numerous, obscure; color, light brown, the sap-wood hardly distinguishable; specific gravity, 0.4253; ash, 0.62; generally confounded with that of *Tilia Americana*, and used for similar purposes.

MALPIGHIACEÆ.

19.—*Byrsonima lucida*, HBK.

Nov. Gen. & Spec. v, 147.—De Candolle, Prodr. i, 580.—Jussieu, Mon. Malpig. ii, 40.—Walpers, Rep. v, 168.—Richard, Fl. Cuba, 115, t. 28.—Grisebach, Fl. British West Indies, 115.—Chapman, Fl. S. States, 82.

Malpighia lucida, Swartz, Fl. Ind. Occ. ii, 852.

TALLOWBERRY. GLAMBERRY.

Semi-tropical Florida, on the southern keys (Boca Chica, No-Name Key, etc.); through the West Indies.

A small tree, sometimes 6 to 8 meters in height, with a trunk 0.15 to 0.25 meter in diameter, or often branching from the ground, and frutescent in habit.

Wood light, soft, weak, close-grained, compact; medullary rays numerous, thin; color, light red, the sap-wood a little lighter; specific gravity, 0.5888; ash, 2.46.

Fruit edible.

ZYGOHYLLACEÆ.

20.—*Guaiacum sanctum*, Linnæus,

Spec. 1 ed. 382.—De Candolle, Prodr. i, 707.—Nuttall, Sylva, iii, 16, t. 86; 2 ed. ii, 86, t. 86.—Gray, Genera, ii, 123, t. 148.—Schnizlein, Icon. t. 253, f. 21.—Cooper in Smithsonian Rep. 1858, 264.—Grisebach, Fl. British West Indies, 134.—Chapman, Fl. S. States, 64.—Wood, Bot. & Fl. 67.—Vasey, Cat. Forest Trees, 7.

G. verticale, Richard, Fl. Cuba, 321.

LIGNUM-VITÆ.

Semi-tropical Florida, Upper Metacombe and Lignum-Vitæ Keys, common; Lower Metacombe and Umbrella Keys, rare; in the Bahamas, St. Domingo, Cuba, Porto Rico, etc.

A low, gnarled tree, not exceeding, within the limits of the United States, 8 meters in height, with a trunk sometimes 0.30 meter in diameter.

Wood exceedingly heavy, very hard, strong, brittle, close-grained, compact, difficult to work, splitting irregularly, containing many evenly-distributed resinous ducts; medullary rays numerous, obscure; color, rich yellow-brown, varying in older specimens to almost black, the sap-wood light yellow; specific gravity, 1.1432; ash, 0.82; used in turnery and for the sheaves of ships' blocks, for which it is preferred to other woods.

Lignum Guaiaci, *Guaiacum wood*, the heart of this and the allied *G. officinale*, Linnæus, formerly largely used in the treatment of syphilis, is now only retained in the *Materia Medica* as an ingredient in the compound decoction of sarsaparilla.

Guaiac, the resinous gum obtained from these species, is a stimulating diaphoretic and alterative, or in large doses cathartic, and is still employed in cases of chronic rheumatism, gout, etc. (*Flückiger & Hanbury, Pharmacographia*, 92.—*U. S. Dispensatory*, 14 ed. 456.—*Nat. Dispensatory*, 2 ed. 696.—*Guibourt, Hist. Drogues*, 7 ed. iii, 551.—*Berg, Pharm. Anat. Atl.* 53, t. 27).

21.—*Porlieria angustifolia*, Gray,

Smithsonian Contrib. iii, 28.—Torrey, Bot. Mex. Boundary Survey, 42.

Guaiacum angustifolium, Engelm., Wislizenus' Rep. 29.—Gray in Jour. Boston Soc. Nat. Hist. vi, 158; Genera, ii, 123, t. 149.—Walpers, Ann. iii, 840.—Watson in Proc. Am. Acad. xvii, 334.

Western Texas, valley of the Colorado river to the Rio Grande (Austin, Matagorda bay, New Braunfels, San Antonio, Brownsville, Fort McIntosh), extending west to the Rio Pecos (*Havard*); in northern Mexico.

A small tree, 8 to 10 meters in height, with a trunk 0.15 to 0.20 meter in diameter, or toward its eastern, northern, and western limits reduced to a low shrub; reaching its greatest development in the United States on the calcareous hillsides bordering the valley of the Gualalupe river.

Wood exceedingly heavy, very hard, close-grained, compact, the open ducts smaller and less regularly distributed than in *Guaiacum*; medullary rays very thin, numerous; color, rich dark brown, turning green with exposure, the sap-wood bright yellow; specific gravity, 1.1101; ash, 0.51; probably possessing medicinal properties similar to those of *lignum-vitæ*.

R U T A C E Æ.

22.—*Xanthoxylum Americanum*, Miller,

Dict. No. 2.—Du Roi, Obs. Bot. 57.—Wangenheim, Amer. 116.—Torrey & Gray, Fl. N. America, i, 214.—Torrey in Nicolle's Rep. 147.—Emerson, Trees Massachusetts, 509; 2 ed. ii, 581.—Gray, Genera, ii, 148, t. 156; Pacific R. R. Rep. xii³, 41; Manual N. States, 5 ed. 110.—Richardson, Arctic Exped. 423.—Parry in Owen's Rep. 610.—Darby, Bot. S. States, 253.—Cooper in Smithsonian Rep. 1858, 250.—Wood, Cl. Book, 282; Bot. & Fl. 70.—Engelmann in Trans. Am. Phil. Soc. new ser. xii, 187.—Koch, Dendrologie, i, 563.—Vasey, Cat. Forest Trees, 8.

X. Olava-Herculis, Lamarck, Dict. ii, 38; Ill. t. 811, f. 3 [not Linnaeus].—Aiton, Hort. Kew. iii, 399.—Mench, Meth. 340.

X. fraxinifolium, Marshall, Arbustum, 167.—B. S. Barton, Coll. i, 52; ii, 38.

X. fraxineum, Willdenow, Spec. iv, 757; Enum. 1013; Berl. Baumz. 413.—Persoon, Syn. ii, 615.—Desfontaines, Hist. Arb. ii, 343.—Aiton, Hort. Kew. 2 ed. v, 383.—Pursh, Fl. Am. Sept. 1, 210.—Nuttall, Genera, ii, 236.—Nouveau Duhamel, vii, 3, t. 2.—Hayne, Dend. Fl. 197.—Bigelow, Med. Bot. iii, 156, t. 59; Fl. Boston. 3 ed. 405.—De Candolle, Prodr. i, 726.—Sprengel, Syst. i, 945.—Torrey, Compend. Fl. N. States, 373.—Rafinesque, Med. Bot. ii, 113, f. 96.—Don, Miller's Dict. i, 802.—Eaton, Manual, 6 ed. 399.—Beck, Bot. 70.—Spach, Hist. Veg. ii, 364.—Lindley, Fl. Med. 216.—London, Arboretum, i, 488, f. 158 & t.—Dietrich, Syn. ii, 1000.—Hooker, Fl. Bor.-Am. i, 118.—Eaton & Wright, Bot. 482.—Nees, Pl. Wied. 5.—Griffith, Med. Bot. 195, f. 103.—Browne, Trees of America, 150.—Agardh, Theor. & Syst. Pl. t. 19, f. 9.—Schnizlein, Icon. t. 250, f. 1-14.—Maout & Decaisne, Bot. English ed. 324 & figs.—Baillon, Hist. Pl. iv, 398, f. 433-438.

X. mite, Willdenow, Enum. 1013.—Poirot, Suppl. v, 622.—De Candolle, Prodr. i, 727.—Don, Miller's Dict. i, 802.—London, Arboretum, i, 489.

X. ramiflorum, Michaux, Fl. Bor.-Am. ii, 235.

X. tricarpum, Hooker, Fl. Bor.-Am. i, 118 [not Michaux].

Thylax fraxineum, Rafinesque, Med. Bot. ii, 114.

PRICKLY ASH. TOOTHACHE TREE.

Eastern Massachusetts, west to northern Minnesota, eastern Nebraska, and eastern Kansas, south to the mountains of Virginia and northern Missouri.

A small tree, not often 7 meters in height, with a trunk 0.15 to 0.20 meter in diameter; or, reduced to a shrub, 1.50 to 1.80 meter in height; common and reaching its greatest development in the region of the great lakes; rocky hillsides, or more often along streams and rich river bottoms.

Wood light, soft, coarse-grained; medullary rays numerous, thin; color, light brown, the sap-wood lighter; specific gravity, 0.5654; ash, 0.57.

The bark of *Xanthoxylum*, an active stimulant, is used in decoction to produce diaphoresis in cases of rheumatism, syphilis, etc., and as a popular remedy for toothache (*U. S. Dispensatory*, 14 ed. 940.—*Bentley in London Pharm. Jour.* 2 ser. v, 399.—*Guibourt, Hist. Drogues*, 7 ed. iii, 562.—*Nat. Dispensatory*, 2 ed. 1535).

23.—*Xanthoxylum Clava-Herculis*, Linnaeus,

Spec. 1 ed. 270, in part.—B. S. Barton, Coll. i, 25, 52; ii, 38.—Willdenow, Spec. iv, 754, in part.—Aiton, Hort. Kew. 2 ed. v, 382.—Elliott, Sk. ii, 690.—Planchon & Triana in Ann. Sci. Nat. 5 ser. xiv, 312.

X. fraxinifolium, Walter, Fl. Caroliniana, 243 [not Marshall].

Fagara fraxinifolia, Lamarck, Ill. i, 334.

X. Carolinianum, Lamarck, Dict. ii, 39; Ill. 403, t. 811, f. 1.—Torrey & Gray, Fl. N. America, i, 214.—Engelmann & Gray in Jour. Boston Soc. Nat. Hist. v, 213.—Gray, Genera, ii, 148, t. 156, f. 13, 14; Manual N. States, 5 ed. 110; Hall's Fl. Texas, 5.—Scheele in Roemer, Texas, 432.—Nuttall, Sylva, iii, 8, t. 83; 2 ed. ii, 78, t. 83.—Darby, Bot. S. States, 253.—Cooper in Smithsonian Rep. 1858, 250.—Chapman, Fl. S. States, 66.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 103.—Wood, Cl. Book, 292; Bot. & Fl. 70.—Young, Bot. Texas, 194.—Vasey, Cat. Forest Trees, 8.

X. aromaticum, Willdenow, Spec. iv, 755 (excl. syn.).—Jacquin f. Eclogæ, i, 103, t. 70.

X. tricarpum, Michaux, Fl. Bor.-Am. ii, 235.—Poiret, Suppl. ii, 294.—Aiton, Hort. Kew. 2 ed. v, 383.—Pursh, Fl. Am. Sept. i, 210.—De Candolle, Prodr. i, 726.—Elliott, Sk. ii, 690.—A. de Jussieu in Mem. Mus. xii, t. 26, f. 38.—Sprengel, Syst. i, 945.—Don, Miller's Dict. i, 803.—Spach, Hist. Veg. ii, 365.—London, Arboretum, i, 488.—Eaton, Manual, 6 ed. 399.—Eaton & Wright, Bot. 482.—Dietrich, Syn. ii, 1000.

Kampmania fraxinifolia, Rafinesque, Med. Rep. v, 354.

Pseudopetalon glandulosum, Rafinesque, Fl. Ludoviciana, 108; Med. Bot. ii, 114.

Pseudopetalon tricarpum, Rafinesque, Fl. Ludoviciana, 108; Med. Bot. ii, 114.

X. Catesbianum, Rafinesque, Med. Bot. ii, 114.

TOOTHACHE TREE. PRICKLY ASH. SEA ASH. PEPPEE WOOD. WILD ORANGE.

Southern Virginia, southward near the coast to bay Biscayne and Tampa bay, Florida, westward through the Gulf states to northwestern Louisiana, southern Arkansas (south of the Arkansas river), and the valley of the Brazos river, Texas.

A small tree, rarely 12 to 14 meters in height, with a trunk 0.30 meter in diameter, of very rapid growth; usually along streams and low, rich river bottoms, reaching its greatest development in southern Arkansas, Louisiana, and eastern Texas.

A form with trifoliate leaves is—

X. macrophyllum, Nuttall, Sylva, iii, 10; 2 ed. ii, 80.—Lesquereux in Owen's 2d Rep. Arkansas, 353.

X. Clava-Herculis, var. Watson in Proc. Am. Acad. xvii, 335.

Wood light, hard, not strong, soft, coarse-grained, not durable, containing many scattered open ducts; medullary rays numerous, thin; color, light brown, the sap-wood lighter; specific gravity, 0.5056; ash, 0.82.

X. Clava-Herculis probably possesses similar medicinal properties to those of the last species (*Nat. Dispensatory* 2 ed. 1535).

Var. *fruticosum*, Gray,

Smithsonian Contrib. iii, 30.—Torrey & Gray in Pacific R. R. Rep. ii, 161.—Torrey, Bot. Mex. Boundary Survey, 43.—Chapman, Fl. S. States, 66?—Wood, Bot. & Fl. 71.

X. hirsutum, Buckley in Proc. Philadelphia Acad. 1861, 450; 1870, 136 (see Gray in same, 1862, 162).—Young, Bot. Texas, 195.

Western Texas, Corpus Christi (*Buckley*), mouth of the Colorado river (*Mohr*), near Austin, and west to Devil's river and Eagle pass; Florida (?) (*Chapman l. c.*).

A low shrub, or on the Texas coast a small tree, 6 to 8 meters in height, with a trunk 0.20 to 0.30 meter in diameter.

Wood light, soft, close-grained, compact; medullary rays numerous, thin; color, light brown, the sap-wood yellow; specific gravity, 0.5967; ash, 0.76.

24.—*Xanthoxylum Caribæum*, Lamarck,

Dict. ii, 40.—Gærtner, Fruct. i, 333, t. 68, f. 8.—Descourtilz, Fl. Med. Antilles, ii, 58.—Planchon & Triana in Ann. Sci. Nat. 5 ser. xiv, 315.—Guibourt, Hist. Drogues, 7 ed. iii, 562.

X. Clava-Herculis, Linnaeus, Spec. 1 ed. 270, in part.—De Candolle, Prodr. i, 727.—Macfadyen, Fl. Jamaica, 194.—Grisebach, Fl. British West Indies, 138.

X. lanceolatum, Poiret, Suppl. ii, 293.—De Candolle, Prodr. i, 727.

X. Floridanum, Nuttall, Sylva, iii, 14, t. 85; 2 ed. ii, 85, t. 85.—Chapman, Fl. S. States, 66.—Wood, Bot. & Fl. 70.—Young, Bot. Texas, 194.—Vasey, Cat. Forest Trees, 8.

SATIN WOOD.

Semi-tropical Florida, south Bahia Honda and Boca Chica Keys; in the West Indies.

A small tree, 6 to 10 meters in height, with a trunk 0.30 to 0.40 meter in diameter.

Wood very heavy, exceedingly hard, not strong, brittle, fine-grained, compact, satiny, susceptible of a beautiful polish; medullary rays numerous, thin, conspicuous; color, light orange, the sap-wood lighter; specific gravity, 0.9002; ash, 2.02.

25.—*Xanthoxylum Pterota*, HBK.

Nov. Gen. & Spec. vi, 3.—Kunth, Syn. iii, 325.—De Candolle, Prodr. i, 725.—Torrey & Gray, Fl. N. America, i, 680.—Macfadyen, Fl. Jamaica, 190.—Nuttall, Sylva, iii, 11, t. 84; 2 ed. ii, 81, t. 84.—Seemann, Bot. Herald, 275.—Torrey, Bot. Mex. Boundary Survey, 43.—Cooper in Smithsonian Rep. 1858, 264.—Chapman, Fl. S. States, 66.—Young, Bot. Texas, 195.—Planchon & Triana in Ann. Sci. Nat. 5 ser. xiv, 311.—Engler in Martius, Fl. Brasil. xiii, 154.—Vasey, Cat. Forest Trees, 8.—Hemsley, Bot. Am.-Cent. i, 169.—Watson in Proc. Am. Acad. xvii, 335.

Fagara Pterota, Linnaeus, Amoen, v, 393, in part.—Lamarck, Dict. ii, 444; Ill. i, 335, t. 84.—Willdenow, Spec. i, 666.—Aiton, Hort. Kew. 2 ed. i, 263.—Titford, Hort. Bot. Am. 40.—Turpin, Dict. Sci. Nat. xvi, 107, t. 127.

Fagara lentiscifolia, Willdenow, Enum. i, 166.—Grisebach, Fl. British West Indies, 137.

WILD LIME.

Semi-tropical Florida, Mosquito inlet to the southern keys, on the west coast from about latitude 29° to cape Sable; southwestern Texas, and southward through Mexico to Brazil.

A small tree, sometimes 8 meters in height, with a trunk rarely exceeding 0.15 meter in diameter, or often reduced to a slender shrub; in Florida common, and reaching its greatest development on the keys of the west coast; in Texas not common, but widely distributed as a small shrub, or on the shores of Matagorda bay, west of the Nueces river, and in the valley of the Rio Grande a low tree.

Wood heavy, hard, close-grained, compact; medullary rays thin, numerous; color, brown tinged with red, the sap-wood yellow; specific gravity, 0.7444; ash, 0.78.

26.—*Ptelia trifoliata*, Linnaeus,

Spec. 1 ed. 118.—Medicus, Bot. Beobacht. 215.—Marshall, Arbustum, 115.—Walter, Fl. Caroliniana, 88.—Aiton, Hort. Kew. i, 162; 2 ed. i, 264.—Lamarck, Ill. i, 336, t. 84.—Mönch, Meth. 55.—Willdenow, Spec. i, 670; Enum. i, 116.—Nouveau Duhamel, i, 252, t. 57.—Michaux, Fl. Bor. Am. i, 99.—Schkuhr, Handb. 83, t. 83.—Poiret in Lamarck, Dict. v, 706.—Persoon, Syn. i, 145.—Desfontaines, Hist. Arb. ii, 343.—Robin, Voyages, iii, 509.—Pursh, Fl. Am. Sept. i, 107.—Nuttall, Genera, i, 104.—Guimpel, Otto & Hayne, Abb. Holz. 94, t. 74.—Hayne, Dend. Fl. 8.—Elliott, Sk. i, 201.—Römer & Schultes, Syst. iii, 291.—Torrey, Fl. U. S. 189; Compend. Fl. N. States, 86.—Fl. N. York, i, 133; Pacific R. R. Rep. iv, 73; Bot. Mex. Boundary Survey, 43.—De Candolle, Prodr. ii, 82.—Sprengel, Syst. i, 441.—Turpin, Dict. Sci. Nat. xlv, 2, t. 128.—A. de Jussieu in Mem. Mus. xii, t. 26, f. 42.—Beck in Am. Jour. Sci. 1 ser. x, 264; Bot. 71.—Don, Miller's Dict. i, 806.—Spach, Hist. Veg. ii, 369.—Hooker, Jour. Bot. i, 202.—Lindley, Fl. Med. 215.—London, Arboretum, i, 489 & t.—Eaton, Manual, 6 ed. 288.—Torrey & Gray, Fl. N. America, i, 215.—Eaton & Wright, Bot. 379.—Dietrich, Syn. i, 497.—Browne, Trees of America, 153.—Scheele in Römer, Texas, 432.—Gray, Genera, ii, 150, t. 157; Manual N. States, 5 ed. 110.—Richardson, Arctic Exped. 423.—Parry in Owen's Rep. 610.—Agardh, Theor. & Syst. Pl. t. 19, f. 7, 8.—Cooper in Smithsonian Rep. 1858, 250.—Darby, Bot. S. States, 254.—Chapman, Fl. S. States, 66.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 107.—Lesquereux in Owen's 2d Rep. Arkansas, 353.—Wood, Cl. Book, 283; Bot. & Fl. 71.—Schnizlein, Icon. t. 250, f. 15-26.—Young, Bot. Texas, 195.—Baillon, Hist. Pl. iv, 395, f. 445, 446.—Koch, Dendrologie, i, 566.—Vasey, Cat. Forest Trees, 8.—Hemsley, Bot. Am.-Cent. i, 171.—Burgess in Coulter's Bot. Gazette, vii, 95.

Amyris elemifera, Linnaeus, Spec. 2 ed. 295.—St. Hilaire, Fam. Nat. i, 263.

P. viticifolia, Salisbury, Prodr. 68.

HOP TREE. SHRUBBY TREFOIL. WAFER ASH.

Ontario and New York (banks of the Niagara river), Pennsylvania southward to northern Florida, west to Minnesota and the headwaters of the Canadian river; through western Texas to the valley of the Mimbres river, New Mexico (*Bigelow*), and southward into northern Mexico.

A small tree, sometimes 4 to 6 meters in height, with a trunk 0.15 to 0.20 meter in diameter, or more often reduced to a slender shrub; shady, rocky hillsides.

A variety with more or less pubescent leaves, not rare on the south Atlantic coast, and the common form of western Texas, is—

var. *mollis*, Torrey & Gray, Fl. N. America, i, 680.—Engelmann & Gray in Jour. Boston Soc. Nat. Hist. v, 213.—Torrey in Marcey's Rep. 282.—Gray in Smithsonian Contrib. iii, 31; Hall's Pl. Texas, 5.—Wood, Bot. & Fl. 71.—Watson in Proc. Am. Acad. xvii, 335.

P. mollis, Curtis in Am. Jour. Sci. 2 ser. vii, 406; Rep. Geological Surv. N. Carolina, 1860, iii, 107.—Walpers, Ann. ii, 259.—Chapman, Fl. S. States, 67.—Young, Bot. Texas, 196.

Wood heavy, hard, close-grained, compact, satiny, the annual growths clearly marked by two or three rows of open ducts; medullary rays few, thin; color, yellow-brown, the sap-wood hardly distinguishable; specific gravity, 0.8319; ash, 0.30.

The bark of the root possesses tonic properties and is employed by herbalists in the form of tinctures and fluid extracts in cases of dyspepsia, debility, etc. (*Am. Jour. Pharm.* 1862, 198; 1867, 337.—*U. S. Dispensatory*, 14 ed. 1740.—*Nat. Dispensatory*, 2 ed. 1179); the bitter fruit is occasionally used domestically as a substitute for hops.

27.—*Canotia holocantha*, Torrey,

Pacific R. R. Rep. iv, 68.—Gray in *Ives' Rep.* 15; *Proc. Am. Acad.* xii, 159.—Baillon, *Adansonia*, x, 18; *Hist. Veg.* vi, 7, 42.—Brewer & Watson, *Bot. California*, i, 190.—Rothrock in *Wheeler's Rep.* 24, 81, t. 1.—Maximowicz in *Act. Hort. St. Petersburg* v, 256.—Rusby in *Bull. Torrey Bot. Club*, ix, 106.

Arizona, White Mountain region, valley of the Gila river (*Rothrock*), valley of Bill Williams Fork (*Bigelow*).

A small tree, 6 to 8 meters in height, with a trunk sometimes 0.30 meter in diameter, or often a large shrub; dry, rocky *mesas*. Wood heavy, hard, close-grained, compact; medullary rays numerous, not prominent; color light brown, the sap-wood lighter; specific gravity, 0.6885; ash, 5.33.

SIMARUBEÆ.

28.—*Simaruba glauca*, De Candolle,

Diss. in Ann. Mus. xvii, 323; *Prodr.* i, 733.—Humboldt, *Bonpland & Kunth*, *Nov. Gen. et Spec.* vi, 16.—Descourtilz, *Fl. Med. Antilles*, i, 66, t. 14.—Planchon in *London Jour. Bot.* v, 567.—Gray, *Genera*, ii, 152.—Nuttall, *Sylva*, iii, 20, t. 87; 2 ed. ii, 88, t. 87.—Cooper in *Smithsonian Rep.* 1858, 264.—Grisebach, *Fl. British West Indies*, 139.—Chapman, *Fl. S. States*, 67.—Wood, *Bot. & Fl.* 72.—Planchon & Triana in *Ann. Sci. Nat.* 5 ser. xv, 357.—Engler in *Martius*, *Fl. Brasil.* xii³, 223.—Vasey, *Cat. Forest Trees*, 8.—Hemsley, *Bot. Am.-Cent.* i, 173.

Quassia Simaruba, Linnæus, *Suppl.* 234.—Wright, *Trans. Edinburgh Soc.* ii, 73, t. 1, 2; *Bot. & Med. Account of Q. Simaruba*.—Gærtner, *Fruct.* i, 340, t. 70.—Lamarck, *Ill.* ii, 478, t. 343, f. 2.—Willdenow, *Spec.* ii, 568.—Aiton, *Hort. Kew.* 2 ed. iii, 42.—Descourtilz, *Fl. Med. Antilles*, i, 23, t. 5.

Quassia dioica, Bergius, *Mat. Med.* 355.

S. amara, Aublet, *Guian.* t. 331.—Hayne, *Arzn.* iv, t. 15.—Schnizlein, *Icon.* t. 249, f. 1-6.

S. medicinalis, Endlicher, *Medz. Pf.* 525.—Berg, *Handb.* i, 373.—Berg & Schmidt, *Off. Gew.* ii, t. 13.

PARADISE TREE.

Semi-tropical Florida, cape Canaveral to the southern keys; through the West Indies to Brazil.

A tree sometimes 15 meters in height, with a trunk 0.60 meter in diameter; within the United States not common, and reaching its greatest development on the shores of bay Biscayne.

Wood light, soft, not strong, coarse-grained, containing many large scattered open ducts; medullary rays few, thin; color, light brown, the sap-wood a little darker; specific gravity, 0.4136; ash, 0.93.

The bark of this species has been occasionally used as a substitute for that of *S. officinalis*, DC. as an aromatic, bitter tonic (*U. S. Dispensatory*, 14 ed. 838.—*Nat. Dispensatory*, 2 ed. 1294).

BURSERACEÆ.

29.—*Bursera gummifera*, Jacquin,

Am. Piet. t. 65.—Linnæus, *Spec.* 2 ed. 741.—Lamarck, *Ill.* ii, 392, t. 256.—Willdenow, *Spec.* iv, 1119.—Aiton, *Hort. Kew.* 2 ed. v, 481.—Titford, *Hort. Bot. Am.* 107.—De Candolle, *Prodr.* ii, 78.—Descourtilz, *Fl. Med. Antilles*, ii, t. 97.—Spach, *Hist. Veg.* ii, 239.—Macfadyen, *Fl. Jamaica*, 229.—Nuttall, *Sylva*, ii, 117, t. 79; 2 ed. ii, 64, t. 79.—Richard, *Fl. Cuba*, 390.—Browne, *Trees of America*, 189.—Grisebach, *Fl. British West Indies*, 173.—Cooper in *Smithsonian Rep.* 1858, 264; 1860, 440.—Chapman, *Fl. S. States*, 68.—Wood, *Bot. & Fl.* 72.—Planchon & Triana in *Ann. Sci. Nat.* 5 ser. xv, 302.—Vasey, *Cat. Forest Trees*, 8.—Hemsley, *Bot. Am.-Cent.* i, 177.—Engler in *De Candolle*, *Suites*, iv, 39.

B. acuminata, Willdenow, *Spec.* iv, 1120.—De Candolle, *Prodr.* ii, 78.

Blaphrium integerrimum, Tulasne in *Ann. Sci. Nat.* 3 ser. vi, 369. (*Fide Engler, l. c.*)

GUM ELEMI. GUMBO LIMBO. WEST-INDIAN BIRCH.

Semi-tropical Florida, cape Canaveral to the southern keys, west coast Caloosa river and Oaximbas bay; through the West Indies.

A tree often 18 meters in height, with a trunk 0.50 to 0.70 meter in diameter; one of the largest and most common trees of southern Florida, of very rapid growth and decay.

Wood very light, exceedingly soft and weak, spongy, containing many scattered open ducts; medullary rays numerous, thin; color, light brown or gray, quickly discoloring with decay; specific gravity, 0.3003; ash, 2.04; used in making live-fences, pieces of the trunk when planted in the coral rock of the keys throwing out roots and growing rapidly.

The aromatic resin obtained from this species was formerly somewhat used in various forms, under the name of *Caranna*, as a remedy for gout (*Watts, Chem. Dict.* i, 749.—*Guibourt, Hist. Drogues*, 7 ed. iii, 525, f. 749); and in the West Indies is manufactured into a valuable varnish. An infusion of the leaves is occasionally used as a domestic substitute for tea.

30.—*Amyris sylvatica*, Jacquin,

Am. Pict. t. 108.—Willdenow, Spec. ii, 333.—Aiton, Hort. Kew. 2 ed. ii, 351.—De Candolle, Prodr. ii, 81.—Dietrich, Syn. ii, 1271.—Macfadyen, Fl. Jamaica, 231.—Richard, Fl. Cuba, 393.—Grisebach, Fl. British West Indies, 174.—Planchon & Triana in Ann. Sci. Nat. 5 ser. xv, 321.—Vasey, Cat. Forest Trees, 8.

Toxicodendron arborescens, Miller, Dict. No. 9.

A. dyatripa, Sprengel, Neue Entdeck. iii, 48.—De Candolle, Prodr. ii, 81.

Rhus arborescens, De Candolle, Prodr. ii, 73.

A. Plumieri, De Candolle, Prodr. ii, 81.

A. Floridana, Nuttall in Am. Jour. Sci. v, 294; Sylva ii, 114, t. 78; 2 ed. ii, 61, t. 78.—De Candolle, Prodr. ii, 81.—Torrey & Gray, Fl. N. America, i, 221.—Eaton, Manual, 6 ed. 16.—Eaton & Wright, Bot. 123.—London, Arboretum, ii, 561.—Cooper in Smithsonian Rep. 1858, 264.—Chapman, Fl. S. States, 68.—Wood, Bot. & Fl. 72.—Vasey, Cat. Forest Trees, 8.

A. cymosa, Reichenbach in Sieb. Pl. Trin. No. 29 f.

A. maritima, Richard, Fl. Cuba, 392 [not Jacquin].

TORCH WOOD.

Semi-tropical Florida, Mosquito inlet to the southern keys; in the West Indies.

A small tree sometimes 7 meters in height, with a trunk 0.20 to 0.25 meter in diameter; common.

Wood very heavy, exceedingly hard and strong, close-grained, compact, resinous, exceedingly durable, susceptible of a beautiful polish; medullary rays obscure; color, light orange, the sap-wood lighter; specific gravity, 1.0459; ash, 0.59.

MELIACEÆ.

31.—*Swietenia Mahagoni*, Linnæus,

Spec. 2 ed. 548.—Jacquin, Stirp. Am. t. 127.—Cavanilles, Diss. ii, 365, t. 209.—Gærtner, Fruct. ii, 89, t. 96.—Lamarck, Dict. iii, 678.—Willdenow, Spec. ii, 557.—Aiton, Hort. Kew. 2 ed. ii, 338.—Titford, Hort. Bot. Am. 64.—Descourtilz, Fl. Med. Antilles, ii, 125, t. 99.—De Candolle, Prodr. i, 625.—Turpin in Dict. Sci. Nat. Atlas, t. 170.—Tussac, Fl. Antilles, iv, t. 23.—Hayne, Arzn. i, t. 19.—Hooker, Bot. Misc. i, 21, t. 16, 17.—A. de Jussieu in Mem. Mus. xix, 248, t. 11.—Don, Miller's Dict. i, 687, f. 116.—Woodville, Med. Bot. 3 ed. iii, 620, t. 220.—Spach, Hist. Veg. iii, 164, t. 21.—Lindley, Fl. Med. 155.—Macfadyen, Fl. Jamaica, 175.—Torrey & Gray, Fl. N. America, i, 242.—Eaton, Manual, 6 ed. 360.—Eaton & Wright, Bot. 447.—Walpers, Rep. i, 436.—Nuttall, Sylva, ii, 98, t. 75; 2 ed. ii, 46, t. 75.—Richard, Fl. Cuba, 304.—Schnizlein, Icon. t. 226, f. 1.—Cooper in Smithsonian Rep. 1858, 264.—Darby, Bot. S. States, 263.—Chapman, Fl. S. States, 62.—Grisebach, Fl. British West Indies, 131.—Wood, Bot. & Fl. 66.—Baillon, Hist. Pl. v, 478, f. 472-476.—Guibourt, Hist. Drogues, 7 ed. iii, 596.—Tippel & Bollevar, Ausland. Cult. Pfl., Atlas, i, t. 2, f. 1.—C. De Candolle, Suites, i, 723.—Hemsley, Bot. Am.-Cent. i, 183.

S. Senegalensis, Desrousseaux in Lamarck, Dict. iii, 678.

Cedrus Mahagoni, Miller, Dict. No. 2.

3 FOR

MAHOGANY. MADEIRA.

Semi-tropical Florida, on the southern keys (Key Largo, Elliott's Key); through the West Indies, and in Central America.

A large tree, on the Florida keys rarely exceeding 15 meters in height, with a trunk sometimes 0.90 meter in diameter.

Wood heavy, exceedingly hard, very strong, brittle, very close-grained, compact, very durable, susceptible of a high polish; medullary rays numerous, obscure; color, rich reddish-brown, turning darker with age, the thin sap-wood yellow; specific gravity, 0.7282; ash, 1.09; varying greatly in quality in different regions; largely used and preferred to all other woods for cabinet-making of all sorts, interior finish, etc.; formerly somewhat employed in ship-building.

OLACINÆ.

32.—*Ximenia Americana*, Linnæus,

Spec. 1 ed. Appx. 1193.—Bartram, Travels, 2 ed. 112.—Lamarck, Ill. ii, 435, t. 297.—Willdenow, Spec. ii, 338.—Aiton, Hort. Kew. 2 ed. ii, 352.—De Candolle, Prodr. i, 533.—Nuttall, Sylva, i, 124, t. 36; 2 ed. i, 138, t. 36.—Schnizlein, Icon. t. 223, f. 1-9, 30, 31.—Cambessedes in St. Hilaire, Fl. Brasil. i, 341.—Wight & Walker-Arnott, Prodr. Fl. Penins. Or. i, 89.—Walpers, Rep. i, 377; Ann. vi, 565.—Richard, Fl. Cuba, 304.—Cooper in Smithsonian Rep. 1858, 264.—Grisebach, Fl. British West Indies, 310.—Baillon, Adansonia, ii, t. 9, f. 5, 6.—Chapman, Fl. S. States, 61.—Engler in Martius, Fl. Brasil. xii, 9, t. 2, f. 1.—Vasey, Cat. Forest Trees, 8.—Hemsley, Bot. Am.-Cent. i, 185.

Heymassoli spinosa, Aublet, Guian. i, 324, t. 125.—Lamarck, Ill. ii, 435.

X. multiflora, Jacquin, Stirp. Am. 106, t. 177, f. 31.—Lamarck, Ill. ii, 435, t. 297, f. 1, 2.—Spach, Hist. Veg. xiii, 264.

X. montana, Macfadyen, Fl. Jamaica, i, 121.

WILD LIME. TALLOW NUT. HOG PLUM. MOUNTAIN PLUM.

Florida, east coast from the Saint John's river to the southern keys, west coast Caloosa river to Caximbas bay; through the West Indies to Brazil, and on the coast of the Indian peninsula (introduced?, *A. De Candolle*, Geog. Bot. ii, 1027).

A small, low, wide-spreading tree, rarely exceeding 4 meters in height, with a trunk 0.15 meter in diameter, or in pine-barren soil and toward its northern limits reduced to a low shrub; common and reaching its greatest development in Florida on the west coast.

Wood very heavy, tough, hard, close-grained, compact, containing numerous regularly-distributed open ducts; medullary rays few, thin; color, brown, tinged with red, the sap-wood lighter; specific gravity, 0.9196; ash, 0.73.

Hydrocyanic acid has been obtained from the edible plum-shaped fruit (*Flückiger & Hanbury, Pharmacographia*, 222).

ILICINÆ.

33.—*Ilex opaca*, Aiton,

Hort. Kew. i, 169; 2 ed. i, 277.—Willdenow, Spec. i, 708; Enum. 172; Berl. Baumz. 190.—Nouveau Duhamel, i, 8.—Michaux, Fl. Bor.-Am. ii, 223.—Persoon, Syn. i, 151.—Poirer, Suppl. iii, 65.—Michaux f. Hist. Arb. Am. ii, 191, t. 11; N. American Sylva, 3 ed. ii, 122, t. 84.—Barton, Prodr. Fl. Philadelph. 95; Compend. Fl. Philadelph. 94.—Pursh, Fl. Am. Sept. i, 117.—Rafinesque, Fl. Ludoviciana, 111; Med. Bot. ii, 7, t. 53.—Nuttall, Genera, i, 109.—Rœmer & Schultes, Syst. iii, 487.—Link, Enum. 147.—James, Cat. 176; Long's Exped. ii, 294.—Hayne, Dend. Fl. 10.—Torrey in Ann. Lyc. N. York, ii, 173; Fl. U. S. 194; Compend. Fl. N. States, 87; Fl. N. York, ii, 2.—Elliott, Sk. ii, 679.—De Candolle, Prodr. ii, 14.—Sprengel, Syst. i, 495.—Watson, Dend. Brit. i, t. 3.—Beck, Bot. 230.—Eaton, Manual, 6 ed. 186.—London, Arboretum, ii, 516 & t.—Hooker, Fl. Bor.-Am. i, 121; Jour. Bot. i, 201.—Eaton & Wright, Bot. 232.—Bigelow, Fl. Boston. 3 ed. 64.—Don, Miller's Dict. ii, 17.—Spach, Hist. Veg. ii, 427.—Dietrich, Syn. i, 554.—Griffith, Med. Bot. 432.—Emerson, Trees Massachusetts, 341; 2 ed. ii, 385 & t.—Browne, Trees of America, 167.—Darby, Bot. S. States, 426.—Darlington, Fl. Cestrica, 3 ed. 17.—Cooper in Smithsonian Rep. 1858, 253.—Chapman, Fl. S. States, 269.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 58.—Lesquereux in Owen's 2d Rep. Arkansas, 373.—Wood, Cl. Book, 496; Bot. & Fl. 207.—Gray, Manual N. States, 5 ed. 306.—Young, Bot. Texas, 372.—Vasey, Cat. Forest Trees, 8.—Maximowicz in Mem. Acad. St. Petersburg, xxix, No. 3, 29.—Mellichamp in Bull. Torrey Bot. Club, viii, 113.

I. aquifolium, Marshall, Arbustum, 63 [not Linnæus].—Walter, Fl. Caroliniana, 241.

I. Canadensis, Marshall, Arbustum, 64.

I. laxiflora, Lamarck, Dict. iii, 147; Ill. i, 355.—Pursh, Fl. Am. Sept. i, 117.—Rømer & Schultes, Syst. iii, 494; Mant. 334.—De Candolle, Prodr. ii, 14.—Sprengel, Syst. i, 495.—Don, Miller's Dict. ii, 17.—Spach, Hist. Veg. ii, 427.—Dietrich, Syn. i, 555.—London, Arboretum, ii, 517.—Eaton, Manual, 6 ed. 186.—Eaton & Wright, Bot. 282.

I. quercifolia, Meerburgh, Icon. ii, t. 5.

Ageria opaca, Rafinesque, Sylva Telluriana, 47.

AMERICAN HOLLY.

Quincy, Massachusetts, southward, near the coast, to Mosquito inlet and Charlotte harbor, Florida, valley of the Mississippi river, southern Indiana southward to the gulf of Mexico, and southwest through Missouri, Arkansas, and eastern Texas to the valley of the Colorado river.

An evergreen tree, sometimes 15 meters in height, with a trunk 0.30 to 1.20 meter in diameter, or toward its northern limits reduced to a shrub; generally in low, rather moist soil; most common and reaching its greatest development in the rich bottoms of southern Arkansas and eastern Texas.

Wood light, soft, not strong, tough, rather hard, close-grained, very compact, easily worked; medullary rays numerous, inconspicuous; color, nearly white, turning to light brown with exposure, the sap-wood still lighter; specific gravity, 0.5818; ash, 0.76; used and admirably adapted for cabinet work, interior finish, and turnery of the highest class.

A bitter principle (*Ilicin*), common to other species of the genus, has been obtained from the fruit of this tree (*Am. Jour. Pharm.* xxviii, 314.—*U. S. Dispensatory*, 14 ed. 1670.—*Nat. Dispensatory*, 2 ed. 754).

34.—*Ilex Dahoon*, Walter,

Fl. Caroliniana, 241.—Michaux, Fl. Bor.-Am. ii, 228.—Pursh, Fl. Am. Sept. i, 117.—Nuttall, Genera, i, 109.—Rømer & Schultes, Syst. iii, 489; Mant. 332.—De Candolle, Prodr. ii, 14.—Elliott, Sk. ii, 680.—Watson, Dend. Brit. ii, t. 114.—Sprengel, Syst. i, 495.—Audubon, Birds, t. 43.—Don, Miller's Dict. ii, 19.—Hooker, Jour. Bot. i, 202.—Eaton, Manual, 6 ed. 186.—Eaton & Wright, Bot. 282.—Spach, Hist. Veg. ii, 428.—Dietrich, Syn. i, 554.—London, Arboretum, ii, 519.—Griffith, Med. Bot. 433.—Darby, Bot. S. States, 426.—Chapman, Fl. S. States, 269.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 58.—Wood, Bot. & Fl. 207.—Gray, Manual N. States, 5 ed. 306.—Vasey, Cat. Forest Trees, 8.—Maximowicz in Mem. Acad. St. Petersburg, xxix, No. 3, 29.—*Nat. Dispensatory*, 2 ed. 755.

I. Cassine, Linnæus, Spec. 125, in part.—Marshall, Arbustum, 64.—Aiton, Hort. Kew. i, 170, in part; 2 ed. i, 279.—Lamarck, Dict. iii, 147; Ill. i, 355.—Willdenow, Spec. i, 709; Enum. i, 172; Hort. Berol. i, t. 31.—Nouveau Duhamel, i, 9.—Persoon, Syn. 151.—Desfontaines, Hist. Arb. ii, 362.—Poiret, Suppl. iii, 65.—Pursh, Fl. Am. Sept. i, 117.—Rømer & Schultes, Syst. iii, 490.—Hayne, Dend. Fl. 10.—De Candolle, Prodr. ii, 14.—Sprengel, Syst. i, 495.—Don, Miller's Dict. ii, 17.—Spach, Hist. Veg. ii, 428.—Dietrich, Syn. i, 544.—London, Arboretum, ii, 517, f. 184.—Eaton & Wright, Bot. 282.—Gœppert in Del. Sem. Vratisl. 1885 (*Linnæa*, xxvi, 746).

I. Cassine, var. *latifolia*, Aiton, Hort. Kew. 2 ed. i, 278.

I. cassinoides, Link, Enum. i, 148.—Rømer & Schultes, Syst. iii; Mant. 332.

I. laurifolia, Nuttall in Am. Jour. Sci. 1 ser. v, 289.—Eaton, Manual, 6 ed. 186.—Eaton & Wright, Bot. 282.

Ageria palustris, Rafinesque, Sylva Telluriana, 47.

Ageria obovata, Rafinesque, Sylva Telluriana, 47.

Ageria heterophylla, Rafinesque, Sylva Telluriana, 48.

DAHOON. DAHOON HOLLY.

Southern Virginia, southward near the coast to Mosquito inlet and Tampa bay, Florida, west along the Gulf coast to the prairie region of western Louisiana.

A small tree, sometimes 8 meters in height, with a trunk from 0.20 to 0.30 meter in diameter; low, wet soil; not common, and running into numerous forms, of which the best marked are—

var. *angustifolia*, Torrey & Gray, Fl. N. America, *ined.*

I. Cassine, var. *angustifolia*, Willdenow, Spec. i, 709.—Aiton, Hort. Kew. 2 ed. i, 278.—Nouveau Duhamel, i, 9, t. 3.

I. angustifolia, Willdenow, Enum. i, 172.—Pursh, Fl. Am. Sept. i, 118.—Nuttall, Genera, i, 109.—Rømer & Schultes, Syst. iii, 489.—De Candolle, Prodr. ii, 14.—Watson, Dend. Brit. i, t. 4.—Sprengel, Syst. i, 495.—Don, Miller's Dict. ii, 17.—Hooker, Jour. Bot. i, 201.—Spach, Hist. Veg. ii, 428.—Dietrich, Syn. i, 554.—London, Arboretum, ii, 517, f. 185.

I. ligustrina, Elliott, Sk. ii, 708 [not Jacquin].—Spach, Hist. Veg. ii, 429.—Eaton, Manual, 6 ed. 187.—Eaton & Wright, Bot. 282.—Darby, Bot. S. States, 123.

? *I. Watsoniana*, Spach, Hist. Veg. ii, 429.

var. *myrtifolia* (only in low cypress swamps and ponds), Chapman, Fl. S. States, 269.—Nat. Dispensatory, 2 ed. 755.

I. myrtifolia, Walter, Fl. Caroliniana, 214.—Nouveau Duhamel, i, 10, t. 4.—Michaux, Fl. Bor.-Am. ii, 229.—Poirét, Suppl. iii, 65.—Willdenow, Enum. Suppl. 8.—Rømer & Schultes, Syst. iii, 489.—Link, Enum. 148.—Spach, Hist. Veg. ii, 429.—Eaton, Manual, 6 ed. 187.—Eaton & Wright, Bot. 282.—Darby, Bot. S. States, 426.—Gray, Manual N. States, 5 ed. 306.—Maximowicz in Mem. Acad. St. Petersburg, xxix, No. 3, 26.

I. rosmarifolia, Lamarck, Ill. i, 356.—Persoon, Syn. i, 151.—Poirét, Suppl. iii, 65.

I. ligustrifolia, Don, Miller's Dict. ii, 19.—Eaton, Manual, 6 ed. 187.—Wood, Cl. Book, 497; Bot. & Fl. 207.

Wood light, soft, not strong, close-grained, compact; medullary rays numerous, thin; color, light brown, the sap-wood nearly white; specific gravity, 0.4806; ash, 0.91; that of var. *myrtifolia* heavier, nearly white; specific gravity, 0.5873; ash, 0.90.

35.—*Ilex Cassine*, Walter,

Fl. Caroliniana, 241.—Aiton, Hort. Kew. i, 170, in part.—James, Cat. 176; Long's Exped. ii, 294.—Hooker, Jour. Bot. i, 202.—Eaton, Manual, 6 ed. 186.—Chapman, Fl. S. States, 269.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 59.—Lesquereux in Owen's 2d Rep. Arkansas, 373.—Wood, Bot. & Fl. 208.—Gray, Manual N. States, 5 ed. 306.—Young, Bot. Texas, 373.—Maximowicz in Mem. Acad. St. Petersburg, xxix, No. 3, 22.

I. Cassine, β . Linnæus, Spec. 1 ed. 125.

Cassine Peragua, Linnæus, Mant. ii, 220.—Marshall, Arbustum, 26.—Plenck, Icon. t. 239.

Cassine Caroliniana, Lamarck, Dict. i, 652

I. vomitoria, Aiton, Hort. Kew. i, 170; 2 ed. i, 278.—Salisbury, Prodr. 70.—Willdenow, Spec. i, 709.—Enum. Suppl. 8.—B. S. Barton, Coll. i, 36, 56.—Nouveau Duhamel, i, 10.—Persoon, Syn. i, 151.—Desfontaines, Hist. Arb. ii, 362.—Titford, Hort. Bot. Am. 41.—Pursh, Fl. Am. Sept. i, 118.—Nuttall, Genera, i, 109.—Rømer & Schultes, Syst. iii, 491; Mant. 333.—De Candolle, Prodr. ii, 14.—Sprengel, Syst. i, 495.—Torrey in Ann. Lyc. N. York, ii, 173.—Don, Miller's Dict. ii, 17.—Hooker, Jour. Bot. i, 202.—Spach, Hist. Veg. ii, 430.—Lindley, Fl. Med. 393.—Dietrich, Syn. i, 555.—London, Arboretum, ii, 518, f. 186.—Eaton, Manual, 6 ed. 187.—Eaton & Wright, Bot. 282.—Griffith, Med. Bot. 433.—Browne, Trees of America, 169.—Guibourt, Hist. Drogues, 7 ed. iii, 544.

I. ligustrina, Jacquin, Coll. iv, 105; Icon. Rar. ii, 9, t. 310 [not Elliott].—Lamarck, Ill. i, 356.

I. Floridana, Lamarck, Ill. i, 356.

I. Cassena, Michaux, Fl. Bor.-Am. ii, 229.—Poirét, Suppl. iii, 65.—Rømer & Schultes, Syst. iii, 490.—Elliott, Sk. ii, 681.—Darby, Bot. S. States 426.—Wood, Cl. Book, 497.

I. religiosa, Barton, Fl. Virginica, 66.

Cassine ramulosa, Rafinesque, Fl. Ludoviciana, 363.

Hierophyllus Cassine, Rafinesque, Med. Bot. ii, 8.

Emetila ramulosa, Rafinesque, Sylva Telluriana, 45.

Ageria Cassena, Rafinesque, Sylva Telluriana, 47.

Ageria geminata, Rafinesque, Sylva Telluriana, 48.

CASSENA. YAUPON. YOPON.

Southern Virginia, southward, near the coast, to the Saint John's river and Cedar Keys, Florida, west along the Gulf coast to southern Arkansas, and the valley of the Colorado river, Texas.

A small tree, 6 to 8 meters in height, with a trunk 0.10 to 0.15 meter in diameter, or more often a shrub, sending up many slender stems and forming dense thickets; sandy, moist soil, along ponds and streams, reaching its greatest development in the river bottoms of eastern Texas.

Wood heavy, hard, close-grained, liable to check in drying; medullary rays numerous, conspicuous; color, nearly white, becoming yellow with exposure, the sap-wood lighter; specific gravity, 0.7270; ash, 0.87.

The leaves possess powerful emetic properties, and were employed by the southern Indians, together perhaps with those of *I. Dahoon*, in the preparation of their "black drink" (*Am. Jour. Pharm.* xlv, 217.—*U. S. Dispensatory*, 14 ed. 1670.—*Nat. Dispensatory*, 2 ed. 754).

36.—*Ilex decidua*, Walter,

Fl. Caroliniana, 241.—Poiret, Suppl. iii, 65.—Chapman, Fl. S. States, 269.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 59.—Lesquereux in Owen's 2d Rep. Arkansas, 373.—Wood, Cl. Book, 497; Bot. & Fl. 208.—Gray, Manual N. States, 5 ed. 306.—Young, Bot. Texas, 373.—Vasey, Cat. Forest Trees, 8.—Maximowicz in Mem. Acad. St. Petersburg, xxix, No. 3, 30.—Watson in Proc. Am. Acad. xvii, 335.

I. prinoides, Aiton, Hort. Kew. i, 169; 2 ed. i, 278.—Lamarck, Ill. i, 355.—Willdenow, Spec. i, 709.—Nouveau Duhamel, i, 11.—Michaux, Fl. Bor.-Am., ii, 229.—Persoon, Syn. i, 151.—Desfontaines, Hist. Arb. ii, 362.—Pursh, Fl. Am. Sept. i, 118.—Nuttall, Genera, i, 109.—Römer & Schultes, Syst. iii, 488; Mant. 332.—Watson, Dend. Brit. i, t. 15.—Sprengel, Syst. i, 495.—Audubon, Birds, t. 89.—Eaton, Manual, 6 ed. 187.—Eaton & Wright, Bot. 282.—Darby, Bot. S. States, 426.

I. æstivalis, Lamarck, Dict. iii, 147; Ill. i, 356.

Prinos deciduus, De Candolle, Prodr. ii, 16.—Don, Miller's Dict. ii, 20.—Hooker, Jour. Bot. i, 202.—London, Arboretum, ii, 520.

I. ambiguus, Elliott, Sk. ii, 705.

Southern Virginia, southward, through the middle districts, to western Florida, valley of the Mississippi river, southern Illinois southward to the Gulf of Mexico, and through southeastern Missouri, Arkansas, and eastern Texas to the valley of the Colorado river.

A small tree, 8 to 9 meters in height, with a trunk 0.15 to 0.20 meter in diameter, or in the Atlantic states a tall, straggling shrub; low, wet woods along streams, reaching its greatest development in the Iron Mountain region of Missouri and in southern Arkansas.

Wood heavy, hard, close-grained, compact; medullary rays numerous, thin; color, creamy-white, the sap-wood lighter; specific gravity, 0.7420; ash, 0.70.

CYRILLACEÆ.

37.—*Cyrilla racemiflora*, Linnæus,

Mant. i, 50; Syst. 14 ed. 241.—Jacquin, Icon. Rar. t. 47; Coll. i, 162.—Walter, Fl. Caroliniana, 103.—Lamarck, Dict. ii, 245; Ill. ii, 144, t. 147, f. 2.—Nouveau Duhamel, i, 215, t. 46.—Desfontaines, Hist. Arb. i, 255.—Elliott, Sk. i, 294.—Eaton, Manual, 6 ed. 119.—Eaton & Wright, Bot. 218.—Torrey & Gray, Fl. N. America, i, 256.—Nuttall, Sylva, ii, 96, t. 74; 2 ed. ii, 43, t. 74.—Planchon in Hooker's Jour. Bot. v, 254.—Schnizlein, Icon. t. 240, f. 1-4, 6, 17, 19, 21.—Darby, Bot. S. States, 417.—Cooper in Smithsonian Rep. 1858, 253.—Chapman, Fl. S. States, 272.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 105.—Porcher, Resources S. Forests, 130.—Maout & Decaisne, Bot. English ed. 540 & f.—Baillon, Adansonia, i, 203, t. 4.—Wood, Cl. Book, 493; Bot. & Fl. 205.—Vasey, Cat. Forest Trees, 18.

Andromeda plumata, Bartram, Cat.—Marshall, Arbustum, 9.

C. Caroliniana, Michaux, Fl. Bor.-Am. i, 158.—Gærtner, f. Fruct. Suppl. 147, t. 209, f. 8.—Persoon, Syn. i, 175.—Pursh, Fl. Am. Sept. i, 170.—Nuttall, Genera, i, 145.—Poiret, Suppl. ii, 436.—Römer & Schultes, Syst. v, 408.—Bot. Mag. t. 2456.—Walpers, Rep. vi, 421.—Dietrich, Syn. i, 805.

Itea Cyrilla, L'Heritier, Stirp. i, 137, t. 66.—Swartz, Prodr. 50; Fl. Ind. Occ. i, 506; Obs. 94, t. 4.—Willdenow, Spec. i, 1146.—Aiton, Hort. Kew. 2 ed. ii, 37.

C. racemosa, London, Arboretum, iv, 2577, f. 2503.

C. polystachia, *C. parvifolia*, *C. fuscata*, Rafinesque, Aulikon Botanikon, 8.

IRON WOOD.

North Carolina southward, near the coast, to middle Florida (latitude 30°), westward, along the Gulf coast, to the valley of the Pearl river, Mississippi.

A small tree, sometimes 8 to 10 meters in height, with a trunk 0.15 to 0.20 meter in diameter, or often a tall shrub, sending up many stems from the root; open swamps and low thickets; a variety (*Chapman, Curtiss*) with narrower, persistent leaves, and thicker spongy bark, in pond holes and wet depressions of the pine barrens of the Apalachicola region of western Florida, forms dense, impenetrable thickets.

Wood heavy, weak, hard, close-grained, compact; medullary rays thin, not conspicuous; color, brown tinged with red, the sap-wood a little lighter; specific gravity, 0.6784; ash, 0.42.

38.—*Cliftonia ligustrina*, Banks,

Ex. Gärtner f. Fruct. Suppl. 246, t. 225.—Bartram, Travels, 2 ed. 31.—Torrey & Gray, Fl. N. America, i, 256.—Nuttall, Sylva, ii, 92, t. 73; 2 ed. ii, 39, t. 73.—Planchon in Hooker's Jour. Bot. v, 255.—Walpers, Rep. vi, 422.—Dietrich, Syn. ii, 1412.—Schnizlein. Icon. t. 240^{xx}, f. 5, 7-10, 20.—Cooper in Smithsonian Rep. 1858, 251.—Chapman, Fl. S. States, 273.—Porcher, Resources S. Forests, 130.—Baillon in Adansonia, i, 202, t. 4, f. 3-6.—Vasey, Cat. Forest Trees, 18.

Mylocaryum ligustrinum, Willdenow, Enum. i, 454.—Bot. Mag. t. 1625.—Lamarck, Ill. iii, 616, t. 952, f. 1.—Pursh, Fl. Am. Sept. i, 302, t. 14.—Poiret, Suppl. iv, 41.—Elliott, Sk. i, 508.—Eaton, Manual, 6 ed. 231.—Eaton & Wright, Bot. 323.—Darby, Fl. S. States, 417.—Wood, Cl. Book, 493; Bot. & Fl. 205.

TITL. IRON WOOD. BUCKWHEAT TREE.

Valley of the Savannah river, Georgia, southward to the Chattahoochee region of west Florida, westward along the Gulf coast to the valley of the Pearl river, Louisiana.

A small tree, sometimes 12 meters in height, with a trunk 0.30 to 0.40 meter in diameter, or toward its southern limits in Florida reduced to a shrub; margins of pine-barren ponds and streams.

Wood heavy, soft, not strong, close-grained, compact; medullary rays numerous, thin; color, brown tinged with red, the sap-wood lighter; specific gravity, 0.6249; ash, 0.42; largely used as fuel, burning with a clear flame.

CELASTRACEÆ.

39.—*Euonymus atropurpureus*, Jacquin,

Hort. Vind. ii, 155, t. 120.—Lamarck, Dict. ii, 573; Ill. ii, 98.—Aiton, Hort. Kew. i, 274; 2 ed. ii, 29.—Willdenow, Spec. i, 1132; Enum. i, 256.—Michaux, Fl. Bor.-Am. i, 155.—Persoon, Syn. i, 243.—Nouveau Duhamel, iii, 26.—Desfontaines, Hist. Arb. ii, 356.—Pursh, Fl. Am. Sept. i, 168.—Turpin, Dict. Sci. Nat. xvii, 532, t. 272.—Eaton, Manual, 28; 6 ed. 140.—Nuttall, Genera, 155.—Römer & Schultes, Syst. v, 466.—Hayne, Dend. Fl. 24.—Elliott, Sk. i, 293.—De Candolle, Prodr. ii, 4.—Torrey in Ann. Lye. N. York, ii, 173; Fl. U. S. 261; Compend. Fl. N. States, 120; Fl. N. York, i, 141; Nicolle's Rep. 147.—Sprengel, Syst. i, 788.—Don, Miller's Dict. ii, 5.—Beck, Bot. 72.—Hooker, Jour. Bot. i, 201.—Spach, Hist. Veg. ii, 405.—Rafinesque, New Fl. 60.—London, Arboretum, ii, 499, f. 167.—Torrey & Gray, Fl. N. America, i, 257.—Dietrich, Syn. i, 819.—Eaton & Wright, Bot. 240.—Griffith, Med. Bot. 219, f. 112.—Gray, Genera, ii, 188; Manual N. States, 5 ed. 116.—Richardson, Arctic Exped. 423.—Parry in Owen's Rep. 610.—Darby, Bot. S. States, 268.—Darlington, Fl. Cestrica, 3 ed. 48.—Baillon in Bull. Soc. Bot. France, v, 314.—Chapman, Fl. S. States, 76.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 102.—Lesquereux in Owen's 2d Rep. Arkansas, 354.—Wood, Cl. Book, 289; Bot. & Fl. 76.—Porcher, Resources S. Forests, 129.—Engelmann in Trans. Am. Phil. Soc. new ser. xii, 187.—Koch, Dendrologie, i, 629.—Young, Bot. Texas, 205.—Vasey, Cat. Forest Trees, 9.

E. Carolinensis, Marshall, Arbustum, 43.

E. latifolius, Marshall, Arbustum, 44 [not Aiton].—Agardh, Theor. & Syst. Pl. t. 22, f. 4.

BURNING BUSH. WAHOO. SPINDLE TREE. ARROW WOOD.

Western New York, west to the valley of the upper Missouri river (*Fort Union*), Montana, southward to northern Florida, southern Arkansas, and eastern Kansas.

A small tree, rarely 6 to 8 meters in height, with a trunk 0.15 meter in diameter, or more often a shrub 2 to 3 meters in height; low, rich woods, reaching its greatest development west of the Mississippi river.

Wood heavy, very close-grained, liable to check badly in seasoning; medullary rays hardly discernible; color, white tinged with orange; specific gravity, 0.6592; ash, 0.58.

Wahoo bark, a mild but rather uncertain purgative, is used by herbalists in the form of decoctions, tinctures, fluid extracts, etc. (*Am. Jour. Pharmacy*, xx, 80.—*U. S. Dispensatory*, 14 ed. 402.—*Nat. Dispensatory*, 2 ed. 559).

40.—*Myginda pallens*, Smith,

Rees' Cycl. xxv, No. 4.—De Candolle, Prodr. ii, 13.—Dietrich, Syn. i, 554.—Grisebach, Fl. British West Indies, 146.—Chapman in Coulter's Bot. Gazette, iii, 3; Fl. S. States, Suppl. 612.

Semi-tropical Florida, Upper Metacombe Key; in the West Indies.

A small tree, rarely exceeding 4 meters in height, with a trunk 0.15 meter in diameter.

Wood very heavy, hard, very close-grained, compact, satiny; layers of annual growth and numerous medullary rays hardly distinguishable; color, dark brown or nearly black, the thick sap-wood lighter brown tinged with red; specific gravity, 0.9048; ash, 3.42.

41.—*Schæfferia frutescens*, Jacquin,

Stirp. Am. 259.—Gærtner f. Fruct. Suppl. 249, t. 225, f. 7.—Lamarck, Ill. iii, 402, t. 809.—Poiret in Lamarck, Dict. vi, 727.—De Candolle, Prodr. ii, 41.—Karsten, Fl. Columbiae, i, t. 91.—Chapman, Fl. S. States, 76.—Grisebach, Fl. British West Indies, 146.—Walpers, Ann. vii, 581.

S. completa, Swartz, Fl. Ind. Occ. i, 327, t. 7, f. A.—Willdenow, Spec. iv, 741.—Aiton, Hort. Kew. 2 ed. v, 371.—Macfadyen, Fl. Jamaica, 207.

S. buxifolia, Nuttall, Sylva, ii, 42, t. 56; 2 ed. i, 190, t. 56.—Cooper in Smithsonian Rep. 1858, 264.

YELLOW WOOD. BOX WOOD.

Semi-tropical Florida, southern keys from Metacombe Key eastward, Caloosa river and sparingly on the Reef Keys; in the West Indies.

A small tree, occasionally 10 meters in height, with a trunk 0.15 to 0.20 meter in diameter, generally hollow and defective.

Wood heavy, hard, close-grained, compact, susceptible of a high polish; medullary rays numerous, obscure; color, light bright yellow, the sap-wood a little lighter; specific gravity, 0.7745; ash, 2.54.

RHAMNACEÆ.

42.—*Reynosia latifolia*, Grisebach,

Cat. Pl. Cuba, 34.—Eggers, Videnskab, Medd. fra. Nat. For. 173 & t.; Bull. U. S. Nat. Mus. xiii, 40.—Gray in Coulter's Bot. Gazette, iv, 208.—Chapman, Fl. S. States, Suppl. 612.

? *Rhamnus lævigatus*, Vahl, Symbolæ, iii, 41.

Ceanothus lævigatus, De Candolle, Prodr. ii, 30.

Scutia ferrea, Chapman, Fl. S. States, 72 [not Brongniart].

? *Rhamindium revolutum*, Chapman, Fl. S. States, Suppl. 612.

RED IRON WOOD. DARLING PLUM.

Semi-tropical Florida, Miami (*Garber*), bay Biscayne, and on the southern keys (*Ourtiss*); in the West Indies.

A small tree, sometimes 8 meters in height, with a trunk 0.15 to 0.20 meter in diameter.

Wood heavy, exceedingly hard, strong, close-grained, compact; medullary rays numerous, thin; color, rich dark brown, the sap-wood light brown; specific gravity, 1.0715; ash, 3.20.

The edible fruit, ripening in April and May, of agreeable flavor.

43.—*Condalia ferrea*, Grisebach,

Fl. British West Indies, 100.—Walpers, Ann. vii, 588.—Gray in Coulter's Bot. Gazette, iv, 208.—Chapman, Fl. S. States, Suppl. 612.

Rhamnus ferrea, Vahl, Symbolæ, iii, 41, t. 58.

Zizyphus emarginatus, Swartz, Fl. Ind. Occ. iii, 1954.

Ceanothus ferreus, De Candolle, Prodr. ii, 30.

Scutia ferrea, Brongniart in Ann. Sci. Nat. 1 ser. x, 363 [not Chapman, Fl. S. States, 72].—Vasey, Cat. Forest Trees, 9.

BLACK IRON WOOD.

Semi-tropical Florida, cape Canaveral to bay Biscayne, on the southern keys; in the West Indies.

A small tree, sometimes 11 meters in height, with a trunk 0.25 to 0.38 meter in diameter, generally hollow and defective; common.

Wood exceedingly heavy and hard, strong, brittle, close-grained, compact, difficult to work; remarkable for the large percentage of ash; medullary rays very numerous, thin; color, rich orange-brown, the sap-wood lighter; specific gravity, 1.3020; ash, 8.31.

44.—*Condalia obovata*, Hooker,

Icon. t. 287.—Torrey & Gray, Fl. i, 685.—Gray in Jour. Boston Soc. Nat. Hist. vi, 169; Genera, ii, 172, t. 164; Smithsonian Contrib. iii, 32; v, 27; Hall's Pl. Texas, 5.—Torrey, Bot. Mex. Boundary Survey, 47.—Watson in Proc. Am. Acad. xvii, 336.

BLUE WOOD. LOGWOOD. PURPLE HAW.

Eastern and southwestern Texas, westward through southern New Mexico to southern Arizona; probably extending into northern Mexico.

A small tree, 6 to 10 meters in height, with a trunk 0.15 to 0.20 meter in diameter, or often a shrub; reaching its greatest development along the streams of eastern Texas; one of the common "chaparral" plants of western Texas, here forming dense, impenetrable thickets.

Wood very heavy, hard, close-grained, liable to check in seasoning, containing many groups of large irregularly-arranged open ducts; medullary rays numerous, obscure; color, light red, the sap-wood light yellow; specific gravity, 1.1999; ash, 7.03.

45.—*Rhamnus Caroliniana*, Walter,

Fl. Caroliniana, 101.—Lamarck, Ill. ii, 88; Dict. iv, 476.—Michaux, Fl. Bor.-Am. i, 153.—Nouveau Duhamel, iii, 47.—Persoon, Syn. i, 239.—Pursh, Fl. Am. Sept. i, 166.—Nuttall, Genera, i, 153.—Römer & Schultes, Syst. v, 285.—Elliott, Sk. i, 289.—De Candolle, Prodr. ii, 26.—Sprengel, Syst. i, 768.—Torrey in Ann. Lyc. N. York, ii, 174.—Don, Miller's Dict. ii, 32.—Hooker, Jour. Bot. i, 202.—Torrey & Gray, Fl. N. America, i, 262.—Dietrich, Syn. i, 807.—London, Arboretum, ii, 537.—Eaton, Manual, 6 ed. 300.—Eaton & Wright, Bot. 390.—Scheele in Römer, Texas, 432.—Nuttall, Sylva, ii, 50, t. 59; 2 ed. i, 193, t. 59.—Darby, Bot. S. States, 269.—Lesquereux in Owen's 2d Rep. Arkansas, 354.—Wood, Cl. Book, 219; Bot. & Fl. 77.—Koch, Dendrologie, i, 610.—Gray, Hall's Pl. Texas, 5.

? *Frangula fragillis*, Rafinesque, Fl. Ludoviciana, 320; Sylva Telluriana, 27.

Sarcophalus Carolinianus, Rafinesque, Sylva Telluriana, 29.

Frangula Caroliniana, Gray, Genera, ii, 178, t. 167; Manual N. States, 5 ed. 115.—Torrey, Bot. Mex. Boundary Survey, 46.—Cooper in Smithsonian Rep. 1858, 251.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 92.—Chapman, Fl. S. States, 73.—Vasey, Cat. Forest Trees, 9.

INDIAN CHERRY.

Long Island, New York, west along the valley of the Ohio river to southern Illinois, Missouri south of the Meramec river, eastern Kansas, and the Indian territory, south to northern Florida (latitude 30°), and through the Gulf states to western Texas.

A small tree, 6 to 10 meters in height, with a trunk 0.20 to 0.30 meter in diameter, or in the Atlantic states generally a tall shrub; rich woods along streams and river bottoms, reaching its greatest development in southern Arkansas and eastern Texas.

Wood light, hard, not strong, coarse-grained, compact; medullary rays numerous, thin; color, light brown, the sap-wood lighter; specific gravity, 0.5462; ash, 0.64.

The edible fruit sweet and agreeable.

46.—*Rhamnus Californica*, Eschscholtz,

Mem. Acad. St. Petersburg, x, 281 (*Linnaea* Litt.-Ber. 1823, 149.—Presl, Rep. Bot. i, 197).—Don, Miller's Dict. ii, 32.—Torrey & Gray, Fl. N. America, i, 263.—Dietrich, Syn. i, 806.—Eaton & Wright, Bot. 390.—Brewer & Watson, Bot. California, i, 101.—Hemsley, Bot. Am.-Cent. i, 197.

R. oleifolius, Hooker, Fl. Bor.-Am. i, 123, t. 44.—Hooker & Arnott, Bot. Beechey, 136, 328.—Torrey & Gray, Fl. N. America, i, 260.—Eaton & Wright, Bot. 390.—Bentham, Bot. Sulphur, 10; Pl. Hartweg. 302.—Durand in Jour. Philadelphia Acad. 1855, 85.—Carrière in Rev. Hort. xlvii, 354, f. 47-49.

Endotropis oleifolia, Rafinesque, Sylva Telluriana, 31.

R. laurifolius, Nuttall in Torrey & Gray, Fl. N. America, i, 260.—Eaton & Wright, Bot. 390.

Frangula Californica, Gray, Genera, ii, 178; Jour. Boston Soc. Nat. Hist. vi, 146.—Torrey in Sitgreaves' Rep. 157. Pacific R. R. Rep. iv, 74; Bot. Mex. Boundary Survey, 46; Bot. Wilkes Exped. 261.—Newberry in Pacific R. R. Rep. vi, 69.—Bolander in Proc. California Acad. iii, 78.

California, west of the Sierra Nevadas, from the valley of the upper Sacramento river southward to Santa Barbara and fort Tejon.

A small tree, rarely 7 to 9 meters in height, with a trunk 0.30 to 0.37 meter in diameter (*Pringle*), or commonly a shrub, along the sea-coast and at high elevations often prostrate; common and reaching its greatest development in the valleys of the Santa Cruz mountains. A low shrubby form, densely white-tomentose, especially on the under side of the leaves, of southern California, Arizona, and New Mexico, is—

var. *tomentella*, Brewer & Watson, Bot. California, i, 101.

R. tomentellus, Benth. Pl. Hartweg. 303.—Seemann, Bot. Herald, 275.—Walpers, Ann. ii, 267.

Frangula Californica, var. *tomentella*, Gray in Smithsonian Contrib. vi, 28.—Torrey in Pacific R. R. Rep. iv, 74; vii, 9.

Wood light, soft, rather coarse-grained, checking in drying; layers of annual growth marked by many rows of open ducts; medullary rays narrow, obscure; color, brown or light yellow, the sap-wood lighter; specific gravity, 0.6000; ash, 0.58.

47.—*Rhamnus Purshiana*, De Candolle,

Prodr. ii, 25.—London, Arboretum, ii, 538, f. 211.—Hooker, Fl. Bor.-Am. i, 123, t. 43; London Jour. Bot. vi, 78.—Don, Miller's Dict. ii, 32.—Torrey & Gray, Fl. N. America, i, 262.—Dietrich, Syn. i, 807.—Nuttall, Sylva, ii, 52; 2 ed. i, 200.—Richardson, Arctic Exped. 423.—Newberry in Pacific R. R. Rep. vi, 69.—Koch, Dendrologie, i, 610.—Gray in Proc. Am. Acad. viii, 379.—Brewer & Watson, Bot. California, i, 101.—Hall in Coulter's Bot. Gazette, ii, 86.

R. alnifolius, Pursh, Fl. Am. Sept. i, 166 [not L'Heritier].

Cardiolepis obtusa, Rafinesque, Sylva Telluriana, 28.

Frangula Purshiana, Cooper in Smithsonian Rep. 1858, 259; Pacific R. R. Rep. xii^a, 29, 57.—Vasey, Cat. Forest Trees, 9.—Torrey, Bot. Wilkes Exped. 262.

BEARBERRY. BEAR WOOD. SHITTIM WOOD.

Puget sound, east along the mountain ranges of northern Washington territory to the Bitter Root mountain, Idaho (Mullan pass, *Watson*), and the shores of Flathead lake, Montana (*Canby & Sargent*), southward through western Washington territory, Oregon, and California, west of the Sierra Nevada, to about latitude 40°.

A small tree, often 12 meters in height, with a trunk 0.30 to 0.45 meter in diameter; depressions and along the sides and bottoms of cañons in the coniferous forests, reaching its greatest development along the western slope of the Coast Range of southern Oregon.

Wood light, very hard, not strong, close-grained, compact, satiny; medullary rays numerous, thin; color, light brown tinged with yellow, the sap-wood somewhat lighter; specific gravity, 0.5672; ash, 0.67.

The bark, like that of other species of the genus, possesses powerful cathartic properties, and, under the name of *Cascara sagrada*, has recently been introduced by herbalists in the form of fluid extracts, tinctures, etc., immense quantities being gathered for this purpose in the Oregon forests (*Nat. Dispensatory*, 2 ed. 659).

48.—*Ceanothus thyrsiflorus*, Eschscholtz,

Mem. Acad. St. Petersburg, x, 285.—Hooker, Fl. Bor.-Am. i, 125.—Don, Miller's Dict. ii, 37.—Hooker & Arnott, Bot. Beechey, 136, 328.—Torrey & Gray, Fl. N. America, i, 266.—Dietrich, Syn. i, 813.—London, Arboretum, ii, 540.—Eaton & Wright, Bot. 185.—Lindley, Bot. Reg. xxx, t. 38.—Nuttall, Sylva, ii, 44, t. 57; 2 ed. i, 193, t. 57.—Benth. Bot. Sulphur, 10; Pl. Hartweg. 302.—Ann. Gand. 1847, t. 107.—Torrey in Pacific R. R. Rep. iv, 14; Bot. Mex. Boundary Survey, 45; Bot. Wilkes Exped. 263.—Newberry in Pacific R. R. Rep. vi, 69.—Cooper in Pacific R. R. Rep. xii^a, 57.—Bolander in Proc. California Acad. iii, 78.—Koch, Dendrologie, i, 621.—Watson in Proc. Am. Acad. x, 334.—Brewer & Watson, Bot. California, i, 102.—Vasey, Cat. Forest Trees, 9.

BLUE MYRTLE.

California Coast ranges, from Mendocino county south to the valley of the San Louis Rey river (*Pala, Parish Brothers*).

A small tree, 8 to 10 meters in height, with a trunk 0.10 to 0.15 meter in diameter, or toward the southern limits reduced to a low shrub; common and reaching its greatest development in the *Sequoia* forests near Santa Cruz.

Wood light, soft, close-grained, compact; medullary rays very obscure; color, light brown, the sap-wood darker; specific gravity, 0.5750; ash, 0.69.

The bark of the root may be expected to possess similar astringent properties to that of the shrubby *C. Americana*, used with advantage in cases of diarrhea and dysentery, and as a domestic remedy in throat troubles (*U. S. Dispensatory*, 14 ed. 1609.—*Nat. Dispensatory*, 2 ed. 373).

49.—*Colubrina reclinata*, Brongniart,

Ann. Sci. Nat. 1 ser. x, 369.—Richard, Fl. Cuba, 359.—Grisebach, Fl. British West Indies, 101.—Eggers in Bull. U. S. Nat. Mus. No. 13, 40.

Rhamnus ellipticus, Aiton, Hort. Kew. i, 265; 2 ed. ii, 17.—Willdenow, Spec. i, 1098.—Swartz, Prodr. 50; Fl. Ind. Occ. i, 497.

Zizyphus Dominicensis, Nouveau Duhamel, iii, 56.

Ceanothus reclinatus, L'Heritier, Sert. 6.—Römer & Schultes, Syst. v, 288.—De Candolle, Prodr. ii, 31.—Macfadyen, Fl. Jamaica, 211.

NAKED WOOD.

Semi-tropical Florida, Umbrella Key, on the north end of Key Largo, and sparingly on the small islands south of Elliott's Key; through the West Indies.

One of the largest trees of the region, deciduous, 12 to 18 meters in height, with a trunk 0.60 to 1.25 meter in diameter; reaching its greatest development within the United States on Umbrella Key, here forming a dense forest.

Wood heavy, hard, very strong, brittle, close-grained, compact, satiny, susceptible of a good polish, containing many small open ducts; medullary rays numerous, thin; color, dark brown tinged with yellow, the sap-wood light yellow; specific gravity, 0.8208; ash, 1.75.

"The trunk attains a size of over 1 meter and is most extraordinary. When 0.152 meter thick it becomes furrowed, and the furrows and ridges multiply and extend in all directions; trunks 0.75 to 1 meter in diameter appear like a mass of braided serpents. On small trunks the bark breaks up into flakes which curl up and drop off. Between the ridges where the bark persists the edges of dozens of papery layers may be seen" (*Curtiss in let*).

SAPINDACEÆ.

50.—*Æsculus glabra*, Willdenow,

Enum. 405.—Pursh, Fl. Am. Sept. i, 255.—Nuttall, Genera, i, 241.—De Candolle, Prodr. i, 597.—Torrey, Fl. U. S. 384; Compend. Fl. N. States, 164.—Guimpel, Otto & Hayne, Abb. Holz. 28, t. 24.—Hayne, Dend. Fl. 44.—Sprengel, Syst. ii, 166.—Don, Miller's Dict. i, 652.—Beck, Bot. 65.—London, Arboretum, i, 467, f. 133.—Torrey & Gray, Fl. N. America, i, 251.—Dietrich, Syn. ii, 1225.—Eaton & Wright, Bot. 115.—Walpers, Rep. i, 424.—Gray, Genera, ii, 207, t. 176, 177; Manual N. States, 5 ed. 118.—Cooper in Smithsonian Rep. 1858, 251.—Chapman, Fl. S. States, 79.—Wood, Cl. Book, 288; Bot. & Fl. 85.—Engelmann in Trans. Am. Phil. Soc. new ser. xii, 187.—Koch, Dendrologie, i, 508.—Vasey, Cat. Forest Trees, 9.—Ridgway in Proc. U. S. Nat. Mus. 1882, 61.

Æ. pallida, Willdenow, Enum. 406.—Nuttall, Genera, i, 242.—De Candolle, Prodr. i, 597.—Guimpel, Otto & Hayne, Abb. Holz. 29, t. 25.—Sprengel, Syst. ii, 166.—Don, Miller's Dict. i, 650.—Eaton, Manual, 6 ed. 6.—Lindley, Bot. Reg. xxiv, t. 51.—London, Arboretum, i, 468, f. 134.

Æ. echinata, Muhlenberg, Cat. 38.

Æ. Ohioensis, Michaux f. Hist. Arb. Am. iii, 242; N. American Sylva, 3 ed. ii, 156, t. 92.—Poiret, Suppl. iii, 593.—De Candolle, Prodr. i, 597.—Don, Miller's Dict. i, 652.—Eaton, Manual, 6 ed. 6.—Riddell, Syn. Fl. W. States, 34.—Lindley, Bot. Reg. xxiv, 51, t. 51.—Nuttall, Sylva, ii, 71; 2 ed. ii, 17.

? *Æ. carnea*, Guimpel, Otto & Hayne, Abb. Holz. 25, t. 22.—Hayne, Dend. Fl. 43.—Lindley, Bot. Reg. xiii, t. 1056.—Watson, Dend. Brit. ii, t. 121.—Don, Miller's Dict. i, 652.—Torrey & Gray, Fl. N. America, i, 253.—Walpers, Rep. i, 425.

Pavia glabra, Spach in Ann. Sci. Nat. 2 ser. ii, 54; Hist. Veg. iii, 23.

Pavia pallida, Spach in Ann. Sci. Nat. 2 ser. ii, 54; Hist. Veg. iii, 23.

? *Pavia carnea*, Spach in Ann. Sci. Nat. 2 ser. ii, 54; Hist. Veg. iii, 23.—Don in Sweet's Brit. Fl. Gard. 2 ser. t. 301.

? *Pavia Watsoniana*, Spach in Ann. Sci. Nat. 2 ser. ii, 54; Hist. Veg. iii, 23.—Torrey & Gray, Fl. N. America, i, 253.

? *Æ. Watsoniana*, Dietrich, Syn. ii, 1225.—Walpers, Rep. i, 425.

Æ. Hippocastanum, var. *Ohioensis*, London, Arboretum, i, 467.—Browne, Trees of America, 110.

Æ. Hippocastanum, var. *glabra*, London, Arboretum, i, 467.—Browne, Trees of America, 111.

Æ. Hippocastanum, var. *pallida*, London, Arboretum, i, 468.—Browne, Trees of America, 111.

OHIO BUCKEYE. FETID BUCKEYE.

Western slopes of the Alleghany mountains, Pennsylvania to northern Alabama, westward through southern Michigan (rare) to southern Iowa, eastern Kansas to about longitude 97° west, and the Indian territory.

A small tree, 8 to 15 meters in height, with a trunk 0.30 to 0.60 meter in diameter; rich soil along streams and river bottoms, reaching its greatest development in the high valleys of the southern Alleghany mountains.

Wood light, soft, not strong, close-grained, compact, difficult to split, often blemished by dark lines of decay; medullary rays obscure; color, white, the sap-wood darker; specific gravity, 0.4542; ash, 0.86; largely used in

common with that of the other species of the genus in the manufacture of woodenware, artificial limbs (for which the wood of *Æsculus* is preferred to that of all other American trees), paper-pulp, wooden hats, less commonly for the bearings of shafting and machinery, and occasionally manufactured into lumber.

The bark of the allied old world species *Æ. Hippocastanum* occasionally has been found efficacious as a substitute for cinchona bark in the treatment of intermittent fevers (*U. S. Dispensatory*, 14 ed. 1565.—*Nat. Dispensatory*, 2 ed. 712), and similar properties may be looked for in the bark of *Æ. glabra*.

51.—*Æsculus flava*, Aiton,

Hort. Kew. i, 494; 2 ed. ii, 335.—B. S. Barton, Coll. i, 13; Bot. Appx. 26, t. 15, f. 2.—Willdenow, Spec. ii, 226; Enum. i, 405; Berl. Baumz. 13.—Desfontaines, Hist. Arb. i, 385.—Pursh, Fl. Am. Sept. i, 255.—Nuttall, Genera, i, 242.—James in Long's Exped. i, 22.—Guimpel, Otto & Hayne, Abb. Holz. 27, t. 23.—Hayne, Dend. Fl. 44.—Elliott, Sk. i, 436.—Watson, Dend. Brit. ii, t. 163.—Loddiges, Bot. Cab. t. 1280.—Torrey & Gray, Fl. N. America, i, 252.—Dietrich, Syn. ii, 1225.—Eaton, Manual, 6 ed. 7.—Eaton & Wright, Bot. 116.—Walpers, Rep. i, 424.—Darby, Bot. S. States, 266.—Torrey in Pacific R. R. Rep. iv, 74.—Browne, Trees of America, 118.—Schnizlein, Icon. t. 230^{xx}, f. 3.—Cooper in Smithsonian Rep. 1858, 251.—Chapman, Fl. S. States, 80.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 48.—Lesquereux in Owen's 2d Rep. Arkansas, 354.—Wood, Cl. Book, 288; Bot. & Fl. 75.—Gray, Manual N. States, 5 ed. 118.—Vasey, Cat. Forest Trees, 9.

Æ. octandra, Marshall, Arbustum, 4.—Miller's Dict. No. 1.

Pavia flava, Mönch, Meth. 66.—De Candolle, Prodr. i, 598.—Don, Miller's Dict. i, 653.—Spach in Ann. Sci. Nat. 2 ser. ii, 55; Hist. Veg. iii, 25.—London, Arboretum, i, 471 & t.

Æ. lutea, Wangerheim in Schrift. Gesell. Nat. Fr. Berlin, viii, 133, t. 6.—Michaux, Fl. Bor.-Am. i, 219.—Persoon, Syn. i, 403.—Koch, Dendrologie, i, 509.

Pavia lutea, Poiret in Lamarck, Dict. v, 94.—Nouveau Duhamel, iii, 155, t. 38.—Michaux f. Hist. Arb. Am. iii, 237, t. 11; N. American Sylva, 3 ed. ii, 153, t. 91.

Æ. neglecta, Lindley, Bot. Reg. xii, t. 1009.

Pavia neglecta, Don, Miller's Dict. i, 653.—Spach in Ann. Sci. Nat. 2 ser. ii, 55; Hist. Veg. iii, 24.—London, Arboretum, i, 472.

SWEET BUCKEYE.

Allegheny county, Pennsylvania (*T. O. Porter*), southward along the Alleghany mountains to northern Georgia (Augusta) and Alabama, west along the valley of the Ohio river to southern Iowa, the Indian territory, and the valley of the Brazos river, eastern Texas.

A tree 18 to 28 meters in height, with a trunk 0.60 to 0.90 meter in diameter, or toward its southwestern limits reduced to a shrub; rich woods and along streams, reaching its greatest development on the slopes of the Alleghany mountains of North Carolina and Tennessee.

A variety with purple or flesh-colored flowers, the leaflets pubescent beneath, is—

var. *purpurascens*, Gray, Manual N. States, 5 ed. 118.

Æ. hybrida, De Candolle, Hort. Monsp. 1813, 75.—Poiret, Suppl. iv, 334.

Æ. discolor, Pursh, Fl. Am. Sept. i, 255.—Nuttall, Genera, i, 242.—Bot. Reg. iv, t. 310.—Elliott, Sk. i, 436.—Sprengel, Syst. ii, 167.—Sertum Botanicum, iv & t.—Eaton & Wright, Bot. 116.—Walpers, Ann. iv, 381.

Pavia discolor, Poiret, Suppl. v, 769.—Don, Miller's Dict. i, 653.—Eaton, Manual, 6 ed. 7.—Spach in Ann. Sci. Nat. 2 ser. ii, 57; Hist. Veg. iii, 28.—London, Arboretum, i, 472.

Pavia hybrida, De Candolle, Prodr. i, 598.—Don, Miller's Dict. i, 653.—Eaton, Manual, 6 ed. 6.—Spach in Ann. Sci. Nat. 2 ser. ii, 56; Hist. Veg. iii, 27.—London, Arboretum, i, 472.—Eaton & Wright, Bot. 116.—Koch, Dendrologie, i, 512.

Æ. Pavia, var. *discolor*, Torrey & Gray, Fl. N. America, i, 252.—Walpers, Rep. i, 424.—Gray in Jour. Boston Soc. Nat. Hist. vi, 167.

Wood light, soft, close-grained, compact, difficult to split; medullary rays numerous, obscure; color, creamy-white, the sap-wood hardly distinguishable; specific gravity, 0.4274; ash, 1.00.

52.—*Æsculus Californica*, Nuttall;

Torrey & Gray, Fl. N. America, i, 251; Sylva, ii, 69, t. 64; 2 ed. ii, 16, t. 64.—Hooker & Arnott, Bot. Beechey, 327.—Dietrich, Syn. ii, 1225.—Eaton & Wright, Bot. 116.—Walpers, Rep. i, 424.—Bentham, Bot. Sulphur, 9; Pl. Hartweg. 301.—Durand in Jour. Philadelphia Acad. 1855, 85.—Rev. Hort. iv, 150, f. 10, 11.—Torrey in Pacific R. R. Rep. iv, 74; Bot. Mex. Boundary Survey, 48; Bot. Wilkes Exped. 260.—Newberry in Pacific R. R. Rep. vi, 20, 69, f. 1.—Bot. Mag. t. 5077.—Fl. des Serres, xiii, 39, t. 1312.—London Gard. Chronicle, 1858, 844.—Belge, Hort. ix, 121 & t.—Gray in Proc. Boston Soc. Nat. Hist. vii, 146.—Bolander in Proc. California Acad. iii, 78.—Walpers, Ann. 624.—Koch, Dendrologie, i, 513.—Brewer & Watson, Bot. California, i, 106.—Vasey, Cat. Forest Trees, 9.

Calothyrsus Californica, Spach in Ann. Sci. Nat. 2 ser. ii, 62; Hist. Veg. iii, 35.

Pavia Californica, Hartweg in Jour. Hort. Soc. London, ii, 123.—Carrière in Rev. Hort. 1862, 369 & f.

CALIFORNIA BUCKEYE.

California, valley of the upper Sacramento river and Mendocino county, southward along the Coast ranges to San Luis Obispo, and along the western foot-hills of the Sierra Nevada to the San Bernardino mountains.

A low, widely-branching tree, 8 to 12 meters in height, with a short trunk 0.60 to 0.90 meter in diameter, often greatly expanded at the base, or more often a much-branched shrub 3 to 5 meters in height; borders of streams, reaching its greatest development in the cañons of the Coast Range, north of San Francisco bay.

Wood light, soft, not strong, very close-grained, compact; medullary rays numerous, obscure; color, white slightly tinged with yellow, the sap-wood hardly distinguishable; specific gravity, 0.4980; ash, 0.70.

53.—*Ungnadia speciosa*, Endlicher,

Atacta Bot. t. 36; Nov. Stirp. Desc. ix, 75.—Torrey & Gray, Fl. N. America, i, 684; Pacific R. R. Rep. ii, 162.—Walpers, Rep. i, 423; v, 371; Ann. vii, 625.—Gray in Jour. Boston Soc. Nat. Hist. vi, 167; Genera, ii, 211, t. 178, 179; Smithsonian Contrib. iii, 38; v, 30; Mem. Am. Acad. new ser. v, 299; Hall's Pl. Texas, 5.—Fl. des Serres, x, 217, t. 1059.—Torrey, Bot. Mex. Boundary Survey, 48.—Schnitzlein, Icon. t. 230, f. 2, 8.—Cooper in Smithsonian Rep. 1858, 265.—Koch, Dendrologie, i, 515.—Baillon, Hist. Pl. v, 423.—Vasey, Cat. Forest Trees, 9.—Watson in Proc. Am. Acad. xvii, 337.

U. heterophylla, Scheele in Linnæa, xxi, 589; Rømer, Texas, 589.

U. heptaphylla, Scheele in Linnæa, xxii, 352; Rømer, Texas, 432.

SPANISH BUCKEYE.

Valley of the Trinity river (Dallas, *Reverchon*) through western Texas to the cañons of the Organ mountains, New Mexico (*Bigelow*); southward into Mexico.

A small tree, sometimes 6 to 8 meters in height, with a trunk 0.15 to 0.20 meter in diameter, or toward its eastern and western limits reduced to a low shrub; common west of the Colorado river; bottoms and rich hillsides, reaching its greatest development in the valley of the Guadalupe river, between New Braunfels and the coast.

Wood heavy, soft, not strong, close-grained, compact, satiny, containing numerous evenly-distributed open ducts; medullary rays numerous, inconspicuous; color, red tinged with brown, the sap-wood lighter; specific gravity, 0.6332; ash, 1.17.

Fruit reputed poisonous.

54.—*Sapindus marginatus*, Willdenow,

Enum. i, 432.—Muhlenberg, Cat. 41.—De Candolle, Prodr. i, 607.—Sprengel, Syst. ii, 250.—Don, Miller's Dict. i, 665.—Spach, Hist. Veg. iii, 54.—Torrey & Gray, Fl. N. America, i, 255, 685; Pacific R. R. Rep. ii, 162.—Eaton, Manual, 6 ed. 323.—Eaton & Wright, Bot. 411.—Nuttall, Sylva, ii, 72, t. 65; 2 ed. ii, 19, t. 65.—Leavenworth in Am. Jour. Sci. i, 49, 130.—Engelmann & Gray in Jour. Boston Soc. Nat. Hist. v, 241.—Gray in Jour. Boston Soc. Nat. Hist. vi, 169; Genera, ii, 214, t. 180; Smithsonian Contrib. iii, 38; Hall's Pl. Texas, 5.—Engelmann in Wislizenus' Rep. 12.—Torrey in Emory's Rep. 138; Marcy's Rep. 282; Pacific R. R. Rep. iv, 2, 74; Bot. Mex. Boundary Survey, 47.—Scheele in Rømer, Texas, 433.—Schnitzlein, Icon. t. 230, f. 22.—Chapman, Fl. S. States, 79.—Lesquereux in Owen's 2d Rep. Arkansas, 354.—Wood, Cl. Book, 288; Bot. & Fl. 75.—Porcher, Resources S. Forests, 85.—Young, Bot. Texas, 208.—Vasey, Cat. Forest Trees, 9.—Hemsley, Bot. Am.-Cent. i, 214.—Watson in Proc. Am. Acad. xvii, 337.

S. saponaria, Lamarck, Ill. ii, 441, t. 307 [not Linnæus].—Michaux, Fl. Bor.-Am. i, 242.—Poiret in Lamarck, Dict. vi, 663, in part.—Persoon, Syn. i, 444.—Pursh, Fl. Am. Sept. i, 274.—Nuttall, Genera, i, 257.—Elliott, Sk. i, 460.—Torrey in Ann. Lyc. N. York, ii, 172.—Darby, Bot. S. States, 267.

? *S. inæqualis*, De Candolle, Prodr. i, 608.

S. falcatus, Rafinesque, Med. Bot. ii, 261.

S. acuminata, Rafinesque, New Fl. 22.

S. Drummondii, Hooker & Arnott, Bot. Beechey, 281 (excl. var.).—Walpers, Rep. i, 417.

WILD CHINA. SOAPBERRY.

Atlantic coast, Savannah river to the Saint John's river, Florida, and on Cedar Keys; southern Arkansas, valley of the Washita river (Prescott, *Letterman*) through western Louisiana and Texas to the mountain valleys of southern New Mexico and Arizona; southward into Mexico, and in the West Indies (? *S. inæqualis*).

A tree, sometimes 15 to 18 meters in height, with a trunk rarely 0.60 meter in diameter; west of the Colorado river much smaller, rarely 9 meters in height; along streams or toward the western limits of its distribution only in mountain valleys, reaching its greatest development along the river bottoms of eastern Texas.

Wood heavy, strong, hard, close-grained, compact, easily split into thin strips; layers of annual growth clearly marked by several rows of large open ducts; medullary rays thin, obscure; color, light brown tinged with yellow, the sap-wood lighter; specific gravity, 0.8126; ash, 1.50; largely used in Texas in the manufacture of cotton-baskets, and in New Mexico for the frames of pack-saddles.

Saponin, common in several species of the genus, and affording a substitute for soap, may be looked for in the fruit and roots of this tree.

55.—*Sapindus Saponaria*, Linnæus,

Spec. 1 ed. 367; Swartz, Obs. 152.—Lamarck, Ill. ii, 441, t. 307.—Willdenow, Spec. ii, 468.—Aiton, Hort. Kew, 2 ed. ii, 424.—Tittford, Hort. Bot. Am. 61.—Poiret in Lamarck, Dict. vi, 663.—Desconrtitz, Fl. Med. Antilles, iv, 121, t. 261.—De Candolle, Prodr. i, 607.—Spach, Hist. Veg. iii, 53.—Eaton, Manual, 6 ed. 323.—Macfadyen, Fl. Jamaica, 159.—Rafinesque, New Fl. 22.—Nuttall, Sylva, ii, 72; 2 ed. 20.—Richard, Fl. Cuba, 280.—Grisebach, Fl. British West Indies, 126.—Baillon, Hist. Pl. v, 349, f. 353.—Vasey, Cat. Forest Trees, 10.—Chapman in Coulter's Bot. Gazette, iii, 3; Fl. S. States, Suppl. 613.

SOAPBERRY.

Semi-tropical Florida, bay Biscayne, cape Sable, Caximbas bay, Thousand Islands, Key Largo, Elliott's Key; in the West Indies.

A small tree, 6 to 10 meters in height, with a trunk sometimes 0.38 meter in diameter; common on cape Sable, and reaching its greatest development within the United States on the Thousand Islands and along the shores of Caximbas bay.

Wood heavy, rather hard, close-grained, compact; medullary rays numerous, thin; color, light brown tinged with yellow, the sap-wood yellow; specific gravity, 0.8367; ash, 4.34.

The fruit and roots rich in *saponin* and used in the West Indies as a substitute for soap (*Guibourt, Hist. Drogues*, 7 ed. iii, 598.—*U. S. Dispensatory*, 14 ed. 1751); the round, black seeds for beads, buttons, and small ornaments.

56.—*Hypelate paniculata*, Cambessedes,

Mem. Mus. xviii, 32.—Don, Miller's Dict. i, 671.—Richard, Fl. Cuba, 295.—Grisebach, Fl. British West Indies, 127.—Chapman, Fl. S. States, 79.—Vasey, Cat. Forest Trees, 10.

Melicocca paniculata, Jussieu in Mem. Mus. iii, 187, t. 5.—De Candolle, Prodr. i, 615.—Nuttall, Sylva, ii, 74, t. 66; 2 ed. ii, 21, t. 66.

Exothea oblongifolia, Macfadyen, Fl. Jamaica, 232.

H. oblongifolia, Hooker in London Jour. Bot. iii, 226, t. 7.

INK WOOD. IRON WOOD.

Semi-tropical Florida, east coast, Mosquito inlet to the southern keys; in the West Indies.

A tree often 12 meters in height, with a trunk 0.45 meter in diameter.

Wood very heavy, exceedingly hard, very strong, close-grained, susceptible of a good polish, checking in drying; medullary rays obscure; color, bright reddish-brown, the sap-wood lighter; specific gravity, 0.9533; ash, 1.25; used in ship-building, for the handles of tools, and piles; resisting the attacks of the teredo.

57.—*Hypelate trifoliata*, Swartz,

Fl. Ind. Occ. ii, 655, t. 14.—Delessert, Icon. iii, t. 39.—De Candolle, Prodr. i, 614.—Chapman, Fl. S. States, 78.—Grisebach, Fl. British West Indies, 127; Cat. Pl. Cuba, 46.

WHITE IRON WOOD.

Semi-tropical Florida, Upper Metacombe and Umbrella Keys; in the West Indies.

A tree sometimes 12 meters in height, with a trunk 0.45 to 0.60 meter in diameter.

Wood very heavy, hard, close-grained, compact, susceptible of a fine polish, durable in contact with the soil; medullary rays thin, obscure; color, rich light brown, the sap-wood darker; specific gravity, 0.9102; ash, 1.38; used in ship-building, for the handles of tools, posts, etc.

58.—*Acer Pennsylvanicum*, Linnæus,

Spec. 1 ed. 1055.—Aiton, Hort. Kew. iii, 435.—Michaux, Fl. Bor.-Am. ii, 252.—Willdenow, Spec. iv, 989; Enum. i, 1045.—Desfontaines, Hist. Arb. i, 391.—Nouveau Duhamel, iv, 32.—Trattinick, Archiv. i, t. 11.—Hayne, Dend. Fl. 210.—Elliott, Sk. i, 451.—Torrey, Fl. U. S. 397; Compend. Fl. N. States, 170; Fl. N. York, i, 135.—Sprengel, Syst. ii, 224.—Eaton, Manual, 6 ed. 2.—Torrey & Gray, Fl. N. America, i, 246.—Hooker, Fl. Bor.-Am. i, 111.—Emerson, Trees Massachusetts, 496; 2 ed. ii, 566 & t.—Gray, Genera, ii, 200, t. 174, f. 1-3; Manual N. States, 5 ed. 119.—Richardson, Arctic Exped. 422.—Darby, Bot. S. States, 265.—Cooper in Smithsonian Rep. 1858, 251.—Chapman, Fl. S. States, 80.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 52.—Buchenau in Bot. Zeit. xix, 285, t. 2, f. 24.—Wood, Cl. Book, 286; Bot. & Fl. 74.—Koch, Dendrologie, i, 521.—Baillon, Hist. Pl. v, 373, f. 418-420.—Vasey, Cat. Forest Trees, 10.—Sears in Bull. Essex Inst. xiii, 175.—Bell in Geological Rep. Canada, 1879-'80, 53c.

A. Canadense, Marshall, Arbustum, 4.

A. striatum. Du Roi, Diss. 58; Harbk. i, 8, t. 1.—Wangenheim, Amer. 29, t. 12, f. 2.—Lamarck, Dict. ii, 381.—Ehrhart, Beitr. iv, 25.—Mœnch, Meth. 56.—Persoon, Syn. i, 417.—Michaux f. Hist. Arb. Am. ii, 242, t. 17; N. American Sylva, 3 ed. ii, 175, t. 47.—Pursh, Fl. Am. Sept. i, 267.—Nuttall, Genera, i, 253.—De Candolle, Prodr. i, 593.—Watson, Dend. Brit. i, t. 70.—Don, Miller's Dict. i, 648.—Beck, Bot. 64.—London, Arboretum, i, 407 & t.—Spach, Hist. Veg. iii, 85; Ann. Sci. Nat. 2 ser. ii, 162.—Dietrich, Syn. 1281.—Eaton & Wright, Bot. 112.—Bigelow, Fl. Boston. 3 ed. 407.—Browne, Trees of America, 76.

STRIPED MAPLE. MOOSE WOOD. STRIPED DOGWOOD. GOOSE-FOOT MAPLE. WHISTLE WOOD.

Valley of the Saint Lawrence river (Ha-Ha bay), northern shores of lake Ontario, islands of lake Huron, south through the northern Atlantic states, and along the Alleghany mountains to northern Georgia, west through the lake region to northeastern Minnesota.

A small tree, 6 to 10 meters in height, with a trunk 0.15 to 0.20 meter in diameter; cool ravines and mountain sides.

Wood light, soft, close-grained, compact, satiny; medullary rays numerous, thin; color, light brown, the sap-wood lighter; specific gravity, 0.5299; ash, 0.36.

59.—*Acer spicatum*, Lamarck,

Dict. ii, 381.—Aiton, Hort. Kew. iii, 485.—Persoon, Syn. i, 417.—De Candolle, Prodr. i, 593.—Don, Miller's Dict. i, 648.—Audubon, Birds, t. 134.—Penn. Cycl. i, 77.—Eaton, Manual, 6 ed. 2.—Beck, Bot. 64.—Spach, Hist. Veg. 87; Ann. Sci. Nat. 2 ser. ii, 163.—London, Arboretum, i, 406, t. 26.—Torrey & Gray, Fl. N. America, i, 246.—Dietrich, Syn. ii, 1281.—Eaton & Wright, Bot. 112.—Torrey, Fl. N. York, i, 185.—Browne, Trees of America, 74.—Emerson, Trees Massachusetts, 497; 2 ed. ii, 567 & t.—Parry in Owen's Rep. 610.—Richardson, Arctic Exped. 422.—Chapman, Fl. S. States, 80.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 52.—Wood, Cl. Book, 287; Bot. & Fl. 74.—Gray, Manual N. States, 5 ed. 119.—Koch, Dendrologie, i, 522.—Macoun in Geological Rep. Canada, 1875-'76, 192.—Sears in Bull. Essex Inst. xiii, 175.—Bell in Geological Rep. Canada, 1879-'80, 54c.—Nicholson in London Gard. Chronicle, 1881, 172.

A. Pennsylvanicum, Du Roi, Diss. 61; Harbk. i, 22, t. 1 [not Linnæus].—Wangenheim, Amer. 32, t. 12, f. 30.—Marshall, Arbustum, 2.

A. parviflorum, Ehrhart, Beitr. iv, 25; vi, 40.—Mœnch, Meth. 56.

A. montanum, Aiton, Hort. Kew. iii, 435; 2 ed. v, 447 (excl. syn. *striatum*).—Michaux, Fl. Bor.-Am. ii, 253.—Willdenow, Spec. iv, 988; Enum. i, 1045.—Desfontaines, Hist. Arb. i, 391.—Nouveau Duhamel, iv, 33.—Trattinick, Archiv. i, t. 13.—Pursh, Fl. Am. Sept. i, 267.—Nuttall, Genera, i, 253.—Guimpel, Otto & Hayne, Abb. Holz. 59, t. 48.—Hayne, Dend. Fl. 213.—Elliott, Sk. i, 452.—Torrey, Fl. U. S. 398; Compend. Fl. N. States, 170.—Sprengel, Syst. ii, 224.—Hooker, Fl. Bor.-Am. i, 111.—Bigelow, Fl. Boston. 3 ed. 408.—Darby, Bot. S. States, 265.

MOUNTAIN MAPLE.

Valley of the Saint Lawrence river, west along the northern shores of the great lakes to northern Minnesota and the Saskatchewan region, south through the northern states, and along the Alleghany mountains to northern Georgia.

A small tree, sometimes 8 to 10 meters in height, with a trunk 0.15 to 0.20 meter in diameter, or often a tall shrub; cool woods and mountain ravines, reaching its greatest development on the western slopes of the Alleghany mountains of North Carolina and Tennessee.

Wood light, soft, close-grained, compact; medullary rays inconspicuous; color, light brown tinged with red, the sap-wood lighter; specific gravity, 0.5330; ash, 0.43.

60.—*Acer macrophyllum*, Pursh,

Fl. Am. Sept. i, 267.—Poiret, Suppl. v, 669.—Nuttall, Genera, i, 253; Sylva, ii, 77, t. 67; 2 ed. ii, 24, t. 67.—De Candolle, Prodr. i, 594.—Sprengel, Syst. ii, 225.—Penn. Cycl. i, 78.—Eaton, Manual, 6 ed. 2.—Hooker, Fl. Bor.-Am. i, 112, t. 38.—Don, Miller's Dict. i, 648.—Spach in Ann. Sci. Nat. 2 ser. ii, 165.—Torrey & Gray, Fl. N. America, i, 246.—Hooker & Arnott, Bot. Beechey, 327.—Dietrich, Syn. ii, 1281.—London, Arboretum, i, 408, t. 28, f. 117, 118.—Eaton & Wright, Bot. 112.—Bentham, Fl. Hartweg. 301.—Browne, Trees of America, 78.—Richardson, Arctic Exped. 423.—Durand in Jour. Philadelphia Acad. 1855, 84.—Torrey in Pacific R. R. Rep. iv, 74; Bot. Mex. Boundary Survey, 47; Bot. Wilkes Exped. 258.—Newberry in Pacific R. R. Rep. vi, 21, 67.—Cooper in Pacific R. R. Rep. xii, 28, 57; Smithsonian Rep. 1858, 258.—Lyll in Jour. Linnæan Soc. vii, 134, 144.—Bolander in Proc. California Acad. iii, 78.—Wood, Cl. Book, 287; Bot. & Fl. 74.—Rothrock in Smithsonian Rep. 1867, 334.—Koch, Dendrologie, i, 528.—Gray in Proc. Am. Acad. viii, 379.—Brewer & Watson, Bot. California, i, 107.—Vasey, Cat. Forest Trees, 10.—Macoun in Geological Rep. Canada, 1875-76, 192.—G. M. Dawson in Canadian Nat. new ser. ix, 330.—Nicholson in London Gard. Chronicle, 1881, 10.

A. palmatum, Rafinesque, New Fl. & Bot. i, 48 [not Thunberg].

BROAD-LEAVED MAPLE.

Coast of Alaska, from latitude 55° south along the islands and coast of British Columbia, through western Washington territory and Oregon, and along the California Coast ranges and western slopes of the Sierra Nevada to the San Bernardino mountains and Hot Spring valley, San Diego county (*Parish Brothers*), not ascending above 4,000 feet altitude.

A tree 24 to 30 meters in height, with a trunk 1.20 to 1.50 meter in diameter; along streams and river bottoms, reaching its greatest development on the rich bottom lands of the Coquille and other rivers of southern Oregon, where, with the California laurel, it forms dense, heavy forests.

Wood light, soft, not strong, close-grained, compact, easily worked, susceptible of a good polish; medullary rays numerous, thin; color, rich light brown tinged with red, the sap-wood lighter, often nearly white; specific gravity, 0.4909; ash, 0.54; largely used in Oregon in the manufacture of furniture, for ax and broom handles, frames of snow-shoes, etc.; specimens with the grain beautifully curled and contorted are common and valued in cabinet-making.

61.—*Acer circinatum*, Pursh,

Fl. Am. Sept. i, 266.—Poiret, Suppl. v, 669.—Nuttall, Genera, i, 253; Jour. Philadelphia Acad. vii, 16 (excl. syn.); Sylva, ii, 80, t. 67; 2 ed. ii, 27, t. 67.—De Candolle, Prodr. i, 595.—Sprengel, Syst. ii, 225.—Penn. Cycl. i, 79.—Eaton, Manual, 6 ed. 2.—Don, Miller's Dict. i, 651.—Spach in Ann. Sci. Nat. 2 ser. ii, 166; Hist. Veg. iii, 97.—London, Arboretum, i, 422, f. 112, 127.—Torrey & Gray, Fl. N. America, i, 247.—Hooker, Fl. Bor.-Am. i, 112, t. 39.—Eaton & Wright, Bot. 112.—Dietrich, Syn. ii, 1282.—Browne, Trees of America, 91.—Richardson, Arctic Exped. 422.—Lindley in Paxton's Fl. Gard. ii, 156, f. 210 (London Gard. Chronicle, 1851, 791, f. 211).—Newberry in Pacific R. R. Rep. vi, 21, 69.—Cooper in Pacific R. R. Rep. xii, 28, 57; Smithsonian Rep. 1858, 258.—Lyll in Jour. Linnæan Soc. vii, 134.—Gray in Proc. Am. Acad. viii, 379.—Wood, Cl. Book, 287, Bot. & Fl. 74.—Koch, Dendrologie, i, 523.—Torrey, Bot. Wilkes Exped. 258.—Brewer & Watson, Bot. California, i, 107.—Vasey, Cat. Forest Trees, 10.—Hall in Coulter's Bot. Gazette, ii, 85.—Macoun in Geological Rep. Canada, 1875-76, 192.—G. M. Dawson, Canadian Nat. new ser. ix, 330.—Nicholson in London Gard. Chronicle, 1881, 10.

A. virgatum, Rafinesque, New Fl. & Bot. i, 48.

VINE MAPLE.

British Columbia, valley of the Fraser river (Yale) and probably farther north, southward through Washington territory and Oregon, west of the Cascade mountains to the Mount Shasta region of northern California, rarely ascending to 4,000 feet altitude.

A small tree, sometimes 8 to 12 meters in height, with a trunk 0.20 to 0.30 meter in diameter; along streams; the stems often prostrate and forming dense, impenetrable thickets.

Wood heavy, hard, not strong, close-grained, compact; medullary rays numerous, thin; color, light brown or often nearly white, the sap-wood lighter; specific gravity, 0.6660; ash, 0.39; used as fuel; by lumbermen for ax and shovel handles, and by the coast Indians for the bows of fishing nets.

62.—*Acer glabrum*, Torrey,

Ann. Lyc. N. York, ii, 172; Bot. Wilkes Exped. 259.—Don, Miller's Dict. i, 650.—Eaton, Manual, 6 ed. 2.—Torrey & Gray, Fl. N. America, i, 247, 684.—Eaton & Wright, Bot. 112.—Walpers, Rep. i, 409.—Nuttall, Sylva, ii, 86; 2 ed., ii, 33.—Newberry in Pacific R. R. Rep. vi, 69.—Cooper in Smithsonian Rep. 1858, 258; Pacific R. R. Rep. xii, 51, 57; Am. Nat. iii, 406.—Engelmann in Trans. Am. Phil. Soc. new ser. xii, 187.—Gray in Am. Jour. Sci. 2 ser. xxxiv, 259; Proc. Philadelphia Acad. 1863, 59.—Porter in Hayden's Rep. 1870, 474; 1871, 480.—Watson in King's Rep. v, 52.—Porter & Coulter, Fl. Colorado; Hayden's Surv. Misc. Pub. No. 4, 19.—Coulter in Hayden's Rep. 1872, 763.—Macoun in Geological Rep. Canada, 1875-76, 192.—Brewer & Watson, Bot. California, i, 107.—Rothrock in Wheeler's Rep. vi, 83.—Nicholson in London Gard. Chronicle, 1881, 750.

A. barbatum, Douglas in Hooker, Fl. Bor.-Am. i, 113.—London, Arboretum, i, 420, f. 125 (excl. syn.).

A. Douglasii, Hooker in London Jour. Bot. vi, 77, t. 6.

A. tripartitum, Nuttall in Torrey & Gray, Fl. N. America, i, 247.—Dietrich, Syn. ii, 1281.—Eaton & Wright, Bot. 112.—Walpers, Rep. i, 409.—Nuttall, Sylva, ii, 85, t. 71; 2 ed. ii, 33, t. 71.—Gray in Mem. Am. Acad. new ser. iv¹, 28; Pacific R. R. Rep. iv, 73.—Newberry in Pacific R. R. Rep. vi, 69.

DWARF MAPLE.

British Columbia, valley of the Fraser river and probably farther north, south through Washington territory, Oregon, and along the Sierra Nevada of California to the Yosemite valley; east along the mountain ranges of Idaho and Montana to the eastern base of the Rocky mountains, south through Colorado and Utah, in the east Humboldt Range, Nevada, and in the mountain ranges of western New Mexico and eastern Arizona.

A small tree, 8 to 12 meters in height, with a trunk sometimes 0.30 meter in diameter, or more often reduced to a low shrub 1 to 2 meters in height; borders of streams, reaching its greatest development in the mountain cañons of western New Mexico and eastern Arizona.

Wood heavy, hard, close-grained, compact; medullary rays numerous, thin; color, light brown, or often nearly white, the sap-wood lighter; specific gravity, 0.6028; ash, 0.30.

63.—*Acer grandidentatum*, Nuttall;

Torrey & Gray, Fl. N. America, i, 247.—Dietrich, Syn. ii, 1283.—Eaton & Wright, Bot. 112.—Walpers, Rep. i, 409.—Nuttall, Sylva, ii, 82, t. 69; 2 ed. ii, 29, t. 69.—Watson in King's Rep. v, 52; Pl. Wheeler, 7.—Porter in Hayden's Rep. 1871, 480.—Vasey, Cat. Forest Trees, 10.—Parry in Am. Nat. ix, 201, 268.—Rothrock in Wheeler's Rep. vi, 83.—Rusby in Bull. Torrey Bot. Club, ix, 106.—Watson in Proc. Am. Acad. xvii, 338.—Nicholson in London Gard. Chronicle, 1881, 172.

Western Montana, headwaters of the Columbia river (*Nuttall*), cañons of the Wahsatch mountains, Utah, and south through eastern Arizona to southwestern New Mexico (Mogollon mountains, *E. L. Greene*), and reported in the ranges east of the Rio Grande; southward into Coahuila (*Palmer*).

A small tree, rarely exceeding 10 meters in height, with a trunk 0.20 to 0.25 meter in diameter; along streams; not common.

Wood heavy, hard, close-grained, compact; medullary rays numerous, thin, distinct; color, light brown, or often nearly white; specific gravity, 0.6902; ash, 0.64.

64.—*Acer saccharinum*, Wangenheim,

Amer. 36, t. 11, f. 26.—Lamarck, Dict. ii, 379.—Walter, Fl. Caroliniana, 251.—Aiton, Hort. Kew. iii, 434; 2 ed. v, 447.—Ehrhart, Beitr. iv, 24.—Persoon, Syn. i, 417.—Nouveau Duhamel, iv, 29, t. 8.—Willdenow, Spec. iv, 985; Enum. ii, 1044.—Desfontaines, Hist. Arb. i, 392.—Trattinick, Archiv. i, t. 3.—Michaux f. Hist. Arb. Am. ii, 218, t. 15; N. American Sylva, 3 ed. i, 153, t. 42.—Titford, Hort. Bot. Am. 105.—Pursh, Fl. Am. Sept. i, 266.—Eaton, Manual, 44; 6 ed. 2.—Nuttall, Genera, i, 253.—Hayne, Dend. Fl. 214.—Elliott, Sk. i, 450.—Richardson, Franklin Jour. 26; Arctic Exped. 422.—De Candolle, Prodr. i, 595.—Torrey, Fl. U. S. 396; Compend. Fl. N. States, 170; Fl. N. York, i, 135.—Sprengel, Syst. ii, 225.—Penn. Cycl. i, 79.—Hooker, Fl. Bor.-Am. i, 113.—Don, Miller's Dict. i, 650.—Beck, Bot. 63.—Bigelow, Fl. Boston. 3 ed. 406.—Spach, Hist. Veg. iii, 170; Ann. Sci. Nat. 2 ser. ii, 99.—London, Arboretum, i, 411, t. 31, f. 122.—Torrey & Gray, Fl. N. America, i, 248.—Eaton & Wright, Bot. 112.—Dietrich, Syn. ii, 1282.—Walpers, Rep. i, 410.—Nees, Pl. Med. 5.—Nuttall, Sylva, ii, 88; 2 ed. ii, 35.—Browne, Trees of America, 83.—Emerson, Trees Massachusetts, 489; 2 ed. ii, 258 & t.—Gray, Genera, ii, 200, t. 174; Manual N. States, 5 ed. 119.—Darlington, Fl. Cestrica, 3 ed. 45.—Darby, Bot. S. States, 265.—Parry in Owen's Rep. 610.—Chapman, Fl. S. States, 80.—Lesquereux in Owen's 2d Rep. Arkansas, 354.—Wood, Cl. Book, 286; Bot. & Fl. 74.—Porcher, Resources S. Forests, 80.—Engelmann in Trans. Am. Phil. Soc. new ser. xii, 187.—Young, Bot. Texas, 206.—Vasey, Cat. Forest Trees, 10.—Guibourt, Hist. Drogues, 7 ed. iii, 606.—Ward in Bull. U. S. Nat. Mus. No. 22, 73.—Sears in Bull. Essex Inst. xiii, 175.—Bell in Geological Rep. Canada, 1879-'80, 51^c.—Ridgway in Proc. U. S. Nat. Mus. 1882, 62.

A. saccharum, Marshall, Arbustum, 4.

A. barbatum, Michaux, Fl. Bor.-Am. ii, 253.—Willdenow, Spec. iv, 989.—Poiret, Suppl. ii, 575.—Pursh, Fl. Am. Sept. i, 266.—Nuttall, Genera, i, 255.—Elliott, Sk. i, 451.—De Candolle, Prodr. i, 595.—Torrey, Fl. U. S. 396; Compend. Fl. N. States, 169.—Eaton, Manual, 6 ed. 2.—Sprengel, Syst. ii, 224.—Don, Miller's Dict. i, 649.—Beck, Bot. 63.—Spach, Hist. Veg. iii, 178; Ann. Sci. Nat. 2 ser. ii, 118.—Torrey & Gray, Fl. N. America, i, 249, 684.—Eaton & Wright, Bot. 112.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 51.

SUGAR MAPLE. SUGAR TREE. HARD MAPLE. ROCK MAPLE.

Southern Newfoundland, valleys of the Saint Lawrence and Saguenay rivers, shores of lake Saint John, west along the northern shores of the great lakes to Lake of the Woods; south through the northern states and along the Alleghany mountains to northern Alabama and the Chattahoochee region of west Florida (var. *Floridanum*, *Chapman*, l. c.); west to Minnesota, eastern Nebraska, eastern Kansas (rare), and eastern Texas.

A tree of great economic value, 24 to 36 meters in height, with a trunk 0.60 to 1.20 meter in diameter, or toward its southwestern limits greatly reduced in size; rich woods, often forming extensive forests, and reaching its greatest development in region of the great lakes.

Wood heavy, hard, strong, tough, close-grained, compact, susceptible of a good polish; medullary rays numerous, thin; color, light brown tinged with red, the sap-wood lighter; specific gravity, 0.6912; ash, 0.54; largely used in the manufacture of furniture, shoe lasts and pegs, saddle-trees, in turnery, for interior finish, and flooring; in ship-building for keels, keelsons, shoes, etc., and furnishing valuable fuel; "curled" maple and "bird's-eye" maple, accidental forms in which the grain is beautifully curled and contorted, are common and highly prized in cabinet-making.

Maple sugar is principally made from this species; the ashes of the wood, rich in alkali, yield large quantities of potash.

Var. nigrum, Torrey & Gray,

Fl. N. America, i, 248.—Torrey, Fl. N. York, i, 136.—London, Arboretum, i, 411.—Browne, Trees of America, 84.—Gray, Manual N. States, 5 ed. 119.—Vasey, Cat. Forest Trees, 10.—Bell in Geological Rep. Canada, 1879-'80, 54c.

A. saccharinum, Michaux, Fl. Bor.-Am. ii, 252 [not Wangenheim].

A. nigrum, Michaux f. Hist. Arb. Am. ii, 238, t. 16; N. American Sylva, 3 ed. i, 163, t. 43.—Pursh, Fl. Am. Sept. i, 266.—Poiret, Suppl. v, 669.—Nuttall, Genera, i, 253.—Elliott, Sk. i, 450.—De Candolle, Prodr. i, 595.—Torrey, Fl. U. S. 397; Compend. Fl. N. States, 170.—Sprengel, Syst. ii, 225.—Don, Miller's Dict. i, 650.—Beck, Bot. 63.—Eaton, Manual, 6 ed. 2.—Spach, Hist. Veg. iii, 104; Ann. Sci. Nat. 2 ser. ii, 170.—Dietrich, Syn. ii, 1282.—Eaton & Wright, Bot. 112.—Koch, Dendrologie, i, 532.—Gray in Am. Nat. vi, 767; vii, 422.—Wood, Cl. Book, 286; Bot. & Fl. 74.

BLACK SUGAR MAPLE.

Western Vermont, shores of lake Champlain, westward to southern Missouri, south through Tennessee to northern Alabama, the valley of the Chickasaw river, Mississippi (*Mohr*), and southwestern Arkansas (Fulton, *Letterman*).

A large tree along streams and river bottoms, in lower ground than the species with which it is connected by numerous intermediate forms.

Wood heavier than that of the species; specific gravity, 0.6915; ash, 0.71.

65.—*Acer dasycarpum*, Ehrhart.

Beitr. iv, 24.—Möench, Meth. 56.—Persoon, Syn. i, 417.—Willdenow, Spec. iv, 985; Enum. ii, 1044.—Aiton, Hort. Kew. 2 ed. v, 446.—Pursh, Fl. Am. Sept. i, 266.—Nuttall, Genera, i, 252; Sylva, ii, 87; 2 ed. ii, 35.—Hayne, Dend. Fl. 213.—Elliott, Sk. i, 449.—Torrey, Fl. U. S. 396; Compend. Fl. N. States, 169; Fl. N. York, i, 136, t. 18; Nicotlet's Rep. 147.—Sprengel, Syst. ii, 225.—Tausch, Regensb. Fl. xii³, 553.—Eaton, Manual, 6 ed. 2.—London, Arboretum, i, 423, fig. 129 & t.—Hooker, Fl. Bor.-Am. i, 113; Jour. Bot. i, 200.—Bigelow, Fl. Boston. 3 ed. 407.—Torrey & Gray, Fl. N. America, i, 248.—Eaton & Wright, Bot. 112.—Emerson, Trees Massachusetts, 487; 2 ed. ii, 556 & t.—Parry in Owen's Rep. 610.—Darlington, Fl. Cestrica, 3 ed. 46.—Richardson, Arctic Exped. 423.—Darby, Bot. S. States, 265.—Cooper in Smithsonian Rep. 1858, 251.—Chapman, Fl. S. States, 81.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 51.—Lesquereux in Owen's 2d Rep. Arkansas, 354.—Wood, Cl. Book, 286; Bot. & Fl. 74.—Engelmann in Trans. Am. Phil. Soc. new ser. xii, 187.—Buchenau in Bot. Zeit. xix, 285, t. 11.—Gray, Manual N. States, 5 ed. 119.—Vasey, Cat. Forest Trees, 10.—Coulter's Bot. Gazette, v, 88.—Koch, Dendrologie, i, 541.—Sears in Bull. Essex Inst. xiii, 3.—Bell in Geological Rep. Canada, 1879-'80, 53c.—Nicholson in London Gard. Chronicle, 1881, 136, f. 24.—Ridgway in Proc. U. S. Nat. Mus. 1882, 62.

A. saccharinum, Linnaeus, Spec. 1 ed. 1055.

A. rubrum, var. *pallidum*, Aiton, Hort. Kew. iii, 434.

A. eriocarpum, Michaux, Fl. Bor.-Am. ii, 253.—Desfontaines in Ann. Mus. vii, 412, t. 25, f. 1; Hist. Arb. i, 392.—Poiret, Suppl. ii, 573.—Trattinick, Archiv. i, t. 8.—Michaux f. Hist. Arb. Am. ii, 205, t. 13; N. American Sylva, 3 ed. i, 146, t. 40.—Nouveau Duhamel, iv, 30.—De Candolle, Prodr. i, 595.—Don, Miller's Dict. i, 650.—Penn. Cycl. i, 79.—Beck, Bot. 63.—Spach, Hist. Veg. iii, 116; Ann. Sci. Nat. 2 ser. ii, 177.—Darlington, Fl. Cestrica, 2 ed. 245.—Dietrich, Syn. ii, 1282.—Browne, Trees of America, 95.—Meehan in Proc. Philadelphia Acad. 1868, 140.

SOFT MAPLE. WHITE MAPLE. SILVER MAPLE.

Valley of the Saint John's river, New Brunswick, to Ontario, south of latitude 45°, south to western Florida; west to eastern Dakota, eastern Nebraska, the valley of the Blue river, Kansas, and the Indian territory.

A large tree, 18 to 30 or, exceptionally, 36 meters in height, with a trunk 1.20 to 1.80 meter in diameter; along streams and intervalles, in rich soil; most common west of the Alleghany mountains, and reaching its greatest development in the basin of the lower Ohio river.

Wood light, hard, strong, brittle, close-grained, compact, easily worked; medullary rays numerous, thin; specific gravity, 0.5269; ash, 0.33; somewhat used in the manufacture of cheap furniture, for flooring, etc.; maple sugar is occasionally made from this species.

66.—*Acer rubrum*, Linnaeus.

Spec. 1 ed. 1055.—Du Roi, Diss. 59.—Marshall, Arbustum, 3.—Lamarck, Dict. ii, 300; Ill. iii, 438, t. 844, f. 3.—Ehrhart, Beitr. iv, 23.—Abbot, Insects Georgia, ii, 93.—Aiton, Hort. Kew. iii, 434 (excl. var.); 2 ed. v, 446.—Mönch, Meth. 56.—Michaux, Fl. Bor.-Am. ii, 253.—Persoon, Syn. i, 417.—Robin, Voyages, iii, 471.—Nouveau Duhamel, iv, 31.—Willdenow, Spec. iv, 984; Enum. ii, 1044.—Desfontaines in Ann. Mus. vii, 413, t. 25, f. 2; Hist. Arb. i, 391.—Poiret, Suppl. ii, 574.—Trattinick, Archiv. i, t. 9.—Michaux f. Hist. Arb. Am. ii, 210, t. 14; N. American Sylva, 3 ed. i, 149, t. 41.—Pursh, Fl. Am. Sept. i, 265.—Bigelow, Fl. Boston. 377.—Nuttall, Genera, i, 252.—Eaton, Manual, 44; 6 ed. 2.—Hayne, Dend. Fl. 213.—Elliott, Sk. i, 449.—Torrey, Fl. U. S. 395; Compend. Fl. N. States, 169; Fl. N. York, i, 137.—Watson, Dend. Brit. ii, t. 169.—Sprengel, Syst. ii, 225.—Audubon, Birds, t. 54, 67.—Tansch, Regensb. Fl. xii^a, 552.—Penn. Cycl. i, 79.—Hooker, Fl. Bor.-Am. i, 114; Jour. Bot. i, 199.—Don, Miller's Dict. i, 650.—Beck, Bot. 63.—Spach, Hist. Veg. iii, 113; Ann. Sci. Nat. 2 ser. ii, 176.—London, Arboretum, i, 424, f. 130 & t.—Torrey & Gray, Fl. N. America, i, 249, 684.—Dietrich, Syn. ii, 1282.—Eaton & Wright, Bot. 112.—Bigelow, Fl. Boston. 3 ed. 405.—Walpers, Rep. i, 409.—Reid in London Gard. Chronicle, 1844, 276.—Emerson, Trees Massachusetts, 483; 2 ed. ii, 551 & t.—Parry in Owen's Rep. 610.—Richardson, Arctic Exped. 422.—Nuttall, Sylva, ii, 87; 2 ed. ii, 34.—Darlington, Fl. Cestrica, 3 ed. 46.—Darby, Bot. S. States, 265.—Cooper in Smithsonian Rep. 1858, 251.—Chapman, Fl. S. States, 81.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 50.—Lesquereux in Owen's 2d Rep. Arkansas, 354.—Wood, Cl. Book, 286; Bot. & Fl. 74.—Engelmann in Trans. Am. Phil. Soc. new ser. xii, 187.—Porcher, Resources S. Forests, 79.—Buchenau in Bot. Zeit. xix, 285, t. 11.—Gray, Manual N. States, 5 ed. 119.—Koch, Dendrologie, i, 542.—Young, Bot. Texas, 206.—Vasey, Cat. Forest Trees, 10.—Macoun in Geological Rep. Canada, 1875-'76, 192.—Sears in Bull. Essex Inst. xiii, 176.—Bell in Geological Rep. Canada, 1879-'80, 54^c.—Nicholson in London Gard. Chronicle, 1881, 172, f. 30, 31.—Ridgway in Proc. U. S. Nat. Mus. 1882, 62.

? *A. glaucum*, Marshall, Arbustum, 2.

? *A. Caroliniana*, Walter, Fl. Caroliniana, 251.

A. coccineum, Michaux f. Hist. Arb. Am. ii, 203; N. American Sylva, 3 ed. i, 142.

A. sanguineum, Spach, Hist. Veg. iii, 115; Ann. Sci. Nat. 2 ser. ii, 176.—Dietrich, Syn. ii, 1282.

RED MAPLE. SWAMP MAPLE. SOFT MAPLE. WATER MAPLE.

New Brunswick, Quebec and Ontario, south of latitude 49°, north and west to the Lake of the Woods, south to Indian and Caloosa rivers, Florida, west to eastern Dakota, eastern Nebraska, the Indian territory, and the valley of the Trinity river, Texas.

A large tree, 20 to 30 or, exceptionally, 32 meters in height, with a trunk 0.90 to 1.50 meter in diameter; borders of streams and low, wet swamps, reaching its greatest development in the valleys of the lower Wabash and Yazoo rivers.

Wood heavy, hard, not strong, close-grained, compact, easily worked; medullary rays numerous, obscure; color, brown, often tinged with red, the sap-wood lighter; specific gravity, 0.6178; ash, 0.37; largely used in cabinet-making, turnery, and for woodenware, gun stocks, etc.; an accidental variety with undulating grain is highly valued.

Ink is occasionally made, domestically, by boiling the bark of this species in soft water and combining the tannin with sulphate of iron; formerly somewhat used in dyeing.

Var. *Drummondii*.

A. Drummondii, Hooker & Arnott in Hooker, Jour. Bot. i, 199.—Nuttall, Sylva, ii, 83, t. 70; 2 ed. ii, 30, t. 70.

Southern Arkansas, eastern Texas, western Louisiana, and sparingly through the Gulf states to southern Georgia.

Well characterized by its obovate or truncate leaves, the base entire or slightly crenulate-toothed, densely covered, as well as the petioles and young shoots, with a thick white tomentum; fruit convergent, the wings bright red, even when fully ripe.

A large tree, in deep, wet swamps, connected with the species by numerous intermediate forms of Georgia, Florida, and Alabama.

Wood lighter than that of the species; specific gravity, 0.5459; ash, 0.34.

67.—*Negundo aceroides*, Mönch.

Meth. 334.—Torrey & Gray, Fl. N. America, i, 250.—Eaton & Wright, Bot. 327.—Torrey in Nicolle's Rep. 147; Fremont's Rep. 88; Pacific R. R. Rep. iv, 73.—Nuttall, Sylva, ii, 92; 2 ed. ii, 38.—Gray in Jour. Boston Soc. Nat. Hist. vi, 166; Mem. Am. Acad. new ser. iv, 29; v, 309; Genera, ii, 202, t. 175; Pacific R. R. Rep. xii, 41; Manual N. States, 5 ed. 120.—Richardson, Arctic Exped. 423.—Parry in Owen's Rep. 610.—Darlington, Fl. Cestrica, 3 ed. 46.—Cooper in Smithsonian Rep. 1858, 251; Am. Nat. iii, 306.—Chapman, Fl. S. States, 81.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 53.—Wood, Cl. Book, 287; Bot. & Fl. 74.—Engelmann in Trans. Am. Phil. Soc. new ser. xii, 188.—Porter in Hayden's Rep. 1870, 474.—Watson in King's Rep. v, 52; Pl. Wheeler, 7.—Porter & Coulter, Fl. Colorado; Hayden's Surv. Misc. Pub. No. 4, 19.—Macoun & Gibson in Trans. Bot. Soc. Edinburgh, xii, 319.—Young, Bot. Texas, 207.—Vasey, Cat. Forest Trees, 10.—Macoun in Geological Rep. Canada, 1875-'76, 192.—Brewer & Watson, Bot. California, i, 108.—Rothrock in Wheeler's Rep. vi, 84.—Hemsley, Bot. Am.-Cent. i, 214.—Sears in Bull. Essex Inst. xiii, 176.—Bell in Geological Rep. Canada, 1879-'80, 48^c.—Nicholson in London Gard. Chronicle, 1881, 815.—Ridgway in Proc. U. S. Nat. Mus. 1882, 63.—Watson in Proc. Am. Acad. xvii, 338.

Acer Negundo, Linnaeus, Spec. 1 ed. 1056.—Wangenheim, Amer. 30, t. 12, f. 29.—Marshall, Arbustum, 2.—Lamarck, Diet. ii, 380.—Walter, Fl. Caroliniana, 250.—Aiton, Hort. Kew. iii, 436; 2 ed. v, 448.—Michaux, Fl. Bor.-Am. ii, 253.—Persoon, Syn. i, 418.—Desfontaines, Hist. Arb. i, 391.—Willdenow, Spec. iv, 992; Entm. ii, 1046.—Nouveau Duhamel, iv, 27, t. 7.—Trattinick, Archiv. i, t. 40.—Michaux f. Hist. Arb. Am. ii, 247, t. 18; N. American Sylva, 3 ed. i, 172, t. 46.—Pursh, Fl. Am. Sept. i, 268.—Hayne, Dend. Fl. 216.—Elliott, Sk. i, 452.—James in Long's Exped. ii, 69.—Torrey, Fl. U. S. 298; Compend. Fl. N. States, 170; Ann. Lyc. N. York, ii, 172; Emory's Rep. 407.—Sprengel, Syst. ii, 225.—Guimpel, Otto & Hayne, Abb. Holz. 119, t. 95.—Eaton, Manual, 6 ed. 2.—Dietrich, Syn. ii, 1283.—London, Arboretum, i, 460, t. 46, 47.—Darby, Bot. S. States, 265.—Buchenau in Bot. Zeit. xiv, 285, t. 11 & figs.—Koch, Dendrologie, i, 544.—Baillon, Hist. Pl. v, 374, f. 426.

Negundium fraxinifolium, Rafinesque, Med. Rep. v, 354.—Desvaux, Jour. Bot. v, 170.

Negundo fraxinifolium, Nuttall, Genera, i, 253.—De Candolle, Prodr. i, 596.—Hooker, Fl. Bor.-Am. i, 114; Jour. Bot. i, 200.—Don, Miller's Diet. i, 651.—Beck, Bot. 64.—Spach, Hist. Veg. iii, 119.—Rafinesque, New Fl. & Bot. i, 48.—Browne, Trees of America, 106.—Scheele in Rœmer, Texas, 433.—Schinzlein, Icon. t. 227, f. 2, 18.

? *N. Mexicanum*, De Candolle, Prodr. i, 596.—Hemsley, Bot. Am.-Cent. i, 214.

N. trifoliatum, Rafinesque, New Fl. & Bot. i, 48.

N. lobatum, Rafinesque, New Fl. & Bot. i, 48.

N. Californicum, Scheele in Rœmer, Texas, 433 [not Torrey & Gray].

BOX ELDER. ASH-LEAVED MAPLE.

Shores of the Winooski river and lake Champlain, Vermont, near Ithaca, New York, eastern Pennsylvania, and south to Hernando county, Florida (not detected in northeastern Florida); northwest through the lake region of the United States and Manitoba to the Dog's Head, lake Winnipeg, and along the southern branch of the Saskatchewan to the eastern base of the Rocky mountains; west in the United States to the eastern slopes of the Rocky mountains of Montana, through Colorado to the Wahsatch mountains, Utah; southwest through the basin of the Mississippi river, western Texas, and New Mexico to the Mogollon mountains, eastern Arizona; southward into Mexico.

A tree 15 to 22 meters in height, with a trunk 0.60 to 0.90 or, exceptionally, 1.20 meter in diameter; moist soil, borders of streams, etc.; in the Rocky Mountain region in high valleys, between 5,000 and 6,000 feet elevation; one of the most widely distributed trees of the American forest, reaching its greatest development in the valleys of the Wabash and Cumberland rivers.

Wood light, soft, not strong, close-grained, compact; medullary rays numerous, thin; color, creamy-white, the sap-wood hardly distinguishable; specific gravity, 0.4328; ash, 1.07; occasionally used in the interior finish of houses, for woodenware, cooperage, and paper-pulp.

Small quantities of maple sugar are sometimes obtained from this species.

68.—*Negundo Californicum*, Torrey & Gray,

Fl. N. America, i, 250, 684.—Hooker & Arnott, Bot. Beechey, 327, t. 77.—Eaton & Wright, Bot. 327.—Walpers, Rep. i, 410.—Bentham, Pl. Hartweg. 301.—Nuttall, Sylva, ii, 90, t. 72; 2 ed. ii, 37, t. 72.—Cooper in Smithsonian Rep. 1858, 258, in part.—Koch, Dendrologie, i, 545.—Brewer & Watson, Bot. California, i, 108.—Vasey, Cat. Forest Trees, 10.—Nicholson in London Gard. Chronicle, 1881, 815.

Acer Californicum, Dietrich, Syn. ii, 1283.

N. aceroides, Torrey in Pacific R. R. Rep. iv, 74; Bot. Mex. Boundary Survey, 47; Bot. Wilkes Exped. 259 [not Mönch].—Bolander in Proc. California Acad. iii, 78.

BOX ELDER.

California, valley of the lower Sacramento river (Sacramento, and in Marin and Contra Costa counties), southward in the interior valleys of the Coast ranges to about latitude 35°, cañons of the western slopes of the San Bernardino mountains (*Parish Brothers*).

A small tree, 6 to 12 meters in height, with a trunk 0.30 to 0.60 meter in diameter; borders of streams.

Wood light, soft, not strong, close-grained, compact; medullary rays numerous, thin; color, nearly white, or slightly tinged with yellow; specific gravity, 0.4821; ash, 0.54; occasionally used in the manufacture of cheap furniture.

ANACARDIACEÆ.

69.—*Rhus cotinoides*, Nuttall,

Mss. in Herb. Philadelphia Acad.; Travels, 177.—Cooper in Smithsonian Rep. 1858, 250.—Chapman, Fl. S. States, 70.—Wood, Cl. Book, 285; Bot. & Fl. 73.—Buckley in Proc. Philadelphia Acad. 1881, 125.—Mohr in Proc. Philadelphia Acad. 1881, 217.

R. cotinus? Torrey & Gray, Fl. N. America, i, 216.—Wood, Cl. Book, 285.

Cotinus Americanus, Nuttall, Sylva, iii, 1, t. 81; 2 ed. ii, 71, t. 81.

Cotinus coggygia, Engler in De Candolle, Suites, iv, 351, in part.

Indian territory, "on the light, broken, calcareous, rocky banks of the Grand river, a large tributary of the Arkansas, at a place then known as the Eagle's Nest," (Nuttall, l. c.); Alabama, north of the Tennessee river on southern slopes of the Cumberland mountains (on a hill near Bailie's farm, twelve miles from Huntsville, on the Madison road, Buckley, Mohr), and doubtfully reported north of the Alabama line, in Tennessee.

CHITTAM WOOD.

In Alabama, a small wide-branching tree, 9 to 10 meters in height, with a trunk sometimes 0.30 meter in diameter; on limestone benches from 700 to 900 feet elevation, in dense forests of oak, ash, maple, etc.; local and very rare; not rediscovered in Arkansas or the Indian territory; in Alabama nearly exterminated.

Wood light, soft, rather coarse-grained, checking badly in drying, very durable in contact with the soil; layers of annual growth marked by several rows of large open ducts; medullary rays, numerous, very obscure; color, bright, clear, rich orange, the thin sap-wood nearly white; specific gravity, 0.6425; ash, 0.50; largely used locally for fencing, and yielding a clear orange dye.

70.—*Rhus typhina*, Linnæus,

Amen. iv, 311.—Medicus, Bot. Beobacht. 1782, 228.—Wangenheim, Amer. 95.—Marshall, Arbustum, 129.—Walter, Fl. Caroliniana, 255.—Aiton, Hort. Kew. i, 365; 2 ed. ii, 162.—Ehrhart, Beitr. vi, 89.—Mœnch, Meth. 72.—Willdenow, Spec. i, 1478; Enum. i, 323.—B. S. Barton, Coll. i, 51.—Schkuhr, Handb. 237.—Michaux, Fl. Bor.-Am. i, 182.—Nouveau Duhamel, ii, 164, t. 47.—Persoon, Syn. i, 324.—Desfontaines, Hist. Arb. ii, 325.—Poiret in Lamarck, Dict. vii, 503.—Barton, Prodr. Fl. Philadelph. 39; Compend. Fl. Philadelph. i, 153.—Pursh, Fl. Am. Sept. i, 204.—Eaton, Mannal, 35; 6 ed. 302.—Nuttall, Genera, i, 203.—Rœmer & Schultes, Syst. vi, 643.—Hayne, Dend. Fl. 33.—Elliott, Sk. i, 360.—Torrey, Fl. U. S. 322; Compend. Fl. N. States, 140; Fl. N. York, i, 128.—De Candolle, Prodr. ii, 67.—Sprengel, Syst. i, 936.—Watson, Dend. Brit. i, t. 17, 18.—Hooker, Fl. Bor.-Am. i, 126.—Don, Miller's Dict. ii, 70.—Beck, Bot. 76.—Spach, Hist. Veg. ii, 212.—Bennett, Pl. Jav. Rar. 80.—Loudon, Arboretum, ii, 550, f. 224.—Torrey & Gray, Fl. N. America, i, 217, 680.—Eaton & Wright, Bot. 392.—Bigelow, Fl. Boston. 3 ed. 126.—Dietrich, Syn. ii, 1002.—Emerson, Trees Massachusetts, 501; 2 ed. ii, 571 & t.—Browne, Trees of America, 184.—Griffith, Med. Bot. 186.—Parry in Owen's Rep. 610.—Darlington, Fl. Cestrica, 3 ed. 43.—Richardson, Arctic Exped. 424.—Darby, Bot. S. States, 254.—Cooper in Smithsonian Rep. 1858, 250.—Chapman, Fl. S. States, 69.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 93.—Lesquereux in Owen's 2d Rep. Arkansas, 353.—Wood, Cl. Book, 384; Bot. & Fl. 73.—Porcher, Resources S. Forests, 208.—Gray, Manual N. States, 5 ed. 111.—Koch, Dendrologie, i, 576.—Young, Bot. Texas, 197.—Vasey, Cat. Forest Trees, 10.—Guibourt, Hist. Drogues, 7 ed. iii, 488.—Nat. Dispensatory, 2 ed. 1230.—Ridgway in Proc. U. S. Nat. Mus. 1882, 63.—Engler in De Candolle, Suites, iv, 377.

Datisca hirta, Linnæus, Spec. 1 ed. 1037.—Don, Miller's Dict. i, 290.

R. hypselodendron, Mœnch, Meth. 73.

R. Canadense, Miller, Dict. No. 5.—Nouveau Duhamel, ii, 163.

R. viridiflora, Nouveau Duhamel, ii, 163.—Poiret in Lamarck, Dict. vii, 504.—De Candolle, Prodr. ii, 67.—Nuttall, Genera, i, 203.—Don, Miller's Dict. ii, 70.—Dietrich, Syn. ii, 1002.—Loudon, Arboretum, ii, 551.—Browne, Trees of America, 184.

R. typhina, var. *viridiflora*, Engler in De Candolle, Suites, iv, 378.

STAGHOEN SUMACH.

New Brunswick, west through the valley of the Saint Lawrence river to southern Ontario and Minnesota, south through the northern states and along the Alleghany mountains to northern Georgia, central Alabama and Mississippi.

A small tree, rarely 9 meters in height, with a trunk 0.15 to 0.30 meter in diameter, or often a shrub; dry hillsides or often along streams in sandy, moist soil. A variety with laciniate leaves occurs near Hanover, New Hampshire, var. *laciniata*, Wood, *Cl. Book*, 284.—*Bot. & Fl.* 73).

Wood light, brittle, soft, coarse-grained, compact, satiny, susceptible of a good polish; layers of annual growth clearly marked by four to six rows of large open ducts; medullary rays numerous, obscure; color, yellow streaked with green, the sap-wood nearly white; specific gravity, 0.4357; ash, 0.50; occasionally used for inlaying cabinet work; the young shoots for "sap quills" in drawing the sap of the sugar maple.

Bark and leaves astringent, rich in tannin, and somewhat used locally as a dye and in dressing skins (*Special Rep. No. 26, U. S. Ag. Dep.* 22, t. 3); an infusion of the berries used domestically as a gargle in cases of catarrhal sore throat.

71.—*Rhus copallina*, Linnaeus.

Spec. 1 ed 266.—Medicus, *Bot. Beobacht.* 1782, 224.—Marshall, *Arbustum*, 128.—Wangenheim, *Amer.* 96.—Walter, *Fl. Caroliniana*, 255.—Gärtner, *Fruct.* i, 205, t. 44.—Aiton, *Hort. Kew.* i, 366; 2 ed. ii, 163.—Plenck, *Icon.* t. 233.—Lamarck, *Ill.* ii, 346, t. 207, f. 3.—Jacquin, *Hort. Schön.* iii, 50, t. 341.—Willdenow, *Spec.* i, 1490; *Enum.* i, 324.—Michaux, *Fl. Bor.-Am.* i, 182.—Schkuhr, *Handb.* 237.—Nouveau Duhamel, ii, 160.—Persoon, *Syn.* i, 324.—Desfontaines, *Hist. Arb.* ii, 325.—Poiret in Lamarck, *Dict.* vii, 506.—Barton, *Prodr. Fl. Philadelph.* 39.—Pursh, *Fl. Am. Sept.* i, 205.—Eaton, *Manual*, 34; 6 ed. 302.—Nuttall, *Genera*, i, 203.—Römer & Schultes, *Syst.* vi, 647.—Hayne, *Dend. Fl.* 34.—Elliott, *Sk.* i, 362.—Torrey, *Fl. U. S.* 323; *Compend. Fl. N. States*, 140; *Fl. N. York*, 129.—De Candolle, *Prodr.* ii, 68.—Sprengel, *Syst.* i, 936.—Don, *Miller's Dict.* ii, 72.—Beck, *Bot.* 75.—Hooker in *Jour. Bot.* i, 202.—Spach, *Hist. Veg.* ii, 214.—Torrey & Gray, *Fl. N. America*, i, 217.—Eaton & Wright, *Bot.* 392.—Bigelow, *Fl. Boston*. 3 ed. 126.—Dietrich, *Syn.* ii, 1003.—Loudon, *Arboretum*, ii, 554.—Emerson, *Trees Massachusetts*, 503; 2 ed. ii, 574.—Griffith, *Med. Bot.* 188.—Gray in *Mem. Am. Acad. new ser.* vi, 28; *Manual N. States*, 5 ed. 111; *Hall's Pl. Texas*, 5.—Scheele in Römer, *Texas*, 431.—Darlington, *Fl. Cestrica*, 3 ed. 43.—Darby, *Bot. S. States*, 255.—Chapman, *Fl. S. States*, 69.—Curtis in *Rep. Geological Surv. N. Carolina*, 1860, iii, 92.—Lesquereux in Owen's 2d *Rep. Arkansas*, 352.—Wood, *Cl. Book*, 284; *Bot. & Fl.* 73.—Engelmann in *Trans. Am. Phil. Soc. new ser.* xii, 187.—Porcher, *Resources S. Forests*, 207.—Koch, *Dendrologie*, 575.—Young, *Bot. Texas*, 197.—Vasey, *Cat. Forest Trees*, 11.—*Nat. Dispensatory*, 2 ed. 1236.—Ward in *Bull. U. S. Nat. Mus.* No. 22, 73.—Ridgway in *Proc. U. S. Nat. Mus.* 1882, 63.—Engler in De Candolle, *Suites*, iv, 384.

? *R. copallina*, vars. *latifolia*, *latialata*, *angustifolia*, and *serrata*, Engler in De Candolle, *Suites*, iv, 384.

DWARF SUMACH.

Northern New England, south to Manatee and Caximbas bay, Florida, west to Missouri, Arkansas, and the valley of the San Antonio river, Texas.

A small tree, 6 to 9 meters in height, with a trunk 0.15 to 0.20 meter in diameter, or at the north a low shrub 1 to 2 meters in height; dry hills and ridges, reaching its greatest development in southern Arkansas and eastern Texas; running into various forms. The best marked is—

var. *leucantha*, De Candolle, *Prodr.* ii, 68.—Gray in *Jour. Boston Soc. Nat. Hist.* vi, 158.

R. leucantha, Jacquin, *Hort. Schön.* iii, 50, t. 342.—Spach, *Hist. Veg.* ii, 215.

R. copallina, var. *angustialata*, Engler in De Candolle, *Suites*, iv, 384.

Shrubby, leaflets lanceolate, flowers white.

Wood light, soft, not strong, coarse-grained, compact, satiny, susceptible of a good polish; layers of annual growth clearly marked by several rows of large open ducts; medullary rays thin, not prominent; color, light brown streaked with green, or often tinged with red; the sap-wood lighter; specific gravity, 0.5273; ash, 0.60.

Leaves and bark astringent, rich in tannin; the leaves largely collected, principally in Maryland, Virginia, West Virginia, and Tennessee, and ground for tanning and dyeing (*Special Rep. No. 26, U. S. Ag. Dep.* 26, t. 5); the fruit, acid and astringent, used, as well as that of the shrubby *Rhus glabra*, by herbalists in the form of decoctions, fluid extracts, etc., as a gargle in the treatment of sore throat.

Var. *lanceolata*, Gray,

Jour. Boston Soc. Nat. Hist. vi, 158.—Torrey, *Bot. Mex. Boundary Survey*, 44.—Watson in *Proc. Am. Acad.* xvii, 338.

R. copallina, var. *integrifolia*, Engler in De Candolle, *Suites*, iv, 384.

Western Texas, Dallas (*Reverchon*) to the Rio Grande.

A small tree, with lanceolate, elongated leaflets, 5 to 6 meters in height, with a trunk 0.12 to 0.15 meter in diameter; calcareous soil; common; specific gravity, 0.5184; ash, 0.85.

72.—*Rhus venenata*, De Candolle,

Prodr. ii, 68.—Hooker, Fl. Bor.-Am. i, 126.—Don, Miller's Dict. ii, 71.—Beck, Bot. 76.—Spach, Hist. Veg. ii, 215.—Lindley, Fl. Med. 284.—London, Arboretum, ii, 552, f. 226.—Torrey & Gray, Fl. N. America, i, 218, 681.—Eaton & Wright, Bot. 392.—Dietrich, Syn. ii, 1003.—Torrey, Fl. N. York, i, 130.—Browne, Trees of America, 186.—Griffith, Med. Bot. 185.—Emerson, Trees Massachusetts, 504; 2 ed. ii, 575 & t.—Darlington, Fl. Cestrica, 3 ed. 44.—Richardson, Arctic Exped. 424.—Cooper in Smithsonian Rep. 1858, 250.—Chapman, Fl. S. States, 69.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 93.—Lesquereux in Owen's 2d Rep. Arkansas, 353.—Wood, Cl. Book, 284; Bot. & Fl. 73.—Gray, Manual N. States, 5 ed. 111.—Vasey, Cat. Forest Trees, 11.—Bailey in Am. Nat. vii, 5, f. 3.—Ward in Bull. U. S. Nat. Mus. No. 22, 73.—Engler in De Candolle, Suites, iv, 397.

R. vernix, Linnæus, Spec. 1 ed. 265, in part.—Kalm, Travels, English ed. 177.—Medicus, Bot. Beobacht. 1782, 223.—Marshall, Arbustum, 130.—Wangenheim, Amer. 92.—Aiton, Hort. Kew. i, 366; 2 ed. ii, 163.—Plenck, Icon. t. 234.—Lamarck, Ill. ii, 346, t. 207, f. 2.—Willdenow, Spec. i, 1479; Enum. i, 323.—B. S. Barton, Coll. i, 23, 50.—Schkuhr, Handb. 236.—Michaux, Fl. Bor.-Am. i, 183.—Nouveau Duhamel, ii, 165.—Persoon, Syn. i, 324.—Desfontaines, Hist. Arb. ii, 325.—Poirot in Lamarck, Dict. vii, 505.—Nuttall, Genera, i, 203.—Barton, Prodr. Fl. Philadelph. 39; Compend. Fl. Philadelph. 154.—Pursh, Fl. Am. Sept. i, 205.—Eaton, Manual, 34; 6 ed. 302.—Bigelow, Med. Bot. i, 96, t. 10; Fl. Boston. 3 ed. 126.—Rœmer & Schultes, Syst. vi, 646.—Hayne, Dend. Fl. 34.—Elliott, Sk. i, 362.—Torrey, Fl. U. S. 323; Compend. Fl. N. States, 203.—Sprengel, Syst. i, 936.—Hooker, Jour. Bot. i, 202.—Darby, Bot. S. States, 255.—Porcher, Resources S. Forests, 206.

POISON SUMACH. POISON ELDER.

Northern New England, south to northern Georgia, Alabama, and western Louisiana, west to northern Minnesota, Missouri, and Arkansas.

A small tree, 6 to 8 meters in height, with a trunk sometimes 0.15 to 0.20 meter in diameter, or more often a tall shrub; low, wet swamps or, more rarely, on higher ground.

Wood light, soft, coarse-grained, moderately compact; layers of annual growth clearly marked by three or four rows of large open ducts; medullary rays thin, very obscure; color, light yellow streaked with brown, the sap-wood lighter; specific gravity, 0.4382; ash, 0.64.

The whole plant, as well as the allied *R. Toxicodendron*, to most persons exceedingly poisonous to the touch, owing to the presence of a volatile principle, *Toxicodendric acid* (*U. S. Dispensatory*, 14 ed. 908.—*Nat. Dispensatory*, 2 ed. 1464); the white milky sap turning black in drying and yielding a valuable lacquer (*Bigelow, Med. Bot. l. c.*)

73.—*Rhus Metopium*, Linnæus,

Amœn. v, 395.—Titford, Hort. Bot. Am. 51.—Descourtilz, Fl. Med. Antilles, ii, 49, t. 79.—De Candolle, Prodr. ii, 67.—Macfadyen, Fl. Jamaica, 225.—Nuttall, Sylva, ii, 121, t. 80; 2 ed. ii, 68, t. 80.—Richard, Fl. Cuba, 381.—Cooper in Smithsonian Rep. 1858, 264.—Grisebach, Fl. British West Indies, 175.—Chapman, Fl. S. States, 69.—Wood, Bot. & Fl. 73.—Vasey, Cat. Forest Trees, 11.

Metopium Linnæi, Engler in De Candolle, Suites, iv, 367.

POISON WOOD. CORAL SUMACH. MOUNTAIN MANCHINEEL. BUM WOOD. HOG PLUM. DOCTOR GUM.

Semi-tropical Florida, bay Biscayne to the southern keys; in the West Indies.

A tree 12 to 15 meters in height, with a trunk sometimes 0.60 meter in diameter, reaching in the United States its greatest development on the shores of bay Biscayne, near Miami; one of the most common trees of the region, the large specimens generally decayed.

Wood heavy, hard, not strong, close-grained, checking badly in drying, containing many evenly-distributed open ducts; medullary rays numerous, thin; color, rich dark brown streaked with red, the sap-wood light brown or yellow; specific gravity, 0.7917; ash, 2.39; little esteemed.

A resinous gum, emetic, purgative, and diuretic, is obtained from incisions made in the bark of this species (*Pharm. Jour.* vii, 270.—*Guibourt, Hist. Drogues*, 7 ed. iii, 489).

74.—*Pistacia Mexicana*, HBK.

Nov. Gen. & Spec. vii, 22, t. 608.—De Candolle, Prodr. ii, 64.—Gray in Smithsonian Contrib. v, 27.—Torrey, Bot. Mex. Boundary Survey, 44.—Cooper in Smithsonian Rep. 1858, 265.—Brewer & Watson, Bot. California, i, 109.—Vasey, Cat. Forest Trees, 11.—Hemsley, Bot. Am.-Cent. i, 221.—Watson in Proc. Am. Acad. xvii, 338.

Texas, valley of the Rio Grande (near the mouth of the Pecos river, *Bigelow*); southward into Mexico (Saltillo, *Palmer*, etc.).

Wood not collected.

LEGUMINOSÆ.

75.—*Eysenhardtia orthocarpa*, Watson,

Proc. Am. Acad. xvii, 339.

E. amorphoides, var. *orthocarpa*, Gray in Smithsonian Contrib. iii, 46; v, 237.

E. amorphoides, Torrey, Bot. Mex. Boundary Survey, 51, in part.

Western Texas, valleys of the upper Guadalupe and Rio Grande, west to the Santa Rita and Santa Catalina mountains, Arizona (*Pringle*); southward into northern Mexico.

A small tree, 5 to 6 meters in height, with a trunk 0.09 to 0.15 meter in diameter, or more often a low shrub; dry, gravelly soil, reaching its greatest development near the summit of the Santa Catalina mountains, at 3,000 feet altitude.

Wood heavy, hard, close-grained, very compact; layers of annual growth clearly defined by numerous rows of open ducts; medullary rays numerous, thin; color, light reddish-brown, sap-wood clear yellow; specific gravity, 0.8740; ash, 1.28.

76.—*Dalea spinosa*, Gray,

Mem. Am. Acad. new ser. v, 315; Ives' Rep. 10.—Torrey, Pacific R. R. Rep. iv, 78; vii, 9, t. 3.—Bot. Mex. Boundary Survey, 53.—Walpers, Ann. iv, 485.—Cooper in Smithsonian Rep. 1858, 266.—Watson in Proc. Am. Acad. xi, 132.—Brewer & Watson, Bot. California, i, 143.—Hemsley, Bot. Am.-Cent. 249.

Asagraea spinosa, Baillon in Adansonia, ix, 232; Hist. Pl. ii, 288.

Colorado desert, southern California (Agua Caliente, Toras, etc.), and eastward to the valley of the lower Gila river, Arizona.

A small tree, sometimes 6 meters in height, with a short, stout trunk 0.45 to 0.50 meter in diameter (*Parry*, *Parish Brothers*), or often a low shrub; dry, gravelly, rocky soil.

Wood light, soft, rather coarse-grained, containing many evenly-distributed open ducts; medullary rays numerous, thin; color, walnut-brown, the sap-wood nearly white; specific gravity, 0.5536; ash, 4.04.

77.—*Robinia Pseudacacia*, Linnæus,

Spec. 1 ed. 722.—Marshall, Arbustum, 133.—Wangenheim, Amer. 16, t. 7.—L'Heritier, Stirp. Nov. 158.—Walter, Fl. Caroliniana, 186.—Aiton, Hort. Kew. iii, 53; 2 ed. iv, 323.—Gärtner, Fruct. ii, 307, t. 145.—Willdenow, Spec. iii, 1131; Enum. i, 769.—Michaux, Fl. Bor.-Am. ii, 65.—Nouveau Duhamel, ii, 60, t. 16.—Poiret in Lamarck Diet. vi, 222; Ill. iii, 163, t. 606.—Persoon, Syn. ii, 311.—Desfontaines, Hist. Arb. ii, 302.—Michaux f. Hist. Arb. Am. iii, 245, t. 1; N. American Sylva, 3 ed. ii, 92, t. 76.—Pursh, Fl. Am. Sept. ii, 487.—Eaton, Manual, 82; 6 ed. 306.—Thomas in Am. Month. Mag. & Crit. Rev. ii, 90.—Nuttall, Genera, ii, 118.—Hayne, Dend. Fl. 140.—Elliott, Sk. ii, 242.—De Candolle, Prodr. ii, 261.—Sprengel, Syst. iii, 247.—Torrey in Ann. Lyc. N. York, ii, 178; Compend. Fl. N. States, 271; Fl. N. York, i, 165; Emory's Rep. 408.—Hooker, Fl. Bor.-Am. i, 140.—Audubon, Birds, t. 104.—Don, Miller's Diet. ii, 237.—Beck, Bot. 52.—Spach, Hist. Veg. i, 258.—Torrey & Gray, Fl. N. America, i, 294.—London, Arboretum, ii, 609, f. 305 & t.—Eaton & Wright, Bot. 397.—Bigelow, Fl. Boston. 3 ed. 295.—Browne, Trees of America, 197.—Emerson, Trees, Massachusetts, 460; 2 ed. ii, 522 & t.—Griffith, Med. Bot. 238, f. 123.—Dietrich, Syn. iv, 1053.—Darlington, Fl. Cestrica, 3 ed. 65.—Darby Bot. S. States, 280.—Cooper in Smithsonian Rep. 1858, 251.—Chapman, Fl. S. States, 94.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 48.—Lesquereux in Owen's 2d Rep. Arkansas, 356.—Wood, Cl. Book, 319; Bot. & Fl. 95.—Lemaire, Ill. Hort. xii, t. 427.—Porcher, Resources S. Forests, 188.—Gray, Manual N. States, 5 ed. 131.—Koch, Dendrologie, i, 55.—Verlot in Rev. Hort. 1873, 152 & f.—Young, Bot. Texas, 228.—Vasey, Cat. Forest Trees, 11.—Ridgway in Proc. U. S. Nat. Mus. 1882, 65.

Pseudacacia odorata, Moench, Meth. 145.

R. fragilis, Salisbury, Prodr. 336.

LOCUST. BLACK LOCUST. YELLOW LOCUST.

Alleghany mountains, Pennsylvania (Locust ridge, Monroe county, *Porter*) to northern Georgia; widely and generally naturalized throughout the United States east of the Rocky mountains, and possibly indigenous in northeastern (Crowley's ridge) and western Arkansas and the prairies of eastern Indian territory.

A tree 22 to 25 meters in height, with a trunk 0.90 to 1.20 meter in diameter; west of the Mississippi river much smaller or often a low shrub 1.80 to 3 meters in height, reaching its greatest development on the western slopes of the mountains of West Virginia.

Wood heavy, exceedingly hard and strong, close-grained, compact, very durable in contact with the ground; layers of annual growth clearly marked by two or three rows of large open ducts; color, brown or, more rarely, light green, the sap-wood yellow; specific gravity, 0.7333; ash, 0.51 (*Trecul in Am. Jour. Sci.* 3 ser. xix, 182, t. 2, f. 1; t. 6, 7, f. 10.); largely used in ship-building, for posts of all sorts, construction, and in turnery; preferred to other American woods for treenails, and in this form largely exported.

The bark of the root tonic, or in large doses purgative and emetic (*U. S. Dispensatory*, 14 ed. 1746.—*Nat. Dispensatory*, 2 ed. 1233); formerly widely planted as a timber tree (*Cobbett, Woodlands*, par. 323); its cultivation in the United States now generally abandoned on account of the destructive attacks of the locust borer (*Oyllene picta*, *Packard in Bull. U. S. Entomological Com.* No. 7, 95).

78.—*Robinia viscosa*, Ventenat,

Hort. Cels. 4, t. 4.—Bot. Mag. t. 560.—Willdenow, Spec. iii, 1131; Enum. ii, 769.—Michaux, Fl. Bor.-Am. ii, 65.—Nouveau Duhamel, ii, 64, t. 17.—Poir. in Lamarck, Dict. vi, 222.—B. S. Barton, Bot. Appx. 29, t. 21.—Persoon, Syn. ii, 311.—Desfontaines, Hist. Arb. ii, 302.—Aiton, Hort. Kew. 2 ed. iv, 323.—Michaux f. Hist. Arb. Am. iii, 262, t. 2; N. American Sylva, ii, 104, t. 77.—Pursh, Fl. Am. Sept. ii, 488.—Nuttall, Genera, ii, 118.—Hayne, Dend. Fl. 140.—Elliott, Sk. ii, 242.—DeCandolle, Prodr. ii, 262.—Guimpel, Otto & Hayne, Abb. Holz. 81, t. 65.—Sprengel, Syst. iii, 247.—Don, Miller's Dict. ii, 236.—Eaton, Manual, 6 ed. 306.—Spach, Hist. Veg. i, 260.—Torrey & Gray, Fl. N. America, i, 295.—Loudon, Arboretum, ii, 626, t. 87, f. 306.—Eaton & Wright, Bot. 397.—Browne, Trees of America, 209.—Dietrich, Syn. iv, 1053.—Darby, Bot. S. States, 280.—Cooper in Smithsonian Rep. 1858, 251.—Chapman, Fl. S. States, 94.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 49.—Wood, Cl. Book, 319; Bot. & Fl. 95.—Porcher, Resources S. Forests, 193.—Gray, Manual N. States, 5 ed. 131.—Vasey, Cat. Forest Trees, 11.

R. glutinosa, Curtis, Bot. Mag. t. 560.—Koch, Dendrologie, i, 59.

CLAMMY LOCUST.

"High Alleghany mountains south of latitude 35°" (*Michaux*). "Open woods, slopes of Buzzard ridge, altitude 4,500 feet, near Highland, Macon county, North Carolina" (*J. Donnell Smith*).

A small tree, 9 to 12 meters in height, with a trunk not exceeding 0.30 meter in diameter; very rare, and not rediscovered until 1882 by the numerous botanists who have visited, during the last thirty years, the localities where the Michauxs, father and son, discovered this species; widely cultivated and now occasionally naturalized in the Atlantic states.

Wood (of a cultivated specimen) heavy, hard, close-grained, compact; layers of annual growth clearly marked by many rows of open ducts; medullary rays numerous, thin; color, brown, the sap-wood light yellow; specific gravity, 0.8094; ash, 0.20.

79.—*Robinia Neo-Mexicana*, Gray,

Mem. Am. Acad. new ser. v, 314.—Torrey in Pacific R. R. Rep. iv, 79; Bot. Mex. Boundary Survey, 53.—Walpers, Ann. iv, 491.—Cooper in Smithsonian Rep. 1858, 265.—Watson in King's Rep. v, 419.—Porter & Coulter, Fl. Colorado; Hayden's Surv. Misc. Pub. No. 4, 23.—Vasey, Cat. Forest Trees, 11.

LOCUST.

Colorado, valley of the Purgatory river (near Trinidad), headwaters of the Canadian river, through western and southwestern New Mexico to the Santa Catalina and Santa Rita mountains (*Lemmon, Pringle*), Arizona (4,500 to 7,000 feet altitude), southern Utah, Mount Zion cañon, west fork of the Rio Virgin, and near Kanah.

A small tree, sometimes 6 to 8 meters in height, with a trunk 0.15 to 0.25 meter in diameter, or toward its upper limits of growth reduced to a low shrub; reaching its greatest development in the valley of the Purgatory river, Colorado.

Wood heavy, exceedingly hard, strong, close-grained, compact, satiny, containing many evenly-distributed open ducts; medullary rays, thin, conspicuous; color, yellow streaked with brown, the sap-wood light yellow; specific gravity, 0.8034; ash, 0.60.

80.—*Olneya Tesota*, Gray,

Mem. Am. Acad. new ser. v, 328; Ives' Rep. 11.—Torrey in Pacific R. R. Rep. iv, 11, 82; vii, 10, t. 5; Bot. Mex. Boundary Survey, 58.—Walpers, Ann. iv, 479, 587.—Cooper in Smithsonian Rep. 1858, 265.—Brewer & Watson, Bot. California, i, 157.—Vasey, Cat. Forest Trees, 11.—Hemsley, Bot. Am.-Cent. i, 260.

IRON WOOD. ARBOL DE HIERRO.

California, valley of the Colorado river south of the Mohave mountains, valley of the lower Gila river, southwestern Arizona; southward in Sonora.

A small tree in the United States, rarely 9 meters in height, with a trunk sometimes 0.45 meter in diameter; dry *arroyos* and cañons; in Sonora more common and of larger size.

Wood very heavy and hard, strong, brittle, close-grained, compact, the grain generally contorted, difficult to cut and work, susceptible of a high polish; medullary rays numerous, thin; color, rich dark brown streaked with red, the sap-wood clear bright yellow; specific gravity, 1.0602; ash, 2.29 (the heart-wood, 1.1486; ash, 2.59; sap-wood, 0.8958; ash, 1.85); occasionally manufactured into canes.

81.—*Piscidia Erythrina*, Linnæus,

Spec. 2 ed. 993.—Jacquin, Amer. 206.—Swartz, Obs. 277.—Lamarck, Dict. i, 443; Ill. iii, 163, t. 605.—Titford, Hort. Bot. Am. 84.—Lunan, Hort. Jam. i, 269.—Humboldt, Bonpland & Kunth, Nov. Gen. & Spec. vi, 382.—De Candolle, Prodr. ii, 267.—Descourtilz, Fl. Med. Antilles, iii, 203, t. 196.—Macfadyen, Fl. Jamaica, i, 258.—Nuttall, Sylva, ii, 31, t. 52; 2 ed. i, 180.—Bentham in Jour. Linnæan Soc. iv, Suppl. 116; Bot. Sulphur, 81.—Cooper in Smithsonian Rep. 1858, 264.—Chapman, Fl. S. States, 110.—Grisebach, Fl. British West Indies, 200.—Porcher, Resources S. Forests, 175.—Vasey, Cat. Forest Trees, 11.—Hemsley, Bot. Am.-Cent. i, 319.

Erythrina piscipula, Linnæus, Spec. 1 ed. 107.

P. Carthagenensis, De Candolle, Prodr. ii, 267.

JAMAICA DOGWOOD.

Semi-tropical Florida, bay Biscayne, west coast, Pease creek to cape Sable, and on the southern keys; in the West Indies and southern Mexico.

A tree 12 to 15 meters in height, with a trunk 0.45 to 0.75 meter in diameter.

Wood heavy, very hard, not strong, close-grained, compact, susceptible of a high polish, containing few large scattered open ducts; medullary rays thin, not conspicuous; color, yellowish-brown, the sap-wood lighter; specific gravity, 0.8734; ash, 3.38; one of the favorite woods of the region for boat-building, fire-wood, and charcoal.

The bark, especially of the root, narcotic, occasionally administered in the form of tinctures, or used, as well as the young branches and leaves, to poison or stupefy fish.

82.—*Cladrastis tinctoria*, Rafinesque,

Fl. Kent. 1824; Neog. 1825; Med. Bot. ii, 210; New Sylva, iii, 83.—Torrey & Gray, Fl. N. America, i, 390.—Walpers, Rep. i, 807.—Browne, Trees of America, 192.—Darby, Bot. S. States, 294.—Cooper in Smithsonian Rep. 1858, 251.—Chapman, Fl. S. States, 113.—Porcher Resources S. Forests, 175.—Wood, Cl. Book, 301; Bot. & Fl. 84.—Gray, Manual N. States, 5 ed. 143.—Vasey, Cat. Forest Trees, 11.

Virgilia lutea, Michaux f. Hist. Arb. Am. iii, 266, t. 3; Travels, 289; N. American Sylva, 3 ed. ii, 106, t. 78.—Parsh, Fl. Am. Sept. i, 309.—Nuttall, Genera, i, 284.—Hayne, Dend. Fl. 53.—Loiseleur, Herb. Amat. t. 297.—De Candolle, Prodr. ii, 93.—Sprengel, Syst. iv³, i, 171.—Don, Miller's Dict. ii, 112.—Eaton, Manual, 6 ed. 397.—Spach, Hist. Veg. i, 163.—Eaton & Wright, Bot. 480.—Dietrich, Syn. ii, 1501.—Loudon, Arboretum, ii, 565, t. 78.

O. lutea, Koch, Dendrologie, i, 6.

YELLOW WOOD. YELLOW ASH. GOPHER WOOD.

Central Kentucky, cliffs of the Kentucky and Dick's rivers; middle Tennessee, mountains of east Tennessee to Cherokee county, North Carolina.

A tree 9 to 15 meters in height, with a trunk sometimes 0.90 or, exceptionally, 1.20 meter in diameter; rich hillsides; in Kentucky on the Trenton limestones, and reaching its best development in middle Tennessee; rare and very local, the large trees generally hollow or defective.

Wood heavy, very hard, strong, close-grained, compact, susceptible of a good polish; layers of annual growth clearly marked by several rows of open ducts, and containing many evenly-distributed similar ducts; color, bright, clear yellow, changing with exposure to light brown, the sap-wood nearly white; specific gravity, 0.6278; ash, 0.28; used for fuel, occasionally for gunstocks, and yielding a clear yellow dye.

83.—*Sophora secundiflora*, Lagasca;

De Candolle, Cat. Hort. Mousp. 148; Prodr. ii, 96.—Don, Miller's Dict. ii, 110.—Gray in Smithsonian Contrib. iii, 54.—Rev. Hort. 4 ser. iii, 201, t. 11.—Bentham & Hooker, Genera, i, 555.—Hemsley, Bot. Am.-Cent. i, 321.—Watson in Proc. Am. Acad. xvii, 347.

Broussonetia secundiflora, Ortega, Dec. v, 61, t. 7.

Virgilia secundiflora, Cavanilles, Icon. t. 401.

Agastianis secundiflora, Rafinesque, New Sylva, iii, 86.

Dermatophyllum speciosum, Scheele in Linnæus, xxi, 458.

S. speciosa, Bentham in Jour. Boston Soc. Nat. Hist. vi, 176.—Gray in Mem. Am. Acad. new ser. iv², 38; Smithsonian Contrib. iii, 54; Hall's Pl. Texas, 7.—Walpers, Ann. ii, 439.—Torrey, Bot. Mex. Boundary Survey, 58.—Young, Bot. Texas, 242.—Vasey, Cat. Forest Trees, 12.

FRIGOLITO.

Matagorda bay, Texas, west to the mountains of New Mexico (*Havard*).

A small tree, sometimes 9 meters in height, with a trunk 0.15 to 0.20 meter in diameter, or often, especially west of the San Antonio river, a tall shrub, rarely exceeding 2 meters in height, forming dense thickets; borders of streams, generally in a low, rather moist soil.

Wood very heavy, hard, close-grained, compact, susceptible of a high polish; medullary rays numerous, thin; color, orange streaked with red, the heavier sap-wood brown or yellow; specific gravity, 0.9842; ash, 1.59; furnishing valuable fuel.

The seeds contain an exceedingly poisonous alkaloid, *Sophoria* (*H. C. Wood in Philadelphia Med. Times*, August 4, 1877.—*Rothrock in Coulter's Bot. Gazette*, ii, 133.—*Nat. Dispensatory*, 2 ed. 1333).

84.—*Sophora affinis*, Torrey & Gray,

Fl. N. America, i, 390.—Leavenworth in *Am. Jour. Sci.* 1 ser. ix, 130.—Gray in *Jour. Boston Soc. Nat. Hist.* vi, 178; Hall's *Fl. Texas*, 7.—Scheele in *Römer*, Texas, 428.—Vasey, *Cat. Forest Trees*, 12.

Styphnolobium affine, Walpers, *Rep.* i, 807.

Arkansas, valley of the Arkansas river (*Letterman*) to the valley of the San Antonio river, Texas.

A small tree, 5 to 7 meters in height, with a trunk sometimes 0.15 to 0.25 meter in diameter; borders of streams and prairies.

Wood heavy, very hard, strong, coarse-grained, compact; layers of annual growth clearly marked by several rows of large open ducts; medullary rays thin, conspicuous; color, light red, the sap-wood bright, clear yellow; specific gravity, 0.8509; ash, 0.73.

Ink is occasionally made domestically from the resinous exudations of the pod.

85.—*Gymnocladus Canadensis*, Lamarck,

Dict. i, 733; *Ill.* iii, 412, t. 823.—Michaux, *Fl. Bor.-Am.* ii, 241, t. 51.—Willdenow, *Spec.* iv, 460; *Enum.* ii, 1019; *Berl. Baumz.* 169.—Persoon, *Syn.* ii, 626.—Desfontaines, *Hist. Arb.* ii, 250.—Aiton, *Hort. Kew.* 2 ed. v, 400.—Michaux *f. Hist. Arb. Am.* ii, 272, t. 23; *N. American Sylva*, 3 ed. i, 182, t. 50.—Pursh, *Fl. Am. Sept.* i, 304.—Nuttall, *Genera*, ii, 243.—Hayne, *Dend. Fl.* 203.—James in *Long's Exped.* i, 138.—Reichenbach, *Mag. Bot.* t. 40.—De Candolle, *Prodr.* ii, 480.—Sprengel, *Syst.* ii, 327.—Torrey in *Ann. Lye. N. York*, ii, 193; *Compend. Fl. N. States*, 376; *Fl. N. York*, i, 196; *Emory's Rep.* 407.—Hooker, *Fl. Bor.-Am.* i, 166.—Don, *Miller's Dict.* 429.—Eaton, *Manual*, 6 ed. 162.—Beck, *Bot.* 93.—Spach, *Hist. Veg.* i, 89.—London, *Arboretum*, ii, 256 & t.—Torrey & Gray, *Fl. N. America*, i, 398.—Eaton & Wright, *Bot.* 258.—Richardson, *Arctic Exped.* 424.—Walpers, *Rep.* i, 809.—Browne, *Trees of America*, 218.—Cooper in *Smithsonian Rep.* 1858, 251.—Lesquereux in *Owen's 2d Rep. Arkansas*, 358.—Wood, *Cl. Book*, 300; *Bot. & Fl.* 83.—Engelmann in *Trans. Am. Phil. Soc.* new ser. xii, 190.—Gray, *Manual N. States*, 5 ed. 145.—Briot in *Rev. Hort.* 1870, 436.—Vasey, *Cat. Forest Trees*, 12.—Bell in *Geological Rep. Canada*, 1879-'80, 54^c.—Ridgway in *Proc. U. S. Nat. Mus.* 1882, 63.—Chapman, *Fl. S. States*, *Suppl.* 618.

Guilandina dioica, Linnæus, *Spec.* 1 ed. 381.—Marshall, *Arbustum*, 56.—Aiton, *Hort. Kew.* ii, 56.—James in *Long's Exped.* i, 138.

Hyperanthera dioica, Vahl, *Symbolæ*, i, 31.

G. dioica, Koch, *Dendrologie*, i, 5.—Baillon, *Hist. Pl.* ii, 87, f. 52, 53.

KENTUCKY COFFEE TREE. COFFEE NUT.

Conococheague creek, Franklin county, Pennsylvania (*Porter*); western New York, shores of Cayuga and Seneca lakes, west through southern Ontario and southern Michigan to the valley of the Minnesota river, Minnesota, eastern Nebraska, eastern Kansas, southwestern Arkansas, and the Indian territory, to about longitude 96° west, south to middle Tennessee.

A tree 25 to 33 meters in height, with a trunk 0.60 to 0.90 meter in diameter; rich woods and bottoms; not common.

Wood heavy, not hard, strong, coarse-grained, durable in contact with the ground, liable to check in drying, easily worked, susceptible of a high polish; layers of annual growth clearly marked by one or two rows of open ducts; medullary rays numerous, thin; color, rich light brown tinged with red, the thin sap-wood lighter; specific gravity, 0.6934; ash, 0.67; occasionally used in cabinet-making, for posts, rails, &c.

The fresh leaves, macerated and sweetened, are used in Tennessee as a poison for house-flies; the seeds formerly as a domestic substitute for coffee.

86.—*Gleditschia triacanthos*, Linnæus,

Spec. 1 ed. 1056 (excl. var.).—Medicus, Bot. Beobacht. 1782, 230.—Lamarck, Dict. ii, 465; Ill. iii, 446, t. 857, f. 1.—Aiton, Hort. Kew. iii, 444 (excl. vars.); 2 ed. v, 474.—Mönch, Meth. 69.—Abbot, Insects Georgia, ii, t. 285.—Michaux, Fl. Bor.-Am. ii, 257.—Schkuhr, Handb. iii, 554, t. 356.—Robin, Voyages, iii, 497.—Persoon, Syn. ii, 123.—Desfontaines, Hist. Arb. ii, 246.—Willdenow, Spec. iv, 1097; Enum. 1058; Berl. Baumz. 163.—Nouveau Duhamel, iv, 100, t. 25.—Michaux f. Hist. Arb. Am. iii, 164, t. 10; N. American Sylva, 3 ed. 108, t. 79.—Pursh, Fl. Am. Sept. i, 221.—Nuttall, Genera, ii, 239.—James in Long's Exped. i, 138.—Hayne, Dend. Fl. 218.—Elliott, Sk. ii, 709.—Guimpel, Otto & Hayne, Abb. Holz. 157, t. 132.—De Candolle, Prodr. ii, 479.—Sprengel, Syst. iii, 918.—Torrey, Compend. Fl. N. States, 375; Fl. N. York, i, 192.—Audubon, Birds, t. 42, 146, 150.—Römer & Schultes, Syst. vii, 78.—Don, Miller's Dict. ii, 428.—Beck, Bot. 93.—Eaton, Manual, 6 ed. 158.—Spach, Hist. Veg. i, 92.—Torrey & Gray, Fl. N. America, i, 398.—Loudon, Arboretum, ii, 650, t. 90, 91.—Eaton & Wright, Bot. 254.—Browne, Trees of America, 212.—Dietrich, Syn. iv, 539.—Darby, Bot. S. States, 295.—Cooper in Smithsonian Rep. 1858, 251.—Gray in Pacific R. R. Rep. xii², 42; Manual N. States, 5 ed. 145.—Chapman, Fl. S. States, 115.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 49.—Lesquereux in Owen's 2d Rep. Arkansas, 358.—Wood, Cl. Book, 300; Bot. & Fl. 83.—Engelmann in Trans. Am. Phil. Soc. new ser. xii, 190.—Porcher, Resources S. Forests, 195.—Koch, Dendrologie, i, 8.—Hunt in Am. Nat. i, 433.—Young, Bot. Texas, 246.—Vasey, Cat. Forest Trees, 12.—Ridgway in Proc. U. S. Nat. Mus. 1882, 64.—Burgess in Coulter's Bot. Gazette, vii, 95.

G. spinosa, Marshall, Arbustum, 54.

G. Meliloba, Walter, Fl. Caroliniana, 254.

G. macrantha, Willdenow, Berl. Baumz. 164.

G. elegans, Salisbury, Prodr. 323.

Melilobus heterophylla, Rafinesque, Sylva Telluriana, 121.

HONEY LOCUST. BLACK LOCUST. THREE-THORNED ACACIA. SWEET LOCUST. HONEY SHUCKS.

Pennsylvania, western slopes of the Alleghany mountains, west through southern Michigan to eastern Nebraska, eastern Kansas, and the Indian territory to about longitude 96° west; south to Tampa bay, Florida (not detected in eastern Florida), northern Alabama, northern Mississippi, and the valley of the Brazos river, Texas.

A tree, 25 or 30 meters, or exceptionally 40 meters, in height, with a trunk 0.60 to 1.20 meter in diameter; low, rich bottom lands, or more rarely on dry, sterile hills; the characteristic tree of the "barrens" of middle Kentucky and Tennessee, reaching its greatest development in the bottoms of the lower Ohio River basin; widely cultivated for shade and as a hedge plant, and now somewhat naturalized in the Atlantic states east of the Alleghany mountains.

A not uncommon form, nearly destitute of thorns, is—

var. *inermis*, Pursh, Fl. Am. Sept. i, 221.—De Candolle, Mem. Leg. t. 22, f. 109; Prodr. ii, 479.—Eaton, Manual, 6 ed. 158.—Torrey & Gray, Fl. N. America, i, 398.—Loudon Arboretum, ii, 650, t. 92, 93.—Browne, Trees of America, 213.

G. inermis, Linnæus, Spec. 1509, in part.—Nouveau Duhamel, iv, 100.—Bentham in Trans. Linnæan Soc. xxx³, 557.

A form with spines and fruit shorter than those of the type is—

var. *brachycarpus*, Michaux, Fl. Bor.-Am. ii, 257.—Torrey & Gray, Fl. N. America, i, 398.—Browne, Trees of America, 213.

G. brachycarpa, Pursh, Fl. Am. Sept. 221.—De Candolle, Prodr. ii, 479.—Sprengel, Syst. iii, 919.—Don, Miller's Dict. ii, 428.—Eaton, Manual, 6 ed. 158.—Eaton & Wright, Bot. 254.—Loudon, Arboretum, ii, 653.—Dietrich, Syn. iv, 539.

Wood heavy, hard, strong, coarse-grained, moderately compact, very durable in contact with the soil, susceptible of a high polish; layers of annual growth strongly marked by many rows of open ducts; medullary rays numerous, conspicuous; color, bright brown or red, the sap-wood lighter; specific gravity, 0.6740; ash, 0.80; used for fence posts and rails, wagon hubs, construction, etc.; its value hardly appreciated.

Beer is sometimes made domestically by fermenting the sweet, unripe fruit (*Porcher l. c.*).

87.—*Gleditschia monosperma*, Walter,

Fl. Caroliniana, 254.—Michaux, Fl. Bor.-Am. ii, 257.—Schkuhr, Handb. iii, 555.—Persoon, Syn. ii, 623.—Desfontaines, Hist. Arb. ii, 24.—Willdenow, Spec. iv, 1097; Enum. 1058; Berl. Baumz. 165.—Nouveau Duhamel, iv, 101.—Aiton, Hort. Kew. 2 ed. v, 474.—Michaux f. Hist. Arb. Am. iii, 169, t. 11; N. American Sylva, 3 ed. ii, 111, t. 80.—Pursh, Fl. Am. Sept. 221.—Poiret, Suppl. ii, 641.—Nuttall, Genera, ii, 239.—Hayne, Dend. Fl. 218.—Elliott, Sk. ii, 709.—De Candolle, Prodr. ii, 479.—Sprengel, Syst. iii, 919.—Don, Miller's Dict. 428.—Eaton, Manual, 6 ed. 158.—Spach, Hist. Veg. i, 98.—Torrey & Gray, Fl. N. America, i, 398.—Eaton & Wright, Bot. 254.—Loudon, Arboretum, ii, 653, f. 364.—Browne, Trees of America, 215.—Dietrich, Syn. iv, 539.—Darby, Bot. S. States, 295.—Chapman, Fl. S. States, 115.—Wood, Cl. Book, 300; Bot. & Fl. 83.—Gray, Manual N. States, 5 ed. 145.—Vasey, Cat. Forest Trees, 12.—Ridgway in Proc. U. S. Nat. Mus. 1882, 64.

G. triacanthos, var. *monosperma*, Linnæus, Spec. 1 ed. 1057.—Aiton, Hort. Kew. iii, 444.

G. aquatica, Marshall, Arbustum, 54.

G. Carolinensis, Lamarck, Dict. ii, 465; Ill. iii, 447, t. 857, f. 2.—Römer & Schultes, Syst. vii, 74.

G. triacantha, Gärtner, Fruct. ii, 311, t. 146, f. 3 [not Linnæus].

G. inermis, Koch, Dendrologie, i, 9 [not Linnæus].

WATER LOCUST.

South Carolina to Matanzas inlet and Tampa bay, Florida, through the Gulf states to the valley of the Brazos river, Texas, and through Arkansas to middle Kentucky and Tennessee, southern Indiana and Illinois.

A tree 12 to 18 meters in height, with a trunk sometimes 0.60 or, exceptionally, 0.90 meter in diameter; deep swamps; rare in the south Atlantic and Gulf states; common and reaching its greatest development in the bottom lands of southern Arkansas, Louisiana, and eastern Texas, here often covering extensive areas.

Wood heavy, very hard, strong, rather coarse-grained, compact, susceptible of a high polish; layers of annual growth clearly marked by one to three rows of open ducts; medullary rays thin, conspicuous; color, rich bright brown tinged with red, the thick heavier sap-wood clear light yellow; specific gravity, 0.7342; ash, 0.73.

88.—*Parkinsonia Torreyana*, Watson,

Proc. Am. Acad. xi, 135.—Brewer & Watson, Bot. California, i, 162.

Cercidium floridum, Torrey in Pacific R. R. Rep. iv, 11, 82; v, 360, t. 3; Bot. Mex. Boundary Survey, 59.—Gray in Ives' Rep. 11.—Vasey, Cat. Forest Trees, 12.—James in Am. Nat. xv, 982.—Hemsley, Bot. Am.-Cent. i, 327.

GREEN-BARK ACACIA. PALO VERDE.

Colorado desert, southern California (Inio, Toras, etc., *Parish Brothers*), east to the valley of the lower Gila river, Arizona.

A low, much-branched tree, 8 to 10 meters in height, the short trunk sometimes 0.45 to 0.50 meter in diameter; low cañons and depressions in the sandhills of the desert; common and reaching its greatest development in the valleys of the lower Colorado and Gila rivers.

Wood heavy, not strong, soft, close-grained, compact, satiny, susceptible of a beautiful polish, containing many small evenly-distributed open ducts; medullary rays very numerous, thin; color, light brown, the sap-wood clear light yellow; specific gravity, 0.6531; ash, 1.12.

89.—*Parkinsonia microphylla*, Torrey,

Pacific R. R. Rep. iv, 82; Bot. Mex. Boundary Survey, 59.—Walpers, Ann. vii, 812.—Gray in Ives' Rep. 11.—Bentham in Martius, Fl. Brasil. xv², 78.—Watson, Pl. Wheeler, 8; Proc. Am. Acad. xi, 136.—Brewer & Watson, Bot. California, i, 162.—Hemsley, Bot. Am.-Cent. i, 327.

Valley of the lower Colorado and Bill Williams rivers, eastward through southern Arizona.

A small, much-branched tree, 6 to 7 meters in height, with a trunk 0.25 to 0.30 meter in diameter (*Wickenburg, Pringle*), or often a low shrub 1 to 3 meters in height.

Wood heavy, hard, coarse-grained, compact, containing numerous large, scattered, open ducts; medullary rays numerous, thin, conspicuous; color, rich dark brown streaked with red, the sap-wood light brown or yellow; specific gravity, 0.7449; ash, 3.64.

90.—*Parkinsonia aculeata*, Linnaeus,

Spec. 1 ed. 375.—Jacquin, Stirp. Am. 121, t. 80.—Lamarck, Ill. ii, 475, t. 336.—Willdenow, Spec. ii, 513.—Aiton, Hort. Kew. 2 ed. iii, 24.—De Candolle, Mem. Leg. ii, t. 21; Prodr. ii, 486.—Descourtilz, Fl. Med. Antilles, i, 54, t. 12.—Macfadyen, Fl. Jamaica, 334.—Bentham, Bot. Sulphur, 87; Martius, Fl. Brasil. xv², 78, t. 26.—Cooper in Smithsonian Rep. 1858, 265.—Torrey, Bot. Mex. Boundary Survey, 59.—Grisebach, Fl. British West Indies, 204; Pl. Lorentz. 81.—Gray, Hall's Pl. Texas, 8.—Brewer & Watson, Bot. California, i, 162.—Vasey, Cat. Forest Trees, 12.—Hemsley, Bot. Am.-Cent. i, 327.—Watson in Proc. Am. Acad. xvii, 348.

Corpus Christi, Texas, west along the Mexican boundary to the valley of the Colorado river, Arizona (Yuma); and southward into Mexico; probably of American origin, but now widely naturalized throughout the tropical and warmer regions of the globe (*A. De Candolle, Geog. Bot.* ii, 719, 770, 793).

A small tree, 6 to 12 meters in height, with a trunk sometimes 0.30 meter in diameter.

Wood heavy, hard, very close-grained, inclined to check in drying, containing many evenly-distributed small open ducts; medullary rays very numerous, thin, conspicuous; color, light brown, the very thick sap-wood lighter, often tinged with yellow; specific gravity, 0.6116; ash, 2.32.

91.—*Cercis Canadensis*, Linnaeus,

Spec. 1 ed. 374.—Du Roi, Obs. Bot. 10.—Marshall, Arbustum, 32.—Lamarck, Dict. ii, 586.—Wangenheim, Amer. 84.—Walter, Fl. Caroliniana, 135.—Aiton, Hort. Kew. ii, 47; 2 ed. iii, 22.—Willdenow, Spec. ii, 508; Enum. 439; Berl. Baumz. 84.—Nouveau Duhamel, i, 19.—Michaux, Fl. Bor. Am. i, 265.—Schkuhr, Handb. 354.—Persoon, Syn. i, 454.—Desfontaines, Hist. Arb. ii, 254.—Pursh, Fl. Am. Sept. i, 308.—Eaton, Manual, 46; 6 ed. 89.—Nuttall, Genera, i, 283.—Hayne, Dend. Fl. 53.—Elliott, Sk. i, 470.—Torrey in Ann. Lyc. N. York, ii, 194; Fl. U. S. 441; Compend. Fl. N. States, 188; Fl. N. York, i, 188; Nicotlet's Rep. 149; Emory's Rep. 408.—De Candolle, Prodr. ii, 518.—Sprengel, Syst. ii, 346.—Guimpel, Otto & Hayne, Abb. Holz. 116, t. 92.—Hooker, Fl. Bor. Am. i, 167; Companion Bot. Mag. i, 24.—Don, Miller's Dict. ii, 468.—Beck, Bot. 94.—Spach, Hist. Veg. i, 129.—Torrey & Gray, Fl. N. America, i, 392.—Loudon, Arboretum, ii, 659 & t.—Eaton & Wright, Bot. 190.—Dietrich, Syn. ii, 155.—Browne, Trees of America, 221.—Gray in Mem. Am. Acad. new ser. iv, 38; Manual N. States, 5 ed. 144.—Richardson, Arctic Exped. 424.—Parry in Owen's Rep. 611.—Darlington, Fl. Cestrica, 3 ed. 67.—Darby, Bot. S. States, 294.—Cooper in Smithsonian Rep. 1858, 251.—Chapman, Fl. S. States, 114.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 50.—Lesqueureux in Owen's 2d Rep. Arkansas, 357.—Wood, Cl. Book, 301; Bot. & Fl. 84.—Engelmann in Trans. Am. Phil. Soc. new ser. xii, 190.—Porcher, Resources S. Forests, 197.—Koch, Dendrologie i, 14.—Baillon, Hist. Pl. ii, 121.—Vasey, Cat. Forest Trees, 12.—Ridgway in Proc. U. S. Nat. Mus. 1882, 65.

Siliquastrum cordatum, Mönch, Meth. 54.

C. Canadensis, var. *pubescens*, Pursh. Fl. Am. Sept. i, 308.—Loudon, Arboretum, ii, 659.

REDBUD. JUDAS TREE.

Western Pennsylvania, southward to Tampa bay, Florida, northern Alabama and Mississippi, westward through southern Michigan and Minnesota to eastern Nebraska; southwest through Missouri and Arkansas to the eastern portions of the Indian territory, Louisiana, and the valley of the Brazos river, Texas.

A small tree, 12 to 16 meters in height, with a trunk sometimes 0.30 meter in diameter; rich woods, borders of streams and swamps; most common and reaching its greatest development in southern Arkansas, the Indian territory, and eastern Texas, here, when in bloom, a conspicuous feature of the forest.

Wood heavy, hard, not strong, rather coarse-grained, compact, susceptible of a good polish; layers of annual growth clearly marked by one to three rows of open ducts; medullary rays exceedingly numerous, thin; color, rich dark brown tinged with red, the sap-wood lighter; specific gravity, 0.6363; ash, 0.72.

92.—*Cercis reniformis*, Engelmann;

Scheele in Roemer, Texas, 428.—Watson in Proc. Am. Acad. xvii, 348.

C. occidentalis, var. Gray in Jour. Boston Soc. Nat. Hist. vi, 177.—Walpers, Ann. ii, 440.—Torrey, Bot. Mex. Boundary Survey, 58.—Brewer & Watson, Bot. California, i, 161.

C. occidentalis, Gray, Hall's Pl. Texas, 7 [not Torrey].—Hemsley, Bot. Am.-Cent. i, 340, in part.

C. occidentalis, var. *Texensis*, Watson, Index, i, 209.

REDBUD.

Middle and western Texas west of the Colorado river; in northern Mexico.

A small tree, 6 to 8 meters in height, with a trunk 0.15 to 0.20 meter in diameter, or often a shrub forming dense thickets; limestone hills; formerly often confounded with the shrubby *C. occidentalis* of the California coast region.

Wood heavy, hard, close-grained, compact; layers of annual growth clearly marked by one to three rows of open ducts; medullary rays numerous, not conspicuous; color, brown streaked with yellow, the sap-wood lighter; specific gravity, 0.7513; ash, 0.77.

93.—*Prosopis juliflora*, De Candolle,

Prodr. ii, 447.—Descourtilz, Fl. Med. Antilles, viii, 107, t. 550.—Lindley, Fl. Med. 270.—Walpers, Rep. i, 861.—Bentham, Rev. Mim. in Trans. Linnæan Soc. xxx, 377.—Schnizlein, Icon. t. 277, f. 13.—Brewer & Watson, Bot. California, i, 163.—Kothrock in Wheeler's Rep. vi, 42, 107.—Hemsley, Bot. Am.-Cent. i, 344.

P. glandulosa, Torrey in Ann. Lyc. N. York, ii, 192, t. 2; Emory's Rep. 139; Pacific R. R. Rep. iv, 82.—Don, Miller's Dict. ii, 400.—Dietrich, Syn. ii, 1424.—Eaton & Wright, Bot. 376.—Walpers, Rep. i, 861.—Bentham in Hooker's Jour. Bot. iv, 348; London Jour. Bot. v, 81.—Grisebach, Fl. British West Indies, 217.—Watson in King's Rep. v, 420; Pl. Wheeler, 8.—Gray, Hall's Pl. Texas, 7.—Vasey, Cat. Forest Trees, 12.

Algarobia glandulosa, Torrey & Gray, Fl. N. America, i, 399; Pacific R. R. Rep. ii, 164.—Engelmann & Gray in Jour. Boston Soc. Nat. Hist. v, 242.—Engelmann in Wislizenus' Rep. 10.—Scheele in Roemer, Texas, 427.—Gray in Jour. Boston Soc. Nat. Hist. vi, 181; Smithsonian Contrib. iii, 60; v, 51; Mem. Am. Acad. new ser. v, 304; Ives' Rep. 11.—Torrey in Sitgreaves' Rep. 158; Pacific R. R. Rep. iv, 20, 82; vii, 10; Bot. Mex. Boundary Survey, 60.—Cooper in Smithsonian Rep. 1858, 259; Scientific Press, San Francisco, Nov. 1871, & f.—Palmer in Am. Nat. xii, 594.

P. odorata, Torrey in Fremont's Rep. 313, t. 1 (excl. fruit).

MESQUIT. ALGAROBÁ. HONEY LOCUST. HONEY POD.

Texas, valley of the Trinity river (Dallas, etc.) to the northern and western limits of the state; west through New Mexico and Arizona to the *mesas* west of the San Bernardino mountains, California, reaching southern Colorado, southern Utah (Saint George), and southern Nevada; southward through southern Mexico; in Jamaica.

A tree of the first economic value, sometimes 9 to 15 meters in height, with a trunk 0.90 meter in diameter, or much smaller, often reduced to a low shrub; on dry prairies and high rocky plains, or west of the Rocky mountains, along desert streams, here often forming open forests, and reaching its greatest development within the United States in the valley of the Santa Cruz and other streams of southern Arizona; in western Texas (Fort Stockton, etc.), on account of the annual burning of the prairies, rarely 1 meter in height, the roots then enormously developed, often weighing several hundred pounds, forming, as they are here locally known, "underground forests" and furnishing the best and cheapest fuel of the region.

Wood heavy, very hard, not strong, close-grained, compact, difficult to work, almost indestructible in contact with the soil, containing many evenly-distributed, rather large, open ducts; medullary rays numerous, distinct; color, rich dark brown or often red, the sap-wood clear yellow; specific gravity, 0.7652; ash, 2.18; of the root, specific gravity, 0.8493; ash, 3.02; exclusively used for the beams and underpinnings of the adobe houses of New Mexico, Arizona, and northern Mexico; for posts and fencing, and occasionally in the manufacture of furniture, the felloes of heavy wheels, etc.; the best and often the only fuel of the region, burning slowly with a clear flame, and producing valuable charcoal, but unsuited for the generation of steam on account of its destructive action upon boilers.

A gum resembling gum arabic is yielded by this species; the unripe and pulpy pods rich in grape sugar, edible, and furnishing valuable and important fodder.

94.—*Prosopis pubescens*, Benth.

London Jour. Bot. v, 82; Rev. Mim. in Trans. Linnæan Soc. xxx, 380.—Walpers, Ann. i, 259.—Watson in King's Rep. v, 420; Pl. Wheeler, 8.—Brewer & Watson, Bot. California, i, 163.—Rothrock in Wheeler's Rep. vi, 42, 107.—Hemsley, Bot. Am.-Cent. i, 344.

P. odorata, Torrey in Fremont's Rep. 313, t. 1 (for fruit).

P. Emoryi, Torrey in Emory's Rep. 139.

Strombocarpa pubescens, Gray in Smithsonian Contrib. iii, 60; v, 51; Ives' Rep. 9.—Torrey & Gray in Pacific R. R. Rep. ii, 163.—Torrey in Pacific R. R. Rep. iv, 11, 20, 82; v, 360, t. 4; vii, 10; Bot. Mex. Boundary Survey, 60.—Cooper in Smithsonian Rep. 1858, 259; Scientific Press, San Francisco, Nov. 1871 & f.—Vasey, Cat. Forest Trees, 12.

Strombocarpa odorata, Torrey in Sitgreaves' Rep. 158.

SCREW BEAN. SCREW-POD MESQUIT. TORNILLA.

Valley of the Rio Grande (Presidio), western Texas, westward through New Mexico and Arizona (valley of the Gila and Colorado rivers) to southern California (White Water, *Parish Brothers*, Vallecito, *Thurber*), and southward into Mexico; southern Utah (Saint George), and southern Nevada (Ash Meadows).

A small tree, rarely 9 meters in height, with a trunk sometimes 0.30 to 0.45 meter in diameter, or often a tall, much-branched shrub; sandy or gravelly bottom lands, reaching its greatest development within the United States in the valleys of the lower Colorado and Gila rivers.

Wood heavy, exceedingly hard, not strong, brittle, close-grained, compact, containing many evenly-distributed open ducts; medullary rays numerous, thin; color, light brown, the sap-wood somewhat lighter; specific gravity, 0.7609; ash, 0.95; used for fuel and fencing.

The pods used as fodder, and sometimes made into flour by the Indians.

95.—*Leucæna glauca*, Benth.

Hooker's London Jour. Bot. iv, 417; Rev. Mim. in Trans. Linnæan Soc. xxx, 443.—Walpers, Rep. i, 884.—Grisebach, Fl. British West Indies, 220.—Hemsley, Bot. Am.-Cent. i, 351.—Watson in Proc. Am. Acad. xvii, 350.—Chapman, Fl. S. States, Suppl. 619.

Mimosa glauca, Linnæus, Spec. 2 ed. 1504.

Acacia glauca, Willdenow, Spec. iv, 1075.—De Candolle, Prodr. ii, 467.

Acacia frondosa, Willdenow, Spec. iv, 1076.—De Candolle, Prodr. ii, 468.

Acacia biceps, Willdenow, Spec. iv, 1075.—De Candolle, Prodr. ii, 467.

Mimosa leucocephala, Lamarck, Dict. i, 12.

Acacia leucocephala, Link, Enum. Hort. Berl. ii, 444.—De Candolle, Prodr. ii, 467.

Mimosa biceps, Poiret, Suppl. i, 75.

Mimosa frondosa, Klein in Poiret, Suppl. i, 76.

Western Texas, San Saba to Devil's river (*Buckley*); southward into Mexico; semi-tropical Florida (introduced, *Curtiss*), and through the West Indies.

A small tree, 7 to 9 meters in height, with a trunk 0.10 to 0.15 meter in diameter, or often a tall or, in Florida, low shrub, sending up many stems from the ground.

Wood heavy, hard, close-grained, compact, containing many small, regularly-distributed open ducts; layers of annual growth and medullary rays hardly distinguishable; color, rich brown streaked with red, the sap-wood clear yellow; specific gravity, 0.9235; ash, 3.29.

96.—*Leucæna pulverulenta*, Bentham,

Hooker's London Jour. Bot. iv, 417; Rev. Mim. in Trans. Linnæan Soc. xxx, 443.—Hemsley, Bot. Am.-Cent. i, 351.

Acacia pulverulenta, Schlechtendal in Linnæa, xii, 571.

Acacia esculenta, Martens & Galeotti in Bull. Acad. Brux. x³, 312.

Southern Texas, valley of the lower Rio Grande; southward into Mexico.

A small tree, 6 to 8 meters in height, with a trunk 0.10 to 0.15 meter in diameter, often forming dense thickets; rich, sandy loam.

Wood heavy, hard, very close-grained, compact, containing many small, regularly-distributed open ducts; medullary rays very numerous, thin, conspicuous; color, rich dark brown, the sap-wood clear yellow; specific gravity, 0.6732; ash, 1.01.

97.—*Acacia Wrightii*, Bentham,

Smithsonian Contrib. iii, 64; Rev. Mim. in Trans. Linnæan Soc. xxx, 521.—Gray, Smithsonian Contrib. v, 53.—Walpers, Ann. iv, 626.—Torrey, Bot. Mex. Boundary Survey, 161.—Brewer & Watson, Bot. California, i, 61.—Watson in Proc. Am. Acad. xvii, 351.

OAT'S CLAW.

Western Texas, valley of the Guadalupe river (New Braunfels), westward and southward to the valley of the Rio Grande; in northern Mexico.

A small tree, rarely 9 meters in height, with a trunk sometimes exceeding 0.30 meter in diameter, or often a low, much-branched shrub.

Wood very heavy, hard, very close-grained, compact; layers of annual growth marked by one or two rows of small open ducts, and containing many scattered smaller ducts; medullary rays hardly distinguishable; color, bright, clear brown streaked with red and yellow, the sap-wood clear yellow; specific gravity, 0.9392; ash, 0.63.

98.—*Acacia Greggii*, Gray,

Smithsonian Contrib. iii, 65; v, 53; Ives' Rep. 11.—Torrey in Sitgreaves' Rep. 158; Pacific R. R. Rep. vii, 10; Bot. Mex. Boundary Survey, 61.—Walpers, Ann. iv, 625.—Bentham, Rev. Mim. in Trans. Linnæan Soc. xxx, 521.—Cooper in Smithsonian Rep. 1860, 442.—Brewer & Watson, Bot. California, i, 164.—Rothrock in Wheeler's Rep. vi, 108.—Hemsley, Bot. Am.-Cent. i, 353.—James in Am. Nat. xv, 981.

OAT'S CLAW.

Western Texas, valley of the Rio Grande, westward through southern New Mexico and Arizona to San Diego, California; southward into northern Mexico.

A low, much-branched tree, sometimes 9 meters in height, with a trunk rarely 0.45 meter in diameter, or often a shrub; dry *mesas* and in low cañons; common; the large specimens generally hollow and defective.

Wood heavy, exceedingly hard, strong, brittle, close-grained, compact; layers of annual growth marked by numerous rows of rather large open ducts; medullary rays numerous, thin; color, rich brown or red, the sap-wood light yellow; specific gravity, 0.8550; ash, 0.91; used for fuel.

A resinous gum resembling gum arabic is produced by this species (*Am. Jour. Pharm.* lii, 419).

99.—*Acacia Berlandieri*, Bentham,

London Jour. Bot. i, 522; Rev. Mim. in Trans. Linnæan Soc. xxx, 529.—Walpers, Rep. i, 919.—Dietrich, Syn. iv, 500.

A. tephroloba, Gray in Smithsonian Contrib. iii, 65; v, 54.—Walpers, Ann. iv, 625.—Torrey, Bot. Mex. Boundary Survey, 61.—Hemsley, Bot. Am.-Cent. i, 352.—Watson in Proc. Am. Acad. xvii, 351.

Southern Texas, valley of the Nueces (La Salle county) to Devil's river; southward into Mexico.

A small tree, sometimes 6 to 8 meters in height, with a trunk 0.15 to 0.20 meter in diameter, or more often a tall shrub, sending up many stems from the ground; the large specimens usually hollow and defective.

Wood not examined.

100.—*Lysiloma latisiliqua*, Benth.

Rev. Mim. in Trans. Linnæan Soc. xxx, 534.—Chapman, Fl. S. States, Suppl. 619.

Mimosa latisiliqua, Linnæus, Spec. 2 ed. 1504.

Acacia latisiliqua, Willdenow, Spec. iv, 1067.—Persoon, Syn. ii, 255.—De Candolle, Prodr. ii, 467.—Macfadyen, Fl. Jamaica, 318.—Nuttall, Sylva, ii, 34, t. 53; 2 ed. i, 183, t. 53.—Cooper in Smithsonian Rep. 1858, 264.

L. Bahamensis, Benth. in Hooker's London Jour. Bot. iii, 82.

Acacia Bahamensis, Grisebach, Fl. British West Indies, 221.

WILD TAMARIND.

Semi-tropical Florida, southern keys (Key Largo, Elliott's, Plantation, and Boca Chica Keys); through the West Indies.

A tree sometimes 15 meters in height, with a trunk 0.60 to 0.90 meter in diameter; bark of the young, vigorous trees smooth; the older trees generally decayed and defective, with rough, dark bark (*Curtiss*).

Wood heavy, hard, not strong, tough, close-grained, compact, susceptible of a fine polish, containing many scattered, open ducts; medullary rays numerous, not conspicuous; color, rich dark brown tinged with red, the sap-wood white; specific gravity, 0.6418; ash, 2.12; somewhat used locally in boat and ship-building, and considered equal to mahogany for this purpose.

101.—*Pithecolobium Unguis-cati*, Benth.

Hooker's London Jour. Bot. iii, 200; Rev. Mim. in Trans. Linnæan Soc. xxx, 572, 648.—Grisebach, Fl. British West Indies, 276.—Chapman, Fl. S. States, 116.—Vasey, Cat. Forest Trees, 13.

Mimosa Unguis-cati, Linnæus, Spec. 2 ed. 1497.—Jacquin, Hort. Schœnb. iii, 74, t. 392.—Descourtilz, Fl. Med. Antilles, i, t. 11.

Inga Unguis-cati, Willdenow, Spec. iv, 1006.—De Candolle, Prodr. ii, 436.—Nuttall, Sylva, ii, 37, t. 54; 2 ed. i, 86, t. 54.

Mimosa rosea, Vahl, Ecol. iii, 33, t. 25.

Inga rosea, Stendel in De Candolle, Prodr. ii, 437.

Inga forfex, Kunth, Mim. 12, t. 16.

P. forfex, Benth. in Hooker's London Jour. Bot. iii, 199.

Inga Guadalupensis, Desvaux, Jour. i, 70.

Mimosa Guadalupensis, Persoon, Syn. ii, 262.

Inga microphylla, Humboldt & Bonpland in Willdenow, Spec. iv, 1004.

P. microphyllum, Benth. in Hooker's London Jour. Bot. iii, 200.

P. Guadalupensis, Chapman, Fl. S. States, 116.

CAT'S CLAW.

Semi-tropical Florida, Caximbas bay, and on the southern keys; through the West Indies.

A small tree, sometimes 6 meters in height, with a trunk rarely exceeding 0.15 meter in diameter, or often throwing out many spreading, vine-like stems from the ground.

Wood very heavy, hard, close-grained, checking badly in drying; medullary rays numerous, inconspicuous; color, rich red varying to purple, sap-wood clear yellow; specific gravity, 0.9049; ash, 2.46.

ROSACEÆ.

102.—*Chrysobalanus Icaco*, Linnæus,

Spec. 1 ed. 513.—Jacquin, Stirp. Am. 154, t. 94.—Lamarck, Dict. iii, 224; Ill. ii, 542, t. 428.—Poiret, Suppl. iii, 135.—Aiton, Hort. Kew. 2 ed. iii, 200.—De Candolle, Prodr. ii, 525.—Lindley in Trans. Hort. Soc. London, v, 98.—Turpin, Dict. Sci. Nat. 236.—Tussac, Fl. Antilles, iv, 91, t. 31.—Spach, Hist. Veg. i, 369, t. 5, f. 4.—Torrey & Gray, Fl. N. America, i, 406.—Walpers, Rep. ii, 1; Ann. iv, 642.—Benth. Bot. Sulphur, 91; Fl. Nigritiana, 336.—Sprengel, Icon. t. 274, f. 1-13.—Cooper in Smithsonian Rep. 1860, 439.—Chapman, Fl. S. States, 119.—Grisebach, Fl. British West Indies, 229.—Baillon in Adansonia, vii, 221; Hist. Pl. i, 427, f. 486, 487.—Hooker f. in Martius, Fl. Brasil. ii, 7.—Guibourt, Hist. Drogues, 7 ed. iii, 287.—Hemsley, Bot. Am.-Cent. i, 365.

COCOA PLUM.

Semi-tropical Florida, cape Canaveral to bay Biscayne, west coast Caximbas bay, and on the southern keys; through the West Indies and tropical America to Brazil.

A small tree, 7 to 10 meters in height, with a trunk 0.15 to 0.30 meter in diameter, or along sandy beaches a low, prostrate shrub 1.08 to 2.16 meters in height; reaching its greatest development within the United States on the borders and islands of the Everglades, near bay Biscayne.

Wood heavy, hard, strong, close-grained, compact, containing few irregularly-distributed, not large, open ducts; medullary rays numerous, thin; color, light brown often tinged with red, the sap wood lighter; specific gravity, 0.7709; ash, 0.87.

Varieties are distinguished by *A. H. Curtiss* with the skin of the edible fruit white or black, the latter more ovate with narrower, softer stones (? var. *pellocarpa*, *Hooker f. l. c.*—*C. pellocarpa*, *Miquel, Prim. Esseq.* 193.—*Grisebach, l. c.*).

103.—*Prunus Americana*, Marshall,

Arbustum, iii.—Darlington in Ann. Lyc. N. York, iii, 87, t. 1; Fl. Cestrica, 3 ed. 72.—Eaton, Manual, 6 ed. 285.—Beck, Bot. 95.—Torrey & Gray, Fl. N. America, i, 407; Pacific R. R. Rep. ii, 164.—Eaton & Wright, Bot. 377.—Nuttall, Sylva, ii, 19, t. 48; 2 ed. i, 169, t. 48.—Torrey, Fl. N. York, i, 194; Emory's Rep. 408; Pacific R. R. Rep. iv, 82.—Emerson, Trees Massachusetts, 449; 2 ed. ii, 511.—Hooker in London Jour. Bot. vi, 217.—Römer, Syn. Mon. iii, 59.—Gray in Mem. Am. Acad. new ser. iv¹, 40; Manual N. States, 5 ed. 148.—Scheele in Römer, Texas, 430.—Richardson, Arctic Exped. 424.—Parry in Owen's Rep. 611.—Chapman, Fl. S. States, 119.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 56.—Lesquereux in Owen's 2d Rep. Arkansas, 358.—Wood, Cl. Book, 327; Bot. & Fl. 102.—Engelmann in Trans. Am. Phil. Soc. new ser. xiii, 190.—Koch, Dendrologie, i, 101.—Porter & Coulter, Fl. Colorado; Hayden's Surv. Misc. Pub. No. 4, 33.—Vasey, Cat. Forest Trees, 13.—Macoun in Geological Rep. Canada, 1875-'76, 194.—Broadhead in Coulter's Bot. Gazette, iii, 52.—Bell in Geological Rep. Canada, 1879-'80, 54^c.—Ridgway in Proc. U. S. Nat. Mus. 1882, 65.

P. Mississippi, Marshall, Arbustum, 112.

P. spinosa, Walter, Fl. Caroliniana, 146 [not Linnæus].

P. nigra, Aiton, Hort. Kew. ii, 165; 2 ed. iii, 198.—Willdenow, Spec. ii, 993; Berl. Baumz, 311.—Poiret in Lamarck, Dict. v, 674.—Persoon, Syn. ii, 35.—Bot. Mag. t. 1117.—Pursh, Fl. Am. Sept. i, 331.—Torrey, Fl. U. S. 469; Compend. Fl. N. States, 199.—Sprengel, Syst. ii, 477.—Römer, Syn. Mon. iii, 59.

Cerasus nigra, Loiseleur in Nouveau Duhamel, v, 32.—Seringe in De Candolle, Prodr. ii, 538.—Hooker, Fl. Bor.-Am. i, 167; Companion Bot. Mag. i, 24.—Don, Miller's Dict. ii, 513.—Beck, Bot. 96.—Spach, Hist. Veg. i, 399.—London, Arboretum, ii, 704, f. 411, 412.

P. hiemalis, Elliott, Sk. i, 542 [not Michaux].

P. coccinea, Rafinesque, Fl. Ludoviciana, 135.

WILD PLUM. CANADA PLUM. HORSE PLUM.

Valley of the Saint Lawrence (Quebec) to the valley of Rainy and Assinaboine rivers and southern shores of lake Manitoba; northern Vermont, western New England, and southward through the Atlantic states to the Chattahoochee region of western Florida, west to the valley of the upper Missouri river, Dakota, and Cheyenne cañon, Pike's Peak region, Colorado, southwest through Arkansas, the Indian territory, to about longitude 102°, and the valley of the lower Concho river, Texas.

A small tree, 6 to 12 meters in height, with a trunk rarely exceeding 0.30 meter in diameter; rich woods, or along streams and borders of ponds and swamps, reaching its greatest development on the bottom lands of eastern Texas.

A form with the young leaves and pedicles pubescent is—

var. *mollis*, Torrey & Gray, Fl. N. America, i, 407.

P. hiemalis, Michaux, Fl. Bor.-Am. i, 284.—Poiret in Lamarck, Dict. v, 679.—Persoon, Syn. ii, 35.—Desfontaines, Hist. Arb. ii, 206.—Nouveau Duhamel, v, 184.—Hayne, Dend. Fl. 73.—Sprengel, Syst. ii, 477.—Spach, Hist. Veg. i, 398.—Römer, Syn. Mon. iii, 59.

P. mollis, Torrey, Fl. U. S. 470; Compend. Fl. N. States, 199.—Beck, Bot. 95.

Cerasus hiemalis, Seringe in De Candolle, Prodr. ii, 538.—Hooker, Fl. Bor.-Am. i, 168.—Beck, Bot. 96.—London, Arboretum, ii, 704.—Don, Miller's Dict. ii, 504.

Cerasus Americana, Hooker, Companion Bot. Mag. i, 24.

Wood heavy, very hard, strong, very close-grained, compact, satiny, susceptible of a beautiful polish; medullary rays numerous, thin; color, rich bright brown or often red, the sap-wood lighter; specific gravity, 0.7215; ash, 0.18; used for the handles of tools, etc.

Often cultivated for the yellow, red, or rarely nearly black, acid or rarely sweet fruit, and furnishing an excellent stock on which to graft the varieties of the domestic plum.

104.—*Prunus angustifolia*, Marshall,

Arbustum, iii.—Koch, *Dendrologie*, i, 103.

P. Chicasa, Michaux, *Fl. Bor.-Am.* i, 284.—Poiret in *Lamarck, Dict.* v, 680.—Persoon, *Syn.* ii, 35.—Nuttall, *Genera*, i, 302.—Nouveau Duhamel, v, 183.—Elliott, *Sk.* i, 542.—Torrey in *Ann. Lyc. N. York*, ii, 194; *Pacific R. R. Rep.* iv, 82.—Sprengel, *Syst.* ii, 476.—Audubon, *Birds*, t. 53.—Eaton, *Manual*, 6 ed. 285.—Spach, *Hist. Veg.* i, 397.—Torrey & Gray, *Fl. N. America*, i, 407; *Pacific R. R. Rep.* ii, 164.—Eaton & Wright, *Bot.* 377.—Römer, *Syn. Mon.* iii, 58.—Darlington, *Fl. Cestrica*, 3 ed. 73.—Darby, *Bot. S. States*, 299.—Browne, *Trees of America*, 250.—Cooper in *Smithsonian Rep.* 1858, 251.—Chapman, *Fl. S. States*, 119.—Curtis in *Rep. Geological Surv. N. Carolina* 1860, ii, 56.—Lesquereux in *Owen's 2d Rep. Arkansas*, 858.—Wood, *Cl. Book*, 328; *Bot. & Fl.* 102.—Gray, *Manual N. States*, 5 ed. 148; *Hall's Pl. Texas*, 9.—Young, *Bot. Texas*, 1251.—Porter & Coulter, *Fl. Colorado*; *Hayden's Surv. Misc. Pub. No. 4*, 33.—Vasey, *Cat. Forest Trees*, 13.—Ridgway in *Proc. U. S. Nat. Mus.* 1882, 65.

P. insititia, Walter, *Fl. Caroliniana*, 146.—Abbot, *Insecta Georgia*, ii, t. 60.

Cerasus Chicasa, Seringe in *De Candolle, Prodr.* ii, 538.—Hooker, *Fl. Bor.-Am.* i, 168; *Companion Bot. Mag.* i, 24.—Don, *Miller's Dict.* ii, 514.—Loudon, *Arboretum*, ii, 705.

CHICKASAW PLUM. HOG PLUM.

Probably native of the eastern slopes of the southern Rocky mountains, where it is found at an altitude of 7,000 feet, and of the high plateau east and southeast of them; now widely naturalized by early cultivation throughout the Atlantic forests south of Pennsylvania, and west of the Alleghany mountains extending as far north as southern Michigan.

A small tree, 6 to 8 meters in height, with a trunk, 0.15 to 0.20 meter in diameter, or often a low shrub; generally along streams or borders of prairies, in rich soil.

Wood heavy, soft, not strong, close-grained, compact; medullary rays numerous, thin; color, light brown or red, the sap-wood lighter; specific gravity, 0.6884; ash, 0.28; often cultivated for its globose red or yellow fruit.

105.—*Prunus Pennsylvanica*, Linnæus f.

Suppl. 252.—Willdenow, *Spec.* ii, 992; *Enum.* 518; *Berl. Baumz.* 310.—Abbot, *Insecta Georgia*, i, t. 45.—Poiret in *Lamarck, Dict.* v, 673.—Persoon, *Syn.* ii, 35.—Nouveau Duhamel, v, 9.—Aiton, *Hort. Kew.* 2 ed. iii, 198.—Pursh, *Fl. Am. Sept.* i, 331.—Nuttall, *Genera*, i, 302.—Torrey, *Fl. U. S.* 468; *Compend. Fl. N. States*, 198.—Sprengel, *Syst.* ii, 477.—Hayne, *Dend. Fl.* 73.—Eaton, *Manual*, 6 ed., 235.—Beck in *Am. Journal Sci.* 1 ser. xiv, 112.—Dietrich, *Syn.* iii, 42.—Chapman, *Fl. S. States*, 130.—Curtis in *Rep. Geological Surv. N. Carolina*, 1860, iii, 57.—Wood, *Bot. & Fl.* 102.—Gray in *Proc. Philadelphia Acad.* 1863, 61; *Manual N. States*, 5 ed. 148.—Koch, *Dendrologie*, i, 117.—Porter & Coulter, *Fl. Colorado*; *Hayden's Surv. Misc. Pub. No. 4*, 33.—Emerson, *Trees Massachusetts*, 2 ed. ii, 513.—Vasey, *Cat. Forest Trees*, 13.—Macoun in *Geological Rep. Canada*, 1875-'76, 194.—Bell in *Geological Rep. Canada*, 1879-'80, 54.—Sears in *Bull. Essex Inst.* xiii, 176.

? *P. lanceolata*, Willdenow, *Berl. Baumz.* 240, t. 3, f. 3.

Cerasus borealis, Michaux, *Fl. Bor.-Am.* i, 286.—Nouveau Duhamel, v, 32.—Michaux f. *Hist. Arb. Am.* iii, 159, t. 8; *N. American Sylva*, 3 ed. ii, 152, t. 90.—Seringe in *De Candolle, Prodr.* ii, 558.—Don, *Miller's Dict.* ii, 513.—Beck, *Bot.* 97.—Loudon, *Arboretum*, ii, 703, f. 410.—Römer, *Syn. Mon.* iii, 78.

P. borealis, Poiret in *Lamarck, Dict.* v, 674.—Pursh, *Fl. Am. Sept.* i, 538.—Eaton, *Manual*, 54.—Barton, *Compend. Fl. Philadelph.* i, 223.—Nuttall, *Genera*, i, 302.—Loddiges, *Bot. Cab.* t. 1598.—Bigelow, *Fl. Boston.* 3 ed. 205.

? *P. persicifolia*, Desfontaines, *Hist. Arb.* ii, 205.

? *Cerasus persicifolia*, Loiseleur in *Nouveau Duhamel*, v, 9.—Seringe in *De Candolle, Prodr.* ii, 537.—Don, *Miller's Dict.* ii, 512.—Spach, *Hist. Veg.* i, 411.—Römer, *Syn. Mon.* iii, 81.

Cerasus Pennsylvanica, Seringe in *De Candolle, Prodr.* ii, 538.—Hooker, *Fl. Bor.-Am.* i, 168.—Don, *Miller's Dict.* ii, 514.—Beck, *Bot.* 97.—Torrey & Gray, *Fl. N. America*, i, 409.—Loudon, *Arboretum*, ii, 705.—Eaton & Wright, *Bot.* 189.—Torrey, *Fl. N. York*, i, 196.—Nuttall, *Sylva*, ii, 15; 2 ed. i, 165.—Browne, *Trees of America*, 265.—Emerson, *Trees Massachusetts*, 1 ed. 451.—Römer, *Syn. Mon.* iii, 57.—Gray, *Manual N. States*, 1 ed. 115.—Parry in *Owen's Rep.* 611.—Richardson, *Arctic Exped.* 425.—Cooper in *Smithsonian Rep.* 1858, 251.—Wood, *Cl. Book*, 327.

WILD RED CHERRY. PIN CHERRY. PIGEON CHERRY.

Labrador, shores of Hudson's bay, and west through the Saskatchewan region to the valley of the upper Fraser river (Soda creek, *Macoun*); south through the northern states to Pennsylvania, central Michigan, northern Illinois, central Iowa, and along the high Alleghany mountains of North Carolina and Tennessee, and the Rocky mountains of Colorado.

A small tree, rarely exceeding 12 meters in height, with a trunk sometimes 0.60 meter in diameter, or in the Rocky Mountain region reduced to a low shrub; common in all the northern forests, in northern New England taking possession of ground cleared by fire of the coniferous forests.

Wood light, soft, close-grained, compact; medullary rays numerous, thin; color, light brown, sap-wood clear yellow; specific gravity, 0.5023; ash, 0.40.

The small acid fruit used domestically and by herbalists in the preparation of cough mixtures, etc.

106.—*Prunus umbellata*, Elliott,

Sk. i, 541.—Eaton, Manual, 6 ed. 286.—Dietrich, Syn. iii, 44.—Chapman, Fl. S. States, 119.—Wood, Cl. Book, 328; Bot. & Fl. 102.—Young, Bot. Texas, 251.—Vasey, Cat. Forest Trees, 13.

P. pumila, Walter, Fl. Caroliniana, 146 [not Linnæus].

Cerasus umbellata, Torrey & Gray, Fl. N. America, i, 409.—Eaton & Wright, Bot. 190.—Rømer, Syn. Mon. iii, 78.

SLOE. BLACK SLOE.

South Carolina, south near the coast to Mosquito inlet and Tampa bay, Florida, and through central Alabama to eastern Mississippi (Holly Springs and Enterprise, *Mohr*).

A small tree, 5 to 6 meters in height, with a trunk 0.25 to 0.38 meter in diameter; dry, sandy soil.

Wood heavy, hard, close-grained, compact; medullary rays numerous, thin; color, dark reddish-brown, the sap-wood much lighter; specific gravity, 0.8202; ash, 0.12.

The black or red pleasantly acid fruit used as a preserve.

107.—*Prunus emarginata*, Walpers,

Rep. ii, 9.—Dietrich, Syn. iii, 42.—London, Arboretum, ii, 714.—Watson in King's Rep. v, 79.—Torrey, Bot. Wilkes Exped. 284.—Brewer & Watson, Bot. California, i, 167.

Cerasus emarginata, Douglas in Hooker, Fl. Bor.-Am. i, 169.—Don, Miller's Dict. ii, 515.—Torrey & Gray, Fl. N. America, i, 410.—Eaton & Wright, Bot. 189.—Rømer, Syn. Mon. iii, 79.—Torrey in Pacific R. R. Rep. iv, 83.—Bolander in Proc. California Acad. iii, 79.

Cerasus erecta, Presl, Epimel. Bot. 194.—Walpers, Ann, iii, 854.

Cerasus glandulosa, Kellogg in Proc. California Acad. i, 59.

Vancouver's island and the valley of the lower Fraser river, south through western Washington territory and Oregon, east to the western slopes of the Bitter Root mountain, Idaho (Lolo trail, *Watson*), and the valley of the Jocko river, Montana (*Canby & Sargent*). California along the western slopes of the Sierra Nevadas and on the Coast ranges, from San Francisco bay to the Santa Lucia mountains (*G. R. Vasey*), reaching an elevation of from 3,000 to 4,000 feet.

A tree often 12 to 15 meters in height, with a trunk sometimes exceeding 0.30 meter in diameter; at high elevations and throughout central California reduced to a shrub 2 to 3 meters in height, or in the Santa Lucia mountains 15 to 18 meters in height, with a trunk 0.60 to 0.90 meter in diameter (*Vasey*); generally along streams or in low, rich woods.

The wood of the type not collected.

Var. *mollis*, Brewer,

Bot. California, i, 167.—Hall in Coulter's Bot. Gazette, ii, 86.

Cerasus mollis, Douglas in Hooker, Fl. Bor.-Am. i, 169.—Hooker, London Jour. Bot. vi, 217.—Don, Miller's Dict. ii, 515.—Torrey & Gray, Fl. N. America, i, 410.—London, Arboretum, ii, 417.—Eaton & Wright, Bot. 189.—Nuttall, Sylva, ii, 14, t. 46; 2 ed. i, 164, t. 46.—Rømer, Syn. Mon. iii, 79.—Richardson, Arctic Exped. 425.—Newberry in Pacific R. R. Rep. vi, 73.—Cooper in Pacific R. R. Rep. xii, 29, 59; Am. Nat. iii, 406.—Lyal in Jour. Linnæan Soc. vii, 131.—Gray in Proc. Am. Acad. viii, 381.

P. mollis, Walpers, Rep. ii, 9.—Dietrich, Syn. iii, 42.—Torrey, Bot. Wilkes Exped. 284.—Vasey, Cat. Forest Trees, 13.—Macoun in Geological Rep. Canada, 1875-76, 194.

The common northern and Idaho form, more or less wooly pubescent, especially on the under side of the leaves.

Wood light, soft, not strong, brittle, close-grained, compact; medullary rays numerous, thin; color, brown streaked with green; specific gravity, 0.4502; ash, 0.21.

108.—*Prunus serotina*, Ehrhart,

Beitr. iii, 20.—Willdenow, Spec. ii, 986; Enum. 517; Berl. Baumz. 301.—Persoon, Syn. ii, 34.—Desfontaines, Hist. Arb. ii, 204.—Aiton, Hort. Kew. 2 ed. iii, 196.—Eaton, Manual, 54; 6 ed. 284.—Nuttall, Genera, i, 302.—Barton, Compend. Fl. Philadelph. 54.—Guimpel, Otto & Hayne, Abb. Holz. 45, t. 37.—Hayne, Dend. Fl. 70.—Sprengel, Syst. ii, 478.—Nees, Pl. Neuwied, 9.—Hooker f. in Trans. Linnæan Soc. xxii^a, 327.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 56.—Lesquereux in Owen's 2d Rep. Arkansas, 358.—Wood, Bot. & Fl. 102.—Engelmann in Trans. Am. Phil. Soc. new ser. xii, 190.—Chapman, Fl. S. States, 120.—Gray, Manual N. States, 5 ed. 149; Hall's Pl. Texas, 9.—Koch, Dendrologie, i, 122.—Torrey, Bot. Wilkes Exped. 284.—Emerson, Trees Massachusetts, 2 ed. ii, 515 & t.—Brewer & Watson, Bot. California, i, 167.—Vasey, Cat. Forest Trees, 13.—Bentley & Trimen, Med. Pl. ii, 97, t. 97.—Sears in Bull. Essex Inst. xiii, 176.—Bell in Geological Rep. Canada, 1879-'80, 54^c.—Ridgway in Proc. U. S. Nat. Mus. 1882, 66.

P. Virginiana, Miller, Dict. No. 3 [not Linnæus].—Du Roi, Obs. Bot. 12; Harbk. ii, 191.—Wangenheim, Amer. 34, t. 14.—Medicus, Bot. Beobacht. 1782, 345.—Marshall, Arbustum, 112.—Walter, Fl. Caroliniana, 146.—Aiton, Hort. Kew. ii, 163.—Poiret in Lamarck, Dict. v, 664.—Pursh, Fl. Am. Sept. i, 329.—Elliott, Sk. i, 540.—Torrey, Fl. U. S. 467; Compend. Fl. N. States, 189.—Bigelow, Fl. Boston. 3 ed. 204.

Cerasus Virginiana, Michaux, Fl. Bor.-Am. i, 285.—Michaux f. Hist. Arb. Am. iii, 151, t. 6; N. American Sylva, 3 ed. ii, 147, t. 88.—Hooker, Fl. Bor.-Am. i, 169 (excl. syn.).—Don, Miller's Dict. ii, 515.—Beck, Bot. 97.—Darlington, Fl. Cestrica, 2 ed. 289.—London, Arboretum, ii, 710, f. 418.—Browne, Trees of America, 268.

Cerasus serotina, Loiseleur in Nouveau Duhamel, v, 3.—Seringe in De Candolle, Prodr. ii, 540.—Spach, Hist. Veg. i, 416.—Torrey & Gray, Fl. N. America, i, 410.—London, Arboretum, ii, 712, f. 419 & t.—Eaton & Wright, Bot. 189.—Torrey, Fl. N. York, i, 196; Pacific R. R. Rep. vii, 11.—Penn. Cycl. vi, 432.—Carson, Med. Bot. i, 41, t. 35.—Griffith, Med. Bot. 238.—Emerson, Trees Massachusetts, 1 ed. 453.—Gray, Manual N. States, 1 ed. 115; Jour. Boston Soc. Nat. Hist. vi, 186.—Darlington, Fl. Cestrica, 3 ed. 75.—Darby, Bot. S. States, 299.—Cooper in Smithsonian Rep. 1858, 252.—Porcher, Resources S. Forests, 169.—Richardson, Arctic Exped. 425.—Wood, Cl. Book, 326.—Bolander in Proc. California Acad. iii, 79.

P. cartilaginea, Lehmann, Ind. Sem. Hamburg, 1833.

Padus serotina, Agardh, Theor. & Syst. Pl. t. 14, f. 8.

Padus Virginiana, Rømer, Syn. Mon. iii, 86.

Padus cartilaginea, Rømer, Syn. Mon. iii, 86.

WILD BLACK CHERRY. RUM CHERRY.

Southern Ontario, southward through the Atlantic forests to Matanzas inlet and Tampa bay, Florida, west to the valley of the Missouri river, Dakota, eastern Kansas, the Indian territory, and the valley of the upper San Antonio River, Texas.

A tree 18 to 30 meters in height, with a trunk 0.90 to 1.20 or, exceptionally, 1.50 meter in diameter; rich, generally elevated woodlands; common and reaching its greatest development on the western slopes of the Alleghany mountains from West Virginia southward; not common and of small size in the Gulf region and Texas.

Wood light, hard, strong, close, straight-grained, compact, easily worked; medullary rays numerous, thin; color, light brown or red, growing darker with exposure, the thin sap-wood yellow; specific gravity, 0.5822; ash, 0.15; largely used and esteemed in cabinet work, interior finish, etc., and now becoming scarce.

The bark contains a bitter tonic principle, and infused with cold water generates a small percentage of hydrocyanic acid; employed as a tonic and sedative in cases of pulmonary consumption in the form of cold infusions, sirups, and fluid extracts (*Proc. Am. Phar. Assoc.* xxiii, 209.—*Globley in Jour. Pharm. et Chimie*, xv, 40.—*Guibourt, Hist. Drogues*, 7 ed. iii, 317.—*Pharm. Jour.* 3 ser. iv, 44.—*Flückiger & Hanbury, Pharmacographia*, 224.—*U. S. Dispensatory*, 14 ed. 749.—*Nat. Dispensatory*, 2 ed. 1177); the bitter fruit used domestically in the preparation of cherry brandy.

NOTE.—The closely-allied *P. Virginiana* of the north Atlantic region, a tall shrub, sometimes 6 to 8 meters in height, does not assume arborescent habit.

109.—*Prunus Capuli*, Cavanilles,

Sprengel, Syst. ii, 477.—Schlechtendal in Linnæa, xiii, 69, 404.—Koch, Dendrologie, i, 123.—Hemslay, Bot. Am.-Cent. i, 367.—Watson in Proc. Am. Acad. xvii, 352.

Cerasus Capollin, De Candolle, Prodr. ii, 539.—Don, Miller's Dict. ii, 515.—London, Arboretum, ii, 713, f. 420.—Bentham, Pl. Hartweg. 10.—Lindley, Fl. Med. 232.—Penn. Cycl. vi, 432.—Torrey & Gray, Fl. N. America, i, 412.—Gray in Smithsonian Contrib. v, 54.

Cerasus Capuli, Seringe in De Candolle, Prodr. ii, 541.—Don, Miller's Dict. ii, 516.—Spach, Hist. Veg. i, 422.

P. Capollin, Zuccarini in Abhandl. Acad. Munich, ii, 345, t. 8.—Rømer, Syn. Mon. iii, 87.—Torrey, Bot. Mex. Boundary Survey, 62.—Rusby in Bull. Torrey Bot. Club, ix, 53.

P. Canadensis, Mocino & Sessé, Pl. Mex. Icon. ined.

WILD CHERRY.

Apache and Guadalupe mountains, Texas, west through southern New Mexico and Arizona to the southern slopes of the San Francisco mountains; southward through northern New Mexico, and in Peru.

A small tree, in the United States, rarely 12 meters in height, with a trunk often 0.30 meter in diameter; bottoms of cañons and mountain valleys, generally between 5,000 and 7,000 feet elevation.

Wood heavy, moderately hard, close-grained, compact; medullary rays very numerous, thin; color, brown, or often bright, clear red, the sap-wood nearly white; specific gravity, 0.7879; ash, 0.20.

110.—*Prunus demissa*, Walpers,

Rep. ii, 10.—Dietrich, Syn. iii, 43.—Bentham, Pl. Hartweg, 307.—Torrey, Bot. Mex. Boundary Survey, 63.—Watson in King's Rep. v, 80; Pl. Wheeler, 8.—Porter in Hayden's Rep. 1871, 481.—Coulter in Hayden's Rep. 1872, 764.—Rothrock, Pl. Wheeler, 37.—Brandege in Hayden's Rep. 1875, 236.—Brewer & Watson, Bot. California, i, 167.—Vasey, Cat. Forest Trees, 13.—Hall in Coulter's Bot. Gazette, ii, 86.—Macoun in Geological Rep. Canada, 1875-'76, 194.—Hemsley, Bot. Am.-Cent. i, 368.

Cerasus serotina, Hooker, Fl. Bor.-Am. i, 169, in part.

Cerasus demissa, Nuttall in Torrey & Gray, Fl. N. America, i, 411.—Gray in Mem. Am. Acad. new ser. iv¹, 40.—Durand in Jour. Philadelphia Acad. 1855, 87.—Torrey in Pacific R. R. Rep. iv, 83.—Newberry in Pacific R. R. Rep. vi, 73.—Cooper in Smithsonian Rep. 1858, 259; Pacific R. R. Rep. xii³, 59.

Padus demissa, Rømer, Syn. Mon. iii, 87.

P. Virginiana, var. *demissa*, Torrey, Bot. Wilkes Exped. 284.—Gray in Proc. Am. Acad. viii, 381.

WILD CHERRY.

Vancouver's island east to the western slopes of the Rocky mountains of Montana, south through the Pacific region; in Sonora.

A small tree, sometimes 7 to 10 meters in height, with a trunk 0.30 to 0.45 meter in diameter, or more often a low shrub; reaching its greatest development in the rich valleys of southern Oregon and northern California, near the coast; in southern California, and east of the Cascade and Sierra Nevada ranges, a low shrub confined to high, mountain valleys.

Wood heavy, hard, not strong, close-grained, compact; medullary rays numerous, conspicuous; color, light brown, the sap-wood lighter; specific gravity, 0.6951; ash, 0.50.

111.—*Prunus Caroliniana*, Aiton,

Hort. Kew. ii, 163; 2 ed. iii, 196.—Willdenow, Spec. ii, 987.—Poiret in Lamarck, Dict. v, 667.—Persoon, Syn. ii, 34.—Desfontaines, Hist. Arb. ii, 203.—Nuttall, Genera, i, 302.—Sprengel, Nene Entdeck. i, 304; Syst. ii, 478.—Hayne, Dend. Fl. 71.—Elliott, Sk. i, 540.—Audubon, Birds, t. 159, 190.—Eaton, Manual, 6 ed. 286.—Schlechtendal in Linnæa, xiii, 89.—Dietrich, Syn. iii, 43.—Chapman, Fl. S. States, 120.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 57.—Wood, Bot. & Fl. 103.—Koch, Dendrologie, i, 124.—Young, Bot. Texas, 252.—Gray, Hall's Pl. Texas, 9.—Vasey, Cat. Forest Trees, 13.

P. Carolina, Miller, Dict.—Du Roi, Harbk. ii, 196.

P. serratifolia, Marshall, Arbustum, 114.

P. Lusitanica, Walter, Fl. Caroliniana, 146.

Cerasus Caroliniana, Michaux, Fl. Bor.-Am. i, 285.—Nouveau Duhamel, v, 5.—Michaux f. Hist. Arb. Am. iii, 156, t. 7; N. American Sylva, 3 ed. ii, 150, t. 89.—Seringe in De Candolle, Prodr. ii, 540.—Don, Miller's Dict. ii, 516.—Spach, Hist. Veg. i, 420.—Penn. Cycl. vi, 432.—London, Arboretum, ii, 720, f. 423.—Torrey & Gray, Fl. N. America, i, 411.—Eaton & Wright, Bot. 190.—Browne, Trees of America, 272.—Darby, Bot. S. States, 299.—Griffith, Med. Bot. 291.—Cooper in Smithsonian Rep. 1858, 252.—Porcher, Resources S. Forests, 171.—Wood, Cl. Book, 326.

P. sempervirens, Willdenow, Enum. Suppl. 33.

? *Bumelia serrata*, Pursh, Fl. Am. Sept. 155.—Rømer & Schultes, Syst. iv, 498.

? *Achras serrata*, Poiret, Suppl. v, 36.

Leptocarpa Caroliniana, Nuttall, Sylva, ii, 18; 2 ed. i, 167.

Ohimanthus amygdalinus, Rafinesque, Fl. Ludoviciana, 159.

Laurocerasus Caroliniana, Rømer, Syn. Mon. iii, 90.

WILD ORANGE. MOCK ORANGE. WILD PEACH.

North Carolina, south, near the coast, to bay Biscayne, Florida, and southern Alabama, west, along the Gulf coast, to the valley of the Guadalupe river, Texas.

A small tree, evergreen, 10 to 12 meters in height, with a trunk rarely exceeding 0.30 meter in diameter; common and reaching its greatest development in the rich, light, deep soil of the bottoms of eastern Texas, here often covering extensive tracts known as "peach brakes"; not common in the eastern Gulf states.

Wood heavy, hard, strong, close-grained, checking badly in seasoning, susceptible of a good polish; medullary rays numerous, thin; color, light reddish-brown, or, more rarely, rich dark brown, the sap-wood lighter; specific gravity, 0.8688; ash, 0.41.

Generally planted in the southern states as an ornamental and hedge plant; foliage, bark, and fruit contain prussic acid, the leaves, especially when partly withered, often proving fatal to animals browsing upon them.

112.—*Prunus sphærocarpa*, Swartz,

Prodr. 81; Fl. Ind. Occ. ii, 927 [not Michaux].—Willdenow, Spec. ii, 987.—Poir. in Lamarck, Dict. v, 666.—Persoon, Syn. ii, 34.—Don, Miller's Dict. ii, 516.—Schlechtendal in Linnæa, xiii, 87.—Walpers, Rep. ii, 10.—Grisebach, Fl. British West Indies, 231.—Chapman, Fl. S. States, Suppl. 620.

Cerasus sphærocarpa, Loiseleur in Nouveau Duhamel, v, 4.—Seringe in De Candolle, Prodr. ii, 540.—London, Arboretum ii, 721.—Bot. Mag. t. 3141.—Spach, Hist. Veg. i, 421.

Semi-tropical Florida, western shores of bay Biscayne (*Curtiss*); in the West Indies.

A small tree, in Florida not exceeding 6 meters in height, with a trunk 0.10 to 0.15 meter in diameter; high rocky woods or, more rarely, along the borders of streams and ponds; rare.

Wood heavy, hard, close-grained, checking badly in drying, containing many very small open ducts; layers of annual growth and medullary rays obscure; color, light, clear red, the sap-wood pale yellow; specific gravity, 0.8998; ash, 0.87.

113.—*Prunus ilicifolia*, Walpers,

Rep. ii, 10.—Dietrich, Syn. iii, 43.—Torrey, Bot. Mex. Boundary Survey, 63; Bot. Wilkes Exped. 285.—Brewer & Watson, Bot. California, i, 168; ii, 443.—Vasey, Cat. Forest Trees, 13.

Cerasus ilicifolia, Nuttall in Hooker & Arnott, Bot. Beechey, 340, t. 83.—Torrey & Gray, Fl. N. America, i, 411.—Nuttall, Sylva, ii, 16, t. 47; 2 ed. i, 165, t. 47.—Torrey in Emory's Rep. 139; Pacific R. R. Rep. iv, 83.—Paxton, Brit. Fl. Garden, iii, 44, f. 254.—Walpers, Ann. iv, 654.—Cooper in Smithsonian Rep. 1858, 259.—Kellogg in Proc. California Acad. ii, 22.—Bolander in Proc. California Acad. iii, 79; iv, 22.—London Garden, 1873, 131 & fig.

Laurocerasus ilicifolia, Roemer, Syn. Mon. iii, 92.

ISLAY.

California, Coast ranges from San Francisco bay south to the southern boundary of the state, extending to the western slopes of the San Bernardino and San Jacinto mountains.

A small tree, evergreen, often 9 to 12 meters in height, with a trunk 0.30 to 0.60 meter in diameter, or when distant from the coast often reduced to a low shrub.

Wood very heavy, hard, strong, close-grained, checking in seasoning, satiny, susceptible of a beautiful polish, containing many regularly-distributed rather small open ducts; medullary rays numerous, thin; color, bright reddish-brown, the sap-wood much lighter; specific gravity, 0.9803; ash, 0.78; furnishing valuable fuel.

114.—*Vauquelinia Torreyi*, Watson,

Proc. Am. Acad. xi, 147.—Brewer & Watson, Bot. California, i, 169.—Maximowicz in Act. Hort. St. Petersburg, v⁸, 237.—Hemsley, Bot. Am.-Cent. i, 370.

Spiræa Californica, Torrey in Emory's Rep. 140.

V. corymbosa, Torrey, Bot. Mex. Boundary Survey, 64 [not Correa].

Arizona, high mountains near the Gila (*Emory*), summits of the Santa Catalina mountains (*Pringle*, *Lemmon*); in Sonora.

A small tree in the Santa Catalina mountains, 4 to 6 meters in height, with a trunk 0.10 to 0.20 meter in diameter; dry slopes and rocky bluffs at 2,700 to 4,000 feet elevation, granitic soil; generally hollow and decayed.

Wood very heavy, hard, very close-grained, compact, susceptible of a beautiful polish; medullary rays numerous, thin; color, rich dark brown streaked with red, the sap-wood yellow; specific gravity, 1.1374; ash, 1.45.

115.—*Cercocarpus ledifolius*, Nuttall;

Torrey & Gray, Fl. N. America, i, 427.—Hooker, Icon. t. 324.—Nuttall, Sylva, ii, 28, t. 51; 2 ed. i, 178, t. 51.—Walpers, Rep. ii, 46.—Dietrich, Syn. iii, 119.—Watson in King's Rep. v, 83, 420; Pl. Wheeler, 8.—Porter in Hayden's Rep. 1871, 481.—Coulter in Hayden's Rep. 1872, 765.—Parry in Am. Nat. ix, 201, 270; Proc. Davenport Acad. i, 146.—Engelmann in Simpson's Rep. 435.—Brewer & Watson, Bot. California, i, 174.—Vasey, Cat. Forest Trees, 13.—Sargent in Am. Jour. Sci. 3 ser. xvii, 421.—Rothrock in Wheeler's Rep. vi, 43, 111, 360.

MOUNTAIN MAHOGANY.

Cœur d'Alène mountains, Idaho, southward along the western slopes of the Rocky mountains of Montana and Wyoming; eastern extremities of the Blue mountains of Washington territory and Oregon, Wahsatch mountains, Utah, and west along the mountain ranges of the Great Basin to the western slope of the Sierra Nevada of California, extending southward into Arizona and New Mexico.

A small, low tree, rarely 12 meters in height, with a trunk sometimes 0.60 to 0.90 meter in diameter, or north of Utah and Nevada reduced to a low shrub; dry, rocky mountain slopes, between 6,000 and 8,000 feet elevation, reaching its greatest development on the high ranges of central Nevada.

A shrubby variety of the Wahsatch mountain and other ranges of Utah, characterized by its rigid, intricately branched growth, short, revolute leaves and smaller flowers and fruit, is—

var. *intricatus*, M. E. Jones in herb.

C. intricatus, Watson in Proc. Am. Acad. x, 346.—Parry in Am. Nat. ix, 270; Proc. Davenport Acad. i, 147.

C. brevifolius, Watson in King's Rep. v, 83 [not Gray].

Wood very heavy, hard, close-grained, compact, brittle, difficult to work, susceptible of a beautiful polish; medullary rays very numerous, thin; color, bright, clear red, or often rich dark brown, the sap-wood clear yellow; specific gravity, 1.0731; ash, 1.04; furnishing the most valuable fuel of the region, and largely manufactured into charcoal.

116.—*Cercocarpus parvifolius*, Nuttall;

Hooker & Arnott, Bot. Beechey, 337.—Torrey & Gray, Fl. N. America, i, 427; Pacific R. R. Rep. ii, 164.—Hooker, Icon. t. 323.—Walpers, Rep. ii, 45.—Torrey in Fremont's Rep. 89; Emory's Rep. 139; Sitgreaves' Rep. 158; Pacific R. R. Rep. iv, 83; Bot. Mex. Boundary Survey, 63; Bot. Wilkes Exped. 287.—Dietrich, Syn. iii, 119.—Gray in Mem. Am. Acad. new ser. iv¹, 41; Smithsonian Contrib. iii, 68; v, 54; Proc. Boston Soc. Nat. Hist. vii, 146; Am. Jour. Sci. 2 ser. xxxiii, 411; Proc. Philadelphia Acad. 1863, 61.—Engelmann in Trans. Am. Phil. Soc. new ser. xii, 190.—Bolander in Proc. California Acad. iii, 79.—Porter in Hayden's Rep. 1870, 475; 1871, 481.—Watson in King's Rep. v, 82.—Porter & Coulter, Fl. Colorado; Hayden's Surv. Misc. Pub. No. 4, 34.—Rothrock, Pl. Wheeler, 37; Wheeler's Rep. vi, 111, 359.—Brewer & Watson, Bot. California, i, 174; ii, 444.—Vasey, Cat. Forest Trees, 13.—M. E. Jones, Excur. Bot. 12, 15, 20, 21.—Hemslay, Bot. Am.-Cent. i, 374.—Watson in Proc. Am. Acad. xvii, 353.

MOUNTAIN MAHOGANY.

California, valley of the Klamath river, southward through the Coast ranges to the San Bernardino and San Jacinto mountains, and in Lower California; Rocky mountains of Wyoming, Colorado, and New Mexico, mountains of southern Arizona, and southward into Sonora.

A small tree, rarely 6 to 9 meters in height, with a trunk sometimes 0.30 meter in diameter, or more often a shrub; dry, gravelly soil, reaching its greatest development on the mountains of southern New Mexico and Arizona, at an elevation of 6,000 to 8,000 feet.

A glabrous variety of southern California, with dark green leaves, is—

var. *glaber*, Watson, Bot. California, i, 175.

C. betulæfolius, Nuttall in Hooker, Icon. t. 322.—Walpers, Rep. ii, 46.

C. betuloides, Nuttall in Torrey & Gray, Fl. N. America, i, 427.—Hooker in London Jour. Bot. vi, 218.

A form with small entire or sparingly toothed leaves, of northern Mexico, is—

var. *paucidentatus*, Watson in Proc. Am. Acad. xvii, 353.

Wood very heavy, hard, close-grained, compact, difficult to work, susceptible of a beautiful polish; medullary rays numerous, thin; color, bright reddish-brown, the sap-wood light brown; specific gravity, 0.9365; ash, 0.45; furnishing valuable fuel.

117.—*Pyrus coronaria*, Linnæus,

Spec. 1 ed. 480.—Kalm, Travels, English ed. ii, 166.—Du Roi, Harbk. i, 229.—Marshall, Arbustum, 118.—Aiton, Hort. Kew. ii, 176; 2 ed. iii, 209.—Willdenow, Spec. ii, 1019; Enum. 527; Berl. Baumz. 330.—Persoon, Syn. ii, 40.—Pursh, Fl. Am. Sept. i, 340.—Eaton, Manual, 56; 6 ed. 291.—Nuttall, Genera, i, 307.—Barton, Compend. Fl. Philadelph. i, 228.—Hayne, Dend. Fl. 86.—Torrey, Fl. U. S. i, 180; Compend. Fl. N. States, 203; Fl. N. York, i, 223.—Bot. Mag. t. 2009.—Elliott, Sk. i, 559.—Bot. Reg. viii, 651.—Sprengel, Syst. ii, 510.—De Candolle, Prodr. ii, 635.—Don, Miller's Dict. ii, 647.—Beck, Bot. 113.—Hooker, Companion Bot. Mag. i, 25.—Reichenbach, Fl. Exot. t. 240.—Torrey & Gray, Fl. N. America, i, 223.—Dietrich, Syn. iii, 154.—Loudon, Arboretum, ii, 908 & t.—Browne, Trees of America, 297.—Richardson, Arctic Exped. 428.—Perry in Owen's Rep. 612.—Darby, Bot. S. States, 307.—Cooper in Smithsonian Rep. 1858, 252.—Chapman, Fl. S. States, 128.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 69.—Lesquereux in Owen's 2d Rep. Arkansas, 359.—Wood, Cl. Book, 332; Bot. & Fl. 112.—Porcher, Resources S. Forests, 149.—Gray, Manual N. States, 5 ed. 161.—Koch, Dendrologie, i, 214.—Wenzig in Linnæa, xxxviii, 40 (excl. var.).—Macoun & Gibson in Trans. Bot. Soc. Edinburgh, xii, 325.—Vasey, Cat. Forest Trees, 13.—London Garden, xix, 400, t. 280.—Ward in Bull. U. S. Nat. Mus. No. 22, 78.—Ridgway in Proc. U. S. Nat. Mus. 1882, 66.

Malus coronaria, Miller, Dict. No. 2.—Moench, Meth. 682.—Michaux, Fl. Bor.-Am. i, 292.—Poiret in Lamarck, Dict. v, 562.—Desfontaines, Hist. Arb. ii, 140.—Nouveau Duhamel, vi, 139, t. 44, f. 1.—Michaux f. Hist. Arb. Am. iii, 65, t. 10; N. American Sylva, 3 ed. ii, 58, t. 65.—Barton, Prodr. Fl. Philadelph. 55.—Spach, Hist. Veg. ii, 136, t. 8.—Römer, Syn. Mon. iii, 191.—Decaisne in Nouv. Arch. Mus. x, 154.—Carrière in Rev. Hort. 1877, 410 & t.

Crataegus coronaria, Salisbury, Prodr. 357.

Malus microcarpa coronaria, Carrière in Rev. Hort. 1884, 104, f. 24.

AMERICAN CRAB. SWEET-SCENTED CRAB.

Ontario, valley of the Humber river, shores of lake Erie, southward through western New York and Pennsylvania to the District of Columbia, and along the Alleghany mountains to central Alabama and northern Mississippi; west to southern Minnesota, Iowa, eastern Kansas, the Indian territory, and northern Louisiana.

A small tree, rarely 6 to 9 meters in height, with a trunk often 0.30 meter in diameter; rich, rather low woods, reaching its greatest development in the valleys of the lower Ohio region.

Wood heavy, rather soft, not strong, very close-grained, checking badly in drying; medullary rays numerous, obscure; color, brown varying to light red, the sap-wood yellow; specific gravity, 0.7048; ash, 0.52; used for levers, handles of tools, and in turnery.

Often planted for ornament on account of its fragrant blossoms; the small, yellow-green austere fruit used for preserves, and occasionally made into cider.

118.—*Pyrus angustifolia*, Aiton,

Hort. Kew. ii, 176; 2 ed. iii, 209.—Willdenow, Spec. ii, 1020.—Poiret in Lamarck, Dict. v, 455.—Persoon, Syn. ii, 40.—Pursh, Fl. Am. Sept. i, 341.—Elliott, Sk. i, 559.—Torrey, Fl. U. S. 480; Compend. Fl. N. States, 203.—Sprengel, Syst. ii, 509.—De Candolle, Prodr. ii, 635.—Watson, Dend. Brit. ii, t. 132.—Bot. Reg. xiv, 1207.—Don, Miller's Dict. 647.—Beck, Bot. 113.—Hooker, Companion Bot. Mag. i, 25.—Torrey & Gray, Fl. N. America, i, 471.—Loudon, Arboretum, ii, 909 & t.—Eaton & Wright, Bot. 382.—Dietrich, Syn. iii, 154.—Nuttall, Sylva, ii, 24; 2 ed. i, 174.—Darby, Bot. S. States, 307.—Cooper in Smithsonian Rep. 1858, 252.—Chapman, Fl. S. States, 128.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 69.—Lesquereux in Owen's 2d Rep. Arkansas, 359.—Wood, Cl. Book, 333; Bot. & Fl. 112.—Gray, Manual N. States, 5 ed. 161.—Koch, Dendrologie, i, 213.—Vasey, Cat. Forest Trees, 14.—Ridgway in Proc. U. S. Nat. Mus. 1882, 66.

P. coronaria, Wangenheim, Amer. 61, t. 21, f. 47 [not Linnæus].—Walter, Fl. Caroliniana, 148.

Malus angustifolia, Michaux, Fl. Bor.-Am. i, 292.—Decaisne in Nouv. Arch. Mus. x, 155.

Malus sempervirens, Desfontaines, Hist. Arb. ii, 141.—Nouveau Duhamel, vi, 638, t. 43, f. 1.—Poiret, Suppl. iv, 524.—Spach, Hist. Veg. ii, 135, t. 8, figs.—Römer, Syn. Mon. iii, 491.

P. coronaria, var. *angustifolia*, Wenzig in Linnæa, xxxviii, 41.

Ohloromeles sempervirens, Decaisne in Fl. des Serres, xxiii, 126.

AMERICAN CRAB APPLE. SOUTHERN CRAB APPLE.

Pennsylvania †, southern Delaware, and the valley of the lower Wabash river, Illinois, south to the Chattahoochee region of western Florida.

A small tree, 6 to 9 meters in height, with a trunk rarely 0.30 meter in diameter; low, rich woods; most common and reaching its greatest development along the river bottoms of the south Atlantic states; less common west of the Alleghany mountains.

Wood heavy, hard, close-grained, checking badly in drying; medullary rays numerous, obscure; color, light brown tinged with red, the sap-wood yellow; specific gravity, 0.6895; ash, 0.33; used for levers, handles of tools, etc.

The austere fruit used for preserves and made into cider.

119.—*Pyrus rivularis*, Douglas;

Hooker, Fl. Bor.-Am. i, 203, t. 68.—Don, Miller's Dict. ii, 647.—Torrey & Gray, Fl. N. America, i, 471.—Eaton & Wright, Bot. 383.—Walpers, Rep. ii, 53.—Dietrich, Syn. iii, 154.—Ledebour, Fl. Rossica, ii, 99.—Nuttall, Sylva, ii, 22, t. 49; 2 ed. i, 172, t. 49.—Richardson, Arctic Exped. 428.—Torrey in Pacific R. R. Rep. iv, 85; Bot. Wilkes Exped. 292.—Newberry in Pacific R. R. Rep. vi, 73.—Cooper in Smithsonian Rep. 1858, 259; Pacific R. R. Rep. xii, 29, 60.—Rothrock in Smithsonian Rep. 1867, 435, 446.—Koch, Dendrologie, i, 212.—Gray in Proc. Am. Acad. viii, 382.—Wenzig in Linnæa, xxxviii, 38.—Brewer & Watson, Bot. California, i, 188.—Vasey, Cat. Forest Trees, 14.—Hall in Coulter's Bot. Gazette, ii, 67.—Macoun in Geological Rep. Canada, 1875-'76, 195.—Dawson in Canadian Nat. new ser. ix, 330.

P. diversifolia, Bongard in Mem. Acad. Sci. St. Petersburg, 6 ser. ii, 133.

P. fusca, Rafinesque, Med. Bot. ii, 254.

P. subcordata, Ledebour, Fl. Rossica, ii, 95.

Malus rivularis, Rømer, Syn. Mon. iii, 215.—Decaisne in Nouv. Arch. Mus. x, 155.

Malus diversifolia, Rømer, Syn. Mon. iii, 215.—Decaisne in Nouv. Arch. Mus. x, 155.

Malus subcordata, Rømer, Syn. Mon. iii, 192.

OREGON ORAB APPLE.

Coast of Alaska, southward along the coast and islands of British Columbia, through Washington territory and Oregon, west of the Cascade mountains, to Sonoma county, California.

A small tree, sometimes 9 meters in height, with a trunk 0.30 to 0.45 meter in diameter; rich, low woods, generally along streams, often forming dense thickets.

Wood heavy, hard, very close-grained, liable to check badly in drying, susceptible of a beautiful polish; medullary rays numerous, obscure; color, light brown tinged with red, the sap-wood lighter; specific gravity, 0.8316; ash, 0.41; used for mallets, mauls, bearings of machinery, etc.

The small, black, pleasantly acid fruit occasionally used as a preserve, and prized by the Indians as food.

120.—*Pyrus Americana*, De Candolle,

Prodr. ii, 637.—Watson, Dend. Brit. i, t. 54.—Sprengel, Syst. ii, 511.—Hooker, Fl. Bor.-Am. i, 204.—Don, Miller's Dict. ii, 648.—Beck, Bot. 113.—Audubon, Birds, t. 363.—Torrey & Gray, Fl. N. America, i, 472.—London, Arboretum, iii, 920 & t.—Eaton & Wright, Bot. 383.—Torrey, Fl. N. York, i, 224.—Dietrich, Syn. iii, 155.—Nuttall, Sylva, ii, 25, t. 50; 2 ed. i, 175, t. 50.—Browne, Trees of America, 326.—Emerson, Trees Massachusetts, 439; 2 ed. ii, 499.—Parry in Owen's Rep. 612.—Richardson, Arctic Exped. 428.—Lange, Pl. Grœnl. 134.—Cooper in Smithsonian Rep. 1858, 252.—Chapman, Fl. S. States, 129.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 70.—Wood, Cl. Book, 333; Bot. & Fl. 112.—Porcher, Resources S. Forests, 168.—Gray, Manual N. States, 5 ed. 161.—Koch, Dendrologie, i, 190.—Brewer & Watson, Bot. California, i, 189.—Vasey, Cat. Forest Trees, 14.—Macoun in Geological Rep. Canada, 1875-'76, 195.—Sears in Bull. Essex Inst. xiii, 176.—Bell in Geological Rep. Canada, 1879-'80, 54°.

Sorbus Americana, Marshall, Arbustum, 145.—Willdenow, Enum. 520.—Pursh, Fl. Am. Sept. i, 341.—Poiret, Suppl. v, 164.—Eaton, Manual, 55; 6 ed. 351.—Nuttall, Genera, i, 305.—Hayne, Dend. Fl. 75.—Torrey, Fl. U. S. 477; Compend. Fl. N. States, 202.—Spach, Hist. Veg. ii, 95.—Bigelow, Fl. Boston. 3 ed. 207.—Rømer, Syn. Mon. iii, 138.—Maximowicz in Bull. Acad. St. Petersburg, xix, 174.—Wenzig in Linnæa, xxxviii, 71.—Decaisne in Nouv. Arch. Mus. x, 158.

Sorbus aucuparia, Poiret in Lamarck, Dict. vii, 234, in part.—Bigelow, Fl. Boston. 1. ed. 119.—Decaisne in Nouv. Arch. Mus. x, 158, in part.

Sorbus aucuparia, var. *Americana*, Persoon, Syn. ii, 38 & addend.

P. aucuparia, Meyer, Pl. Labrador, 81, in part.—Schlechtendal in Linnæa, x, 99.—Hooker f. in Trans. Linnæan Soc. xxii^a, 290, 327, in part.

Sorbus humifusa, Rafinesque, Med. Bot. ii, 265.

MOUNTAIN ASH.

Greenland?, Labrador, Newfoundland, Anticosti island, and westward along the southern shore of James' bay to the valley of the Nelson river (White Mud falls), southward through all mountainous regions of the northeastern states, and along the high mountains of Virginia and North Carolina; in northern Michigan, Wisconsin, and Minnesota.

A small tree, 6 to 9 meters in height, with a trunk 0.30 to 0.45 meter in diameter; borders of swamps and in moist, rocky woods, reaching its greatest development on the northern shores of lakes Huron and Superior.

A form with smaller fruit, peculiar to the high southern Alleghany mountains, is—

var. *microcarpa*, Torrey & Gray, Fl. N. America, i, 472.

Sorbus aucuparia, var. *a.* Michaux, Fl. Bor.-Am. i, 290.

Sorbus microcarpa, Pursh, Fl. Am. Sept. i, 341.—Poiret, Suppl. v, 164.—Elliott, Sk. i, 555.—Torrey, Fl. U. S. 477.—Eaton, Manual, 6 ed. 351.—Spach, Hist. Veg. ii, 95.—Römer, Syn. Mon. iii, 138.

P. microcarpa, Sprengel, Syst. ii, 511.—De Candolle, Prodr. ii, 636.—Don, Miller's Dict. ii, 648.—Beck, Bot. 113.—Eaton & Wright, Bot. 383.—Loudon, Arboretum, ii, 921.

Sorbus Americana, var. *microcarpa*, Wenzig in Linnæa, xxxviii, 71.

Sorbus riparia, Rafinesque, New Sylva, 15.

Wood light, soft, close-grained, compact; medullary rays numerous, obscure; color, light brown, the sap-wood lighter; specific gravity, 0.5451; ash, 0.83.

Often planted for ornament.

121.—*Pyrus sambucifolia*,

Chamisso & Schlechtendal in Linnæa, ii, 36.—Bongard in Mem. Acad. Sci. St. Petersburg, 6 ser. ii, 133.—Don, Miller's Dict. ii, 648.—Torrey & Gray, Fl. N. America, i, 472.—Walpers, Rep. ii, 53.—Dietrich, Syn. iii, 155.—Ledebour, Fl. Rossica, ii, 99.—Trautvetter & Meyer, Fl. Ochot. 37.—Maximowicz, Prim. Fl. Amurensis, 103.—Rothrock in Smithsonian Rep. 1867, 446.—Gray, Manual N. States, 5 ed. 161; Proc. Am. Acad. viii, 382.—Porter in Hayden's Rep. 1870, 475.—Watson in King's Rep. v, 92.—Porter & Coulter, Fl. Colorado; Hayden's Surv. Misc. Pub. No. 4, 38.—Brewer & Watson, Bot. California, i, 189.—Macoun in Geological Rep. Canada, 1875-'76, 195.—Hall in Coulter's Bot. Gazette, ii, 87.—G. M. Dawson in Canadian Nat. new ser. ix, 10.—Sears in Bull. Essex Inst. xiii, 176.

Sorbus aucuparia, var. *β.* Michaux, Fl. Bor.-Am. i, 290.

Sorbus aucuparia, Schrank, Pl. Labrador, 25, in part [not Linnæus].

P. Americana, Newberry in Pacific R. R. Rep. vi, 73 [not De Candolle].—Cooper in Pacific R. R. Rep. xii^a, 60.—Torrey, Bot. Wilkes Exped. 292.

P. aucuparia, Meyer, Pl. Labrador, 81, in part.—Schlechtendal in Linnæa, x, 99, in part.—Hooker in Trans. Linnæan Soc. xxii^a, 290, 327, in part.

Sorbus sambucifolia, Römer, Syn. Mon. iii, 139.—Maximowicz in Bull. Acad. Sci. St. Petersburg, xix, 174.—Wenzig in Linnæa, xxxviii, 73.—Decaisne in Nouv. Arch. Mus. x, 159.

Sorbus Sitchensis, Römer, Syn. Mon. iii, 139.

MOUNTAIN ASH.

Labrador to northern New England and the shores of lake Superior; high mountain ranges of the Pacific region from Alaska to southern New Mexico; in Kamtchatka.

A small tree, 9 to 12 meters in height, with a trunk sometimes 0.30 meter in diameter, or in the Pacific forests generally reduced to a low shrub; cold, wet swamps or borders of streams, reaching its greatest development in northern New England and Minnesota.

Wood light, soft, weak, close-grained, compact; medullary rays numerous, obscure; color, light brown, the sap-wood nearly white; specific gravity, 0.5928; ash, 0.35.

The bark and unripe fruit of the American mountain ashes, like those of the nearly-allied *P. aucuparia* of Europe, are extremely astringent, and occasionally used, domestically, in infusions, decoctions, etc., in the treatment of diarrhea (*Nat. Dispensatory*, 2 ed. 1333).

122.—*Crataegus rivularis*, Nuttall;

Torrey & Gray, Fl. N. America, i, 464.—Dietrich, Syn. iii, 161.—Walpers, Rep. ii, 58.—Nuttall, Sylva, ii, 9; 2 ed. i, 160.—Cooper in Smithsonian Rep. 1858, 258; Am. Nat. iii, 407.—Regel in Act. Hort. St. Petersburg, i, 107.—Watson in King's Rep. v, 92.—Porter in Hayden's Rep. 1871, 482.—Coulter in Hayden's Rep. 1872, 765.—Brandegge in Hayden's Rep. 1875, 236.—Vasey, Cat. Forest Trees, 14.—Macoun in Geological Rep. Canada, 1875-'76, 195.—Engelmann in Coulter's Bot. Gazette, vii, 128.

C. sanguinea, var. *Douglasii*, Coulter in Hayden's Rep. 1872, 765 [not Torrey & Gray].

British Columbia, south through eastern Oregon and Washington territory, east and southeast along the mountain ranges of Idaho, Montana, Utah, and Colorado, to the Pinos Altos mountains, New Mexico (*Greene*).

A small tree, 6 to 8 meters in height, with a trunk rarely exceeding 0.30 meter in diameter, or often a tall, much-branched shrub, forming dense, impenetrable thickets along borders of streams and swamps.

Wood heavy, hard, close-grained, compact; medullary rays numerous, thin; color, bright reddish-brown, the sap-wood nearly white; specific gravity, 0.7703; ash, 0.35.

123.—*Cratægus Douglasii*, Lindley,

Bot. Reg. xxi, t. 1810.—London, Arboretum, ii, 823, f. 584 & t.—Koch, Dendrologie, i, 147.—Kaleniczenko in Bull. Soc. Imp. Nat. Moscow, xlviii, 26.—Brewer & Watson, Bot. California, i, 189.—Macoun in Geological Rep. Canada, 1875-76, 195.—Engelmann in Coulter's Bot. Gazette, vii, 128.

? *C. glandulosa*, Pursh, Fl. Am. Sept. i, 337, in part.

C. punctata, var. *brevispina*, Douglas in Hooker, Fl. Bor.-Am. i, 201.

C. sanguinea, var. *Douglasii*, Torrey & Gray, Fl. N. America, i, 464.—Walpers, Rep. ii, 58.—Dietrich, Syn. iii, 160.—Torrey, Bot. Wilkes Exped. 292.—Regel in Act. Hort. St. Petersburg, i, 116.

C. sanguinea, Nuttall, Sylva, ii, 6, t. 44; 2 ed. i, 157, t. 44 [not Pallas].—Cooper in Smithsonian Rep. 1858, 259; Am. Nat. iii, 407.—Vasey, Cat. Forest Trees, 14.

Anthomeles Douglasii, Rømer, Syn. Mon. iii, 140.

C. rivularis, Brewer & Watson, Bot. California, i, 189 [not Nuttall].

British Columbia, valley of the Parsnip river, in about latitude 55° N., south through Washington territory and Oregon to the valley of the Pitt river, California, extending east through Idaho and Montana to the western base of the Rocky mountains (valley of the Flathead river, *Canby & Sargent*).

A small tree, sometimes 12 meters in height, with a trunk 0.30 to 0.45 meter in diameter, or often a tall shrub throwing up many stems from the ground and forming impenetrable thickets; rather wet, sandy soil along streams, and reaching its greatest development in the valleys west of the Cascade mountains; toward its eastern limits a low shrub.

Wood heavy, hard, tough, close-grained, compact, satiny, susceptible of a beautiful polish; medullary rays numerous, thin; color, nearly white tinged with rose, the sap-wood lighter; specific gravity, 0.6950; ash, 0.33; used for wedges, mauls, etc.

The small, sweet, black fruit, ripening in August, is largely collected by the Indians.

124.—*Cratægus brachyacantha*, Sargent & Englemann;

Engelmann in Coulter's Bot. Gazette, vii, 128.

HOGS' HAW.

New Orleans?, (*Drummond* in herb. *Gray*); Minden, Louisiana (*Mohr*); Concord, Texas (*Sargent*); Longview, Texas (in fruit, *Letterman*).

A tree 9 to 12 meters in height, with a trunk sometimes 0.60 meter in diameter; borders of streams in low, very rich soil; the largest North American representative of the genus.

Wood heavy, hard, very close-grained, compact, susceptible of a beautiful polish; medullary rays numerous, very obscure; color, light brown tinged with rose, the sap-wood lighter; specific gravity, 0.6793; ash, 0.42.

The large blue-black fruit greedily eaten by hogs and other animals.

125.—*Cratægus arborescens*, Elliott,

Sk. i, 550.—Eaton, Manual, 6 ed. 112.—Torrey & Gray, Fl. N. America, i, 466.—Eaton & Wright, Bot. 212.—Dietrich, Syn. iii, 160.—Walpers, Rep. ii, 58.—Nuttall, Sylva, ii, 10, t. 45; 2 ed. i, 160, t. 45.—Darby, Bot. S. States, 306.—Cooper in Smithsonian Rep. 1858, 252.—Chapman, Fl. S. States, 127.—Wood, Cl. Book, 331; Bot. & Fl. 111.—Young, Fl. Texas, 259.—Vasey, Cat. Forest Trees, 14.—Engelmann in Bull. Torrey Bot. Club, ix, 4.

Phænopyrum arborescens, Rømer, Syn. Mon. iii, 153.

C. Crus-galli, var. *pyracanthifolia*, Regel in Act. Hort. St. Petersburg, i, 109, in part.

Valley of the Savannah river, South Carolina (*Aiken, Ravenel*), south to the Chattahoochee region of western Florida; valley of the Mississippi river, near Saint Louis (*Engelmann*), south and southwest to western Louisiana, and the valley of the lower Colorado river, Texas.

A small tree, 6 to 9 meters in height, with a trunk sometimes 0.45 to 0.60 meter in diameter; borders of streams and in rather low, wet swamps.

Wood heavy, hard, not strong, close-grained, compact, susceptible of a beautiful polish; medullary rays very numerous, obscure; color, light brown tinged with red, the sap-wood lighter; specific gravity, 0.6491; ash, 0.57.

The small globular fruit bright red or, more rarely, orange.

126.—*Crataegus Crus-galli*, Linnaeus,

Spec. 1 ed. 476.—Kalm, Travels, English ed. i, 115.—Medicus, Bot. Beobacht. ii, 344.—Walter, Fl. Caroliniana, 147.—Aiton, Hort. Kew. ii, 170; 2 ed. iii, 202.—Willdenow, Spec. ii, 1004.—Micaux, Fl. Bor.-Am. i, 298.—Persoon, Syn. ii, 37.—Pursh, Fl. Am. Sept. i, 338.—Eaton, Manual, 55; 6 ed. 111.—Nuttall, Genera, i, 305.—Barton, Compend. Fl. Philadelph. i, 225; Prodr. Fl. Philadelph. 54.—Elliott, Sk. i, 548.—Torrey, Fl. U. S. 476; Compend. Fl. N. States, 202; Fl. N. York, i, 221.—Watson, Dend. Brit. i, t. 56.—De Candolle, Prodr. ii, 626.—Hooker, Fl. Bor.-Am. i, 260; Companion Bot. Mag. i, 25.—Don, Miller's Dict. ii, 598.—Beck, Bot. 111.—Torrey & Gray, Fl. N. America, i, 463.—London, Arboretum, ii, 820, f. 574, 575 & t.—Eaton & Wright, Bot. 212.—Bigelow, Fl. Boston. 3 ed. 206.—Dietrich, Syn. iii, 158.—Browne, Trees of America, 278.—Emerson, Trees Massachusetts, 433; 2 ed. ii, 492 & t.—Römer, Syn. Mon. iii, 117.—Parry in Owen's Rep. 612.—Darlington, Fl. Cestrica, 3 ed. 83.—Darby, Bot. S. States, 305.—Cooper in Smithsonian Rep. 1858, 252.—Chapman, Fl. S. States, 127.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 83.—Lesquereux in Owen's 2d Rep. Arkansas, 359.—Wood, Cl. Book, 331; Bot. & Fl. 111.—Porcher, Resources S. Forests, 148.—Gray, Manual N. States, 5 ed. 160; Hall's Pl. Texas, 9.—Young, Bot. Texas, 258.—Regel in Act. Hort. St. Petersburg, i, 108.—Kaleniczenko in Bull. Soc. Imp. Nat. Moscow, xlviii, 19.—Vasey, Cat. Forest Trees, 14.—Bell in Geological Rep. Canada, 1879-'80, 54c.—Ridgway in Proc. U. S. Nat. Mus. 1882, 66.

C. lucida, Du Roi, Obs. Bot. 13.—Wangenheim, Amer. 53, t. 17, f. 42.—Sprengel, Syst. ii, 506.—De Candolle, Prodr. ii, 629.—Eaton, Manual, 6 ed. 112.—Don, Miller's Dict. ii, 599.—Eaton & Wright, Bot. 212.

Mespilus Crus-galli, Marshall, Arbustum, 88.—Lamarck, Dict. iv, 441.—Desfontaines, Hist. Arb. ii, 157.—Nouveau Duhamel, iv, 149.—Willdenow, Enum. 522; Berl. Baumz. 244.—Hayne, Dend. Fl. 80.—Koch, Dendrologie, i, 142.

? *Mespilus cuneiformis*, Marshall, Arbustum, 88.

Mespilus lucida, Ehrhart, Beitr. iv, 17.—Mönch, Meth. 685.—Spach, Hist. Veg. ii, 57.

Mespilus cuneifolia, Mönch, Meth. 684.

C. Crus-galli, var. *splendens*, Aiton, Hort. Kew. 2 ed. iii, 202.

Mespilus Watsoniana, Spach, Hist. Veg. ii, 57.

C. Watsoniana, Römer, Syn. Mon. iii, 117.

COCKSPUR THORN. NEWCASTLE THORN.

Valley of the Saint Lawrence river, west through southern Ontario to Manitoba, south through the Atlantic forests to the valley of the Chipola river, western Florida, and the valley of the Colorado river, Texas.

A small tree, 4 to 10 meters in height, with a trunk sometimes 0.30 meter in diameter, running into various forms. The best marked are—

var. *pyracanthifolia*, Aiton, Hort. Kew. ii, 170; 2 ed. iii, 202.—De Candolle, Prodr. ii, 626.—Torrey & Gray, Fl. N. America, i, 464.—London, Arboretum, ii, 820, t. 128, f. 580.—Browne, Trees of America, 278.—Regel in Act. Hort. St. Petersburg, i, 109, in part.

C. salicifolia, Medicus, Bot. Beobacht. ii, 345.—Römer, Syn. Mon. iii, 117.

C. Crus-galli, var. *salicifolia*, Aiton, l. c.; 2 ed. l. c.—Willdenow, Berl. Baumz. 244.—De Candolle, l. c.—London, l. c. f. 551-553, 578 & t.—Browne, l. c.—Regel, l. c. 110.

Mespilus Crus-galli, var. *salicifolia*, Hayne, Dend. Fl. 80.

Mespilus Crus-galli, var. *pyracanthifolia*, Hayne, l. c.

Mespilus salicifolia, Koch, Dendrologie, i, 144.

C. Coursetiana, Römer, Syn. Mon. iii, 117.

var. *ovalifolia*, Lindley, Bot. Reg. xxii, t. 1860.—Torrey & Gray, Fl. N. America, i, 464.—Dietrich, Syn. iii, 159.—London, Arboretum, ii, 821, f. 579 & t.—Regel in Act. Hort. St. Petersburg, i, 109.

Mespilus ovalifolia, Hornemann, Hort. Hafn. Suppl. 52.—Koch, Dendrologie, i, 143.

Mespilus prunellifolia, Poiret, Suppl. iv, 72.

C. ovalifolia, De Candolle, Prodr. ii, 627.—Don, Miller's Dict. ii, 598.—Römer, Syn. Mon. iii, 117.

C. prunellifolia, De Candolle, l. c.—Don, l. c.—Römer, l. c.

Mespilus elliptica, Guimpel, Otto & Hayne, Abb. Holz. 170, t. 144 [not Lamarck].—Spach, Hist. Veg. ii, 68.

var. *linearis*, De Candolle, Prodr. ii, 626.—Torrey & Gray, Fl. N. America, i, 464.—Dietrich, Syn. iii, 159.—London, Arboretum, ii, 821, f. 577.—Browne, Trees of America, 278.—Regel in Act. Hort. St. Petersburg, i, 110.

Mespilus lucida, var. *angustifolia*, Ehrhart, Beitr. iv, 18.

O. linearis, Persoon, Syn. ii, 37.—Rømer, Syn. Mon. iii, 118.

Mespilus linearis, Desfontaines, Hist. Arb. ii, 156.—Poiret, Suppl. iv, 70.—Spach, Hist. Veg. ii, 57.

var. *prunifolia*, Torrey & Gray, Fl. N. America, i, 464.—Dietrich, Syn. iii, 159.—London, Arboretum, ii, 821, f. 576 & t.—Regel in Act. Hort. St. Petersburg, i, 110.

Mespilus prunifolia, † Marshall, Arbustum, 90.—Lamarck, Dict. iv, 443.—Nouveau Duhamel, iv, 150, t. 40.—Sprengel, Syst. ii, 506.

Mespilus rotundifolia, Ehrhart, Beitr. iii, 20.

C. prunifolia, Persoon, Syn. ii, 37.—Bosc in De Candolle, Prodr. ii, 627.—Don, Miller's Dict. ii, 598.—Lindley, Bot. Reg. xxii, t. 1868.—Eaton, Manual, 6 ed. 112.—Eaton & Wright, Bot. 212.

Mespilus Bosciana, Spach, Hist. Veg. ii, 58.

C. Bosciana, Rømer, Syn. Mon. iii, 118.

Wood heavy, hard, not strong, close-grained, compact, satiny, susceptible of a fine polish; medullary rays numerous, very obscure; color, brown tinged with red, the sap-wood rather lighter; specific gravity, 0.7194; ash, 0.56.

The long, strong spines are occasionally collected and used to fasten sacks and for similar purposes.

127.—*Cratægus coccinea*, Linnaeus,

Spec. 1 ed. 476.—Walter, Fl. Caroliniana, 147.—Aiton, Hort. Kew. ii, 167; 2 ed. iii, 200.—Willdenow, Spec. ii, 1000 (excl. syn.).—Michaux, Fl. Bor.-Am. i, 288.—Persoon, Syn. ii, 36.—Pursh, Fl. Am. Sept. i, 337.—Eaton, Manual, 55; 6 ed. 111.—Nuttall, Genera, i, 305.—Schrunk, Fl. Labrador, 26.—Barton, Compend, Fl. Philadelph. i, 226.—Hayne, Dend. Fl. 77.—Elliott, Sk. i, 553.—Torrey, Fl. U. S. 474; Compend. Bot. N. States, 201; Fl. N. York, i, 221; Emory's Rep. 408.—De Candolle, Prodr. ii, 627.—Hooker, Fl. Bor.-Am. i, 201; Bot. Mag. t. 3432.—Don, Miller's Dict. ii, 599.—Meyer, Fl. Labrador, 82.—Beck, Bot. 112.—Lindley, Bot. Reg. 23, t. 1957.—Torrey & Gray, Fl. N. America, i, 465.—Bigelow, Fl. Boston. 3 ed. 206.—Eaton & Wright, Bot. 211.—Dietrich, Syn. iii, 160.—Walpers, Rep. ii, 58.—London, Arboretum, ii, 816, f. 564-566, t. 121.—Schnizlein, Icon. t. 270, f. 18-20, 22.—Emerson, Trees Massachusetts, 434; 2 ed. ii, 493 & t.—Richardson, Arctic Exped. 427.—Darlington, Fl. Cestrica, 3 ed. 83.—Darby, Bot. S. States, 305.—Cooper in Smithsonian Rep. 1858, 252.—Gray in Pacific R. R. Rep. xii, 43; Manual N. States, 5 ed. 160.—Chapman, Fl. S. States, 127.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 82.—Lesquereux in Owen's 2d Rep. Arkansas, 309.—Wood, Cl. Book, 331; Bot. & Fl. 111.—Kaleniczenko in Bull. Soc. Imp. Nat. Moscow, xlviii, 9.—Vasey, Cat. Forest Trees, 14.—Sears in Bull. Essex Inst. xiii, 177.—Bell in Geological Rep. Canada, 1879-'80, 55c.—Ridgway in Proc. U. S. Nat. Mus. 1882, 66.

Mespilus coccinea, Marshall, Arbustum, 87.—Mönch, Meth. 684.—Lamarck, Dict. iv, 442.—Desfontaines, Hist. Arb. ii, 156.—Willdenow, Enum. 523; Berl. Baumz. 238.—Wendland in Regensb. Fl. 1823, 699.—Sprengel, Syst. ii, 507.—Spach, Hist. Veg. ii, 64.

Mespilus rotundifolia, Ehrhart, Beitr. iii, 20.—Wendland in Regensb. Fl. 1823, 700.—Watson, Dend. Brit. i, t. 58.—Koch, Dendrologie, i, 148.

Pyrus glandulosa, Mönch, Meth. 680.

C. glandulosa, Willdenow, Spec. ii, 1002 (excl. syn.).—Pursh, Fl. Am. Sept. i, 337, in part.—Torrey, Fl. U. S. 475; Compend. Fl. N. States, 201.—De Candolle, Prodr. ii, 627.—Loddiges, Bot. Cab. t. 1012.—Hooker, Fl. Bor.-Am. i, 201.—Don, Miller's Dict. ii, 599.—Eaton, Manual, 6 ed. 111.—Beck, Bot. 112.—Eaton & Wright, Bot. 211.—London, Arboretum, ii, 817, f. 550, 567, 568 & t.—Richardson, Arctic Exped. 427.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 84.—Regel in Act. Hort. St. Petersburg, i, 120.

Mespilus glandulosa, Willdenow, Enum. 523.—Sprengel, Syst. ii, 507.—Spach, Hist. Veg. ii, 62.—Koch, Dendrologie, i, 145.

Mespilus pubescens, Wendland in Regensb. Fl. 1823, 700.

C. Crus-galli, Bigelow, Fl. Boston. 2 ed. 194 [not Linnaeus].

† *Mespilus Wendlandii*, Opiz in Regensb. Fl. 1834, 590.

C. macracantha, Loddiges in London, Arboretum, ii, 819, f. 572, 573 & t.

C. glandulosa, var. *macracantha*, Lindley in Bot. Reg. xxii, t. 1912.

Mespilus flabellata, Spach, Hist. Veg. ii, 63.—Koch, Dendrologie, i, 148.

Halmia flabellata, Rømer, Syn. Mon. iii, 136.

Anthomeles rotundifolia, Rømer, Syn. Mon. iii, 140.

Phænopyrum coccineum, Rømer, Syn. Mon. iii, 156.

Phænopyrum Wendlandii, Rømer, Syn. Mon. iii, 156.

SCARLET HAW. RED HAW. WHITE THORN.

West coast of Newfoundland, west along the valley of the Saint Lawrence river and the northern shores of the great lakes to Manitoba, south through the Atlantic forests to northern Florida and eastern Texas.

A small tree, sometimes 9 meters in height, with a trunk 0.30 meter in diameter; open upland woods or along streams and borders of prairies; very common at the north, rare at the south; running into many forms, varying in the size and shape of the leaves, size of the fruit, etc. The best marked are—

var. *viridis*, Torrey & Gray, Fl. N. America, i, 465.—Torrey in Nicolle's Rep. 149.

O. viridis, Linnaeus, Spec. 1 ed. 476.—Willdenow, Spec. ii, 1001.—Persoon, Syn. ii, 36.—Elliott, Sk. i, 551.—De Caudolle, Prodr. ii, 630.—Don, Miller's Dict. ii, 601.—Eaton, Manual, 6 ed. 112.—Darlington, Fl. Cestrica, 2 ed. 293.—Eaton & Wright, Bot. 212.—Beck, Bot. 305.—Darby, Bot. S. States, 305.—Wood, Cl. Book, 332; Bot. & Fl. 111.

? *Phænopyrum viride*, Rømer, Syn. Mon. iii, 156.

Mespilus viridis, Koch, Dendrologie, i, 149.

O. glandulosa, var. *rotundifolia*, Regel in Act. Hort. St. Petersburg, i, 120.

var. *populifolia*, Torrey & Gray, Fl. N. America, i, 465.

O. populifolia, Elliott, Sk. i, 553 [not Walter].—Nuttall, Genera, i, 305.—Eaton, Manual, 6 ed. 112.—Beck, Bot. 305.—Eaton & Wright, Bot. 212.—Darby, Bot. S. States, 305.

Mespilus populifolia, Lamarck, Dict. iv, 447.

Phænopyrum populifolium, Rømer, Syn. Mon. iii, 156.

O. coccinea, var. *typica*, Regel in Act. Hort. St. Petersburg, i, 121.

var. *oligandra*, Torrey & Gray, Fl. N. America, i, 465.

Wood heavy, hard, close-grained, compact; medullary rays thin, very obscure; color, brown tinged with red, the sap-wood a little lighter; specific gravity, 0.8618; ash, 0.38.

128.—*Crataegus subvillosa*, Schrader,

Ind. Sem. Hort. Gœtt.—Torrey in Pacific R. R. Rep. iv, 35.—Ridgway in Proc. U. S. Nat. Mus. 1882, 66.

O. coccinea, var. *mollis*, Torrey & Gray, Fl. N. America, i, 465.—Gray in Jour. Boston Soc. Nat. Hist. vi, 186.—Parry in Owen's Rep. 612.—Regel in Act. Hort. St. Petersburg, i, 121.

Phænopyrum subvillosum, Rømer, Syn. Mon. iii, 154.

O. mollis, Scheele in Linnaea, xxi, 569; Rømer, Texas, Appx. 473.—Walpers, Ann. ii, 523.

O. sanguinea, var. *villosa*, Ruprecht & Maximowicz, Prim. Fl. Amurensis, 101.

O. Texana, Buckley in Proc. Philadelphia Acad. 1861, 454 (see Gray in same, 1862, 163).—Young, Fl. Texas, 258.

O. tomentosa, var. *mollis*, Gray, Manual N. States, 5 ed. 160.—Wood, Cl. Book, 330; Bot. & Fl. 121.—Vasey, Cat. Forest Trees, 14.

Mespilus tiliæfolia, Koch, Dendrologie, i, 151.

SCARLET HAW.

Eastern Massachusetts (possibly introduced); central Michigan to eastern Nebraska, south to middle Tennessee, and southwest through Missouri, Arkansas, the Indian territory, and Texas to the valley of the San Antonio river.

A small tree, 7 to 9 meters in height, with a trunk rarely 0.45 meter in diameter; rich woods and along borders of streams and prairies.

Wood heavy, hard, not strong, close-grained, compact; medullary rays numerous, very obscure; color, light brown or light red, the sap-wood lighter; specific gravity, 0.7953; ash, 0.69.

The large red fruit often downy, edible, and of agreeable flavor.

129.—*Crataegus tomentosa*, Linnæus,

Spec. 1 ed. 476 (excl. syn. Gronovius).—Kalm, Travels, English ed. ii, 151.—Du Roi, Harbk. i, 183.—Torrey & Gray, Fl. N. America, i, 466.—Dietrich, Syn. iii, 160.—Torrey, Fl. N. York, i, 222.—Emerson, Trees Massachusetts, 1 ed. 435; 2 ed. ii, 494 & t.—Parry in Owen's Rep. 612.—Cooper in Smithsonian Rep. 1858, 252.—Chapman, Fl. S. States, 127.—Lesquereux in Owen's 2d Rep. Arkansas, 359.—Wood, Cl. Book, 330.—Engelmann in Trans. Am. Phil. Soc. new ser. xii, 191.—Gray, Manual N. States, 5 ed. 160.—Young, Bot. Texas, 258.—Vasey, Cat. Forest Trees, 14.—Macoun in Geological Rep. Canada, 1875-'76, 195.—Ridgway in Proc. U. S. Nat. Mus. 1882, 66.

C. leucophlæos, Mönch, Hort. Weiss. 31, t. 2.—Regel in Act. Hort. St. Petersburg, i, 106.

Mespilus Calpodendron, Ehrhart, Beitr. ii, 67.

C. pyrifolia, Aiton, Hort. Kew. ii, 168; 2 ed. iii, 200.—Willdenow, Spec. ii, 1001.—Persoon, Syn. ii, 36.—Nouveau Duhamel, iv, 131.—Poiret, Suppl. i, 292.—Pursh, Fl. Am. Sept. i, 337.—Nuttall, Genera, i, 305.—Elliott, Sk. i, 550.—Torrey, Fl. U. S. 475; Compend. Fl. N. States, 201.—De Candolle, Prodr. ii, 627.—Hooker, Fl. Bor.-Am. i, 201.—Don, Miller's Dict. ii, 899.—Eaton, Manual, 6 ed. 111.—Lindl'y, Bot. Reg. xxii, t. 1877.—London, Arboretum, ii, 819, f. 571 & t.—Eaton & Wright, Bot. 211.

* *Mespilus latifolia*, Lamarck, Dict. iv, 444.—Desfontaines, Hist. Arb. ii, 156.—Nouveau Duhamel, iv, 150.—Spach, Hist. Veg. ii, 60.

C. latifolia, Persoon, Syn. ii, 36.—Don, Miller's Dict. ii, 598.—Eaton, Manual, 6 ed. 112.—Eaton & Wright, Bot. 212.—Rømer, Syn. Mon. 119.

Mespilus pyrifolia, Willdenow, Enum. 523; Berl. Baumz. 240.—Kaleniczenko in Bull. Soc. Imp. Nat. Moscow, xlvi, 15.—Sprengel, Syst. ii, 507.—Hayne, Dend. Fl. 78.

Mespilus lobata, Poiret, Suppl. iv, 71.

Mespilus odorata, Wendland in Regensb. Fl. 1823, 700.

Mespilus pruinosa, Wendland in Regensb. Fl. 1823, 700.

C. lobata, Bosc in De Candolle, Prodr. ii, 628.

C. flava, Hooker, Fl. Bor.-Am. i, 202 (excl. syn.).

Halmia tomentosa, Rømer, Syn. Mon. 135.

Halmia lobata, Rømer, Syn. Mon. 135.

Phænopyrum pruinosa, Rømer, Syn. Mon. 155.

? *C. coccinea*, var. *viridis*, Torrey in Pacific R. R. Rep. iv, 86 [not Torrey & Gray].

C. tomentosa, var. *pyrifolia*, Gray, Manual N. States, 5 ed. 160.—Wood, Bot. & Fl. 111.

C. coccinea, Brandegee in Hayden's Rep. 1875, 236 [not Linnæus].

C. leucocephalus, Lavallée, Arboretum Segrez. 78, t. 22 [not Mönch].

C. coccinea, var. *cordata*, Lavallée, Arboretum Segrez. 81, t. 22.

BLACK THORN. PEAR HAW.

New Brunswick, westward along the valley of the Saint Lawrence river and the northern shores of the great lakes to the Saskatchewan region, southward through the Atlantic forests to the Chattahoochee region of western Florida, and eastern Texas west to the mountains of eastern Washington territory and Oregon, southwestern Colorado, and southwestern New Mexico.

A small tree, 6 to 9 meters in height, with a trunk rarely 0.45 meter in diameter, or often, especially west of the Rocky mountains, reduced to a low shrub, here forming dense thickets along mountain streams; the most widely-distributed of the North American *Cratægi*, varying greatly in the size, shape, and color of the fruit, form of the leaves, amount of pubescence, etc.

Wood heavy, hard, not strong, close-grained, compact; medullary rays numerous, thin; color, bright reddish-brown, the sap-wood lighter; specific gravity, 0.7633; ash, 0.50.

Var. *punctata*, Gray,

Manual N. States, 2 ed. 124.—Cooper in Smithsonian Rep. 1858, 252.—Chapman, Fl. S. States, 127.—Porter in Hayden's Rep. 1871, 481.—Vasey, Cat. Forest Trees, 14.

O. punctata, Jacquin, Hort. Vindob. i, 10, t. 28.—Aiton, Hort. Kew. ii, 169; 2 ed. iii, 202.—Willdenow, Spec. ii, 1004.—Michaux, Fl. Bor.-Am. i, 289.—Persoon, Syn. i, 37.—Pursh, Fl. Am. Sept. i, 338.—Elliott, Sk. i, 548.—Torrey, Fl. U. S. 476; Compend. Fl. N. States, 202; Fl. N. York, i, 222.—De Candolle, Prodr. ii, 627.—Hooker, Fl. Bor.-Am. i, 201 (excl. var.); Companion Bot. Mag. i, 25.—Don, Miller's Dict. ii, 589.—Eaton, Manual, 6 ed. 111.—Beck, Bot. 111.—Torrey & Gray, Fl. N. America, i, 466.—London, Arboretum, ii, 818, f. 569, 570 & t.—Eaton & Wright, Bot. 211.—Dietrich, Syn. iii, 159.—Browne, Trees of America, 277.—Emerson, Trees Massachusetts, 435; 2 ed. ii, 495.—Gray, Manual N. States, 1 ed. 128.—Richardson, Arctic Exped. 427.—Darlington, Fl. Cestrica, 3 ed. 84.—Darby, Bot. S. States, 306.—Lesquereux in Owen's 2d Rep. Arkansas, 359.—Wood, Cl. Book, 330; Bot. & Fl. 111.—Engelmann in Trans. Am. Phil. Soc. new ser. xii, 191.—Kaleniczenko in Bull. Soc. Imp. Nat. Moscow, xlviii, 14.

Mespilus cornifolia, Muenchhausen, Hausv. v, 145.—Lamarck, Dict. iv, 444.—Koch, Dendrologie, i, 134.—Spach, Hist. Veg. ii, 60, t. 10, f. c.

C. Crus-galli, Wangenheim, Amer. 52.—Du Roi, Harbk. i, 195 [not Linnaeus].

Mespilus cuneifolia, Ehrhart, Beitr. iii, 21.—Sprengel, Syst. ii, 506.—Spach, Hist. Veg. ii, 61.

Mespilus punctata, Loiseleur in Nouveau Duhamel, iv, 152.—Willdenow, Enum. 524; Berl. Baumz. 243.—Poiret, Suppl. iv, 70.—Hayne, Dend. Fl. 79.—Watson, Dend. Brit. i, t. 57.—Spach, Hist. Veg. ii, 61.—Wenzig in Linnaea, xxxviii, 128.

Mespilus pyrifolia, Desfontaines, Hist. Arb. ii, 155.

C. punctata, var. *rubra* and *aurea*, Aiton, Hort. Kew. 2 ed. iii, 202.

C. latifolia, De Candolle, Prodr. ii, 627.

? *C. flexuosa*, Schweinitz in Long's 2d Exped. ii, Appx. 112.

C. flava, Darlington, Fl. Cestrica, 2 ed. 292 [not Aiton].

C. cuneifolia, Rømer, Syn. Mon. iii, 118.

C. obovatifolia, Rømer, Syn. Mon. iii, 120.

Halmia punctata, Rømer, Syn. Mon. iii, 134.

Halmia cornifolia, Rømer, Syn. Mon. iii, 134.

C. tomentosa, var. *plicata*, Wood, Cl. Book, 330; Bot. & Fl. 111.

C. punctata, var. *xanthocarpa*, Lavallée, Arboretum Segrez. i, 53, t. 16.

Fruit larger than that of the species, dull red or yellow.

130.—*Crataegus cordata*, Aiton,

Hort. Kew. ii, 168; 2 ed. iii, 200.—Willdenow, Spec. ii, 1000.—Persoon, Syn. ii, 36.—Eaton, Manual, 55; 6 ed. 111.—Elliott, Sk. i, 554.—Torrey, Fl. U. S. 474; Compend. Fl. N. States, 201.—De Candolle, Prodr. ii, 628.—Watson, Dend. Brit. i, t. 63.—Lindley, Bot. Reg. xiv, t. 1151.—Hooker, Fl. Bor.-Am. i, 201.—Don, Miller's Dict. ii, 599.—Beck, Bot. 112.—Torrey & Gray, Fl. N. America, i, 467.—London, Arboretum, ii, 825 & t.—Eaton & Wright, Bot. 211.—Dietrich, Syn. iii, 160.—Browne, Trees of America, 280.—Richardson, Arctic Exped. 427.—Darlington, Fl. Cestrica, 3 ed. 83.—Darby, Bot. S. States, 306.—Cooper in Smithsonian Rep. 1858, 252.—Chapman, Fl. S. States, 127.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 82.—Wood, Cl. Book, 331; Bot. & Fl. 111.—Gray, Manual N. States, 5 ed. 159.—Young, Bot. Texas, 257.—Regel in Act. Hort. St. Petersburg, i, 114.—Kaleniczenko in Bull. Soc. Imp. Nat. Moscow, xlviii, 31.—Vasey, Cat. Forest Trees, 14.

Mespilus Phænopyrum, Ehrhart in Linnaeus f. Suppl. 254; Beitr. i, 181; ii, 67.—Mönch, Meth. 685.—Lamarck, Dict. iv, 446.

C. populifolia, Walter, Fl. Caroliniana, 147 [not Elliott].—Pursh, Fl. Am. Sept. i, 337.

Mespilus acerifolia, Burgsdorf in Lamarck, Dict. iv, 442.—Nouveau Duhamel, iv, 151.—Spach, Hist. Veg. ii, 65.

Mespilus cordata, Miller, Icon. t. 179.—Willdenow, Enum. 523; Berl. Baumz. 239.—Hayne, Dend. Fl. 77.—Sprengel, Syst. ii, 507.—Koch, Dendrologie, i, 138.

Phænopyrum cordatum, Rømer, Syn. Mon. iii, 157.

Phænopyrum acerifolium, Rømer, Syn. Mon. iii, 157.

WASHINGTON THORN.

Valley of the upper Potomac river, Virginia, southward along the Alleghany mountains to northern Georgia and Alabama, extending west through eastern and middle Kentucky and Tennessee to the valley of the lower Wabash river, Illinois.

A small tree, 6 to 8 meters in height, with a trunk rarely 0.30 meter in diameter; generally along banks of streams.

Wood heavy, hard, close-grained, compact; medullary rays numerous, obscure; color, brown tinged with red, the sap-wood lighter; specific gravity, 0.7293; ash, 0.46.

Formerly widely planted as a hedge plant.

131.—*Cratægus apiifolia*, Michaux.

Fl. Bor.-Am. i, 287.—Persoon, Syn. ii, 38.—Pursh, Fl. Am. Sept. i, 336.—Nuttall, Genera, i, 305.—Elliott, Sk. i, 552.—De Candolle, Prodr. ii, 627.—Don, Miller's Dict. ii, 599.—Audubon, Birds, t. 192.—Eaton, Manual, 6 ed. 112.—Hooker, Companion Bot. Mag. i, 25.—Torrey & Gray, Fl. N. America, i, 467.—London, Arboretum, ii, 824, f. 588, 589 & t.—Eaton & Wright, Bot. 212.—Dietrich, Syn. iii, 160.—Darby, Bot. S. States, 306.—Rœmer, Syn. Mon. iii, 121.—Cooper in Smithsonian Rep. 1858, 252.—Chapman, Fl. S. States, 127.—Wood, Cl. Book, 331; Bot. & Fl. 111.—Gray, Manual N. States, 5 ed. 159; Hall's Pl. Texas, 9.—Young, Bot. Texas, 257.—Kaleniczzenko in Bull. Soc. Imp. Nat. Moscow, xlviii, 29.—Vasey, Cat. Forest Trees, 14.

C. oxyacantha, Walter, Fl. Caroliniana, 147 [not Linnaeus].

Mespilus apiifolia, Marshall, Arbustum, 89.—Poiret, Suppl. iv, 68.—Sprengel, Syst. ii, 508.—Spach, Hist. Veg. ii, 67.

Mespilus monogyna, var. *apiifolia*, Koch, Dendrologie, i, 160.

C. oxyacantha, var. *apiifolia*, Regel in Act. Hort. St. Petersburg, 119.

PARSLEY HAW.

Southern Virginia, southward near the coast to about latitude 28°, extending west through the Gulf states to southern Arkansas and the valley of the Trinity river, Texas.

A small tree, rarely 6 to 9 meters in height, with a slender stem rarely exceeding 0.08 to 0.10 meter in diameter, or more often a low shrub, throwing up many stems from the ground; low, rich soil, reaching its greatest development in the pine barren hummocks of central Florida.

Wood heavy, hard, very close-grained, compact, susceptible of a beautiful polish; medullary rays thin, very obscure; color, bright brown tinged with red or rose, the sap-wood much lighter; specific gravity, 0.7453; ash, 0.97.

132.—*Cratægus spathulata*, Michaux.

Fl. Bor.-Am. i, 228.—Persoon, Syn. ii, 37.—Barton, Compend. Fl. Philadelph. i, 226.—Elliott, Sk. i, 552.—Loddiges, Bot. Cab. t. 1261.—Don, Miller's Dict. ii, 599.—Hooker, Companion Bot. Mag. i, 25.—Gray in Lindley, Bot. Reg. xxiii under t. 1957; Manual N. States, 5 ed. 159.—Eaton, Manual, 6 ed. 112.—Torrey & Gray, Fl. N. America, i, 467.—London, Arboretum, ii, 825, f. 591 & t.—Eaton & Wright, Bot. 212.—Dietrich, Syn. iii, 160.—Darby, Bot. S. States, 306.—Chapman, Fl. S. States, 126.—Lesquereux in Owen's 2d Rep. Arkansas, 359.—Wood, Cl. Book, 331; Bot. & Fl. 111.—Young, Bot. Texas, 257.—Kaleniczzenko in Bull. Soc. Imp. Nat. Moscow, xlviii, 31.—Ridgway in Am. Nat. vi, 728.

Mespilus Azarolus, Marshall, Arbustum, 89 [not Linnaeus].

Mespilus spathulata, Poiret, Suppl. iv, 68.—Desfontaines, Hist. Arb. ii, 157.—Sprengel, Syst. ii, 507.—Spach, Hist. Veg. ii, 66.—Koch, Dendrologie, i, 137.

C. microcarpa, Lindley, Bot. Reg. xxii, t. 1846.

Phænopyrum spathulatum, Rœmer, Syn. Mon. iii, 355.

SMALL-FRUITED HAW.

Virginia, southward to the Chattahoochee region of western Florida, west through the Gulf states to the valley of the Washita river, Arkansas (Hot Springs, *Letterman*), and the Colorado river, Texas.

A small tree, 6 to 8 meters in height, with a trunk 0.20 to 0.25 meter in diameter, or often reduced to a low shrub; margins of streams and prairies; common and reaching its greatest development along the bottom lands of western Louisiana and eastern Texas.

Wood heavy, hard, not strong, close-grained, compact; medullary rays very numerous, obscure; color, light brown or red, the sap-wood lighter; specific gravity, 0.7159; ash, 0.66.

133.—*Cratægus berberifolia*, Torrey & Gray,

Fl. N. America, i, 469.—Dietrich, Syn. iii, 159.—Walpers, Rep. ii, 59.—Rœmer, Syn. Mon. iii, 115.—Wood, Cl. Book, 332.—Regel in Act. Hort. St. Petersburg, i, 123.—Engelmann in Coulter's Bot. Gazette, vii, 128.

Mespilus berberifolia, Wenzig in Linnæa, xxxviii, 125.

Phænopyrum ellipticum, Rœmer, Syn. Mon. iii, 155.

Phænopyrum Virginicum, Rœmer, Syn. Mon. iii, 155.

New Orleans? (*Drummond*, No. 105¹); Opelousas, Louisiana (*Carpenter*, *Sargent*).

A small tree, 6 to 8 meters in height, with a trunk 0.20 to 0.25 meter in diameter; borders of prairies, in low ground; the fruit and wood not yet collected.

134.—*Cratægus æstivalis*, Torrey & Gray,

Fl. N. America, i, 468.—Walpers, Rep. ii, 58.—Dietrich, Syn. iii, 162.—Nuttall, Sylva, ii, 12; 2 ed. i, 162.—Darby, Bot. S. States, 306.—Chapman, Fl. S. States, 127.—Lesquereux in Owen's 2d Rep. Arkansas, 359.—Wood, Cl. Book, 331; Bot. & Fl. 111.—Regel in Act. Hort. St. Petersburg, i, 124.—Vasey, Cat. Forest Trees, 14.

Mespilus æstivalis, Walter, Fl. Caroliniana, 148.—Lamarck, Dict. iv, 447.

C. elliptica, Elliott, Sk. i, 548 [not Aiton].

C. lucida, Elliott, Sk. i, 549 [not Ehrhart].

C. opaca, Hooker & Arnott in Companion Bot. Mag. i, 25.—London, Arboretum, iv, 2663.

Anthomeles æstivalis, Rœmer, Syn. Mon. iii, 141.

MAY HAW. APPLE HAW.

South Carolina, south to northern Florida, west through the Gulf states to southern Arkansas and the valley of the Sabine river, Texas.

A small tree, 6 to 9 meters in height, with a trunk 0.15 to 0.20 meter in diameter; generally in sandy soil along the margins of streams and ponds; common and reaching its greatest development in the bottom lands of western Louisiana and eastern Texas.

Wood heavy, hard, not strong, close-grained, compact; medullary rays numerous, obscure; color, light brown or red, the sap-wood lighter; specific gravity, 0.6564; ash, 0.57.

The large, globular, fragrant, red fruit, of agreeable subacid flavor, used as a preserve, in jellies, etc.; ripening in May.

135.—*Cratægus flava*, Aiton,

Hort. Kew. ii, 169; 2 ed. iii, 201.—Willdenow, Spec. ii, 1002.—Persoon, Syn. ii, 37.—Pursh, Fl. Am. Sept. i, 338.—Nuttall, Genera, i, 305.—De Candolle, Prodr. ii, 628.—Watson, Dend. Brit. i, t. 59.—Don, Miller's Dict. ii, 600.—Lindley, Bot. Reg. xxiii, t. 1939.—Torrey & Gray, Fl. N. America, i, 468.—Eaton, Manual, 6 ed. 112.—London, Arboretum, ii, 823, f. 585 & t.—Eaton & Wright, Bot. 211.—Dietrich, Syn. iii, 160.—Darby, Bot. S. States, 306.—Cooper in Smithsonian Rep. 1858, 252.—Chapman, Fl. S. States, 28.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 83.—Lesquereux in Owen's 2d Rep. Arkansas, 359.—Wood, Cl. Book, 332; Bot. & Fl. 111.—Gray, Manual N. States, 5 ed. 160.—Regel in Act. Hort. St. Petersburg, i, 122.—Kaleniczenko in Bull. Soc. Imp. Nat. Moscow, xlviii, 27.—Vasey, Cat. Forest Trees, 14.

Mespilus flexispina, Mœnch, Verz. Baum. 62, t. 4.—Koch, Dendrologie, i, 139.

C. glandulosa, Aiton, Hort. Kew. ii, 168; 2 ed. iii, 201 [not Michaux].—Persoon, Syn. ii, 37.—Poiret, Suppl. iv, 69, in part.

Mespilus Caroliniana, Poiret in Lamarck, Dict. iv, 442.—Desfontaines, Hist. Arb. ii, 156.—Sprengel, Syst. ii, 507.

C. Caroliniana, Persoon, Syn. ii, 36.—Elliott, Sk. i, 554.—Eaton, Manual, 6 ed. 112.—Eaton & Wright, Bot. 212.

Mespilus flava, Willdenow, Enum. 523.—Poiret, Suppl. iv, 70.—Watson, Dend. Brit. i, t. 59.—Spach, Hist. Veg. ii, 59.

C. turbinata, Pursh, Fl. Am. Sept. Addend. 735.—Poiret, Suppl. v, 543.—Elliott, Sk. i, 549.—De Candolle, Prodr. ii, 627.—Don, Miller's Dict. ii, 599.—Eaton & Wright, Bot. 212.

Mespilus turbinata, Sprengel, Syst. ii, 506.—Spach, Hist. Veg. ii, 66.

C. flava, var. *lobata*, Lindley, Bot. Reg. xxiii, t. 1932.

C. lobata, Bosc in De Candolle, Prodr. ii, 628.—Don, Miller's Dict. ii, 599.—London, Arboretum, ii, 824, f. 554, 586.

Phænopyrum Carolinianum, Rœmer, Syn. Mon. iii, 152.

Anthomeles flava, *glandulosa*, and *turbinata*, Rœmer, Syn. Mon. iii, 141.

SUMMER HAW. YELLOW HAW.

Virginia, southward, generally near the coast, to Tampa bay, Florida, west through the Gulf states to eastern Texas and southern Arkansas.

A small tree, rarely 7 meters in height, with a trunk 0.30 meter in diameter, or reduced to a much-branched shrub 2 to 3 meters in height; borders of streams, in low, sandy soil subject to overflow.

Wood heavy, hard, close-grained, checking badly in drying, satiny, susceptible of a good polish; medullary rays very numerous, obscure; color, light brown tinged with red or rose, the sap-wood lighter; specific gravity, 0.7809; ash, 0.79.

Fruit small, red or yellow, acid.

Var. *pubescens*, Gray,

Manual N. States, 5 ed. 160.

Mespilus hiemalis, Walter, Fl. Caroliniana, 148.—Lamarck, Dict. iv, 447.

C. viridis, Walter, Fl. Caroliniana, 147 [not Linnæus].—Elliott, Sk. i, 551.

C. elliptica, Aiton, Hort. Kew. ii, 168; 2 ed. iii, 201.—Waldenow, Spec. ii, 1002.—Persoon, Syn. ii, 37.—Pursh, Fl. Am. Sept. i, 337.—Nuttall, Genera, i, 305.—Torrey, Fl. U. S. 475; Compend. Fl. N. States, 201.—De Candolle, Prodr. ii, 627.—Hooker, Fl. Bor.-Am. i, 201.—Don, Miller's Dict. ii, 598.—Beck, Bot. 33.—Eaton, Manual, 6 ed. 111.—Torrey & Gray, Fl. N. America, i, 469.—Eaton & Wright, Bot. 211.—Dietrich, Syn. iii, 109.—Darby, Bot. S. States, 306.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 84.—Regel in Act. Hort. St. Petersburg, i, 122.

Mespilus elliptica, Lamarck, Dict. iv, 447.—Wenzig in Linnæa, xxxviii, 125.—Koch, Dendrologie, i, 140.

C. glandulosa, Michaux, Fl. Bor.-Am. i, 288 [not Aiton].—Nuttall, Genera, i, 305.—Chapman, Fl. S. States, 128.—Vasey, Cat. Forest Trees, 14.

C. Michauxii, Persoon, Syn. ii, 38.

C. spathulata, Pursh, Fl. Am. Sept. i, 336 [not Michaux].—De Candolle, Prodr. ii, 627.—Lindley, Bot. Reg. xxii, t. 1890; xxiii, under t. 1957.

Mespilus Michauxii, Hornemann, Hort. Hafn. 455.—Poiret, Suppl. iv, 69.

C. flava, Elliott, Sk. i, 551 [not Aiton].

C. Virginica, Loddiges in London, Arboretum, ii, 842, f. 560, 615.—Kaleniczenko in Bull. Soc. Imp. Nat. Moscow, xlviii, 58.

SUMMER-HAW. RED HAW.

Virginia, southward to Tampa bay, Florida, and sparingly through the Gulf states to western Louisiana.

A low tree growing with the species, from which it is distinguished by the pubescence of the calyx and young branches, the smaller flowers, and larger, bright red or yellow, globular or pear-shaped fruit.

Wood heavy, hard, not strong, close-grained, compact; medullary rays numerous, very obscure; color, bright red or rose, the sap-wood lighter; specific gravity, 0.7683; ash, 0.91.

The large, edible fruit used in the south Atlantic states in preserves, jellies, etc.

NOTE.—*Crataegus parvifolia*, Aiton, of the south Atlantic region, a low shrub, is not included in this catalogue.

136.—*Heteromeles arbutifolia*, Rømer,

Syn. Mon. iii, 105.—Decaisne in Nouv. Arch. Mus. x, 144, t. 9.—Brewer & Watson, Bot. California, i, 188; ii, 444.

Crataegus arbutifolia, Poiret in Nouveau Duhamel, iv, 131; Dict. Suppl. i, 292.—Aiton, Hort. Kew. 2 ed. iii, 202.—Loddiges, Bot. Cab. t. 201.

Aronia arbutifolia, Nuttall, Genera, i, 306.

Photinia arbutifolia, Lindley in Trans. Linnæan Soc. xiii, 103; Bot. Reg. vi, 491 & under t. 1956.—Sprengel, Syst. ii, 508.—De Candolle, Prodr. ii, 631.—Chamisso & Schlechtendal in Linnæa, ii, 542.—Don, Miller's Dict. ii, 602.—Spach, Hist. Veg. ii, 80.—Hooker & Arnott, Bot. Beechey, 139, 340.—Torrey & Gray, Fl. N. America, i, 473.—Dietrich, Syn. iii, 162.—London, Arboretum, ii, 868, f. 619.—Bentham, Bot. Sulphur, 14; Pl. Hartweg. 307.—Torrey in Emory's Rep. 140; Sitgreaves' Rep. 119; Pacific R. R. Rep. iv, 86; Bot. Mex. Boundary Survey, 64; Bot. Wilkes Exped. 291.—Wood, Cl. Book, 329.—Bolander in Proc. California Acad. iii, 80.—Vasey, Cat. Forest Trees, 14.—Palmer in Am. Nat. xii, 599.—Maximowicz in Bull. Acad. Sci. St. Petersburg, xix, 180.—Wenzig in Linnæa, xxxviii, 96.

Mespilus arbutifolia, Link, Enum. Hort. Berol. ii, 36.

Photinia salicifolia, Presl, Epimel. Bot. 204.—Walpers, Ann. iii, 858.

H. Fremontiana, Decaisne in Nouv. Arch. Mus. x, 144.

TOYON. TOLLON. CALIFORNIA HOLLY.

California Coast ranges, Mendocino to San Diego county, extending east to the foot-hills of the Sierra Nevada and San Bernardino mountains.

A small, low-branched evergreen tree, rarely exceeding 9 meters in height, the short trunk sometimes 0.30 to 0.45 meter in diameter, or more often a low, much-branched shrub.

Wood very heavy, hard, close-grained, inclined to check in drying, satiny, susceptible of a beautiful polish; medullary rays numerous, very obscure; color, dark reddish-brown, the sap-wood lighter; specific gravity, 0.9326; ash, 0.54.

137.—*Amelanchier Canadensis*, Torrey & Gray,

Fl. N. America, i, 473.—Walpers, Rep. ii, 55.—Dietrich, Syn. iii, 158.—Torrey, Fl. N. York, i, 225.—Browne, Trees of America, 282.—Emerson, Trees Massachusetts, i, 443; 2 ed. ii, 503 & t.—Parry in Owen's Rep. 612.—Darlington, Fl. Cestrica, 3 ed. 86.—Richardson, Arctic Exped. 428.—Seemann, Bot. Herald, 52.—Hooker f. in Trans. Linnæan Soc. xxii², 290, 327.—Cooper in Smithsonian Rep. 1858, 252.—Chapman, Fl. S. States, 129.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 68.—Lesquereux in Owen's 2d Rep. Arkansas, 359.—Wood, Cl. Book, 329; Bot. & Fl. 110.—Engelmann in Trans. Am. Phil. Soc. new ser. xii, 191.—Porcher, Resources S. Forests, 168.—Gray, Manual N. States, 5 ed. 162.—Koch, Dendrologie, i, 180.—Vasey, Cat. Forest Trees, 14.—Maximowicz in Bull. Acad. St. Petersburg, xix, 175.—Ridgway in Proc. U. S. Nat. Mus. 1882, 66.

Mespilus Canadensis, Linnæus, Spec. 1 ed. 478 (excl. syn. Gronovius).—Walter, Fl. Caroliniana, 148.—Aiton, Hort. Kew. ii, 173.

Cratægus tomentosa, Linnæus, Spec. 1 ed. 476 (excl. syn. Gronovius).

Pyrus Botryapium, Linnæus f. Suppl. 255.—Wangenheim, Amer. 90, t. 28, f. 65.—Ehrhart, Beitr. i, 183; ii, 68.—Willdenow, Spec. ii, 1013; Enum. 525; Berl. Baumz. 322.—Aiton, Hort. Kew. 2 ed. iii, 207.—Pursh, Fl. Am. Sept. i, 339.—Hayne, Dend. Fl. 83.—Guimpel, Otto & Hayne, 100, t. 79.—Sprengel, Syst. ii, 509.—Audubon, Birds, t. 60.—Bigelow, Fl. Boston. 3 ed. 308.

Cratægus racemosa, Lamarck, Dict. i, 84.—Desfontaines, Hist. Arb. ii, 148.—Nouveau Duhamel, iv, 133.—Poiret, Suppl. i, 292.

Mespilus nivea, Marshall, Arbustum, 90.

Mespilus Canadensis, var. *cordata*, Michaux, Fl. Bor.-Am. i, 291.

Aronia Botryapium, Persoon, Syn. ii, 39.—Nuttall, Genera, i, 557.—Elliott, Sk. i, 557.—Torrey, Fl. U. S. 479; Compend. Fl. N. States, 203.—Eaton, Manual, 6 ed. 29.—Eaton & Wright, Bot. 135.

Mespilus arborea, Michaux f. Hist. Arb. Am. iii, 68, t. 11; N. American Sylva, 3 ed. ii, 60, t. 66.—Barton, Prodr. Fl. Philadelph. 55.

A. Botryapium, Lindley in Trans. Linnæan Soc. xiii, 100.—De Candolle, Prodr. ii, 632.—Hooker, Fl. Bor.-Am. i, 202.—Don, Miller's Dict. ii, 604.—Beck, Bot. 112.—Spach, Hist. Veg. ii, 84.—London, Arboretum, ii, 874, f. 627-629 & t.—Rœmer, Syn. Mon. iii, 145.—Darby, Bot. S. States, 307.—Wenzig in Linnæa, xxxiii, 110.—Decaisne in Nouv. Arch. Mus. x, 135.

Aronia arborea, Barton, Compend. Philadelph. i, 228.

Aronia cordata, Rafinesque, Med. Bot. ii, 196.

A. ovalis, Hooker, Fl. Bor.-Am. i, 202, in part.

Pyrus Bartramiana, Tausch, Fl. xxi, 715.

Pyrus Wangenheimiana, Tausch, Fl. xxi, 715.

A. Bartramiana, Rœmer, Syn. Mon. iii, 145.

A. Wangenheimiana, Rœmer, Syn. Mon. 146.

JUNE BERRY. SHAD BUSH. SERVICE TREE. MAY CHERRY.

Newfoundland and Labrador, west along the southern shores of Hudson bay to the Saskatchewan region, south through the Atlantic forests to northern Florida, southwestern Arkansas, and the Indian territory.

A small tree, 9 to 15 meters in height, with a trunk 0.30 to 0.45 meter in diameter, or in some forms reduced to a low shrub (var. *rotundifolia*, Torrey & Gray; var. *oligocarpa*, Torrey & Gray); common at the north, rare at the south, and reaching its greatest development on the high slopes of the southern Alleghany mountains; varying greatly in the shape of the leaves, size of the flowers, amount of pubescence on the leaves and young shoots, etc.

The best marked arborescent variety is—

var. *oblongifolia*, Torrey & Gray, Fl. N. America, i, 473.—Walpers, Rep. ii, 55.—Dietrich, Syn. iii, 158.—Torrey, Fl. N. York, i, 225; Nicolle's Rep. 149.—Emerson, Trees Massachusetts, i, 444; 2 ed. ii, 504 & t.—Wood, Cl. Book, 330; Bot. & Fl. 110.—Gray, Manual N. States, 5 ed. 162.—Macoun in Geological Rep. Canada, 1875-76, 195.

Crataegus spicata, Lamarck, Dict. i, 84.—Desfontaines, Hist. Arb. ii, 148.—Nouveau Duhamel, iv, 132.—Poiret, Suppl. i, 292.

Mespilus Canadensis, var. *obovalis*, Michaux, Fl. Bor.-Am. i, 291.

Pyrus ovalis, Willdenow, Spec. ii, 1014; Berl. Baumz. 323.—Pursh, Fl. Am. Sept. i, 340.—Schränk, Fl. Labrador, 26.—Bigelow, Fl. Boston. 3 ed. 207.

Aronia ovalis, Torrey, Fl. U. S. 479; Compend. Fl. N. States, 203.—Eaton, Manual, 6 ed. 29.—Eaton & Wright, Bot. 135.

A. ovalis, De Candolle, Prodr. ii, 632.—Meyer, Fl. Labrador, 81.—Hooker, Fl. Bor.-Am. i, 202, in part.—Don, Miller's Dict. ii, 604.—Beck, Bot. 112.—Spach, Hist. Veg. ii, 85.—London, Arboretum, ii, 876, f. 632.

A. intermedia, Spach, Hist. Veg. ii, 85.—Wenzig in Linnæa, xxxiii, 112.

A. oblongifolia, Rømer, Syn. Mon. iii, 147.

A. spicata, Decaisne in Nouv. Arch. Mus. x, 135, t. 9, f. 5.

Wood heavy, exceedingly hard, strong, close-grained, checking somewhat in seasoning, satiny, susceptible of a good polish; medullary rays very numerous, obscure; color, dark brown often tinged with red, the sap-wood much lighter; specific gravity, 0.7838; ash, 0.55; the small fruit sweet and edible.

NOTE.—The closely allied *Amelanchier alnifolia*, Nuttall, a low shrub, is widely distributed over the mountain ranges of the interior Pacific region.

HAMAMELACEÆ.

138.—*Hamamelis Virginica*, Linnæus,

Spec. 2 ed. 124.—Marshall, Arbustum, 58.—Du Roi, Harbk. i, 423.—Wangenheim, Amer. 89, t. 29, f. 62.—Lamarck, Dict. iii, 68; Ill. i, 350, t. 88.—Aiton, Hort. Kew. i, 167; 2 ed. i, 275.—Schkuhr, Handb. i, 88, t. 27.—Willdenow, Spec. i, 701; Enum. 171; Berl. Baumz. 172.—Michaux, Fl. Bor. Am. i, 100.—Persoon, Syn. i, 150.—Desfontaines, Hist. Arb. ii, 29.—Pursh, Fl. Am. Sept. i, 116.—Nuttall, Genera, i, 107.—Nouveau Duhamel, vii, 207, t. 60.—Elliott, Sk. i, 219.—Rømer & Schultes, Syst. iii, 483.—Loddiges, Bot. Cab. t. 598.—Barton, Fl. N. America, iii, 21, t. 78.—Torrey, Fl. U. S. 192; Compend. Fl. N. States, 86; Fl. N. York, i, 260.—Guimpel, Otto & Hayne, Abb. Holz. 95, t. 75.—Sprengel, Syst. i, 491.—Rafinesque, Med. Bot. i, 227, f. 45.—De Candolle, Prodr. iv, 268.—Hooker, Fl. Bor.-Am. i, 275; Companion Bot. Mag. i, 48.—Don, Miller's Dict. iii, 396, f. 69.—Beck, Bot. 152.—Eaton, Manual 6 ed. 164.—Spach, Hist. Veg. viii, 79.—Dietrich, Syn. i, 550.—Torrey & Gray, Fl. N. America, i, 597.—London, Arboretum, ii, 1007, f. 756, 757.—Eaton & Wright, Bot. 260.—Bigelow, Fl. Boston. 3 ed. 63.—Emerson, Trees Massachusetts, 416; 2 ed. ii, 473 & t.—Darby, Bot. S. States, 328.—Darlington, Fl. Cestrica, 3 ed. 98.—Agardh, Theor. & Syst. Pl. t. 13, f. 7.—Schnizlein, Icon. t. 167, f. 18-25, 27-29.—Gray in Am. Jour. Sci. 2 ser. xxiv, 438; 3 ser. v, 144; Manual N. States, 5 ed. 173.—Chapman, Fl. S. States, 157.—Curtis in Rep. Geological Surv. N. Carolina, iii, 105.—Leaquereux in Owen's 2d Rep. Arkansas, 362.—Wood, Cl. Book, 375; Bot. & Fl. 120.—Engelmann in Trans. Am. Phil. Soc. new ser. xii, 193.—Porcher, Resources S. Forests, 58.—Koch, Dendrologie, ii, 458.—Baillon in Adansonia, x, 123; Hist. Pl. iii, 389, f. 462-464.—Young, Bot. Texas, 291.—Maout & Decaisne, Bot. English ed. 408 & f.

H. dioica, Walter, Fl. Caroliniana, 255.—Gmelin, Syst. Veg. i, 281.

H. androgyna, Walter, Fl. Caroliniana, 255.—Gmelin, Syst. Veg. i, 282.

H. corylifolia, Mönch, Meth. 273.

H. macrophylla, Pursh, Fl. Am. Sept. i, 116.—Poiret, Suppl. v, 698.—Elliott, Sk. i, 220.—Rømer & Schultes, Syst. iii, 483.—Rafinesque, Med. Bot. i, 230.—Eaton, Manual, 6 ed. 164.—Don, Miller's Dict. iii, 396.—Eaton & Wright, Bot. 261.

Trilopus Virginiana, *nigra*, *rotundifolia*, and *dentata*, Rafinesque, New Sylva, 15-17.

H. Virginiana, var. *parvifolia*, Nuttall, Genera, i, 107.—Torrey, Fl. U. S. 193; Compend. Fl. N. States, 87.—Don, Miller's Dict. iii, 396.—Beck, Bot. 152.—Torrey & Gray, Fl. N. America, i, 597.

H. parvifolia, Rafinesque, Med. Bot. i, 230.

Trilopus parvifolia, Rafinesque, New Sylva, 17.

WITCH HAZEL.

Northern New England and southern Ontario to Wisconsin, south through the Atlantic region to northern Florida and eastern Texas.

A small tree, exceptionally 7 to 9 meters in height, with a trunk 0.30 to 0.37 meter in diameter, or more often a tall shrub throwing up many stems from the ground; common; rich, rather damp woodlands, reaching its greatest development in the region of the southern Alleghany mountains.

Wood heavy, hard, very close-grained, compact; layers of annual growth hardly distinguishable; medullary rays numerous, thin, obscure; color, light brown tinged with red, the sap-wood nearly white; specific gravity, 0.6856; ash, 0.37.

The bark and leaves rich in tannin, and largely used by herbalists in the form of fluid extracts, decoctions, etc., in external applications, and as a reputed remedy in hemorrhoidal affections (*New York Jour. Med.* x, 208.—*Trans. Am. Med. Assoc.* i, 350.—*U. S. Dispensatory*, 14 ed. 1661.—*Nat. Dispensatory*, 2 ed. 704).

139.—Liquidambar Styraciflua, Linnæus,

Spec. 1 ed. 999.—Marshall, *Arbustum*, 77.—Wangenheim, *Amer.* 49, t. 16, f. 40.—Walter, *Fl. Caroliniana*, 237.—Lamarck, *Dict.* iii, 533; III. iii, 367, t. 783.—Aiton, *Hort. Kew.* iii, 365; 2 ed. v, 306.—Gærtner, *Fruct.* ii, 57, t. 90.—Moench, *Meth.* 340.—Abbot, *Insecta Georgia*, i, 48.—B. S. Barton, *Coll.* i, 16.—Willdenow, *Spec.* iv, 475; *Enum.* 985; *Berl. Baumz.* 214.—Michaux, *Fl. Bor.-Am.* ii, 202.—Persoon, *Syn.* ii, 573.—Desfontaines, *Hist. Arb.* ii, 541.—Titford, *Hort. Bot. Am.* 97.—Schkuhr, *Handb.* iii, 275, t. 307.—Nouveau Duhamel, ii, 42, t. 10; vii, 207, t. 60.—Michaux f. *Hist. Arb. Am.* iii, 194, t. 4; *N. American Sylva*, 3 ed. ii, 42, t. 64.—Barton, *Prodr. Fl. Philadelph.* 92; *Compend. Fl. Philadelph.* ii, 177.—Pursh, *Fl. Am. Sept.* ii, 635.—Eaton, *Manual*, 110; 6 ed. 208.—Rafinesque, *Fl. Ludoviciana*, 116.—Nuttall, *Genera*, ii, 219; *Trans. Am. Phil. Soc.* 2 ser. v, 168.—Nees, *Fl. Offic.* t. 95.—Elliott, *Sk.* ii, 621.—Sprengel, *Syst.* iii, 864.—Humboldt, Bonpland & Kunth, *Nouv. Gen. & Spec.* vii, 273.—Audubon, *Birds*, t. 44.—Torrey, *Compend. Fl. N. States*, 357; *Fl. N. York*, ii, 217.—Beck, *Bot.* 326.—Hooker, *Companion Bot. Mag.* ii, 64.—Eaton & Wright, *Bot.* 302.—Spach, *Hist. Veg.* x, 84.—London, *Arboretum*, iv, 2049, f. 1961 & t.—Lindley, *Fl. Med.* 322.—Griffith, *Med. Bot.* 581, f. 254.—Broomfield in *London Jour. Bot.* vii, 144.—Schnizlein, *Icon.* t. 98, f. 5-21.—Seemann, *Bot. Herald*, 346.—Darby, *Bot. S. States*, 509.—Cooper in *Smithsonian Rep.* 1858, 252.—Chapman, *Fl. S. States*, 157.—Curtis in *Rep. Geological Surv. N. Carolina*, 1860, iii, 77.—Lesquereux in *Owen's 2d Rep. Arkansas*, 362.—Wood, *Cl. Book*, 375; *Bot. & Fl.* 120.—Porcher, *Resources S. Forests*, 344.—De Candolle, *Prodr.* xvi^a, 157.—Oliver in *Hooker f. Icon.* xi, 13.—Gray, *Manual N. States*, 5 ed. 174.—Koch, *Dendrologie*, ii, 464.—Young, *Bot. Texas*, 291.—Vasey, *Cat. Forest Trees*, 15.—Maout & Decaisne, *Bot. English ed.* 412 & figs.—Baillon, *Hist. Pl.* iii, 397, f. 471-474.—Guibourt, *Hist. Drogues*, 7 ed. ii, 300, f. 445.—Ridgway in *Am. Nat.* vi, 664; *Proc. U. S. Nat. Mus.* 1882, 67.—Broadhead in *Coulter's Bot. Gazette*, iii, 53.—Hemslay, *Bot. Am.-Cent.* i, 400.

L. Styraciflua, var. *Mexicana*, Örsted, *Am.-Cent.* xvi, t. 11.

L. macrophylla, Örsted, *Am.-Cent.* xvi, t. 10.

SWEET GUM. STAR-LEAVED GUM. LIQUIDAMBER. RED GUM. BILSTED.

Fairfield county, Connecticut, to the valleys of the lower Ohio, White, and Wabash rivers, south to cape Canaveral and Tampa bay, Florida, southwest through southern Missouri, Arkansas, and the Indian territory to the valley of the Trinity river, Texas; in central and southern Mexico.

A large tree, often 30 to 36 or, exceptionally, 48 meters in height, with a trunk 1.20 to 1.80 meter in diameter; in low, wet soil; very common and reaching its greatest development in the bottom lands of the Mississippi basin, here, with the cotton gum, forming a large proportion of the heavy forest growth.

Wood heavy, hard, not strong, rather tough, close-grained, compact, inclined to shrink and warp badly in seasoning, susceptible of a beautiful polish; medullary rays numerous, very obscure; color, bright brown tinged with red, the sap-wood nearly white; specific gravity, 0.5910; ash, 0.61; manufactured into lumber and used in the construction of buildings for plates, boarding, and clapboards, in cabinet work as a substitute for black walnut, and for veneering and street pavements; its great economic value hardly appreciated on account of the difficulty experienced in properly seasoning it.

The balsamic exudation obtained from this species at the south collected by herbalists and sometimes used in the form of a sirup as a substitute for storax in the treatment of catarrhal affections, or externally as an ointment in dressing frost-bite, abscess, etc., and in the manufacture of chewing gums (*Flückiger & Hanbury, Pharmacographia*, 246.—*Nat. Dispensatory*, 2 ed. 834).

RHIZOPHORACEÆ.

140.—Rhizophora Mangle, Linnæus,

Spec. 1 ed. 443.—Jacquin, *Amer.* 141, t. 89.—Gærtner, *Fruct.* i, 212, t. 45, f. 1.—Lamarck, *III.* ii, 517, t. 396; *Dict.* vi, 160.—Willdenow, *Spec.* ii, 844.—Persoon, *Syn.* ii, 2.—Decourtilz, *Fl. Med. Antilles*, i, 45, t. 10.—Vellozo, *Fl. Flum.* t. 1.—De Candolle, *Prodr.* iii, 32.—Eaton, *Manual*, 6 ed. 301.—Spach, *Hist. Veg.* iv, 332, t. 34.—Torrey & Gray, *Fl. N. America*, i, 484.—Nuttall in *Am. Jour. Sci.* 1 ser. v, 295.—Hooker & Arnott, *Bot. Beechey*, 290.—Arnott in *Ann. Nat. Hist.* i, 361.—Walpers, *Rep.* ii, 70.—Bentham, *Bot. Sulphur*, 14.—Darby, *Bot. S. States*, 312.—Porcher, *Resources S. Forests*, 55.—Grisebach, *Fl. British West Indies*, 274.—Schnizlein, *Icon.* t. 263, f. 1-7.—Maout & Decaisne, *Bot. English ed.* 419.—Eichler in *Martius, Fl. Brasil.* xii^a, 426, t. 90.—Vasey, *Cat. Forest Trees*, 15.—Baillon, *Hist. Pl.* vi, 284, f. 253-259.

R. racemosa, Meyer, *Prim. Fl. Esseq.* 185.—De Candolle, *Prodr.* iii, 32.

R. Americana, Nuttall, *Sylva*, i, 95, t. 24; 2 ed. i, 112, t. 24.—Cooper in *Smithsonian Rep.* 1858, 264.

MANGROVE.

Semi-tropical Florida, Mosquito inlet and Cedar Keys to the southern keys; delta of the Mississippi river and coast of Texas; southward through the West Indies and tropical America; now widely naturalized throughout the tropics of the old world (*A. De Candolle, Geog. Bot.* ii, 772).

A tree 12 to 18, or, exceptionally, 27 meters in height, with a trunk 0.30 to 0.60 meter in diameter, or more commonly not exceeding 4 to 7 meters in height; low saline shores, reaching in the United States its greatest development on bay Biscayne and cape Sable; south of latitude 29°, bordering with almost impenetrable thickets the coast of the Florida peninsula, ascending the rivers for many miles, especially those flowing from the Everglades, and entirely covering many of the southern keys.

Wood exceedingly heavy, hard and strong, close-grained, checking in drying, satiny, susceptible of a beautiful polish, containing many evenly-distributed rather small open ducts; medullary rays numerous, thin; color, dark reddish brown streaked with lighter brown, sap-wood lighter; specific gravity, 1.1617; ash, 1.82; furnishing valuable fuel; not greatly affected by the teredo, and used for piles.

COMBRETACEÆ.

141.—*Conocarpus erecta*, Linnæus,

Spec. 1 ed. 176.—Lamarck, Dict. ii, 96; Ill. i, 126, f. 1.—Jacquin, Amer. t. 78.—Gærtner, Fruct. ii, 470, t. 177, f. 3.—Swartz, Obs. 79.—Willdenow, Sp. i, 994.—Aiton, Hort. Kew. 2 ed. i, 381.—Titford, Hort. Bot. Am. 47.—De Candolle, Prodr. iii, 16.—Decourtilz, Fl. Med. Antilles, vi, 68, t. 399.—Spach, Hist. Veg. iv, 304.—Torrey & Gray, Fl. N. America, i, 485.—Nuttall, Sylva, i, 113, t. 33; 2 ed. i, 128, t. 33.—Richard, Fl. Cuba, 526.—Cooper in Smithsonian Rep. 1858, 264.—Chapman, Fl. S. States, 136.—Grisebach, Fl. British West Indies, 277.—Eichler in Martius, Fl. Brasil. xiv², 101, t. 35, f. 2.—Vasey, Cat. Forest Trees, 15.

BUTTON WOOD

Semi-tropical Florida, cape Canaveral to the southern keys, west coast, Tampa bay to cape Sable; through the West Indies to Brazil.

A low tree, often 8 or, exceptionally, 15 to 18 meters in height, with a trunk sometimes 0.60 meter in diameter; common and reaching its greatest development in the United States on Lost Man's river, north of cape Sable; or reduced to a low under shrub (var. *procumbens*, De Candolle, l. c.—Eichler, l. c.; *O. procumbens*, Linnæus, Spec. 1 ed. 177.—Jacquin l. c. 79, t. 52, f. 2.—Gærtner, l. c. iii, 205, f. 4.—Grisebach, l. c.; *O. acutifolia*, Willdenow in Roemer & Schultes, Syst. v, 574).

Wood very heavy and hard, strong, close-grained, very compact, susceptible of a beautiful polish; medullary rays numerous, obscure; color, dark yellow brown, the sap-wood lighter; specific gravity, 0.9900; ash, 0.32; burning slowly like charcoal, and highly valued for fuel.

142.—*Laguncularia racemosa*, Gærtner f.

Fruct. Suppl. 209, t. 217.—De Candolle, Prodr. iii, 17.—Spach, Hist. Veg. iv, 305.—Nuttall, Sylva, i, 117, t. 34; 2 ed. i, 132, t. 34.—Bentham, Bot. Sulphur, 14, 92.—Richard, Fl. Cuba, 527.—Eichler in Martius, Fl. Brasil. xiv², 102, t. 35, f. 3.—Cooper in Smithsonian Rep. 1858, 264.—Chapman, Fl. S. States, 136.—Grisebach, Fl. British West Indies, 276.—Vasey, Cat. Forest Trees, 15.—Baillon, Hist. Pl. vi, 278.

Conocarpus racemosa, Linnæus, Spec. 2 ed. 251; Syst. 181.—Jacquin, Amer. 80, t. 53.—Swartz, Obs. 79.—Willdenow, Spec. i, 995.

Schousboea commutata, Sprengel, Syst. ii, 332.

Bucida Buceras, Vellozo, Fl. Flum. iv, t. 87 [not Linnæus].

L. glabrifolia, Presl, Reil. Hænk, ii, 22.—Walpers, Rep. ii, 63.—Chapman, Fl. S. States, 136.

WHITE BUTTON WOOD. WHITE MANGROVE.

Semi-tropical Florida, cape Canaveral to the southern keys, west coast, Cedar Keys to cape Sable; through the West Indies and tropical America; coast of tropical Africa.

A small tree, sometimes 6 or, exceptionally, 22 meters in height (Shark river, Florida, *Curtiss*), with a trunk 0.30 to 0.60 meter in diameter, or toward its northern limits reduced to a low shrub; very common; saline shores of lagoons and bays.

Wood very heavy and hard, strong, close-grained, very compact; susceptible of a beautiful polish; medullary rays numerous, obscure; color, dark yellow-brown, the sap-wood much lighter; specific gravity, 0.7137; ash, 1.62.

MYRTACEÆ.

143.—*Calyptanthus Chytraculia*, Swartz,

Prodr. 79; Fl. Ind. Occ. ii, 921.—Willdenow, Spec. ii, 975.—Aiton, Hort. Kew. 2 ed. iii, 192.—De Candolle, Prodr. iii, 237.—Nuttall, Sylva, i, 101, t. 26; 2 ed. i, 117, t. 26.—Berg in Linnæa, xxvii, 26.—Cooper in Smithsonian Rep. 1858, 264.—Chapman, Fl. S. States, 131.—Grisebach, Fl. British West Indies, 232.—Hemsley, Bot. Am.-Cent. i, 408.

Myrtus Chytraculia, Linnæus, Amœn. v, 398.—Swartz, Obs. 202.

Eugenia pallens, Poiret, Suppl. iii, 122.

Semi-tropical Florida, shores of bay Biscayne, Key Largo; in the West Indies.

A small tree, sometimes 8 meters in height, with a trunk 0.10 to 0.15 meter in diameter.

Wood very heavy, hard, close-grained, compact, containing many evenly-distributed rather large open ducts; medullary rays numerous, thin; color, brown tinged with red, the sap-wood a little lighter; specific gravity, 0.8992; ash, 3.32.

144.—*Eugenia buxifolia*, Willdenow,

Spec. ii, 960.—Persoon, Syn. ii, 28.—De Candolle, Prodr. iii, 275.—Nuttall, Sylva, i, 108, t. 29; 2 ed. i, 123, t. 29.—Cooper in Smithsonian Rep. 1858, 264.—Chapman, Fl. S. States, 131.—Grisebach, Fl. British West Indies, 236.—Vasey, Cat. Forest Trees, 15.

Myrtus buxifolia, Swartz, Prodr. 78; Fl. Ind. Occ. ii, 899.

Myrtus axillaris, Poiret in Lamarck, Dict. iv, 412.

E. myrtoides, Poiret, Suppl. iii, 125.

Myrtus Poireti, Sprengel, Syst. ii, 483.

E. triplinervia, Berg in Linnæa, xxvii, 190, in part.

GURGEON STOPPER. SPANISH STOPPER.

Semi-tropical Florida, cape Canaveral to the southern keys, west coast, Caloosa river to cape Romano; in the West Indies.

A small tree, rarely 6 to 9 meters in height, with a trunk sometimes 0.30 meter in diameter, reaching its greatest development on the rich hummocks of the Everglades.

Wood very heavy, exceedingly hard, very strong, close-grained, very compact; medullary rays numerous, thin; color, dark brown shaded with red, the sap-wood a little lighter; specific gravity, 0.9360; ash, 1.50; somewhat used for fuel.

145.—*Eugenia dichotoma*, De Candolle,

Prodr. iii, 278.—Nuttall, Sylva, i, 103, t. 27; 2 ed. i, 120, t. 27.—Berg in Linnæa, xxvii, 261.—Cooper in Smithsonian Rep. 1858, 264.—Chapman, Fl. S. States, 131.—Vasey, Cat. Forest Trees, 15.

E. divaricata, Lamarck, Dict. i, 202.

? *Myrtus dichotoma*, Vahl in Poiret, Suppl. iv, 53.

Anamomis punctata, Grisebach, Fl. British West Indies, 240.

NAKED WOOD.

Semi-tropical Florida, Mosquito inlet to cape Canaveral, common; west coast, Caloosa river to cape Romano; in the West Indies.

A small tree, sometimes 6 to 8 meters in height, with a trunk rarely 0.15 meter in diameter.

A form with the leaves, buds, and calyx more or less pubescent (*E. dichotoma*, var. *fragrans*, Nuttall, l. c.; *E. pungens*, Willdenow, Spec. ii, 964; Bot. Mag. t. 1242; *E. montana*, Aublet, Guian. i, 495, t. 195), not rare in West Indies, and, according to Nuttall, collected by Mr. Baldwin in the vicinity of New Smyrna, Florida, has not been rediscovered within the limits of the United States.

Wood very heavy, hard, close-grained, compact; medullary rays numerous, thin; color, light brown or red, sap-wood yellow; specific gravity, 0.8983; ash, 0.74.

The small, edible fruit of agreeable aromatic flavor, and greatly improved by cultivation in rich soil.

146.—*Eugenia monticola*, De Candolle,

Prodr. iii, 275.—Chapman, Fl. S. States, 131.—Grisebach, Fl. British West Indies, 236.—Vasey, Cat. Forest Trees, 15.

Myrtus monticola, Swartz, Fl. Ind. Occ. ii, 898.

E. triplinervia, Berg in Linnæa, xxvii, 193, in part.

E. axillaris, Berg in Linnæa, xxvii, 201, in part.

STOPPER. WHITE STOPPER.

Florida, Saint John's river to Umbrella Key; rare; in the West Indies.

A small tree, rarely 7 meters in height, with a trunk 0.30 meter in diameter, or in northern Florida reduced to a low shrub.

Wood very heavy, hard, strong, very close-grained, compact; medullary rays numerous, thin; color, brown, often tinged with red, the sap-wood darker; specific gravity, 0.9156; ash, 1.89.

147.—*Eugenia longipes*, Berg.

Linnæa, xxvii, 150.—Chapman, Fl. S. States, Suppl. 620.

STOPPER.

Semi-tropical Florida, No-Name Key; in the West Indies.

A small tree, 4 to 7 meters in height, with a trunk 0.15 to 0.20 meter in diameter; rare.

Wood very heavy, hard, close-grained, checking badly in drying, containing many evenly-distributed open ducts; medullary rays numerous, very obscure; color, dark brown or nearly black, the sap-wood brown tinged with red; specific gravity, 1.1235; ash, 3.48.

The small red fruit with the flavor of cranberries.

148.—*Eugenia procera*, Poirét,

Suppl. ii, 129.—De Candolle, Prodr. iii, 268.—Nuttall, Sylva, i, 106, t. 28; 2 ed. i, 122, t. 28.—Berg in Linnæa, xxvii, 207.—Cooper in Smithsonian Rep. 1858, 264.—Chapman, Fl. S. States, 131.—Grisebach, Fl. British West Indies, 238.—Vasey, Cat. Forest Trees, 15.

Myrtus procera, Swartz, Prodr. 77; Fl. Ind. Occ. ii, 887.—Willdenow, Spec. ii, 968.

E. Baruensis, Grisebach, Cat. Pl. Cub. [not Jacquin], 87.

RED STOPPER.

Semi-tropical Florida, shores of bay Biscayne, Key Largo, Elliott's Key; in the West Indies.

A tree, 12 to 18 meters in height, with a trunk 0.30 to 0.45 meter in diameter; often forming extensive groves, and reaching its greatest development in the United States in the neighborhood of Miami, bay Biscayne.

Wood very heavy, exceedingly hard, very strong and close-grained, compact; medullary rays numerous, hardly distinguishable; color, light yellow-brown, the sap-wood darker; specific gravity, 0.9453; ash, 2.62; probably valuable in cabinet-making and as a substitute for box-wood for coarse wood-engraving.

NOTE.—*Psidium Guaiava*, Raddi, the Guava, widely cultivated in the tropics for its fruit, is now sparingly naturalized in semi-tropical Florida.

C A C T A C E Æ.

149.—*Cereus giganteus*, Engelmann;

Emory's Rep. 158; Am. Jour. Sci. 2 ser. xiv, 335; xvii, 231; Proc. Am. Acad. iii, 287; Bot. Mex. Boundary Survey, Cactaceæ, 42, t. 61, 62 & front.; Brewer & Watson, Bot. California, i, 247.—Thurber in Mem. Am. Acad. new ser. v, 302, 305.—Fl. des Serres, x, 24, & t.; xv, 187, t. 1600.—Bigelow in Pacific R. R. Rep. iv, 12.—Engelmann & Bigelow in Pacific R. R. Rep. iv, 36.—Walpers, Ann. v, 46.—Cooper in Smithsonian Rep. 1858, 259.—Lemaire, Ill. Hort. ix, Misc. 95.—Marcou in Jour. Hort. Soc. France, 2 ser. iii, 676.—Lindley, Treasury Bot. 256, t. 17.—Vasey, Cat. Forest Trees, 15.—Rothrock in Wheeler's Rep. vi, front.—Hemsley, Bot. Am.-Cent. i, 343.—James in Am. Nat. xv, 982, f. 3.

Pilocereus Engelmanni, Lemaire, Ill. Hort. ix, Misc. 95.

SUWARROW. SAGUARO. GIANT CACTUS.

Valley of Bill Williams river, Arizona, south and east through central Arizona to the valley of the San Pedro river; southward in Sonora.

A tall, columnar tree, 8 to 18 meters in height, with a trunk sometimes 0.60 meter in diameter; dry, stony *mesas* or low hills rising from the desert.

Wood of the large, strong ribs, very light, soft, rather coarse-grained, solid, satiny, susceptible of a fine polish, almost indestructible in contact with the ground; medullary rays very numerous, broad; color, light brown tinged with yellow; specific gravity, 0.3188; ash, 3.45; used in the region almost exclusively for the rafters of adobe houses, for fencing, and by the Indians for lances, bows, etc.

The edible fruit largely collected and dried by the Indians.

CORNACEÆ.

150.—*Cornus alternifolia*, Linnæus f.

Suppl. 125.—Lamarck, Dict. ii, 116; Ill. i, 303.—L'Heritier, Cornus, 10, t. 6.—Ehrhart, Beitr. iii, 19.—Aiton, Hort. Kew. i, 159; 2 ed. i, 262.—Willdenow, Spec. i, 664; Enum. 165; Berl. Baumz. 104.—Michaux, Fl. Bor.-Am. i, 93.—Persoon, Syn. i, 144.—Desfontaines, Hist. Arb. i, 351.—Nouveau Duhamel, ii, 157, t. 45.—Pursh, Fl. Am. Sept. i, 109.—Nuttall, Genera, i, 99.—Rømer & Schultes, Syst. iii, 323; Mant. 251.—Elliott, Sk. i, 210.—Guimpel, Otto & Hayne, Abb. Holz. 53, t. 43.—Hayne, Dend. Fl. 8.—Torrey, Fl. U. S. 180; Compend. Fl. N. States, 83; Fl. N. York, i, 288.—Sprengel, Syst. i, 451.—De Candolle, Prodr. iv, 271.—Hooker, Fl. Bor.-Am. i, 275.—Don, Miller's Dict. iii, 398.—Beck, Bot. 154.—Eaton, Manual, 6 ed. 109.—Tausch in Regensb. Fl. xxi, 732.—Spach, Hist. Veg. viii, 92.—Dietrich, Syn. i, 503.—Torrey & Gray, Fl. N. America, i, 649.—London, Arboretum, ii, 1010, f. 760.—Eaton & Wright, Bot. 210.—Bigelow, Fl. Boston. 3 ed. 60.—C. A. Meyer in Mem. Acad. Sci. St. Petersburg, v, 6, 13.—Walpers, Rep. v, 932.—Emerson, Trees Massachusetts, 409; 2 ed. ii, 463 & t.—Parry in Owen's Rep. 613.—Darlington, Fl. Cestrica, 3 ed. 110.—Cooper in Smithsonian Rep. 1858, 252.—Chapman, Fl. S. States, 167.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 167.—Wood, Cl. Book, 391; Bot. & Fl. 143.—Gray, Manual N. States, 5 ed. 201.—Koch, Dendrologie, i, 690.—Young, Bot. Texas, 303.

C. alternata, Marshall, Arbustum, 35.

DOGWOOD.

New Brunswick, west along the valley of the Saint Lawrence river to the northern shores of lake Superior, south through the northern states and along the Alleghany mountains to northern Georgia and Alabama.

A small tree, 4 to 8 meters in height, with a trunk 0.15 to 0.20 meter in diameter; low, rich woods and borders of streams and swamps.

Wood heavy, hard, close-grained, checking badly in drying; medullary rays numerous, thin; color, brown tinged with red, the sap-wood light yellow; specific gravity, 0.6696; ash, 0.41.

151.—*Cornus florida*, Linnæus,

Spec. 1 ed. 117.—Marshall, Arbustum, 35.—Lamarck, Dict. ii, 114; Ill. i, 302.—Wangenheim, Amer. 51, t. 17, f. 41.—Walter, Fl. Caroliniana, 88.—L'Heritier, Cornus, 4.—Aiton, Hort. Kew. i, 157; 2 ed. i, 261.—Willdenow, Spec. i, 661; Enum. 164; Berl. Baumz. 100.—Abbot, Insects Georgia, ii, t. 73.—B. S. Barton, Coll. i, 12, 45; ii, 17, 19.—Bot. Mag. t. 526.—Michaux, Fl. Bor.-Am. i, 91.—Persoon, Syn. i, 143.—Desfontaines, Hist. Arb. i, 350.—Schkuhr, Handb. 82.—Titford, Hort. Bot. Am. 41, t. 16, f. 7.—Nouveau Duhamel, ii, 153.—Michaux f. Hist. Arb. Am. iii, 138, t. 3; N. American Sylva, 3 ed. i, 176, t. 48.—Pursh, Fl. Am. Sept. i, 108.—Bigelow, Med. Bot. ii, 69, t. 73; Fl. Boston. 3 ed. 59.—Eaton, Manual, 19; 6 ed. 108.—Nuttall, Genera, i, 98.—Barton, Med. Bot. i, 43, t. 3.—Rømer & Schultes, Syst. iii, 319.—Hayne, Dend. Fl. 6.—Guimpel, Otto & Hayne, Abb. Holz. 21, t. 19.—Elliott, Sk. i, 207.—Torrey in Ann. Lyc. N. York, ii, 208; Fl. U. S. 178; Compend. Fl. N. States, 82; Fl. N. York, i, 290; Nicolle's Rep. 151; Emory's Rep. 408.—Sprengel, Syst. i, 451.—Beck in Am. Jour. Sci. 1 ser. x, 264; Bot. 153.—Audubon, Birds, t. 8, 73, 122.—Rafinesque, Med. Bot. i, 131, f. 28.—De Candolle, Prodr. iv, 273.—Hooker, Fl. Bor.-Am. i, 277, in part; Companion Bot. Mag. i, 48.—Don, Miller's Dict. iii, 400.—Lindley, Fl. Med. 81.—Dietrich, Syn. i, 504.—Torrey & Gray, Fl. N. America, i, 652.—London, Arboretum, ii, 1017, f. 769.—Eaton & Wright, Bot. 209.—Reid in London Gard. Chronicle, 1844, 276.—Browne, Trees of America, 350.—Emerson, Trees Massachusetts, 413; 2 ed. ii, 467 & t.—Griffith, Med. Bot. 347, f. 164.—Carson, Med. Bot. i, 50, t. 42.—Richardson, Arctic Exped. 429.—Darlington, Fl. Cestrica, 3 ed. 111.—Darby, Bot. S. States, 339.—Cooper in Smithsonian Rep. 1858, 252.—Chapman, Fl. S. States, 168.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 60.—Lesquereux in Owen's 2d Rep. Arkansas, 364.—Wood, Cl. Book, 391; Bot. & Fl. 143.—Blakie in Canadian Nat. vi, 1.—Engelmann in Trans. Am. Phil. Soc. new ser. xii, 194.—Porcher, Resources S. Forests, 59.—Gray, Manual N. States, 5 ed. 200; Hall's Pl. Texas, 11.—Koch, Dendrologie, i, 694.—Young, Bot. Texas, 303.—Vasey, Cat. Forest Trees, 16.—Baillon, Hist. Pl. vii, 68, f. 46.—Broadhead in Coulter's Bot. Gazette, iii, 53.—Bentley & Trimen, Med. Pl. ii, 136, t. 136.—Bell in Geological Rep. Canada, 1879-'80, 55c.—Ridgway in Proc. U. S. Nat. Mus. 1882, 67.

Benthamidia florida, Spach, Hist. Veg. viii, 107.

FLOWERING DOGWOOD. BOX WOOD.

Southern New England, southern Ontario, southern Minnesota, and through the Atlantic forests to latitude 28° 50' in Florida, and the valley of the Brazos river, Texas.

A small tree, 9 to 12 meters in height, with a trunk 0.30 to 0.45 meter in diameter, or toward its northern limits reduced to a low shrub; rich woods; very common, especially at the south.

Wood heavy, hard, strong, close-grained, tough, checking badly in drying, satiny, susceptible of a beautiful polish; medullary rays numerous, conspicuous; color, brown, changing in different specimens to shades of green and red, the sap-wood lighter; specific gravity, 0.8153; ash, 0.67; used in turnery, for wood engravings and the bearings of machinery, hubs of wheels, barrel hoops, etc.

The bark, especially of the root, in common with that of the other species of the genus, possesses bitter tonic properties, and is used in decoctions, etc., in the treatment of intermittent and malarial fevers (*Am. Jour. Pharm.* vii, 109.—*Maisch in Proc. Am. Pharm. Assoc.* 315.—*U. S. Dispensatory*, 14 ed. 352.—*Nat. Dispensatory*, 2 ed. 467).

152.—*Cornus Nuttallii*, Audubon,

Birds, t. 467.—Torrey & Gray, *Fl. N. America*, i, 652.—Walpers, *Rep.* ii, 435.—Bentham, *Pl. Hartweg*, 312.—Nuttall, *Sylva*, iii, 51, t. 97; 2 ed. ii, 117, t. 97.—Durand in *Jour. Philadelphia Acad.* 1855, 89.—Torrey in *Pacific R. R. Rep.* iv, 94; *Bot. Mex. Boundary Survey*, 71; *Bot. Wilkes Exped.* 326.—Newberry in *Pacific R. R. Rep.* vi, 24, 75.—Cooper in *Smithsonian Rep.* 1858, 259; *Pacific R. R. Rep.* xii³, 29, 63.—Lyll in *Jour. Linnæan Soc.* vii, 134.—Gray in *Proc. Am. Acad.* viii, 387.—Brewer & Watson, *Bot. California*, i, 274; ii, 452.—Vasey, *Cat. Forest Trees*, 16.—Hall in *Coulter's Bot. Gazette*, ii, 88.—Macoun in *Geological Rep. Canada*, 1875-76, 198.—G. M. Dawson in *Canadian Nat. new ser.* ix, 331.

C. florida, Hooker, *Fl. Bor.-Am.* i, 277, in part.

FLOWERING DOGWOOD.

Vancouver's island and along the coast of southern British Columbia, through western Washington territory and Oregon, and southward through the Coast ranges of California and along the western slope of the Sierra Nevada to the San Bernardino mountains.

A small, slender tree, sometimes 18 to 24 meters in height, with a trunk rarely 0.45 meter in diameter; ascending the Cascade mountains to 3,000 feet, and the San Bernardino mountains to from 4,000 to 5,000 feet elevation; common; rich, rather damp soil, generally in the dense shade of coniferous forests.

Wood heavy, exceedingly hard, strong, close-grained, compact, satiny, susceptible of a good polish; medullary rays numerous, obscure; color, light brown tinged with red, the sap-wood lighter; specific gravity, 0.7481; ash, 0.50; somewhat used in cabinet-making, for mauls, handles, etc.

153.—*Nyssa capitata*, Walter,

Fl. Caroliniana, 253.—Lamarck, *Dict.* iv, 508.—Michaux f. *Hist. Arb. Am.* ii, 257, t. 20; *N. American Sylva*, 3 ed. iii, 37, t. 113.—Aiton, *Hort. Kew.* 2 ed. v, 480.—Poiret, *Suppl.* v, 740.—Elliott, *Sk.* ii, 685.—Hooker, *Companion Bot. Mag.* ii, 62.—Eaton, *Manual*, 6 ed. 236.—Eaton & Wright, *Bot.* 329.—Spach, *Hist. Veg.* x, 464.—Darby, *Bot. S. States*, 493.—Cooper in *Smithsonian Rep.* 1858, 253.—Chapman, *Fl. S. States*, 168.—Lesquereux in *Owen's 2d Rep. Arkansas*, 364.—Wood, *Cl. Book*, 392; *Bot. & Fl.* 143.—Koch, *Dendrologie*, ii, 456.—Vasey, *Cat. Forest Trees*, 16.

N. Ogeche, Marshall, *Arbustum*, 97.

N. coccinea, Bartram, *Travels*, 2 ed. 17.

N. tomentosa, Poiret in Lamarck, *Dict.* iv, 508.

N. candicans, Michaux, *Fl. Bor.-Am.* ii, 259.—Persoon, *Syn.* ii, 614.—Desfontaines, *Hist. Arb.* i, 37.—Willdenow, *Spec.* iv, 1113.—Pursh, *Fl. Am. Sept.* i, 117.—Poiret, *Suppl.* iv, 116.—Nuttall, *Genera*, ii, 236; *Trans. Am. Phil. Soc.* v, 167.—Rømer & Schultes, *Syst.* v, 557.—Sprengel, *Syst.* i, 832.—Dietrich, *Syn.* i, 879.—London, *Arboretum*, iii, 1318, f. 1199.—Browne, *Trees of America*, 426.

N. montana, Gärtner, *Fruct.* iii, 201, t. 216.

OGEECHEE LIME. SOUR TUPELO. GOPHER PLUM.

Georgia, from the valley of the Ogeechee to the Saint Mary's river, west Florida (near Vernon, *Mohr*), and in southern Arkansas.

A tree 9 to 18 meters in height, with a trunk 0.30 to 0.90 meter in diameter; deep swamps and river bottoms; rare and local.

Wood light, soft, not strong, tough, rather coarse-grained, compact, unwedgeable, containing many regularly-distributed open ducts; medullary rays numerous, thin; color, white, the sap-wood hardly distinguishable; specific gravity, 0.4613; ash, 0.34.

A conserve, under the name of "Ogeechee limes", is made from the large, acid fruit.

154.—*Nyssa sylvatica*, Marshall,

Arbustum, 97.—Michaux f. Hist. Arb. Am. ii, 260, t. 21; N. American Sylva, 3 ed. iii, 29, t. 110.—Poirer, Suppl. iv, 116.—Barton, Prodr. Fl. Philadelph. 97; Compend. Fl. Philadelph. ii, 193.

N. aquatica, Linnæus, Spec. 1 ed. 1058, in part.—St. Hilaire, Fam. Nat. ii, 152.—Persoon, Syn. ii, 614.—Michaux f. Hist. Arb. Am. ii, 165, t. 22; N. American Sylva, 3 ed. iii, 31, t. 111.—Rømer & Schultes, Syst. v, 576.—Barton, Prodr. Fl. Philadelph. 97; Compend. Fl. Philadelph. ii, 192.—Sprengel, Syst. i, 832.—Audubon, Birds, t. 133.—Elliott, Sk. ii, 684.—Dietrich, Syn. i, 878.—Eaton, Manual, 6 ed. 236.—Eaton & Wright, Bot. 329.—Spach, Hist. Veg. x, 464.—Darby, Bot. S. States, 492.—Chapman, Fl. S. States, 168.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 168.—Porcher, Resources S. Forests, 347.—Koch, Dendrologie, ii, 455.—Young, Bot. Texas, 304.—Vasey, Cat. Forest Trees, 16.

N. multiflora, Wangenheim, Amer. 46, t. 16, f. 39.—Elliott, Sk. ii, 684.—Walter, Fl. Caroliniana, 253.—Beck, Bot. 307.—Eaton, Manual, 6 ed. 236.—Eaton & Wright, Bot. 329.—Spach, Hist. Veg. x, 463.—Torrey, Fl. N. York, ii, 161, t. 95.—Emerson, Trees Massachusetts, 312, t. 17; 2 ed. ii, 353 & t.—Schnizlein, Icon. t. 108, f. 1, 2.—Darlington, Fl. Cestrica, 3 ed. 254.—Darby, Bot. S. States, 492.—Cooper in Smithsonian Rep. 1858, 252.—Chapman, Fl. S. States, 168.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 62.—Lesquereux in Owen's 2d Rep. Arkansas, 364.—Wood, Cl. Book, 392; Bot. & Fl. 143.—Gray, Manual N. States, 5 ed. 201.—Koch, Dendrologie, ii, 554.—Young, Bot. Texas, 304.—Vasey, Cat. Forest Trees, 16.—Broadhead in Coulter's Bot. Gazette, iii, 53.—Bessey in Am. Nat. xv, 134.—Bell in Geological Rep. Canada, 1879-'80, 55.—Ridgway in Proc. U. S. Nat. Mus. 1882, 68.—Burgess in Coulter's Bot. Gazette, vii, 95.

N. Caroliniana, Poirer in Lamarck, Dict. iv, 507; Lamarck, Ill. iii, 442, t. 851, f. 1.

N. biflora, Walter, Fl. Caroliniana, 253.—Lamarck, Dict. iv, 508.—Michaux, Fl. Bor.-Am. ii, 259.—Willdenow, Spec. iv, 1113; Enum. 1061; Berl. Baumz. 256.—Desfontaines, Hist. Arb. i, 37.—Gærtner f. Fruct. Suppl. 203, t. 216.—Aiton, Hort. Kew. 2 ed. v, 479.—Pursh, Fl. Am. Sept. i, 177.—Nuttall, Genera, ii, 236; Trans. Am. Phil. Soc. v, 167.—Poirer, Suppl. iv, 115.—Torrey in Ann. Lyc. N. York, ii, 200; Compend. Bot. N. States, 372.—Hayne, Dend. Fl. 229.—Eaton, Manual, 116.—Beck, Bot. 307.—London, Arboretum, iii, 1317, f. 1195, 1196.—Browne, Trees of America, 423.—Baillon, Hist. Pl. v, 266, f. 241-244.

N. integrifolia, Aiton, Hort. Kew. iii, 446.—Persoon, Syn. ii, 614.

N. Canadensis, Poirer in Lamarck, Dict. iv, 507.

N. villosa, Michaux, Fl. Bor.-Am. ii, 258.—Willdenow, Spec. iv, 1112.—Desfontaines Hist. Arb. i, 37.—Aiton, Hort. Kew. 2 ed. v, 479.—Bigelow, Fl. Boston. 3 ed. 380.—Pursh, Fl. Am. Sept. i, 117.—Nuttall, Genera, ii, 276.—Rømer & Schultes, Syst. v, 575.—Sprengel, Syst. i, 832.—Torrey, Compend. Bot. N. States, 372.—Dietrich, Syn. i, 878.—London, Arboretum, iii, 1317, f. 1197, 1198.

N. multiflora, var. *sylvatica*, Watson, Index, 442.

TUPELO. SOUR GUM. PEPPERIDGE. BLACK GUM.

Valley of the Kennebec river, Maine (Kent's Hill, *Prof. Stone*), West Milton, Vermont, west to central Michigan, south to Tampa bay, Florida, and the valley of the Brazos river, Texas.

A tree 15 to 36 meters in height, with a trunk 0.60 to 1.50 meter in diameter, or at the north much smaller; borders of swamps, or on rather high, rich hillsides and pine uplands; at the south often in pine-barren ponds and deep swamps, the base of the trunk then greatly enlarged and swollen (*N. aquatica*).

Wood heavy, rather soft, strong, very tough, unwedgeable, difficult to work, inclined to check unless carefully seasoned, not durable in contact with the soil, containing numerous regularly-distributed small open ducts; medullary rays numerous, thin; color, light yellow or often nearly white, the sap-wood hardly distinguishable; specific gravity, 0.6353; ash, 0.52; now largely used for the hubs of wheels, rollers in glass factories, ox yokes, and on the gulf coast for wharf piles.

NOTE.—Various forms of *Nyssa*, which at different times have been considered by botanists as entitled to specific rank, are connected by so many intermediate forms, and offer so few distinctive characters, that they are here united into one polymorphous species, which thus enlarged may properly bear Marshall's earlier name of *Nyssa sylvatica*, rather than the more familiar *Nyssa multiflora* of Wangenheim.

155.—*Nyssa uniflora*, Wangenheim,

Amer. 83, t. 27, f. 57.—Walter, Fl. Caroliniana, 253.—Elliott, Sk. ii, 686.—Eaton & Wright, Bot. 329.—Darby, Bot. S. States, 493.—Cooper in Smithsonian Rep. 1858, 253.—Chapman, Fl. S. States, 168.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 62.—Wood, Cl. Book, 392; Bot. & Fl. 143.—Gray, Manual N. States, 5 ed. 201.—Koch, Dendrologie, ii, 455.—Young, Bot. Texas, 304.—Vasey, Cat. Forest Trees, 16.

N. aquatica, Linnæus, Spec. 1058, in part.—Marshall, Arbustum, 96.—Lamarck, Dict. iv, 507.—Desfontaines, Hist. Arb. i, 36.

N. denticulata, Aiton, Hort. Kew. iii, 446; 2 ed. v, 480.—Persoon, Syn. ii, 615.—Willdenow, Spec. iv, 1114.—Gærtner f. Fruct. Suppl. 203, t. 216.—Pursh, Fl. Am. Sept. i, 178.—Poirer, Suppl. iv, 115.—Nuttall, Genera, ii, 236.—Hayne, Dend. Fl. 229.—Rømer & Schultes, Syst. v, 577.—Sprengel, Syst. i, 832.—Dietrich, Syn. i, 879.

N. angulosa, Poiret in Lamarek, Dict. iv, 507; Ill. iii, 442, t. 851, f. 2.—Rœmer & Schultes, Syst. v, 578.

N. palustris, Salisbury, Prodr. 175.

N. tomentosa, Michaux, Fl. Bor.-Am. ii, 259.—Persoon, Syn. ii, 615.—Willdenow, Spec. iv, 1113.—Pursh, Fl. Am. Sept. i, 177.—Nuttall, Genera, ii, 236.—Rœmer & Schultes, Syst. v, 577.—Elliott, Sk. ii, 685.—Sprengel, Syst. i, 832.—Audubon, Birds, t. 13.—Dietrich, Syn. i, 879.—Eaton & Wright, Bot. 329.—Darby, Bot. S. States, 498.

N. angulisans, Michaux, Fl. Bor.-Am. ii, 259.—Dietrich, Syn. i, 879.—Spach, Hist. Veg. x, 465.

N. grandidentata, Michaux f. Hist. Arb. Am. ii, 252, t. 19; N. American Sylva, 3 ed. ii, 34, t. 112.—London, Arboretum, iii, 1319, f. 1200, 1201.—Lesquereux in Owen's 2d Rep. Arkansas, 364.

N. capitata var. *grandidentata*, Browne, Trees of America, 426.

LARGE TUPELO. COTTON GUM. TUPELO GUM.

Southern Virginia, south near the coast to the valley of the Saint Mary's river, Georgia, through the Gulf states to the valley of the Neches river, Texas, and through Arkansas and southern and southeastern Missouri to the valley of the lower Wabash river, Illinois.

A large tree, 21 to 30 meters in height, with a trunk 0.90 to 1.20 meter in diameter; deep swamps and river bottoms subject to frequent overflow; one of the largest and most common trees of the bottom lands of the lower Mississippi river basin, and reaching its greatest development in the cypress swamps of western Louisiana and eastern Texas, near the coast.

Wood light, soft, not strong, close-grained, compact, unwedgeable; medullary rays numerous, thin; color, light brown, or often nearly white; specific gravity, 0.5194; ash, 0.70; used in turnery, largely for woodenware, broom handles, and wooden shoes; that of the root for the floats of nets, etc., as a substitute for cork.

CAPRIFOLIACEÆ.

156.—*Sambucus glauca*, Nuttall;

Torrey & Gray, Fl. N. America, ii, 13.—Walpers, Rep. ii, 453.—Torrey in Pacific R. R. Rep. vi, 12; Ives' Rep. 15; Bot. Mex. Boundary Survey, 71.—Gray in Smithsonian Contrib. v, 66; Proc. Am. Acad. vii, 387; Syn. Fl. N. America, i², 9.—Watson in King's Rep. v, 134.—Vasey, Cat. Forest Trees, 16.—Brewer & Watson, Bot. California, i, 278.—Hall in Coulter's Bot. Gazette, 88.—Rothrock in Wheeler's Rep. vi, 135, 363.

S. Californica, Hort.—Koch, Dendrologie, ii, 72.

? *S. Mexicana*, Newberry in Pacific R. R. Rep. vi, 75 [not Presl].

ELDER.

Valley of the Fraser river and Vancouver's island, British Columbia, southward through California to the Mexican boundary, extending west to the Blue mountains of Oregon and the Wahsatch range, Utah.

A small tree, sometimes 8 to 9 meters in height, with a trunk 0.30 to 0.45 meter in diameter, or toward its northern limits reduced to a large shrub; confined to valleys, in dry, gravelly soil.

Wood light, soft, weak, coarse-grained, checking in drying; medullary rays numerous, rather conspicuous; color, yellow tinged with brown, the sap-wood lighter; specific gravity, 0.5937; ash, 1.57.

The large blue-black fruit edible and sometimes cooked.

157.—*Sambucus Mexicana*, Presl,

Hort. Hænk.—De Candolle, Prodr. iv, 322.—Don, Miller's Dict. iii, 437.—London, Arboretum, ii, 1030.—Gray in Smithsonian Contrib. v, 66; Syn. Fl. N. America, i², 9.—Torrey in Pacific R. R. Rep. iv, 95; Bot. Mex. Boundary Survey, 71.—Brewer & Watson, Bot. California, i, 278.—Rothrock in Wheeler's Rep. vi, 135.—Hemsley, Bot. Am.-Cent. ii, 1.

S. glauca, Benthams, Pl. Hartweg. 313 [not Nuttall].

S. velatina, Durand & Hilgard in Jour. Philadelphia Acad. new ser. iii, 39.

ELDER.

Valley of the Nueces river (San Patricio), south and west along the southern boundary of the United States to Posa creek, Kern county, California, and southward into Mexico.

A small tree, sometimes 6 meters in height, with a trunk 0.15 to 0.25 meter in diameter; bottom lands, in moist, gravelly loam.

Wood light, soft, rather coarse-grained, compact; medullary rays numerous, thin, conspicuous; color, light brown, the sap-wood lighter; specific gravity, 0.4614; ash, 2.00.

158.—*Viburnum Lentago*, Linnaeus,

Spec. 1 ed. 268.—Marshall, *Arbustum*, 160.—Wangenheim, *Amer.* 100.—Walter, *Fl. Caroliniana*, 116.—Aiton, *Hort. Kew.* i, 372; 2 ed. ii, 168.—Willdenow, *Spec.* i, 1491; *Enum.* 327; *Berl. Baumz.* 531.—Nouveau Duhamel, ii, 129.—Schkuhr, *Handb.* 234.—Michaux, *Fl. Bor.-Am.* i, 178.—Persoon, *Syn.* i, 327.—Desfontaines, *Hist. Arb.* i, 344.—Poiret in Lamarck, *Dict.* viii, 658.—Pursh, *Fl. Am. Sept.* i, 201.—Barton, *Prodr. Fl. Philadelph.* 40.—Eaton, *Manual*, 34; 6 ed. 387.—Nuttall, *Genera*, i, 202.—Hayne, *Dend. Fl.* 37.—Roemer & Schultes, *Syst.* vi, 637.—Elliott, *Sk.* i, 365.—Torrey, *Fl. U. S.* i, 318; *Compend. Fl. N. States*, 138; *Fl. N. York*, i, 305.—Watson, *Dend. Brit.* i, t. 21.—Sprengel, *Syst.* i, 934.—Guimpel, Otto & Hayne, *Abb. Holz.* 125, t. 102.—De Candolle, *Prodr.* iv, 325.—Hooker, *Fl. Bor.-Am.* i, 279.—Beck, *Bot.* 156.—Don, *Miller's Dict.* iii, 440.—Spach, *Hist. Veg.* viii, 311.—London, *Arboretum*, ii, 1033, f. 780.—Dietrich, *Syn.* ii, 1011.—Eaton & Wright, *Bot.* 473.—Torrey & Gray, *Fl. N. America*, ii, 15.—Bigelow, *Fl. Boston.* 3 ed. 123.—Penn. *Cycl.* xxvii, 294.—Emerson, *Trees Massachusetts*, 364; 2 ed. ii, 412.—Darlington, *Fl. Cestrica*, 3 ed. 115.—Darby, *Bot. S. States*, 342.—Chapman, *Fl. S. States*, 171.—Wood, *Cl. Book*, 398; *Bot. & Fl.* 147.—Engelmann in *Trans. Am. Phil. Soc. new ser.* xii, 194; *Trans. St. Louis Acad.* ii, 269.—Gray, *Manual N. States*, 5 ed. 206; *Syn. Fl. N. America*, i², 12.—Koch, *Dendrologie*, ii, 62.—Young, *Bot. Texas*, 309.—Vasey, *Cat. Forest Trees*, 16.—Macoun in *Rep. Geological Surv. Canada*, 1875-76, 198.—Ridgway in *Proc. U. S. Nat. Mus.* 1882, 68.

SHEEPBERRY. NANNYBERRY.

Southern shores of Hudson bay west in British America to about longitude 102°, south through the northern states to southern Indiana and Saint Louis county, Missouri, and along the Alleghany mountains to northern Georgia.

A small tree, 6 to 9 meters in height, with a trunk sometimes 0.15 to 0.25 meter in diameter; rocky ridges and along borders of streams and swamps, in rich, moist soil; most common and reaching its greatest development far north.

Wood heavy, hard, close-grained, compact, emitting a disagreeable odor; medullary rays thin, barely distinguishable; color, dark orange-brown, the sap-wood nearly white; specific gravity, 0.7303; ash, 0.29.

159.—*Viburnum prunifolium*, Linnaeus,

Spec. 1 ed. 268.—Marshall, *Arbustum*, 160.—Wangenheim, *Amer.* 98.—Walter, *Fl. Caroliniana*, 116.—Aiton, *Hort. Kew.* i, 371; 2 ed. ii, 167.—Willdenow, *Spec.* i, 1487; *Enum.* 326; *Berl. Baumz.* 530.—Abbot, *Insects Georgia*, ii, 53.—Nouveau Duhamel, ii, 128, t. 38.—Schkuhr, *Handb.* 233.—Michaux, *Fl. Bor.-Am.* i, 178.—Persoon, *Syn.* i, 326.—Desfontaines, *Hist. Arb.* i, 344.—Poiret in Lamarck, *Dict.* viii, 653.—Pursh, *Fl. Am. Sept.* i, 201.—Barton, *Prodr. Fl. Philadelph.* 39; *Compend. Fl. Philadelph.* i, 151.—Nuttall, *Genera*, i, 202.—Roemer & Schultes, *Syst.* vi, 631.—Hayne, *Dend. Fl.* 37.—Torrey, *Fl. U. S.* i, 318; *Compend. Fl. N. States*, 138.—Elliott, *Sk.* i, 365.—Sprengel, *Syst.* i, 933.—Guimpel, Otto & Hayne, *Abb. Holz.* 125, t. 101.—Watson, *Dend. Brit.* i, t. 23.—Audubon, *Birds*, t. 23.—De Candolle, *Prodr.* iv, 325.—Beck, *Bot.* 156.—Don, *Miller's Dict.* iii, 440.—Spach, *Hist. Veg.* viii, 312.—London, *Arboretum*, ii, 1034, t. 193.—Hooker, *Fl. Bor.-Am.* ii, 279.—Torrey & Gray, *Fl. N. America*, ii, 14.—Walpers, *Rep.* ii, 451.—Darlington, *Fl. Cestrica*, 3 ed. 115.—Darby, *Bot. S. States*, 342.—Chapman, *Fl. S. States*, 171.—Wood, *Cl. Book*, 398; *Bot. & Fl.* 147.—Gray, *Manual N. States*, 5 ed. 206; *Syn. Fl. N. America*, i², 12.—Engelmann in *Trans. St. Louis Acad.* ii, 269.—Koch, *Dendrologie*, ii, 62.—Young, *Bot. Texas*, 309.—Vasey, *Cat. Forest Trees*, 16.—Ridgway in *Proc. U. S. Nat. Mus.* 1882, 68.—Watson in *Proc. Am. Acad.* xviii, 96.

V. pyrifolium, Poiret in Lamarck, *Dict.* v, 658.—Pursh, *Fl. Am. Sept.* i, 201.—Nuttall, *Genera*, i, 202.—Barton, *Compend. Fl. Philadelph.* i, 152.—Roemer & Schultes, *Syst.* vi, 631.—Hayne, *Dend. Fl.* 37.—Watson, *Dend. Brit.* i, t. 22.—Desfontaines, *Hist. Arb.* i, 345; *Cat. Hort. Paris*, 3 ed. 404.—De Candolle, *Prodr.* iv, 325.—Beck, *Bot.* 156.—London, *Arboretum*, ii, 1034, f. 781, 782.—Bigelow, *Fl. Boston*, 3 ed. 123.

V. prunifolium, var. *ferrugineum*, Torrey & Gray, *Fl. N. America*, ii, 15.

BLACK HAW. STAG BUSH.

Fairfield county, Connecticut, valley of the lower Hudson river (Fishkill landing), south to Hernando county, Florida, and the valley of the Colorado river, Texas, west to Missouri, Arkansas, and the Indian territory.

A small tree, sometimes 6 to 9 meters in height, with a trunk rarely exceeding 0.15 meter in diameter, or at the north generally reduced to a low, much-branched shrub; usually on rocky hillsides, in rich soil.

Wood heavy, very hard, strong, brittle, close-grained, liable to check in drying; medullary rays numerous, very obscure; color, brown tinged with red, the sap-wood nearly white; specific gravity, 0.8332; ash, 0.52.

The edible fruit sweet and insipid; the tonic and astringent bark somewhat used in the treatment of uterine disorders in the form of decoctions or fluid extracts (*Boston Med. and Surg. Jour.* October 10, 1867.—*U. S. Dispensatory*, 14 ed. 1783.—*Nat. Dispensatory*, 2 ed. 1821).

RUBIACEÆ.

160.—*Exostemma Caribæum*, Rømer & Schultes,

Syst. v, 18.—Sprengel, Syst. i, 705.—De Candolle, Prodr. iv, 359.—Don, Miller's Dict. iii, 481.—Dietrich, Syn. i, 722.—Spach, Hist. Veg. viii, 395.—Torrey & Gray, Fl. N. America, ii, 36.—Chapman, Fl. S. States, 180.—Grisebach, Fl. British West Indies, 324.—Guibourt, Hist. Drogues, 7 ed. iii, 187, f. 628.—Gray, Syn. Fl. N. America, i², 23.

Cinchona Caribæa, Jacquin, Stirp. Amer. t. 176, f. 65.—Gærtner, Fruct. i, 169, t. 33.—Aiton, Hort. Kew. i, 228; 2 ed. i, 372.—Lambert, Cinchona, 38, t. 12 (excl. syn.).—Andrews, Bot. Rep. vii, t. 481.

Cinchona Jamaicensis, Wright in Trans. Royal Soc. lxvii, 504, t. 10.

Semi-tropical Florida, on the southern keys; through the West Indies.

A small tree, sometimes 7 meters in height, with a trunk 0.20 to 0.30 meter in diameter.

Wood very heavy, exceedingly hard, strong, close-grained, checking in drying, satiny, susceptible of a beautiful polish; medullary rays numerous, very obscure; color, light brown, beautifully streaked with different shades of yellow and brown, the sap-wood clear, rich yellow; specific gravity, 0.9310; ash, 0.23.

161.—*Pinckneya pubens*, Michaux,

Fl. Bor.-Am. i, 103, t. 13.—Willdenow, Enum. Suppl. 30.—Aiton, Hort. Kew. 2 ed. i, 372.—Michaux f. Hist. Arb. Am. ii, 276, t. 24; N. American Sylva, i, 180, t. 49.—Pursh, Fl. Am. Sept. i, 158.—Nuttall, Genera, ii, 37.—Barton, Fl. N. America, i, 25, t. 7.—Sprengel, Syst. i, 705.—Elliott, Sk. i, 269.—Rafinesque, Med. Bot. ii, 57, t. 72.—De Candolle, Prodr. iv, 366.—Audubon, Birds, t. 165.—Eaton, Manual, 6 ed. 263.—Don, Miller's Dict. iii, 486.—Lindley, Fl. Med. 433.—Spach, Hist. Veg. viii, 400.—Eaton & Wright, Bot. 357.—Torrey & Gray, Fl. N. America, ii, 37.—Browne, Trees of America, 354.—Griffith, Med. Bot. 365, f. 174.—Darby, Bot. S. States, 347.—Cooper in Smithsonian Rep. 1858, 253.—Chapman, Fl. S. States, 179.—Wood, Cl. Book, 401; Bot. & Fl. 150.—Porcher, Resources S. Forests, 404.—Vasey, Cat. Forest Trees, 17.—Gray, Syn. Fl. N. America, i², 23.

Cinchona Caroliniana, Poiret in Lamarck, Dict. vi, 40.

P. pubescens, Persoon, Syn. i, 197.—Gærtner f. Fruct. Suppl. 81, t. 194, f. 3.

GEORGIA BARK.

South Carolina, near the coast; basin of the upper Apalachicola river in Georgia and Florida.

A small tree, 6 to 9 meters in height, with a trunk 0.15 to 0.30 meter in diameter; borders of streams, in low, sandy swamps; rare.

Wood light, soft, weak, close-grained, checking badly in drying; layers of annual growth clearly marked by four to six rows of large open ducts; medullary rays few, obscure; color, brown, the sap-wood lighter; specific gravity, 0.5350; ash, 0.41.

Infusions of the bark are successfully used in the treatment of intermittent fever, as a substitute for cinchona (*U. S. Dispensatory*, 14 ed. 1734).

162.—*Genipa clusiæfolia*, Grisebach,

Fl. British West Indies, 317.—Gray, Syn. Fl. N. America, i², 29.

Gardenia clusiæfolia, Jacquin, Coll. Appx. 37, t. 4, f. 3.—Persoon, Syn. i, 199.—De Candolle, Prodr. iv, 381; Dietrich, Syn. i, 796.

Randia clusiæfolia, Chapman, Fl. S. States, 179.—Vasey, Cat. Forest Trees, 17.

SEVEN-YEAR APPLE.

Semi-tropical Florida, on the southern keys; in the West Indies.

A small, much-branched, knotty tree, sometimes 6 meters in height, with a trunk rarely exceeding 0.10 meter in diameter, or in Florida more often a shrub; saline shores.

Wood very heavy, hard, close-grained, compact, susceptible of a beautiful polish; medullary rays numerous, thin; color, rich dark brown shaded with orange, the sap-wood light yellow; specific gravity, 1.0316; ash, 1.06.

The large insipid fruit popularly but incorrectly supposed to require seven years in which to ripen.

163.—*Guettarda elliptica*, Swartz,

Prodr. 59; Fl. Ind. Occ. i, 634.—Lamarck, Ill. ii, 218.—Persoon, Syn. i, 200.—Poiret, Suppl. ii, 859.—Römer & Schultes, Syst. iv, 412.—De Candolle, Prodr. iv, 457.—Dietrich, Syn. i, 787.—Don, Miller's Dict. iii, 551.—Torrey & Gray, Fl. N. America, ii, 35.—Grisebach, Fl. British West Indies, 332.—Gray, Syn. Fl. N. America, i³, 30.

G. Blodgettii, Shuttleworth in herb.—Chapman, Fl. S. States, 172.—Vasey, Cat. Forest Trees, 17.

Semi-tropical Florida, on the southern keys; through the West Indies.

A small tree, 4 to 7 meters in height, with a trunk rarely exceeding 0.20 meter in diameter.

Wood heavy, hard, very close-grained, checking in drying, satiny, susceptible of a beautiful polish, containing numerous scattered small open ducts; medullary rays numerous, thin; color, light brown tinged with red; specific gravity, 0.8337; ash, 1.05.

ERICACEÆ.

164.—*Vaccinium arboreum*, Marshall,

Arbustum, 157.—Michaux, Fl. Bor.-Am. i, 230.—Persoon, Syn. i, 479.—Desfontaines, Hist. Arb. i, 270.—Pursh, Fl. Am. Sept. i, 285.—Nuttall, Genera, i, 263.—Elliott, Sk. i, 495.—Don, Miller's Dict. iii, 853.—London, Arboretum, ii, 1159.—De Candolle, Prodr. vii, 567.—Dietrich, Syn. ii, 1264.—Darby, Bot. S. States, 414.—Loddiges, Bot. Cab. t. 1885.—Walpers, Ann. ii, 1096.—Chapman, Fl. S. States, 259.—Wood, Cl. Book, 482; Bot. & Fl. 198.—Lesquereux in Owen's 2d Rep. Arkansas, 373.—Young, Bot. Texas, 369.—Gray, Hall's Pl. Texas, 15; Syn. Fl. N. America, iii, 20.—Vasey, Cat. Forest Trees, 71.

V. mucronatum, Walter, Fl. Caroliniana, 139 [not Linnæus].

V. diffusum, Aiton, Hort. Kew. ii, 356.—Bot. Mag. t. 1607.—Koch, Dendrologie, ii, 96.

Batodendron arboreum, Nuttall in Trans. Am. Phil. Soc. 2 ser. viii, 261; Sylva, iii, 43; 2 ed. ii, 111.

PARKLEBERRY.

North Carolina, south near the coast to Hernando county, Florida, through the Gulf states, and from southern Illinois and southern Missouri south through Arkansas and eastern Texas to the shores of Matagorda bay.

A small tree, 7 to 9 meters in height, with a trunk rarely 0.25 meter in diameter, or toward its northern limits often reduced to a low shrub; very common throughout the pine belt of the Gulf states along the larger ponds and streams, in moist, sandy soil, and reaching its greatest development in eastern Texas, near the coast.

Wood heavy, hard, very close-grained, compact, liable to twist in drying, satiny, susceptible of a beautiful polish; medullary rays numerous, broad, conspicuous; color, light brown tinged with red, the sap-wood hardly distinguishable; specific gravity, 0.7610; ash, 0.39; somewhat used in turnery in the manufacture of small handles, etc.

165.—*Andromeda ferruginea*, Walter,

Fl. Caroliniana, 138.—Aiton, Hort. Kew. ii, 67; 2 ed. iii, 52.—Willdenow, Sp. ii, 609.—Michaux, Fl. Bor.-Am. i, 252.—Nouveau Duhamel, i, 190.—Ventenat, Hort. Malmaison, 80, t. 80.—Persoon, Syn. i, 480.—Desfontaines, Hist. Arb. i, 257.—Pursh, Fl. Am. Sept. i, 292.—Elliott, Sk. i, 489.—Darby, Bot. S. States, 420.—Chapman, Fl. S. States, 263.—Wood, Cl. Book, 488; Bot. & Fl. 202.—Gray, Syn. Fl. N. America iii, 33.

A. rhomboidalis, Nouveau Duhamel, i, 192.

A. ferruginea, var. *arborescens*, Michaux, Fl. Bor.-Am. i, 252.

A. ferruginea, var. *fruticosa*, Michaux, Fl. Bor. Am. i, 252.

A. rigida, Pursh, Fl. Am. Sept. i, 292.—Loddiges, Bot. Cab. t. 430.

Lyonia ferruginea, Nuttall, Genera, i, 266.—Don, Miller's Dict. iii, 830.—London, Arboretum, ii, 1109.—Dietrich, Syn. ii, 1399.—De Candolle, Prodr. vii, 600.—Koch, Dendrologie, ii, 122.

Lyonia rigida, Nuttall, Genera, i, 266.—Don, Miller's Dict. iii, 830.—De Candolle, Prodr. vii, 600.

South Carolina to northern Florida, near the coast.

A small tree, in rich hummocks, 6 to 9 meters in height, with a trunk 0.15 to 0.25 meter in diameter, often crooked or semi-prostrate; or in sandy pine-barren soil reduced to a low shrub, 0.60 to 0.90 meter in height; the leaves varying greatly in shape, venation, etc.

Wood heavy, hard, not strong, very close-grained, checking in drying, satiny, susceptible of a beautiful polish; medullary rays numerous, thin; color, bright brown tinged with red, the sap-wood a little lighter; specific gravity, 0.7500; ash, 0.46.

166.—*Arbutus Menziesii*, Pursh,

Fl. Am. Sept. i, 282.—Sprengel, Syst. ii, 286.—Don, Miller's Dict. iii, 834.—London, Arboretum, ii, 1122.—De Candolle, Prodr. vii, 582.—Dietrich, Syn. ii, 1387.—Hooker, Fl. Bor.-Am. ii, 36.—Hooker & Arnott, Bot. Beechey, 143.—Nuttall, Sylva, iii, 42, t. 95; 2 ed. ii, 109, t. 95.—Torrey in Pacific R. R. Rep. iv, 116; Bot. Wilkes Exped. 378.—Newberry in Pacific R. R. Rep. vi, 23, 79, f. 22.—Cooper in Smithsonian Rep. 1858, 260; Pacific R. R. Rep. xii^a, 29, 66.—Lyll in Jour. Linnæan Soc. vii, 131.—Gray in Proc. Am. Acad. vii, 393; Bot. California, i, 452, in part; Syn. Fl. N. America, ii¹, 27, in part.—Vasey, Cat. Forest Trees, 17.—Hall in Coulter's Bot. Gazette, ii, 88.—Macoun in Geological Rep. Canada, 1875-76, 203.—G. M. Dawson in Canadian Nat. new ser. ix, 331.—Hemsley, Bot. Am.-Cent. ii, 276.

A. procera, Douglas in Lindley's Bot. Reg. xxi, t. 1753.—London, Arboretum, ii, 1121.—De Candolle, Prodr. vii, 582.—Dietrich, Syn. ii, 1387.—Paxton, Mag. Bot. ii, 147 & t.—Walpers, Rep. vi, 416.

A. laurifolia, Lindley, Bot. Reg. xxx, t. 67.—Hooker, Fl. Bor.-Am. ii, 36.

MADROÑA.

Islands of British Columbia, from Seymour narrows southward through Washington territory and Oregon, near the coast, and through the Coast ranges of California to the Santa Lucia mountains.

A small tree, sometimes 15 to 25 meters in height, with a trunk 0.90 to 1.20 meter in diameter, or, exceptionally, much larger (the great specimen near San Rafael, Marin county, California, 6.85 meters in circumference 2 meters from the ground); south of San Francisco bay much smaller, often reduced to a low shrub; hillsides, in rich soil.

Wood heavy, hard, strong, close-grained, checking in drying; medullary rays numerous, conspicuous; color, light brown shaded with red, the sap-wood lighter; specific gravity, 0.7052; ash, 0.40; largely used in the manufacture of gunpowder, the bark in tanning.

167.—*Arbutus Kalapensis*, HBK.

Nov. Gen. & Spec. iii, 281.—Sprengel, Syst. ii, 286.—Don, Miller's Dict. iii, 834.—Hooker, Icon. i, t. 27.—Bentham, Pl. Hartweg. 66.—De Candolle, Prodr. vii, 583.—Dietrich, Syn. ii, 1388.—Walpers, Ann. ii, 1105.—Jour. Hort. Soc. London, v, 192 & t.

?*A. variens*, Bentham, Pl. Hartweg. 77.—Paxton, Brit. Fl. Gard. ii, 118.—Hemsley, Bot. Am.-Cent. ii, 277.

?*A. macrophylla*, Martens & Galeotti in Bull. Acad. Brux. ix, 9.—Walpers, Rep. ii, 725.

A. Menziesii, Gray in Bot. California, i, 452, in part; Syn. Fl. N. America, ii¹, 27, in part.—Rothrock in Wheeler's Rep. vi, 25, 183 [not Pursh].

Southern Arizona, Santa Rita mountains, between 4,500 and 7,000 feet elevation; southward through northern Mexico.

A small tree, with white, scaly bark, 9 to 12 meters in height, with a trunk 0.45 to 0.60 meter in diameter; dry, gravelly slopes; large specimens generally hollow and defective.

Wood heavy, soft, not strong, brittle, close-grained, checking badly in drying, susceptible of a good polish; medullary rays numerous, obscure; color, light brown tinged with red, the sap-wood lighter; specific gravity, 0.7099; ash, 0.25.

168.—*Arbutus Texana*, Buckley,

Proc. Philadelphia Acad. 1861, 460.—Gray in Proc. Philadelphia Acad. 1862, 165.—Young, Bot. Texas, 370.

A. Menziesii, Gray in Bot. California, i, 452, in part; Syn. Fl. N. America, ii¹, 27, in part.

?*A. Kalapensis*, Watson in Proc. Am. Acad. xviii, 111.

Western Texas, Hays and Travis counties (*Buckley*), west to the Guadalupe and Eagle mountains (*Havard*), and southward, probably into northern Mexico.

A small tree, 5 to 6 meters in height, with a trunk 0.15 to 0.25 meter in diameter; dry limestone hills and ridges; rare.

Wood heavy, hard, close-grained, compact; medullary rays numerous, obscure; color, brown, the sap-wood lighter, tinged with red; specific gravity, 0.7500; ash, 0.51; used in turnery, the manufacture of mathematical instruments, etc.

NOTE.—The synonymy and specific position of the Mexican species of *Arbutus* which reach the southern boundary of the United States are still obscure, and cannot be well elucidated with the existing knowledge of the Mexican flora.

169.—*Oxydendrum arboreum*, De Candolle,

Prodr. vii, 601.—Dietrich, Syn. ii, 1389.—Cooper in Smithsonian Rep. 1858, 253.—Chapman, Fl. S. States, 263.—Lesquereux in Owen's 2d Rep. Arkansas, 372.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 79.—Wood, Cl. Book, 489; Bot. & Fl. 203.—Gray, Manual N. States, 5 ed. 296; Syn. Fl. N. America, ii¹, 33.—Koch, Dendrologie, ii, 128.—Vasey, Cat. Forest Trees, 17.—Nat. Dispensatory, 2 ed. 798.

Andromeda arborea, Linnæus, Spec. 1 ed. 394.—Lamarck, Dict. i, 158.—Marshall, Arbustum, 7.—Wangenheim, Amer. 105.—Walter, Fl. Caroliniana, 138.—Aiton, Hort. Kew. ii, 69; 2 ed. iii, 53.—Willdenow, Spec. ii, 612; Enum. 452; Berl. Baumz. 31.—Michaux, Fl. Bor.-Am. i, 255.—Nouveau Duhamel, i, 178.—Bot. Mag. t. 905.—Desfontaines, Hist. Arb. i, 257.—Michaux f. Hist. Arb. Am. iii, 222, t. 7; N. American Sylva, 3 ed. ii, 126, t. 85.—Pursh, Fl. Am. Sept. i, 295.—Nuttall, Genera, i, 265.—Elliott, Sk. i, 491.—Barton, Fl. N. America, i, 105, t. 30.—Hayne, Dend. Fl. 59.—Torrey, Fl. U. S. i, 420; Compend. Fl. N. States, 182.—Sprengel, Syst. ii, 291.—Gray, Manual N. States, 1 ed. 266.—Darby, Bot. S. States, 419.—Porcher, Resources S. Forests, 379.

Andromeda arborescens, Persoon, Syn. i, 480.—Willdenow, Enum. 453.—Loddiges, Bot. Cab. t. 1210.

Lyonia arborea, Don in Edinburgh Phil. Jour. xvii, 159.—Don, Miller's Dict. iii, 831.—London, Arboretum, ii, 1111.—Spach, Hist. Veg. ix, 486.—Browne, Trees of America, 356.

SORREL TREE. SOUR WOOD.

Western Pennsylvania, southward along the Alleghany mountains to western Florida and the eastern shores of Mobile bay, west to middle Tennessee and through the upper regions of the Gulf states to western Louisiana.

A small tree, 12 to 18 meters in height, with a trunk 0.25 to 0.35 meter in diameter; usually in rather dry, gravelly soil.

Wood heavy, hard, very close-grained, compact, susceptible of a beautiful polish; medullary rays numerous, thin; color, brown tinged with red, the sap-wood somewhat lighter; specific gravity, 0.7458; ash, 0.37; used for the handles of tools, bearings of machinery, etc.

170.—*Kalmia latifolia*, Linnæus,

Spec. 1 ed. 301.—Kalm, Travels, English ed. i, 335.—Marshall, Arbustum, 72.—Lamarck, Dict. ii, 345; Ill. ii, 487, t. 363, f. 1.—Gartner, Fruct. i, 305, t. 63, f. 7.—Wangenheim, Amer. 64, t. 24, f. 50.—Walter, Fl. Caroliniana, 138.—Aiton, Hort. Kew. ii, 64; 2 ed. iii, 47.—Lamarck, Ill. 487, t. 363, f. 1.—Abbot, Insects Georgia, i, t. 87.—Willdenow, Spec. ii, 600; Enum. 450; Berl. Baumz. 202.—Schkuhr, Handb. 359, t. 116.—Michaux, Fl. Bor.-Am. i, 258.—Persoon, Syn. i, 477.—Desfontaines, Hist. Arb. i, 220.—Robin, Voyages, iii, 419.—Michaux f. Hist. Arb. Am. iii, 144, t. 4; N. American Sylva, 3 ed. ii, 62, t. 67.—Pursh, Fl. Am. Sept. i, 296.—Barton, Prodr. Fl. Philadelph. 49.—Eaton, Manual, 47; 6 ed. 195.—Bigelow, Med. Bot. i, 113, t. 13; Fl. Boston. 3 ed. 179.—Nuttall, Genera, i, 267.—Hayne, Dend. Fl. 54.—Elliott, Sk. i, 481.—Torrey, Fl. U. S. i, 422; Compend. Fl. N. States, 182.—Sprengel, Syst. ii, 293.—Audubon, Birds, t. 55.—Rafinesque, Med. Bot. ii, 16, t. 57.—Sertum Botanicum, iv & t.—Beck, Bot. 219.—Don, Miller's Dict. iii, 850.—Lindley, Fl. Med. 380.—London, Arboretum, ii, 1151, f. 959.—De Candolle, Prodr. vii, 729.—Spach, Hist. Veg. ix, 498, t. 139.—Hooker, Fl. Bor.-Am. ii, 41.—Dietrich, Syn. ii, 1407.—Browne, Trees of America, 363.—Emerson, Trees Massachusetts, 392; 2 ed. ii. 443 & t.—Griffith, Med. Bot. 428, f. 192.—Darlington, Fl. Cestrica, 3 ed. 172.—Darby, Bot. S. States, 420.—Cooper in Smithsonian Rep. 1858, 253.—Chapman, Fl. S. States, 264.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 99.—Lesquereux in Owen's 2d Rep. Arkansas, 373.—Wood, Cl. Book, 484; Bot. & Fl. 200.—Porcher, Resources S. Forests, 381.—Gray, Manual N. States, 5 ed. 296; Syn. Fl. N. America, ii¹, 38.—Koch, Dendrologie, ii, 152.—Vasey, Cat. Forest Trees, 17.—London Garden, xxii, 6, t. 343.

LAUREL. CALICO BUSH. SPOON WOOD. IVY.

New Brunswick and the northern shores of lake Erie, south to western Florida, and through the Gulf states to western Louisiana and the valley of the Red river, Arkansas (Hot Springs, *Letterman*).

A small tree, sometimes 9 to 12 meters in height, with a trunk 0.30 to 0.60 meter in diameter, or more often a low shrub; rich woodlands; most common and reaching its greatest development in the southern Alleghany mountains, here often forming dense, impenetrable thickets.

Wood heavy, hard, strong, brittle, close-grained, compact; principal medullary rays broad, dark brown, conspicuous, intermediate rays numerous, thin, inconspicuous; color, brown tinged with red, the sap-wood somewhat lighter; specific gravity, 0.7160; ash, 0.41; used for tool handles, in turnery, and for fuel.

The leaves, buds, and fruit, reputed poisonous to cattle, are occasionally used medicinally (*U. S. Dispensatory*, 14 ed. 1682.—*Nat. Dispensatory*, 2 ed. 798).

171.—*Rhododendron maximum*, Linnæus,

Spec. 1 ed. 391.—Marshall, Arbustum, 127.—Gärtner, Fruct. i, 304, t. 63, f. 6.—Wangenheim, Amer. 63, t. 22, f. 49.—Aiton, Hort. Kew. ii, 67; 2 ed. iii, 50.—Mönch, Meth. 45.—Lamarck, Dict. vi, 365; Ill. ii, 448, t. 364, f. 1.—B. S. Barton, Coll. i, 18.—Willdenow, Spec. ii, 606; Enum. 451; Berl. Baumz. 357.—Nouveau Duhamel, ii, 141.—Michaux, Fl. Bor.-Am. i, 259.—Schkuhr, Handb. 362.—Persoon, Syn. i, 478.—Desfontaines, Hist. Arb. i, 221.—Bot. Mag. t. 951.—Michaux f. Hist. Arb. Am. iii, 144, t. 4; N. American Sylva, 3 ed. ii, 64, t. 68.—Pursh, Fl. Am. Sept. i, 297.—Eaton, Manual, 47; 6 ed. 301.—Nuttall, Genera, i, 268.—Bigelow, Med. Bot. iii, 101, t. 51; Fl. Boston. 3 ed. 178.—Elliott, Sk. i, 483.—Hayne, Dend. Fl. 57.—Torrey, Fl. U. S. i, 426; Compend. Fl. N. States, 184.—Sprengel, Syst. ii, 292.—Audubon, Birds, t. 103.—Beck, Bot. 220.—Don, Miller's Dict. iii, 843.—London, Arboretum, ii, 1134, f. 932.—De Candolle, Prodr. vii, 722.—Hooker, Fl. Bor.-Am. ii, 43.—Spach, Hist. Veg. ix, 503.—Dietrich, Syn. ii, 1404.—Eaton & Wright, Bot. 391.—Browne, Trees of America, 359.—Emerson, Trees Massachusetts, 384; 2 ed. ii, 435 & t.—Griffith, Med. Bot. 428.—Darlington, Fl. Cestrica, 3 ed. 171.—Darby, Bot. S. States, 421.—Cooper in Smithsonian Rep. 1858, 253.—Chapman, Fl. S. States, 265.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 97.—Lesquereux in Owen's 2d Rep. Arkansas, 373.—Wood, Cl. Book, 491; Bot. & Fl. 204.—Porscher, Resources S. Forests, 380.—Gray, Manual N. States, 5 ed. 300; Syn. Fl. N. America iii, 42.—Koch, Dendrologie, ii, 169.—Vasey, Cat. Forest Trees, 17.

R. procerum, Salisbury, Prodr. 287.

R. maximum, var. *roseum*, Pursh, Fl. Am. Sept. i, 297.—Elliott, Sk. i, 484.

R. maximum, var. *album*, Pursh, Fl. Am. Sept. i, 297.—Elliott, Sk. i, 484.

R. maximum, var. *purpureum*, Pursh, Fl. Am. Sept. i, 297.—Elliott, Sk. i, 484.

R. purpureum, Don, Miller's Dict. iii, 843.—London, Arboretum, ii, 1134.—Dietrich, Syn. ii, 1404.

R. Purshii, Don, Miller's Dict. iii, 843.—London, Arboretum, ii, 1135.—Dietrich, Syn. ii, 1404 (var. *album*, Pursh, l. c.).

GREAT LAUREL. ROSE BAY.

Nova Scotia and the northern shores of lake Erie, south through New England, New York, and along the Alleghany mountains to northern Georgia.

A small tree, sometimes 10 to 12 meters in height, with a trunk rarely exceeding 0.30 meter in diameter, or often a tall, straggling shrub; at the north in cold swamps; rare; very common and reaching its greatest development in the southern Alleghany mountains, steep, rocky banks of streams, etc.; never on limestone.

Wood heavy, hard, strong, brittle, close-grained, compact; medullary rays numerous, thin; color, light clear brown, the sap-wood lighter; specific gravity, 0.6303; ash, 0.36; occasionally used in turnery for the handles of tools, etc., and a possible substitute for box-wood in engraving. A decoction of the leaves is occasionally used domestically in the treatment of rheumatism, sciatica, etc.

MYRSINACEÆ.

172.—*Myrsine Rapania*, Roemer & Schultes,

Syst. iv, 509.—Don, Miller's Dict. iv, 10.—Dietrich, Syn. i, 618.—A. De Candolle, Prodr. viii, 97.—Miquel in Martius, Fl. Brasil. ix, 307, t. 50-52.—Gray, Syn. Fl. N. America ii, 65.

Rapania Guyanensis, Aublet, Guian. i, 121, t. 46.—Swartz, Obs. 51; Fl. Ind. Occ. i, 262.—Lamarck, Ill. ii, 48, t. 122, f. 1.

Samara pentandra, Swartz, Obs. 51; Fl. Ind. Occ. i, 262 [not Aiton].

Samara floribunda, Willdenow, Spec. i, 665.—Lamarck, Ill. ii, 46, t. 122, f. 1.

Oaballeria coriacea, Meyer, Prim. Fl. Esseq. 118.

M. Floridana, A. De Candolle in Trans. Linnæan Soc. xvii, 107; Prodr. viii, 98.—Dietrich, Syn. i, 98.—Chapman, Fl. S. States, 277.

M. floribunda, Grisebach, Fl. British West Indies, 393.

Semi-tropical Florida, Indian river southward to the southern keys; through the West Indies to Brazil.

A small tree, in Florida rarely exceeding 8 meters in height, with a trunk 0.10 to 0.15 meter in diameter, or often a shrub; borders of ponds and fresh-water creeks; in the West Indies much larger.

Wood heavy, hard, very close-grained, compact; medullary rays numerous, very conspicuous; color, brown tinged with red and beautifully striped with the darker medullary rays, the sap-wood hardly distinguishable; specific gravity, 0.8341; ash, 0.81.

173.—*Ardisia Pickeringia*, Nuttall,

Sylva, iii, 69, t. 102; 2 ed. ii, 133, t. 102.—A. De Candolle, *Prodr.* viii, 124.—Cooper in *Smithsonian Rep.* 1858, 264.—Chapman, *Fl. S. States*, 277.—Vasey, *Cat. Forest Trees*, 19.—Gray, *Syn. Fl. N. America*, ii¹, 65.—Hemsley, *Bot. Am.-Cent.* ii, 294.

Cyrilla paniculata, Nuttall in *Am. Jour. Sci.* v, 290.

Pickeringia paniculata, Nuttall in *Jour. Philadelphia Acad.* vii, 1.

MARLBERRY. CHERRY.

Semi-tropical Florida, Mosquito inlet to the southern keys, west coast, Caloosa river to cape Romano; in the West Indies and southern Mexico.

A small tree, sometimes 8 meters in height, with a trunk rarely 0.15 meter in diameter, or often a shrub; reaching its greatest development in Florida on the shores of bay Biscayne.

Wood heavy, hard, very close-grained, compact, susceptible of a beautiful polish; medullary rays very numerous, conspicuous; color, rich brown, beautifully marked with the darker medullary rays, the sap-wood a little lighter; specific gravity, 0.8602; ash, 1.85.

174.—*Jacquinia armillaris*, Jacquin,

Amer. 53, t. 39.—Linnaeus, *Spec.* 2 ed. 272.—Aiton, *Hort. Kew.* i, 257; 2 ed. ii, 5.—Lamarck, *Ill.* ii, 46, t. 39.—Vahl, *Eclog.* i, 26.—Swartz, *Obs.* 85.—Willdenow, *Spec.* i, 1064; *Enum.* 246.—Persoon, *Syn.* i, 234.—Römer & Schultes, *Syst.* iv, 490.—Sprengel, *Syst.* i, 668.—Don, *Miller's Dict.* iv, 24.—Dietrich, *Syn.* i, 638.—Bentham, *Bot. Sulphur*, 123.—A. De Candolle, *Prodr.* viii, 149.—Miquel in *Martius, Fl. Brasil.* ix, 282, t. 27.—Cooper in *Smithsonian Rep.* 1858, 265.—Chapman, *Fl. S. States*, 276.—Grisebach, *Fl. British West Indies*, 397.—Seemann, *Jour. Bot.* iii, 279.—Vasey, *Cat. Forest Trees*, 19.—Gray, *Syn. Fl. N. America*, ii¹, 66.—Hemsley, *Bot. Am.-Cent.* ii, 294.

Chrysophyllum Barbasco, Læfing, *Iter.* 204, 277.

JOE WOOD.

Semi-tropical Florida, on the southern keys; rare; through the West Indies to Brazil.

A low, rigid tree, rarely exceeding in Florida 4 meters in height, with a trunk sometimes 0.15 meter in diameter; in the Bahamas and other West Indian islands probably much larger.

Wood heavy, hard, coarse-grained, checking and shrinking badly in drying, containing many scattered large open ducts; medullary rays numerous, broad, conspicuous; color, light clear brown tinged with yellow; specific gravity, 0.6948; ash, 3.45.

The saponaceous leaves sometimes used as a substitute for soap.

SAPOTACEÆ.

175.—*Chrysophyllum oliviforme*, Lamarck,

Dict. i, 552; *Ill.* ii, 42.—Descourtilz, *Fl. Med. Antilles*, ii, 71.—A. De Candolle, *Prodr.* viii, 158.—Grisebach, *Fl. British West Indies*, 396.—Gray, *Syn. Fl. N. America*, ii¹, 67.—Chapman, *Fl. S. States*, *Suppl.* 634.

C. Caneto, *β.* Linnaeus, *Sp.* 3 ed. 278 (excl. syn. Læfing).

C. monopyrenum, Swartz, *Prodr.* 49; *Fl. Ind. Occ.* i, 480.—Persoon, *Syn.* i, 236.—Römer & Schultes, *Syst.* iv 703.—Sprengel, *Syst.* i, 666.—*Bot. Mag.* t. 3303.—Dietrich, *Syn.* i, 638.—Miquel in *Martius, Fl. Brasil.* vii, 94.

C. ferrugineum, Gärtner f. *Fruct. Suppl.* 120, t. 202, f. 1.

C. microphyllum, Chapman in *Coulter's Bot. Gazette*, iii, 9.—Vasey, *Cat. Forest Trees*, 18 [not A. De Candolle].

Semi-tropical Florida, cape Canaveral to the southern keys (Elliott's Key, No-Name Key, Key Largo), west coast, Caloosa river to cape Sable; rare; through the West Indies to Brazil.

A small tree, sometimes 9 meters in height, with a trunk 0.25 to 0.30 meter in diameter.

Wood very heavy, hard, strong, close-grained, checking in drying; medullary rays numerous, not conspicuous; color, light brown shaded with red, the thin sap-wood a little lighter; specific gravity, 0.9360; ash, 1.24.

176.—*Sideroxylon Mastichodendron*, Jacquin,

Coll. ii, t. 17, f. 5.—Lamarck, Ill. ii, 41, t. 120, f. 2.—Gærtner f. Fruct. Suppl. 125.—Sprengel, Syst. i, 666.—Dietrich, Syn. i, 622.—A. De Candolle, Prodr. viii, 181.—Grisebach, Fl. British West Indies, 399.—Gray, Syn. Fl. N. America ii¹, 67.

Bumelia pallida, Swartz, Prodr. 40; Fl. Ind. Occ. 489.

Achras pallida, Poiret in Lamarck, Dict. vi, 533.

Bumelia Mastichodendron, Römer & Schultes, Syst. iv, 493.

S. pallidum, Sprengel, Syst. i, 666.—A. De Candolle, Prodr. viii, 180.—Chapman, Fl. S. States, 274.—Vasey, Cat. Forest Trees, 18.

Bumelia foetidissima, Nuttall, Sylva, iii, 39, t. 94; 2 ed. ii, 108, t. 94.—Cooper in Smithsonian Rep. 266.

MASTIC.

Semi-tropical Florida, cape Canaveral to the southern keys, west coast, cape Romano to cape Sable; in the West Indies.

A tree often 18 meters in height, with a trunk 0.60 to 0.90 meter in diameter; the largest and most valuable tree of semi-tropical Florida; common.

Wood very heavy, exceedingly hard, strong, close-grained, checking in drying, containing few scattered small open ducts; medullary rays numerous, not conspicuous; color, bright orange, the sap-wood yellow; specific gravity, 1.0109; ash, 5.14; not affected by the teredo; largely used in ship- and boat-building.

The dry fruit, of a pleasant subacid flavor, eagerly eaten by animals.

177.—*Dipholis salicifolia*, A. De Candolle,

Prodr. viii, 188 (Delessert, Icon. Mex. ined. t. 40).—Richard, Fl. Cuba, t. 54².—Miquel in Martius, Fl. Brasil. vii, 45, t. 18.—Chapman, Fl. S. States, 274.—Grisebach, Fl. British West Indies, 401.—Vasey, Cat. Forest Trees, 18.—Gray, Syn. Fl. N. America ii¹, 67.

Achras salicifolia, Linnæus, Spec. 2 ed. 470.

Bumelia salicifolia, Swartz, Prodr. 50; Fl. Ind. Occ. i, 491.—Lamarck, Ill. ii, 42.—Willdenow, Spec. i, 1086.—Aiton, Hort. Kew. 2 ed. ii, 12.—Römer & Schultes, Syst. iv, 494.—Dietrich, Syn. i, 621.

Sideroxylon salicifolium, Gærtner f. Fruct. Suppl. 124, t. 202.—Lamarck, Ill. ii, 42.

BUSTIC. CASSADA.

Semi-tropical Florida, bay Biscayne to the southern keys; through the West Indies to Brazil.

A tree sometimes 15 meters in height, with a trunk rarely 0.60 meter in diameter; the large trees hollow and defective; rare.

Wood very heavy, exceedingly hard, very strong, close-grained, compact, checking in drying, susceptible of a beautiful polish, containing many scattered large open ducts; color, dark brown or red, the sap-wood lighter; specific gravity, 0.9316; ash, 0.32.

178.—*Bumelia tenax*, Willdenow,

Spec. i, 1088; Enum. 248; Berl. Baumz. 67.—Aiton, Hort. Kew. 2 ed. ii, 12.—Römer & Schultes, Syst. iv, 496.—Elliott, Sk. i, 288.—Persoon, Syn. i, 237.—Hayne, Dend. Fl. 18.—Sprengel, Syst. i, 664.—Eaton, Manual, 6 ed. 60.—Don, Miller's Dict. iv, 30.—London, Arboretum, ii, 1193, f. 1017.—Dietrich, Syn. i, 621.—Spach, Hist. Veg. ix, 388.—Eaton & Wright, Bot. 162.—Nuttall, Sylva, iii, 35, t. 92; 2 ed. ii, 104, t. 92.—A. De Candolle, Prodr. viii, 196.—Darby, Bot. S. States, 428.—Cooper in Smithsonian Rep. 1858, 253.—Chapman, Fl. S. States, 275.—Wood, Cl. Book, 501; Bot. & Fl. 210.—Vasey, Cat. Forest Trees, 19.—Gray, Syn. Fl. N. America, ii¹, 68.

Sideroxylon tenax, Linnæus, Mant. 48.—Jacquin, Coll. ii, 252.—Lamarck, Dict. i, 245; Ill. ii, 42.—Aiton, Hort. Kew, i, 262.—Swartz, Obs. 91.—Desfontaines, Hist. Arb. i, 204.—Robin, Voyages, iii, 461.

Sideroxylon Carolinense, Jacquin, Obs. iii, 3, t. 54.

Sideroxylon sericeum, Walter, Fl. Caroliniana, 100.

Sideroxylon chrysophylloides, Michaux, Fl. Bor.-Am. i, 123.—Rafinesque, Fl. Ludoviciana, 53.

B. chrysophylloides, Pursh, Fl. Am. Sept. i, 155.—Nuttall, Genera, i, 135.—Watson, Dend. Brit. i, t. 10.

?*B. reclinata*, Chapman, Fl. S. States, 275 [not Ventenat].

North Carolina, south near the coast to cape Canaveral and Cedar Keys, Florida.

A small tree, 6 to 9 meters in height, with a trunk sometimes 0.15 meter in diameter; sandy soil.

Wood heavy, hard, not strong, very close-grained, compact, susceptible of a beautiful polish; well characterized, as in all the North American species, by large open ducts, defining, with several rows, the rings of annual growth, and connected by conspicuous branching groups of similar ducts, giving to a cross-section a beautifully reticulated appearance; medullary rays numerous, thin; color, light brown streaked with white, the sap-wood lighter; specific gravity, 0.7293; ash, 0.78.

179.—*Bumelia lanuginosa*, Persoon.

Syn. i, 237.—Pursh, Fl. Am. Sept. i, 155.—Nuttall, Genera, i, 135.—Roemer & Schultes, Syst. iv, 497.—Elliott, Sk. i, 288.—Eaton, Manual, 6 ed. 60.—Don, Miller's Dict. iv, 30.—London, Arboretum, ii, 1194.—Eaton & Wright, Bot. 162.—A. De Candolle, Prodr. viii, 190.—Darby, Bot. S. States, 428.—Cooper in Smithsonian Rep. 1858, 253.—Chapman, Fl. S. States, 275.—Lesquereux in Owen's 2d Rep. Arkansas, 374.—Wood, Cl. Book, 501; Bot. & Fl. 210.—Gray, Manual N. States, 5 ed. 308; Hall's Pl. Texas, 15; Syn. Fl. N. America, ii, 68.—Young, Bot. Texas, 377.—Vasey, Cat. Forest Trees, 19.

?*Sideroxylon tenax*, Walter, Fl. Caroliniana, 100.

Sideroxylon lanuginosum, Michaux, Fl. Bor.-Am. i, 122.

?*B. oblongifolia*, Nuttall, Genera, i, 135; Sylva, iii, 33; 2 ed. ii, 102.—Sprengel, Syst. i, 664.—Eaton, Manual, 6 ed. 60.—Eaton & Wright, Bot. 162.—Don, Miller's Dict. iv, 30.—London, Arboretum, ii, 1194.—Dietrich, Syn. i, 621.—A. De Candolle, Prodr. viii, 190.—Lesquereux in Owen's 2d Rep. Arkansas, 374.

B. ferruginea, Nuttall, Sylva, iii, 34; 2 ed. ii, 103.

B. tomentosa, A. De Candolle, Prodr. viii, 190.

B. arborea, Buckley in Proc. Philadelphia Acad. 1861, 461.

GUM ELASTIC. SHITTIM WOOD.

Georgia and northern Florida to Mobile bay, Alabama; southern Illinois and southern Missouri, through Arkansas to the valley of the Rio Grande, Texas (Eagle pass, *Havard*) (*B. oblongifolia*).

An evergreen tree, sometimes 18 meters in height, with a trunk 0.90 meter in diameter, or in the Atlantic states much smaller, rarely exceeding 6 meters in height; common and reaching its greatest development in the rich bottom lands of eastern Texas.

A low, depressed form of the sand-hills of the Altamaha river, Georgia, still to be rediscovered, with small leaves and "edible fruit as large as a small date", is var. *macrocarpa*, Gray, Syn. Fl. N. America, iii, 68 (*B. macrocarpa*, Nuttall, Sylva, iii, 37; 2 ed. ii, 106).

Wood heavy, soft, weak, close-grained, very compact, the open ducts conspicuous; medullary rays numerous, thin; color, light brown or yellow, the sap-wood lighter; specific gravity, 0.6544; ash, 1.23; somewhat used in cabinet-making, for which it is well suited.

A clear, very viscid gum exuded from the freshly-cut wood is sometimes used domestically.

180.—*Bumelia spinosa*, A. De Candolle,

Prodr. viii, 191 (Delessert, Icon. Mex. ined. t. 75).—Hemsley, Bot. Am.-Cent. ii, 299.—Watson in Proc. Am. Acad. xviii, 112.

Santa Catalina mountains, Arizona, at an elevation of 2,700 feet (*Pringle*); Parras and Saltillo, Mexico (*Palmer*, No. 787).

A small tree, 6 to 7 meters in height, with a trunk 0.20 to 0.25 meter in diameter; dry, gravelly soil, near water-courses.

Wood heavy, hard, very close-grained, compact, the open ducts conspicuous; medullary rays thin, obscure; color, light rich brown or yellow, the sap-wood lighter; specific gravity, 0.6603; ash, 1.24.

181.—*Bumelia lycioides*, Gartner f.

Fruct. Suppl. 127, t. 120.—Persoon, Syn. i, 237.—Willdenow, Enum. 249; Berl. Baumz. 68.—Pursh, Fl. Am. Sept. i, 237.—Nuttall, Genera, i, 135; Sylva, iii, 31, t. 91; 2 ed. ii, 101, t. 91.—Roemer & Schultes, Syst. iv, 495.—Hayne, Dend. Fl. 19.—Elliott, Sk. i, 287.—Sprengel, Syst. i, 664.—Eaton, Manual, 6 ed. 60.—Don, Miller's Dict. iv, 30.—London, Arboretum, ii, 1193, f. 1016.—Dietrich, Syn. i, 621.—Spach, Hist. Veg. ix, 388.—Eaton & Wright, Bot. 162.—A. De Candolle, Prodr. viii, 189.—Griffith, Med. Bot. 441.—Darby, Bot. S. States, 427.—Cooper in Smithsonian Rep. 1858, 253.—Chapman, Fl. S. States, 275.—Lesquereux in Owen's 2d Rep. Arkansas, 374.—Wood, Cl. Book, 501; Bot. & Fl. 210.—Gray, Manual N. States, 5 ed. 308; Syn. Fl. N. America, iii, 68.—Young, Bot. Texas, 376.—Vasey, Cat. Forest Trees, 19.—Hemsley, Bot. Am.-Cent. ii, 298.

Sideroxylon lycioides, Linnæus, Hort. Cliff. 488 (excl. hab.).—Lamarck, Dict. i, 246; Ill. ii, 42.—Aiton, Hort. Kew. i, 262; 2 ed. ii, 13.—Willdenow, Spec. i, 1090.—Michaux, Fl. Bor.-Am. i, 122.—Pursh, Fl. Am. Sept. i, 155.—Jaume St. Hilaire, Fl. & Pom. Am. Franc. t. 81.

Sideroxylon decandrum, Linnæus, Mant. 48.—Willdenow, Spec. i, 1091.

Syderoxylon læve, Walter, Fl. Caroliniana, 100.

IRON WOOD. SOUTHERN BUCKTHORN.

Coast of Virginia and southern Illinois, south to Mosquito inlet and Caloosa river, Florida, and through southern Missouri, Arkansas, and Texas to the valley of the Rio Concho, Texas.

A small tree, sometimes 9 to 12 meters in height, with a trunk rarely exceeding 0.15 meter in diameter; low, rich soil, or often, in the Atlantic and Gulf states, a low, semi-prostrate shrub, described as—

var. *reclinatum*, Gray, Syn. Fl. N. America, ii¹, 68.

Sideroxylon reclinatum, Michaux, Fl. Bor.-Am. i, 122.

B. reclinata, Ventenat, Choix, t. 22.—Persoon, Syn. i, 237.—Pursh, Fl. Am. Sept. i, 155.—Roemer & Schultes, Syst. iv, 496.—Elliott, Sk. i, 287.—Eaton, Manual, 6 ed. 60.—Dietrich, Syn. i, 621.—Don, Miller's Dict. iv, 30.—Loudon, Arboretum, ii, 1193.—A. De Candolle, Prodr. viii, 190.—Darby, Bot. S. States, 428.—Wood, Cl. Book, 501; Bot. & Fl. 210.

Wood heavy, hard, not strong, close-grained, compact; medullary rays numerous, thin; color, light brown or yellow, the sap-wood lighter; specific gravity, 0.7467; ash, 0.81.

182.—*Bumelia cuneata*, Swartz,

Fl. Ind. Occ. i, 496.—Persoon, Syn. i, 237.—Roemer & Schultes, Syst. iv, 498.—Sprengel, Syst. i, 665.—Don, Miller's Dict. iv, 30.—Dietrich, Syn. i, 621.—Grisebach, Fl. British West Indies, 401.—Gray, Syn. Fl. N. America, ii¹, 68.—Hemslay, Bot. Am.-Cent. ii, 297.

Achras cuneifolia, Poir. in Lamarck, Dict. vi, 534.

B. angustifolia, Nuttall, Sylva, iii, 38, t. 93; 2 ed. ii, 106, t. 93.—Cooper in Smithsonian Rep. 1858, 265.

Sideroxylon cuneatum, A. De Candolle, Prodr. viii, 181.

B. parvifolia, A. De Candolle, Prodr. viii, 190.—Chapman, Fl. S. States, 275.—Vasey, Cat. Forest Trees, 19.

B. myrsinifolia, A. De Candolle, Prodr. viii, 192.

B. reclinata, Torrey, Bot. Mex. Boundary Survey, 109 [not Ventenat].

ANTS' WOOD. DOWNWARD PLUM. SAFFRON PLUM.

A small tree, rarely exceeding 4 meters in height, with a trunk sometimes 0.30 meter in diameter.

Semi-tropical Florida, Merritt's island, Indian river, and southward to the southern keys, not rare; west coast, Cedar Keys to cape Romano, rare; rocky shores and in the interior of low, barren keys; Texas, valley of the lower Rio Grande, Ross to Laredo, and southward into northern Mexico; in the West Indies.

Wood heavy, hard, not strong, very close-grained, compact, satiny, susceptible of a beautiful polish; medullary rays numerous, thin; color, light brown or orange, the sap-wood lighter; specific gravity, 0.7959; ash, 1.90.

183.—*Mimusops Sieberi*, A. De Candolle,

Prodr. viii, 204.—Chapman, Fl. S. States, 275.—Vasey, Cat. Forest Trees, 18.—Gray, Syn. Fl. N. America, ii¹, 69.

Achras Zapotilla, var. *parviflora*, Nuttall, Sylva, iii, 28, t. 90; 2 ed. ii, 97, t. 90.

M. dissecta, Grisebach, Fl. British West Indies, 400, in part.

Achras mammosa, Sieber, Pl. Trin. No. 33 [not Linnæus nor Bonpland].

WILD DILLY.

Semi-tropical Florida, on the southern keys, common; in the West Indies.

A small, low, gnarled tree, sometimes 9 meters in height, with a trunk 0.30 to 0.40 meter in diameter; generally hollow and defective.

Wood very heavy, hard, strong, close-grained, inclined to check in drying, susceptible of a beautiful polish; medullary rays numerous, very obscure; color, rich, very dark brown, the sap-wood lighter; specific gravity, 1.0838; ash, 2.61.

EBENACEÆ.

184.—*Diospyros Virginiana*, Linnæus,

Spec. 1 ed. 1057.—Kalm, Travels, English ed. i, 127, 345.—Marshall, Arbustum, 40.—Wangenheim, Amer. 84, t. 28, f. 58.—Walter, Fl. Caroliniana, 253.—Aiton, Hort. Kew. iii, 446; 2 ed. v, 478.—Abbot, Insects Georgia, ii, t. 61, 74.—B. S. Barton, Coll. i, 11, 45; ii, 52.—Michaux, Fl. Bor.-Am. ii, 258.—Gærtner f. Fruct. Suppl. 138, t. 207.—Willdenow, Spec. iv, 1107; Enum. 1061; Berl. Baumz. 127.—Poiret in Lamarck, Dict. v, 528.—Persoon, Syn. ii, 1806.—Desfontaines, Hist. Arb. i, 208.—Titford, Hort. Bot. Am. 106.—Michaux f. Hist. Arb. Am. ii, 195, t. 12; N. American Sylva, 3 ed. ii, 157, t. 93.—Pursh, Fl. Am. Sept. ii, 265.—Nouveau Duhamel, vi, 84.—Barton, Prodr. Fl. Philadelph. 97; Compend. Fl. Philadelph. ii, 198.—Eaton, Manual, 117; 6 ed. 126.—Nuttall, Genera, ii, 240.—Hayne, Dend. Fl. 228.—Elliott, Sk. ii, 712.—Collin, Förslag af några Nord-Americas Träd. 23.—Torrey, Compend. Fl. N. States, 375.—Audubon, Birds, t. 87.—Sprengel, Syst. ii, 202.—Watson, Dend. Brit. ii, 146.—Rafinesque, Med. Bot. i, 153, t. 32.—Beck, Bot. 229.—Don, Miller's Dict. iv, 39.—London, Arboretum, ii, 1195, t. 200, 201.—Eaton & Wright, Bot. 225.—A. De Candolle, Prodr. iv, 228.—Browne, Trees of America, 368.—Griffith, Med. Bot. 435, f. 196.—Dietrich, Syn. v, 437.—Belg. Hort. iv, 118 & t.—Darby, Bot. S. States, 425.—Darlington, Fl. Cestrica, 3 ed. 176.—Cooper in Smithsonian Rep. 1858, 253.—Chapman, Fl. S. States, 273.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 70.—Lesquereux in Owen's 2d Rep. Arkansas, 374.—"Ettingah. Blatt-Skel. Dikot. 89, t. 38, f. 12."—Wood, Cl. Book, 500; Bot. & Fl. 209.—Porcher, Resources S. Forests, 385.—Engelmann in Trans. Am. Phil. Soc. new ser. xii, 200.—Gray, Manual N. States, 5 ed. 308; Hall's Pl. Texas, 15; Syn. Fl. N. America, ii¹, 69.—Koch, Dendrologie, ii, 204.—Hiern in Trans. Cambridge Phil. Soc. xii¹, 224.—Vasey, Cat. Forest Trees, 18.—Broadhead in Coulter's Bot. Gazette, iii, 59.—Ridgway in Proc. U. S. Nat. Mus. 1882, 68.

D. concolor, Mönch, Meth. 471.

D. Guaiacana, Robin, Voyages, iii, 417.

D. pubescens, Pursh, Fl. Am. Sept. i, 265 [not Persoon].—Rafinesque, Fl. Ludoviciana, 139.—Don, Miller's Dict. iv, 38.—London, Arboretum, ii, 1196.

D. Virginiana, var. *pubescens*, Nuttall, Genera, ii, 240.—Elliott, Sk. ii, 713.

D. Virginiana, var. *microcarpa*, Rafinesque, Med. Bot. i, 115.

D. Virginiana, var. *concolor*, Rafinesque, Med. Bot. i, 155.

D. Virginiana, var. *macrocarpa*, Rafinesque, Med. Bot. i, 155.

D. Persimon, Wikström, Jahr. Schwed. 1830, 92.

D. ciliata, Rafinesque, New Fl. & Bot. i, 25 [not A. De Candolle].

D. calycina, Audibert, Cat. Hort. Tonn. (ex. Spach).—London, Gard. Mag. 1841, 394.

D. angustifolia, Audibert, Cat. Hort. Tonn. (ex. Spach).—London, Gard. Mag. 1841, 394.

D. lucida, Hort.—London, Gard. Mag. 1841, 394.

D. intermedia, Hort.—London, Gard. Mag. 1841, 394.

PERSIMMON.

Light-house point, New Haven, Connecticut, Long Island, New York, and southward to bay Biscayne and the Caloosa river, Florida, southern Alabama and Mississippi; southern Ohio to southeastern Iowa, southern Missouri, Arkansas, eastern Kansas, the Indian territory, and the valley of the Colorado river, Texas.

A tree 10 to 20 or, exceptionally, 30 to 35 meters in height (*Ridgway*), with a trunk sometimes 0.60 meter in diameter; very common and often entirely occupying abandoned fields throughout the middle and lower regions of the southern Atlantic and Gulf states, reaching its greatest development in the rich bottom lands of the lower Ohio basin.

Wood heavy, hard, strong, very close-grained, compact, susceptible of a high polish, containing few scattered, open ducts, the rings of annual growth marked by one or more rows of similar ducts; medullary rays numerous, conspicuous; color, dark brown, or often nearly black, the thick sap-wood light brown, often containing numerous darker spots; specific gravity of the sap-wood, 0.7908; ash, 0.96; used in turnery for shoe-lasts, plane-stocks, etc., and preferred for shuttles; the dark heart-wood only developed in very old specimens and rarely seen.

The yellow edible fruit exceedingly austere until after frost, then becoming sweet and luscious, or in the Gulf states ripening in August without austerity; sometimes used domestically, fermented with hops, corn-meal, or wheat bran, as a beverage under the name of "simmon beer".

A decoction of the bitter and astringent unripe fruit and inner bark occasionally used in the treatment of diarrhoea, sore throat, hemorrhage, etc. (*B. R. Smith in Am. Jour. Pharm.* October, 1846, 215.—*J. E. Bryan in same*, May, 1860, 215.—*U. S. Dispensatory*, 14 ed. 380.—*Nat. Dispensatory*, 2 ed. 514).

185.—*Diospyros Texana*, Scheele,

Linnaea, xxii, 145; Roemer, Texas, 441; Appx. 763.—Walpers, Ann. iii, 14.—Torrey, Bot. Mex. Boundary Survey, 109.—Cooper in Smithsonian Rep. 1858, 266.—Young, Bot. Texas, 376.—Hiern in Trans. Cambridge Phil. Soc. xii¹, 238.—Gray, Hall's Pl. Texas, 15; Syn. Fl. N. America, ii¹, 70.—Vasey, Cat. Forest Trees, 18.—Hemsley, Bot. Am.-Cent. ii, 300.

BLACK PERSIMMON. MEXICAN PERSIMMON. CHAPOTE.

Western Texas, Matagorda bay to the valley of the Concho river; southward into northern Mexico.

A small tree, 4 to 10 meters in height, with a trunk sometimes 0.30 meter in diameter, or more often a low shrub; not rare, and reaching its greatest development in Texas along the rich bottoms of the Guadalupe river; borders of prairies, in rich soil; in Mexico more common and of larger size.

Wood heavy, hard, very close-grained, compact, satiny, taking a beautiful polish, containing few minute, scattered, open ducts; medullary rays numerous, thin; color, nearly black, often streaked with yellow, the thick sap-wood clear bright yellow; specific gravity, 0.8460; ash, 3.33; used in turnery for the handles of tools, etc., suitable for wood-engraving, and probably the best substitute among American woods for box-wood.

The small black fruit sweet and insinid.

STYRACACEÆ.

186.—*Symplocos tinctoria*, L'Heritier,

Trans. Linnæan Soc. i, 176.—Willdenow, Spec. iii, 1436.—Aiton, Hort. Kew. iv, 419.—Sprengel, Syst. iii, 339.—Don, Miller's Dict. iv, 2.—A. De Candolle, Prodr. viii, 254.—Cooper in Smithsonian Rep. 1858, 253.—Chapman, Fl. S. States, 272.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 65.—Lesquereux in Owen's 2d Rep. Arkansas, 374.—Wood, Cl. Book, 499; Bot. & Fl. 209.—Gray, Manual N. States, 5 ed. 310; Syn. Fl. N. America, ii¹, 71.—Young, Bot. Texas, 374.—Vasey, Cat. Forest Trees, 18.

Hopea tinctoria, Linnaeus, Mant. 105.—Walter, Fl. Caroliniana, 189.—Michaux, Fl. Bor.-Am. ii, 42.—Persoon, Syn. ii, 72.—Desfontaines, Hist. Arb. i, 217.—Gærtner f. Fruct. Suppl. 146, t. 209, f. 2.—Robin, Voyages, iii, 419.—Michaux f. Hist. Arb. Am. iii, 61, t. 9; N. American Sylva, 3 ed. iii, 45, t. 117.—Pursh, Fl. Am. Sept. ii, 451.—Nuttall, Genera, ii, 83.—Elliott, Sk. ii, 173.—Eaton, Manual, 6 ed. 176.—Spach, Hist. Veg. ix, 420.—Eaton & Wright, Bot. 272.—Darby, Bot. S. States, 425.—Porcher, Resources S. Forests, 388.

HORSE SUGAR. SWEET LEAF.

Southern Delaware, south to about latitude 30° in Florida, and west through the Gulf states to western Louisiana and southern Arkansas (Malvern, Texarkana, *Letterman*).

A small tree, 6 to 10 meters in height, with a trunk 0.20 to 0.25 meter in diameter, or often a low shrub; borders of cypress swamps or in deep, damp, shaded woods.

Wood light, soft, not strong, close-grained, checking in drying; medullary rays numerous, thin; color, light red, or often nearly white, the sap-wood lighter; specific gravity, 0.5325; ash, 0.68.

Leaves sweet, greedily eaten by cattle and horses, and yielding, as does also the bark, a yellow dye.

187.—*Halesia diptera*, Linnaeus,

Spec. 2 ed. 636.—Marshall, Arbustum, 57.—Lamarck, Dict. ii, 66.—Willdenow, Spec. ii, 849; Enum. 496; Berl. Baumz. 171.—Cavanilles, Diss. vi, 338, t. 187.—Michaux, Fl. Bor.-Am. ii, 40.—Persoon, Syn. ii, 4.—Aiton, Hort. Kew. 2 ed. iii, 143.—Nouveau Duhamel, v, 144.—Pursh, Fl. Am. Sept. ii, 450.—Nuttall, Genera, ii, 83.—Elliott, Sk. i, 508.—Hayne, Dend. Fl. 66.—Loddiges, Bot. Cab. t. 1172.—Sprengel, Syst. iii, 84.—Eaton, Manual, 6 ed. 164.—Don, Miller's Dict. iv, 7.—Loudon, Arboretum, ii, 1191, f. 1014.—Spach, Hist. Veg. ix, 426.—Eaton & Wright, Bot. 260.—A. De Candolle, Prodr. viii, 270.—Miers, Contrib. i, 193.—Darby, Bot. S. States, 425.—Cooper in Smithsonian Rep. 1858, 253.—Chapman, Fl. S. States, 271.—Wood, Cl. Book, 499; Bot. & Fl. 209.—Koch, Dendrolgie, ii, 201.—Vasey, Cat. Forest Trees, 18.—Gray, Syn. Fl. N. America, ii¹, 71.

H. reticulata, Buckley in Proc. Philadelphia Acad. 1860, 444.

SNOW-DROP TREE. SILVER-BELL TREE.

South Carolina to northern Florida, near the coast, and west through the lower region of the Gulf states to eastern Texas and Garland county, Arkansas (*Harvey*).

A small tree, sometimes 6 to 10 meters in height, with a trunk 0.10 to 0.20 meter in diameter, or often a shrub sending up many clustered stems from the root; borders of swamps, in low, wet woods.

Wood light, soft, strong, very close-grained, compact; medullary rays numerous, thin; color, light brown, the sap-wood lighter; specific gravity, 0.5705; ash, 0.42.

188.—*Halesia tetraptera*, Linnaeus,

Spec. 2 ed. 636.—Marshall, Arbustum, 57.—Gärtner, Fruct. i, 160, t. 32, f. 2.—Lamarck, Dict. ii, 66; Ill. ii, 521, t. 404, f. 1.—Aiton, Hort. Kew. ii, 125; 2 ed. iii, 143.—Mönch, Meth. 507.—Abbot, Insects Georgia i, t. 46.—Willdenow, Spec. ii, 849; Enum. 496; Berl. Baumz. 170.—Cavanilles, Diss. vi, 338, t. 186.—Michaux, Fl. Bor.-Am. ii, 40.—Persoon, Syn. ii, 4.—Desfontaines, Hist. Arb. i, 216.—Nouveau Duhamel, v, 143, t. 45.—Pursh, Fl. Am. Sept. ii, 449.—Nuttall, Genera, ii, 82.—Bot. Mag. t. 910.—Elliott, Sk. i, 507.—Hayne, Dend. Fl. 66.—Loddiges, Bot. Cab. t. 1173.—Sprengel, Syst. iii, 84.—Guimpel, Otto & Hayne, Abb. Holz. 43, t. 35.—Eaton, Manual, 6 ed. 164.—Don, Miller's Dict. iv, 6.—London, Arboretum, ii, 1190, f. 1012, t. 196, 197.—Spach, Hist. Veg. ix, 426.—Eaton & Wright, Bot. 260.—A. De Candolle, Prodr. viii, 270.—Browne, Trees of America, 366.—Miers, Contrib. i, 191, t. 93.—Darby, Bot. S. States, 425.—Cooper in Smithsonian Rep. 1858, 253.—Agardh, Theor. & Syst. Pl. t. 22, f. 16, 17.—Chapman, Fl. S. States, 271.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 80.—Wood, Cl. Book, 499; Bot. & Fl. 209.—Ørsted in Saerakitt. Afttryk. af Nat. For. Viden. Meddelt. Nos. 1-6, 1866, 89, f. 2.—Gray, Manual N. States, 5 ed. 310; Syn. Fl. N. America, ii, 71.—Koch, Dendrologie, ii, 199.—Young, Bot. Texas, 374.—Vasey, Cat. Forest Trees, 18.

RATTLEBOX. SNOW-DROP TREE. SILVER-BELL TREE. CALICO WOOD.

Mountains of West Virginia to southern Illinois, south to middle Florida, central Alabama and Mississippi, and through Arkansas to western Louisiana and eastern Texas.

A tree 10 to 15 meters in height, with a trunk rarely 0.60 meter in diameter, or often a tall shrub; generally along streams, in rich soil; most common and reaching its greatest development in the southern Alleghany mountains; common in cultivation.

Wood light, soft, close-grained, compact; medullary rays numerous, thin; color, light brown, the sap-wood lighter; specific gravity, 0.5628; ash, 0.40.

NOTE.—*Halesia parviflora*, Michaux, of southern Georgia, and Florida, does not attain the size or habit of a tree.

OLEACEÆ.

189.—*Fraxinus Greggii*, Gray,

Proc. Am. Acad. vii, 64; Syn. Fl. N. America, ii, 74.—Hemsley, Bot. Am.-Cent. ii, 305.

F. Schiedeana, var. *parvifolia*, Torrey, Bot. Mex. Boundary Survey, 166.

Western Texas, valley of the Rio Grande, from the San Pedro to the Pecos river; southward into Mexico.

A small tree, sometimes 7 to 9 meters in height, with a trunk 0.10 to 0.15 meter in diameter (Lampasas mountains, Mexico, *Buckley*), or often a graceful shrub; limestone soil.

Wood heavy, hard, very close-grained, compact; layers of annual growth and medullary rays obscure; color, brown, the sap-wood lighter; specific gravity, 0.7904; ash, 0.93.

190.—*Fraxinus anomala*, Torrey;

Watson in King's Rep. v, 283.—Parry in Am. Nat. ix, 203.—Vasey, Cat. Forest Trees, 20.—Gray, Syn. Fl. N. America, ii, 74.

Southwestern Colorado, McElmo river (*Brandegge*), southern Utah, Kanawa, Leeds, Silver Leaf, Labyrinth cañon of the Colorado river, valley of the Rio Virgen, near Saint George.

A small tree, sometimes 6 meters in height, with a trunk 0.15 to 0.20 meter in diameter, with the habit of a dwarf pear tree; common on elevated sandstone *mesas* and plateaus.

Wood heavy, hard, coarse-grained, containing many large, open, scattered ducts, the layers of annual growth marked by several rows of similar ducts; medullary rays numerous, thin; color, light brown, the sap-wood lighter; specific gravity, 0.6597; ash, 0.85.

191.—*Fraxinus pistaciæfolia*, Torrey,

Pacific R. R. Rep. iv, 128; Bot. Mex. Boundary Survey, 166.—Cooper in Smithsonian Rep. 1858, 260.—Gray, Hall's Pl. Texas, 19; Syn. Fl. N. America, ii, 74.—Vasey, Cat. Forest Trees, 20.—Rusby in Bull. Torrey Bot. Club. ix, 54.—Hemsley, Bot. Am.-Cent. ii, 305.—Watson in Proc. Am. Acad. xviii, 113.

F. velutina, Torrey in Emory's Rep. 149.

F. coriacea, Watson in Am. Nat. vii, 302, in part.—Rothrock in Wheeler's Rep. vi, 186, t. 22.—Vasey, Cat. Forest Trees, 20.

F. pistaciæfolia, var. *coriacea*, Gray, Syn. Fl. N. America, ii, 74.

ASH.

Mountains of western Texas, through southern New Mexico, southern and eastern Arizona, to southern Nevada (Ash Meadows, *Rothrock*); in northern Mexico.

A small tree, 10 to 12 meters in height, with a trunk rarely 0.45 meter in diameter; generally along borders of streams, in elevated cañons, less commonly in dry soil, the foliage then thick and coriaceous or, more rarely, velvety tomentose (var. *coriacea*, Gray, l. c.); the large specimens generally hollow and defective.

Wood heavy, soft, not strong, coarse-grained, compact; medullary rays numerous, thin; color, light brown, the sap-wood lighter; specific gravity, 0.6810; ash, 0.62; occasionally used in wagon-building, for ax handles, etc.

192.—*Fraxinus Americana*, Linnæus,

Spec. 2 ed. 1510.—Walter, Fl. Caroliniana, 254.—Aiton, Hort. Kew. iii, 445; 2 ed. v, 476.—Willdenow, Spec. iv, 1102; Enum. 1060; Berl. Baumz. 145.—Muhlenberg & Willdenow in Neue Schriften Gesell. Nat. Fr. Berlin, iii, 393.—Vahl Enum. i, 49.—Persoon, Syn. ii, 604.—Desfontaines, Hist. Arb. i, 102.—Nouveau Duhamel, iv, 63.—Michaux f. Hist. Arb. Am. iii, 106, t. 8; N. American Sylva, 3 ed. iii, 49, t. 118 (excl. fruit).—Barton, Prodr. Fl. Philadelph. 97; Compend. Fl. Philadelph. ii, 192.—Eaton, Manual, 114.—Hayne, Dend. Fl. 221.—Cobbett, Woodlands, 131.—Sprengel, Syst. i, 95.—Beck, Bot. 232.—London, Arboretum, ii, 1232, f. 1055 & t.—Penn. Cycl. x, 455.—Bigelow, Fl. Boston. 3 ed. 408.—Hooker, Fl. Bor.-Am. ii, 51.—Torrey, Fl. N. York, ii, 125, t. 89.—A. De Candolle, Prodr. viii, 177.—Browne, Trees of America, 394.—Darlington, Fl. Cestrica, 3 ed. 238.—Cooper in Smithsonian Rep. 1858, 253.—Chapman, Fl. S. States, 369.—Curtis in Geological Rep. N. Carolina, 1860, iii, 54.—Wood, Cl. Book, 597; Bot. & Fl. 277.—Lesquereux in Owen's 2d Rep. Arkansas, 382.—Engelmann in Trans. Am. Phil. Soc. new ser. xii, 206.—Porcher, Resources S. Forests, 494.—Gray, Manual N. States, 5 ed. 401; Hall's Pl. Texas, 19; Syn. Fl. N. America, ii, 74.—Koch, Dendrologie, ii, 252.—Young, Bot. Texas, 452.—Vasey, Cat. Forest Trees, 20.—Macoun in Geological Rep. Canada, 1875-'76, 207.—Sears in Bull. Essex Inst. xiii, 177.—Bell in Geological Rep. Canada, 1879-'80, 52.—Ridgway in Proc. U. S. Nat. Mus. 1882, 68.

F. Caroliniensis, Wangenheim, Amer. 81.

F. alba, Marshall, Arbustum, 51.—Hayne, Dend. Fl. 223.

F. acuminata, Lamarck, Dict. ii, 542.—Bosc in Mem. Inst. 1808, 205.—Pursh, Fl. Am. Sept. ii, 9.—Nuttall, Genera, ii, 231; Sylva, iii, 64; 2 ed. ii, 129.—Hayne, Dend. Fl. 220.—Elliott, Sk. ii, 672.—Sprengel, Syst. i, 95.—Torrey, Compend. Fl. N. States, 371; Nicolle's Rep. 154.—Römer & Schultes, Syst. iii, 277.—Darlington, Fl. Cestrica, 2 ed. 8.—Eaton, Manual, 6 ed. 148.—Beck, Bot. 232.—Don, Miller's Dict. iv, 56.—Eaton & Wright, Bot. 247.—Emerson, Trees Massachusetts, 333; 2 ed. ii, 376 & t.—Darby, Bot. S. States, 429.—Porcher, Resources S. Forests, 494.

? *F. juglandifolia*, Lamarck, Dict. ii, 542.—Bosc in Mem. Inst. 1808, 208.—Desfontaines, Hist. Arb. i, 103.—Hayne, Dend. Fl. 221.—Beck, Bot. 232.—Don, Miller's Dict. iv, 55.

F. Canadensis, Gärtner, Fruct. i, 222, t. 49.

F. epiptera, Michaux, Fl. Bor.-Am. ii, 256.—Vahl, Enum. i, 50.—Willdenow, Spec. iv, 1102; Berl. Baumz. 147.—Persoon, Syn. ii, 603.—Desfontaines, Hist. Arb. i, 103.—Poiret, Suppl. ii, 671.—Nuttall, Genera, ii, 231.—Pursh, Fl. Am. Sept. i, 8.—Elliott, Sk. ii, 672.—Sprengel, Syst. i, 96.—Römer & Schultes, Syst. 278.—Eaton, Manual, 6 ed. 148.—Don, Miller's Dict. iv, 55.—London, Arboretum, ii, 1237.—Penn. Cycl. x, 455.—Eaton & Wright, Bot. 247.—Hooker, Fl. Bor.-Am. ii, 50.—A. De Candolle, Prodr. viii, 277.—Darby, Bot. S. States, 429.—Cooper in Smithsonian Rep. 1858, 399.

F. lancea, Bosc in Mem. Inst. 1808, 209 (*vide* London, Arboretum, ii, 1237).

F. discolor, Muhlenberg, Cat. 111.—Rafinesque, Fl. Ludoviciana, 37.—Spach, Hist. Veg. viii, 297.

F. Americana, var. *latifolia*, London, Arboretum, ii, 1232.—Browne, Trees of America, 396.

? *F. juglandifolia*, var. *serrata*, Hayne, Dend. Fl. 221.

? *F. juglandifolia*, var. *subserrata*, Hayne, Dend. Fl. 221.

WHITE ASH.

Nova Scotia, New Brunswick, southern Ontario to northern Minnesota, south to northern Florida, central Alabama and Mississippi, and west to eastern Nebraska, eastern Kansas, the Indian territory, and the valley of the Trinity river, Texas.

A large tree of the first economic value, 15 to 30 or, exceptionally, 42 meters (*Ridgway*) in height, with a trunk 1.20 to 1.80 meter in diameter; low, rich, rather moist soil, reaching its greatest development in the bottom lands of the lower Ohio River basin; toward its western and southwestern limits smaller, of less economic value, and generally replaced by the green ash (*Fraxinus viridis*).

A form of the southern states with remarkably small fruit has been described as—

var. *microcarpa*, Gray, Syn. Fl. N. America, ii¹, 75.

F. albicans, Buckley in Proc. Philadelphia Acad. 1862, 4, in part.

F. Curtissii, Vasey, Cat. Forest Trees, 20.

Wood heavy, hard, strong, ultimately brittle, coarse-grained, compact; layers of annual growth clearly marked by several rows of large open ducts, occupying in slowly-grown specimens nearly the entire width of the annual rings; medullary rays numerous, obscure; color, brown, the sap-wood much lighter, often nearly white; specific gravity, 0.6543; ash, 0.42; specific gravity of the heavier sap-wood, 0.7180; largely used in the manufacture of agricultural implements, carriages, handles, oars, and for interior and cabinet work.

Var. *Texensis*,

Gray, Syn. Fl. N. America, ii¹, 75.

F. albicans, Buckley in Proc. Philadelphia Acad. 1862, 4, in part.

F. coriacea, Watson in Am. Nat. vii, 302, in part.

F. pistaciifolia, Gray, Hall's Pl. Texas, 19 [not Torrey].

Western Texas, Dallas (*Reverchon*), to the valley of the Devil's river.

A small tree, 10 to 12 meters in height, with a trunk sometimes 0.60 meter in diameter; dry, rocky hills and ridges.

Wood heavy, hard, strong, rather close-grained, compact; layers of annual growth marked by one or more rows of open ducts; medullary rays numerous, obscure; color, light brown, the sap-wood lighter; specific gravity, 0.7636; ash, 0.70; used for the same purposes as that of the species.

193.—*Fraxinus pubescens*, Lamarck,

Dict. ii, 548.—Walter, Fl. Caroliniana, 254.—Willdenow, Spec. iv, 1103; Enum. 1060; Berl. Baumz. 148.—Muhlenberg & Willdenow in Neue Schriften Gesell. Nat. Fr. Berlin, iii, 393.—Vahl, Enum. i, 51.—Persoon, Syn. ii, 604.—Desfontaines, Hist. Arb. i, 102.—Nouveau Duhamel, iv, 62.—Aiton, Hort. Kew. 2 ed. v, 476.—Pursh, Fl. Am. Sept. i, 9.—Römer & Schultes, Syst. 279.—Nuttall, Genera, ii, 231.—Hayne, Dend. Fl. 223.—Elliott, Sk. ii, 673.—Sprengel, Syst. i, 95.—Torrey, Compend. Fl. N. States, 371; Fl. N. York, ii, 126.—Beck, Bot. 232.—Eaton, Manual, 6 ed. 148.—Don, Miller's Dict. iv, 55.—London, Arboretum, ii, 1233, f. 1056.—Penn. Cycl. x, 455.—Eaton & Wright, Bot. 247.—Hooker, Fl. Bor.-Am. ii, 51.—A. De Candolle, Prodr. viii, 278.—Emerson, Trees Massachusetts, 337; 2 ed. ii, 380.—Darlington, Fl. Cestrica, 3 ed. 239.—Darby, Bot. S. States, 429.—Cooper in Smithsonian Rep. 1858, 253.—Chapman, Fl. S. States, 370.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 54.—Wood, Cl. Book, 597; Bot. & Fl. 277.—Gray, Manual N. States, 5 ed. 402; Syn. Fl. N. America, ii¹, 75.—Young, Bot. Texas, 452.—Vasey, Cat. Forest Trees, 20.—Sears in Bull. Essex Inst. xiii, 177.—Ridgway in Proc. U. S. Nat. Mus. 1882, 69.

F. Pennsylvanica, Marshall, Arbustum, 51.—Koch, Dendrologie, ii, 253.

F. nigra, Du Roi, Harbk. 2 ed. i, 398 [not Marshall].

F. pubescens, var. *longifolia*, Willdenow, Spec. iv, 1104.—Vahl, Enum. i, 52.—Pursh, Fl. Am. Sept. ii, 9.—Loddiges, Cat. ed. 1836.—London, Arboretum, ii, 1233.—A. De Candolle, Prodr. viii, 278.

F. pubescens, var. *latifolia*, Willdenow, Spec. iv, 1104.—Vahl, Enum. i, 52.—Pursh, Fl. Am. Sept. i, 9.—Hayne, Dend. Fl. 223.—Eaton, Manual, 6 ed. 148.—London, Arboretum, ii, 1233.—A. De Candolle, Prodr. viii, 278.

F. pubescens, var. *subpubescens*, Persoon, Syn. ii, 605.—Pursh, Fl. Am. Sept. i, 9.—Eaton, Manual, 6 ed. 148.—London, Arboretum, ii, 1234.—A. De Candolle, Prodr. viii, 278.—Browne, Trees of America, 395.

F. longifolia, Bosc in Mem. Inst. 1808, 209.

F. subvillosa, Bosc in Mem. Inst. 1808, 209.

F. tomentosa, Michaux f. Hist. Arb. Am. iii, 112, t. 9; N. American Sylva, 3 ed. iii, 53, t. 119.—Barton, Compend. Fl. Philadelph. ii, 192.

F. Americana, var. *pubescens*, Browne, Trees of America, 395.

F. oblongocarpa, Buckley in Proc. Philadelphia Acad. 1864, 4.

RED ASH.

New Brunswick to southern Ontario and northern Minnesota, south to northern Florida and central Alabama.

A tree 12 to 15 meters in height, with a trunk rarely exceeding 0.60 meter in diameter; borders of streams and swamps, in low ground; common and reaching its greatest development in the north Atlantic states; rare west of the Alleghany mountains, probably not extending west of the Mississippi river.

Wood heavy, hard, strong, brittle, coarse-grained, compact; medullary rays numerous, thin; color, rich brown, the sap-wood light brown streaked with yellow; specific gravity, 0.6251; ash, 0.26; specific gravity of the lighter sap-wood, 0.5609; somewhat used as a substitute for the more valuable white ash, with which it is often confounded.

194.—*Fraxinus viridis*, Michaux f.

Hist. Arb. Am. iii, 115, t. 10; N. American Sylva, 3 ed. iii, 54, t. 120 (excl. fruit).—Hayne, Dend. Fl. 222.—Cooper in Smithsonian Rep. 1858, 253.—Chapman, Fl. S. States, 370.—Gray in Pacific R. R. Rep. xii³, 46; Manual N. States, 5 ed. 402; Hall's Fl. Texas, 19; Syn. Fl. N. America, ii¹, 75.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 54.—Lesquereux in Owen's 2d Rep. Arkansas, 382.—Wood, Cl. Book, 598; Bot. & Fl. 277.—Watson in King's Rep. v, 284.—Young, Bot. Texas, 453.—Vasey, Cat. Forest Trees, 20.—Macoun in Geological Rep. Canada, 1875-'76, 207.—Bell in Geological Rep. Canada, 1879-'80, 49.—Hemslay, Bot. Am.-Cent. ii, 305.—Burgess in Coulter's Bot. Gazette, vii, 95.

F. juglandifolia, Willdenow, Spec. iv, 1104; Enum. 1060; Berl. Baumz. 140 [not Lamarek].—Vahl, Enum. i, 50.—Persoon, Syn. ii, 604.—Nouveau Duhamel, iv, 63, t. 16.—Aiton, Hort. Kew. 2 ed. v, 476.—Pursh, Fl. Am. Sept. i, 9.—Römer & Schultes, Syst. i, 278; iii, Suppl. 255.—Eaton, Manual, 114.—Sprengel, Syst. i, 95.—Torrey, Compend. Fl. N. States, 371.—Beck, Bot. 233.—Don, Miller's Dict. iv, 55.—London, Arboretum, ii, 1236, f. 1061, 1062 & t.—Eaton & Wright, Bot. 247.—Gray, Manual N. States, 1 ed. 373.

F. Caroliniana, Willdenow, Spec. iv, 1103; Enum. 1060; Berl. Baumz. 148.—Vahl, Enum. i, 51.—Du Roi, Harbk. 2 ed. i, 400.—Persoon, Syn. ii, 605.—Desfontaines, Hist. Arb. i, 103.—Nouveau Duhamel, iv, 62.—Pursh, Fl. Am. Sept. i, 9.—Nuttall, Genera, ii, 231.—Elliott, Sk. ii, 673.—Hayne, Dend. Fl. 223.—Sprengel, Syst. i, 95.—Eaton, Manual, 6 ed. 148.—Don, Miller's Dict. iv, 55.—Eaton & Wright, Bot. 147.—Darby, Bot. S. States, 429.

F. juglandifolia, var. *subintegerrima*, Vahl, Enum. i, 50.

F. expansa, Willdenow, Berl. Baumz. 150.—Römer & Schultes, Syst. i, 279.—Don, Miller's Dict. iv, 55.—London, Arboretum, ii, 1238.—A. De Candolle, Prodr. viii, 278.—Browne, Trees of America, 399.

F. Americana, var. *juglandifolia*, Browne, Trees of America, 398.

F. Novæ-Angliæ, Koch, Dendrologie, ii, 251 [not Miller nor Wangenheim]

GREEN ASH.

Shores of lake Champlain, Tiverton, Rhode Island, and southward to northern Florida, west to the valley of the Saskatchewan, the eastern ranges of the Rocky mountains of Montana, the Wahsatch mountains of Utah, and the ranges of eastern and northern Arizona.

A tree 15 to 18 meters in height, with a trunk rarely exceeding 0.60 meter in diameter; borders of streams or in low, rather moist soil; at the west confined to the bottom lands of the large streams and to high mountain cañons.

Wood heavy, hard, strong, brittle, rather coarse-grained, compact, satiny, containing numerous scattered, small, open ducts, the layers of annual growth marked by several rows of larger ducts; medullary rays numerous, obscure; color, brown, the sap-wood lighter; specific gravity, 0.7117; ash, 0.65; inferior in quality, although often used as a substitute for white ash.

Var. *Berlandieriana*, Torrey,

Bot. Mex. Boundary Survey, 166.—Gray, Syn. Fl. N. America, iii¹, 75.—Hemslay, Bot. Am.-Cent. ii, 305.—Watson in Proc. Am. Acad. xviii, 113.

F. Berlandieriana, De Candolle, Prodr. viii, 278.

F. trialata, Buckley in Proc. Philadelphia Acad. 1862, 5.

Texas, west of the Colorado river; southward into northern Mexico.

A small tree, 9 to 12 meters in height, with a trunk rarely exceeding 0.30 meter in diameter; borders of streams, in low, rich soil.

Wood light, soft, rather close-grained, compact, containing few small, scattered, open ducts, the layers of annual growth clearly marked by one or two rows of larger ducts; medullary rays numerous, obscure; color, light brown, the sap-wood lighter; specific gravity, 0.5780; ash, 0.54.

195.—*Fraxinus platycarpa*, Michaux,

Fl. Bor.-Am. ii, 256.—Vahl, Enum. i, 49.—Persoon, Syn. ii, 605.—Desfontaines, Hist. Arb. i, 103.—Nouveau Duhamel, iv, 64.—Michaux f. Hist. Arb. Am. iii, 128, t. 13; N. American Sylva, 3 ed. iii, 63, t. 124.—Poiret, Suppl. ii, 671.—Pursh, Fl. Am. Sept. i, 9.—Römer & Schultes, Syst. i, 278.—Nuttall, Genera, ii, 231.—Hayne, Dend. Fl. 225.—Elliott, Sk. ii, 673.—Sprengel, Syst. i, 96.—Eaton, Manual, 6 ed. 149.—Don, Miller's Dict. iv, 55.—Eaton & Wright, Bot. 247.—A. De Candolle, Prodr. viii, 277.—Darby, Bot. S. States, 429.—Cooper in Smithsonian Rep. 1858, 253.—Chapman, Fl. S. States, 370.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 53.—Lesquereux in Owen's 2d Rep. Arkansas, 382.—Wood, Cl. Book, 598; Bot. & Fl. 277.—Gray, Manual N. States, 5 ed. 402; Syn. Fl. N. America, ii, 75.—Young, Bot. Texas, 453.—Vasey, Cat. Forest Trees, 20.

?*F. Caroliniana*, Miller, Dict. No. 6.—Lamarck, Dict. ii, 518.—Römer & Schultes, Syst. i, 278.—Don, Miller's Dict. iv, 55.—London, Arboretum, ii, 1237.—Koch, Dendrologie, ii, 258.

F. excelsior, Walter, Fl. Caroliniana, 254 [not Linnaeus].

F. Americana, Marshall, Arbustum, 50 [not Linnaeus].

F. pallida, Bosc in Mem. Inst. 1808, 209.

F. pubescens, Bosc in Mem. Inst. 1808, 210 [not Lamarck].

F. triptera, Nuttall, Genera, ii, 232; Sylva, iii, 62, t. 100; 2 ed. 127, t. 100.—Elliott, Sk. ii, 674.—Don, Miller's Dict. iv, 56.—London, Arboretum, ii, 1240.—A. De Candolle, Prodr. viii, 274.—Darby, Bot. S. States, 429.

F. curvidens, Hoffmannsegg, Verz. d. Pflanzenkult. 29.

F. pauciflora, Nuttall, Sylva, iii, 61, t. 100; 2 ed. ii, 126, t. 100.

F. Americana, var. *Caroliniana*, Browne, Trees of America, 398.

F. Americana, var. *triptera*, Browne, Trees of America, 399.

F. Nuttallii, Buckley in Proc. Philadelphia Acad. 1860, 444.

F. nigrescens, Buckley in Proc. Philadelphia Acad. 1862, 5.

WATER ASH.

Southeastern Virginia, south near the coast to cape Canaveral and the Caloosa river, Florida, west through the Gulf states to the valley of the Sabine river, Texas, and the Washita river, southwestern Arkansas; in the West Indies.

A small tree, 9 to 12 meters in height, with a trunk rarely exceeding 0.30 meter in diameter; deep river swamps.

Wood very light, soft, not strong, brittle, close-grained, compact, the open ducts not conspicuous; medullary rays few, obscure; color, nearly white, or sometimes tinged with yellow, the sap-wood lighter; specific gravity, 0.3541; ash, 0.73.

196.—*Fraxinus quadrangulata*, Michaux,

Fl. Bor.-Am. ii, 255.—Willdenow, Spec. iv, 1104.—Vahl, Enum. i, 50.—Persoon, Syn. ii, 605.—Bosc in Mem. Inst. 1808, 211.—Desfontaines, Hist. Arb. i, 103.—Nouveau Duhamel, iv, 64.—Michaux f. Hist. Arb. Am. iii, 118, t. 11; 2 ed. iii, 61, t. 123.—Poiret, Suppl. ii, 671.—Pursh, Fl. Am. Sept. i, 8.—Römer & Schultes, Syst. i, 278.—Nuttall, Genera, ii, 231.—Hayne, Dend. Fl. 223.—Sprengel, Syst. i, 96.—Eaton, Manual, 6 ed. 149.—Don, Miller's Dict. iv, 55.—London, Arboretum, ii, 1235, f. 1059, 1060.—Spach, Hist. Veg. viii, 286.—Penn. Cycl. x, 455.—Eaton & Wright, Bot. 247.—A. De Candolle, Prodr. viii, 278.—Cooper in Smithsonian Rep. 1858, 254.—Chapman, Fl. S. States, 370.—Lesquereux in Owen's 2d Rep. Arkansas, 382.—Wood, Cl. Book, 598; Bot. & Fl. 277.—Gray, Manual N. States, 5 ed. 402; Syn. Fl. N. America, ii, 75.—Koch, Dendrologie, ii, 259.—Young, Bot. Texas, 453.—Vasey, Cat. Forest Trees, 20.—Engelmann in Coulter's Bot. Gazette, v, 63.—Ridgway in Proc. U. S. Nat. Mus. 1882, 69.—Burgess in Coulter's Bot. Gazette, vii, 95.

F. tetragona, Cels in Nouv. Cours, Agr. vii, 73.

F. quadrangularis, Loddiges, Cat. 1836.

F. nervosa, Loddiges, Cat. 1836.

F. quadrangulata, var. *nervosa*, London, Arboretum, ii, 1235.

F. Americana, var. *quadrangulata*, Browne, Trees of America, 397.

F. Americana, var. *quadrangulata nervosa*, Browne, Trees of America, 397.

BLUE ASH.

Southern Michigan to central Minnesota, south to northern Alabama, and through Iowa and Missouri to northeastern Arkansas (Duvall's bluff, *Letterman*).

A tree 18 to 25 or, exceptionally, 37 meters in height, with a trunk rarely exceeding 0.60 meter in diameter; generally on limestone hills, rarely extending into the bottom lands, and reaching its greatest development in the basin of the lower Wabash river.

Wood heavy, hard, not strong, brittle, close-grained, compact, satiny; layers of annual growth clearly marked by one to three rows of large open ducts; medullary rays numerous, obscure; color, light yellow streaked with brown, the sap-wood lighter; specific gravity, 0.7184; ash, 0.78; largely used for flooring, in carriage-building, etc. The inner bark, macerated, dyes blue.

197.—*Fraxinus Oregana*, Nuttall,

Sylva, iii, 59, t. 99; 2 ed. ii, 124, t. 99.—Torrey in *Pacific R. R. Rep.* iv, 128.—Newberry in *Pacific R. R. Rep.* vi, 25, 87.—Cooper in *Smithsonian Rep.* 1858, 260; *Pacific R. R. Rep.* xii, 28, 68; *Am. Nat.* iii, 407.—Koch, *Dendrologie*, ii, 260.—Gray in *Bot. California*, i, 472; *Syn. Fl. N. America*, ii, 76.—Vasey, *Cat. Forest Trees*, 20.

F. pubescens, var. Hooker, *Fl. Bor.-Am.* ii, 51.

F. grandifolia, Benthams, *Bot. Sulphur*, 33.

OREGON ASH.

Shores of Puget sound, south through Washington territory and Oregon west of the eastern valleys of the Cascade mountains, along the California Coast ranges to San Francisco bay and the western slopes of the Sierra Nevada to the San Bernardino and Hot Spring mountains, California.

A tree sometimes 24 meters in height, with a trunk rarely exceeding 0.60 meter in diameter; moist soil, generally along streams, and reaching its greatest development in the bottom lands of southwestern Oregon.

Wood light, hard, not strong, brittle, coarse-grained, compact, containing many large, open, scattered ducts, the layers of annual growth strongly marked with several rows of similar ducts; medullary rays numerous, thin; color, brown, the sap-wood lighter; specific gravity, 0.5731; ash, 0.34; specific gravity of the lighter sap-wood, 0.5630; used in the manufacture of furniture, for the frames of carriages and wagons, in cooperage, for fuel, etc.

198.—*Fraxinus sambucifolia*, Lamareck,

Dict. ii, 549.—Muhlenberg & Willdenow in *Neue Schriften Gesell. Nat. Fr. Berlin*, iii, 393.—Willdenow, *Spec.* iv, 1099; *Enum.* 1059; *Berl. Baumz.* 150.—Vahl, *Enum.* i, 51.—Persoon, *Syn.* ii, 605.—Desfontaines, *Hist. Arb.* i, 103.—Bosc in *Mem. Inst.* 1808, 211.—Nouveau Duhamel, iv, 60.—Aiton, *Hort. Kew.* v, 475.—Michaux f. *Hist. Arb. Am.* iii, 122, t. 12; *N. American Sylva*, 3 ed. iii, 159, t. 122.—Pursh, *Fl. Am. Sept.* i, 8.—Römer & Schultes, *Syst.* i, 279.—Nuttall, *Genera*, ii, 231.—Barton, *Compend. Fl. Philadelph.* ii, 192.—Hayne, *Dend. Fl.* 224.—Torrey, *Compend. Fl. N. States*, 371; *Fl. N. York*, ii, 126.—Beck, *Bot.* 232.—Eaton, *Manual*, 6 ed. 148.—Don, *Miller's Dict.* iv, 54.—London, *Arboretum*, ii, 1234, f. 1057, 1058.—Spach, *Hist. Veg.* viii, 299.—Hooker, *Fl. Bor.-Am.* ii, 50.—Eaton & Wright, *Bot.* 147.—A. De Candolle, *Prodr.* viii, 278.—Emerson, *Trees Massachusetts*, 338; 2 ed. ii, 381 & t.—Darlington, *Fl. Cestrica*, 3 ed. 239.—Cooper in *Smithsonian Rep.* 1858, 253.—Lesquereux in *Owen's 2d Rep. Arkansas*, 382.—Wood, *Cl. Book*, 598; *Bot. & Fl.* 277.—Gray, *Manual N. States*, 5 ed. 402; *Syn. Fl. N. America*, ii, 76.—Vasey, *Cat. Forest Trees*, 20.—Ridgway in *Proc. U. S. Nat. Mus.* 1892, 69.—Bell in *Geological Rep. Canada*, 1879-'80, 48.

F. nigra, Marshall, *Arbustum*, 51.

F. Nova-Angliae, Wangenheim, *Amer.* 51.

F. crispa, Hort.

F. sambucifolia, var. *crispa*, Loddiges, *Cat.* 1836.—London, *Arboretum*, ii, 1234.

F. Americana, var. *sambucifolia*, Browne, *Trees of America*, 393.

BLACK ASH. HOOP ASH. GROUND ASH.

Southern Newfoundland, along the northern shores of the gulf of Saint Lawrence, southwesterly to the eastern shores of lake Winnipeg, south through the northern states to New Castle county, Delaware, the mountains of Virginia, southern Illinois, and northwestern Arkansas.

A tree 25 to 30 meters in height, with a trunk 0.30 to 0.60 meter in diameter; swamps and low river banks; the most northern representative of the genus in America.

Wood heavy, soft, not strong, tough, rather coarse-grained, compact, durable, separating easily into thin layers; layers of annual growth strongly marked by several rows of large open ducts; medullary rays numerous,

thin; color, dark brown, the sap-wood light brown, or often nearly white; specific gravity, 0.6318; ash, 0.72; specific gravity of the heavier sap-wood, 0.7465; largely used for interior finish, fencing, barrel hoops, in cabinet-making, and the manufacture of baskets.

NOTE.—*Frazinus dipetala*, Hooker & Arnott, of the California Coast ranges and the western slopes of the southern Sierra Nevada, and *F. cuspidata*, Torrey, of the valley of the Rio Grande, do not attain arborescent habit or dimensions.

The following, characterized by Bosc in Mem. Inst. 1808, mainly from the foliage of garden specimens of supposed North American origin, cannot be safely referred to our species: *F. alba*, *cinerea*, *elliptica*, *fusca*, *mixta*, *nigra*, *ovata*, *pannosa*, *pulverulenta*, *Richardi*, *rubicunda*, and *rufa*.

199.—*Forestiera acuminata*, Poiret,

Suppl. ii, 664.—Hayne, Dend. Fl. 194.—Nuttall in Trans. Am. Phil. Soc. new ser. v, 176.—Torrey in Nicolle's Rep. 154.—Engelmann & Gray in Jour. Boston Soc. Nat. Hist. v, 262.—Chapman, Fl. S. States, 370.—Lesquereux in Owen's 2d Rep. Arkansas, 382.—Wood, Cl. Book, 600; Bot. & Fl. 277.—Gray, Manual N. States, 5 ed. 402; Proc. Am. Acad. iv, 363 (excl. var.); Syn. Fl. N. America, ii, 76.—Koch, Dendrologie, ii, 224.—Vasey, Cat. Forest Trees, 20.

Adelia acuminata, Michaux, Fl. Bor.-Am. ii, 225, t. 48.

Borya acuminata, Willdenow, Spec. iv, 711.—Aiton, Hort. Kew. 2 ed. 366.—Elliott, Sk. ii, 675.—Eaton, Manual, 6 ed. 57.—Eaton & Wright, Bot. 159.

Borya ligustrina, Willdenow, Spec. iv, 711, in part.—Aiton, Hort. Kew. 2 ed. 366, in part.—Gray, Manual N. States, 2 ed. 358, in part.

Borya nitida, Willdenow, Enum. Suppl. 66.

Bigelovia acuminata, Smith in Rees' Cycl. xxxix, No. 4.

PRIVET.

Western Georgia, western Florida, through the Gulf states to the valley of the Colorado river, Texas, and northward through Arkansas to southern Missouri and Cahokia creek, Illinois (opposite Saint Louis).

A small tree, 6 to 8 meters in height, with a trunk rarely 0.20 meter in diameter; borders of swamps and streams, in low, wet soil; common in the Gulf region, near the coast, and reaching its greatest development in southern Arkansas.

Wood heavy, soft, not strong, brittle, close-grained, compact; medullary rays numerous, thin, rather conspicuous; color, light yellow streaked with brown; the sap-wood lighter; specific gravity, 0.6345; ash, 0.72.

200.—*Chionanthus Virginica*, Linnaeus,

Spec. 1 ed. 8.—Marshall, Arbustum, 33.—Walter, Fl. Caroliniana, 60.—Wangenheim, Amer. 92.—Aiton, Hort. Kew. iii, 14; 2 ed. i, 23.—Lamarck, Ill. i, 30, t. 9, f. 1.—Willdenow, Spec. i, 46; Enum. 14; Berl. Baumz. 87.—Abbot, Insects Georgia, ii, t. 98.—Michaux, Fl. Bor.-Am. i, 3.—Vahl, Enum. i, 44.—Persoon, Syn. i, 9.—Desfontaines, Hist. Arb. i, 111.—Pursh, Fl. Am. Sept. i, 7.—Römer & Schultes, Syst. i, 72.—Nuttall, Genera, i, 5; Sylva, iii, 56, t. 88; 2 ed. ii, 122, t. 88.—Elliott, Sk. i, 6.—Hayne, Dend. Fl. 2.—Torrey, Fl. U. S. i, 7; Compend. Fl. N. States, 17.—Sprengel, Syst. i, 34.—Loddiges, Bot. Cab. t. 1264.—Guimpel, Otto & Hayne, Abb. Holz. 93, t. 73.—Beck, Bot. 232.—Eaton, Manual, 6 ed. 92.—Don, Miller's Dict. iv, 50.—London, Arboretum, ii, 1206, f. 1029, 1030.—Spach, Hist. Veg. viii, 259.—Dietrich, Syn. i, 37.—Eaton & Wright, Bot. 193.—A. De Candolle, Prodr. viii, 295.—Browne, Trees of America, 371.—Darlington, Fl. Cestrica, 3 ed. 238.—Darby, Bot. S. States, 429.—Cooper in Smithsonian Rep. 1858, 253.—Chapman, Fl. S. States, 349.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 95.—Lesquereux in Owen's 2d Rep. Arkansas, 382.—Wood, Cl. Book, 599; Bot. & Fl. 276.—Porcher, Resources S. Forests, 494.—Gray, Manual N. States, 5 ed. 401; Hall's Pl. Texas, 19; Syn. Fl. N. America, ii, 77.—Koch, Dendrologie, ii, 262.—Young, Bot. Texas, 452.—Vasey, Cat. Forest Trees, 20.

O. trifida, Mönch, Meth. 437.

O. Virginica, var. *latifolia*, Vahl, Enum. i, 44.—Aiton, Hort. Kew. 2 ed. i, 23.—Pursh, Fl. Am. Sept. i, 8.—Hayne, Dend. Fl. 2.—Don, Miller's Dict. iv, 50.

O. Virginica, var. *angustifolia*, Vahl, Enum. i, 44.—Aiton, Hort. Kew. 2 ed. i, 23.—Hayne, Dend. Fl. 2.—Watson, Dend. Brit. i, t. 1.—Don, Miller's Dict. iv, 50.

O. Virginica, var. *montana*, Pursh, Fl. Am. Sept. i, 8.—Torrey, Fl. U. S. i, 7; Compend. Fl. N. States, 17.—Beck, Bot. 232.—Eaton, Manual, 6 ed. 92.—Eaton & Wright, Bot. 194.—A. De Candolle, Prodr. viii, 295.

O. Virginica, var. *maritima*, Pursh, Fl. Am. Sept. i, 8.—Torrey, Fl. U. S. i, 7; Compend. Fl. N. States, 17.—Beck, Bot. 232.—Eaton, Manual, 6 ed. 92.—Don, Miller's Dict. iv, 50.—Eaton & Wright, Bot. 194.—A. De Candolle, Prodr. viii, 295.—Regel, Gartenflora, xvi, t. 564.

O. maritima, Loddiges, Cat. 1836.

O. heterophylla, Rafinesque, New Fl. & Bot. i, 86.

O. longifolia, Rafinesque, New Fl. & Bot. i, 87.

O. montana, Rafinesque, New Fl. & Bot. i, 88.

O. angustifolia, Rafinesque, New Fl. & Bot. i, 88.

FRINGE TREE. OLD MAN'S BEARD.

Lancaster county and the banks of the Brandywine, Chester county, Pennsylvania, south to Tampa bay, Florida, and through the Gulf states to southern Arkansas and the valley of the Brazos river, Texas.

A small tree, 6 to 10 meters in height, with a trunk 0.15 to 0.20 meter in diameter; generally along streams in low, rich soil; very common in cultivation.

Wood heavy, hard, close-grained, compact; layers of annual growth marked by several rows of large open ducts, connected as in that of *Bumelia* by branching groups of similar ducts; medullary rays numerous, obscure; color, light brown, the sap-wood lighter; specific gravity, 0.6372; ash, 0.51.

A decoction of the tonic and anti-periodic bark of the root sometimes employed in the treatment of intermittent fevers (*Am. Jour. Pharm.* xlv, 398.—*U. S. Dispensatory*, 14 ed. 1612).

201.—*Osmanthus Americanus*, Benthams & Hooker,

Genera, ii, 667.—Gray, *Syn. Fl. N. America*, ii¹, i, 78.

Olea Americana, Linnaeus, *Mant.* 24.—Marshall, *Arbustum*, 98.—Lamarck, *Dict.* iv, 543; *Ill.* i, 28.—Aiton, *Hort. Kew.* i, 14; 2 ed. i, 22.—Willdenow, *Spec.* i, 45; *Enum.* 13.—Michaux, *Fl. Bor.-Am.* ii, 222.—Vahl, *Enum.* i, 41.—Persoon, *Syn.* i, 9.—Desfontaines, *Hist. Arb.* i, 112.—Nouveau Duhamel, v, 67.—Michaux f. *Hist. Arb. Am.* iii, 50, t. 6; *N. American Sylva*, ii, 3 ed. 128, t. 86.—Pursh, *Fl. Am. Sept.* i, 7.—Römer & Schultes, *Syst.* i, 70.—Rafinesque, *Fl. Ludoviciana*, 38.—Nuttall, *Genera*, i, 5.—Elliott, *Sk.* i, 5.—Sprengel, *Syst.* i, 34.—Croom in *Am. Jour. Sci.* 1 ser. xxvi, 315.—Dietrich, *Syn.* i, 37.—Don, *Miller's Dict.* iv, 48.—Spach, *Hist. Veg.* viii, 267.—Eaton, *Manual*, 6 ed. 239.—Dietrich, *Syn.* i, 37.—Eaton & Wright, *Bot.* 333.—A. De Candolle, *Prodr.* viii, 286.—Browne, *Trees of America*, 381.—Darby, *Bot. S. States*, 429.—Cooper in *Smithsonian Rep.* 1858, 253.—Chapman, *Fl. S. States*, 369.—Curtis in *Rep. Geological Surv. N. Carolina*, 1860, iii, 57.—Lesquereux in *Owen's 2d Rep. Arkansas*, 382.—Wood, *Cl. Book*, 599; *Bot. & Fl.* 276.—Porcher, *Resources S. Forests*, 493.—Gray, *Manual N. States*, 5 ed. 401.—Young, *Bot. Texas*, 451.—Vasey, *Cat. Forest Trees*, 20.

DEVIL WOOD.

Southern Virginia, south to cape Canaveral and Tampa bay, Florida, and through the Gulf states to eastern Louisiana, near the coast.

A small tree, 10 to 15 meters in height, with a trunk sometimes 0.30 meter in diameter; borders of streams and pine-barren swamps, in moist, rich soil.

Wood heavy, very hard and strong, close-grained, unwedgeable, difficult to work, containing many radiating groups of open cells parallel to the thin, obscure, medullary rays; color, dark brown, the thick sap-wood light brown or yellow; specific gravity, 0.8111; ash, 0.46.

BORRAGINACEÆ.

202.—*Cordia Sebestena*, Linnaeus,

Spec. 1 ed. 190.—Jacquin, *Amer. t.* 42.—Lamarck, *Ill.* i, 421, t. 96, f. 1.—Willdenow, *Spec.* i, 1073; *Enum.* 248.—Andrews, *Bot. Rep.* iii, 157, t. 157.—Poiret in Lamarck, *Dict.* vii, 45.—Persoon, *Syn.* i, 166.—Trattinick, *Archiv.* t. 354.—Römer & Schultes, *Syst.* iv, 452.—Sprengel, *Syst.* i, 649.—*Bot. Mag.* t. 794.—Aiton, *Hort. Kew.* 2 ed. ii, 8.—Descourtilz, *Fl. Antilles*, iv, 205, t. 277.—Chamisso in Linnaeus, vi, 755.—Audubon, *Birds*, t. 177.—Don, *Miller's Dict.* iv, 375.—Dietrich, *Syn.* i, 611.—Nuttall, *Sylva*, iii, 81, t. 106; 2 ed. ii, 145, t. 106.—Cooper in *Smithsonian Rep.* 1858, 265.—Grisebach, *Fl. British West Indies*, 478.—Gray, *Syn. Fl. N. America*, ii¹, 180.

? *C. juglandifolia*, Jacquin, *Amer. t.* 43.

C. speciosa, Willdenow in Römer & Schultes, *Syst.* iv, 799.—A. De Candolle, *Prodr.* ix, 476.

Sebestena scabra, Rafinesque, *Sylva Telluriana*, 38.

GEIGER TREE.

Semi-tropical Florida, on the southern keys; rare; in the West Indies.

A small tree, sometimes 8 meters in height, with a trunk 0.06 to 0.08 meter in diameter; rich hummock soil; ornamental and becoming a large tree in cultivation.

Wood heavy, hard, close-grained, compact, satiny, containing few scattered, small, open ducts; medullary rays very numerous, thin, conspicuous; color, dark brown, the thick sap-wood light brown or yellow; specific gravity, 0.7108; ash, 4.22.

203.—*Cordia Boissieri*, A. De Candolle,

Prodr. ix, 478.—Torrey, Bot. Mex. Boundary Survey, 135.—Cooper in Smithsonian Rep. 1860, 442.—Gray, Syn. Fl. N. America, iii¹, 180.

Texas, valley of the Rio Grande, westward to New Mexico and southward into Mexico.

A small tree, rarely 8 meters in height, with a trunk 0.12 to 0.15 meter in diameter, or more often reduced to a low shrub.

Wood light, rather soft, close-grained, compact, containing many small scattered open ducts; medullary rays very numerous, thin, conspicuous; color, dark brown, the sap-wood light brown; specific gravity, 0.6790; ash, 3.53.

204.—*Bourreria Havanensis*, Miers,

Bot. Contrib. ii, 238.—Gray, Syn. Fl. N. America, iii¹, 181.

Ehretia Havanensis, Willdenow in Roemer & Schultes, Syst. iv, 805.—Humboldt, Bonpland & Kunth, Nov. Gen. & Spec. vii, 206.—A. De Candolle, Prodr. ix, 508.

Ehretia tomentosa, Lamarck, Ill. i, 425.—Poiret, Suppl. ii, 1.—Sprengel, Syst. i, 648.—Dietrich, Syn. i, 630.

B. tomentosa, Don, Miller's Dict. iv, 390.

B. recurva, Miers, Bot. Contrib. ii, 238.

B. ovata, Miers, Bot. Contrib. ii, 238.

Ehretia Bourreria, Chapman, Fl. S. States, 329 [not Linnaeus].—Vasey, Cat. Forest Trees, 19.

B. tomentosa, var. *Havanensis*, Grisebach, Fl. British West Indies, 482.

STRONG BARK.

Semi-tropical Florida, southern keys (Key Largo, Elliott's Key, etc.); in the West Indies.

A small tree, 10 or, exceptionally, 15 meters (Key Largo, *Curtiss*) in height, with a trunk 0.20 to 0.25 meter in diameter; the large specimens generally hollow and defective.

A form (generally shrubby in Florida) with scabrous or hispidulous leaves is—

var. *radula*, Gray, Syn. Fl. N. America, iii¹, 181.

Ehretia radula, Poiret, Suppl. ii, 2.—Dietrich, Syn. i, 630.—A. De Candolle, Prodr. ix, 506.—Chapman, Fl. S. States, 329.

B. radula, Don, Miller's Dict. iv, 390.—Chamisso in Linnaea, viii, 120.—Miers, Bot. Contrib. ii, 238.

Cordia Floridana, Nuttall, Sylva, iii, 83, t. 107; 2 ed. ii, 147, t. 107.—Cooper in Smithsonian Rep. 1858, 265.

Wood heavy, very hard, strong, very close-grained, compact, susceptible of a beautiful polish; medullary rays numerous, obscure; color, brown streaked with orange, the sap-wood not distinguishable; specific gravity, 0.8073; ash, 2.79.

205.—*Ehretia elliptica*, De Candolle,

Prodr. ix, 503.—Torrey, Bot. Mex. Boundary Survey, 136.—Cooper in Smithsonian Rep. 1858, 266.—Miers, Bot. Contrib. ii, 228, t. 85.—Gray, Syn. Fl. N. America, iii¹, 181.

KNACKAWAY. ANAQUA.

Texas, Corpus Christi to New Braunfels (*Mohr*), and southward to the valley of the lower Rio Grande.

A tree 10 to 15 meters in height, with a trunk sometimes 0.50 meter in diameter; generally along borders of streams, in rich loam, and reaching its greatest development between the Guadalupe and Nueces rivers, 50 to 75 miles from the Gulf coast.

Wood heavy, hard, not strong, very close-grained, compact, unwedgeable, containing many small open ducts arranged in numerous concentric rings within the layers of annual growth, these marked by several rows of larger ducts; medullary rays numerous, thin; color, light brown, the sap-wood a little lighter; specific gravity, 0.6440; ash, 1.31.

BIGNONIACEÆ.

206.—*Catalpa bignonioides*, Walter,

Fl. Caroliniana, 64.—De Candolle, Prodr. ix, 226.—Darlington, Fl. Cestrica, 3 ed. 182.—Cooper in Smithsonian Rep. 1858, 253.—Chapman, Fl. S. States, 285.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 50.—Wood, Cl. Book, 513; Bot. & Fl. 218.—Bureau, Mon. Bignoniaceæ, t. 25.—Gray, Manual N. States, 5 ed. 321, in part; Syn. Fl. N. America, ii¹, 319, in part.—Koch, Dendrologie, ii, 302.—Young, Bot. Texas, 385.—Vasey, Cat. Forest Trees, 19, in part.—Guibourt, Hist. Drogues, 7 ed. ii, 548.

Bignonia Catalpa, Linnæus, Spec. 1 ed. 622 (excl. syn.).—Lamarck, Dict. i, 417.—Marshall, Arbustum, 21.—Wangenheim, Amer. 58, t. 20, f. 45.—Willdenow, Spec. iii, 289; Enum. 649.—Michaux, Fl. Bor.-Am. ii, 25.—Desfontaines, Hist. Arb. i, 189.—Michaux f. Hist. Arb. Am. iii, 217, t. 6; N. American Sylva, 3 ed. ii, 55, t. 64.—Barton, Prodr. Fl. Philadelph. 66.—Rafinesque, Fl. Ludoviciana, 159.—Porcher, Resources S. Forests, 460.—Maout & Decaisne, Bot. English ed. 602 & f.

C. cordifolia, Jaume St. Hilaire in Nouveau Duhamel, ii, 13, in part (excl. t. 5).—Barton, Compend. Fl. Philadelph. i, 9.—Nuttall, Genera, i, 10.—Elliott, Sk. i, 24.—Torrey, Fl. U. S. i, 16; Compend. Fl. N. States, 20.—Beck, Bot. 245.—Eaton, Manual, 6 ed. 85.—Darlington, Fl. Cestrica, 2 ed. 363.—Spach, Hist. Veg. ix, 132.—Eaton & Wright, Bot. 184.—Darby, Bot. S. States, 439.

C. syriacæfolia, Sims, Bot. Mag. t. 1094.—Schkuhr, Handb. t. 175.—Aiton, Hort. Kew. 2 ed. i, 24.—Pursh, Fl. Am. Sept. i, 10.—Eaton, Manual, 8; 6 ed. 85.—Meyer, Prim. Fl. Esseq. 3.—Hayne, Dend. Fl. 2.—Loddiges, Bot. Cab. t. 1285.—Sprengel, Syst. i, 70.—Sertum Botanicum, i, t.—Lindley, Fl. Med. 499; Penn. Cycl. vi, 363.—Don, Miller's Diet. iv, 230.—London, Arboretum, iii, 1261 & t.—Dietrich, Syn. i, 82.—Nuttall, Sylva, iii, 77; 2 ed. ii, 140.—Torrey, Fl. N. York, ii, 25.—Browne, Trees of America, 406.

C. communis, Du Mont, Bot. Cult. 2 ed. iii, 242.

CATALPA. CATAWBA. BEAN TREE. CIGAR TREE. INDIAN BEAN.

Southwestern Georgia, valleys of the Little and Apalachicola rivers, western Florida, and through central Alabama and Mississippi.

A low, much-branched tree, 12 to 15 meters in height, with a trunk 0.50 to 0.75 meter in diameter; borders of streams and swamps, in rich loam; rare and local; long cultivated for ornament, and now extensively naturalized throughout the middle and southern Atlantic states.

Wood light, soft, not strong, coarse-grained, compact, very durable; layers of annual growth clearly marked by many rows of large open ducts; medullary rays numerous, obscure; color, light brown, the thin (one or two years') sap-wood lighter, often nearly white; specific gravity, 0.4474; ash, 0.38; used and highly valued for fence posts, rails, etc.; a reputed emetic.

A decoction of the seeds and dried bark occasionally used in cases of asthma and bronchitis (*Am. Jour. Pharm.* xlii, 204.—*U. S. Dispensatory*, 14 ed. 1608.—*Nat. Dispensatory*, 2 ed. 367).

207.—*Catalpa speciosa*, Warder;

Engelmann in Coulter's Bot. Gazette, v, 1.—Sargent in London Gard. Chronicle 1879, 784.—Ridgway in Proc. U. S. Nat. Mus. 1882, 70.—Barnes in Coulter's Bot. Gazette, ix, 74.

C. cordifolia, Jaume St. Hilaire in Nouveau Duhamel, ii, 13, in part, t. 5.—Nuttall in Trans. Am. Phil. Soc. 2 ser. v, 183.

C. bignonioides, Lesquereux in Owen's 2d Rep. Arkansas, 375 [not Walter.].—Gray, Manual N. States, 5 ed. 321, in part; Syn. Fl. N. America, ii¹, 319, in part.—Vasey, Cat. Forest Trees, 19, in part.—Broadhead in Coulter's Bot. Gazette, iii, 59.

WESTERN CATALPA.

Valley of the Vermilion river, Illinois, through southern Illinois and Indiana, western Kentucky and Tennessee, southeastern Missouri and western Arkansas.

A tree 20 to 35 or, exceptionally, 45 meters in height (*Ridgway*), with a trunk 1 to 2 meters in diameter; borders of streams and swamps, in rich bottom lands; common and reaching its greatest development in the valley of the lower Wabash river; cultivated and now widely naturalized through southern Arkansas, western Louisiana, and eastern Texas.

Wood light, soft, not strong, coarse-grained, compact, very durable in contact with the soil; layers of annual growth clearly marked by several rows of large open ducts; medullary rays numerous, obscure; color, brown, the thin sap-wood lighter; specific gravity, 0.4165; ash, 0.39; largely used for railway ties, fence posts, rails, etc., and adapted for cabinet work and interior finish.

208.—*Chilopsis saligna*, D. Don,

Edinburgh Phil. Jour. ix, 261.—Don, Miller's Dict. iv, 223.—Dietrich, Syn. iii, 566.—Gray in Bot. California, i, 587; Syn. Fl. N. America, ii, 320.—Vasey, Cat. Forest Trees, 19.—Rothrock in Wheeler's Rep. vi, 217.—Hemslay, Bot. Am.-Cent. ii, 494.—Rusby in Bull. Torrey Bot. Club, ix, 54.

Bignonia linearis, Cavanilles, Icon. iii, 35, t. 269.

C. linearis, De Candolle, Prodr. ix, 227.—Cooper in Smithsonian Rep. 1858, 266.

C. glutinosa, Engelmann in Wislizenus' Rep. 10.

DESERT WILLOW.

Valley of the Rio Grande, Texas (Laredo, *Letterman*), west through southern New Mexico and Arizona to the San Geronio pass and the San Felipe cañon, San Diego county, California; southward into northern Mexico.

A small tree, 6 to 8 meters in height, with a trunk sometimes 0.30 meter in diameter; *mesas* and banks of depressions and water-courses in the desert; the large specimens generally hollow and defective.

Wood light, soft, not strong, close-grained, checking in drying, containing many scattered, small, open ducts, the layers of annual growth marked by several rows of larger ducts; medullary rays numerous, obscure; color, brown streaked with yellow, the sap-wood much lighter; specific gravity, 0.5902; ash, 0.37.

209.—*Crescentia cucurbitina*, Linnæus,

Mant. 2 ed. 250.—Swartz, Obs. 234.—Willdenow, Spec. iii, 311.—Persoon, Syn. ii, 168.—Aiton, Hort. Kew. 2 ed. iv, 37.—Gærtner f. Fruct. Suppl. 230, t. 223.—Dietrich, Syn. iii, 567.—Don, Miller's Dict. iv, 232.—De Candolle, Prodr. ix, 246.—Seemann in Jour. Bot. & Kew Gard. Misc. vi, 274; ix, 142.—Walpers, Ann. v, 524.—Grisebach, Fl. British West Indies, 445.—Hemslay, Fl. Am. Cent. ii, 489.

C. ovata, Burmann, Fl. Ind. 132.

C. latifolia, Lamarck, Dict. i, 558; Ill. iii, 96, t. 547.—Descourtilz, Fl. Antilles, iii, 143, t. 182.

C. lethifera, Tussac, Fl. Antilles, iv, 50, t. 17.

C. toxicaria, Tussac, Fl. Antilles, iv, 50, t. 17.

C. obovata, Benthams, Bot. Sulphur, 130, t. 46.

BLACK CALABASH TREE.

Semi-tropical Florida, near Miami, and on Little river (*Garber*, *Curtiss*); in the West Indies.

A small tree, in Florida rarely exceeding 6 meters in height, with a trunk 0.10 to 0.12 meter in diameter.

Wood heavy, hard, very close-grained, compact, containing many small, regularly-distributed, open ducts; medullary rays thin, hardly distinguishable; color, light brown tinged with orange, the sap-wood lighter; specific gravity, 0.6319; ash, 1.35.

VERBENACEÆ.

210.—*Citharexylum villosum*, Jacquin,

Coll. i, 72; Icon. Rar. t. 118.—Persoon, Syn. ii, 142.—Aiton, Hort. Kew. 2 ed. iv, 36.—Dietrich, Syn. iii, 614.—Schauer in De Candolle, Prodr. xi, 610.—Walpers, Rep. iv, 76.—Chapman, Fl. S. States, 309.—Vasey, Cat. Forest Trees, 19.—Gray, Syn. Fl. N. America, ii, 340.—Hemslay, Bot. Am.-Cent. ii, 537.

FIDDLE WOOD.

Semi-tropical Florida, cape Canaveral to the southern keys (Pumpkin Key, *Curtiss*); and through the West Indies to Mexico.

A small tree, rarely exceeding in Florida 6 meters in height, with a trunk 0.10 to 0.15 meter in diameter, or north of bay Biscayne reduced to a low, much-branched shrub; common and reaching within the United States its greatest development on the shores of bay Biscayne, Lost Man's river, etc.

Wood heavy, exceedingly hard, strong, close-grained, compact, susceptible of a fine polish, containing numerous small, regularly-distributed, open ducts; color, clear bright red, the sap-wood lighter; specific gravity, 0.8710; ash, 0.52.

211.—*Avicennia nitida*, Jacquin,

Amer. 177, t. 112, f. 1.—Persoon, Syn. ii, 143.—Chamisso in Linnæa, vii, 370.—Sprengel, Syst. ii, 768.—Martius, Mat. Med. Brasil. 49; Bot. Brasil. ix, 303.—Dietrich, Syn. iii, 619.—Schauer in De Candolle, Prodr. xi, 699.—Grisebach, Fl. British West Indies, 502.—Gray, Syn. Fl. N. America, ii¹, 341.

A. tomentosa, Meyer, Prim. Fl. Esseq. 221 [not Jacquin].—Nuttall, Sylva, iii, 79, t. 105; 2 ed. ii, 143, t. 105.—Cooper in Smithsonian Rep. 1858, 265.—Chapman, Fl. S. States, 310.—Vasey, Cat. Forest Trees, 19.

A. oblongifolia, Nuttall?; Chapman, Fl. S. States, 310.—Vasey, Cat. Forest Trees, 19.

BLACK MANGROVE. BLACK TREE. BLACK WOOD.

Florida coast, Saint Augustine to the southern keys, and from Cedar Keys to cape Sable; deltas of the Mississippi river; through the West Indies to Brazil.

A tree 6 to 9 meters in height, with a trunk 0.25 to 0.30 meter in diameter, or, exceptionally, 20 to 23 meters in height, with a trunk 0.60 meter in diameter; north of Mosquito inlet reduced to a low shrub; common along saline shores and swamps, throwing up many leafless, corky stems, and forming, with the red mangrove (*Rhizophora*), impenetrable thickets, or, more rarely, scattered and round-headed; reaching its greatest development in the United States on the west coast of Florida, north of cape Sable.

Wood very heavy, hard, rather coarse-grained, compact, the eccentric layers of annual growth marked by several rows of large open ducts; medullary rays numerous, thin; color, dark brown or nearly black, the sap-wood brown; specific gravity, 0.9138; ash, 2.51.

NYCTAGINACEÆ.

212.—*Pisonia obtusata*, Swartz,

Fl. Ind. Occ. 1960.—Jacquin, Hort. Schönbr. iii, 36, t. 314.—Lamarck, Ill. iii, 449, t. 861.—Dietrich, Syn. ii, 1226.—Choisy in De Candolle, Prodr. xiii², 443.—Chapman, Fl. S. States, 374.—Grisebach, Fl. British West Indies, 71.—Vasey, Cat. Forest Trees, 21.

PIGEON WOOD. BEEF WOOD. CORK WOOD. PORK WOOD.

Semi-tropical Florida, cape Canaveral to the southern keys; through the West Indies.

A tree 9 to 15 meters in height, with a trunk 0.25 to 0.45 meter in diameter; saline shores and beaches, reaching its greatest development in Florida on Elliott's and Old Rhodes Keys.

Wood heavy, rather soft, weak, coarse-grained, compact, containing numerous large open ducts; layers of annual growth and medullary rays hardly distinguishable; color, yellow tinged with brown, the sap-wood darker; specific gravity, 0.6529; ash, 7.62; probably of little value.

NOTE.—The semi-prostrate and vine-like trunks of *P. aculeata*, Linnæus, of the same region, although attaining a considerable size, cannot be properly considered arborescent.

POLYGONACEÆ.

213.—*Coccoloba floridana*, Meisner;

De Candolle, Prodr. xiv, 165.—Chapman, Fl. S. States, 392.—Porcher, Resources S. Forests, 376.—Vasey, Cat. Forest Trees, 21.

C. parvifolia, Nuttall, Sylva, iii, 25, t. 89; 2 ed. ii, 95, t. 89 [not Poiret].—Cooper in Smithsonian Rep. 1858, 265.

PIGEON PLUM.

Semi-tropical Florida, cape Canaveral to the southern keys, and from cape Romano to cape Sable.

A tree 15 to 18 meters in height, with a trunk 0.30 to 0.60 meter in diameter; one of the largest and most common trees of the region.

Wood very heavy, exceedingly hard, strong, brittle, very close-grained, inclined to check in drying, containing few small, scattered, open ducts; layers of annual growth and numerous medullary rays obscure; color, rich dark brown tinged with red, the sap-wood lighter; specific gravity, 0.9835; ash, 5.03; valuable and somewhat used for cabinet-making.

The edible and abundant grape-like fruit, ripening in February and March, is eagerly devoured by raccoons and other animals.

214.—*Coccoloba uvifera*, Jacquin,

Amer. 112, t. 73.—Gärtner, Fruct. i, 214, t. 45, f. 3.—Aiton, Hort. Kew. ii, 34; 2 ed. ii, 421.—Lamarck, Ill. ii, 445, t. 316, f. 2.—Willdenow, Spec. ii, 457; Enum. 431.—Poirer in Lamarck, Dict. vi, 61.—Persoon, Syn. i, 442.—Titford, Hort. Bot. Am. 61.—Aiton, Hort. Kew. 2 ed. ii, 421.—Sprengel, Syst. ii, 252.—Descourtilz, Fl. Antilles, ii, 41, t. 77.—Bot. Mag. t. 3130.—Rafinesque, Fl. Telluriana, ii, 34.—Spach, Hist. Veg. x, 542.—Dietrich, Syn. Fl. ii, 1326.—Nuttall, Sylva, iii, 23, t. 88; 2 ed. ii, 93, t. 88.—Carson, Med. Bot. ii, 21, t. 67.—Meisner in De Candolle, Prodr. xiv, 152; Bot. Brasil. v¹, 42.—Cooper in Smithsonian Rep. 1858, 265.—Chapman, Fl. S. States, 391.—Porcher, Resources S. Forests, 376.—Grisebach, Fl. British West Indies, 161.

Polygonum uvifera, Linnæus, Spec. 1 ed. 365.

SEA GRAPE.

Semi-tropical Florida, Mosquito inlet to the southern keys, west coast, Tampa bay to cape Sable; through the West Indies to Brazil.

A low tree, rarely exceeding in Florida 4 meters in height, with a gnarled and contorted trunk often 0.90 to 1.20 meter in diameter, or reduced to a low, generally prostrate shrub; saline shores and beaches; common.

West Indian forms, differing in the shape of the leaves, etc., are—

var. *ovalifolia*, Meisner, l. c.

var. *Læganensis*, Meisner, l. c.

C. Læganensis, Jacquin, Amer. 113, t. 178, f. 33.

Wood very heavy, hard, very close-grained, inclined to check in drying, susceptible of a beautiful polish, containing few scattered, rather small, open ducts; layers of annual growth and numerous medullary rays hardly distinguishable; color, rich dark brown or violet, the sap-wood lighter; specific gravity, 0.9635; ash, 1.37; valuable for cabinet-making.

The edible fruit of agreeable subacid flavor.

L A U R A C E Æ .

215.—*Persea Carolinensis*, Nees,

Syst. Laurinarum, 150.—Spach, Hist. Veg. x, 492.—Dietrich, Syn. ii, 1339.—Cooper in Smithsonian Rep. 1858, 254.—Chapman, Fl. S. States, 63.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 63.—Wood, Cl. Book, 620; Bot. & Fl. 290.—Meisner in De Candolle, Prodr. xv¹, 50.—Gray, Manual N. States, 5 ed. 422; Hall's Pl. Texas, 473.—Young, Bot. Texas, 473.—Vasey, Cat. Forest Trees, 21.

Laurus Borbonia, Linnæus, Spec. 1 ed. 370, in part.—Marshall, Arbustum, 73.—Walter, Fl. Caroliniana, 133.—Aiton, Hort. Kew. ii, 39; 2 ed. ii, 429.—Lamarck, Dict. iii, 450.—Willdenow, Spec. ii, 481.—Desfontaines, Hist. Arb. i, 65.—Nouveau Duhamel, ii, 163.

Laurus Carolinensis, Catesby, Carol. i, 63, t. 63.—Michaux, Fl. Bor.-Am. i, 245.—Persoon, Syn. i, 449.—Desfontaines, Hist. Arb. i, 65.—Poirer, Suppl. iii, 321.—Willdenow, Enum. Suppl. 22.—Michaux f. Hist. Arb. Am. iii, 180, t. 2; N. American Sylva, 3 ed. ii, 116, t. 82.—Pursh, Fl. Am. Sept. i, 276.—Elliott, Sk. i, 461.—Sprengel, Syst. ii, 665.—Torrey, Compend. Fl. N. States, 174.—Beck, Bot. 305.—Eaton, Manual, 6 ed. 199.—London, Arboretum, iii, 1299, f. 1168, 1169.—Eaton & Wright, Bot. 293.—Browne, Trees of America, 414.—Darby, Bot. S. States, 491.—Schnizlein, Icon. t. 106, f. 5-12.

Laurus Carolinensis, var. *glabra*, Pursh, Fl. Am. Sept. i, 276.

Laurus Carolinensis, var. *obtus*a, Pursh, Fl. Am. Sept. i, 276.

Laurus Caroliniana, Poirer, Suppl. iii, 323.—Nuttall, Genera, i, 258.

P. Borbonia, Sprengel, Syst. ii, 268.

P. Carolinensis, var. *glabriuscula*, Meisner in De Candolle, Prodr. xv¹, 51.

RED BAY.

Southern Delaware?, south to bay Biscayne and cape Romano, Florida, and through the Gulf states to southern Arkansas and the valley of the Trinity river, Texas, near the coast.

A tree 15 to 20 meters in height, with a trunk 0.60 to 0.90 meter in diameter; borders of streams and swamps, in low, rich soil.

Wood heavy, hard, very strong, brittle, very close-grained, compact, susceptible of a beautiful polish, containing many evenly-distributed open ducts; medullary rays numerous, thin; color, bright red, the sap-wood much lighter; specific gravity, 0.6429; ash, 0.76; formerly somewhat used in ship-building, interior finish, and for cabinet work.

Var. *palustris*, Chapman,

Fl. S. States, 393.

Laurus Carolinensis, var. *pubescens*, Pursh, Fl. Am. Sept. i, 276.*P. Carolinensis*, var. *pubescens*, Meisner in De Candolle, Prodr. xv¹, 51.

North Carolina to Alabama, generally near the coast.

A small tree, 9 to 12 meters in height, with a trunk rarely exceeding 0.30 meter in diameter; low, sandy banks of pine-barren streams and swamps; well distinguished from the species by the longer peduncles densely clothed, as are the young shoots and under sides of the leaves, with short, brown tomentum, and by the somewhat coarser-grained orange-colored wood.

Wood heavy, soft, strong, close-grained, compact, containing numerous rather large open ducts; medullary rays numerous, thin; color, orange streaked with brown; the sap-wood light brown or gray; specific gravity, 0.6396; ash, 0.37.

216.—*Nectandra Willdenoviana*, Nees,Syst. Laurinarum, 290, 321.—Meisner in De Candolle, Prodr. xvi², 165.*Laurus sanguinea*, Swartz, Fl. Ind. Occ. ii, 707.

Laurus Catesbyana, Michaux, Fl. Bor.-Am. i, 244.—Poiret, Suppl. iii, 321.—Pursh, Fl. Am. Sept. i, 275.—Elliott, Sk. i, 462.—Sprengel, Syst. ii, 265.—Eaton, Manual, 6 ed. 199.—Eaton & Wright, Bot. 294.—Darby, Bot. S. States, 491.

Laurus Catesbæi, Persoon, Syn. i, 499.—Nuttall, Genera, i, 258.*Gymnobalanus Catesbyana*, Nees, Syst. Laurinarum, 483.*N. Bredemeieriana*, Nees in Linnæa, xxi, 505.*Persea Catesbyana*, Chapman, Fl. S. States, 393.—Vasey, Cat. Forest Trees, 21.

LANOE WOOD.

Semi-tropical Florida, cape Canaveral and cape Romano to the southern keys; through the West Indies to Central America.

A small tree, 6 to 9 meters in height, with a trunk rarely exceeding 0.15 meter in diameter; common and reaching its greatest development in Florida on the shores of bay Biscayne and in the neighborhood of cape Romano.

Wood heavy, hard, close-grained, checking in drying, containing many small, regularly-distributed, open ducts; medullary rays numerous, thin; color, rich dark brown, the sap-wood bright yellow; specific gravity, 0.7693; ash, 0.60.

217.—*Sassafras officinale*, Nees,

Handb. der Med. Pharm. Bot. ii, 418; Syst. Laurinarum, 488.—Hayne, Arzn. i, 12, t. 19.—Lindley, Fl. Med. 338.—Dietrich, Syn. ii, 1357.—Spach, Hist. Veg. x, 503.—Torrey, Fl. N. York, ii, 158.—Emerson, Trees Massachusetts, 359; 2 ed. ii, 359 & t.—Griffith, Med. Bot. 551.—Darlington, Fl. Cestrica, 3 ed. 251.—Spruce in Hooker's London Jour. Bot. vii, 278.—Cooper in Smithsonian Rep. 1858, 254.—Chapman, Fl. S. States, 394.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 63.—Lesquereux in Owen's 2d Rep. Arkansas, 384.—Wood, Cl. Book, 620; Bot. & Fl. 290.—Porcher, Resources S. Forests, 350.—Meisner in De Candolle, Prodr. xv¹, 171.—Gray, Manual N. States, 5 ed. 423; Hall's Pl. Texas, 19.—Koch, Dendrologie, ii, 364.—Young, Bot. Texas, 473.—Vasey, Cat. Forest Trees, 21.—Broadhead in Coulter's Bot. Gazette, iii, 59.—Bentley & Trimen, Med. Pl. iii, 220, t. 220.—Ridgway in Proc. U. S. Nat. Mus. 1882, 70.—Bell in Geological Rep. Canada, 1879-'80, 55^c.

Laurus Sassafras, Linnæus, Spec. 1 ed. 371.—Du Roi, Harbk. i, 356.—Kalm, Travels, English ed. i, 146, 341.—Marshall, Arbustum, 74.—Wangenheim, Amer. 82, t. 27, f. 56.—Walter, Fl. Caroliniana, 134.—Aiton, Hort. Kew. ii, 40; 2 ed. ii, 429.—Lamarck, Dict. iii, 454.—Abbot, Insects Georgia, i, t. 11.—B. S. Barton, Coll. 11, 19; ii, 27.—Willdenow, Spec. ii, 485; Enum. 435; Berl. Baumz. 208.—Michaux, Fl. Bor.-Am. i, 243.—Schkuhr, Handb. 349.—Persoon, Syn. i, 450.—Robin, Voyages, iii, 361.—Desfontaines, Hist. Arb. i, 66.—Titford, Hort. Bot. Am. 130.—Michaux f. Hist. Arb. Am. iii, 173, t. 1; N. American Sylva, 3 ed. ii, 113, t. 81.—Pursh, Fl. Am. Sept. i, 277.—Rafinesque, Fl. Ludoviciana, 25.—Bigelow, Med. Bot. ii, 142, t. 35; Fl. Boston. 3 ed. 170.—Nuttall, Genera, i, 259; Sylva, i, 88; 2 ed. i, 104.—Elliott, Sk. i, 464.—Nees, Pl. Offic. t. 131.—Torrey, Fl. U. S. i, 408; Compend. Fl. N. States, 174.—Descourtilz, Fl. Antilles, vii, 51, t. 464.—Audubon, Birds, t. 144.—Stephenson & Churchill, Med. Bot. iii, t. 126.—Beck, Bot. 305.—Eaton, Manual, 6 ed. 199.—Darlington, Fl. Cestrica, 2 ed. 254.—Eaton & Wright, Bot. 293.—Browne, Trees of America, 416.—Darby, Bot. S. States, 492.

Persea Sassafras, Sprengel, Syst. ii, 270.—Schnizlein, Icen. t. 106, f. 15-23.

SASSAFRAS.

Eastern Massachusetts, southwestern Vermont, and west through southern Ontario and central Michigan to southeastern Iowa, eastern Kansas, and the Indian territory; south to Hernando county, Florida, and the valley of the Brazos river, Texas.

A tree 12 to 15 meters in height, with a trunk 0.60 to 0.90 meter in diameter, exceptionally 24 to 27 meters in height, with a trunk 1.80 to 2.25 meters in diameter, or toward its northern limits reduced to a small tree or shrub; rich, sandy loam, reaching its greatest development in southwestern Arkansas and the Indian territory; at the south often taking possession, with the persimmon, of abandoned fields in the middle districts.

Wood light, soft, not strong, brittle, coarse-grained, very durable in contact with the soil, slightly aromatic, checking in drying; layers of annual growth clearly marked with three or four rows of large open ducts; medullary rays numerous, thin; color, dull orange-brown, the thin sap-wood light yellow; specific gravity, 0.5042; ash, 0.10; used for light skiffs, ox yokes, etc., and largely for fence posts and rails, and in cooperage.

The root, and especially its bark, enters into commerce, affording a powerful aromatic stimulant; the oil of sassafras, distilled from the root, is largely used in imparting a pleasant flavor to many articles of domestic use; the pith of the young branches infused with water furnishes a mucilage used as a demulcent in febrile and inflammatory affections (*Sharpe in Am. Jour. Pharm.* 1863, 53.—*Proctor in Proc. Am. Pharm. Assoc.* 1866, 217.—*U. S. Dispensatory*, 14 ed. 814.—*Nat. Dispensatory*, 2 ed. 1274; *Flückiger & Hanbury, Pharmacographia*, 483).

"*Gumbo filet*," a powder prepared by the Choctaw Indians of Louisiana from the mucilaginous leaves, is used at the south in the preparation of "gumbo" soup.

218.—*Umbellularia Californica*, Nuttall,

Sylva, i, 87; 2 ed. i, 102.—*Watson, Bot. California*, ii, 61.

Laurus regia, Douglas in *Companion Bot. Mag.* ii, 137.

Oreodaphne Californica, Nees, *Syst. Laurinarum*, 463.—*Bentham, Pl. Hartweg*, 334; *Bot. Sulphur*, 49.—*Dietrich, Syn.* ii, 1356.—*Hooker & Arnott, Bot. Beechey*, 389.—*Torrey in Pacific R. R. Rep.* iv, 133; v, 364; *Mex. Boundary Survey*, 184.—*Newberry in Pacific R. R. Rep.* vi, 24, 88, f. 3.—*Cooper in Smithsonian Rep.* 1858, 260.—*Bot. Mag.* t. 5320.

Tetranthera Californica, *Hooker & Arnott, Bot. Beechey*, 159.—*Meisner in De Candolle, Prodr.* xv¹, 192.—*Torrey in Bot. Wilkes Exped.* 451.

Drimophyllum pauciflorum, Nuttall, *Sylva*, i, 85, t. 22; 2 ed. i, 102, t. 22.

MOUNTAIN LAUREL. CALIFORNIA LAUREL. SPICE TREE. CAGIPUT. CALIFORNIA OLIVE. CALIFORNIA BAY TREE.

Rogue River valley, Oregon, south through the California coast ranges to San Diego county, and along the western slopes of the Sierra Nevada to the San Bernardino mountains.

An evergreen tree, 24 to 30 meters in height, with a trunk 1.20 to 1.80 meter in diameter, or toward its southern limits and at high elevations a small tree or shrub; most common and reaching its greatest development in the rich valleys of southwestern Oregon.

Wood heavy, hard, strong, close-grained, compact, susceptible of a beautiful polish, containing numerous small, regularly-distributed, open ducts; medullary rays numerous, thin; color, rich light brown, the sap-wood lighter; specific gravity, 0.6517; ash, 0.39; used on the Oregon coast in ship-building, for jaws, bitts, cleats, cross-trees, etc.; the most valuable material produced by the Pacific forests for interior and cabinet work.

The leaves yield a volatile oil, *Oreodaphne* (*Am. Jour. Pharm.* xlvii, 105).

EUPHORBIAEÆ.

219.—*Drypetes crocea*, Poiteau,

Mem. Mus. i, 159, t. 8.—*Nuttall, Sylva*, ii, 66, t. 63; 2 ed. ii, 12, t. 63.—*Cooper in Smithsonian Rep.* 1858, 265.—*Chapman, Fl. S. States*, 410.—*Grisebach, Fl. British West Indies*, 32; *Cat. Pl. Cuba*, 15.—*Müller in De Candolle, Prodr.* xv², 456.

Schæfferia lateriflora, Swartz, *Fl. Ind. Occ.* i, 329.

D. sessiliflora, Baillon, *Etud. Gen. Euphorbiaceæ*, Atlas, 45, t. 24, f. 34–40.

D. glauca, Grisebach in *Mem. Am. Acad. new ser.* viii, 157 [not Vahl].

D. crocea, var. *longipes*, Müller in De Candolle, *Prodr.* xv², 456.

GUIANA PLUM. WHITE WOOD.

Semi-tropical Florida, bay Biscayne to the southern keys; in the West Indies.

A small tree, sometimes 9 meters in height, with a trunk 0.12 to 0.17 meter in diameter.

Wood heavy, hard, not strong, brittle, close-grained, checking in drying; medullary rays numerous, thin; color, rich dark brown, the sap-wood yellow; specific gravity, 0.9209; ash, 6.14.

Var. *latifolia*, Müller,

De Candolle, Prodr. xv², 456.

D. glauca, Nuttall, Sylva, ii, 68; 2 ed. ii, 14.—Chapman, Fl. S. States, 410.

D. alba, var. *latifolia*, Grisebach in Nachrich. d. Konigl. Gesell. Wiss. Univ. Götting. 1865, 165, in part.

Semi-tropical Florida, bay Biscayne to the southern keys; in the West Indies.

A tree sometimes 12 meters in height, with a trunk 0.30 to 0.35 meter in diameter.

Wood heavy, hard, not strong, brittle, very close-grained, checking in drying; medullary rays numerous, obscure; color, brown streaked with bright yellow, the sap-wood dull brown; specific gravity, 0.9346; ash, 8.29.

Perhaps a distinct species, the fruit and flowers not recently collected.

220.—*Sebastiania lucida*, Müller;

De Candolle, Prodr. xv², 1181.

Gymnanthes lucida, Swartz, Prodr. 96.

Excoecaria lucida, Swartz, Fl. Ind. Occ. ii, 1122.—Willdenow, Spec. iv, 865.—Poiret, Suppl. i, 155.—Persoon, Syn. ii, 634.—Nuttall, Sylva, ii, 60, t. 61; 2 ed. ii, 6, t. 61.—A. de Jussieu, Tent. Euphorb. t. 16, f. 55.—Richard, Fl. Cuba, 199.—Dietrich, Syn. v, 256.—Cooper in Smithsonian Rep. 1858, 265.—Chapman, Fl. S. States, 405.—Grisebach, Fl. British West Indies, 50.—Vasey, Cat. Forest Trees, 21.

ORAB WOOD. POISON WOOD.

Semi-tropical Florida, bay Biscayne to the southern keys; common; in the West Indies.

A small tree, sometimes 9 meters in height, with a trunk 0.15 to 0.20 meter in diameter; the large specimens generally hollow and decayed.

Wood very heavy, hard, very close-grained, compact, susceptible of a beautiful polish; medullary rays numerous, obscure; color, rich dark brown streaked with yellow, the sap-wood bright yellow; specific gravity, 1.0905; ash, 2.78; now largely manufactured into canes, and furnishing valuable fuel.

221.—*Hippomane Mancinella*, Linnaeus,

Spec. 1 ed. 1191.—Jacquin, Amer. 250, t. 159.—Lamarck, Dict. ii, 694.—Aiton, Hort. Kew. iii, 378; 2 ed. v, 333.—Swartz, Obs. 369.—Willdenow, Spec. iv, 571.—Persoon, Syn. ii, 589.—Titford, Hort. Bot. Am. Suppl. 9, t. 12, f. 5.—Lamarck, Ill. iii, 374, t. 793, f. 1.—Sprengel, Syst. iii, 805.—Spach, Hist. Veg. ii, 524.—Nuttall, Sylva, ii, 54, t. 60; 2 ed. i, 202, t. 60.—Bentham, Bot. Sulphur, 163.—Richard, Fl. Cuba, 200.—Dietrich, Syn. v, 224.—Cooper in Smithsonian Rep. 1858, 265.—Baillon, Etud. Gen. Euphorbiaceæ, t. 6, f. 12-20.—Chapman, Fl. S. States, 404.—Porcher, Resources S. Forests, 120.—Grisebach, Fl. British West Indies, 50.—Regel, Gartenflora, xv, 163, t. 510.—Müller in De Candolle, Prodr. xv², 1201.—Schnizlein, Icon. t. 243, f. 3.—Maout & Decaisne, Bot. English ed. 693 & f.—Vasey, Cat. Forest Trees, 21.

Mancinella venenata, Tussac, Fl. Antilles, iii, 21, t. 5.

MANCHINEEL.

Semi-tropical Florida, on the southern keys; common; through the West Indies and Central America to the Pacific.

A small tree, in Florida rarely exceeding 4 meters in height, with a trunk 0.12 to 0.17 meter in diameter; abounding in white, milky, exceedingly caustic poisonous sap. "Rain washing the leaves becomes poisonous, and the smoke of the burning wood injures or destroys the eyes."—(A. H. Curtiss).

Wood light, soft, close-grained, compact, containing numerous evenly-distributed, small, open ducts; medullary rays numerous, obscure; color, dark brown, the thick sap-wood light brown or yellow; specific gravity (sap-wood), 0.5172; ash, 5.16.

URTICACEÆ.

222.—*Ulmus crassifolia*, Nuttall,

Trans. Am. Phil. Soc. 2 ser. v, 169.—Planchon in Ann. Sci. Nat. 3 ser. x, 279; De Candolle, Prodr. xvii, 162.—Walpers, Ann. iii, 426.—Cooper in Smithsonian Rep. 1858, 254.—Lesquereux in Owen's 2d Rep. Arkansas, 386.—Wood, Cl. Book, 633.—Gray, Hall's, Pl. Texas, 21.—Vasey, Cat. Forest Trees, 23.

U. opaca, Nuttall, Sylva, i, 35, t. 11; 2 ed. i, 51, t. 11.—Browne, Trees of America, 503.

CEDAR ELM.

Arkansas, south of the valley of the Arkansas river to the valley of the Rio Grande, Texas, extending west to Eagle Pass.

A tree 18 to 20 meters in height, with a trunk 0.60 to 0.90 meter in diameter, or toward its southern or southwestern limits much smaller; borders of streams, in rich soil; one of the most common and valuable timber trees of Texas west of the Trinity river, and reaching its greatest development in the valleys of the Guadalupe and Trinity rivers.

Wood heavy, hard, not strong, brittle, very close-grained, compact; layers of annual growth and medullary rays obscure; marked, in common with that of all the North American species, by concentric circles of irregularly-arranged groups of small open ducts; color, light brown tinged with red, the heavier sap-wood lighter; specific gravity, 0.7245; ash, 1.20; used in the manufacture of wagon hubs, saddle-trees, chairs, etc., and very largely for fencing.

223.—*Ulmus fulva*, Michaux,

Fl. Bor.-Am. i, 172.—Persoon, Syn. i, 291.—Willdenow, Enum. Suppl. 14.—Pursh, Fl. Am. Sept. i, 200.—Smith in Rees' Cycl. xxxix, No. 10.—Eaton, Manual, 31; 6 ed. 376.—Nuttall, Genera, i, 201.—Römer & Schultes, Syst. vi, 301.—Elliott, Sk. i, 333.—Hayne, Dend. Fl. 32.—Torrey, Fl. U. S. i, 299; Compend. Fl. N. States, 132; Fl. N. York, ii, 166; Fremont's Rep. 97.—Sprengel, Syst. i, 931.—Rafinesque, Med. Bot. ii, 271.—Beck, Bot. 333.—Hooker, Fl. Bor.-Am. ii, 142.—Bigelow, Fl. Boston. 3 ed. 114.—Eaton & Wright, Bot. 464.—London, Arboretum, iii, 1407, f. 1247.—Dietrich, Syn. ii, 992.—Spach in Ann. Sci. Nat. xv, 363; Hist. Veg. xi, 107.—Emerson, Trees Massachusetts, 297; 2 ed. ii, 334 & t.—Browne, Trees of America, 501.—Griffith, Med. Bot. 551.—Planchon in Ann. Sci. Nat. 3 ser. x, 276.—De Candolle, Prodr. xvii, 161.—Scheele in Römer, Texas, 446.—Walpers, Ann. iii, 426.—Richardson, Arctic Exped. 436.—Darlington, Fl. Cestrica, 3 ed. 255.—Darby, Bot. S. States, 502.—Cooper in Smithsonian Rep. 1858, 254.—Chapman, Fl. S. States, 416.—Curtis in Rep. Geological Surv. N. Carolina, iii, 1860, 55.—Lesquereux in Owen's 2d Rep. Arkansas, 386.—Wood, Cl. Book, 633; Bot. & Fl. 299.—Porcher, Resources S. Forests, 310.—Engelmann in Trans. Am. Phil. Soc. new ser. xii, 208.—Gray, Manual N. States, 5 ed. 442.—Koch, Dendrologie, ii, 422.—Young, Bot. Texas, 496.—Hayden in Warren's Rep. Nebraska & Dakota, 2 ed. 121.—Vasey, Cat. Forest Trees, 22.—Bentley & Trimen, Med. Pl. iv, 233, t. 233.—Ridgway in Proc. U. S. Nat. Mus. 1882, 72.—Bell in Geological Rep. Canada, 1879-'80, 55c.

U. pubescens, Walter, Fl. Caroliniana, 111.

U. Americana, var. *rubra*, Aiton, Hort. Kew. i, 319; 2 ed. ii, 107.—Willdenow, Spec. i, 1325.—Hayne, Dend. Fl. 31.

? *U. crispa*, Willdenow, Enum. 295; Berl. Baumz. 520.

U. rubra, Michaux f. Hist. Arb. Am. iii, 278, t. 6; N. American Sylva, 3 ed. iii, 73, t. 128.

RED ELM. SLIPPERY ELM. MOOSE ELM.

Valley of the lower Saint Lawrence river to Ontario and northern Dakota, south to the Chattahoochee region of northern Florida, central Alabama and Mississippi, and the valley of the San Antonio river, Texas.

A tree 15 to 20 meters in height, with a trunk 0.45 to 0.60 meter in diameter; borders of streams and hillsides, in rich soil.

Wood heavy, hard, strong, very close-grained, compact, durable in contact with the ground, splitting readily when green; layers of annual growth clearly marked by several rows of large open ducts; medullary rays numerous, thin; color, dark brown or red, the thin sap-wood lighter; specific gravity, 0.6956; ash, 0.83; largely used for wheel stock, fence posts, rails, railway ties, sills, etc.

The inner bark mucilaginous, nutritious, and extensively used in various medicinal preparations (*Am. Jour. Pharm.* xxiv, 180.—*Philadelphia Med. Times*, 1874, 303.—*U. S. Dispensatory*, 14 ed. 913.—*Nat. Dispensatory*, 2 ed. 1480.—*Flückiger & Hanbury, Pharmacographia*, 501).

224.—*Ulmus Americana*, Linnæus,

Spec. 1 ed. 226.—Kalm, Travels, English ed. ii, 298.—Marshall, Arbustum, 156.—Wangenheim, Amer. 46.—Gærtner, Fruct. i, 225, t. 49, f. 5.—Walter, Fl. Caroliniana, 111.—Aiton, Hort. Kew. i, 319; 2 ed. ii, 107.—Willdenow, Spec. i, 1325; Enum. 295; Suppl. 14; Berl. Baumz. 519.—Nouveau Duhamel, ii, 147.—Schkuhr, Handb. 179.—Michaux, Fl. Bor.-Am. i, 173.—Persoon, Syn. ii, 191.—Desfontaines, Hist. Arb. ii, 442.—Michaux f. Hist. Arb. Am. iii, 269, t. 4; N. American Sylva, 3 ed. iii, 67, t. 126.—Pursh, Fl. Am. Sept. i, 199.—Smith in Rees' Cycl. xxxix, No. 7.—Eaton, Manual, 31; 6 ed. 376.—Barton, Compend. Fl. Philadelph. i, 150.—Nuttall, Genera, i, 201.—Rømer & Schultes, Syst. vi, 300.—Elliott, Sk. i, 333.—Hayne, Dend. Fl. 31.—Torrey, Fl. U. S. i, 298; Compend. Fl. N. States, 132; Fl. N. York, ii, 165; Nicotlet's Rep. 160; Emory's Rep. 412.—Sprengel, Syst. i, 930.—Beck, Bot. 333.—London, Arboretum, iii, 1406, f. 1246.—Hooker, Fl. Bor.-Am. ii, 142.—Bigelow, Fl. Boston. 3 ed. 114.—Dietrich, Syn. ii, 992.—Eaton & Wright, Bot. 464.—Spach in Ann. Sci. Nat. 2 ser. xv, 364; Hist. Veg. xi, 108.—Emerson, Trees Massachusetts, 286; 2 ed. ii, 322 & t.—Browne, Trees of America, 499.—Planchon in Ann. Sci. Nat. 3 ser. x, 268; De Candolle, Prodr. xvii, 155.—Scheele in Rømer, Texas, 446.—Walpers, Ann. iii, 424.—Buckley in Am. Jour. Sci. 2 ser. xiii, 398.—Richardson, Arctic Exped. 436.—Darlington, Fl. Cestrica, 3 ed. 255.—Darby, Bot. S. States, 502.—Cooper in Smithsonian Rep. 1858, 254.—Chapman, Fl. S. States, 416.—Curtis in Rep. Geological Surv. N. Carolina, iii, 1860, 55.—Lesquereux in Owen's 2d Rep. Arkansas, 386.—Wood, Cl. Book, 633; Bot. & Fl. 298.—Porcher, Resources S. Forests, 311.—Engelmann in Trans. Am. Phil. Soc. new ser. xii, 208.—Gray, Manual N. States, 5 ed. 442.—Hall's Pl. Texas, 21.—Koch, Dendrologie, ii, 421.—Young, Bot. Texas, 496.—Winchell in Ludlow's Rep. Black Hills, 68.—Vasey, Cat. Forest Trees, 22.—Hayden in Warren's Rep. Nebraska & Dakota, 2 ed. 121.—Macoun in Geological Rep. Canada, 1875-'76, 209.—Sears in Bull. Essex Inst. xiii, 177.—Ridgway in Proc. U. S. Nat. Mus. 1882, 71.—Bell in Geological Rep. Canada, 1879-'80, 48°.

U. mollifolia, Marshall, Arbustum, 156.

U. Americana, var. *pendula*, Aiton, Hort. Kew. i, 320; 2 ed. ii, 107.—Willdenow, Spec. i, 1326.—Pursh, Fl. Am. Sept. i, 200.—Eaton, Manual, 31.—Spach in Ann. Sci. Nat. 2 ser. xv, 364; Hist. Veg. xi, 109.

U. Americana, var. *alba*, Aiton, Hort. Kew. i, 320; 2 ed. ii, 107.—Hayne, Dend. Fl. 32.

U. pendula, Willdenow, Berl. Baumz. 519.—Hayne, Dend. Fl. 33.

U. alba, Rafinesque, Fl. Ludoviciana, 115; New Fl. & Bot. i, 38.

U. Americana, var. *scabra*, Spach in Ann. Sci. Nat. 2 ser. xv, 364; Hist. Veg. ix, 109.—Walpers, Ann. iii, 424.

U. Americana, var. *Bartramii*, Walpers, Ann. iii, 424.

U. Americana, var. *aspera*, Chapman, Fl. S. States, 416.

U. Floridana, Chapman, Fl. S. States, 416.

WHITE ELM. AMERICAN ELM. WATER ELM.

Southern Newfoundland to the northern shores of lake Superior and the eastern slope of the Rocky mountains, in about latitude 52° N.; south to cape Canaveral and Pease creek, Florida, extending west in the United States to the Black hills of Dakota, central Nebraska, the Indian territory, in about longitude 100° W., and the valley of the Rio Concho, Texas.

A large tree, 30 to 35 meters in height, with a trunk 1.80 to 2.70 meters in diameter; rich, moist soil, borders of streams, etc.; toward its western and southwestern limits only in river bottoms.

Wood heavy, hard, strong, tough, rather coarse-grained, compact, difficult to split; layers of annual growth clearly marked by several rows of large open ducts; medullary rays numerous, thin; color, light brown, the sapwood somewhat lighter; specific gravity, 0.6506; ash, 0.80; largely used for wheel stock, saddle-trees, flooring, in cooperage, and now largely exported to Great Britain and used in boat- and ship-building.

225.—*Ulmus racemosa*, Thomas,

Ann. Jour. Sci. 1 ser. xix, 170 & t.—Beck, Bot. 334.—Eaton, Manual, 6 ed. 376.—Eaton & Wright, Bot. 464.—Nuttall, Sylva, i, 37, t. 12; 2 ed. i, 53, t. 12.—Torrey, Fl. N. York, ii, 166, t. 96.—Browne, Trees of America, 500.—Cooper in Smithsonian Rep. 1858, 254.—Wood, Cl. Book, 633; Bot. & Fl. 299.—Gray, Manual N. States, 5 ed. 442.—Vasey, Cat. Forest Trees, 22.—Sargent in Rep. Massachusetts Board Ag. 1878, 271.—Bell in Geological Rep. Canada, 1879-'80, 55°.—Chapman, Fl. S. States, Suppl. 649.

U. Americana, Planchon in De Candolle, Prodr. xvii, 155, in part.

ROCK ELM. CORK ELM. HICKORY ELM. WHITE ELM. CLIFF ELM.

Southwestern Vermont (*Robbins*), west through western New York, Ontario, and southern Michigan to northeastern Iowa (*Waverly*, *Bessey*), and south through Ohio to central Kentucky.

A large tree of great economic value, 20 to 30 meters in height, with a trunk sometimes 0.90 meter in diameter; low, wet clay, rich uplands, rocky declivities, or river cliffs; common and reaching its greatest development in southern Ontario and the southern peninsula of Michigan.

Wood heavy, hard, very strong, tough, very close-grained, compact, susceptible of a beautiful polish; layers of annual growth marked with one to two rows of small open ducts; medullary rays numerous, obscure; color, light clear brown often tinged with red, the thick sap-wood much lighter; specific gravity, 0.7263; ash, 0.60; largely used in the manufacture of heavy agricultural implements, wheel stock, and for railway ties, bridge timbers, sills, etc.

226.—*Ulmus alata*, Michaux,

Fl. Bor.-Am. i, 173.—Persoon, Syn. i, 291.—Michaux f. Hist. Arb. Am. iii, 275, t. 5; N. American Sylva, 3 ed. iii, 71, t. 127.—Pursh, Fl. Am. Sept. i, 200.—Nuttall, Genera, i, 201.—Römer & Schultes, Syst. vi, 209.—Elliott, Sk. i, 333.—Sprengel, Syst. i, 931.—Audubon, Birds, t. 18.—Eaton, Manual, 6 ed. 376.—London, Arboretum, iii, 1408, f. 1248.—Dietrich, Syn. ii, 992.—Eaton & Wright, Bot. 464.—Penn. Cycl. xxv, 493.—Browne, Trees of America, 502.—Planchon in Ann. Sci. Nat. 3 ser. x, 270; De Candolle, Prodr. xvii, 155.—Walpers, Ann. iii, 425.—Darby, Bot. S. States, 503.—Cooper in Smithsonian Rep. 1858, 254.—Chapman, Fl. S. States, 417.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 35.—Lesquereux in Owen's 2d Rep. Arkansas, 386.—Wood, Cl. Book, 633; Bot. & Fl. 299.—Porcher, Resources S. Forests, 311.—Gray, Manual N. States, 5 ed. 443; Hall's Pl. Texas, 21.—Young, Bot. Texas, 496.—Vasey, Cat. Forest Trees, 22.—Broadhead in Coulter's Bot. Gazette, iii, 60.—Ridgway in Proc. U. S. Nat. Mus. 1882, 70.

U. pumila, Walter, Fl. Caroliniana, 111 [not Linnaeus].

U. Americana, var. *alata*, Spach in Ann. Sci. Nat. 2 ser. xv, 364; Hist. Veg. xi, 109.

WAHOO. WINGED ELM.

Southern Virginia, south through the middle districts to the Chattahoochee region of western Florida; southern Indiana and Illinois, south to the Gulf coast, and southwest through southern Missouri, Arkansas, the eastern portions of the Indian territory to the valley of the Trinity river, Texas.

A small tree, 7 to 12 meters in height, with a trunk 0.30 to 0.60 meter in diameter; generally in dry, gravelly soil, or, rarely, along the borders of swamps and river bottoms; most common and reaching its greatest development in southern Missouri and Arkansas.

Wood heavy, hard, not strong, very close-grained, compact, unwedgeable; medullary rays distant, not conspicuous; color, brown, the sap-wood lighter; specific gravity, 0.7491; ash, 0.99; largely used for hubs, blocks, etc.

227.—*Planera aquatica*, Gmelin,

Syst. ii, 150.—Willdenow, Spec. iv, 967; Enum. Suppl. 14; Berl. Baumz. 281.—Persoon, Syn. i, 291.—Nuttall, Genera, i, 202.—Hayne, Dend. Fl. 202.—Eaton, Manual, 6 ed. 266.—Eaton & Wright, Bot. 360.—Spach in Ann. Sci. Nat. 2 ser. xv, 355; Hist. Veg. xi, 116.—Planchon in Ann. Sci. Nat. 3 ser. x, 261; De Candolle, Prodr. xvii, 167.—Walpers, Ann. iii, 428.—Cooper in Smithsonian Rep. 1858, 254.—Chapman, Fl. S. States, 417.—Wood, Cl. Book, 633; Bot. & Fl. 299.—Gray, Manual N. States, 5 ed. 443.—Koch, Dendrologie, ii, 424.—Young, Bot. Texas, 497.—Vasey, Cat. Forest Trees, 23.

Anonymos aquatica, Walter, Fl. Caroliniana, 230.

P. Gmelini, Michaux, Fl. Bor.-Am. ii, 248.—Desfontaines, Hist. Arb. ii, 446.—Römer & Schultes, Syst. vi, 305.—Elliott, Sk. i, 334.—Sprengel, Syst. i, 493.—Dietrich, Syn. i, 551.—Penn. Cycl. xxv, 490.—Darby, Bot. S. States, 503.

P. ulmifolia, Michaux f. Hist. Arb. Am. iii, 283, t. 7; N. American Sylva, 3 ed. iii, 80, t. 130.—Poiret, Suppl. iv, 429.—Nouveau Duhamel, vii, 65, t. 21.—London, Arboretum, iii, 1413, f. 1251.—Browne, Trees of America, 515.—Curtis in Rep. Geological Surv. N. Carolina, iii, 1860, 81.

? *Ulmus nemoralis*, Aiton, Hort. Kew. i, 319; 2 ed. ii, 108.—Willdenow, Spec. i, 1326; Berl. Baumz. 520.—Desfontaines, Hist. Arb. ii, 442.—Pursh, Fl. Am. Sept. i, 200.—Smith in Rees' Cycl. xxxix, No. 8.—Nuttall, Genera, i, 201.—Beck, Bot. 334.—Eaton, Manual, 6 ed. 376.—Eaton & Wright, Bot. 464.

Ulmus aquatica, Rafinesque, Fl. Ludoviciana, 165.

P. Richardi, Sprengel, Syst. i, 493, in part.—Torrey & Gray in Pacific R. R. Rep. ii, 175 [not Michaux].

Valley of the Cape Fear river, North Carolina, south to the Chattahoochee region of western Florida, and through central Alabama and Mississippi to western Louisiana and the valley of the Trinity river, Texas, extending north through Arkansas and southern Missouri to the valley of the lower Wabash river and central Kentucky.

A small tree, 9 to 12 meters in height, with a trunk 0.30 to 0.60 meter in diameter; cold, deep, inundated river swamps; rare in the Atlantic and eastern Gulf states; very common and reaching its greatest development in the Red River valley and southern Arkansas.

Wood light, soft, not strong, close-grained, compact, containing few scattered open ducts; medullary rays numerous, thin; color, light brown, the sap-wood nearly white; specific gravity, 0.5294; ash, 0.45.

228.—*Celtis occidentalis*, Linnaeus,

Spec. 2 ed. 1478.—Du Roi, Harbk. i, 141.—Marshall, Arbustum, 29.—Wangenheim, Amer. 48.—Gærtner, Fruct. i, 374, t. 77, f. 3.—Walter, Fl. Caroliniana, 250.—Aiton, Hort. Kew. iii, 437; 2 ed. v, 449.—Lamarck, Dict. iv, 137; Ill. iii, 437, t. 844, f. 1.—Abbot, Insecta Georgia, i, t. 36.—Willdenow, Spec. iv, 944; Enum. 1046; Berl. Baumz. 82.—Nouveau Duhamel, ii, 36, t. 9.—Michaux, Fl. Bor.-Am. ii, 249.—Persoon, Syn. i, 292.—Desfontaines, Hist. Arb. ii, 448.—Michaux f. Hist. Arb. Am. iii, 225, t. 8; N. American Sylva, 3 ed. iii, 38, t. 114.—Pursh, Fl. Am. Sept. i, 200.—Eaton, Manual, 31; 6 ed. 36.—Nuttall, Genera, i, 202.—Römer & Schultes, Syst. vi, 306.—Hayne, Dend. Fl. 216.—Elliott, Sk. ii, 584.—Torrey, Fl. U. S. i, 300; Compend. Fl. N. States, 132; Fl. N. York, ii, 167; Bot. Wilkes Exped. 456.—Guimpel, Otto & Hayne, Abb. Holz. 119, t. 96.—Sprengel, Syst. i, 932.—Watson, Dend. Brit. ii, 147.—Beck, Bot. 334.—Rafinesque, New Fl. & Bot. i, 32.—London, Arboretum, iii, 1417 & t.—Hooker, Fl. Bor.-Am. ii, 142.—Eaton & Wright, Bot. 186.—Spach in Ann. Sci. Nat. 2 ser. xvi, 40; Hist. Veg. xi, 133.—Penn. Cycl. xxv, 490.—Browne, Trees of America, 517.—Emerson, Trees Massachusetts, 306, t. 16; 2 ed. ii, 344 & t.—Planchon in Ann. Sci. Nat. 3 ser. x, 288; De Candolle, Prodr. xvii, 174.—Walpers, Ann. iii, 396.—Richardson, Arctic Exped. 436.—Darlington, Fl. Cestrica, 3 ed. 256.—Darby, Bot. S. States, 503.—Cooper in Smithsonian Rep. 1858, 254.—Chapman, Fl. S. States, 417.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 61.—Lesquereux in Owen's 2d Rep. Arkansas, 386.—Wood, Cl. Book, 634; Bot. & Fl. 299.—Engelmann in Trans. Am. Phil. Soc. new ser. xii, 208.—Porcher, Resources S. Forests, 312.—Gray, Manual N. States, 5 ed. 443; Hall's Pl. Texas, 21.—Koch, Dendrologie, ii, 432.—Hayden in Warren's Rep. Nebraska & Dakota, 2 ed. 121.—Vasey, Cat. Forest Trees, 23.—Burbank in Proc. Boston Soc. Nat. Hist. xviii, 215.—Putzby in Fl. des Serres, xxii, 206.—Macoun in Geological Rep. Canada, 1875-'76, 209.—Ridgway in Proc. U. S. Nat. Mus. 1882, 72.

C. crassifolia, Lamarck, Dict. iv, 133.—Nouveau Duhamel, ii, 37.—Michaux f. Hist. Arb. Am. iii, 228, t. 9; N. American Sylva, 3 ed. iii, 40, t. 115.—Pursh, Fl. Am. Sept. i, 200.—Nuttall, Genera, i, 202.—Römer & Schultes, Syst. vi, 307.—Torrey, Fl. U. S. i, 300; Compend. Fl. N. States, 132; Fremont's Rep. 97; Emory's Rep. 412.—Sprengel, Syst. i, 932.—Beck, Bot. 334.—Eaton, Manual, 6 ed. 86.—Rafinesque, New Fl. & Bot. i, 34.—London, Arboretum, iii, 1418, f. 1254.—Eaton & Wright, Bot. 186.—Spach in Ann. Sci. Nat. 2 ser. xvi, 39; Hist. Veg. xi, 130.—Penn. Cycl. xxv, 490.—Browne, Trees of America, 519.—Emerson, Trees of Massachusetts, 309; 2 ed. ii, 347 & t.

C. obliqua, Mönch, Meth. 344.

C. occidentalis, var. *scabriuscula*, Willdenow, Spec. iv, 995; Berl. Baumz. 2 ed. 82.—Hayne, Dend. Fl. 217.—London, Arboretum, iii, 1417.

C. occidentalis, var. *tenuifolia*, Persoon, Syn. i, 292.

C. cordata, Persoon, Syn. i, 292.—Desfontaines, Hist. Arb. ii, 448.—Du Mont, Cour. Bot. Cult. vi, 389.

C. laevigata, Willdenow, Berl. Baumz. 2 ed. 81; Enum. Suppl. 68.—Römer & Schultes, Syst. vi, 306.—Sprengel, Syst. i, 932.—Rafinesque, New Fl. & Bot. i, 34.—London, Arboretum, iii, 1420.—Koch, Dendrologie, ii, 432.

C. pumila, Pursh, Fl. Am. Sept. i, 200.—Römer & Schultes, Syst. vi, 306.—Torrey, Fl. U. S. i, 300; Compend. Fl. N. States, 132.—Beck, Bot. 334.—Eaton, Manual, 6 ed. 86.—Rafinesque, New Fl. & Bot. i, 35.—London, Arboretum, iii, 1420.—Eaton & Wright, Bot. 186.

C. alba, Rafinesque, Fl. Ludoviciana, 25; New Fl. & Bot. i, 32.—Planchon in De Candolle, Prodr. xvii, 177.

C. canina and *C. maritima*, Rafinesque in Am. Monthly Mag. & Crit. Rev. ii, 43, 44.

C. occidentalis, var. *cordata*, Willdenow, Berl. Baumz. 2 ed. 82.—Hayne, Dend. Fl. 217.—Römer & Schultes, Syst. vi, 306.—London, Arboretum, iii, 1417.

C. tenuifolia, Nuttall, Genera, i, 202; Sylva, i, 135; 2 ed. i, 149.—Rafinesque, New Fl. & Bot. i, 36.

C. occidentalis, var. *integrifolia*, Nuttall, Genera, i, 202.—Chapman, Fl. S. States, 417.—Wood, Cl. Book, 634; Bot. & Fl. 299.

C. Mississippiensis, Bosc, Dict. Ag. new ed. x, 41.—Poiret, Suppl. iii, 688.—Spach in Ann. Sci. Nat. 2 ser. xvi, 42; Hist. Veg. xi, 136.—Planchon in Ann. Sci. Nat. 3 ser. x, 287; De Candolle, Prodr. xvii, 176.—Walpers, Ann. iii, 397.—Cooper in Smithsonian Rep. 1858, 254.—Lesquereux in Owen's 2d Rep. Arkansas, 386.—Gray, Manual N. States, 5 ed. 443; Hall's Pl. Texas, 21.—Vasey, Cat. Forest Trees, 23.—Ridgway in Proc. U. S. Nat. Mus. 1882, 72.

C. integrifolia, Nuttall in Trans. Am. Phil. Soc. new ser. v, 169.—Cooper in Smithsonian Rep. 1858, 254.

C. longifolia, Nuttall in Trans. Am. Phil. Soc. new ser. v, 169; Sylva, i, 134, t. 40; 2 ed. i, 148, t. 40.—Rafinesque, New Fl. & Bot. i, 33.—Planchon in De Candolle, Prodr. xvii, 177.

C. heterophylla, *C. patula*, *C. Floridiana*, *C. fuscata*, *C. salicifolia*, *C. morifolia*, *C. maritima*, Rafinesque, New Fl. & Bot. i, 31-37.

C. occidentalis, var. *grandidentata*, Spach in Ann. Sci. Nat. 2 ser. xvi, 40; Hist. Veg. xi, 133.—Walpers, Ann. iii, 396.

C. occidentalis, var. *serrulata*, Spach in Ann. Sci. Nat. 2 ser. xvi, 41; Hist. Veg. xi, 134.—Walpers, Ann. iii, 396.

C. crassifolia, var. *tiliaefolia*, Spach in Ann. Sci. Nat. 2 ser. xvi, 39; Hist. Veg. xi, 131.—Walpers, Ann. iii, 396.

C. crassifolia, var. *morifolia*, Spach in Ann. Sci. Nat. 2 ser. xvi, 39; Hist. Veg. xi, 131.—Walpers, Ann. iii, 396.

C. crassifolia, var. *eucalyptifolia*, Spach in Ann. Sci. Nat. 2 ser. xvi, 40; Hist. Veg. xi, 131.—Walpers, Ann. iii, 396.

- O. Audibertiana*, Spach in Ann. Sci. Nat. 2 ser. xvi, 41; Hist. Pl. xi, 135.—Planchon in De Candolle, Prodr. xvii, 174.
O. Audibertiana, var. *ovata*, Spach in Ann. Sci. Nat. 2 ser. xvi, 41; Hist. Veg. xi, 135.
O. Audibertiana, var. *oblongata*, Spach in Ann. Sci. Nat. 2 ser. xvi, 41; Hist. Veg. xi, 135.
O. Lindheimeri, Engelmann in herb. A. Braun. (Koch, Dendrologie, ii, 434).
O. Berlandieri, Klotsch in Linnæa, xviii, 541.—Planchon in De Candolle, Prodr. xvii, 178.
O. Texana, Scheele in Linnæa, xx, 146; Ræmer, Texas, 446; Appx. 146.
O. occidentalis, var. *crassifolia*, Gray, Manual N. States, 2 ed. 395; 5 ed. 443.—Wood, Cl. Book, 634; Bot. & Fl. 299.
O. occidentalis, var. *pumila*, Gray, Manual N. States, 2 ed. 397; 5 ed. 443.—Chapman, Fl. S. States, 417.—Curtis in Rep. Geological Surv. N. Carolina, iii, 1860, 62.—Watson in King's Rep. v, 321.

SUGARBERRY. HACKBERRY.

Valley of the Saint Lawrence river west to eastern Dakota, south through the Atlantic region to bay Biscayne and cape Romano, Florida, and the valley of the Devil's river, Texas.

A large tree, 18 to 30 or, exceptionally, 36 to 39 meters (*Ridgway*) in height, with a trunk 0.60 to 1.50 meter in diameter; most common and reaching its greatest development in the Mississippi River basin; rich bottoms or dry hillsides; sometimes reduced to a low shrub (*O. pumila*), and varying greatly in the size, shape, and texture of the leaves (*O. Mississippiensis lævigata*, *integrifolia*, *crassifolia*, etc.); the extremes connected by innumerable intermediate forms, which, thus considered, make one polymorphous species of wide geographical range.

Wood heavy, rather soft, not strong, coarse-grained, compact, satiny, susceptible of a good polish; layers of annual growth clearly marked by several rows of large open ducts, containing many small groups of smaller ducts arranged in intermediate concentric rings; medullary rays numerous, thin; color, clear light yellow, the sap-wood lighter; specific gravity, 0.7287; ash, 1.09; largely used for fencing and occasionally in the manufacture of cheap furniture.

Var. *reticulata*.

- O. reticulata*, Torrey in Ann. Lyc. N. York, ii, 247.—Eaton, Manual, 6 ed. 86.—Rafinesque, New Fl. & Bot. i, 35.—Eaton & Wright, Bot. 186.—Nuttall, Sylva, i, 133, t. 39; 2 ed. i, 146, t. 39.—Browne, Trees of America, 518.—Planchon in Ann. Sci. Nat. 3 ser. x, 293; De Candolle, Prodr. xvii, 178.—Walpers, Ann. iii, 396.—Torrey & Gray in Pacific R. R. Rep. ii, 175.—Cooper in Smithsonian Rep. 1858, 260; Am. Nat. iii, 407.—Gray in Proc. Am. Acad. vii, 401.—Watson in Pl. Wheeler, 16.—Vasey, Cat. Forest Trees, 23.—Hall in Coulter's Bot. Gazette, ii, 91.—Rothrock in Wheeler's Rep. vi, 238.—Rusby in Bull. Torrey Bot. Club, ix, 54.

- O. Douglasii*, Planchon in Ann. Sci. Nat. 3 ser. x, 293; De Candolle, Prodr. xvii, 178.—Walpers, Ann. iii, 396.

- ?*O. occidentalis*, var. *pumila*, Watson in King's Rep. v, 321 [not Gray].

- O. brevipes*, Watson in Proc. Am. Acad. 3 ser. xiv, 297.—Rothrock in Wheeler's Rep. vi, 238.

HACKBERRY. PALO BLANCO.

Western Texas (Dallas, *Ravenel*) to the mountains of southern Arizona, and through the Rocky mountains to eastern Oregon; in the Tehachipi pass, California (*Pringle*).

A small tree, 12 to 15 meters in height, with a trunk rarely 0.60 meter in diameter; borders of streams, generally in high mountain cañons, or in the more arid regions reduced to a low shrub; well characterized by its small, thick, coriaceous leaves, slightly pubescent on the underside along the prominent reticulated veins, and by the light-colored, deeply-furrowed bark, but connected with the typical *O. occidentalis* by intermediate forms not rare in western Texas.

Wood not distinguishable in structure or color from that of the species; specific gravity, 0.7275; ash, 1.22.

229.—*Ficus aurea*, Nuttall,

- Sylva, ii, 4, t. 43; 2 ed. i, 154, t. 43.—Cooper in Smithsonian Rep. 1858, 265.—Chapman, Fl. S. States, 415.—Vasey, Cat. Forest Trees, 22.

- F. aurea*, var. *latifolia*, Nuttall, Sylva, ii, 4; 2 ed. i, 154.

Semi-tropical Florida, Indian river to the southern keys.

A large parasitic tree, germinating on the trunks and branches of other trees, and sending down to the ground long aerial roots, which gradually grow together, kill the inclosed tree, and form a trunk sometimes 0.90 to 1.20 meter in diameter.

Wood exceedingly light, soft, very weak, coarse-grained, compact, not durable; medullary rays thin, hardly distinguishable; color, light brown, the sap-wood lighter; specific gravity, 0.2616; ash, 5.03.

230.—*Ficus brevifolia*, Nuttall,

Sylva, ii, 3, t. 42; 2 ed. i, 153, t. 42.—Cooper in Smithsonian Rep. 1858, 265.—Chapman, Fl. S. States, 415.—Vasey, Cat. Forest Trees, 22.

Semi-tropical Florida, bay Biscayne to the southern keys (Key Largo, Pumpkin Key, *Curtiss*).

A tree sometimes 15 meters in height, with a trunk rarely exceeding 0.30 meter in diameter.

Wood light, soft, close-grained, compact, containing few large, open, scattered ducts and many groups of much smaller ducts arranged in concentric circles; medullary rays numerous, thin, conspicuous; color, light brown or yellow, the sap-wood lighter; specific gravity, 0.6398; ash, 4.36.

231.—*Ficus pedunculata*, Aiton,

Hort. Kew. iii, 450; 2 ed. v, 486.—Chapman, Fl. S. States, 415.—Grisebach, Fl. British West Indies, 151.

F. complicata, Humboldt, Bonpland & Kunth, Nov. Gen. & Spec. ii, 48.

Urostigma pedunculatum, Miquel in Hooker, London Jour. Bot. vi, 450.—Walpers, Ann. i, 677.

WILD FIG. INDIA-RUBBER TREE.

Semi-tropical Florida, bay Biscayne to the southern keys (Key Largo, Umbrella and Boca Chica Keys, etc. *Curtiss*); in the West Indies.

A tree sometimes 12 meters in height, with a trunk rarely exceeding 0.50 meter in diameter, or often shrubby and much branched from the ground; rare.

Wood light, soft, weak, close-grained, compact, containing many large, open, scattered ducts, with many groups of small ducts arranged in concentric circles; medullary rays numerous, obscure; color, light orange-brown, the sap-wood undistinguishable; specific gravity, 0.4739; ash, 4.92.

232.—*Morus rubra*, Linnæus,

Spec. 1 ed. 986.—Marshall, Arbustum, 93.—Wangenheim, Amer. 37, t. 15, f. 35.—Walter, Fl. Caroliniana, 241.—Aiton, Hort. Kew. iii, 343; 2 ed. v, 266.—Mönch, Meth. 343.—Lamarck, Dict. iv, 377.—Abbot, Insects Georgia, ii, t. 70.—Michaux, Fl. Bor.-Am. ii, 179.—Willdenow, Spec. iv, 369; Enum. 967; Berl. Baumz. 252.—Nouveau Duhamel, iv, 91, t. 23.—Persoon, Syn. ii, 558.—Desfontaines, Hist. Arb. ii, 416.—Michaux f. Hist. Arb. Am. iii, 232, t. 10; N. American Sylva, 3 ed. iii, 42, t. 116.—Pursh, Fl. Am. Sept. ii, 639.—Eaton, Manual, 105; 6 ed. 230.—Barton, Prodr. Fl. Philadelph. 89.—Nuttall, Genera, ii, 209.—Hayne, Dend. Fl. 155.—Elliott, Sk. ii, 574.—Sprengel, Syst. i, 492.—Torrey, Compend. Fl. N. States, 352; Nicotlet's Rep. 160; Fl. N. York, ii, 220; Emory's Rep. 412.—Rafinesque, Med. Bot. ii, 243; New Fl. & Bot. i, 43; Am. Manual Mulberry Trees, 13.—Beck, Bot. 316.—Dietrich, Syn. i, 551.—London, Arboretum, iii, 1359 & t.—Seringe, Descr. & Cult. du Mûr. 223, t. 20.—Eaton & Wright, Bot. 323.—Spach, Hist. Veg. xi, 48.—Browne, Trees of America, 457.—Emerson, Trees Massachusetts, 280; 2 ed. i, 314.—Darlington, Fl. Cestrica, 2 ed. 285.—Darby, Bot. S. States, 503.—Cooper in Smithsonian Rep. 1858, 254.—Chapman, Fl. S. States, 415.—Gray in Pacific R. R. Rep. xii², 47; Manual N. States, 5 ed. 444.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 71.—Lesquereux in Owen's 2d Rep. Arkansas, 386.—Wood, Cl. Book, 635; Bot. & Fl. 300.—Porcher, Resources S. Forests, 305.—Engelmann in Trans. Am. Phil. Soc. new ser. xii, 208.—Koch Dendrologie, ii, 447.—Young, Bot. Texas, 494.—Bureau in De Candolle, Prodr. xvii, 245.—Hayden in Warren's Rep. Nebraska & Dakota, 2 ed. 121.—Vasey, Cat. Forest Trees, 22.—Riley in Special Rep. U. S. Dept. Ag. No. 11, 34.—Ridgway in Proc. U. S. Nat. Mus. 1882, 73.—Burgess in Coulter's Bot. Gazette, vii, 95.

M. Canadensis, Lamarck, Dict. iv, 380.—Seringe, Descr. & Cult. du Mûr. 224.

M. scabra, Willdenow, Enum. 967; Berl. Baumz. 152.—Nuttall, Genera, ii, 209.—Rafinesque, Am. Manual Mulberry Trees, 29.—Hayne, Dend. Fl. 154.—Sprengel, Syst. i, 492.—Loddiges, Cat. 1836.

M. tomentosa, Rafinesque, Fl. Ludoviciana, 113; Am. Manual Mulberry Trees, 30.

M. reticulata, *M. Canadensis*, *M. parvifolia*, and *M. riparia*, Rafinesque, Am. Manual Mulberry Trees, 29-31.

M. rubra, var. *Canadensis*, Loudon, Arboretum, iii, 1360.

M. Missouriensis, Audibert, Cat. Jard. Tonnelle.

M. rubra, var. *tomentosa*, Bureau in De Candolle, Prodr. xvii, 246.

M. rubra, var. *incisa*, Bureau in De Candolle, Prodr. xvii, 247.

RED MULBERRY.

Western New England and Long Island, New York, west through southern Ontario and central Michigan to the Black hills of Dakota, eastern Nebraska and Kansas, south to bay Biscayne and cape Romano, Florida, and the valley of the Colorado river, Texas.

A large tree, 18 to 20 meters in height, with a trunk 0.90 to 1.20 meter or, exceptionally, 2.15 meters in diameter (*P. J. Berckmans*, Augusta, Georgia); generally in rich bottom lands; most common and reaching its greatest development in the basins of the lower Ohio and the Mississippi rivers.

Wood light, soft, not strong, rather tough, coarse-grained, compact, very durable in contact with the soil, satiny, susceptible of a good polish; layers of annual growth clearly marked by several rows of large open ducts; medullary rays numerous, thin; color, light orange-yellow, the sap-wood lighter; specific gravity, 0.5898; ash, 0.71; largely used in fencing, cooperage, for snaths, and at the south in ship- and boat-building.

The large dark purple fruit sweet and edible.

233.—*Morus microphylla*, Buckley,

Proc. Philadelphia Acad. 1862, 8.—Gray in Proc. Philadelphia Acad. 1862, 167.—Young, Bot. Texas, 494.

M. parvifolia, Engelmann in herb.—Gray, Hall's Pl. Texas, 21.—Vasey, Cat. Forest Trees, 22.—Riley in Special Rep. U. S. Dept. Ag. No. 11, 34.

MEXICAN MULBERRY.

Valley of the Colorado river, through western Texas to the valley of the Gila river, New Mexico; and southward into Mexico.

A small tree, sometimes 7 meters in height, with a trunk rarely 0.30 meter in diameter, or often reduced to a low shrub; most common and reaching its greatest development in the mountain cañons of southern New Mexico; in Texas generally on limestone formations.

Wood heavy, hard, close-grained, compact; layers of annual growth marked with several rows of small open ducts; medullary rays numerous, thin; color, orange or, rarely, dark brown, the sap-wood light yellow; specific gravity, 0.7715; ash, 0.68.

The small acid fruit hardly edible.

234.—*Maclura aurantiaca*, Nuttall,

Genera, ii, 234; Trans. Am. Phil. Soc. 2 ser. v, 169; Sylva, i, 126, t. 37, 38; 2 ed. i, 140, t. 37, 38.—James in Long's Exped. ii, 158.—Delile in Bull. Soc. Ag. Her. 1835 & t.—Eaton, Manual, 6 ed. 217.—Seringe in Mem. Soc. Ag. Lyon, 1835, 125 & t; Descr. & Cult. du Mûr. 232, t. 273.—Lambert, Pinus, 2 ed. ii, Appx. 4, t. 3.—London, Arboretum, iii, 1342, 1362, f. 1226–1228; Gard. Mag. xi, 312, f. 45–47.—Eaton & Wright, Bot. 311.—Spach, Hist. Veg. xi, 53.—Browne, Trees of America, 465.—Darby, Bot. S. States, 504.—Cooper in Smithsonian Rep. 1858, 254.—Miquel in Martius, Fl. Brasil. iv, 158.—Wood, Cl. Book, 635; Bot. & Fl. 299.—Porcher, Resources S. Forests, 101.—Koch, Dendrologie, ii, 437.—Bureau in De Candolle, Prodr. xvii, 227.—Dumen in Proc. California Acad. v, 398.—Vasey, Cat. Forest Trees, 22.—Guibourt, Hist. Drogues, 7 ed. ii, 325.—Riley in Special Rep. U. S. Dept. Ag. No. 11, 35.

Toxylon Maclura, Rafinesque, New Fl. & Bot. i, 43; Am. Manual Mulberry Trees, 13.

Toxylon pomiferum, Rafinesque in Am. Monthly Mag. and Crit. Rev. ii, 118.

Broussonetia tinctoria, Torrey in Ann. Lyc. N. York, ii, 246 [not Kunth].

OSAGE ORANGE. BOIS D'ARC.

Southwestern Arkansas, south of the valley of the Arkansas river, southeastern portions of the Indian territory, and southward in northern Texas to about latitude 32° 50' N. (Dallas, *Reverchon*, etc.).

A tree, sometimes 15 to 18 meters in height, with a trunk rarely exceeding 0.60 meter in diameter; rich bottom lands; most common and probably reaching its greatest development along the valley of the Red river in the Indian territory.

Wood heavy, exceedingly hard, very strong, flexible, close-grained, compact, very durable in contact with the ground, satiny, susceptible of a beautiful polish, containing numerous small open ducts, layers of annual growth clearly marked by broad bands of larger ducts; medullary rays thin, numerous, conspicuous; color, bright orange, turning brown with exposure, the sap-wood light yellow; specific gravity, 0.7736; ash, 0.68; largely used for fence posts, paving blocks, railway ties, wheel stock; extensively planted for hedges, especially in the western states.

PLATANACEÆ.

235.—*Platanus occidentalis*, Linnæus,

Spec. 1 ed. 999.—Du Roi, Harbk. ii, 134.—Marshall, Arbustum, 105.—Wangenheim, Amer. 31, t. 13, f. 31.—Walter, Fl. Caroliniana, 236.—Aiton, Hort. Kew. iii, 365; 2 ed. v, 305.—Mœnch, Meth. 358.—Abbot, Insects Georgia, ii, t. 55.—Michaux, Fl. Bor.-Am. ii, 163.—Lamarck, Dict. v, 438.—Nouveau Duhamel, ii, 6, t. 2.—Willdenow, Spec. iv, 474; Enum. 984; Berl. Baumz. 284.—Persoon, Syn. ii, 575.—Desfontaines Hist. Arb. ii, 545.—Schkuhr, Handb. iii, 274, t. 306.—Robin, Voyages, iii, 524.—Michaux f. Hist. Arb. Am. iii, 184, t. 3; N. American Sylva, 3 ed. ii, 48, t. 63.—Pursh, Fl. Am. Sept. ii, 635.—Barton, Prodr. Fl. Philadelph. 91; Compend. Fl. Philadelph. 176.—Eaton, Manual, 110; 6 ed. 267.—Nuttall, Genera, ii, 219.—Hayne, Dend. Fl. 171.—James in Long's Exped. i, 23.—Elliott, Sk. ii, 620.—Sprengel, Syst. iii, 865.—Watson, Dend. Brit. i, t. 100.—Torrey, Compend. Fl. N. States, 356; Fl. N. York, ii, 218; Bot. Mex. Boundary Survey, 205.—Audubon, Birds, t. 206.—London, Arboretum, iv, 2043, f. 1959 & t.—Eaton & Wright, Bot. 361.—Hooker, Fl. Bor.-Am. ii, 158.—Bigelow, Fl. Boston. 3 ed. 384.—Emerson, Trees Massachusetts, 227; 2 ed. i, 261 & t.—Scheele in Rœmer, Texas, 446.—Buckley in Am. Jour. Sci. 2 ser. xiii, 399.—Darlington, Fl. Cestrica, 3 ed. 282.—Darby, Bot. S. States, 509.—Agardh, Theor. & Syst. Pl. t. xiii, f. 1, 2.—Cooper in Smithsonian Rep. 1858, 254.—Hartig, Forst. 446, t. 54.—Chapman, Fl. S. States, 418.—Curtis in Rep. Geological Surv. N. Carolina, 76.—Lesquereux in Owen's 2d Rep. Arkansas, 386.—Wood, Cl. Book, 640; Bot. & Fl. 303.—Engelmann in Trans. Am. Phil. Soc. new ser. xii, 209.—A. De Candolle, Prodr. xvi², 159.—Gray, Manual N. States, 5 ed. 447; Hall's Fl. Texas, 21.—Koch, Dendrologie, ii, 468.—Schnizlein, Icon. t. 97, f. 1-24.—Young, Bot. Texas, 498.—Hayden in Warren's Rep. Nebraska & Dakota, 2 ed. 121.—Vasey, Cat. Forest Trees, 22.—Ridgway in Proc. U. S. Nat. Mus. 1882, 73.—Bell in Geological Rep. Canada, 1879-'80, 55^c.

P. lobata, Mœnch, Meth. 358.

P. hybridus, Brotero, Fl. Lus. ii, 487.

P. vulgaris, var. *angulosa*, Spach in Ann. Sci. Nat. 2 ser. xv, 293; Hist. Veg. xi, 79.

SYCAMORE. BUTTON WOOD. BUTTON-BALL TREE. WATER BEECH.

Southern Maine and southeastern New Hampshire to northern Vermont and the northern shores of lakes Ontario and Erie, west to eastern Nebraska and Kansas, south to northern Florida, central Alabama and Mississippi, and the valley of the Nueces river, Texas, extending southwest to the valley of the Devil's river.

The largest tree of the Atlantic forests, often 30 to 40 meters in height, with a trunk 2.40 to 4.20 meters in diameter; generally along streams and river bottoms, in rich, moist soil; very common and reaching its greatest development in the bottom lands of the Ohio and Mississippi rivers; the large specimens generally hollow.

Wood heavy, hard, not strong, very close-grained, compact, difficult to split and work; layers of annual growth clearly marked by broad bands of small ducts; the numerous medullary rays very conspicuous, as in that of all the North American species; color, brown tinged with red, the sap-wood lighter; specific gravity, 0.5678; ash, 0.46; largely used for tobacco boxes (its principal use), ox-yokes, butchers' blocks, and, rarely, in the manufacture of cheap furniture.

236.—*Platanus racemosa*, Nuttall;

Audubon, Birds, t. 362; Sylva, i, 47, t. 15; 2 ed. i, 63, t. 15.—Bentham, Pl. Hartweg. 336.—Newberry in Pacific R. R. Rep. vi, 33, 89, t. 11, f. 10.—Cooper in Smithsonian Rep. 1858, 260.—Torrey, Bot. Mex. Boundary Survey, 204; Ives' Rep. 27; Bot. Wilkes Exped. 457.—A. De Candolle, Prodr. xvi², 160.—Koch, Dendrologie, ii, 469.—Vasey, Cat. Forest Trees, 23.—Watson, Bot. California, ii, 66.

P. occidentalis, Hooker & Arnott, Bot. Beechey, 160, 380 [not Linnæus].

P. Californica, Bentham, Bot. Sulphur, 54.

P. Mexicana, Moricand, Pl. Rar. Amer. t. 13†.—Torrey in Sitgreaves' Rep. 172; Pacific R. R. Rep. vii, 20.

SYCAMORE. BUTTON WOOD.

California, valley of the Sacramento river, south through the interior valleys and coast ranges to the southern boundary of the state.

A large tree, 24 to 30 meters in height, with a trunk 0.90 to 1.20 meter in diameter; borders of streams, in rich soil.

Wood light, soft, not strong, very close-grained, compact, difficult to split; layers of annual growth clearly marked by narrow bands of small ducts; medullary rays numerous, conspicuous; color, light brown tinged with red, the sap-wood lighter; specific gravity, 0.4880; ash, 1.11.

237.—*Platanus Wrightii*, Watson,

Proc. Am. Acad. x, 349.—Vasey, Cat. Forest Trees, 23.—Rusby in Bull. Torrey Bot. Club, ix, 54.

P. Mexicana, Torrey in Emory's Rep. 151 [not Moricand].

P. racemosa, Watson, Fl. Wheeler, 16 [not Nuttall].—Rothrock in Wheeler's Rep. vi, 239.

SYCAMORE.

Valleys of southwestern New Mexico to the valley of the San Pedro river, Arizona; southward into Mexico.

A tree sometimes 15 to 18 meters in height, with a trunk 0.45 to 0.60 meter in diameter; banks of streams and high mountain cañons.

Wood light, soft, weak, very close-grained, compact; layers of annual growth clearly marked by several rows of open ducts; medullary rays numerous, thin, very conspicuous; color, light brown tinged with red, the sap-wood lighter; specific gravity, 0.4736; ash, 1.35.

JUGLANDACEÆ.

238.—*Juglans cinerea*, Linnæus,

Spec. 2 ed. 1415.—Jacquin, Icon. Rar. i, t. 193.—Wangenheim, Amer. 21, t. 9, f. 21.—Walter, Fl. Caroliniana, 235.—Aiton, Hort. Kew. iii, 361; 2 ed. v, 296.—Lamarck, Dict. iv, 503; Ill. iii, 365, t. 781, f. 7.—B. S. Barton, Coll. i, 22, 31; ii, 43.—Muhlenberg & Willdenow in Neue Schriften Gesell. Nat. Fr. Berlin, iii, 388.—Michaux, Fl. Bor.-Am. ii, 191.—Willdenow, Spec. iv, 456; Enum. 978; Berl. Baumz. 193.—Persoon, Syn. ii, 556.—Desfontaines, Hist. Arb. ii, 347.—Pursh, Fl. Am. Sept. ii, 636.—Barton, Prodr. Fl. Philadelph. 92.—Bigelow, Med. Bot. ii, 115, t. 32; Fl. Boston. 3 ed. 378.—Eaton, Manual, 108; 6 ed. 192.—Nuttall, Genera, ii, 220; Sylva, i, 41; 2 ed. i, 37.—Hayne, Dend. Fl. 163.—Elliott, Sk. ii, 622.—Sprengel, Syst. iii, 865.—Torrey, Compend. Fl. N. States, 357; Fl. N. York, ii, 180.—Rafinesque, Med. Bot. ii, 234.—Audubon, Birds, t. 142.—Beck, Bot. 335.—Spach, Hist. Veg. ii, 170.—Lindley, Fl. Med. 307.—London, Arboretum, iii, 1439, f. 1262.—Hooker, Fl. Bor.-Am. ii, 143.—Eaton & Wright, Bot. 287.—Emerson, Trees Massachusetts, 182; 2 ed. i, 207 & t.—Griffith, Med. Bot. 589.—Carson, Med. Bot. ii, 42, t. 86.—Parry in Owen's Rep. 618.—Darlington, Fl. Cestrica, 3 ed. 262.—Darby, Bot. S. States, 513.—Cooper in Smithsonian Rep. 1858, 254.—Chapman, Fl. S. States, 419.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 45.—Lesquereux in Owen's 2d Rep. Arkansas, 387.—Wood, Cl. Book, 640; Bot. & Fl. 304.—C. De Candolle in Ann. Sci. Nat. 4 ser. xviii, 16, t. 4, f. 45; Prodr. xvi³, 137.—Porcher, Resources S. Forests, 317.—Engelmann in Trans. Am. Phil. Soc. new ser. xii, 209.—Gray, Manual N. States, 5 ed. 447.—Koch, Dendrologie, i, 589.—Hayden in Warren's Rep. Nebraska & Dakota, 2 ed. 121.—Vasey, Cat. Forest Trees, 23.—Broadhead in Coulter's Bot. Gazette, iii, 60.—Bentley & Trimen, Med. Pl. iv, 247, t. 247.—Beal in Am. Nat. xv, 36, f. 6.—Sears in Bull. Essex Inst. xiii, 178.—Bell in Geological Rep. Canada, 1878-'80, 536.—Ridgway in Proc. U. S. Nat. Mus. 1882, 76.

J. oblonga, Miller, Dict. No. 3.—Du Roi, Harbk. i, 332.—Möench, Meth. 696.—Retzius, Obs. i, 10.

J. oblonga alba, Marshall, Arbustum, 67.

J. cathartica, Michaux f. Hist. Arb. Am. i, 165, t. 2; N. American Sylva, 3 ed. i, 109, t. 31.

Carya cathartica, Barton, Compend. Fl. Philadelph. ii, 178.

Wallia cinerea, Alefeld in Bonplandia, 1861, 334.

BUTTERNUT. WHITE WALNUT.

Southern New Brunswick, valley of the Saint Lawrence river, Ontario and southern Michigan to northern Minnesota (lake Pokegoma, *Garrison*) and central Iowa, south to Delaware and along the Alleghany mountains to northern Georgia, central Alabama and Mississippi, northern Arkansas, and southeastern Kansas.

A tree 18 to 24 or, exceptionally, 30 to 35 meters (*Ridgway*) in height, with a trunk 0.60 to 0.90 meter in diameter; rich woodlands; rare at the south; most common and reaching its greatest development in the Ohio River basin.

Wood light, soft, not strong, rather coarse-grained, compact, easily worked, satiny, susceptible of a beautiful polish, containing numerous regularly-distributed, large, open ducts; medullary rays distant, thin, obscure; color, bright light brown, turning dark with exposure, the sap-wood lighter; specific gravity, 0.4086; ash, 0.51; largely used for interior finish, cabinet work, etc.

The inner bark, especially that of the root, is employed medicinally as a mild cathartic (*Am. Jour. Pharm.* 1874, 169.—*U. S. Dispensatory*, 14 ed. 526.—*Nat. Dispensatory*, 2 ed. 794), and furnishes a yellow dye.

239.—*Juglans nigra*, Linnaeus,

Spec. 1 ed. 997.—Jacquin, Icon. Rar. i, t. 191.—Wangenheim, Amer. 20, t. 8, f. 20.—Walter, Fl. Caroliniana, 235.—Aiton, Hort. Kew, iii, 360; 2 ed. v, 296.—Mönch, Meth. 696.—Lamarck, Dict. iv, 502; Ill. iii, 365, t. 781, f. 6.—Abbot, Insects Georgia, i, t. 88.—Muhlenberg & Willdenow in Nene Schriften Gesell. Nat. Fr. Berlin, iii, 398.—Michaux, Fl. Bor.-Am. ii, 191.—Willdenow, Spec. iv, 456; Enum. 978; Berl. Baumz. 193.—Smith in Rees' Cycl. xx, No. 3.—Persoon, Syn. ii, 566.—Desfontaines, Hist. Arb. ii, 347.—Neuveau Duhamel, iv, 179, t. 48.—Michaux f. Hist. Arb. Am. i, 158, t. 1; N. American Sylva, 3 ed. i, 140, t. 30.—Pursh, Fl. Am. Sept. ii, 636.—Barton, Prodr. Fl. Philadelph. 92; Compend. Fl. Philadelph. ii, 177.—Eaton, Manual, 108; 6 ed. 192.—Nuttall, Genera, ii, 220; Sylva, i, 41; 2 ed. i, 57.—Hayne, Dend. Fl. 163.—Elliott, Sk. ii, 622.—Sprengel, Syst. iii, 865.—Torrey, Compend. Fl. N. States, 357; Fl. N. York, ii, 179.—Watson, Dend. Brit. ii, t. 158.—Audubon, Birds, t. 84, 156.—Rafinesque, Med. Bot. ii, 233.—Beck, Bot. 335.—Spach, Hist. Veg. ii, 168.—Loudon, Arboretum, iii, 1435, f. 1260 & t.—Eaton & Wright, Bot. 287.—Emerson, Trees Massachusetts, 185; 2 ed. i, 211 & t.—Griffith, Med. Bot. 589.—Parry in Owen's Rep. 618.—Darlington, Fl. Cestrica, 3 ed. 262.—Darby, Bot. 8. States, 513.—Cooper in Smithsonian Rep. 1858, 254.—Chapman, Fl. S. States, 419.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 45.—Lesquereux in Owen's 2d Rep. Arkansas, 387.—Wood, Cl. Book, 640; Bot. & Fl. 304.—C. De Candolle in Ann. Sci. Nat. 4 ser. xviii, 34, t. 1, f. 1, 8-10; Prodr. xvi², 137.—Engelmann in Trans. Am. Phil. Soc. new ser. xii, 209.—Porcher, Resources S. Forests, 318.—Gray, Manual N. States, 5 ed. 447.—Koch, Dendrologie, i, 587.—Schnizlein, Icon. t. 244, f. 1, 8, 12, 13.—Young, Bot. Texas, 500.—Hayden in Warren's Rep. Nebraska & Dakota, 2 ed. 121.—Vasey, Cat. Forest Trees, 23.—Guibourt, Hist. Drogues, 7 ed. ii, 302.—Beal in Am. Nat. xv, 36, f. 5.—Sears in Bull. Essex Inst. xiii, 178.—Bell in Geological Rep. Canada, 1879-'80, 53.—Ridgway in Proc. U. S. Nat. Mus. 1882, 76.—Nicholson in London Gard. Chronicle, 1882, 780.—Watson in Proc. Am. Acad. xviii, 155.

J. nigra oblonga, Marshall, Arbustum, 67.

Wallia nigra, Alefeld in Bonplandia, 1861, 334.

BLACK WALNUT.

Western Massachusetts, west along the southern shores of lake Erie through southern Michigan to southern Minnesota, eastern Nebraska, and eastern Kansas, south to the Chattahoochee region of northern Florida, central Alabama and Mississippi, and the valley of the San Antonio river, Texas.

A large tree, often 30 to 45 meters in height, with a trunk 1.80 to 3 meters in diameter; rich bottom lands and hillsides; most common and reaching its greatest development on the western slopes of the southern Alleghany mountains and in the rich bottoms of southwestern Arkansas and the Indian territory; less common east of the Alleghany mountains, and now everywhere scarce.

Wood heavy, hard, strong, rather coarse-grained, liable to check if not carefully seasoned, easily worked, susceptible of a beautiful polish, durable in contact with the soil, containing numerous large, regularly-distributed, open ducts; medullary rays numerous, thin, not conspicuous; color, rich dark brown, the thin sap-wood much lighter; specific gravity, 0.6115; ash, 0.79; more generally used in cabinet-making, interior finish, and for gun stocks than that of any other North American tree.

240.—*Juglans rupestris*, Engelmann;

Sitgreaves' Rep. 171, t. 15.—Torrey, Bot. Mex. Boundary Survey, 205; Ives' Rep. 27.—Cooper in Smithsonian Rep. 1858, 260.—C. De Candolle in Ann. Sci. Nat. 4 ser. xviii, 28, t. 2, f. 11; Prodr. xvi², 138.—Vasey, Cat. Forest Trees, 24.—Watson, Bot. California, ii, 93; Proc. Am. Acad. xviii, 155.—Rusby in Bull. Torrey Bot. Club. ix, 54.

J. rupestris, var. *major*, Torrey in Sitgreaves' Rep. 171, t. 16; Bot. Mex. Boundary Survey, 205; Pacific R. R. Rep. vii, 20.—C. De Candolle, Prodr. xvi², 138.—Hemsley, Bot. Am.-Cent. iii, 164.

J. Californica, Watson in Proc. Am. Acad. x, 349; Bot. California, ii, 93.—Vasey, Cat. Forest Trees, 24.—Rothrock in Wheeler's Rep. vi, 249.

WALNUT.

Valley of the Colorado river (near Austin), west through western Texas, southern New Mexico, and Arizona from 5,000 to 7,000 feet elevation, and in the California Coast ranges from the San Bernardino mountains to the neighborhood of San Francisco bay and the valley of the Sacramento river.

A tree rarely 15 to 22 meters in height, with a trunk 0.30 to 0.90 meter in diameter, reaching its greatest development in the neighborhood of San Francisco bay; in Texas generally reduced to a low, much-branched shrub; borders of streams and mountain cañons, in rich soil.

Wood heavy, hard, not strong, coarse-grained, checking in drying, susceptible of a good polish, containing numerous regularly-distributed, large, open ducts; medullary rays distant, thin, obscure; color, rich dark brown, the sap-wood lighter; specific gravity, 0.6554; ash, 1.01.

The small nuts sweet and edible.

241.—*Carya olivæformis*, Nuttall,

Genera, ii, 221.—Sprengel, Syst. ii, 849.—Eaton, Manual, 6 ed. 83.—Spach, Hist. Veg. ii, 173.—Penn. Cycl. vi, 331.—London, Arboretum, iii, 1441, f. 1263.—Eaton & Wright, Bot. 183.—Scheele in Roemer, Texas, 447.—Belg. Hort. vi, 223, t. 45, f. 2.—Torrey, Bot. Mex. Boundary Survey 205.—Cooper in Smithsonian Rep. 1858, 255.—Chapman, Fl. S. States, 418.—Lesquereux in Owen's 2d Rep. Arkansas, 387.—Wood, Cl. Book, 641; Bot. & Fl. 304.—C. De Candolle in Ann. Sci. Nat. 4 ser. xviii, 36, t. 1, f. 3, t. 5, f. 59; Prodr. xvi², 144.—Porcher, Resources S. Forests, 333.—Gray, Manual N. States, 5 ed. 448.—Young, Bot. Texas, 499.—Vasey, Cat. Forest Trees, 24.—Broadhead in Coulter's Bot. Gazette, iii, 60.—Ridgway in Proc. U. S. Nat. Mus. 1882, 77.—Hemsley, Bot. Am.-Cent. iii, 163.—Watson in Proc. Am. Acad. xviii, 155.

Juglans Pecan, Marshall, Arbustum, 69.—Walter, Fl. Caroliniana, 236.—Muhlenberg & Willdenow in Neue Schriften Gesell. Nat. Fr. Berlin, iii, 392.

Juglans Illinoensis, Wangenheim, Amer. 54, t. 18, f. 43.

Juglans angustifolia, Aiton, Hort. Kew. iii, 361; 2 ed. v, 296.

Juglans rubra, Gærtner, Fruct. ii, 51, t. 89, f. 1.—Lamarck, Ill. iii, 365, t. 781, f. 4.

Juglans cylindrica, Lamarck, Dict. iv, 505; Ill. iii, 365, t. 781, f. 5.—Nouveau Duhamel, iv, 179.

Juglans olivæformis, Michaux, Fl. Bor.-Am. ii, 192.—Willdenow, Spec. iv, 457; Enum. 979; Berl. Baumz. 194.—Persoon, Syn. ii, 566.—Desfontaines, Hist. Arb. ii, 348.—Michaux f. Hist. Arb. Am. i, 175, t. 3; N. American Sylva, 3 ed. i, 114, t. 32.—Muhlenberg, Cat. 88.—Aiton, Hort. Kew. 2 ed. v, 296.—Pursh, Fl. Am. Sept. ii, 636.—Hayne, Dend. Fl. 163.—Regel, Gartenflora, xviii, 89.

C. angustifolia, Nuttall, Sylva, i, 41; 2 ed. i, 57.

?*C. tetraptera*, Liebmann in Dansk. Vidensk. Selsk. Forhand. 1850, 80.

Hickorea species, LeConte in Proc. Philadelphia Acad. vi, 402.

C. Illinoensis, Koch, Dendrologie, i, 593.

PECAN. ILLINOIS NUT.

Near Davenport, Iowa (*C. C. Parry*), southern Illinois, and Indiana, northwestern Kentucky, south and southwest through Missouri and Arkansas to eastern Kansas, the Indian territory, and through western Louisiana and Texas to the valley of the Concho river.

A tree 30 to 52 meters in height, with a trunk 0.90 to 1.80 meter in diameter; borders of streams in low, rich soil; very common and reaching its greatest development in the bottom lands of Arkansas and the Indian territory; the largest species of the genus and the largest and most important tree of western Texas.

Wood heavy, hard, not strong, brittle, close-grained, compact; layers of annual growth marked by one or two rows of large open ducts; medullary rays numerous, thin; color, light brown tinged with red; the sap-wood lighter brown; specific gravity, 0.7180; ash, 1.13; less valuable than the wood of the other species and hardly used except for fuel.

The sweet, edible nuts are collected in great quantities, affording an important article of commerce.

242.—*Carya alba*, Nuttall,

Genera, ii, 221.—Elliott, Sk. ii, 624.—Watson, Dend. Brit. ii, t. 148.—Sprengel, Syst. ii, 849.—Torrey, Compend. Fl. N. States, 357; Fl. N. York, 181.—Beck, Bot. 336.—Eaton, Manual, 6 ed. 83.—Spach, Hist. Veg. ii, 174.—Penn. Cycl. vi, 332.—London, Arboretum, iii, 1446, f. 1269 & t.—Eaton & Wright, Bot. 183.—Hooker, Fl. Bor.-Am. ii, 143.—Emerson, Trees Massachusetts, 191; 2 ed. i, 217 & t.—Darlington, Fl. Cestrica, 3 ed. 263.—Darby, Bot. S. States, 513.—Belg. Hort. vi, 223, t. 48, f. 8.—Cooper in Smithsonian Rep. 1858, 255.—Chapman, Fl. S. States, 418.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 43.—Lesquereux in Owen's 2d Rep. Arkansas, 387.—Wood, Cl. Book, 641; Bot. & Fl. 304.—C. De Candolle in Ann. Sci. Nat. 4 ser. xviii, 36, t. 2, f. 13, 14, 18, t. 3, f. 24, t. 4, f. 44, 46; Prodr. xvi², 142.—Gray, Manual N. States, 5 ed. 448.—Young, Bot. Texas, 499.—Vasey, Cat. Forest Trees, 24.—Aldrich in Am. Nat. xv, 227.—Sears in Bull. Essex Inst. xiii, 179.—Ridgway in Proc. U. S. Nat. Mus. 1882, 72.—Bell in Geological Rep. Canada, 1879-'80, 55c.

Juglans ovata, Miller, Dict.

Juglans alba ovata, Marshall, Arbustum, 69.

Juglans ovalis, Wangenheim, Amer. 24, t. 10, f. 23.

Juglans compressa, Gærtner, Fruct. ii, 50, t. 89, f. 1.—Muhlenberg & Willdenow in Neue Schriften Gesell. Nat. Fr. Berlin, iii, 300.—Willdenow, Spec. iv, 458; Enum. 979; Berl. Baumz. 195.—Persoon, Syn. ii, 566.—Desfontaines, Hist. Arb. ii, 347.—Aiton, Hort. Kew. 2 ed. v, 297.—Hayne, Dend. Fl. 164.—Lamarck, Ill. iii, 365, t. 781, f. 3.

?*Juglans exaltata*, Bartram, Travels, 2 ed. 38.

Juglans squamosa, Lamarck, Dict. iv, 504.—Desfontaines, Hist. Arb. ii, 348.—Michaux f. Hist. Arb. Am. i, 190, t. 7; N. American Sylva, 3 ed. i, 123, t. 36.—Barton, Prodr. Fl. Philadelph. 92; Compend. Fl. Philadelph. ii, 179.—Bigelow, Fl. Boston. 3 ed. 380.

Juglans alba, Michaux, Fl. Bor. Am. ii 193 [not Linnaeus].—Pursh, Fl. Am. Sept. ii, 637.—Eaton, Manual, 108.

O. microcarpa, Nuttall, Genera, ii, 221; Sylva, i, 38, t. 13; 2 ed. i, 55, t. 13.—Sprengel, Syst. ii, 849.—Penn. Cycl. vi, 332.—London, Arboretum, iii, 1451.—Darlington, Fl. Cestrica, 3 ed. 264.—Cooper in Smithsonian Rep. 1858, 255.—Chapman, Fl. S. States, 419.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 44.—Wood, Cl. Book, 642; Bot. & Fl. 304.—C. De Candolle, Prodr. xvi², 143.—Gray, Manual N. States, 5 ed. 448.—Koch, Dendrologie, i, 596.—Young, Bot. Texas, 499.—Vasey, Cat. Forest Trees, 24.—Ridgway in Proc. U. S. Nat. Mus. 1882, 77.

SHELL-BARK HICKORY. SHAG-BARK HICKORY.

Valley of the Saint Lawrence river, along the northern shores of lakes Ontario and Erie to southern Michigan and southeastern Minnesota, south to the Chattahoochee region of western Florida, central Alabama and Mississippi, and west to eastern Kansas, the Indian territory, and eastern Texas.

A large tree of the first economic value, 24 to 30 or, exceptionally, 39 to 45 meters in height (*Ridgway*), with a trunk 0.90 to 1.20 meter in diameter; rich hillsides and sandy ridges; common and reaching its greatest development west of the Alleghany mountains; varying greatly in the size and shape of the fruit. A form with small, thin-shelled nuts (*O. microcarpa*, Nuttall l. c.) is not rare from Delaware southward, and in Michigan.

Wood heavy, very hard and strong, tough, close-grained, compact, flexible; layers of annual growth clearly marked with one to three rows of large open ducts; medullary rays numerous, thin; color, brown, the thin and more valuable sap-wood nearly white; specific gravity, 0.8372; ash, 0.73; largely used in the manufacture of agricultural implements, carriages, ax handles, baskets, etc.

The sweet and edible nuts afford an important article of commerce.

243.—*Carya sulcata*, Nuttall,

Genera, ii, 221.—Elliott, Sk. ii, 624.—Sprengel, Syst. ii, 849.—Torrey, Compend. Fl. N. States, 357.—Beck, Bot. 336.—Eaton, Manual, 6 ed. 83.—Spach, Hist. Veg. ii, 174.—Penn. Cycl. vi, 332.—London, Arboretum, iii, 1448, f. 1271.—Eaton & Wright, Bot. 183.—Darby, Bot. S. States, 513.—Cooper in Smithsonian Rep. 1858, 255.—Chapman, Fl. S. States, 418.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 43.—Lesquereux in Owen's 2d Rep. Arkansas, 387.—Wood, Cl. Book, 641; Bot. & Fl. 304.—C. De Candolle in Ann. Sci. Nat. 4 ser. xviii, 36, t. 5, f. 51, 52; Prodr. xvi², 143.—Gray, Manual N. States, 5 ed. 449.—Young, Bot. Texas, 499.—Vasey, Cat. Forest Trees, 24.—Ridgway in Proc. U. S. Nat. Mus. 1882, 78.

Juglans sulcata, Willdenow, Berl. Baumz. 1 ed. 154, t. 7; Spec. iv, 457.—Muhlenberg & Willdenow in Neue Schriften Gesell. Nat. Fr. Berlin, iii, 391.—Persoon, Syn. ii, 566.—Desfontaines, Hist. Arb. ii, 348.—Pursh, Fl. Am. Sept. ii, 637.

Juglans mucronata, Michaux, Fl. Bor.-Am. ii, 192.

Juglans laciniosa, Michaux f. Hist. Arb. Am. i, 199, t. 8; N. American Sylva, 3 ed. i, 128, t. 37.—Barton, Prodr. Fl. Philadelph. 92.—Poiret, Suppl. iv, 112.—Audubon, Birds, t. 101.

C. cordiformis, Koch, Dendrologie, i, 597.

BIG SHELL-BARK. BOTTOM SHELL-BARK.

Chester county, Pennsylvania, west to southern Indiana and Illinois, eastern Kansas, and the Indian territory.

A tree 24 to 30 or, exceptionally, 37 (*Ridgway*) meters in height, with a trunk 0.60 to 1.20 meter in diameter; bottom lands, in low, rich soil; rare and local; most common and reaching its greatest development along the streams of southern Arkansas and the Indian territory.

Wood heavy, very hard, strong and tough, very close-grained, compact, flexible; layers of annual growth marked by one or two rows of large open ducts; medullary rays numerous, obscure; color, dark brown, the sap-wood nearly white; specific gravity, 0.8108; ash, 0.90; used for the same purposes as that of the shell-bark hickory.

The large nuts sweet and edible.

244.—*Carya tomentosa*, Nuttall,

Genera, ii, 221.—Barton, Compend. Fl. Philadelph. ii, 179.—Elliott, Sk. ii, 625.—Sprengel, Syst. ii, 849.—Torrey, Compend. Fl. N. States, 357; Fl. N. York, ii, 182.—Beck, Bot. 336.—Eaton, Manual, 6 ed. 83.—Spach, Hist. Veg. ii, 176.—Penn. Cycl. vi, 332.—London, Arboretum, iii, 1444, f. 1267.—Eaton & Wright, Bot. 183.—Emerson, Trees Massachusetts, 194, t. 13; 2 ed. i, 222 & t.—Darlington, Fl. Cestrica, 3 ed. 263.—Darby, Bot. S. States, 513.—Cooper in Smithsonian Rep. 1858, 255.—Chapman, Fl. S. States, 419.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 43.—Lesquereux in Owen's 2d Rep. Arkansas, 387.—Wood, Cl. Book, 641; Bot. & Fl. 304.—C. De Candolle in Ann. Sci. Nat. 4 ser. xviii, 36; Prodr. xvi², 143.—Gray, Manual N. States, 5 ed. 449.—Young, Bot. Texas, 499.—Vasey, Cat. Forest Trees, 24.—Ridgway in Proc. U. S. Nat. Mus. 1882, 78.

Juglans alba, Linnaeus, Spec. 1 ed. 997.—Du Roi, Harbk i, 333.—Kalm in Act. Holm. 1769, 117.—Wangenheim, Amer. 23, t. 10, f. 2.—Walter, Fl. Caroliniana, 235.—Aiton, Hort. Kew. iii, 360; 2 ed. v, 296.—Gærtner, Fruct. ii, 50, t. 89, f. 1.—Mönch, Meth. 696.—Abbot, Insects Georgia, i, t. 29.—Lamarck, Dict. iv, 503; Ill. iii, 364, t. 781, f. 2.—Muhlenberg & Willdenow in Neue Schriften Gesell. Nat. Fr. Berlin, iii, 369.—Smith in Rees' Cycl. xx, No. 2.—Willdenow, Spec. iv, 457; Berl. Baumz. 154.—Desfontaines, Hist. Arb. ii, 347.—Bigelow, Fl. Boston. 3 ed. 379.

Juglans tomentosa, Lamarek, Dict. iv, 504.—Michaux, Fl. Bor.-Am. ii, 192.—Michaux f. Hist. Arb. Am. i, 184, t. 6; N. American Sylva, 3 ed. i, 120, t. 35.—Pursh, Fl. Am. Sept. ii, 637.—Barton, Prodr. Fl. Philadelph. 92.

O. tomentosa, var. *maxima*, Nuttall, Genera, ii, 221; Sylva, i, 40; 2 ed. i, 56.—Sweet, Hort. Brit. ed. 1830.—Beck, Bot. 336.—London, Arboretum, iii, 1445.—C. De Candolle, Prodr. xvi³, 143.

O. alba, Koch, Dendrologie, i, 596 [not Nuttall].

MOCKEE NUT. BLACK HICKORY. BULL NUT. BIG-BUD HICKORY. WHITE-HEART HICKORY. KING NUT.

Valley of the Saint Lawrence river, northern shores of lakes Ontario and Erie to eastern Nebraska, eastern Kansas, and the Indian territory, south to cape Canaveral and Tampa bay, Florida, and the valley of the Brazos river, Texas.

A tree 24 to 30 or, exceptionally, 33 (*Ridgway*) meters in height, with a trunk 0.90 to 1.20 meter in diameter; generally on rich upland hillsides—less commonly in low river bottom lands; very common in the Gulf states, and throughout the south the most widely-distributed species of the genus.

Wood heavy, very hard, strong, tough, very close-grained, checking in drying, flexible, containing few large, regularly-distributed, open ducts; medullary rays numerous, thin, obscure; color, rich dark brown, the thick sap-wood nearly white; specific gravity, 0.8216; ash, 1.06; used for the same purposes as that of the shell-bark hickory.

245.—*Carya porcina*, Nuttall,

Genera, ii, 222.—Barton, Compend. Fl. Philadelph. ii, 180.—Elliott, Sk. ii, 627.—Watson, Dend. Brit. ii, t. 167.—Sprengel, Syst. ii, 849.—Torrey, Compend. Fl. N. States, 358.—Beck, Bot. 336.—Eaton, Manual, 6 ed. 83.—Spach, Hist. Veg. ii, 178.—Penn. Cyl. vi, 332.—Darlington, Fl. Cestrica, 2 ed. 546.—London, Arboretum, iii, 1449, f. 1272-1274.—Eaton & Wright, Bot. 183.—Spach, Hist. Veg. ii, 178.—Emerson, Trees Massachusetts, 197, t. 14; 2 ed. i, 224 & t.—Wood, Bot. & Fl. 304.—C. De Candolle in Ann. Sci. Nat. 4 ser. xviii, 36, t. 1, f. 5, t. 5, f. 54; Prodr. xvi³, 143.—Porcher, Resources S. Forests, 332.—Gray, Manual N. States, 5 ed. 449; Hall's Pl. Texas, 21.—Vasey, Cat. Forest Trees, 24.—Ridgway in Proc. U. S. Nat. Mus. 1882, 78.

Juglans glabra, Miller, Dict. No. 5.—Wangenheim, Amer. 25, t. 10, f. 24.—Muhlenberg & Willdenow in Neue Schriften Gesell. Nat. Fr. Berlin, iii, 391.—Willdenow, Spec. iv, 458; Berl. Baumz. 196.—Persoon, Syn. ii, 566.—Aiton, Hort. Kew. 2 ed. v, 297.—Eaton, Manual, 108.—Hayne, Dend. Fl. 164.

Juglans alba acuminata, Marshall, Arbustum, 68.

Juglans obcordata, Lamarek Dict. iv, 504.—Muhlenberg & Willdenow in Neue Schriften Gesell. Nat. Fr. Berlin, iii, 391.—Willdenow, Spec. iv, 458.—Persoon, Syn. 566.

Juglans porcina, Michaux f. Hist. Arb. Am. i, 206, t. 9; N. American Sylva, 3 ed. i, 132, t. 38.—Pursh, Fl. Am. Sept. ii, 638.—Barton, Prodr. Fl. Philadelph. 92.—Audubon, Birds, t. 91.

Juglans pyriformis, Muhlenberg, Cat. 92.

Juglans porcina, var. *obcordata*, Pursh, Fl. Am. Sept. ii, 638.—Barton, Compend. Fl. Philadelph. ii, 180.—Watson, Dend. Brit. ii, 167.

Juglans porcina, var. *pisiformis*, Pursh, Fl. Am. Sept. ii, 638.—Barton, Compend. Fl. Philadelph. ii, 180.

C. glabra, Torrey, Fl. N. York, ii, 182, t. 101.—Gray, Manual N. States, 1 ed. 412.—Darlington, Fl. Cestrica, 3 ed. 264.—Cooper in Smithsonian Rep. 1858, 255.—Chapman, Fl. S. States, 419.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 44.—Lesquereux in Owen's 2d Rep. Arkansas, 387.—Koch, Dendrologie, i, 594.—Young, Bot. Texas, 499.

C. amara, var. *porcina*, Darby, Bot. S. States, 513.

PIG NUT. BROWN HICKORY. BLACK HICKORY. SWITCH-BUD HICKORY.

Southern Maine to southern Ontario, southern Michigan and Minnesota, eastern Nebraska, eastern Kansas, and the Indian territory, south to cape Canaveral and Pease creek, Florida, and the valley of the Nueces river, Texas.

A tree 24 to 30 or, exceptionally, 40 (*Ridgway*) meters in height, with a trunk 0.90 to 1.50 meter in diameter; dry hills and uplands; common.

Wood heavy, hard, very strong and tough, flexible, close-grained, checking in drying, containing many large open ducts; color, dark or light brown, the thick sap-wood lighter, often nearly white; specific gravity, 0.8217; ash, 0.99; used for the same purposes as that of the shell-bark hickory.

246.—*Carya amara*, Nuttall,

Genera, ii, 222.—Barton, Compend. Fl. Philadelph. ii, 180.—Elliott, Sk. ii, 626.—Sprengel, Syst. ii, 849.—Torrey, Compend. Fl. N. States, 358; Fl. N. York, ii, 183.—Beck, Bot. 336.—Spach, Hist. Veg. ii, 177.—Penn. Cycl. vi, 332.—London, Arboretum, iii, 1443, f. 1264.—Hooker, Fl. Bor.-Am. ii, 144.—Emerson, Trees Massachusetts, 199, t. 15; 2 ed. i, 226 & t.—Darlington, Fl. Cestrica, 3 ed. 264.—Darby, Bot. S. States, 513.—Cooper in Smithsonian Rep. 1858, 255.—Chapman, Fl. S. States, 419.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 44.—Lesquereux in Owen's 2d Rep. Arkansas, 387.—Wood, Cl. Book, 641; Bot. & Fl. 304.—C. De Candolle in Ann. Sci. Nat. 4 ser. xviii, 36, t. 1, f. 2, t. 5, f. 53-55; Prodr. xvi², 144.—Gray, Manual N. States, 5 ed. 449; Hall's Fl. Texas, 21.—Koch, Dendrologie, i, 592.—Young, Bot. Texas, 500.—Vasey, Cat. Forest Trees, 24.—Sears in Bull. Essex Inst. xiii, 178.—Bell in Geological Rep. Canada, 1879-'80, 52.—Ridgway in Proc. U. S. Nat. Mus. 1882, 77.

Juglans alba minima, Marshall, Arbustum, 68.

Juglans cordiformis, Wangenheim, Amer. 25, t. 10, f. 25.

Juglans angustifolia, Lamarck, Dict. iv, 504 [not Aiton].

Juglans amara, Michaux f. Hist. Arb. Am. i, 177, t. 4; 3 ed. i, 116, t. 33.—Pursh, Fl. Am. Sept. ii, 638.

Hickorius amara, Rafinesque, Fl. Ludoviciana, 109.

BITTER NUT. SWAMP HICKORY.

Southern Maine to the valley of the Saint Lawrence river, west through Ontario, central Michigan and Minnesota to eastern Nebraska, eastern Kansas, and the Indian territory, south to the Chattahoochee region of western Florida and the valley of the Trinity river, Texas.

A tree 18 to 24 meters in height, with a trunk 0.60 to 0.90 meter in diameter; borders of streams and swamps, in low ground, or often on dry, rich uplands.

Wood heavy, very hard, strong, tough, close-grained, checking in drying; layers of annual growth marked by several rows of large open ducts; medullary rays numerous, obscure; color, dark brown, the thick sap-wood light brown, or often nearly white; specific gravity, 0.7552; ash, 1.03; largely used for hoops, ox-yokes, etc.

247.—*Carya myristicæformis*, Nuttall,

Genera, ii, 222.—Elliott, Sk. ii, 626.—Sprengel, Syst. ii, 849.—Eaton, Manual, 6 ed. 83.—Spach, Hist. Veg. ii, 179.—Penn. Cycl. v, 332.—London, Arboretum, iii, 1451, f. 1275.—Eaton & Wright, Bot. 1833.—Chapman, Fl. S. States, 419.—C. De Candolle in Ann. Sci. Nat. 4 ser. xviii, 36, t. 6, f. 58; Prodr. xvi², 145.—Koch, Dendrologie, i, 595.—Young, Bot. Texas, 500.—Vasey, Cat. Forest Trees, 24.—Ravenel in Bull. Torrey Bot. Club. vi, 81.

Juglans myristicæformis, Michaux f. Hist. Arb. Am. i, 211, t. 10; N. American Sylva, 3 ed. i, 135, t. 39.—Pursh, Fl. Am. Sept. ii, 638.—Poiret, Suppl. iv, 112.—Rafinesque, Fl. Ludoviciana, 161.

C. amara, var. *myristicæformis*, Cooper in Smithsonian Rep. 1858, 255.

NUTMEG HICKORY.

South Carolina, "Goose creek" (*Michaux*), "Berkeley district" (*Ravenel*); Arkansas, valley of the Arkansas river (Pine Bluff, *Letterman*), south to the Red River valley.

A tree 24 to 30 meters in height, with a trunk 0.60 to 0.90 meter in diameter; sandy ridges along the borders of streams and swamps; rare and very local in South Carolina; more common and reaching its greatest development in southern Arkansas.

Wood heavy, hard, very strong and tough, close-grained, compact, containing numerous small open ducts, layers of annual growth marked by one or two rows of larger ducts; medullary rays numerous, thin, not conspicuous; color, light brown, the sap-wood lighter; specific gravity, 0.8016; ash, 1.06.

248.—*Carya aquatica*, Nuttall,

Genera, ii, 222.—Elliott, Sk. ii, 627.—Sprengel, Syst. ii, 849.—Eaton, Manual, 6 ed. 83.—Spach, Hist. Veg. ii, 179.—Penn. Cycl. vi, 332.—London, Arboretum, iii, 1444, f. 1265, 1266.—Eaton & Wright, Bot. 1833.—Scheele in Ræmer, Texas, 447.—Darby, Bot. S. States, 514.—Chapman, Fl. S. States, 419.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 44.—Lesquereux in Owen's 2d Rep. Arkansas, 387.—Wood, Cl. Book, 641; Bot. & Fl. 304.—C. De Candolle in Ann. Sci. Nat. 4 ser. xviii, 36, t. 1, f. 4, t. 5, f. 56, 57; Prodr. xvi², 144.—Koch, Dendrologie, i, 593.—Young, Bot. Texas, 500.—Vasey, Cat. Forest Trees, 24.

Juglans aquatica, Michaux f. Hist. Arb. Am. i, 182, t. 5; N. American Sylva, 3 ed. i, 119, t. 34.—Pursh, Fl. Am. Sept. ii, 638.—Poiret, Suppl. iv, 112.

Hicorius integrifolia, Rafinesque, Fl. Ludoviciana, 109.

C. integrifolia, Sprengel, Syst. ii, 849.—London, Arboretum, iii, 1451.

WATER HICKORY. SWAMP HICKORY. BITTER PECAN.

North Carolina, in the lower districts, south to cape Malabar and the Caloosa river, Florida (in Florida not detected within 8 or 10 miles of the coast), through the Gulf states to western Louisiana, northeastern Arkansas, and the valley of the Brazos river, Texas.

A tree 18 to 21 meters in height, with a trunk 0.60 to 0.90 meter in diameter, or generally much smaller; low river swamps; most common and reaching its greatest development in the bottom lands of the lower Mississippi and Yazoo rivers.

Wood heavy, soft, strong, rather brittle, very close-grained, compact, containing few scattered, open ducts; layers of annual growth less clearly marked than in the other species of the genus; medullary rays numerous, thin; color, dark brown, the sap-wood light, often nearly white; specific gravity, 0.7407; ash, 1.27; used for fencing, fuel, etc.

MYRICACEÆ.

249.—*Myrica cerifera*, Linnaeus,

Spec. 1 ed. 1024.—Kalm, Travels, English ed. i, 92.—Marshall, Arbustum, 94.—Lamarck, Dict. ii, 592; Ill. iii, 402, t. 809, f. 1.—Gärtner, Fruct. i, 190, t. 39, f. 7.—Walter, Fl. Caroliniana, 242.—Aiton, Hort. Kew. iii, 396; 2 ed. v, 379.—Mönch, Meth. 362.—B. S. Barton, Coll. ii, 4.—Nouveau Duhamel, ii, 190.—Schkuhr, Handb. iii, 465, t. 322.—Michaux, Fl. Bor.-Am. ii, 227.—Willdenow, Spec. iv, 745; Enum. 1011; Berl. Baumz. 254.—Persoon, Syn. ii, 614.—Desfontaines, Hist. Arb. ii, 472.—Titford, Hort. Bot. Am. 100.—Pursh, Fl. Am. Sept. ii, 620.—Nuttall, Genera, ii, 235; Trans. Am. Phil. Soc. 2 ser. v, 167.—Bigelow, Med. Bot. iii, 32, t. 43; Fl. Boston. 3 ed. 394.—Hayne, Dend. Fl. 197.—Elliott, Sk. ii, 678.—Sprengel, Syst. i, 493.—Torrey, Compend. Fl. N. States, 372; Fl. N. York, ii, 197.—Rafinesque, Med. Bot. ii, 244.—Eaton, Manual, 6 ed. 231.—Beck, Bot. 324.—London, Arboretum, iv, 2057, f. 1968.—Lindley, Fl. Med. 305.—Dietrich, Syn. i, 551.—Eaton & Wright, Bot. 324.—Spach, Hist. Veg. xi, 263.—Emerson, Trees Massachusetts, 224; 2 ed. i, 256 & t.—Darby, Bot. S. States, 507.—Chapman, Fl. S. States, 426.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 106.—Lesquereux in Owen's 2d Rep. Arkansas, 389.—Wood, Cl. Book, 650; Bot. & Fl. 309.—Porcher, Resources S. Forests, 312.—C. De Candolle in Ann. Sci. Nat. 4 ser. xviii, 21, t. 3, f. 32; Prodr. xvi³, 148.—Lawson in Trans. Bot. Soc. Edinburgh, viii, 108.—Gray, Manual N. States, 5 ed. 457.—Koch, Dendrologie, ii, 663.—Young, Bot. Texas, 511.—Vasey, Cat. Forest Trees, 28.

M. Pennsylvanica, Lamarck, Dict. ii, 592.—Desfontaines, Hist. Arb. ii, 472.—Nouveau Duhamel, ii, 190, t. 55.—Pursh, Fl. Am. Sept. ii, 620.—Sprengel, Syst. i, 493.—Eaton, Manual, 6 ed. 232.—Eaton & Wright, Bot. 325.—Spach, Hist. Veg. xi, 262.

M. Carolinensis, Miller, Dict. No. 3.—Wangenheim, Amer. 102.—Willdenow, Spec. iv, 746; Enum. 1011.—Aiton, Hort. Kew. 2 ed. v, 379.—Pursh, Fl. Am. Sept. ii, 620.—Nuttall, Genera, ii, 235.—Elliott, Sk. ii, 678.—Eaton, Manual, 6 ed. 232.—Eaton & Wright, Bot. 324.—Darby, Bot. S. States, 507.

M. cerifera humilis, Marshall, Arbustum, 95.

M. cerifera, var. *latifolia*, Aiton, Hort. Kew. iii, 396.

M. cerifera, var. *media*, Michaux, Fl. Bor.-Am. ii, 227.—Chapman, Fl. S. States, 427.

M. cerifera, var. *arborescens*, Michaux, Fl. Bor.-Am. ii, 227.

M. cerifera, var. *pumila*, Michaux, Fl. Bor.-Am. ii, 227.—Pursh, Fl. Am. Sept. ii, 620.—Chapman, Fl. S. States, 427.

M. cerifera, var. *angustifolia*, C. De Candolle, Prodr. xvi³, 148.

M. cerifera sempervirens, Hort.

BAYBERRY. WAX MYRTLE.

Shores of lake Erie; Maine, and south near the coast to the Florida keys and southern Alabama.

A tree sometimes 12 meters in height, with a trunk 0.30 to 0.45 meter in diameter, or, except in the southern states, a low, much-branched shrub; usually on sandy beaches and dry hillsides, reaching its greatest development in the bottoms and rich hummocks of the Georgia and Florida coasts.

Wood light, soft, strong, brittle, very close-grained, compact; medullary rays numerous, thin; color, dark brown, the sap-wood lighter; specific gravity, 0.5637; ash, 0.51.

The leaves and stimulant and astringent bark of the roots sometimes employed by herbalists (*Am. Jour. Pharm.* 1863, 193.—*U. S. Dispensatory*, 14 ed. 257, 1706.—*Nat. Dispensatory*, 2 ed. 944). The wax which covers the small globular fruit, formerly largely collected and made into candles, and now, under the name of myrtle-wax, a popular remedy in the treatment of dysentery.

250.—*Myrica Californica*, Chamisso,

Linnaea, vi, 535.—Bentham, Pl. Hartweg, 336; Bot. Sulphur, 55.—Hooker, Fl. Bor.-Am. ii, 160.—Hooker & Arnott, Bot. Beechey, 390.—Lindley in Jour. London Hort. Soc. vii, 282.—Torrey in Pacific R. R. Rep. iv, 137; Bot. Wilkes Exped. 465.—Newberry in Pacific R. R. Rep. vi, 89.—Cooper in Pacific R. R. Rep. xii⁹, 68.—C. De Candolle, Prodr. xvi⁹, 153.—Gray in Proc. Am. Acad. vii, 401.—Vasey, Cat. Forest Trees, 28.—Hall in Coulter's Bot. Gazette, ii, 91.—Watson, Bot. California, ii, 81.

? *M. Xalapensis*, Hooker & Arnott, Bot. Beechey, 160.

Cape Foulweather, Oregon, south near the coast to the bay of Monterey, California.

A small evergreen tree, rarely exceeding 9 meters in height, with a trunk 0.30 to 0.45 meter in diameter, or toward its northern limits reduced to a low shrub; sandy beaches and gravelly hillsides.

Wood heavy, very hard, strong, brittle, very close-grained, compact; medullary rays numerous, thin, conspicuous; color, light rose, the sap-wood lighter; specific gravity, 0.6703; ash, 0.33.

CUPULIFERÆ.

251.—*Quercus alba*, Linnaeus,

Spec. 1 ed. 996.—Du Roi, Harbk. ii, 270, t. 5, f. 5.—Lamarck, Dict. i, 720.—Marshall, Arbustum, 119.—Wangenheim, Amer. 12, t. 3, f. 6.—Walter, Fl. Caroliniana, 235.—Aiton, Hort. Kew. iii, 358; 2 ed. v, 293.—Abbot, Insects Georgia, ii, t. 80, 87.—Michaux, Fl. Bor.-Am. ii, 195.—Muhlenberg & Willdenow in Neue Schriften Gesell. Nat. Fr. Berlin, iii, 395.—Willdenow, Spec. iv, 448; Enum. 977; Berl. Baumz. 346.—Persoon, Syn. ii, 570.—Desfontaines, Hist. Arb. ii, 508.—Michaux f. Hist. Arb. Am. ii, 13, t. 1; N. American Sylva, 3 ed. i, 22, t. 1.—Pursh, Fl. Am. Sept. ii, 633.—Barton, Prodr. Fl. Philadelph. 91; Compend. Fl. Philadelph. ii, 17.—Eaton, Manual, 108; 6 ed. 293.—Nuttall, Genera, ii, 215; Sylva, i, 14; 2 ed. i, 24.—Nouveau Duhamel, vii, 175.—Hayne, Dend. Fl. 158.—Elliott, Sk. ii, 607.—Sprengel, Syst. iii, 864.—Torrey, Compend. Fl. N. States, 359; Fl. N. York, ii, 192.—Audubon, Birds, t. 107, 147.—Beck, Bot. 330.—London, Arboretum, iii, 1864, f. 1723-1726 & t.—Hooker, Fl. Bor.-Am. ii, 158.—Eaton & Wright, Bot. 385.—Bigelow, Fl. Boston. 3 ed. 375.—Spach, Hist. Veg. xi, 155.—Emerson, Trees Massachusetts, 127, t. 1; 2 ed. i, 145 & t.—Griffith, Med. Bot. 585.—Penn. Cycl. xix, 216.—Richardson, Arctic Exped. 437.—Darlington, Fl. Cestrica, 3 ed. 266.—Darby, Bot. S. States, 511.—Cooper in Smithsonian Rep. 1858, 255.—Brendel in Trans. Illinois Ag. Soc. iii, 613, t. 1.—Chapman, Fl. S. States, 423.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 31.—Lesquereux in Owen's 2d Rep. Arkansas, 387.—Wood, Cl. Book, 645; Bot. & Fl. 306.—Porcher, Resourcee S. Forests, 257.—A. De Candolle, Prodr. xvi⁹, 22.—Örsted in Saerskitt. Aftryk. af Nat. For. Viden. Meddelt. Nos. 1-6, 1866, 66.—Liebmann, Chênes Am. Trop. t. xxxiii, 29, 30, 58, 59.—Gray, Manual N. States, 5 ed. 450; Hall's Fl. Texas, 21.—Koch, Dendrologie, ii⁹, 50.—Young, Bot. Texas, 505.—Vasey, Cat. Forest Trees, 25.—Broadhead in Coulter's Bot. Gazette, iii, 60.—Sears in Bull. Essex Inst. xiii, 179.—Britton in Bull. Torrey Bot. Club, viii, 126.—Bell in Geological Rep. Canada, 1879-'80, 52c.—Ridgway in Proc. U. S. Nat. Mus. 78.

? *Q. sinuata*, Walter, Fl. Caroliniana, 235.

Q. alba, var. *pinnatifida*, Michaux, Hist. Chênes Am. No. 4, t. 5, f. 1; Fl. Bor.-Am. ii, 195.—London, Arboretum, iii, 1864.

Q. alba, var. *repanda*, Michaux, Hist. Chênes Am. No. 4, t. 5, f. 2.—Pursh, Fl. Am. Sept. ii, 633.—Hayne, Dend. Fl. 159.—London, Arboretum, iii, 1864.

Q. alba, var. *pinnatifido-sinuata*, Hayne, Dend. Fl. 158.

Q. alba, var. *sinuata*, Hayne, Dend. Fl. 159.

Q. alba, var. *microcarpa*, A. De Candolle, Prodr. xvi⁹, 22.

WHITE OAK.

Northern Maine, valley of the Saint Lawrence river, Ontario, lower peninsula of Michigan to southeastern Minnesota, south to the Saint John's river and Tampa bay, Florida, west to the valley of Nodaway river, Missouri, western Arkansas, and the valley of the Brazos river, Texas.

A large tree of the first economic value, 24 to 45 meters in height, with a trunk 1.20 to 2.40 meters in diameter; all soils; very common and reaching its greatest development along the western slopes of the Alleghany mountains and in the valley of the Ohio river and its tributaries, here often forming more than half the forest growth.

Wood strong, very heavy, hard, tough, close-grained, liable to check unless carefully seasoned, durable in contact with the soil; layers of annual growth strongly marked by several rows of large open ducts; medullary rays broad, prominent; color, brown, the sap-wood lighter brown; specific gravity, 0.7470; ash, 0.41; largely used in ship-building, construction of all sorts, cooperage, in the manufacture of carriages, agricultural implements, and baskets, and for railway ties, fencing, interior finish, cabinet-making, fuel, etc.

A decoction of the astringent inner bark is employed medicinally in cases of hemorrhage, dysentery, etc. (*U. S. Dispensatory*, 14 ed. 755.—*Nat. Dispensatory*, 2 ed. 1196).

252.—*Quercus lobata*, Née,

Ann. Cienc. Nat. iii, 278.—Smith in Rees' Cycl. xxx, No. 77.—Persoon, Syn. ii, 571.—Nouveau Duhamel, vii, 180.—Poiret, Suppl. ii, 224.—Bentham, Pl. Hartweg. 337.—Liebmann in Dansk. Vidensk. Selsk. Forhandl. 1854, 14; Chênes Am. Trop. 23, t. 42, f. 1-3.—Torrey, Bot. Mex. Boundary Survey, 205; Bot. Wilkes Exped. 461, t. 15.—A. De Candolle, Prodr. xvi^e, 24.—Koch, Dendrologie, ii^e, 53.—Vasey, Cat. Forest Trees, 25.—Engelmann in Trans. St. Louis Acad. iii, 388; Wheeler's Rep. vi, 374; Bot. California. ii, 95.

Q. Hindsii, Bentham, Bot. Sulphur, 55.—Endlicher, Genera, Suppl. iv. 24.—Walpers, Ann. 1, 635.—Torrey in Pacific R. R. Rep. iv, 138; v, 365.—Newberry in Pacific R. R. Rep. vi, 29, 89, t. 1, f. 7.—Cooper in Smithsonian Rep. 1858, 261.—Bolander in Proc. California Acad. iii, 230.—Ørsted in Saerskitt. Aftryk. af Nat. For. Viden. Meddelt. 1866, Nos. 1-6, 66.—Liebmann, Chênes Am. Trop. t. 42, f. 4.—R. Brown Campst. Hors Sylvanæ, 52, f. 1-3.

Q. longiglанда, Torrey in Fremont's Geographical Mem. California, 15, 17.

Q. Ransomi, Kellogg in Proc. California Acad. i, 25.

WHITE OAK. WEEPING OAK.

California, west of the Sierra Nevadas from the valley of the upper Sacramento river south through the foothills and interior valleys to the San Bernardino mountains.

The largest of the Pacific oaks, often 30 meters in height, with a trunk 0.90 to 2.40 meters in diameter; very common through the central part of the state.

Wood moderately hard, fine-grained, compact; layers of annual growth marked by few large open ducts and containing few smaller ducts arranged in lines parallel to the broad, conspicuous medullary rays; color, light brown, the sap-wood lighter; specific gravity, 0.7409; ash, 0.30; of little economic value, and only used for fuel.

253.—*Quercus Garryana*, Douglas;

Hooker, Fl. Bor.-Am. ii, 159.—Hooker & Arnott, Bot. Beechey, 391.—Nuttall, Sylva, i, 1, t. 1; 2 ed. i, 14, t. 1.—Torrey in Pacific R. R. Rep. iv, 138; Bot. Wilkes Exped. 462.—Newberry in Pacific R. R. Rep. vi, 89.—Cooper in Smithsonian Rep. 1858, 260; Pacific R. R. Rep. xii^e, 28, 68; Am. Nat. iii, 407.—Lyll in Jour. Linnæan Soc. vii, 131, 144.—A. De Candolle, Prodr. xvi^e, 24.—Bolander in Proc. California Acad. iii, 229.—Ørsted in Saerskitt. Aftryk. af Nat. For. Viden. Meddelt. 1866, Nos. 1-6, 66.—Rothrock in Smithsonian Rep. 1858, 435.—Liebmann, Chênes Am. Trop. t. 40, f. 3.—Vasey, Cat. Forest Trees, 25.—Engelmann in Trans. St. Louis Acad. iii, 389; Bot. California, ii, 95.—Macoun in Geological Rep. Canada, 1875-76, 210.—G. M. Dawson in Canadian Nat. new ser. ix, 330.

Q. Newi, Liebmann in Dansk. Vidensk. Selsk. Forhandl. 1854, 173; Chênes Am. Trop. 23, t. xli, f. 1, 2.

Q. Douglasii, var. *?Newi*, A. De Candolle, Prodr. xvi^e, 24.

Q. Ørstediana, R. Brown Campst. in Ann. & Mag. Nat. Hist. April, 1871, 2.

Q. Jacobi, R. Brown Campst. in Ann. & Mag. Nat. Hist. April, 1871, 7.

WHITE OAK.

Vancouver's island, shores of Puget sound, south through western Washington territory, Oregon, and California to San Francisco bay; in Washington territory and Oregon extending to the eastern slopes of the Cascade mountains.

A tree 21 to 30 meters in height, with a trunk 0.60 to 0.90 meter in diameter, or at high elevations reduced to a low shrub; dry, gravelly soil; common.

Wood strong, hard, that of the young trees tough, close-grained, compact; layers of annual growth marked by one to three rows of open ducts; medullary rays, varying greatly in width, often conspicuous; color, light brown or yellow, the sap-wood lighter, often nearly white; specific gravity, 0.7453; ash, 0.39; somewhat used for carriage and cooperage stock, in cabinet-making, ship-building, and very largely for fuel; the best substitute for eastern white oak produced in the Pacific forests.

254.—*Quercus obtusiloba*, Michaux,

Hist. Chênes Am. No. 1, t. 1; Fl. Bor.-Am. ii, 194.—Smith in Rees' Cycl. xxx, No. 78.—Michaux f. Hist. Arb. Am. ii, 36, t. 4; N. American Sylva, 3 ed. i, 36, t. 5.—Pursh, Fl. Am. Sept. ii, 632.—Nuttall, Genera, ii, 215.—Barton, Compend. Fl. Philadelph. ii, 171.—Elliott, Sk. ii, 606.—Torrey, Compend. Fl. N. States, 359; Fl. N. York, ii, 190.—Beck, Bot. 329.—Eaton, Manual, 6 ed. 293.—Loudon, Arboretum, iii, 1870, f. 1732 & t.—Hooker, Fl. Bor.-Am. ii, 158.—Eaton & Wright, Bot. 384.—Scheele in Rømer, Texas, 446.—Darlington, Fl. Cestrica, 3 ed. 265.—Darby, Bot. S. States, 511.—Cooper in Smithsonian Rep. 1858, 255.—Brendel in Trans. Illinois Ag. Soc. iii, 615, t. 11.—Chapman, Fl. S. States, 423.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 32.—Leaqueroux in Owen's 2d Rep. Arkansas, 387.—Wood, Cl. Book, 645; Bot. & Fl. 306.—Engelmann in Trans. Am. Phil. Soc. new ser. xii, 209.—Ørsted in Saerskitt. Aftryk. af Nat. For. Viden. Meddelt. 1866, Nos. 1-6, 66.—Liebmann, Chênes Am. Trop. t. H, t. 33, f. 60.—Gray, Manual N. States, 5 ed. 451; Hall's, Pl. Texas, 21.—Young, Bot. Texas, 505.

Q. alba minor, Marshall, Arbustum, 120.—Muhlenberg & Willdenow in Neue Schriften Gesell. Nat. Fr. Berlin, iii, 396.

Q. stellata, Wangenheim, Amer. 78, t. 6, f. 15.—Abbot, Insects Georgia, ii, t. 77.—Willdenow, Spec. iv, 452; Enum, 977; Berl. Baumz. 349.—Persoon, Syn. ii, 570.—Aiton, Hort. Kew. 2 ed. v, 294.—Nouveau Duhamel, vii, 180.—Hayne, Dend. Fl. 161.—Nuttall, Sylva, i, 13; 2 ed. i, 23.—Spach, Hist. Veg. xi, 156.—Emerson, Trees Massachusetts, 133, t. 3; 2 ed. i, 151 & t.—A. De Candolle, Prodr. xvi³, 22.—Koch, Dendrologie, iii, 52.—Vasey, Cat. Forest Trees, 25.—Engelmann in Trans. St. Louis Acad. iii, 389.—Ridgway in Proc. U. S. Nat. Mus. 1882, 84.—Watson in Proc. Am. Acad, xviii, 156.

? *Q. villosa*, Walter, Fl. Caroliniana, 235.

Q. lobulata, Abbot, Insects Georgia, i, 47.

? *Q. Drummondii*, Liebmann in Dansk. Vidensk. Selsk. Forhandl. 1854, 170.—A. De Candolle, Prodr. xvi³, 24.

Q. obtusiloba, var. *parvifolia*, Chapman, Fl. S. States, 423.

Q. stellata, var. *Floridana*, A. De Candolle, Prodr. xvi³, 22.

POST OAK. IRON OAK.

Martha's Vineyard, Massachusetts, south to northern Florida, west through southern Ontario and Michigan to eastern Nebraska, Kansas, the Indian territory, and extending to the one hundredth meridian in central Texas.

A tree rarely exceeding 24 meters in height, with a trunk 0.90 to 1.50 meter in diameter, or on the Florida coast reduced to a low shrub (var. *parvifolia*, etc.); dry, gravelly uplands, clay barrens, or in the southwest on Cretaceous formations; the most common and widely-distributed oak of the Gulf states west of the Mississippi river, forming the principal growth of the Texas "cross-timbers."

Wood heavy, hard, close-grained, compact, checking badly in drying, very durable in contact with the soil; layers of annual growth marked by one to three rows of not large open ducts; medullary rays numerous, conspicuous; color, dark or light brown, the sap-wood lighter; specific gravity, 0.8367; ash, 0.79; largely used, especially in the southwest, for fencing, railway ties, and fuel, and somewhat for carriage stock, cooperage, construction, etc.

255.—*Quercus undulata*, var. *Gambelii*, Engelmann,

Wheeler's Rep. vi, 249.

Q. Gambelii, Nuttall in Jour. Philadelphia Acad. new ser. i, 179.—Torrey in Sitgreaves' Rep. 172, t. 18; Bot. Mex. Boundary Survey, 205.—Cooper in Smithsonian Rep. 1858, 260.—Liebmann, Chênes Am. Trop. 22, t. 40, f. 1.—Hemslay, Bot. Am.-Cent. iii, 171.

Q. alba, var. ? *Gunnisonii*, Torrey in Pacific R. R. Rep. ii, 130.—Watson in King's Rep. v, 321.—Porter in Hayden's Rep. 1871, 493.—Porter & Coulter, Fl. Colorado; Hayden's Surv. Misc. Pub. No. 4, 127.—Macoun in Geological Rep. Canada, 1875-76, 209.

Q. Douglasii, var. *Gambelii*, A. De Candolle, Prodr. xvi³, 23.

Q. stellata, var. *Utahensis*, A. De Candolle, Prodr. xvi³, 22.

? *Q. Emoryi*, Porter & Coulter, Fl. Colorado; Hayden's Surv. Misc. Pub. No. 4, 127 [not Torrey].

SCRUB OAK.

Near the mouth of the Pecos river (*Havard*), through the mountains of western Texas, and New Mexico to the Santa Catalina (*Lemmon, Pringle*) and San Francisco mountains, Arizona, eastern slopes of the Rocky mountains of Colorado north to the valley of the Platte river, and through the Wahsatch mountains of Utah.

A small tree, rarely 15 meters in height, with a trunk sometimes 0.60 meter in diameter, or often a low shrub spreading from underground shoots and forming dense thickets, reaching its greatest development on the high mountains of southern New Mexico and Arizona; the large specimens generally hollow and defective.

Wood heavy, hard, strong, that of young trees quite tough, close-grained, checking badly in drying; layers of annual growth marked by few not large open ducts; medullary rays numerous, conspicuous; color, rich dark brown, the sap-wood lighter; specific gravity, 0.8407; ash, 0.99; largely used for fuel, and in Utah the bark in tanning.

256.—*Quercus macrocarpa*, Michaux,

Hist. Chênes Am. No. 2, t. 2, 3; Fl. Bor.-Am. ii, 194.—Willdenow, Spec. iv, 453; Enum. 977; Berl. Baumz. 350.—Smith in Rees' Cycl. xxx, No. 80.—Persoon, Syn. ii, 570.—Poiret, Suppl. ii, 224.—Michaux f. Hist. Arb. Am. ii, 34, t. 3; N. American Sylva, 3 ed. i, 35, t. 4.—Pursh, Fl. Am. Sept. ii, 632.—Nuttall, Genera, ii, 215.—Nouveau Duhamel, vii, 182.—Hayne, Dend. Fl. 161.—Sprengel, Syst. iii, 863.—Torrey, Compend. Fl. N. States, 359; Nicollet's Rep. 160; Fl. N. York, ii, 191, t. 108.—Beck, Bot. 330.—Eaton, Manual, 6 ed. 293.—London, Arboretum, iii, 1869, f. 1731 & t.—Eaton & Wright, Bot. 385.—Spach, Hist. Veg. xi, 159.—Emerson, Trees Massachusetts, 132, t. 2; 2 ed. i, 149 & t.—Scheele in Rømer, Texas, 446.—Richardson, Arctic Exped. 437.—Cooper in Smithsonian Rep. 1858, 255.—Brendel in Trans. Illinois Ag. Soc. 131, t. 5, f. 21.—Chapman, Fl. S. States, 423.—Lesquereux in Owen's 2d Rep. Arkansas, 387.—Wood, Cl. Book, 645; Bot. & Fl. 306.—Engelmann in Trans. Am. Phil. Soc. new ser. xii, 209; Trans. St. Louis Acad. iii, 389.—A. De Candolle, Prodr. xvi², 20.—Ørsted in Saerskitt. Aftryk. af Nat. For. Viden. Meddelt. Nos. 1-6, 1866, 67.—Liebmann, Chênes Am. Trop. t. G, t. 33, f. 27, 28.—Gray, Manual N. States, 5 ed. 451.—Koch, Dendrologie, ii², 51.—Young, Bot. Texas, 506.—Winchell in Ludlow's Rep. Black Hills, 68.—Hayden in Warren's Rep. Nebraska & Dakota, 2 ed. 121.—Vasey, Cat. Forest Trees, 24.—Broadhead in Coulter's Bot. Gazette, iii, 60.—J. F. James in Jour. Cincinnati Soc. Nat. Hist. iv, 1 & t.—Ridgway in Proc. U. S. Nat. Mus. 1882, 81.—Bell in Geological Rep. Canada, 1879-'80, 49c.—Watson in Proc. Am. Acad. xviii, 156.

Q. olivæformis, Michaux f. Hist. Arb. Am. ii, 32, t. 2; N. American Sylva, 3 ed. i, 33, t. 3.—Smith in Rees' Cycl. xxx, No. 91.—Pursh, Fl. Am. Sept. ii, 632.—Nuttall, Genera, ii, 215; Sylva, i, 14; 2 ed. i, 24.—Nouveau Duhamel, vii, 181.—Sprengel, Syst. iii, 864.—Torrey, Compend. Fl. N. States, 359.—Fl. N. York, ii, 191.—Beck, Bot. 330.—Eaton, Manual, 6 ed. 293.—London, Arboretum, iii, 1869, f. 1730.—Eaton & Wright, Bot. 385.—Spach, Hist. Veg. xi, 159.—Gray, Manual N. States, 1 ed. 414.—A. De Candolle, Prodr. xvi², 20.—Ørsted in Saerskitt. Aftryk. af Nat. For. Viden. Meddelt. Nos. 1-6, 1866, 67.—Engelmann in Trans. St. Louis Acad. iii, 391.

Q. obtusiloba, var. *depressa*, Nuttall, Genera, ii, 215.

Q. macrocarpa, var. *olivæformis*, Gray, Manual N. States, 2 ed. 404; 5 ed. 451.

Q. macrocarpa, var. *abbreviata*, A. De Candolle, Prodr. xvi², 20.

Q. macrocarpa, var. *minor*, A. De Candolle, Prodr. xvi², 20.

Q. stellata, var. *depressa*, A. De Candolle, Prodr. xvi², 23.

BURE OAK. MOSSY-CUP OAK. OVER-CUP OAK.

Nova Scotia, New Brunswick, northern shores of lake Huron to lake Winnipeg, south to the valley of the Penobscot river, Maine (*C. E. Hamlin*), and along the shores of lake Champlain and the valley of the Ware river, Massachusetts, to Lancaster county, Pennsylvania, west to the eastern foot-hills of the Rocky mountains of Montana, central Nebraska and Kansas, southwest to the Indian territory and the valley of the Nueces river, Texas.

A large tree of the first economic value, 24 to 30 or, exceptionally, 50 meters in height, with a trunk 1.20 to 2.10 meters in diameter; rich bottoms and prairies; in the prairie region the principal growth of the "oak openings", and extending farther west and northwest than any oak of the Atlantic forests.

Wood heavy, strong, hard, tough, close-grained, compact, more durable in contact with the soil than that of other American oaks; layers of annual growth marked by one to three rows of small open ducts; medullary rays often broad and conspicuous; color, dark or rich light brown, the sap-wood much lighter; specific gravity, 0.7453; ash, 0.71; generally confounded with the less valuable white oak (*Q. alba*), and employed for the same purposes.

257.—*Quercus lyrata*, Walter,

Fl. Caroliniana, 235.—Abbot, Insects Georgia, ii, t. 83.—Michaux, Hist. Chênes Am. No. 3, t. 4; Fl. Bor.-Am. ii, 195.—Willdenow, Spec. iv, 453.—Smith in Rees' Cycl. xxx, No. 79.—Persoon, Syn. ii, 570.—Poiret, Suppl. ii, 224.—Michaux f. Hist. Arb. Am. ii, 42, t. 5; N. American Sylva, 3 ed. i, 39, t. 6.—Aiton, Hort. Kew. 2 ed. v, 295.—Pursh, Fl. Am. Sept. ii, 632.—Nouveau Duhamel, vii, 181.—Nuttall, Genera, ii, 215.—Elliott, Sk. ii, 607.—Sprengel, Syst. xi, 156.—Eaton, Manual, 6 ed. 295.—London, Arboretum, iii, 1871, f. 1733, 1734.—Eaton & Wright, Bot. 386.—Spach, Hist. Veg. xi, 156.—Darby, Bot. S. States, 511.—Cooper in Smithsonian Rep. 1858, 255.—Chapman, Fl. S. States, 423.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 33.—Lesquereux in Owen's 2d Rep. Arkansas, 387.—Wood, Bot. & Fl. 306.—A. De Candolle, Prodr. xvi², 19.—Ørsted in Saerskitt. Aftryk. af Nat. For. Viden. Meddelt. Nos. 1-6, 1866, 66.—Koch, Dendrologie, ii², 53.—Gray, Hall's Pl. Texas, 21.—Young, Bot. Texas, 506.—Vasey, Cat. Forest Trees, 25.—Engelmann in Trans. St. Louis Acad. iii, 389.—Ridgway in Proc. U. S. Nat. Mus. 1882, 80.

OVER-CUP OAK. SWAMP POST OAK. WATER WHITE OAK.

North Carolina, south near the coast to the Chattahoochee region of northern Florida, west through Alabama, Mississippi, and Louisiana to the valley of the Trinity river, Texas, and through Arkansas and southeastern Missouri (*Allenton, Letterman*) to middle Tennessee, southern Indiana and Illinois.

A tree 24 to 30 meters in height, with a trunk 0.60 to 0.90 meter in diameter; deep, often submerged, river swamps; rare in the Atlantic states; more common and reaching its greatest development in the valley of the Red river and the adjacent portions of Arkansas and Texas.

Wood heavy, hard, strong, tough, very durable in contact with the ground, close-grained, inclined to check in drying; layers of annual growth marked by one to three rows of large open ducts; medullary rays broad, numerous, conspicuous; color, rich dark brown, the sap-wood much lighter; specific gravity, 0.8313; ash, 0.65; used for the same purposes as that of the white oak (*Q. alba*).

258.—*Quercus bicolor*, Willdenow,

Neue Schriften Gesell. Nat. Fr. Berlin, iii, 396; Spec. iv, 440.—Smith in Rees' Cycl. xxx, No. 50.—Persoon, Syn. ii, 569.—Poirer, Suppl. ii, 219.—Pursh, Fl. Am. Sept. ii, 633.—Eaton, Manual, 107; 6 ed. 294.—Barton, Compend. Fl. Philadelph. ii, 172.—Nuttall, Genera, ii, 215; Sylva, i, 13; 2 ed. i, 23.—Nouveau Duhamel, vii, 165.—Sprengel, Syst. iii, 860.—Torrey, Compend. Fl. N. States, 359; Fl. N. York, ii, 192.—Beck, Bot. 331.—Bigelow, Fl. Boston. 3 ed. 375.—Eaton & Wright, Bot. 385.—Emerson, Trees Massachusetts, 135, t. 4; 2 ed. i, 153 & t.—Buckley in Am. Jour. Sci. 2 ser. xiii, 397.—Darlington, Fl. Cestrica, 3 ed. 266.—Lesquereux in Owen's 2d Rep. Arkansas, 387.—Wood, Cl. Book, 646; Bot. & Fl. 306.—A. De Candolle, Prodr. xvi², 20.—Ørsted in Saerskitt. Aftryk. af. Nat. For. Viden. Meddelt. Nos. 1-6, 1866, 67.—Gray, Manual N. States, 5 ed. 451.—Koch, Dendrologie, ii², 47.—Vasey, Cat. Forest Trees, 25.—Engelmann in Trans. St. Louis Acad. iii, 389.—Broadhead in Coulter's Bot. Gazette, iii, 60.—Sears in Bull. Essex Inst. xiii, 179.—Bell in Geological Rep. Canada, 1879-'80, 55c.—Ridgway in Proc. U. S. Nat. Mus. 1882, 79.

? *Q. Prinus platanoides*, Lamarck, Dict. i, 21.

Q. alba palustris, Marshall, Arbustum, 120.—Muhlenberg & Willdenow in Neue Schriften Gesell. Nat. Fr. Berlin, iii, 395.

Q. Prinus tomentosa, Michaux, Hist. Chênes Am. No. 5, t. 9, f. 2; Fl. Bor.-Am. ii, 196.—London, Arboretum, iii, 1876, f. 1739.

Q. Prinus, var. *discolor*, Michaux f. Hist. Arb. Am. ii, 46, t. 6; N. American Sylva, 3 ed. i, 41, t. 7.—Cooper in Smithsonian Rep. 1858, 255.—Brendel in Trans. Illinois Ag. Soc. iii, 617, t. 3.—Chapman, Fl. S. States, 424.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 34.

Q. bicolor, var. *mollis*, Nuttall, Genera, ii, 215.—Torrey, Compend. Fl. N. States, 359.

Q. Prinus, var. *bicolor*, Spach, Hist. Veg. xi, 158.

? *Q. bicolor*, var. *platanoides*, A. De Candolle, Prodr. xvi², 21.

SWAMP WHITE OAK.

Southern Maine, valley of the upper Saint Lawrence river, Ontario, southern peninsula of Michigan to southeastern Iowa and western Missouri, south to Delaware, and along the Alleghany mountains to northern Georgia, northern Kentucky, and northern Arkansas.

A large tree, 24 to 36 meters in height, with a trunk 1.20 to 2.40 or, exceptionally, over 3 meters ("Wadsworth Oak", Genesee, New York) in diameter; borders of streams and swamps, in deep alluvial soil; common and reaching its greatest development in the region south of the great lakes.

Wood heavy, hard, strong, tough, close-grained, inclined to check in seasoning; layers of annual growth marked by one to three rows of large open ducts; medullary rays broad and conspicuous; color, light brown, the sap-wood hardly distinguishable; specific gravity, 0.7662; ash, 0.58; used for the same purposes as that of the white oak (*Q. alba*).

259.—*Quercus Michauxii*, Nuttall,

Genera, ii, 215 (excl. syn.).—Elliot, Sk. ii, 609.—Sprengel, Syst. iii, 860.—Eaton, Manual, 6 ed. 295.—Eaton & Wright, Bot. 386.—Darby, Bot. S. States, 511.—Vasey, Cat. Forest Trees, 25.—Engelmann in Trans. St. Louis Acad. iii, 382.—Ward in Bull. U. S. Nat. Mus. No. 22, 113.—Ridgway in Proc. U. S. Nat. Mus. 1882, 81.

Q. Prinus palustris, Michaux, Hist. Chênes Am. No. 5, t. 6; Fl. Bor.-Am. ii, 196.—Michaux f. Hist. Arb. Am. ii, 51, t. 7; N. American Sylva, 3 ed. i, 44, t. 8.—Barton, Prodr. Fl. Philadelph. 91.—London, Arboretum, iii, 1872, f. 1735 & t.

Q. Prinus, var. *Michauxii*, Chapman, Fl. S. States, 424.

Q. Prinus, Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 33, in part.

Q. bicolor, var. *Michauxii*, Engelmann in Trans. St. Louis Acad. iii, 390.

BASKET OAK. COW OAK.

New Castle county, Delaware, south through the lower and middle districts to northern Florida, through the Gulf states to the valley of the Trinity river, Texas, and through Arkansas and southwestern Missouri to central Tennessee and Kentucky, and the valley of the lower Wabash river.

A tree 24 to 36 meters in height, with a trunk 1.20 to 2.10 meters in diameter; borders of streams and deep, often submerged, swamps; the common and most valuable white oak of the Gulf states, reaching its greatest development in the rich bottom lands of southeastern Arkansas and Louisiana.

Wood heavy, hard, very strong, tough, close-grained, compact, very durable in contact with the soil, easily split; layers of annual growth marked by few rather large open ducts; medullary rays broad, conspicuous; color, light brown, the sap-wood darker; specific gravity, 0.8039; ash, 0.45; largely used in the manufacture of agricultural implements, wheel stocks, baskets, for which it is unsurpassed, for cooperage, fencing, construction, and fuel.

The large, sweet, edible acorns eagerly devoured by cattle and other animals.

260.—*Quercus Prinus*, Linnæus,

Spec. 1 ed. 995.—Du Roi, Harbk. ii, 276, t. 6, f. 3.—Lamarck, Dict. i, 720.—Marshall, Arbustum, 125.—Wangenheim, Amer. 15, t. 4, f. 8.—Aiton, Hort. Kew. iii, 356; 2 ed. v, 290.—Moench, Meth. 348.—Abbot, Insects Georgia, ii, t. 82.—Muhlenberg & Willdenow in Neue Schriften Gesell. Nat. Fr. Berlin, iii, 397.—Michaux, Fl. Bor.-Am. ii, 195.—Willdenow, Spec. iv, 439; Enum. 975; Berl. Baumz. 339.—Smith in Rees' Cycl. xxx, No. 47.—Persoon, Syn. ii, 568.—Desfontaines, Hist. Arb. ii, 509.—Pursh, Fl. Am. Sept. ii, 633.—Barton, Compend. Fl. Philadelph. ii, 171.—Nuttall, Genera, ii, 215.—Nouveau Duhamel, vii, 164.—Hayne, Dend. Fl. 155.—Elliott, Sk. ii, 608.—Sprengel, Syst. iii, 859.—Torrey, Compend. Fl. N. States, 359.—Audubon, Birds, t. 50, 131.—Beck, Bot. 331.—Eaton, Manual, 6 ed. 294.—London, Arboretum, iii, 1872.—Eaton & Wright, Bot. 385.—Spach, Hist. Veg. xi, 157.—Penn. Cycl. xix, 216.—Darlington, Fl. Cestrica, 3 ed. 267.—Darby, Bot. S. States, 511.—Cooper in Smithsonian Rep. 1858, 255.—Chapman, Fl. S. States, 423.—Lesquereux in Owen's 2d Rep. Arkansas, 387.—Wood, Cl. Book, 645; Bot. & Fl. 306.—Porcher, Resources S. Forests, 264.—A. De Candolle, Prodr. xvi³, 21.—Örsted in Saerskitt. Aftryk. af Nat. For. Viden. Meddelt. Nos. 1-6, 67.—Gray, Manual N. States, 5 ed. 451.—Young, Bot. Texas, 506.—Koch, Dendrologie, ii², 48.—Vasey, Cat. Forest Trees, 25.—Engelmann in Trans. St. Louis Acad. iii, 390.

Q. Prinus, var. *monticola*, Michaux, Hist. Chênes Am. No. 5, t. 7; Fl. Bor.-Am. ii, 196.—Michaux f. Hist. Arb. Am. ii, 55, t. 8; N. American Sylva, 3 ed. i, 46, t. 9.—Barton, Prodr. Fl. Philadelph. 91.—London, Arboretum, iii, 1873, f. 1736.—Spach, Hist. Veg. xi, 158.—Cooper in Smithsonian Rep. 1858, 255.—Chapman, Fl. S. States, 424.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 34.—Wood, Cl. Book, 646.—A. De Candolle, Prodr. xvi³, 21.—Gray, Manual N. States, 5 ed. 451.—Vasey, Cat. Forest Trees, 25.—Bailey in Am. Nat. xiv, 892, f. 1-4.

Q. montana, Willdenow, Spec. iv, 440; Enum. 975; Berl. Baumz. 340.—Persoon, Syn. ii, 569.—Smith in Rees' Cycl. xxx, No. 49.—Pursh, Fl. Am. Sept. ii, 634.—Eaton, Manual, 107, 6 ed. 294.—Barton, Compend. Fl. Philadelph. ii, 172.—Nuttall, Genera, ii, 216.—Nouveau Duhamel, vii, 165, t. 47, f. 2.—Hayne, Dend. Fl. 156.—Elliott, Sk. ii, 609.—Sprengel, Syst. iii, 860.—Torrey, Compend. Fl. N. States, 354; Fl. N. York, ii, 192.—Beck, Bot. 331.—Bigelow, Fl. Boston. 3 ed. 377.—Eaton & Wright, Bot. 385.—Emerson, Trees Massachusetts, 138, t. 6; 2 ed. i, 156 & t.—Gray, Manual N. States, 1 ed. 414.—Darlington, Fl. Cestrica, 3 ed. 266.—Darby, Bot. S. States, 511.—Lesquereux in Owen's 2d Rep. Arkansas, 387.—Porcher, Resources S. Forests, 263.—Burgess in Coulter's Bot. Gazette, vii, 95.

Q. Prinus, var. *lata*, Aiton, Hort. Kew. 2 ed. v, 290.

Q. Castanea, Emerson, Trees Massachusetts, 137, t. 5; 2 ed. i, 155 & t. [not Muhlenberg & Willdenow].

CHESTNUT OAK. ROCK CHESTNUT OAK.

Blue hills, eastern Massachusetts, west to the shores of lake Champlain, shores of Quinté bay, Ontario (*Macoun*), and the valley of the Genesee river, New York, south to Delaware, and through the Alleghany Mountain region to northern Alabama, extending west to central Kentucky and Tennessee.

A tree 24 to 30 meters in height, with a trunk 0.90 to 1.20 meter in diameter; rocky banks and hillsides; very common and reaching its greatest development in the southern Alleghany region, here often forming a large portion of the forest growth.

Wood heavy, hard, strong, rather tough, close-grained, inclined to check in drying, durable in contact with the soil, containing few open ducts; medullary rays very broad, conspicuous; color, dark brown, the sap-wood lighter; specific gravity, 0.7499; ash, 0.77; largely used in fencing, for railway ties, etc.

The bark, rich in tannin, is largely used in preference to that of other North American white oaks in tanning leather.

261.—*Quercus prinoides*, Willdenow,

Neue Schriften Gesell. Nat. Fr. Berlin, iii, 397; Spec. iv, 440.—Persoon, Syn. ii, 569.—Poirer, Suppl. ii, 219.—Nouveau Duhamel, vii, 166.—Torrey, Fl. N. York, ii, 193, t. 109.—Gray, Manual N. States, 1 ed. 415.—Darlington, Fl. Cestrica, 3 ed. 267.—Chapman, Fl. S. States, 424.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 35.—Lesquereux in Owen's 2d Rep. Arkansas, 387.—Wood, Cl. Book, 646.—Koch, Dendrologie, ii², 49.—Young, Bot. Texas, 506.—Engelmann in Trans. St. Louis Acad. iii, 391.

Q. Prinus humilis, Marshall, Arbustum, 125.—Gray, Manual N. States, 5 ed. 452.

Q. Castanea, Muhlenberg & Willdenow in Neue Schriften Gesell. Nat. Fr. Berlin, iii, 396 [not Née].—Willdenow, Spec. iv, 441; Enum. 976; Berl. Baumz. 341.—Persoon, Syn. ii, 569.—Pursh, Fl. Am. Sept. ii, 634.—Smith in Rees' Cycl. xxx, No. 51.—Poirer, Suppl. ii, 219.—Eaton, Manual, 107; 6 ed. 294.—Barton, Compend. Fl. Philadelph. ii, 172.—Nuttall, Genera, ii, 216.—Hayne, Dend. Fl. 156.—Elliott, Sk. ii, 610.—Sprengel, Syst. iii, 860.—Torrey, Compend. Fl. N. States, 354; Fl. N. York, ii, 193.—Beck, Bot. 331.—Eaton & Wright, Bot. 385.—Gray, Manual N. States, 1 ed. 415.—Darlington, Fl. Cestrica, 3 ed. 267.—Darby, Bot. S. States, 511.—Brendel in Trans. Illinois Ag. Soc. iii, 619, t. 4.—Chapman, Fl. S. States, 424.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 34.—Lesquereux in Owen's 2d Rep. Arkansas, 387.—Wood, Cl. Book, 646.—Örsted in Saerskitt. Aftryk. af Nat. For. Viden. Meddelt. Nos. 1-6, 1866, 68.—Liebmann, Chênes Am. Trop. t. H, K. & 33, f. 31, 32.—Young, Bot. Texas, 506.

Q. Prinus, var. *acuminata*, Michaux, Hist. Chênes Am. No. 5, t. 8; Fl. Bor.-Am. ii, 196.—Michaux f. Hist. Arb. Am. ii, 61, t. 9; N. American Sylva, 3 ed. i, 49, t. 10.—Nouveau Duhamel, vii, 167.—London, Arboretum, iii, 1875, f. 1637.—Cooper in Smithsonian Rep. 1858, 255.—Wood, Bot. & Fl. 306.—Gray, Manual N. States, 5 ed. 451.—Vasey, Cat. Forest Trees, 25.

Q. Prinus pumila, Michaux, Hist. Chênes Am. No. 5, t. 9, f. 1; Fl. Bor.-Am. ii, 196.—London, Arboretum, iii, 1875, f. 1738

Q. Prinus Chinquapin, Michaux f. Hist. Arb. Am. ii, 65, t. 10; N. American Sylva, 3 ed. i, 50, t. 11.—A. De Candolle, Prodr. xvi², 21.

Q. Chinquapin, Pursh, Fl. Am. Sept. ii, 634.—Smith in Rees' Cycl. xxx, No. 48.—Nuttall, Genera, ii, 216.—Elliott, Sk. ii, 611.—Torrey, Compend. Fl. N. States, 354.—Beck, Bot. 331.—Eaton, Manual, 6 ed. 294.—Darlington, Fl. Cestrica, 2 ed. 536.—Eaton & Wright, Bot. 385.—Bigelow, Fl. Boston. 3 ed. 377.—Emerson, Trees Massachusetts, 140; 2 ed. i, 158 & t.—Darby, Bot. S. States, 511.

Q. Prinus, var. *oblongata*, Aiton, Hort. Kew. v, 290.

Q. Prinus, var. *prinoides*, Wood, Bot. & Fl. 306.

Q. Muhlenbergii, Engelmann in Trans. St. Louis Acad. iii, 591.—G. D. Butler in Coulter's Bot. Gazette, iii, 77.—Ridgway in Proc. U. S. Nat. Mus. 1882, 82.

YELLOW OAK. CHESTNUT OAK. CHINQUAPIN OAK.

Eastern Massachusetts, shores of lake Champlain, west along the northern shores of lakes Ontario and Erie, through southern Michigan to eastern Nebraska, eastern Kansas, and the Indian territory; south to Delaware and through the Alleghany region to northern Alabama and Mississippi, southwest to the Guadalupe mountains, western Texas (*Havard*).

A tree 24 to 30 or, exceptionally, 39 meters (*Ridgway*) in height, with a trunk 0.60 to 0.90 meter in diameter (*Q. Muhlenbergii*), or often, especially toward the eastern and western limits of its range, reduced to a low, slender shrub (*Q. prinoides*); dry hillsides and low, rich bottoms; rare, except as a shrub, east of the Alleghany mountains; very common in the Mississippi River basin, and reaching its greatest development in southern Arkansas.

Wood heavy, hard, very strong, close-grained, checking badly in drying, very durable in contact with the soil; layers of annual growth marked by rows of small open ducts; medullary rays broad, conspicuous; color, dark brown, the sap-wood much lighter; specific gravity, 0.8605; ash, 1.14; used for cooperage, wheel stock, fencing, railway ties, etc.

The small acorns sweet and edible.

NOTE.—Differences in the size and habit of individuals of this species, thus enlarged, seem to be dependent upon soil and climate, numerous intermediate forms connecting the extremes of eastern Massachusetts and the Mississippi valley.

262.—*Quercus Douglasii*, Hooker & Arnett,

Bot. Beechey, 391.—Hooker, Icon. iv, t. 382, 383.—Bentham, Pl. Hartweg. 337; Bot. Sulphur, 55.—Nuttall, Sylva, i, 10, t. 4; 2 ed. i, 20, t. 4.—Torrey in Pacific R. R. Rep. v, 365; Bot. Wilkes Exped. 462.—Cooper in Smithsonian Rep. 1858, 260.—A. De Candolle, Prodr. xvi², 23.—Bolander in Proc. California Acad. iii, 230.—Ørsted in Saerskitt. Aftryk. af. Nat. For. Viden. Meddelt. Nos. 1-6, 66.—Liebmann, Chênes Am. Trop. t. 41, f. 3, 4.—Vasey, Cat. Forest Trees, 25.—Engelmann in Trans. St. Louis Acad. iii, 392; Bot. California, ii, 95.—Hall in Coulter's Bot. Gazette, ii, 91.

Q. oblongifolia, var. *brevilobata*, Torrey in Bot. Wilkes Exped. 460.

MOUNTAIN WHITE OAK. BLUE OAK.

California, from about latitude 39°, south along the western foot-hills of the Sierra Nevadas below 4,000 feet elevation, and through the Coast ranges to the San Gabriel mountains.

A tree 18 to 24 meters in height, with a trunk 0.60 to 1.20 meter in diameter; common on the low foot-hills of the sierras.

Wood very hard, heavy, strong, brittle, inclined to check in drying; layers of annual growth marked by several rows of small open ducts and containing many scattered groups of smaller ducts; medullary rays numerous, varying greatly in width; color, dark brown, becoming nearly black with exposure, the thick sap-wood light brown; specific gravity, 0.8928; ash, 0.84.

263.—*Quercus oblongifolia*, Torrey,

Sitgreaves' Rep. 173; Bot. Mex. Boundary Survey, 206; Ives' Rep. 28.—Cooper in Smithsonian Rep. 1858, 261.—A. De Candolle, Prodr. xvi², 36.—Watson, Pl. Wheeler, 17.—Vasey, Cat. Forest Trees, 26.—Engelmann in Trans. St. Louis Acad. iii, 393; Bot. California, ii, 96.

Q. undulata, var. *oblongata*, Engelmann in Wheeler's Rep. vi, 250.

WHITE OAK.

California, foot-hills of the San Gabriel mountains, and in San Diego county (here occupying a narrow belt, 30 miles in width some 30 miles from the coast, *Parish Brothers*); foot-hills of the mountain ranges of southern Arizona and New Mexico; southward into Mexico.

A small evergreen tree, 12 to 15 meters in height, with a trunk 0.45 to 0.60 meter in diameter; the large specimens generally hollow and defective.

Wood very heavy, hard, strong, brittle, very close-grained, checking badly in drying; layers of annual growth hardly distinguishable, containing few small open ducts arranged in many groups parallel to the broad and very conspicuous medullary rays; color, very dark brown or almost black, the thick sap-wood brown; specific gravity, 0.9441; ash, 2.61; of little economic value except as fuel.

264.—*Quercus grisea*, Liebmann,

Dansk. Vidensk. Selsk. Forhandl. 1854, 13; Chênes Am. Trop. t. 46, f. 1, 2.—A. De Candolle, Prodr. xvi², 35.—Ørsted in Saerskitt. Aftryk. af. Nat. For. Viden. Meddelt. Nos. 1-6, 1866, 69.—Rusby in Bull. Torrey Bot. Club, ix, 78.—Watson in Proc. Am. Acad. xviii, 156.

Q. pungens, Liebmann in Dansk. Vidensk. Selsk. Forhandl. 1854, 13; Chênes Am. Trop. 22, t. 45, f. 1-3.—A. De Candolle, Prodr. xvi², 36.—Ørsted in Saerskitt. Aftryk. af. Nat. For. Viden. Meddelt. Nos. 1-6, 69.—Rusby in Bull. Torrey Bot. Club ix, 78.

Q. undulata, var. *grisea*, Engelmann in Trans. St. Louis Acad. iii, 382; Wheeler's Rep. vi, 250.

Q. undulata, var. *pungens*, Engelmann in Trans. St. Louis Acad. iii, 392; Wheeler's Rep. vi, 250; Bot. California, ii, 96.—Palmer in Am. Nat. xii, 596.

Q. undulata, var. *Wrightii*, Engelmann in Trans. St. Louis Acad. iii, 382, 392.

WHITE OAK.

Mountains of southern Colorado and western Texas (*Havard*), southern New Mexico and Arizona from 5,000 to 10,000 feet elevation, west to the Colorado desert of California; southward into northern Mexico.

A tree 15 to 24 meters in height, with a trunk rarely exceeding 0.60 meter in diameter, or reduced to a low, much-branched shrub; a polymorphous species, varying greatly in habit and in the shape and texture of the leaves, but apparently well characterized by its connate cotyledons; the large specimens generally hollow and defective.

Wood very heavy, strong, hard, close-grained, checking badly in drying; layers of annual growth marked by one or two rows of small open ducts, these connected by rows of similar ducts parallel to the numerous conspicuous medullary rays; color, very dark brown, the thick sap-wood much lighter; specific gravity, 1.0092; ash, 1.82.

265.—*Quercus reticulata*, Humboldt & Bonpland,

Pl. Æquin. ii, 40, t. 86.—Poirot, Suppl. v, 609.—Sprengel, Syst. iii, 860.—Loudon, Arboretum, iii, 1944, f. 1865.—Michaux f. N. American Sylva, 3 ed. i, 90.—A. De Candolle, Prodr. xvi², 33.—Ørsted in Saerskitt. Aftryk. af. Nat. For. Viden. Meddelt. Nos. 1-6, 67.—Liebmann, Chênes Am. Trop. t. H, t. 34, f. 10-16, t. 35, f. 15-22.—Vasey, Cat. Forest Trees, 26.—Engelmann in Trans. St. Louis Acad. iii, 383; Wheeler's Rep. vi, 250.—Hemsley, Bot. Am.-Cent. iii, 176.—Watson in Proc. Am. Acad. xviii, 156. ?

Q. spicata, Humboldt & Bonpland, Pl. Æquin. ii, 46, t. 89.—Bentham, Pl. Hartweg. No. 429.

Q. decipiens, Martens & Galeotti in Bull. Brux. v, 10.

? *Q. reticulata*, var. *Greggii*, A. De Candolle, Prodr. xvi², 34.—Hemsley, Bot. Am.-Cent. iii, 176.

Southeastern Arizona, San Francisco and Santa Rita mountains from 7,000 to 10,000 feet elevation; southward into northern Mexico.

A small tree, 9 to 12 meters in height, with a trunk 0.30 to 0.45 meter in diameter; dry, gravelly slopes.

Wood very heavy, hard, close-grained, checking badly in drying, containing many small, scattered, open ducts; medullary rays numerous, very broad; color, dark brown, the sap-wood lighter; specific gravity, 0.9479; ash, 0.52.

266.—*Quercus Durandii*, Buckley,

Proc. Philadelphia Acad. 1860, 445; 1881, 121.—Gray, Hall's Pl. Texas, 21.—Young, Bot. Texas, 507.—Vasey, Cat. Forest Trees, 26.—Watson in Proc. Am. Acad. xviii, 156.

Q. obtusifolia, var. ? *breviloba*, Torrey, Bot. Mex. Boundary Survey, 206.

Q. annulata, Buckley in Proc. Philadelphia Acad. 1860, 445.

Q. San-Sabeana, Buckley in Young, Bot. Texas, 507.

Q. undulata, Engelmann in Trans. St. Louis Acad. iii, 392, in part [not Torrey].

Alabama, Wilcox county (*Buckley*), valley of the Little Cahaba river, Bibb county (*Mohr*); Shreveport, Louisiana f, (*Buckley*); Texas, Dallas (*Reverchon*), valley of the Colorado river (*Buckley, Mohr, Sargent*), west and south.

A tree 21 to 24 meters in height, with a trunk 0.60 to 1.20 meter in diameter; rich bottom lands or dry *mesas* and limestone hills, then reduced to a low shrub, forming dense, impenetrable thickets of great extent (*Q. San-Sabeana*); rare and local in Alabama; the common and most valuable white oak of western Texas.

Wood very heavy and hard, strong, brittle, close-grained, inclined to check in drying; layers of annual growth marked by few large open ducts; medullary rays numerous, conspicuous; color, brown, the sap-wood lighter; specific gravity, 0.9507; ash, 1.78; used for the same purposes as that of the white oak (*Q. alba*).

267.—*Quercus virens*, Aiton,

Hort. Kew. iii, 356; 2 ed. v, 287.—Bartram, Travels, 2 ed. 82.—Michaux, Hist. Chênes Am. No. 6, t. 10, 11; Fl. Bor.-Am. ii, 196.—Willdenow, Spec. iv, 425; Enum. 974.—Robin, Voyages, iii, 264.—Smith in Rees' Cycl. xxx, No. 5.—Persoon, Syn. ii, 567.—Desfontaines, Hist. Arb. ii, 507.—Poiret, Suppl. ii, 213.—Michaux f. Hist. Arb. Am. ii, 67, t. 11; N. American Sylva, 3 ed. i, 52, t. 12.—Pursh, Fl. Am. Sept. ii, 626.—Nuttall, Genera, ii, 214; Sylva, i, 16; 2 ed. i, 28.—Nouveau Duhamel, vii, 151.—Elliott, Sk. ii, 595.—Sprengel, Syst. iii, 858.—Cobbett, Woodlands, 446.—Eaton, Manual, 6 ed. 294.—London, Arboretum, iii, 1918, f. 1802, 1803 & t.—Eaton & Wright, Bot. 385.—Spach, Hist. Veg. xi, 177.—Engelmann & Gray in Jour. Boston Soc. Nat. Hist. v, 234.—Scheele in Rømer, Texas, 446; Appx. 147.—Penn. Cycl. xix, 216.—Darby, Bot. S. States, 510.—Torrey, Bot. Mex. Boundary Survey, 206.—Cooper in Smithsonian Rep. 1858, 255.—Chapman, Fl. S. States, 421.—Curtis in Rep. Geological Surv. N. Carolina, 35.—Wood, Cl. Book, 643; Bot. & Fl. 305.—Porcher, Resources S. Forests, 263.—A. De Candolle, Prodr. xvi², 37.—Ørsted in Saerskitt. Aftryk. af Nat. For. Viden. Meddelt. Nos. 1-6, 69.—Gray, Manual N. States, 5 ed. 452; Hall's Pl. Texas, 21.—Liebmann, Chênes Am. Trop. t. 33, f. 50-57.—Young, Bot. Texas, 503.—Vasey, Cat. Forest Trees, 26.—Engelmann in Trans. St. Louis Acad. iii, 383; iv, 191.—Hemsley, Bot. Am.-Cent. iii, 178.—Watson in Proc. Am. Acad. xviii, 155.

Q. Virginiana, Miller, Dict. 7 ed. No. 17.—Koch, Dendrologie, ii², 57.

Q. Phellos, var. *sempervirens*, Marshall, Arbustum, 124.

Q. sempervirens, Walter, Fl. Caroliniana, 234.

Q. oleoides, Chamisso & Schlechtendal in Linnæa, v, 79.—Martens & Galeotti in Bull. Brux. x, No. 3.—Ørsted in Saerskitt. Aftryk. af Nat. For. Viden. Meddelt. Nos. 1-6, 1866, 69.

Q. retusa, Liebmann in Dansk. Vidensk. Selsk. Forhandl. 1854, 187.—Ørsted in Saerskitt. Aftryk. af Nat. For. Viden. Meddelt. Nos. 1-6, 1866, 69.

LIVE OAK.

Mojo Jack bay, Virginia, south along the coast to bay Biscayne and cape Romano, Florida, along the Gulf coast to Mexico, extending through western Texas to the valley of the Red river, the Apache and Gaudalupe mountains and the mountains of northern Mexico south of the Rio Grande at 6,000 to 8,000 feet elevation (*Havard*); in Costa Rica (*Q. retusa*).

An evergreen tree of great economic value, 15 to 18 meters in height, with a trunk 1.50 to 2.10 meters in diameter, or in the interior of Texas much smaller, often shrubby; on the coast, rich hummocks and ridges, a few feet above water-level; common and reaching its greatest development in the south Atlantic states.

Wood very heavy, hard, strong, tough, very close-grained, compact, difficult to work, susceptible of a beautiful polish; layers of annual growth obscure, often hardly distinguishable, containing many small open ducts arranged in short broken rows parallel to the broad, conspicuous medullary rays; color, light brown or yellow, the sap-wood nearly white; specific gravity, 0.9501; ash, 1.14; formerly very largely and now occasionally used in ship-building.

268.—*Quercus chrysolepis*, Liebmann,

Dansk. Vidensk. Selsk. Forhandl. 1854, 173; Chênes Am. Trop. 23, t. 47.—Torrey, Bot. Mex. Boundary Survey, 206; Bot. Wilkes Exped. 458.—Cooper in Smithsonian Rep. 1858, 260.—Kellogg in Proc. California Acad. ii, 45.—A. De Candolle, Prodr. xvi³, 37.—Bolander in Proc. California Acad. iii, 231.—Ørsted in Saerskitt. Aftryk. af. Nat. For. Viden. Meddelt. Nos. 1-6, 1866, 69.—Vasey, Cat. Forest Trees, 25.—Engelmann in Trans. St. Louis Acad. iii, 383, 393; Wheeler's Rep. vi, 374; Bot. California, ii, 97.—Watson in Proc. Am. Acad. xi, 119.—Palmer in Am. Nat. xii, 596.

Q. fulvescens, Kellogg in Proc. California Acad. i, 67, 71.—Newberry in Pacific R. R. Rep. vi, 27, 89.

Q. crassipocula, Torrey in Pacific R. R. Rep. iv, 137; v, 365, t. 9.

? *Q. oblongifolia*, R. Brown Campst. in Ann. & Mag. Nat. Hist. April, 1871, 4 [not Torrey].

LIVE OAK. MAUL OAK. VALPARAISO OAK.

Cow Creek valley, Oregon, south through the California Coast ranges and along the western slopes of the Sierra Nevada and San Bernardino mountains between 3,000 and 8,000 feet elevation, and south into Lower California; southeastern Arizona, San Francisco (*Greene*) and Santa Catalina mountains (*Pringle*).

An evergreen tree of great economic value, 18 to 27 meters in height, with a trunk sometimes 1.50 meter in diameter, or at high elevations reduced to a low, narrow-leaved shrub (var. *vaccinifolia*, *Engelmann in Trans. St. Louis Acad.* iii, 393; *Bot. California*, ii, 97.—*Q. vaccinifolia*, *Kellogg in Trans. California Acad.* ii, 96).

Wood heavy, very strong and hard, tough, close-grained, compact, difficult to work, containing many rather small open ducts arranged in wide bands parallel to the broad, conspicuous medullary rays; color, light brown, the sap-wood darker; specific gravity, 0.8493; ash, 0.60; somewhat used in the manufacture of agricultural implements, wagons, etc.; the most valuable oak of the Pacific forests.

269.—*Quercus Emoryi*, Torrey,

Emory's Rep. 151, t. 9; Bot. Mex. Boundary Survey, 206; Pacific R. R. Rep. iv, 138; Ives' Rep. 28.—Watson in Pl. Wheeler, 17.—Vasey, Cat. Forest Trees, 26.—Engelmann in Trans. St. Louis Acad. iii, 382, 387, 394; Wheeler's Rep. vi, 250.—Palmer in Am. Nat. xii, 596.—Hemsley, Bot. Am.-Cent. iii, 170.

Q. hastata, Liebmann in Dansk. Vidensk. Selsk. Forhandl. 1854, 13; Chênes Am. Trop. 22.—A. De Candolle, Prodr. xvi³, 36.—Ørsted in Saerskitt. Aftryk. af. Nat. For. Viden. Meddelt. Nos. 1-6, 1866, 69.

BLACK OAK.

Bexar and Comal counties, Texas, through the mountain ranges of western Texas, of southern New Mexico, and of eastern and southern Arizona.

A tree 12 to 15 meters in height, with a trunk 0.30 to 0.90 meter in diameter, or toward its eastern limits in Texas reduced to a low shrub; common and reaching its greatest development in southwestern New Mexico and southern Arizona between 5,000 and 7,000 feet elevation near streams in open cañons; dry, gravelly soil, the large specimens hollow and defective.

Wood very heavy, not hard, strong, brittle, close-grained, compact; layers of annual growth marked by several rows of small open ducts, these connected by narrow groups of similar ducts parallel to the broad, conspicuous medullary rays; color, dark brown or almost black, the thick sap-wood bright brown tinged with red; specific gravity, 0.9263; ash, 2.36.

270.—*Quercus agrifolia*, Née,

Ann. Cienc. Nat. iii, 271.—Fischer, Misc. Hisp. i, 108.—Willdenow, Spec. iv, 431.—Persoon, Syn. ii, 568.—Smith in Rees' Cycl. xxx, No. 29.—Pursh, Fl. Am. Sept. ii, 627.—Nuttall, Genera, ii, 214; Sylva, i, 5, t. 2; 2 ed. i, 16, t. 2.—Nonveau Duhamel, vii, 156.—Sprengel, Syst. iii, 859.—Eaton, Manual, 6 ed. 292.—London, Arboretum, iii, 1894.—Bentham, Pl. Hartweg. 337; Bot. Sulphur, 55.—Eaton & Wright, Bot. 384.—Hooker, Icon. iv, t. 377.—Hooker & Arnott, Bot. Beechey, 391.—Jour. Hort. Soc. London, vi, 157 & t.—Carrière in Fl. des Serres, vii, 137 & f.—Torrey in Sitgreaves' Rep. 173; Pacific R. R. Rep. iv, 138; v, 365; vii, 20; Bot. Mex. Boundary Survey, 206; Ives' Rep. 28; Bot. Wilkes Exped. 460.—Paxton's Brit. Flower Gard. ii, 44.—Newberry in Pacific R. R. Rep. vi, 32, f. 9.—Bolander in Proc. California Acad. iii, 229.—A. De Candolle, Prodr. xvi³, 37.—Ørsted in Saerskitt. Aftryk. af. Nat. For. Viden. Meddelt. Nos. 1-6, 1866, 69.—Liebmann, Chênes Am. Trop. t. 44.—Vasey, Cat. Forest Trees, 25.—Engelmann in Trans. St. Louis Acad. iii, 383; Wheeler's Rep. vi, 374; Bot. California, ii, 98.—Hemsley, Bot. Am.-Cent. iii, 167.

Q. oxyadenia, Torrey in Sitgreaves' Rep. 172, t. 17.—Cooper in Smithsonian Rep. 1858, 261.

Q. acutiglans, Kellogg in Proc. California Acad. i, 25.

ENCENO. COAST LIVE OAK.

California, Mendocino county, south through the Coast Range valleys to Lower California.

A large evergreen tree, 24 to 30 meters in height, with a trunk 1.20 to 2.10 meters in diameter, or, rarely, reduced to a low shrub (var. *frutescens*, *Engelmann in Bot. California*, ii, 98); rare at the north; common south of San Francisco bay, and the largest and most generally distributed oak in the extreme southwestern part of the state; dry slopes and ridges.

Wood heavy, hard, strong, brittle, close-grained, compact; layers of annual growth hardly distinguishable, containing many large open ducts arranged in several rows parallel to the broad, conspicuous medullary rays; color, light brown or red, the sap-wood darker brown; specific gravity, 0.8253; ash, 1.28; of little value except as fuel.

271.—*Quercus Wislizeni*, A. DeCandolle,

Prodr. xvi², 67.—Ørsted in Saerskitt. Aftryk. af Nat. For. Viden. Meddelt. Nos. 1-6, 1866, 73.—Vasey, Cat. Forest Trees, 27.—Engelmann in Trans. St. Louis Acad. iii, 385, 396; *Bot. California*, ii, 98.

Q. Morehus, Kellogg in Proc. California Acad. ii, 36.

LIVE OAK.

California, mount Shasta region, south along the western slopes of the Sierra Nevadas to Tulare county, and in the Coast ranges south to the Santa Lucia mountains.

An evergreen tree, 15 to 18 meters in height, with a trunk 0.90 to 1.80 meter in diameter, or toward its northeastern limits reduced to a shrub 0.90 to 3 meters in height (var. *frutescens*, *Engelmann in Bot. California*, ii, 99); not common.

Wood heavy, very hard, strong, close-grained, compact, containing numerous large open ducts arranged in irregular bands parallel to the broad, conspicuous medullary rays; color, light brown tinged with red, the sap-wood lighter; specific gravity, 0.7855; ash, 1.02.

272.—*Quercus rubra*, Linnaeus,

Spec. 1 ed. 996.—Du Roi, Harbk. ii, 265.—Lamarck, Dict. i, 720.—Walter, Fl. Caroliniana, 234.—Aiton, Hort. Kew. iii, 357; 2 ed. v, 292.—Mönch, Meth. 348.—Abbot, Insects Georgia, ii, t. 103.—Michaux, Hist. Chênes No. 2, t. 35, 36; Fl. Bor.-Am. ii, 200.—Willdenow, Spec. iv, 445; Enum. 976; Berl. Baumz. 342.—Smith in Rees' Cycl. xxx, No. 60.—Persoon, Syn. ii, 569.—Desfontaines, Hist. Arb. ii, 511.—Michaux f. Hist. Arb. Am. ii, 126, t. 26; N. American Sylva, 3 ed. i, 84, t. 28.—Pursh, Fl. Am. Sept. ii, 630.—Eaton, Manual, 106; 6 ed. 293.—Nuttall, Genera, ii, 214.—Barton, Compend. Fl. Philadelph. ii, 169.—Nouveau Duhamel, vii, 170.—Hayne, Dend. Fl. 157.—Elliott, Sk. ii, 602.—Sprengel, Syst. iii, 863.—Torrey, Compend. Fl. N. States, 358; Nicotlet's Rep. 160; Fl. N. York, 189, t. 106.—Beck, Bot. 329.—London, Arboretum, iii, 1877, f. 1740-1744 & t.—Hooker, Fl. Bor.-Am. ii, 158.—Bigelow, Fl. Boston. 3 ed. 376.—Eaton & Wright, Bot. 384.—Spach, Hist. Veg. xi, 165.—Emerson, Trees Massachusetts, 48, t. 10; 2 ed. i, 163 & t.—Scheele in Roemer, Texas, 446.—Penn. Cycl. xix, 216.—Darlington, Fl. Cestrica, 3 ed. 269.—Darby, Bot. S. States, 510.—Cooper in Smithsonian Rep. 1858, 255.—Brendel in Trans. Illinois Ag. Soc. iii, 369, t. 9.—Chapman, Fl. S. States, 422.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 41.—Lesquereux in Owen's 2d Rep. Arkansas, 388.—Wood, Cl. Book, 644; Bot. & Fl. 306.—Porcher, Resources S. Forests, 262.—Engelmann in Trans. Am. Phil. Soc. new ser. v, 209; Trans. St. Louis Acad. iii, 394.—A. De Candolle, Prodr. xvi², 60.—Ørsted in Saerskitt. Aftryk. af Nat. For. Viden. Meddelt. Nos. 1-6, 1866, 72.—Gray, Manual N. States, 5 ed. 454; Hall's Pl. Texas, 21.—Liebmann, Chênes Am. Trop. t. A, B.—Koch, Dendrologie, ii², 70.—Young, Bot. Texas, 504.—Hayden in Warren's Rep. Nebraska & Dakota, 2 ed. 121.—Vasey, Cat. Forest Trees, 26.—Macoun in Geological Rep. Canada, 1875-'76, 209.—Sears in Bull. Essex Inst. xiii, 179.—Ridgway in Proc. U. S. Nat. Mus. 1882, 83.—Bell in Geological Rep. Canada, 1879-'80, 51^c.

Q. rubra maxima, Marshall, Arbustum, 122.—Muhlenberg & Willdenow in Neue Schriften Gesell. Nat. Fr. Berlin, iii, 395.

Q. rubra, var. *latifolia*, Lamarck, Dict. i, 720.—Aiton, Hort. Kew. 2 ed. v, 292.—London, Arboretum, iii, 1877.

Q. rubra, var. *montana*, Aiton, Hort. Kew. 2 ed. v, 292.—London, Arboretum, iii, 1877.

Q. ambigua, Michaux f. Hist. Arb. Am. ii, 120, t. 24; N. American Sylva, 3 ed. i, 81, t. 26 [not HBK.].—Pursh, Fl. Am. Sept. ii, 630.—Nuttall, Genera, ii, 214.—Eaton, Manual, 6 ed. 293.—London, Arboretum, iii, 1881, f. 1749 & t.—Eaton & Wright, Bot. 384.

Q. coccinea, var. *rubra*, Spach, Hist. Veg. xi, 165.

Q. coccinea, var. *ambigua*, Gray, Manual N. States, 5 ed. 454.

Q. rubra, var. *runcinata*, A. De Candolle, Prodr. xvi², 60.—Engelmann in Trans. St. Louis Acad. iii, 542.

RED OAK. BLACK OAK.

Nova Scotia, southern New Brunswick to eastern Minnesota, western Iowa, eastern Kansas, and the Indian territory, south to northern Florida, southern Alabama and Mississippi, and the valley of the San Antonio river, Texas.

A large tree, 24 to 30 or, exceptionally, 45 meters (*Ridgway*) in height, with a trunk 1.20 to 2.10 meters in diameter; very common, especially at the north, in all soils and extending farther north than any Atlantic oak.

Wood heavy, hard, strong, coarse-grained, inclined to check in drying; layers of annual growth marked by several rows of very large open ducts; medullary rays few, conspicuous; color, light brown or red, the sap-wood somewhat darker; specific gravity, 0.6540; ash, 0.26; now largely used for clapboards, cooperage, and somewhat for interior finish, in the manufacture of chairs, etc.

Var. *Texana*, Buckley,

Proc. Philadelphia Acad. 1881, 123.—Engelmann in Coulter's Bot. Gazette, vii, 14.

Q. palustris, Torrey & Gray in Pacific R. R. Rep. ii, 175 [not Du Roi].

Q. coccinea, var. *microcarpa*, Torrey, Bot. Mex. Boundary Survey, 206.

Q. Texana, Buckley in Proc. Philadelphia Acad. 1860, 445.—Young, Bot. Texas, 507.

RED OAK.

Western Texas, valley of the Colorado river with the species and replacing it south and west, extending to the valley of the Nueces river and the Limpia mountains (*Havard*).

A tree 21 to 24 meters in height, with a trunk rarely exceeding 0.60 meter in diameter.

Wood heavier, harder, much closer-grained than the species, not checking in drying; layers of annual growth marked with fewer and smaller open ducts; specific gravity, 0.9080; ash, 0.85.

273.—*Quercus coccinea*, Wangenheim,

Amer. 44, t. 4. f. 9.—Muhlenberg & Willdenow in Neue Schriften Gesell. Nat. Fr. Berlin, iii, 398.—Michaux, Hist. Chênes Am. No. 18, t. 31, 32; Fl. Bor.-Am. ii, 199.—Willdenow, Spec. iv, 445; Enum. 976; Berl. Baumz. 343.—Smith in Rees' Cycl. xxx, 61.—Persoon, Syn. ii, 569.—Desfontaines, Hist. Arb. ii, 511.—Poiret, Suppl. ii, 221.—Michaux f. Hist. Arb. Am. ii, 116, t. 23; N. American Sylva, 3 ed. i, 79, t. 25.—Aiton, Hort. Kew. 2 ed. v, 292.—Pursh, Fl. Am. Sept. ii, 630.—Eaton, Manual, 108; 6 ed. 292.—Nuttall, Genera, ii, 214.—Barton, Compend. Fl. Philadelph. ii, 169.—Nouveau Duhamel, vii, 171.—Hayne, Dend. Fl. 157.—Elliott, Sk. ii, 602.—Sprengel, Syst. iii, 863.—Torrey, Compend. Fl. N. States, 358; Fl. N. York, ii, 189.—Beck, Bot. 329.—London, Arboretum, iii, 1879, f. 1746-1748 & t.—Eaton & Wright, Bot. 384.—Bigelow, Fl. Boston. 3 ed. 376.—Spach, Hist. Veg. xi, 165.—Emerson, Trees Massachusetts, 144, t. 9; 2 ed. i, 163 & t.—Scheele in Roemer, Texas, 446.—Penn. Cycl. xix, 216.—Darlington, Fl. Cestrica, 3 ed. 268.—Darby, Bot. S. States, 510.—Cooper in Smithsonian Rep. 1858, 255.—Chapman, Fl. S. States, 422.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 40.—Lesquereux in Owen's 2d Rep. Arkansas, 388.—Wood, Cl. Book, 645; Bot. & Fl. 306.—A. De Candolle, Prodr. xvi², 61.—Ørsted in Saerskitt. Aftryk. af. Nat. For. Viden. Meddelt. Nos. 1-6, 1866, 72.—Gray, Manual N. States, 5 ed. 453.—Liebmann, Chênes Am. Trop. t. B.—Koch, Dendrologie, ii², 69.—Young, Bot. Texas, 504.—Vasey, Cat. Forest Trees, 26.—Engelmann in Trans. St. Louis Acad. iii, 385, 394.—Ridgway in Proc. U. S. Nat. Mus. 1882, 80.—Watson in Proc. Am. Acad. xviii, 156.

Q. rubra, β . Linnaeus, Spec. 1 ed. 996.—Aiton, Hort. Kew. iii, 357.

SCARLET OAK.

Southern Maine to northern New York, Ontario, northern Michigan and Minnesota, eastern Iowa and northeastern Missouri, south to Delaware and southern Tennessee, and through the Alleghany region to northern Florida.

A tree 24 to 30 or, exceptionally, 54 meters (*Ridgway*) in height, with a trunk rarely exceeding 0.60 to 1.20 meter in diameter; at the east in dry, sandy soil or, less commonly, in rich, deep soil; in the northwestern prairie region with *Q. macrocarpa* forming the oak-opening growth; not common and reaching its greatest development in the basin of the lower Ohio river.

Wood heavy, hard, strong, coarse-grained; layers of annual growth strongly marked by several rows of large open ducts; medullary rays thin, conspicuous; color, light brown or red, the sap-wood rather darker; specific gravity, 0.7405; ash, 0.19; if used at all, confounded with that of *Q. rubra*.

274.—*Quercus tinctoria*, Bartram,

Travels, 2 ed. 37.—Abbot, Insects Georgia, ii, t. 56.—Michaux, Hist. Chênes Am. No. 13, t. 24, 25; Fl. Bor.-Am. ii, 198.—Willdenow, Spec. iv, 444; Enum. 976; Berl. Baumz. 344.—Desfontaines, Hist. Arb. ii, 509.—Poirot, Suppl. ii, 221.—Michaux f. Hist. Arb. Am. ii, 110, t. 22; N. American Sylva, 3 ed. i, 76, t. 24.—Aiton, Hort. Kew. 2 ed. v, 291.—Pursh, Fl. Am. Sept. ii, 629.—Smith in Rees' Cycl. xxx, No. 58.—Barton, Prodr. Fl. Philadelph. 91; Compend. Fl. Philadelph. ii, 168.—Eaton, Manual, 108; 6 ed. 292.—Nuttall, Genera, ii, 214; Sylva, i, 21; 2 ed. i, 32.—Nonveau Duhamel, vii, 169.—Hayne, Dend. Fl. 156.—Elliott, Sk. ii, 601.—Sprengel, Syst. iii, 862.—Torrey, Compend. Fl. N. States, 357; Fl. N. York, ii, 188.—Audubon, Birds, t. 82.—Beck, Bot. 328.—London, Arboretum, iii, 1884, f. 1753, 1754.—Hooker, Fl. Bor.-Am. ii, 158.—Bigelow, Fl. Boston. 3 ed. 376.—Eaton & Wright, Bot. 384.—Spach, Hist. Veg. xi, 164.—Emerson, Trees Massachusetts, 141, t. 7; 2 ed. i, 160 & t.—Griffith, Med. Bot. 586.—Gray, Manual N. States, 1 ed. 416.—Darlington, Fl. Cestricea, 3 ed. 268.—Darby, Bot. S. States, 510.—Cooper in Smithsonian Rep. 1858, 255.—Brendel in Trans. Illinois Ag. Soc. iii, 627, t. 8.—Chapman, Fl. S. States, 422.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 39.—Lesquereux in Owen's 2d Rep. Arkansas, 388.—Wood, Cl. Book, 645.—Engelmann in Proc. Am. Phil. Soc. new ser. xii, 209; Trans. St. Louis Acad. iii, 395.—Porcher, Resources S. Forests, 238.—Ørsted in Saerskitt. Aftryk. af. Nat. For. Viden. Meddelt. Nos. 1-6, 1866, 45, 72, f. 18.—Liebmann, Chênes Am. Trop. 9, f. 6.—Young, Bot. Texas, 504.—Hayden in Warren's Rep. Nebraska & Dakota, 2 ed. 121.—Guibourt, Hist. Drogues, 7 ed. ii, 288.—Vasey, Cat. Forest Trees, 27.—Bentley & Trimen, Med. Fl. iv, 251, t. 251.—Ridgway in Proc. U. S. Nat. Mus. 1882, 84.

? *Q. velutina*, Lamarck, Dict. i, 172.—Koch, Dendrologie, ii², 68.

Q. nigra, Marshall, Arbustum, 120 [not Linnæus].—Wangenheim, Amer. 79, t. 6, f. 16.

Q. rubra, Wangenheim, Amer. 14, t. 3, f. 7 [not Linnæus].—Muhlenberg & Willdenow in Neue Schriften Gesell. Nat. Fr. Berlin, iii, 399.

Q. discolor, Aiton, Hort. Kew. iii, 358.—Abbot, Insects Georgia, ii, 111.—Willdenow, Spec. iv, 444; Berl. Baumz. 345.—Poirot, Suppl. ii, 221.—Smith in Rees' Cycl. xxx, No. 59.—Nuttall, Genera, ii, 214.—Elliott, Sk. ii, 601.—Sprengel, Syst. iii, 863.—Beck, Bot. 329.—Eaton, Manual, 6 ed. 292.—Eaton & Wright, Bot. 384.

Q. tinctoria, var. *angulosa*, Michaux, Fl. Bor.-Am. ii, 198.—London, Arboretum, iii, 1858.

Q. tinctoria, var. *sinuosa*, Michaux, Fl. Bor.-Am. ii, 198.—London, Arboretum, iii, 1885, f. 1755-1757.—Liebmann, Chênes Am. Trop. t. C.

? *Q. shumardii*, Buckley in Proc. Philadelphia Acad. 1860, 445.

Q. coccinea, var. *tinctoria*, Gray, Manual N. States, 5 ed. 454.—Wood, Cl. Book, 306.—A. De Candolle, Prodr. xvi², 61.

BLACK OAK. YELLOW-BARK OAK. QUERCITRON OAK. YELLOW OAK.

Southern Maine to northern Vermont, Ontario, southern Minnesota, eastern Nebraska, eastern Kansas, and the Indian territory, south to the Chattahoochee region of western Florida, southern Alabama and Mississippi, and eastern Texas.

A large tree, 24 to 36 or, exceptionally, 48 meters (*Ridgway*) in height, with a trunk 0.90 to 1.80 meter in diameter; generally on dry or gravelly uplands; very common.

Wood heavy, hard, strong, not tough, coarse-grained, liable to check in drying; layers of annual growth marked by several rows of very large open ducts; color, bright brown tinged with red, the sap-wood much lighter; specific gravity, 0.7045; ash, 0.28; somewhat used for cooperage, construction, etc.

The bark largely used in tanning; the intensely bitter inner bark yields a valuable yellow dye, and is occasionally used medicinally in the form of decoctions, etc., in the treatment of hemorrhage (*U. S. Dispensatory*, 14 ed. 756.—*Nat. Dispensatory*, 2 ed. 1196).

275.—*Quercus Kelloggii*, Newberry,

Pacific R. R. Rep. vi, 89, 286, f. 6.—Torrey, Bot. Wilkes Exped. 468.—R. Brown Campst. Horæ Sylvanæ, 58, f. 4-6.—Engelmann in Bot. California, ii, 90.

Q. rubra, Bentham, Pl. Hartweg. 337 [not Linnæus].

Q. tinctoria, var. *Californica*, Torrey in Pacific R. R. Rep. iv, 138; Bot. Mex. Boundary Survey, 205; Ives' Rep. 28.

Q. Californica, Cooper in Smithsonian Rep. 1858, 261.

Q. Sonomensis, Bentham in De Candolle Prodr. xvi², 62.—Bolander in Proc. California Acad. iii, 230.—Ørsted in Saerskitt. Aftryk. af. Nat. For. Viden. Meddelt. Nos. 1-6, 1866, 72.—Vasey, Cat. Forest Trees, 27.—Engelmann in Wheeler's Rep. vi, 374.—Palmer in Am. Nat. xii, 596.

BLACK OAK.

Valley of the Mackenzie river, Oregon, south through the Coast ranges and along the western slopes of the Sierra Nevada and San Bernardino mountains to the southern borders of California.

A large tree, 18 to 24 meters in height, with a trunk 0.90 to 1.20 meter in diameter, or at high elevations reduced to a shrub; the most common and important oak of the valleys of southwestern Oregon and the California Sierras.

Wood heavy, hard, strong, very brittle, close-grained, compact; layers of annual growth marked by several rows of large open ducts; medullary rays few, broad, conspicuous; color, light red, the thin sap-wood lighter; specific gravity, 0.6435; ash, 0.26; of little value, except as fuel; the bark somewhat used in tanning.

276.—*Quercus nigra*, Linnæus,

Spec. 1 ed. 995.—Lamarck, Dict. i, 721.—Wangenheim, Amer. 77, t. 5, f. 13.—Walter, Fl. Caroliniana, 234.—Aiton, Hort. Kew. iii, 357; 2 ed. v, 291.—Abbot, Insects Georgia, i, 50; ii, 58.—Michaux, Hist. Chênes Am. No. 17, t. 22, 23; Fl. Bor.-Am. ii, 198.—Muhlenberg & Willdenow in Neue Schriften Gesell. Nat. Fr. Berlin, iii, 399.—Willdenow, Spec. iv, 442.—Smith in Rees' Cycl. xxx, No. 53.—Persoon, Syn. ii, 569.—Desfontaines, Hist. Arb. ii, 509.—Pursh, Fl. Am. Sept. ii, 629.—Eaton, Manual, 108; 6 ed. 292.—Barton, Compend. Fl. Philadelph. ii, 168.—Nouveau Duhamel, vii, 168.—Elliott, Sk. ii, 600.—Sprengel, Syst. iii, 862.—Torrey, Compend. Fl. N. States, 357; Fl. N. York, ii, 188; Bot. Mex. Boundary Survey, 206.—Audubon, Birds, t. 116.—Beck, Bot. 328.—London, Arboretum, iii, 1890, f. 1764, 1765.—Eaton & Wright, Bot. 384.—Spach, Hist. Veg. xi, 162.—Darlington, Fl. Cestrica, 3 ed. 267.—Darby, Bot. S. States, 510.—Cooper in Smithsonian Rep. 1858, 255.—Brendel in Trans. Illinois Ag. Soc. iii, 625, t. 7.—Chapman, Fl. S. States, 421.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 38.—Lesquereux in Owen's 2d Rep. Arkansas, 388.—Wood, Cl. Book, 644; Bot. & Fl. 306.—A. De Candolle, Prodr. xvi^{is}, 63.—Ørsted in Saerskitt. Aftryk. af Nat. For. Viden. Meddelt. Nos. 1-6, 72.—Gray, Manual N. States, 5 ed. 453; Hall's Pl. Texas, 21.—Liebmann, Chênes Am. Trop. t. A.—Koch, Dendrologie, ii^{is}, 61.—Young, Bot. Texas, 503.—Vasey, Cat. Forest Trees, 26.—Ridgway in Proc. Nat. Mus. 1882, 82.—Watson in Proc. Am. Acad. xviii, 156.

Q. nigra, var. *latifolia*, Lamarck, Dict. i, 721.

Q. nigra integrifolia, Marshall, Arbustum, 121.

? *Q. aquatica*, Walter, Fl. Caroliniana, 234.

Q. Marylandica, Muhlenberg & Willdenow in Neue Schriften Gesell. Nat. Fr. Berlin, iii, 399.

BLACK JACK. JACK OAK.

Long island, New York, west through northern Ohio and Indiana to about latitude 55° N. in Wisconsin, southern Minnesota, eastern Nebraska, Kansas, and the Indian territory to about 99° west longitude, south to Matanzas inlet and Tampa bay, Florida, and the valley of the Nueces river, Texas.

A small tree, sometimes 12 or even 18 meters in height, with a trunk rarely exceeding 0.60 meter in diameter, or more often much smaller; dry, barren uplands, or often on heavy clay soils; very common through the southern states, and reaching its greatest development in southwestern Arkansas, Indian territory, and eastern Texas, forming, with the post-oak (*Q. obtusiloba*), the growth of the Texas cross-timbers.

Wood heavy, hard, strong, checking badly in drying; layers of annual growth marked by several rows of large open ducts; medullary rays broad, conspicuous; color, rather dark rich brown, the sap-wood much lighter; specific gravity, 0.7324; ash, 1.16; of little value except as fuel.

277.—*Quercus falcata*, Michaux,

Hist. Chênes Am. No. 16, t. 28; Fl. Bor.-Am. ii, 199.—Persoon, Syn. ii, 569.—Poiret, Suppl. ii, 221.—Michaux f. Hist. Arb. Am. ii, 104, t. 21; N. American Sylva, 3 ed. i, 73, t. 23.—Pursh, Fl. Am. Sept. ii, 630.—Nuttall, Genera, ii, 214.—Barton, Compend. Fl. Philadelph. ii, 170.—Nouveau Duhamel, vii, 169.—Elliott, Sk. ii, 604.—Torrey, Compend. Fl. N. States, 358.—Beck, Bot. 329.—Eaton, Manual, 6 ed. 293.—London, Arboretum, iii, 1882, f. 1750, 1751.—Lindley, Fl. Med. 292.—Eaton & Wright, Bot. 384.—Darlington, Fl. Cestrica, 3 ed. 269.—Darby, Bot. S. States, 510.—Cooper in Smithsonian Rep. 1858, 255.—Chapman, Fl. S. States, 422.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 39.—Lesquereux in Owen's 2d Rep. Arkansas, 388.—Wood, Cl. Book, 644; Bot. & Fl. 306.—Porcher, Resources S. Forests, 256.—A. De Candolle, Prodr. xvi^{is}, 59.—Ørsted in Saerskitt. Aftryk. af Nat. For. Viden. Meddelt. Nos. 1-6, 1866, 72.—Gray, Manual N. States, 5 ed. 453; Hall's Pl. Texas, 21.—Liebmann, Chênes Am. Trop. t. A, t. 22, f. 3.—Young, Bot. Texas, 505.—Vasey, Cat. Forest Trees, 26.—Ridgway in Proc. U. S. Nat. Mus. 1882, 80.

Q. rubra montana, Marshall, Arbustum, 123.

Q. nigra digitata, Marshall, Arbustum, 121.

Q. cuneata, Wangenheim, Amer. 78, t. 5, f. 14.—Koch, Dendrologie, ii^{is}, 64.

Q. elongata, Muhlenberg & Willdenow in Neue Schriften Gesell. Nat. Fr. Berlin, iii, 400.—Willdenow, Spec. iv, 444.—Smith in Rees' Cycl. xxx, 57.—Aiton, Hort. Kew. 2 ed. v, 291.

Q. triloba, Michaux, Hist. Chênes Am. No. 14, t. 26.—Willdenow, Spec. iv, 443; Berl. Baumz. 342.—Smith in Rees' Cycl. xxx, No. 54.—Persoon, Syn. ii, 569.—Poiret, Suppl. ii, 220.—Aiton, Hort. Kew. 2 ed. v, 291.—Pursh, Fl. Am. Sept. ii, 628.—Hayne, Deud. Fl. 156.—Sprengel, Syst. iii, 862.—Torrey, Compend. Fl. N. States, 357.—Beck, Bot. 328.—Eaton, Manual, 6 ed. 292.—Eaton & Wright, Bot. 384.—Wood, Cl. Book, 644; Bot. & Fl. 306.

Q. falcata, var. *triloba*, Nuttall, Genera, ii, 214.—Elliott, Sk. ii, 604.—Darby, Bot. S. States, 511.—A. De Candolle, Prodr. xvi³, 59.

Q. falcata, var. *pagodaefolia*, Elliott, Sk. ii, 605.—Darby, Bot. S. States, 511.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 39.

Q. discolor, var. *triloba*, Spach, Hist. Veg. xi, 163.

Q. falcata, var. *Ludoviciana*, A. De Candolle, Prodr. xvi³, 59.

SPANISH OAK. RED OAK.

Long island, New York, south to Hernando county, Florida, through the Gulf states to the valley of the Brazos river, Texas, and through Arkansas and southeastern Missouri to central Tennessee and Kentucky, southern Illinois and Indiana.

A large tree, 24 to 30 meters in height, with a trunk 0.90 to 1.80 meter in diameter; dry, gravelly uplands and barrens; in the north Atlantic states only near the coast; rare; most common and reaching its greatest development in the south Atlantic and Gulf states, where, in the middle districts, it is the most common forest tree.

Wood heavy, very hard and strong, not durable, coarse-grained, checking badly in drying; layers of annual growth strongly marked by several rows of large open ducts; medullary rays few, conspicuous; color, light red, the sap-wood lighter; specific gravity, 0.6928; ash, 0.25; somewhat used for cooperage, construction, etc., and very largely for fuel.

The bark rich in tannin.

278.—*Quercus Catesbæi*, Michaux,

Hist. Chênes Am. No. 17, t. 29, 30; Fl. Bor.-Am. ii, 199.—Abbot, Insects Georgia, i, 27, t. 14.—Willdenow, Spec. iv, 446.—Smith in Rees Cycl. xxx, No. 62.—Persoon, Syn. 569.—Desfontaines, Hist. Arb. ii, 511.—Poirot, Suppl. ii, 221.—Michaux f. Hist. Arb. Am. ii, 101, t. 20; N. American Sylva, 3 ed. i, 71, t. 22.—Pursh, Fl. Am. Sept. ii, 630.—Nuttall, Genera, ii, 214.—Nouveau Duhamel, vii, 172.—Elliott, Sk. ii, 603.—Sprengel, Syst. iii, 866.—Torrey, Compend. Fl. N. States, 358.—Beck, Bot. 329.—Eaton, Manual, 6 ed. 293.—London, Arboretum, iii, 1889, f. 1762, 1763.—Eaton & Wright, Bot. 384.—Spach, Hist. Veg. xi, 162.—Darby, Bot. S. States, 510.—Cooper in Smithsonian Rep. 1858, 255.—Chapman, Fl. S. States, 422.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 41.—Wood, Cl. Book, 644; Bot. & Fl. 306.—A. De Candolle, Prodr. xvi³, 59.—Örsted in Saerskitt. Aftryk. af Nat. For. Viden. Meddelt. Nos. 1-6, 1866, 72.—Koch, Dendrologie, ii³, 67.—Young, Bot. Texas, 503.—Vasey, Cat. Forest Trees, 26.

? *Q. lævis*, Walter, Fl. Caroliniana, 234.

TURKEY OAK. SCRUB OAK. FORKED-LEAF BLACK JACK. BLACK JACK.

North Carolina, south near the coast to cape Malabar and Pease creek, Florida, and along the coast of Alabama and Mississippi.

A small tree, 7 to 15 meters in height, with a trunk 0.45 to 0.60 meter in diameter; very common in the south Atlantic and east Gulf states upon barren sandy hills and ridges of the maritime pine belt; rare in Mississippi.

Wood heavy, hard, strong, close-grained, compact; layers of annual growth marked by several rows of large open ducts and containing many much smaller ducts arranged in short lines parallel to the broad, conspicuous medullary rays; color, light brown tinged with red, the sap-wood somewhat lighter; specific gravity, 0.7294; ash, 0.87; largely used for fuel.

279.—*Quercus palustris*, Du Roi,

Harbk. ii, 268, t. 5, f. 4.—Wangenheim, Amer. 76, t. 5, f. 10.—Michaux, Hist. Chênes Am. No. 19, t. 33, 34; Fl. Bor.-Am. ii, 200.—Willdenow, Spec. iv, 446; Enum. 976; Berl. Baumz. 343.—Persoon, Syn. ii, 569.—Desfontaines, Hist. Arb. ii, 511.—Poirot, Suppl. ii, 222.—Michaux f. Hist. Arb. Am. ii, 123, t. 25; N. American Sylva, i, 83, t. 27.—Aiton, Hort. Kew. 2 ed. v, 292.—Smith in Rees' Cycl. xxx, No. 6.—Pursh, Fl. Am. Sept. ii, 631.—Barton, Prodr. Fl. Philadelph. 91; Compend. Fl. Philadelph. ii, 170.—Eaton, Manual, 108; 6 ed. 293.—Nuttall, Genera, ii, 214.—Nouveau Duhamel, vii, 172.—Hayne, Dend. Fl. 158.—Sprengel, Syst. iii, 863.—Torrey, Compend. Fl. N. States, 358; Fl. N. York, ii, 190, t. 107.—Beck, Bot. 329.—London, Arboretum, iii, 1887, f. 1758-1761 & t.—Eaton & Wright, Bot. 384.—Spach, Hist. Veg. xi, 166.—Darlington, Fl. Cestrica, 3 ed. 269.—Cooper in Smithsonian Rep. 1858, 255.—Brendel in Trans. Illinois Ag. Soc. iii, 631.—Lesquereux in Owen's 2d Rep. Arkansas, 388.—Wood, Cl. Book, 644; Bot. & Fl. 306.—A. De Candolle, Prodr. xvi³, 60.—Örsted in Saerskitt. Aftryk. af Nat. For. Viden. Meddelt. Nos. 1-6, 1866, 23, 72, f. 4.—Gray, Manual N. States, 5 ed. 454.—Liebmann, Chênes Am. Trop. t. A.—Koch, Dendrologie, ii³, 71.—Emerson, Trees Massachusetts, 2 ed. i, 167 & t.—Vasey, Cat. Forest Trees, 27.—W. E. Stone in Bull. Torrey Bot. Club, ix, 57.—Ridgway in Proc. U. S. Nat. Mus. 1882, 83.—Burgess in Coulter's Bot. Gazette, vii, 95.—Chapman, Fl. S. States, Suppl. 649.

Q. rubra, var. *dissecta*, Lamarck, Dict. i, 120.

Q. rubra ramosissima, Marshall, Arbustum, 122.—Muhlenberg & Willdenow in Neue Schriften Gesell. Nat. Fr. Berlin, 398.

PIN OAK. SWAMP SPANISH OAK. WATER OAK.

Valley of the Connecticut river, Massachusetts (Amherst, *Stone*), to central New York, south to Delaware and the District of Columbia; southern Wisconsin to eastern Kansas, southern Arkansas, and southeastern Tennessee.

A tree 24 to 30 or, exceptionally, 36 meters (*Ridgway*) in height, with a trunk 0.90 to 1.50 meter in diameter; low, rich soil, generally along the borders of streams and swamps; most common and reaching its greatest development west of the Alleghany mountains.

Wood heavy, hard, very strong, coarse-grained, inclined to check badly in drying; layers of annual growth marked by several rows of large open ducts; medullary rays broad, numerous, conspicuous; color, light brown, the sap-wood rather darker; specific gravity, 0.6938; ash, 0.81; somewhat used for shingles, clapboards, construction, and in cooperage.

280.—*Quercus aquatica*, Walter,

Fl. Caroliniana, 234.—Aiton, Hort. Kew. iii, 357; 2 ed. v, 290.—Abbot, Insects Georgia, ii, t. 59, 79.—Michaux, Hist. Chênes Am. No. 11, t. 19, 20, 21; Fl. Bor.-Am. ii, 198.—Muhlenberg & Willdenow in Neue Schriften Gesell. Nat. Fr. Berlin, iii, 399.—Persoon, Syn. ii, 569.—Desfontaines, Hist. Arb. ii, 509.—Poiret, Suppl. ii, 220.—Michaux f. Hist. Arb. Am. ii, 89, t. 17; N. American Sylva, 3 ed. i, 65, t. 19.—Smith in Rees' Cycl. xxx, No. 52.—Pursh, Fl. Am. Sept. ii, 628.—Barton, Compend. Fl. Philadelph. ii, 168.—Nouveau Duhamel, vii, 167.—Elliott, Sk. ii, 599.—Sprengel, Syst. iii, 862.—Torrey, Compend. Fl. N. States, 357.—Audubon, Birds, t. 24.—Beck, Bot. 328.—Eaton, Manual, 6 ed. 292.—Loudon, Arboretum, iii, 1892, f. 1767.—Eaton & Wright, Bot. 384.—Spach, Hist. Veg. xi, 161.—Darby, Bot. S. States, 510.—Cooper in Smithsonian Rep. 1858, 255.—Chapman, Fl. S. States, 421.—Curtis in Rep. Geological Surv. N. Carolina, 37.—Lesquereux in Owen's 2d Rep. Arkansas, 388.—Wood, Cl. Book, 643; Bot. & Fl. 305.—A. De Candolle, Prodr. xvi³, 67.—Örsted in Saerskitt. Aftryk. af Nat. For. Viden. Meddelt. Nos. 1-6, 1866, 72.—Gray, Manual N. States, 5 ed. 452; Hall's Pl. Texas, 21.—Liebmann, Chênes Am. Trop. t. D.—Young, Bot. Texas, 503.—Vasey, Cat. Forest Trees, 26.

Q. nigra aquatica, Lamarck, Dict. i, 721.

Q. nigra trifida, Marshall, Arbustum, 121.

? *Q. uliginosa*, Wangenheim, Amer. 80, t. 6, f. 18.

Q. hemisphærica, Willdenow, Spec. iv, 443.—Poiret, Suppl. ii, 628.—Pursh, Fl. Am. Sept. ii, 628.—Smith in Rees' Cycl. xxx, No. 56, 628.—Nuttall, Genera, ii, 214.—Eaton, Manual, 6 ed. 295.—Eaton & Wright, Bot. 385.—Michaux f. N. American Sylva, 3 ed. 187.

Q. nana, Willdenow, Spec. 448.—Elliott, Sk. ii, 599.

Q. aquatica, vars. *cuneata*, *elongata*, *indivisa*, *attenuata*, Aiton, Hort. Kew. 2 ed. v, 290.

Q. hemisphærica, var. *nana*, Nuttall, Genera, ii, 214.

Q. aquatica, var. *hybrida*, Chapman, Fl. S. States, 421.

Q. nigra, Koch, Dendrologie, ii³, 61, in part.

WATER OAK. DUCK OAK. POSSUM OAK. PUNK OAK.

Sussex county, Delaware, south through the coast and middle districts to cape Malabar and Tampa bay, Florida, through the Gulf states to the valley of the Colorado river, Texas, and through Arkansas to the valley of the Black river, southeastern Missouri (Poplar Bluffs, *Letterman*), middle Kentucky and Tennessee.

A tree 15 to 24 meters in height, with a trunk 0.60 to 1.20 meter in diameter; generally along streams and bottoms in heavy, undrained soil, or, more rarely, upon uplands; very common and reaching its greatest development along the large streams in the maritime pine belt of the eastern Gulf states.

Wood heavy, hard, strong, coarse-grained, compact; layers of annual growth marked by several rows of large open ducts; medullary rays thin, conspicuous; color, rather light brown, the sap-wood lighter; specific gravity, 0.7244; ash, 0.51; probably not used except as fuel.

281.—*Quercus laurifolia*, Michaux,

Hist. Chênes Am. No. 10, t. 17; Fl. Bor.-Am. ii, 197.—Willdenow, Spec. iv, 427.—Persoon, Syn. ii, 567.—Smith in Rees' Cycl. xxx, No. 14.—Aiton, Hort. Kew. 2 ed. v, 288.—Pursh, Fl. Am. Sept. ii, 627.—Nuttall, Genera, ii, 214.—Nouveau Duhamel, vii, 153.—Elliott, Sk. ii, 597.—Sprengel, Syst. iii, 857.—Eaton, Manual, 6 ed. 294.—Loudon, Arboretum, iii, 1897, f. 1775, 1776.—Eaton & Wright, Bot. 385.—Darby, Bot. S. States, 510.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 36.—Liebmann, Chênes Am. Trop. t. D.—Wood, Cl. Book, 643.—Vasey, Cat. Forest Trees, 26.—Engelmann in Trans. St. Louis Acad. iii, 385, 395.

Q. laurifolia hybrida, Michaux, Hist. Chênes Am. No. 10, t. 18.

Q. laurifolia, var. *obtusa*, Willdenow, Spec. iv, 428.—Aiton, Hort. Kew. 2 ed. v, 288.—Wood, Cl. Book, 343.

Q. laurifolia, var. *acuta*, Willdenow, Spec. iv, 428.—Aiton, Hort. Kew. 2 ed. v, 288.

Q. obtusa, Pursh, Fl. Am. Sept. ii, 627.

Q. Phellos, var. *laurifolia*, Chapman, Fl. S. States, 420.—Wood, Bot. & Fl. 305.—Young, Bot. Texas, 502.

Q. aquatica, var. *laurifolia*, A. De Candolle, Prodr. xvi³, 68.

LAUREL OAK.

North Carolina, south near the coast to Mosquito inlet and cape Romano, Florida, and along the Gulf coast to the shores of Mobile bay.

A large tree, 18 to 24 meters in height, with a trunk 0.90 to 1.20 meter in diameter; most common and reaching its greatest development on the rich hummocks of the Florida coast.

Wood heavy, very strong and hard, coarse-grained, inclined to check in drying; layers of annual growth marked by several rows of rather small open ducts; medullary rays broad, conspicuous; color, dark brown tinged with red, the sap-wood lighter; specific gravity, 0.7673; ash 0.82.

282.—*Quercus heterophylla*, Michaux f.

Hist. Arb. Am. ii, 87, t. 16; N. American Sylva, 3 ed. i, 64, t. 18.—Pursh, Fl. Am. Sept. ii, 627.—Barton, Compend. Fl. Philadelph. ii, 167.—Nuttall, Genera, ii, 214; Sylva, i, 15; 2 ed. i, 24.—Green in Universal Herbal, ii, 442.—Torrey, Compend. Fl. N. States, 357.—Sweet, Cat. 2 ed. 466.—Beck, Bot. 328.—Eaton, Manual, 6 ed. 292.—London, Arboretum, iii, 1894.—Eaton & Wright, Bot. 383.—Gale in Proc. Nat. Inst. 1855, 70, f. 1.—Wood, Cl. Book, 645.—Buckley in Proc. Philadelphia Acad. 1862, 361; 1862, 100.—Gray, Hall's Pl. Texas, 21.—Liebmann, Chênes Am. Trop. t. B.—Meehan in Proc. Philadelphia Acad. 1875, 437, 465; Coulter's Bot. Gazette, vii, 10.—Leidy in Proc. Philadelphia Acad. 1875, 415.—Engelmann in Trans. St. Louis Acad. iii, 385, 391.—Martindale, Notes on the Bartram Oak, 3; Coulter's Bot. Gazette, vi, 303.—Ward in Bull. U. S. Nat. Mus. No. 22, 114.

Q. aquatica, var. *heterophylla*, Aiton, Hort. Kew. 2 ed. v, 290.—A. De Candolle, Prodr. xvi^a, 68.

Q. nigra, var. Cooper in Smithsonian Rep. 1858, 255.

Q. Phellos × *tinctoria*, Gray, Manual N. States, 4 ed. 406.

Q. Phellos, var. Gray, Manual N. States, 5 ed. 453.

Q. Phellos × *coccinea*, Engelmann in Trans. St. Louis Acad. iii, 541.

BARTRAM'S OAK.

New Jersey, Salem and Cumberland counties, "restricted to a line or belt bordering extreme tidal points of streams entering the Delaware river where the alluvial terminates and the upland commences," (*Commons*); Delaware, near Townsend station and Wilmington; North Carolina (*M. A. Curtis* in herb. *Canby*); eastern Texas (*E. Hall*); this perhaps *Q. Durandii*.

A small tree, 12 to 15 meters in height, with a trunk 0.45 to 0.60 meter in diameter; rare and very local.

Wood heavy, hard, very strong, close-grained, compact; layers of annual growth marked by several rows of small open ducts; medullary rays numerous, conspicuous; color, light brown tinged with red, the sap-wood somewhat darker; specific gravity, 0.6834; ash, 0.17.

283.—*Quercus cinerea*, Michaux,

Hist. Chênes Am. No. 8, t. 14; Fl. Bor.-Am. ii, 197.—Willdenow, Spec. iv, 425.—Persoon, Syn. ii, 567.—Poiret, Suppl. ii, 212.—Michaux f. Hist. Arb. Am. ii, 82, t. 14; N. American Sylva, 3 ed. i, 61, t. 16.—Aiton, Hort. Kew. 2 ed. v, 288.—Pursh, Fl. Am. Sept. ii, 626.—Smith in Rees' Cycl. xxx, No. 6.—Nuttall, Genera, ii, 214.—Nouveau Duhamel, vii, 151.—Elliott, Sk. ii, 594.—Sprengel, Syst. iii, 857.—Eaton, Manual, 6 ed. 294.—Eaton & Wright, Bot. 6 ed. 294.—Engelmann & Gray in Jour. Boston Soc. Nat. Hist. v, 262.—Scheele in Rømer, Texas, 446.—Cooper in Smithsonian Rep. 1858, 255.—Chapman, Fl. S. States, 421.—Curtis in Rep. Geological Surv. N. Carolina, 37.—Wood, Cl. Book, 643; Bot. & Fl. 305.—A. De Candolle, Prodr. xvi^a, 73.—Ørsted in Saerskitt. Aftryk. af Nat. For. Viden. Meddelt. Nos. 1-6, 1866, 73.—Gray, Manual N. States, 5 ed. 452; Hall's Pl. Texas, 21.—Young, Bot. Texas, 502.—Koch, Dendrologie, ii^a, 58.—Vasey, Cat. Forest Trees, 26.—Engelmann in Trans. St. Louis Acad. iii, 385, 395.

Q. Prinus, β. Linnaeus, Spec. 1 ed. 995.

Q. humilis, Walter, Fl. Caroliniana, 234.

Q. Phellos, var. *cinerea*, Aiton, Hort. Kew. iii, 354.—London, Arboretum, iii, 1895, f. 1773.—Spach, Hist. Veg. xi, 161.

UPLAND WILLOW OAK. BLUE JACK. SAND JACK.

North Carolina, south near the coast to cape Malabar and Pease creek, Florida, west along the Gulf coast to the valley of the Brazos river, Texas, extending north through eastern Texas to about latitude 33°.

A tree 9 to 15 meters in height, with a trunk rarely exceeding 0.20 meter in diameter; sandy barrens and dry upland ridges.

Wood heavy, hard, strong, close grained, compact; layers of annual growth marked by several rows of not large open ducts; medullary rays distant, thin, conspicuous; color, light brown tinged with red, the sap-wood darker; specific gravity, 0.6420; ash, 1.21.

284.—*Quercus hypoleuca*, Engelmann,

Trans. St. Louis Acad. iii, 384; Wheeler's Rep. vi, 251.—Vasey, Cat. Forest Trees, 26.—Rusby in Bull. Torrey Bot. Club, ix, 78.

Q. confertifolia, Torrey, Bot. Mex. Boundary Survey, 207 [not HBK.].—Cooper in Smithsonian Rep. 1858, 261.

Limpia mountains, Texas (*Havard*), valleys of the high mountain ranges of southwestern New Mexico, Santa Rita mountains, Arizona, above 6,000 feet elevation; southward into Sonora.

A small evergreen tree of great beauty, 9 to 15 meters in height, with a trunk sometimes 0.75 meter in diameter; dry, gravelly slopes and summits, the large specimens hollow and defective.

Wood heavy, very strong and hard, close-grained, compact; layers of annual growth marked by few small open ducts; medullary rays broad, conspicuous; color, dark brown, the sap-wood much lighter; specific gravity, 0.8009; ash, 1.34.

285.—*Quercus imbricaria*, Michaux,

Hist. Chênes Am. No. 9, t. 15, 16; Fl. Bor.-Am. ii, 197.—Willdenow, Spec. iv, 428; Enum. Suppl. 64; Berl. Baumz. 338.—Persoon, Syn. ii, 567.—Poiret, Suppl. ii, 214.—Michaux f. Hist. Arb. Am. ii, 78, t. 13; N. American Sylva, 3 ed. i, 60, t. 15.—Aiton, Hort. Kew. 2 ed. v, 288.—Smith in Rees' Cycl. xxx, No. 15.—Pursh, Fl. Am. Sept. ii, 627.—Nuttall, Genera, ii, 214.—Barton, Compend. Fl. Philadelph. ii, 167.—Nouveau Duhamel, vii, 153.—Hayne, Dend. Fl. 155.—Elliott, Sk. ii, 598.—Sprengel, Syst. iii, 857.—Torrey, Compend. Fl. N. States, 357.—Beck, Bot. 328.—Eaton, Manual, 6 ed. 292.—London, Arboretum, iii, 1898, f. 1777.—Eaton & Wright, Bot. 383.—Darby, Bot. S. States, 510.—Torrey & Gray in Pacific R. R. Rep. ii, 130.—Cooper in Smithsonian Rep. 1858, 255.—Brendel in Trans. Illinois Ag. Soc. iii, 623, t. 6.—Chapman, Fl. S. States, 420.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 36.—Lesquereux in Owen's 2d Rep. Arkansas, 388.—Wood, Cl. Book, 643; Bot. & Fl. 305.—A. De Candolle, Prodr. xvi³, 63.—Ørsted in Saerskitt. Aftryk. af Nat. For. Viden. Meddelt. Nos. 1-6, 1860, 73.—Gray, Manual N. States, 5 ed. 452.—Young, Bot. Texas, 502.—Liebmann, Chênes Am. Trop. t. D, t. xxii, f. 5.—Koch, Dendrologie, ii³, 60.—Vasey, Cat. Forest Trees, 26.—Broadhead in Coulter's Bot. Gazette, iii, 60.—Ridgway in Proc. U. S. Nat. Mus. 1882, 80.

Q. Phellos, var. *imbricaria*, Spach, Hist. Veg. xi, 160.

SHINGLE OAK. LAUREL OAK.

Allentown, Lehigh county, Pennsylvania (*Porter*), west through southern Michigan, southern Wisconsin, and southeastern Iowa to southeastern Nebraska and northeastern Kansas, south to northern Georgia and Alabama, middle Tennessee, and northern Arkansas.

A tree 24 to 30 meters in height, with a trunk 0.60 to 0.90 meter in diameter; rich woodlands.

Wood heavy, hard, rather coarse-grained, checking badly in drying; layers of annual growth marked by many rows of large open ducts; medullary rays broad, conspicuous; color, light brown tinged with red, the sap-wood much lighter; specific gravity, 0.7529; ash, 0.43; occasionally used for clapboards, shingles, etc.

286.—*Quercus Phellos*, Linnaeus,

Spec. 1 ed. 994.—Lamarck, Dict. i, 722.—Wangenheim, Amer. 76, t. 5, f. 11.—Walter, Fl. Caroliniana, 234.—Aiton, Hort. Kew. iii, 354; 2 ed. v, 287.—Abbot, Insects Georgia, ii, t. 52, 91.—Michaux, Fl. Bor.-Am. ii, 197.—Willdenow, Spec. iv, 423; Enum. 974; Berl. Baumz. 337.—Smith in Rees' Cycl. xxx, No. 7.—Persoon, Syn. ii, 567.—Desfontaines, Hist. Arb. ii, 507.—Michaux f. Hist. Arb. Am. ii, 75, t. 12; N. American Sylva, 3 ed. i, 58, t. 14.—Pursh, Fl. Am. Sept. ii, 625.—Barton, Prodr. Fl. Philadelph. 91; Compend. Fl. Philadelph. ii, 167.—Nuttall, Genera, ii, 214; Sylva, i, 15; 2 ed. i, 17.—Nouveau Duhamel, vii, 150.—Hayne, Dend. Fl. 155.—Elliott, Sk. ii, 593.—Sprengel, Syst. iii, 857.—Torrey, Compend. Fl. N. States, 357; Fl. N. York, ii, 187.—Beck, Bot. 328.—Eaton, Manual, 6 ed. 383.—London, Arboretum, iii, 1894, f. 1774 & t.—Eaton & Wright, Bot. 383.—Spach, Hist. Veg. xi, 160.—Penn. Cycl. xix, 216.—Darby, Bot. S. States, 509.—Cooper in Smithsonian Rep. 1858, 255.—Chapman, Fl. S. States, 420.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 36.—Lesquereux in Owen's 2d Rep. Arkansas, 388.—Wood, Cl. Book, 643; Bot. & Fl. 305.—A. De Candolle, Prodr. xvi³, 63.—Ørsted in Saerskitt. Aftryk. af Nat. For. Viden. Meddelt. Nos. 1-6, 1866, 73.—Gray, Manual N. States, 5 ed. 452; Hall's Pl. Texas, 21.—Young, Bot. Texas, 502.—Koch, Dendrologie, ii³, 59.—Vasey, Cat. Forest Trees, 26.—Gartenflora, xxix, 221 & f.—Ridgway in Proc. U. S. Nat. Mus. 83.

Q. Phellos angustifolia, Marshall, Arbustum, 124.

Q. Phellos latifolia, Marshall, Arbustum, 124.—Loddiges, Cat. ed. 1836.—London, Arboretum, iii, 1895 & t.

Q. Phellos, var. *viridis*, Aiton, Hort. Kew. iii, 354.

Q. Phellos, var. *humilis*, Pursh, Fl. Am. Sept. ii, 625.

WILLOW OAK. PEACH OAK.

Tottenville, Staten island, New York, south near the coast to northeastern Florida, through the Gulf states to the valley of the Sabine river, Texas, and through Arkansas to southeastern Missouri, Tennessee, and southern Kentucky.

A tree 18 to 24 meters in height, with a trunk sometimes 0.90 meter in diameter; bottom lands or rich sandy uplands.

Wood heavy, strong, not hard, rather close-grained, compact; layers of annual growth marked by several rows of small open ducts; medullary rays few, distant; color, light brown tinged with red, the sap-wood lighter red; specific gravity 0.7472; ash, 0.50; somewhat used for fellies of wheels, clapboards, in construction, etc.

287.—*Quercus densiflora*, Hooker & Arnott,

Bot. Beechey, 391.—Hooker, Icon. iv, t. 380.—Bentham, Pl. Hartweg. 337.—Nuttall, Sylva, i, 11, t. 5; 2 ed. i, 21, t. 5.—Torrey in Pacific R. R. Rep. iv, 138.—Bot. Wilkes Exped. 458.—Newberry in Pacific R. R. Rep. vi, 31, 89, f. 8.—A. De Candolle, Prodr. xvi, 82.—Bolander in Proc. California Acad. iii, 231.—Vasey, Cat. Forest Trees, 25.—Engelmann in Trans. St. Louis Acad. iii, 389; Bot. California, ii, 99.

Q. echinacea, Torrey in Pacific R. R. Rep. iv, 137, t. 14.

Pasania densiflora, Örsted in Saerskitt. Aftryk. af Nat. For. Viden. Meddelt. No. 1-6, 1866, 73.

Q. echinoides, R. Brown Campet. in Ann. & Mag. Nat. Hist. April, 1871, 2.

TANBARK OAK. CHESTNUT OAK. PEACH OAK.

Valley of the Umpqua river, Oregon, south through the Coast ranges to the Santa Lucia mountains, California. A tree 18 to 24 meters in height, with a trunk 0.60 to 0.90 meter in diameter; rich valleys and banks of streams; most common and reaching its greatest development in the redwood forests of the California coast.

Wood heavy, hard, strong, very close-grained, compact, containing broad bands of small open ducts parallel to the thin, dark, conspicuous medullary rays; color, bright reddish-brown, the thick sap-wood darker brown; specific gravity, 0.6827; ash, 1.49; largely used as fuel.

The bark, rich in tannin, very largely used and preferred to that of any other tree of the Pacific forests for tanning.

NOTE.—The following shrubby species of *Quercus* do not properly find a place in this catalogue:

Quercus undulata, Torrey in Ann. Lyc. N. York, ii, 248, t. 4.
Interior Pacific region from Colorado southward.

Quercus Breweri, Engelmann in Bot. California, ii, 96.
Q. lobata, var. *fruticosa*, Engelmann in Trans. St. Louis Acad. iii, 388.
Western slopes of the high Sierra Nevadas, California.

Quercus Georgiana, M. A. Curtis in Chapman's Fl. S. States.
Stone Mountain, Georgia.

Quercus myrtifolia, Willdenow, Sp. iv, 424.
Q. Phellos, var. *arcuaria*, Chapman, Fl. S. States, 420.
Q. aquatica, var. *myrtifolia*, A. De Candolle, Prodr. xvi, 68.

South Atlantic and Gulf coast.

Quercus ilicifolia, Wangenheim, Amer. 79, t. 6, f. 17.
Q. Lanisteri, Michaux, Hist. Chênes Am. t. 27.

North Atlantic region.

Quercus pumila, Walter, Fl. Caroliana, 234.
Q. Phellos pumila, Michaux, Hist. Chênes Am. t. 15, f. 1.
Q. cinerea, var. *pumila*, Chapman, Fl. S. States, 421.—A. De Candolle, Prodr. 16, 74.
Q. cinerea, var. *sericea*, Engelmann in Trans. St. Louis Acad. iii, 394.
Q. sericea, Willdenow, Spec. 424.
Q. Phellos, var. *sericea*, Alton, Hort. Kew. iii, 354.

Pine barrens, South Carolina.

Quercus dumosa, Nuttall, Sylva, i, 7.
Q. berberidifolia, Liebm. in Danek. Vidensk. Selsk. Forhandl. 1854, 172, in part.
Q. dumosa, var. *bullata*, Engelmann in Bot. California, 298.
Q. acutidens, Torrey, Bot. Mex. Boundary Survey, 297, t. 51.

Coast ranges of southern California.

Numerous hybrid or supposed hybrid oaks, variously described by American botanists, are not properly considered here.

288.—*Castanopsis chrysophylla*, A. De Candolle;

Seemann's Jour. Bot. i, 182; Prodr. xvi^s, 109.—Watson in King's Rep. v, 322; Bot. California, ii, 100.—Gray in Proc. Am. Acad. vii, 401.—Torrey, Bot. Wilkes Exped. 463.—Vasey, Cat. Forest Trees, 27.—Hall in Coulter's Bot. Gazette, ii, 91.

Castanea chrysophylla, Douglas in Hooker's London Jour. Bot. ii, 496, t. 16.—Bentham, Pl. Hartweg. 337.—Hooker, Fl. Bor.-Am. ii, 159.—Nuttall, Sylva, i, 21; 2 ed. i, 37.—Bot. Mag. t. 4953.—Torrey in Pacific R. R. Rep. iv, 137; Bot. Mex. Boundary Survey, 205.—Morren in Belg. Hort. vii, 248, t. 240.—Newberry in Pacific R. R. Rep. vi, 26, 89, f. 4.—Fl. des Serres, xii, 3, t. 1184.—Cooper in Smithsonian Rep. 1858, 261.—Kellogg in Proc. California Acad. ii, 280.—Bolander in Proc. California Acad. iii, 231.—Engelmann in Wheeler's Rep. vi, 375.—Shingles in London Gard. Chronicle, 1882, 716.

Castanea chrysophylla, var. *minor*, Bentham, Pl. Hartweg. 337.

Castanea sempervirens, Kellogg in Proc. California Acad. i, 71.

C. chrysophylla, var. *minor*, A. De Candolle, Prodr. xvi^s, 110.

C. chrysophylla, var. *pumila*, Vasey, Cat. Forest Trees, 27.

CHINQUAPIN.

Cascade mountains, Oregon, below 4,000 feet elevation, south along the western slopes of the Sierra Nevada, and through the California Coast ranges to the San Bernardino and San Jacinto mountains.

A tree 15 to 24 meters in height, with a trunk 0.30 to 0.90 meter in diameter, or at high elevations and toward its southern limits reduced to a low shrub; most common and reaching its greatest development in the Coast Range valleys of northern California; at its southern limits rarely below 10,000 feet elevation.

Wood light, soft, not strong, close-grained, compact; layers of annual growth marked by a single row of rather large open ducts; medullary rays numerous, obscure; color, light brown tinged with red, the sap-wood lighter; specific gravity, 0.5574; ash, 0.35; in southern Oregon occasionally used in the manufacture of plows and other agricultural implements.

289.—*Castanea pumila*, Miller,

Diet. No. 2.—Lamarck, Dict. i, 708.—Michaux, Fl. Bor.-Am. ii, 193.—Willdenow, Spec. iv, 461; Enum. 980; Berl. Baumz. 78.—Smith in Rees' Cycl. xiv, No. 2.—Nouveau Duhamel, iii, 79.—Persoon, Syn. ii, 572.—Desfontaines, Hist. Arb. ii, 500.—Michaux f. Hist. Arb. Am. ii, 166, t. 7; N. American Sylva, 3 ed. iii, 16, t. 105.—Aiton, Hort. Kew. 2 ed. v, 298.—Pursh, Fl. Am. Sept. ii, 624.—Rafinesque, Fl. Ludoviciana, 159; New Fl. & Bot. i, 83.—Nuttall, Genera, ii, 217; Am. Phil. Soc. 2 ser. v, 168.—Hayne, Dend. Fl. 165.—James in Long's Exped. ii, 287.—Elliott, Sk. ii, 615.—Torrey, Compend. Fl. N. States, 355; Fl. N. York, ii, 196.—Audubon, Birds, t. 85.—Beck, Bot. 332.—Eaton, Manual, 6 ed. 84.—Penn. Cycl. vi, 350.—London, Arboretum, iii, 2002, f. 1927, 1928.—Eaton & Wright, Bot. 184.—Spach, Hist. Veg. xi, 192.—Darlington, Fl. Cestrica, 3 ed. 270.—Darby, Bot. S. States, 512.—Cooper in Smithsonian Rep. 1858, 256.—Chapman, Fl. S. States, 424.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 47.—Lesquereux in Owen's 2d Rep. Arkansas, 388.—Wood, Cl. Book, 646; Bot. & Fl. 307.—Porcher, Resources S. Forests, 237.—A. De Candolle, Prodr. xvi^s, 115.—Gray, Manual N. States, 5 ed. 455.—Young, Bot. Texas, 508.—Koch, Dendrologie, ii^s, 24.—Vasey, Cat. Forest Trees, 27.—Butler in Coulter's Bot. Gazette, iii, 17.

Fagus pumila, Linnæus, Spec. 1 ed. 998.—Du Roi, Harbk. i, 175.—Wangenheim, Amer. 57, t. 19, f. 44.—Walter, Fl. Caroliniana, 233.—Aiton, Hort. Kew. iii, 361.—Abbot, Insects Georgia, ii, t. 57.

Fagus Castanea pumila, Marshall, Arbustum, 47.

Fagus pumila, var. *præcox*, Walter, Fl. Caroliniana, 233.

C. nana, Muhlenberg, Cat. 86.—Elliott, Sk. ii, 615.—Rafinesque, New Fl. & Bot. i, 83.—Darby, Bot. S. States, 512.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 47.—Lesquereux in Owen's 2d Rep. Arkansas, 388.

C. alnifolia, Nuttall, Genera, ii, 217; Sylva, i, 19, t. 6; 2 ed. i, 36, t. 6.

C. vesca, Lesquereux in Owen's 2d Rep. Arkansas, 388 [not Gärtner].

CHINQUAPIN.

Lancaster county, Pennsylvania, and the valley of the lower Wabash river, Indiana, south and southwest to northern Florida and the valley of the Neches river, Texas.

A tree sometimes 15 meters in height, with a trunk 0.30 to 1.05 meter in diameter, or often, especially in the Atlantic states, reduced to a low shrub; rich hillsides and borders of swamps; most common and reaching its greatest development in southern Arkansas.

Wood light, hard, strong, coarse-grained, durable in contact with the ground, liable to check in drying; layers of annual growth marked by many rows of large open ducts; medullary rays numerous, obscure; color, dark brown, the sap-wood hardly distinguishable; specific gravity, 0.5887; ash, 0.12; used for posts, rails, railway ties, etc.

The small nuts sweet and edible.

290.—*Castanea vulgaris*, var. *Americana*, A. De Candolle,

Prodr. xvi², 114.—Schneck in Coulter's Bot. Gazette, vi, 159.—Bell in Geological Rep. Canada, 1879-'80, 53^c.—Ridgway in Proc. U. S. Nat. Mus. 1882, 84.

Fagus Castanea dentata, Marshall, Arbustum, 46.

Fagus Castanea, Wangenheim, Amer. 47 [not Linnaeus].—Walter, Fl. Caroliniana, 233.—Aiton, Hort. Kew. iii, 361, in part.—Lamarek, Ill. iii, 366, t. 782, in part.

C. vesca, var. *Americana*, Michaux, Fl. Bor.-Am. ii, 193.—Persoon, Syn. ii, 572.—Barton, Prodr. Fl. Philadelph. 90.—Pursh, Fl. Am. Sept. ii, 624.—Eaton, Manual, 109; 6 ed. 84.—Nuttall, Genera, ii, 217.—Elliott, Sk. ii, 614.—Torrey, Compend. Fl. N. States, 355; Fl. N. York, ii, 195, t. 111.—London, Arboretum, iii, 1984.—Eaton & Wright, Bot. 184.—Emerson, Trees Massachusetts, 164, 2 ed. i, 187 & t.—Porcher, Resources S. Forests, 238.—Vasey, Cat. Forest Trees, 27.—Rudkin in Bull. Torrey Bot. Club, vii, 81.

C. Americana, Rafinesque, New Fl. & Bot. i, 82.—Willdenow, Enum. Suppl. 64.—Nuttall, Sylva, i, 24; 2 ed. i, 38.—Spach, Hist. Veg. xi, 191.—Cooper in Smithsonian Rep. 1858, 256.—Koch, Dendrologie, ii², 23.—Martindale in Proc. Philadelphia Acad. 1880, 2.

C. vesca, Willdenow, Spec. iv, 460, in part.—Desfontaines, Hist. Arb. ii, 500, in part.—Michaux f. Hist. Arb. Am. ii, 151, t. 6; N. American Sylva, 3 ed. iii, 11, t. 104 [not Gærtner].—Hayne, Dend. Fl. 165, in part.—Sprengel, Syst. iii, 856, in part.—Beck, Bot. 332.—Penn. Cycl. vi, 350.—Bigelow, Fl. Boston. 3 ed. 224.—Darlington, Fl. Cestricea, 3 ed. 270.—Darby, Bot. S. States, 511.—Chapman, Fl. S. States, 424.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 46.—Wood, Cl. Book, 646; Bot. & Fl. 306.—Gray, Manual N. States, 5 ed. 455.

CHESTNUT.

Southern Maine to the valley of the Winooski river, Vermont, southern Ontario and southern Michigan, south through the northern states to Delaware and southern Indiana, and along the Alleghany mountains to northern Alabama, extending west to middle Kentucky and Tennessee.

A large tree, 24 to 30 meters in height, with a trunk 1.80 to 4 meters in diameter; rich woods and hillsides; very common and reaching its greatest development on the western slopes of the southern Alleghany mountains.

Wood light, soft, not strong, coarse-grained, liable to check and warp in drying, easily split, very durable in contact with the soil; layers of annual growth marked by many rows of large open ducts; medullary rays numerous, obscure; color, brown, the sap-wood lighter; specific gravity, 0.4504; ash, 0.18; largely used in cabinet-making, for railway ties, posts, fencing, etc.

The fruit, although smaller, superior in sweetness and flavor to that of the European chestnut.

An infusion or fluid extract of the dried leaves is successfully employed in the treatment of whooping-cough and other pectoral affections (*U. S. Dispensatory*, 14 ed. 245.—*Nat. Dispensatory*, 2 ed. 364).

291.—*Fagus ferruginea*, Aiton,

Hort. Kew. iii, 362; 2 ed. v, 298.—Abbot, Insects Georgia, ii, t. 75.—Willdenow, Spec. iv, 460; Enum. 980; Berl. Baumz. 140.—Persoon, Syn. ii, 571.—Desfontaines, Hist. Arb. ii, 496.—Michaux f. Hist. Arb. Am. ii, 174, t. 9; N. American Sylva, 3 ed. iii, 21, t. 106.—Smith in Rees' Cycl. xiv, No. 4.—Pursh, Fl. Am. Sept. ii, 624.—Barton, Prodr. Fl. Philadelph. 90; Compend. Fl. Philadelph. ii, 174.—Eaton, Manual, 108; 6 ed. 145.—Sprengel, Syst. iii, 856.—Torrey, Compend. Fl. N. States, 354; Fl. N. York, ii, 194, t. 110.—Beck, Bot. 333.—Eaton, Manual, 6 ed. 145.—London, Arboretum, iii, 1980, f. 1917.—Hooker, Fl. Bor.-Am. ii, 159.—Eaton & Wright, Bot. 244.—Bigelow, Fl. Boston. 3 ed. 374.—Darlington, Fl. Cestricea, 3 ed. 271.—Cooper in Smithsonian Rep. 1858, 256.—Chapman, Fl. S. States, 425.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 47.—Wood, Bot. & Fl. 307.—A. De Candolle, Prodr. xvi², 118.—Gray, Manual N. States, 5 ed. 455.—Koch, Dendrologie, ii², 19.—Vasey, Cat. Forest Trees, 27.—Broadhead in Coulter's Bot. Gazette, iii, 60.—Sears in Bull. Essex Inst. xiii, 179.—Bell in Geological Rep. Canada, 1879-'80, 52^c.—Ridgway in Proc. U. S. Nat. Mus. 1882, 85.

F. sylvatica atropunicea, Marshall, Arbustum, 46.

F. Americana latifolia, Wangenheim, Amer. 80, t. 29, f. 55.—London, Arboretum, iii, 1960, f. 1916.

F. sylvatica, Walter, Fl. Caroliniana, 233 [not Linnaeus].—Pursh, Fl. Am. Sept. ii, 624.—Beck, Bot. 333.—Darlington, Fl. Cestricea, 2 ed. 538.—Darby, Bot. S. States, 512.

F. sylvestris, Michaux, Fl. Bor. Am. ii, 194.—Michaux f. Hist. Arb. Am. ii, 170, t. 8; N. American Sylva, 3 ed. iii, 18, t. 107.—Hooker, Fl. Bor.-Am. ii, 159.—Lesquereux in Owen's 2d Rep. Arkansas, 388.

F. alba, Rafinesque, Fl. Ludoviciana, 131.

F. sylvatica, var. *Americana*, Nuttall, Genera, ii, 216.—Barton, Compend. Fl. Philadelph. ii, 174.—Elliott, Sk. ii, 613.—Eaton, Manual, 6 ed. 145.—London, Arboretum, iii, 1953.—Eaton & Wright, Bot. 244.—Emerson, Trees Massachusetts, 153; 2 ed. i, 180 & t.—Wood, Cl. Book, 647.—Porcher, Resources S. Forests, 235.

F. Americana, Sweet, Hort. Brit.—Spach, Hist. Veg. xi, 201.

F. ferruginea, var. *Caroliniana*, London, Arboretum, iii, 1980, f. 1915.

BEECH.

Nova Scotia and the valley of the Restegouche river to the northern shores of lake Huron and northern Wisconsin, south to the Chattahoochee region of western Florida and the valley of the Trinity river, Texas, west to eastern Illinois, southeastern Missouri, and Madison county, Arkansas (*Letterman*).

A large tree, 24 to 30 or, exceptionally, 34 meters (*Ridgway*) in height, with a trunk 0.90 to 1.20 meter in diameter; rich woods, or at the south sometimes in bottom lands or the dryer portions of swamps, reaching its greatest development upon the "bluff" formations of the lower Mississippi basin; very common.

Wood very hard, strong, tough, very close grained, not durable in contact with the soil, inclined to check in drying, difficult to season, susceptible of a beautiful polish; medullary rays broad, very conspicuous; color, varying greatly with soil and situation, dark red, or often lighter, the sap-wood nearly white; specific gravity, 0.6883; ash, 0.51; largely used in the manufacture of chairs, shoe-last, plane-stocks, handles, etc., and for fuel.

292.—*Ostrya Virginica*, Willdenow,

Spec. iv, 469; Enum. 982; Berl. Baumz. 260.—Persoon, Syn. ii, 573.—Aiton, Hort. Kew. 2 ed. v, 302.—Pursh, Fl. Am. Sept. ii, 623.—Eaton, Manual, 109; 6 ed. 244.—Nuttall, Genera, ii, 219.—Hayne, Dend. Fl. 169.—Elliott, Sk. ii, 618.—Sprengel, Syst. iii, 856.—Torrey, Compend. Fl. N. States, 356; Nicotlet's Rep. 160; Fl. N. York, ii, 185, t. 102.—Audubon, Birds, t. 40.—London, Arboretum, iii, 2015, f. 1940.—Hooker, Fl. Bor.-Am. ii, 160.—Eaton & Wright, Bot. 336.—Bigelow, Fl. Boston. 3 ed. 383.—Spach in Ann. Sci. Nat. 2 ser. xvi, 246; Hist. Veg. xi, 218.—Emerson, Trees Massachusetts, 177; 2 ed. i, 201 & t.—Parry in Owen's Rep. 618.—Darlington, Fl. Cestrica, 3 ed. 274.—Darby, Bot. S. States, 509.—Cooper in Smithsonian Rep. 256.—Chapman, Fl. S. States, 426.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 75.—Lesquereux in Owen's 2d Rep. Arkansas, 388.—Wood, Cl. Book, 647; Bot. & Fl. 307.—Porcher, Resources S. Forests, 233.—A. De Candolle, Prodr. xvi², 125.—Gray, Manual N. States, 5 ed. 456.—Young, Bot. Texas, 510.—Vasey, Cat. Forest Trees, 27.—Sargent in Am. Nat. xi, 683.—Sears in Bull. Essex Inst. xiii, 179.—Ridgway in Proc. U. S. Nat. Mus. 85.

Carpinus Ostrya, Linnaeus, Spec. 1 ed. 998, in part.—Du Roi, Harbk. i, 130.—Wangenheim, Amer. 48.—Marshall, Arbustum, 25.—Möench, Meth. 694.—Abbot, Insecta Georgia, ii, t. 76.—Nouveau Duhamel, ii, 200.—Michaux f. Hist. Arb. Am. iii, 53, t. 7; N. American Sylva, 3 ed. iii, 27, t. 109.

Carpinus Virginiana, Miller, Dict. 7 ed. No. 4.—Lamarck, Dict. i, 708; Wangenheim, Amer. 49.—Nouveau Duhamel, ii, 201.—Desfontaines, Hist. Arb. ii, 493.—Smith in Rees' Cycl. vii, No. 5.

Carpinus triflora, Möench, Meth. 394.

Carpinus Ostrya, var. *Americana*, Michaux, Fl. Bor.-Am. ii, 202.

O. Virginica, var. *glandulosa*, Spach in Ann. Sci. Nat. 2 ser. xvi, 246; Hist. Veg. xi, 218.

O. Virginica, var. *eglandulosa*, Spach in Ann. Sci. Nat. 2 ser. xvi, 246; Hist. Veg. xi, 218.

O. Virginiana, Koch, Dendrologie, ii², 6.

HOP-HORNBEAM. IRON WOOD. LEVER WOOD.

Bay of Chaleur, through the valleys of the Saint Lawrence and the lower Ottawa rivers, along the northern shore of lake Huron to northern Minnesota, south through the northern states and along the Alleghany mountains to the Chattahoochee region of western Florida, and through eastern Iowa, southeastern Missouri, and Arkansas to eastern Kansas, the Indian territory, and eastern Texas.

A small tree, 9 to 15 meters in height, with a trunk 0.30 to 0.60 meter in diameter; generally on dry, gravelly hillsides and knolls; reaching its greatest development in southern Arkansas; common.

Wood heavy, very strong and hard, tough, very close-grained, compact, susceptible of a beautiful polish, very durable in contact with the soil; medullary rays numerous, obscure; color, light brown tinged with red, or, like the sap-wood, often nearly white; specific gravity, 0.8284; ash, 0.50; used for posts, levers, handles of tools, etc.

293.—*Carpinus Caroliniana*, Walter,

Fl. Caroliniana, 238.—A. De Candolle, Prodr. xvi², 126.—Koch, Dendrologie, ii², 4.—Sears in Bull. Essex Inst. xviii, 180.—Ridgway in Proc. U. S. Nat. Mus. 1882, 85.

O. Americana, Lamarck, Dict. iv, 708; Suppl. ii, 202.—Michaux, Fl. Bor.-Am. ii, 201.—Willdenow, Spec. iv, 468; Enum. Suppl. 64; Berl. Baumz. 75.—Persoon, Syn. ii, 573.—Michaux f. Hist. Arb. Am. iii, 57, t. 8; N. American Sylva, 3 ed. iii, 26, t. 108.—Pursh, Fl. Am. Sept. ii, 623.—Aiton, Hort. Kew. 2 ed. v, 301.—Eaton, Manual, 109; 6 ed. 82.—Barton, Prodr. Fl. Philadelph. 91; Compend. Fl. Philadelph. ii, 176.—Nuttall, Genera, ii, 218.—Hayne, Dend. Fl. 168.—Elliott, Sk. ii, 618.—Watson, Dend. Brit. ii, t. 157.—Sprengel, Syst. iii, 854.—Guimpel, Otto & Hayne, Abb. Holz. 107, t. 84.—Torrey, Compend. Fl. N. States, 356; Fl. N. York, ii, 185, t. 103.—Penn. Cycl. iv, 315.—London, Arboretum, iii, 2013, f. 1936.—Hooker, Fl. Bor.-Am. ii, 160.—Eaton & Wright, Bot. 182.—Bigelow, Fl. Boston. 3 ed. 383.—Spach in Ann. Sci. Nat. 2 ser. xvi, 252; Hist. Veg. xi, 224.—Emerson, Trees Massachusetts, 174; 2 ed. i, 198 & t.—Parry in Owen's Rep. 618.—Darlington, Fl. Cestrica, 3 ed. 273.—Darby, Bot. S. States, 508.—Cooper in Smithsonian Rep. 1858, 256.—Chapman, Fl. S. States, 425.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 75.—Lesquereux in Owen's 2d Rep. Arkansas, 388.—Wood, Cl. Book, 648; Bot. & Fl. 307.—Gray, Manual N. States, 5 ed. 457; Hall's Fl. Texas, 21.—Young, Bot. Texas, 509.—Vasey, Cat. Forest Trees, 27.—Broadhead in Coulter's Bot. Gazette, iii, 60.—Bell in Geological Rep. Canada, 1879-'80, 52.

O. Betulus Virginiana, Marshall, Arbustum, 25.

HORNBEAM. BLUE BEECH. WATER BEECH. IRON WOOD.

Nova Scotia, southern New Brunswick, northern shores of Georgian bay, southern peninsula of Michigan to northern Minnesota (lake Pokegama, *Garrison*), south to cape Malabar and Tampa bay, Florida, and the valley of the Trinity river, Texas, west to central Iowa, eastern Kansas, and the valley of the Poteau river, Indian territory.

A small tree, 9 to 15 meters in height, with a trunk sometimes 0.60 to 0.90 meter in diameter, or at the north much smaller and often reduced to a low shrub; borders of streams and swamps, in moist soil; most common and reaching its greatest development along the western slopes of the southern Alleghany mountains and in southern Arkansas and eastern Texas.

Wood heavy, very strong and hard, close-grained, inclined to check in drying; medullary rays numerous, broad; color, light brown, the thick sap-wood nearly white; specific gravity, 0.7286; ash, 0.83; sometimes used for levers, handles of tools, etc.

BETULACEÆ.

294.—*Betula alba*, var. *populifolia*, Spach,

Ann. Sci. Nat. 2 ser. xv, 187; Hist. Veg. xi, 233.—Endlicher, Genera, Suppl. iv^a, 19.—Regel in Mem. Soc. Nat. Moscow, xix, 76, t. 4, f. 19-28; Gray, Manual N. States, 5 ed. 459.—Vasey, Cat. Forest Trees, 28.—Macoun in Geological Rep. Canada, 1879-'80, 55^c.

B. lenta, Du Roi, Harbk. i, 92 [not Linnaeus].—Wangenheim, Amer. 45, t. 29, f. 38.

B. populifolia, Marshall, Arbustum, 19.—Aiton, Hort. Kew. iii, 336; 2 ed. v, 299.—Willdenow, Berl. Baumz. 1 ed. 37, t. 2, f. 5; Spec. iv, 463.—Persoon, Syn. ii, 572.—Desfontaines, Hist. Arb. ii, 476.—Nouveau Duhamel, iii, 204.—Poirot, Suppl. i, 687.—Michaux f. Hist. Arb. Am. ii, 139, t. 2; N. American Sylva, 3 ed. ii, 78, t. 71.—Pursh, Fl. Am. Sept. ii, 620.—Smith in Rees' Cycl. iv, No. 8.—Barton, Prodr. Fl. Philadelph. 92; Compend. Fl. Philadelph. ii, 175.—Eaton, Manual, 109; 6 ed. 53.—Nuttall, Genera, ii, 218; Sylva, i, 25; 2 ed. i, 42.—Hayne, Dend. Fl. 166.—Sprengel, Syst. iii, 854.—Watson, Dend. Brit. ii, 151.—Torrey, Compend. Fl. N. States, 355; Fl. N. York, ii, 199, t. 112.—London, Arboretum, iii, 1707, f. 1560.—Hooker, Fl. Bor.-Am. ii, 155.—Eaton & Wright, Bot. 156.—Bigelow, Fl. Boston. 3 ed. 381.—Emerson, Trees Massachusetts, 213; 2 ed. i, 243 & t.—Gray, Manual N. States, 1 ed. 421.—Cooper in Smithsonian Rep. 1858, 256.—Wood, Cl. Book, 649; Bot. & Fl. 308.—Koch, Dendrologie, ii, 646.

B. acuminata, Ehrhart, Beitr. vi, 98.—Moench, Meth. 693.

B. alba, subspecies *populifolia*, Regel in Bull. Soc. Nat. Moscow, xxxviii^a, 399; De Candolle, Prodr. xvi^a, 164.

WHITE BIRCH. OLD-FIELD BIRCH. GRAY BIRCH.

New Brunswick and the valley of the lower Saint Lawrence river to the southern shores of lake Ontario, south, generally near the coast, to New Castle county, Delaware.

A small, short-lived tree of rapid growth, 6 to 9 meters in height, with a trunk 0.30 to 0.45 meter in diameter; dry, gravelly, barren soil or borders of swamps, now generally springing up upon abandoned or burned land in eastern New England.

Wood light, soft, not strong, close-grained, liable to check in drying, not durable; medullary rays numerous, obscure; color, light brown, the sap wood nearly white; specific gravity, 0.5760; ash, 0.29; largely used in the manufacture of spools, shoe-pegs, wood pulp, etc., for hoop-poles and fuel.

The bark and leaves, as well as those of *B. papyrifera* and *B. lenta*, are popularly esteemed as a remedy for various chronic diseases of the skin, bladder, etc., and for rheumatic and gouty complaints; the empyreumatic oil of birch obtained from the inner bark by distillation is used externally and internally for the same purposes (*U. S. Dispensatory*, 14 ed. 1592.—*Nat. Dispensatory*, 2 ed. 287); the bark occasionally used domestically in the manufacture of ink.

295.—*Betula papyrifera*, Marshall,

Arbustum, 19.—Michaux, Fl. Bor.-Am. ii, 180.

B. papyracea, Aiton, Hort. Kew. iii, 337; 2 ed. v, 300.—Willdenow, Spec. iv, 464; Enum. 981; Berl. Baumz. 58, t. 2, f. 1.—Nouveau Duhamel, iii, 205.—Persoon, Syn. ii, 572.—Desfontaines, Hist. Arb. ii, 477.—Poirot, Suppl. i, 688.—Michaux f. Hist. Arb. Am. ii, 133, t. 1; N. American Sylva, 3 ed. ii, 70, t. 69.—Smith in Rees' Cycl. iv, No. 9.—Pursh, Fl. Am. Sept. ii, 621.—B. S. Barton, Bot. Appx. 34, t. 27, f. 1.—Eaton, Manual, 109; 6 ed. 53.—Barton, Compend. Fl. Philadelph. ii, 175.—Nuttall, Genera, ii, 218; Sylva, i, 25; 2 ed. i, 42.—Hayne, Dend. Fl. 167.—Watson, Dend. Brit. ii, t. 152.—Sprengel, Syst. iii, 854.—Torrey, Compend. Fl. N. States, 355; Fl. N. York, ii, 199.—Audubon, Birds, t. 88.—London, Arboretum, iii, 1708, f. 1561 & t.—Hooker, Fl. Bor.-Am. ii, 155.—Eaton & Wright, Bot. 156.—Bigelow, Fl. Boston. 3 ed. 381.—Penn. Cycl. ii, 349.—Emerson, Trees Massachusetts, 210; 2 ed. i, 239 & t.—Parry in Owen's Rep. 618.—Richardson, Arctic Exped. 437.—Cooper in Smithsonian Rep. 1858, 256.—Hooker f. in Trans. Linnæan Soc. xxiii^a, 300, 339.—Wood, Cl. Book, 649; Bot. & Fl. 308.—Gray, Manual N. States, 5 ed. 459.—Koch, Dendrologie, ii, 645.—Vasey, Cat. Forest Trees, 28.—Macoun in Geological Rep. Canada, 1875-'76, 210.—Sears in Bull. Essex Inst. xiii, 180.—Bell in Geological Rep. Canada, 1879-'80, 45^c.

B. nigra, Loiseleur in Nouveau Duhamel, ii, t. 51 [not Linnæus].

B. grandis, Schrader in Ind. Hort. Goett. 1833, 2.

B. rubra, Loddiges, Cat. ed. 1836.

B. Canadensis, Loddiges, Cat. ed. 1836.

B. alba, var. *papyrifera*, Spach. in Ann. Sci. Nat. 2 ser. xv, 188; Hist. Veg. xi, 234.—Endlicher, Genera, Suppl. iv², 19.—Regel in Mem. Soc. Nat. Moscow, xix, 81, t. 5, f. 5-16.

B. cordifolia, Regel in Mem. Soc. Nat. Moscow, xix, 86, t. 12, f. 29-36.

B. alba, subspecies *papyrifera*, Regel in Bull. Soc. Nat. Moscow, xxxviii⁴, 401; De Candolle, Prodr. xvi², 166.

B. alba, subspecies *papyrifera*, var. *cordifolia*, Regel in Bull. Soc. Nat. Moscow, xxxviii⁴, 401; De Candolle, Prodr. xvi², 166.

B. alba, subspecies *papyrifera*, var. *communis*, Regel in Bull. Soc. Nat. Moscow, xxxviii⁴, 401; De Candolle, Prodr. xvi², 166.

B. alba, subspecies *commutata*, Regel in Bull. Soc. Nat. Moscow, xxxviii⁴, 401; De Candolle, Prodr. xvi², 166.

B. occidentalis, Lyall in Jour. Linnæan Soc. vii, 134 [not Hooker].

B. alba, var. *populifolia*, Winchell in Ludlow's Rep. Black Hills, 67 [not Spach].

CANOE BIRCH. WHITE BIRCH. PAPER BIRCH.

Northern Newfoundland and Labrador to the southern shores of Hudson bay and northwest to the Great Bear lake and the valley of the Yukon river, Alaska, south, in the Atlantic region to Wading river, Long island, the mountains of northern Pennsylvania, Clear lake, Montcalm county, Michigan, northeastern Illinois and Saint Cloud, Minnesota; in the Pacific region south to the Black hills of Dakota (*R. Douglas*), the Mullen trail of the Bitter Root mountains and Flathead lake, Montana, the neighborhood of Fort Colville, Washington territory (*Watson*), and the valley of the lower Fraser river, British Columbia (*Engelmann & Sargent*).

A tree 18 to 24 meters in height, with a trunk 0.60 to 0.90 meter in diameter; rich woodlands and banks of streams; very common in the northern Atlantic region and reaching a higher latitude than any deciduous tree of the American forest.

Wood light, strong, hard, tough, very close-grained, compact; medullary rays numerous, obscure; color, brown tinged with red, the sap-wood nearly white; specific gravity, 0.5955; ash, 0.25; largely used in the manufacture of spools, shoe-last and pegs, in turnery, for fuel, wood-pulp, etc.

The very tough, durable bark easily separated into thin layers, impervious to water, is largely used in the manufacture of canoes, tents, etc.

296.—*Betula occidentalis*, Hooker,

Fl. Bor.-Am. ii, 155.—Spach in Ann. Sci. Nat. 2 ser. xv, 197.—Nuttall, Sylva, i, 22, t. 7; 2 ed. i, 40, t. 7.—Endlicher, Genera, Suppl. iv², 20.—Torrey in Fremont's Rep. 97; Bot. Wilkes Exped. 466.—Nowberry in Pacific R. R. Rep. vi, 89.—Cooper in Smithsonian Rep. 1858, 261; Am. Nat. iii, 408.—Regel in Mem. Soc. Nat. Moscow, xix, 131, t. 15, f. 35.—Porter in Hayden's Rep. 1871, 493.—Watson in King's Rep. v, 323, t. 35; Pl. Wheeler, 17; Bot. California, ii, 79.—Porter & Hayden, Fl. Colorado; Hayden's Surv. Misc. Pub. No. 4, 127.—Rothrock in Pl. Wheeler. 50; Wheeler's Rep. vi, 239.—Vasey, Cat. Forest Trees, 28.—Macoun in Geological Rep. Canada, 1875-'76, 210.—G. M. Dawson in Canadian Nat. new ser. ix, 331.

B. alba, subspecies *occidentalis typica*, Regel in Bull. Soc. Nat. Moscow, xxxviii⁴, 400; De Candolle, Prodr. xvi², 165.

BLACK BIRCH.

British Columbia, south to the Mount Shasta region (Strawberry vale) and the eastern cañons of the Sierra Nevadas above Owen's valley (*Lemmon*), California, and through the interior ranges and the Rocky mountains to Utah and northern New Mexico.

A small tree, 8 to 12 meters in height, with a trunk sometimes 0.30 to 0.45 meter in diameter; mountain cañons and along streams, in moist soil, often throwing up several stems from the ground and forming dense thickets.

Wood soft, strong, brittle, close-grained, compact; medullary rays numerous, obscure; color, light brown, the sap-wood lighter; specific gravity, 0.6030; ash, 0.30; somewhat used for fencing, fuel, etc.

297.—*Betula lutea*, Michaux f.

Hist. Arb. Am. ii, 152, t. 5; N. American Sylva, 3 ed. ii, 82, t. 73.—Spach in Ann. Sci. Nat. 2 ser. xv, 191; Hist. Veg. xi, 243.—Endlicher, Genera, Suppl. iv², 20.—Wood, Bot. & Fl. 308.—Gray, Manual N. States, 5 ed. 459.—Koch, Dendrologie, ii, 640.—Vasey, Cat. Forest Trees, 28.—Sears in Bull. Essex Inst. xiii, 180.

B. excelsa, Pursh, Fl. Am. Sept. ii, 621 [not Aiton].—Nuttall, Genera, ii, 218.—Sprengel, Syst. iii, 854.—Torrey, Compend. Fl. N. States, 355; Fl. N. York, ii, 200.—Eaton, Manual, 6 ed. 53.—London, Arboretum, iii, 1711, f. 1564, 1565 & t.—Hooker, Fl. Bor.-Am. ii, 156.—Eaton & Wright, Bot. 156.—Bigelow, Fl. Boston. 3 ed. 382.—Lindley in Penn. Cycl. ii, 349.—Gray, Manual N. States, 1 ed. 422.—Emerson, Trees Massachusetts, 206; 2 ed. i, 235 & t.—Richardson, Arctic Exped. 438.—Cooper in Smithsonian Rep. 1858, 256.—Chapman, Fl. S. States, 428.—Curtis in Rep. Geological Surv. N. Carolina, 1830, iii, 74.—Wood, Cl. Book, 648.—Bell in Geological Rep. Canada, 1879-'80, 50^c.

B. lenta, Regel in Mem. Soc. Nat. Moscow, xix, 125, in part; Bull. Soc. Nat. Moscow, xxxviii⁴, 417, in part; De Candolle, Prodr. xvi², 179, in part.

YELLOW BIRCH. GRAY BIRCH.

Newfoundland, northern shores of the gulf of Saint Lawrence to Abittibi lake and the western shores of lake Superior and Rainy lake, south through the northern states to Delaware and southern Minnesota, and along the Alleghany mountains to the high peaks of North Carolina and Tennessee.

One of the largest and most valuable deciduous trees of the northern New England and Canadian forests, often 21 to 29 meters in height, with a trunk 0.90 to 1.20 meter in diameter; rich woodlands; common.

Wood heavy, very strong and hard, very close-grained, compact, satiny, susceptible of a beautiful polish; medullary rays numerous, obscure; color, light brown tinged with red, the heavier sap-wood nearly white; specific gravity, 0.6553; ash, 0.31; largely used for fuel, in the manufacture of furniture, button and tassel molds, pill and match boxes, and for the hubs of wheels.

298.—*Betula nigra*, Linnæus,

Spec. 1 ed. 982.—Marshall, Arbustum, 18.—Walter, Fl. Caroliniana, 231.—Aiton, Hort. Kew. iii, 336; 2 ed. v, 299.—Gærtner, Fruct. ii, 54, t. 90, f. 1.—Willdenow, Spec. iv, 464; Ennm. 931; Berl. Baumz. 56.—Nouveau Duhamel, iii, 203, t. 51.—Persoon, Syn. ii, 572.—Desfontaines, Hist. Arb. ii, 477.—Smith in Rees' Cycl. iv, No. 2.—Pursh, Fl. Am. Sept. ii, 621.—Nuttall, Genera, ii, 218.—Hayne, Dend. Fl. 166.—Lamarck, Ill. iii, 350, t. 760, f. 2.—Elliott, Sk. ii, 616.—Watson, Dend. Brit. ii, t. 153.—Sprengel, Syst. ii, 854.—Torrey, Compend. Fl. N. States, 355; Fl. N. York, ii, 201.—Beck, Bot. 325.—London, Arboretum, iii, 1710, f. 1562, 1563 & t.—Penn. Cycl. ii, 149.—Emerson, Trees Massachusetts 208; 2 ed. i, 237.—Darlington, Fl. Cestrica, 3 ed. 275.—Darby, Bot. S. States, 508.—Cooper in Smithsonian Rep. 1858, 256.—Chapman, Fl. S. States, 428.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 73.—Regel in Mem. Soc. Nat. Moscow, xix, 118, t. 12, f. 1-12; Bull. Soc. Nat. Moscow, xxxviii⁴, 412; De Candolle, Prodr. xvi², 175.—Lesquereux in Owen's 2d Rep. Arkansas, 389.—Wood, Cl. Book, 649; Bot. & Fl. 308.—Porcher, Resources S. Forests, 266.—Gray, Manual N. States, 5 ed. 459; Hall's Pl. Texas, 21.—Koch, Dendrologie, ii, 644.—Young, Bot. Texas, 512.—Vasey, Cat. Forest Trees, 28.—Burbank in Proc. Boston Soc. Nat. Hist. xviii, 214.—Broadhead in Coulter's Bot. Gazette, iii, 60.—Ridgway in Proc. U. S. Nat. Mus. 1882, 85.

B. lanulosa, Michaux, Fl. Bor.-Am. ii, 181.—Nouveau Duhamel, iii, 206.

B. rubra, Michaux f. Hist. Arb. Am. ii, 142, t. 3; N. American Sylva, 3 ed. ii, 80, t. 72.—Loddiges, Bot. Cab. t. 1248.—Eaton, Manual, 6 ed. 53.—Eaton & Wright, Bot. 156.—Spach in Ann. Sci. Nat. 2 ser. xv, 185; Hist. Veg. xi, 230.—Endlicher, Genera, Suppl. iv², 19.


B. angulata, Loddiges, Cat. ed. 1836.

RED BIRCH. RIVER BIRCH.

Banks of the Merrimac and Spicket rivers, Middlesex and Essex counties, Massachusetts, Wading river, Long island, south through the coast and middle districts to the Chattahoochee region of western Florida, west to western Iowa, northwestern Missouri, eastern Kansas, the Indian territory, and the valley of the Trinity river, Texas.

A tree 18 to 24 meters in height, with a trunk rarely exceeding 0.75 meter in diameter; banks of streams and ponds; very common and reaching its greatest development in the south Atlantic states and in the basin of the lower Mississippi river.

Wood light, rather hard, strong, close-grained, compact; medullary rays numerous, obscure; color, brown, the sap-wood much lighter; specific gravity, 0.5762; ash, 0.35; used in the manufacture of furniture, woodenware, wooden shoes, ox-yokes, etc.



299.—*Betula lenta*, Linnaeus,

Spec. 1 ed. 983.—Lamarck, Diet. i, 453.—Marshall, Arbustum, 19.—Aiton, Hort. Kew. iii, 337; 2 ed. v, 300.—Willdenow, Spec. iv, 464; Enum. 981; Berl. Baumz. 59.—Persoon, Syn. ii, 572.—Desfontaines, Hist. Arb. ii, 477.—Nouveau Duhamel, iii, 205.—Michaux f. Hist. Arb. Am. ii, 147, t. 4; N. American Sylva, 3 ed. ii, 85, t. 74.—Smith in Rees' Cycl. iv, No. 3.—Pursh, Fl. Am. Sept. ii, 621.—Eaton, Manual, 109; 6 ed. 53.—Barton, Compend. Fl. Philadelph. ii, 175.—Nuttall, Genera, ii, 218.—Hayne, Dend. Fl. 167.—Elliott, Sk. ii, 617.—Watson, Dend. Brit. ii, 144.—Sprengel, Syst. ii, 854.—Torrey, Compend. Fl. N. States, 356; Fl. N. York, ii, 200.—Guimpel, Otto & Hayne, Abb. Holz. 105, t. 83.—London, Arboretum, iii, 1713, f. 1566.—Hooker, Fl. Bor.-Am. ii, 156.—Eaton & Wright, Bot. 156.—Bigelow, Fl. Boston. 3 ed. 381.—Lindley in Penn. Cycl. ii, 349.—Spach in Ann. Sci. Nat. 2 ser. xv, 190; Hist. Veg. xi, 241.—Emerson, Trees Massachusetts, 203; 2 ed. i, 232 & t.—Richardson, Arctic Exped. 438.—Endlicher, Genera, Suppl. iv², 20.—Darlington, Fl. Cestrica, 3 ed. 275.—Darby, Bot. S. States, 508.—Cooper in Smithsonian Rep. 1858, 256.—Chapman, Fl. S. States, 428.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 74.—Regel in Mem. Soc. Nat. Moscow, xxxviii⁴, 125, in part; Bull. Soc. Nat. Moscow, xxxviii, 417, in part; De Candolle, Prodr. xvi², 179, in part.—Wood, Cl. Book, 648; Bot. & Fl. 308.—Porcher, Resources S. Forests, 265.—Gray, Manual N. States, 5 ed. 458.—Koch, Dendrologie, ii, 639.—Vasey, Cat. Forest Trees, 28.—Sears in Bull. Essex Inst. xiii, 180.—Bell in Geological Rep. Canada, 1879-'80, 55.—Ridgway in Proc. U. S. Nat. Mus. 1882, 85.

B. nigra, Du Roi, Harbk. i, 93.—Wangenheim, Amer. 35, t. 15, f. 34.

B. excelsa, Aiton, Hort. Kew. iii, 337; 2 ed. v, 299 [not Pursh].—Willdenow, Spec. iv, 464.—Berl. Baumz. 41, t. 2, f. 2.—Nouveau Duhamel, iii, 203, t. 52.—Persoon, Syn. ii, 572.—Desfontaines, Hist. Arb. ii, 477.—Poiret, Suppl. i, 687.—Smith in Rees' Cycl. iv, No. 10.—Hayne, Dend. Fl. i, 7.—Spach in Ann. Sci. Nat. 2 ser. xv, 188; Hist. Veg. xi, 243.—Endlicher, Genera, iv², 20.

B. carpinifolia, Ehrhart, Beitr. vi, 99.—Willdenow, Enum. 981; Berl. Baumz. 49.

CHERRY BIRCH. BLACK BIRCH. SWEET BIRCH. MAHOGANY BIRCH.

Newfoundland and the valley of the Saguenay river, west through Ontario to the Manitou islands of lake Huron, south to northern Delaware and southern Indiana, and along the Alleghany mountains to the Chattahoochee region of northern Florida, extending west to middle Kentucky and Tennessee.

A tree 18 to 24 meters in height, with a trunk 0.90 to 1.50 meter in diameter; rich woodlands; very common in all northern forests.

Wood heavy, very strong and hard, close-grained, compact, satiny, susceptible of a beautiful polish; medullary rays numerous, obscure; color, dark brown tinged with red, the sap-wood light brown or yellow; specific gravity, 0.7617; ash, 0.26; now largely used in the manufacture of furniture and for fuel; in Nova Scotia and New Brunswick largely in ship-building.

"Birch beer" is obtained by fermenting the saccharine sap of this and perhaps some other species of the genus.

300.—*Alnus maritima*, Muhlenberg,

Mss.—Nuttall, Sylva, i, 34, t. 10²; 2 ed. i, 50, t. 10².—Gray, Manual N. States, 5 ed. 461; Hall's Pl. Texas, 21.—Canby in Coulter's Bot. Gazette, vi, 1881.

Betula-Alnus maritima, Marshall, Arbustum, 20.

A. oblongata, Regel in Mem. Soc. Nat. Moscow, xix, 172, t. vi, f. 3-9 [not Willdenow].

A. maritima typica, Regel in Bull. Soc. Nat. Moscow, xxxviii⁴, 427; De Candolle, Prodr. xvi², 186.

SEASIDE ALDER.

Southern Delaware and eastern Maryland, near the coast; valley of the Red river, Indian territory, in about longitude 96° 30' W. (*E. Hall*); Manchuria and Japan (*A. maritima*, *Japonica* and *arguta*, *Regel in De Candolle, Prodr. xvi², 186*).

A small tree, 6 to 7 meters in height, with a trunk 0.10 to 0.15 meter in diameter; borders of streams and swamps.

Wood light, soft, close-grained, checking badly in drying; medullary rays broad, conspicuous; color, light bright brown, the sap-wood hardly distinguishable, somewhat lighter; specific gravity, 0.4996; ash, 0.39.

301.—*Alnus rubra*, Bongard,

Mem. Acad. St. Petersburg, 6 ser. ii, 162.—Hooker, Fl. Bor.-Am. ii, 158.—Spach in Ann. Sci. Nat. 2 ser. xv, 205.—Endlicher, Genera, Suppl. iv², 21.—Lyll in Jour. Linnæan Soc. vii, 134.—Regel in Bull. Soc. Nat. Moscow, xxxviii⁴, 429; De Candolle, Prodr. xvi², 188.—Torrey, Bot. Wilkes Exped. 467.—Watson, Bot. California, ii, 80.—G. M. Dawson in Canadian Nat. new ser. ix, 331.

?*A. glutinosa*, Pursh, Fl. Am. Sept. ii, 622 [not Willdenow].

A. Oregana, Nuttall, Sylva, i, 28, t. 9; 2 ed. i, 44, t. 9.—Newberry in Pacific R. R. Rep. vi, 25, 89.—Cooper in Smithsonian Rep. 1858, 261; Pacific R. R. Rep. xii², 28, 68.—Vasey, Cat. Forest Trees, 28.—Hall in Coulter's Bot. Gazette, ii, 91.

A. incana, var. *rubra*, Regel in Mem. Soc. Nat. Moscow, xix, 157, t. 17, f. 3-4.

ALDER.

Sitka, south through the islands and Coast ranges of British Columbia, Washington territory, Oregon, and California to Santa Barbara, extending east through the Blue mountains to northern Montana.

A large tree, 24 to 30 meters in height, with a trunk 0.90 to 1.20 meter in diameter, or in British Columbia and the Blue mountains often reduced to a low shrub; river bottom lands and borders of streams; most common and reaching its greatest development along the large streams of western Washington territory and Oregon.

Wood light, soft, not strong, brittle, very close-grained, compact, easily worked, satiny, susceptible of a beautiful polish; medullary rays distant, broad; color, light brown tinged with red, the sap-wood nearly white; specific gravity, 0.4813; ash, 0.42; largely used in Oregon in the manufacture of furniture.

302.—*Alnus rhombifolia*, Nuttall,

Sylva, i, 33; 2 ed. i, 49.—Torrey, Bot. Wilkes Exped. 467.—Vasey, Cat. Forest Trees, 28.—Watson, Bot. California, ii, 80.

A. glutinosa, var. *serrulata*, Regel in Mem. Soc. Nat. Moscow, xix, 164, in part.

A. serrulata, var. *rugosa*, Regel in Bull. Soc. Nat. Moscow, xxxviii⁴, 432, in part; De Candolle, Prodr. xvi², 188, in part.

ALDER.

Valley of the lower Fraser river, British Columbia, south through the Coast ranges to southern California, extending east along the ranges of Washington territory to Clear creek, Idaho (*Watson*), and the valley of the Flathead river, Montana (*Canby & Sargent*).

A small tree, 9 to 15 meters in height, with a trunk sometimes 0.60 to 0.90 meter in diameter, or toward its northern and eastern limits reduced to a shrub; borders of streams; the common alder of the California valleys.

Wood light, soft, not strong, brittle, close-grained, compact; medullary rays numerous, obscure; color, light brown, the sap-wood lighter, often nearly white; specific gravity, 0.4127; ash, 0.31.

303.—*Alnus oblongifolia*, Torrey,

Bot. Mex. Boundary Survey, 204.—Cooper in Smithsonian Rep. 1858, 266.—Watson in Pl. Wheeler, 17; Bot. California, ii, 80.—Rothrock in Wheeler's Rep. vi, 239.—Rusby in Bull. Torrey Bot. Club, ix, 79.

A. serrulata, var. *oblongifolia*, Regel in Bull. Soc. Nat. Moscow, xxxviii⁴, 443; De Candolle, Prodr. xvi², 188.

ALDER.

San Bernardino and Cayumaca mountains, California, through the ranges of southern Arizona and southern New Mexico to the valley of the Rio Grande; southward into Mexico.

A tree 15 to 21 meters in height, with a trunk 0.90 to 1.20 meter in diameter; borders of streams in deep mountain cañons.

Wood light, soft, not strong, brittle, close-grained, compact; medullary rays numerous, very obscure; color, light brown tinged with yellow, the sap-wood nearly white; specific gravity, 0.3981; ash, 0.42.

304.—*Alnus serrulata*, Willdenow,

Spec. iv, 336; Enum. 965; Berl. Baumz. 2 ed. 21.—Nouveau Duhamel, ii, 216.—Persoon, Syn. ii, 550.—Desfontaines, Hist. Arb. ii, 488.—Aiton, Hort. Kew. 2 ed. v, 259.—Michaux f. Hist. Arb. Am. iii, 320, t. 4, f. 1; N. American Sylva, 3 ed. ii, 87, t. 75, f. 1.—Pursh, Fl. Am. Sept. ii, 623.—Barton, Prodr. Fl. Philadelph. 89; Compend. Fl. Philadelph. ii, 158.—Eaton, Manual, 105; 6 ed. 12.—Nuttall, Genera, ii, 206.—Hayne, Dend. Fl. 122.—Elliott, Sk. ii, 567.—Torrey, Compend. Fl. N. States, 350; Fl. N. York, ii, 202, t. 115.—Beck, Bot. 326.—Darlington, Fl. Cestrica, 3 ed. 276.—London, Arboretum, iii, 1688, f. 1544.—Eaton & Wright, Bot. 120.—Bigelow, Fl. Boston. 3 ed. 220.—Spach in Ann. Sci. Nat. 2 ser. xv, 206; Hist. Veg. xi, 251.—Emerson, Trees Massachusetts, 218; 2 ed. i, 248 & t.—Endlicher, Genera, Suppl. iv², 21.—Darby, Bot. S. States, 508.—Chapman, Fl. S. States, 429.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 102.—Lesquereux in Owen's 2d Rep. Arkansas, 389.—Wood, Cl. Book, 650; Bot. & Fl. 308.—Porcher, Resources S. Forests, 266.—Gray, Manual N. States, 5 ed. 461.—Young, Bot. Texas, 513.—Broadhead in Coulter's Bot. Gazette, iii, 60.

Betula rugosa, Du Roi, Harbk. i, 176.—Wangenheim, Amer. 86, t. 29, f. 60.—Ehrhart, Beitr. iii, 21.

?*Betula-Alnus glauca*, Marshall, Arbustum, 20.

Betula serrulata, Aiton, Hort. Kew. iii, 338.—Willdenow, Berl. Baumz. 1 ed. 45.—Abbot, Insects Georgia, ii, 183, t. 92.—Michaux, Fl. Bor.-Am. ii, 181.

A. serrulata, var. *vulgaris*, Spach in Ann. Sci. Nat. 2 ser. xv, 206.

A. serrulata, var. *macrophylla*, Spach in Ann. Sci. Nat. 2 ser. xv, 206.

A. serrulata, var. *oblongata*, Spach, Hist. Veg. xi, 251.

A. serrulata, var. *latifolia*, Spach, Hist. Veg. xi, 251.

A. rubra, Tuckerman in Am. Jour. Sci. 1 ser. xlv, 32.

A. hybrida, Reichenbach, Icon. Fl. Germ. xii, t. 630, f. 1292.

A. glutinosa, var. *serrulata*, Regel in Mem. Soc. Nat. Moscow, xix, 164, t. 11, f. 6, 8, in part.

A. glutinosa, var. *rugosa*, Regel in Mem. Soc. Nat. Moscow, xix, 165, t. 11, f. 9, 10.

A. serrulata genuina and *obtusifolia*, Regel in Bull. Soc. Nat. Moscow, xxxviii⁴, 432; De Candolle, Prodr. xvi², 188.

A. serrulata, var. *rugosa*, Regel in Bull. Soc. Nat. Moscow, xxxviii⁴, 432, in part; De Candolle, Prodr. xvi², 188, in part.

A. rugosa, Koch, Dendrologie, ii, 635.

A. oblongata, *undulata*, *rugosa*, *Canadensis*, and *Americana*, Hort.

BLACK ALDER. SMOOTH ALDER.

Essex county, Massachusetts, west to southern Missouri, south to northern Florida and the valley of the Trinity river, Texas.

A small tree, 6 to 12 meters in height, with a trunk 0.10 to 0.15 meter in diameter, or more often a tall, branching shrub forming dense thickets; borders of streams and swamps, probably reaching its greatest development in southern Arkansas.

Wood light, soft, close-grained, compact; medullary rays numerous, conspicuous; color, light brown, the sap-wood lighter; specific gravity, 0.4666; ash, 0.38.

A decoction of the bark and leaves, as well as those of *A. incana*, is a popular remedy against impurity of the blood and in the treatment of diarrhoea and hæmaturia, etc. (*Nat. Dispensatory*, 2 ed. 135).

305.—*Alnus incana*, Willdenow,

Spec. iv, 335; Enum. 965; Berl. Baumz. 2 ed. 20.—Persoon, Syn. ii, 550.—Aiton, Hort. Kew. 2 ed. v, 259.—Hayne, Dend. Fl. 152.—Eaton, Manual, 6 ed. 12.—London, Arboretum, iii, 1687, f. 1543.—Hooker, Fl. Bor.-Am. ii, 157.—Eaton & Wright, Bot. 120.—Spach in Ann. Sci. Nat. 2 ser. xv, 206; Hist. Veg. xi, 252.—Nuttall, Sylva, i, 30; 2 ed. i, 46.—Tuckerman in Am. Jour. Sci. 1 ser. xlv, 32.—Torrey, Fl. N. York, ii, 202.—Emerson, Trees Massachusetts, 220; 2 ed. i, 251 & t.—Endlicher, Genera, Suppl. iv², 21.—Parry in Owen's Rep. 618.—Cooper in Smithsonian Rep. 1858, 256.—Hooker f. in Trans. Linnæan Soc. xxiii², 301.—Wood, Cl. Book, 649; Bot. & Fl. 308.—Regel in Bull. Soc. Nat. Moscow, xxxviii⁴, 433; De Candolle, Prodr. xvi², 188.—Gray, Manual N. States, 5 ed. 461.—Koch, Dendrologie, ii, 636.—Vasey, Cat. Forest Trees, 28.—Macoun in Geological Rep. Canada, 1875-'76, 210.—Bell in Geological Rep. Canada, 1879-'80, 556.

Betula-Alnus, var. *β. incana*, Linnæus, Spec. 1 ed. 983.—Du Roi, Harbk. i, 109.

Betula incana, Linnæus, Suppl. 417.—Aiton, Hort. Kew. iii, 339.—Willdenow, Berl. Baumz. 1 ed. 45.—Smith in Rees' Cycl. iv, No. 7.

?*Betula-Alnus rubra*, Marshall, Arbustum, 20.

A. glauca, Michaux f. Hist. Arb. Am. iii, 322, t. 4, f. 2; N. American Sylva, 3 ed. 89, t. 75, f. 2.—Bigelow, Fl. Boston. 3 ed. 367.

A. incana, var. *glauca*, Gray, Manual N. States, 1 ed. 423; 3 ed. 461.

A. incana, *Americana*, and *genuina*, Regel in Mem. Soc. Nat. Moscow, xix, 155.

SPECKLED ALDER. HOARY ALDER. BLACK ALDER.

Newfoundland to the eastern base of the Rocky mountains, south to northern New England, Wisconsin, Minnesota, and eastern Nebraska; in Europe.

A small tree, 6 to 7 meters in height, with a trunk 0.10 to 0.15 meter in diameter, or more often a tall, branching shrub; borders of streams and swamps.

A form with leaves green and glabrous on both sides or slightly pubescent, extending through the mountain ranges of the Pacific region from the Saskatchewan and British Columbia to New Mexico and the southern Sierra Nevadas of California, is—

var. *virescens*, Watson, Bot. California, ii, 81.

A. incana, var. *glauca*, Regel in Mem. Soc. Nat. Moscow, xix, 154, in part; Bull. Soc. Nat. Moscow, xxxviii⁴, 433, in part; De Candolle, Prodr. xvi³, 189, in part.—Watson in King's Rep. v, 326 [not Aiton]; Pl. Wheeler, 17.—Rothrock, Pl. Wheeler, 50; Wheeler's Rep. vi, 239.—Macoun in Geological Rep. Canada, 1875-'76, 210.

A. serrulata, var. *rugosa*, Regel in Bull. Soc. Nat. Moscow, xxxviii⁴, 432, in part; De Candolle, Prodr. xvi³, 188, in part.

Wood light, soft, close-grained, checking in drying; medullary rays numerous, broad; color, light brown, the sap-wood nearly white; specific gravity, 0.4607; ash, 0.42; preferred and largely used in northern New England in the final baking of bricks, and occasionally, as well as that of *A. serrulata*, in the manufacture of gunpowder.

SALICACEÆ.

306.—*Salix nigra*, Marshall,

Arbustum, 139.—Muhlenberg in Neue Schriften Gesell. Nat. Fr. Berlin, iv, 237, t. 4, f. 5 (Ann. Bot. ii, 65, t. 5, f. 5).—Willdenow, Spec. iv, 657; Enum. 1003; Berl. Baumz. 2 ed. 426.—Perseon, Syn. ii, 599.—Michaux f. Hist. Arb. Am. iii, 324, t. 5, f. 1; N. American Sylva, 3 ed. iii, 64, t. 125, f. 1.—Pursh, Fl. Am. Sept. ii, 614.—Poiret, Suppl. iv, 61.—Eaton, Manual, 118; 6 ed. 320.—Nuttall, Genera, ii, 231; Sylva, i, 79; 2 ed. i, 94.—Hayne, Dend. Fl. 180.—Elliott, Sk. ii, 670.—Sprengel, Syst. i, 100.—Torrey, Compend. Fl. N. States, 370; Fl. N. York, ii, 209.—Forbes, Sal. Woburn. 280.—W. Koch, Comment. 17.—Beck, Bot. 320.—Trautvetter in Mem. Acad. St. Petersburg, iii, 614.—London, Arboretum, iii, 1529, 1604, f. 8.—Hooker, Fl. Bor.-Am. ii, 148.—Barratt, Sal. Am. No. 19.—Eaton & Wright, Bot. 408.—Dietrich, Syn. v, 419.—Seringe, Fl. Jard. ii, 35.—Emerson Trees Massachusetts, 271; 2 ed. i, 307 & t.—Darlington, Fl. Cestrica, 3 ed. 279.—Andersson in Ofr. af. Vet. Akad. Forh. 1858, 114 (Proc. Am. Acad. iv, 53); Kongl. Sven. Akad. Handl. vi, 19, f. 15; De Candolle, Prodr. xvi³, 200.—Darby, Bot. S. States, 506.—Cooper in Smithsonian Rep. 1858, 256.—Walpers, Ann. v, 744.—Chapman, Fl. S. States, 430.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 75.—Lesquereux in Owen's 2d Rep. Arkansas, 389.—Wood, Cl. Book, 654; Bot. & Fl. 310.—Porcher, Resources S. Forests, 334.—Engelmann in Trans. Am. Phil. Soc. new ser. xii, 209.—Gray, Manual N. States, 5 ed. 460; Hall's Pl. Texas, 21.—Koch, Dendrologie, ii, 513.—Young, Bot. Texas, 514.—Macoun in Geological Rep. Canada, 1875-'76, 210.—Vasey, Cat. Forest Trees, 23.—Bebb in Bot. California, ii, 83.—Sears in Bull. Essex Inst. xiii, 181.—Ridgway in Proc. U. S. Nat. Mus. 1882, 86.—Hemsley, Bot. Am.-Cent. iii, 180

S. pentandra, Walter, Fl. Caroliniana, 243.

S. Caroliniana, Michaux, Fl. Bor.-Am. ii, 226.—Lamarek, Dict. vi, 662.—Poiret, Suppl. v, 62.

S. Houstoniana, Pursh, Fl. Am. Sept. ii, 614.—Poiret, Suppl. v, 68.—Sprengel, Syst. i, 107.—Elliott, Sk. ii, 670.—Trautvetter in Mem. Acad. St. Petersburg, iii, 615.—Forbes, Sal. Woburn. 21, t. 21.—Eaton & Wright, Bot. 409.

S. falcata, Pursh, Fl. Am. Sept. ii, 614 [not HBK.].—Poiret, Suppl. v, 70.—Sprengel, Syst. i, 107.—Forbes, Sal. Woburn. 279.—Eaton, Manual, 6 ed. 320.—Hooker, Fl. Bor.-Am. ii, 149.—Barratt, Sal. Am. No. 21.—Dietrich, Syn. v, 420.

? *S. ambigua*, Pursh, Fl. Am. Sept. ii, 617.—Forbes, Sal. Woburn. 282.—Eaton, Manual, 6 ed. 321.—Eaton & Wright, Bot. 409.

S. ligustrina, Michaux f. Hist. Arb. Am. iii, 326, t. 5, f. 2; N. American Sylva, 3 ed. iii, 65, t. 125, f. 2.—Poiret, Suppl. v, 61.

S. Purshiana, Sprengel, Syst. iii, 608.—Beck, Bot. 320.—Darlington, Fl. Cestrica, 2 ed. 560.

S. flavo-virens, Hornemann in Cat. Hort. Hafn. Suppl. ii, 11.

? *S. cordata*, var. *falcata*, Torrey, Compend. Fl. N. States, 370.

S. nigra, var. *falcata*, Torrey, Fl. N. York, ii, 209.—Carey in Gray, Manual N. States, 1 ed. 429.—Darlington, Fl. Cestrica, 3 ed. 280.

BLACK WILLOW.

Southern New Brunswick and the northern shores of lakes Huron and Superior southward through the Atlantic region to bay Biscayne and the Caloosa river, Florida, and the valley of the Guadalupe river, Texas; Pacific region, valleys of the Sacramento river, California, and the Colorado river, Arizona.

A small tree, sometimes 15 to 18 meters in height, with a trunk rarely 0.60 meter in diameter, or in southern Florida reduced to a low shrub; usually along the banks of streams; most common in the basin of the Mississippi river and reaching its greatest development in the rich bottom lands of the Colorado and other rivers of eastern Texas; varying greatly in the size and shape of the leaves (vars. *angustifolia*, *longifolia*, *latifolia*, etc., *Andersson in Kongl. Sven. Akad. Handl.* vi, 20), length and habit of the aments, etc.

The best marked forms are—

var. *marginata*, *Andersson in Kongl. Sven. Akad. Handl.* vi, 22; *De Candolle, Prodr.* xvi², 201.

S. marginata, *Wimmer in Schedul. Herb. Vindab.*

var. *longipes*, *Andersson in Kongl. Sven. Akad. Handl.* vi, 22; *De Candolle, Prodr.* xvi², 201.

S. longipes, *Shuttleworth in herb. Hooker.—Andersson in Ofv. af. Vet. Akad. Forh.* 1855, 114 (*Proc. Am. Acad.* iv, 53).—*Walpers, Ann.* v, 744.

Forms of var. *longipes* more or less pubescent have been characterized by *Andersson in Kongl. Sven. Akad. Handl.* vi, 22; *De Candolle, Prodr.* xvi², 201, as subvars. *venulosa* and *gongylocarpa* [*Shuttleworth*], (*S. longipes*, var. *pubescens*, *Andersson in Proc. Am. Acad.* iv, 53; *S. subvillosa*, *Elliott in herb. Schweinitz ex. Nuttall, Sylva*, i, 79; 2 ed. i, 94, *vide Gray in Proc. Am. Acad.* iv, 53, note).

var. *Wrightii*, *Andersson in Kongl. Sven. Akad. Handl.* vi, 22; *De Candolle, Prodr.* xvi², 201.—*Hemsley, Bot. Am.-Cent.* iii, 180.

S. Wrightii, *Andersson in Ofv. af. Vet. Akad. Forh.* 1858, 115 (*Proc. Am. Acad.* iv, 55 —*Walpers, Ann.* v, 745.—*Torrey in Bot. Mex. Boundary Survey*, 204.

var. *Wardii*, *Bebb in Bull. U. S. Nat. Mus.* No. 22, 114.

Wood light, soft, weak, close-grained, checking badly in drying; medullary rays obscure; color, brown, the sap-wood nearly white; specific gravity, 0.4456; ash, 0.70.

The tonic and astringent bark used domestically as a popular febrifuge, and containing, in common with that of all the species of the genus, salicylic acid, a powerful anti-pyritic now successfully used in the treatment of acute cases of gout, rheumatism, typhoid fever, etc. (*Am. Jour. Pharm.* 1875, 303.—*U. S. Dispensatory*, 14 ed. 796, 1748.—*Nat. Dispensatory*, 2 ed. 1248).

NOTE.—The closely allied *Salix occidentalis*, *Bosc*, of the West Indies is not perhaps specifically distinct from *S. nigra*, with which some of the forms of var. *longipes* from southern Florida seem to connect it.

307.—*Salix amygdaloides*, *Andersson*,

Ofv. af. Vet. Akad. Forh. 1858, 114 (*Proc. Am. Acad.* iv, 53).—*Walpers, Ann.* v, 744.—*Bebb in Wheeler's Rep.* vi, 240.

? *S. melanopsis*, *Nuttall, Sylva*, i, 78, t. 21; 2 ed. i, 93, t. 21.

S. nigra, var. *amygdaloides*, *Andersson in Kongl. Sven. Akad. Handl.* vi, 21; *De Candolle, Prodr.* xvi², 201.—*Rothrock, Pl. Wheeler*, 50.—*Porter & Coulter, Fl. Colorado*; *Hayden's Surv. Misc. Pub.* No. 4, 128.

WILLOW.

Shores of the great lakes (Wayne county, New York, *Hankenson*; Painesville, Ohio, *Beardslee*), westward to the valley of the Saskatchewan, and southward through the Rocky Mountain region to southern New Mexico; banks of the lower Columbia river, Oregon (*Howells*).

A small tree, rarely 9 to 12 meters in height, with a trunk 0.15 to 0.30 meter in diameter; along streams.

Wood light, soft, not strong, close-grained, checking in drying; the heart-wood light brown, sap-wood nearly white; specific gravity, 0.4509; ash, 0.92.

308.—*Salix lævigata*, Bebb,

Am. Nat. viii, 202; Bot. California, ii, 83.

WILLOW.

California, Sierra county (*Lemmon*) and the valley of the Sacramento river to the southern boundary of the state.

A tree sometimes 15 meters in height, with a trunk 0.30 to 0.60 meter in diameter; borders of streams and bottom lands.

A form with narrower falcate leaves (*Yreka*, *E. L. Greene*) is—

var. *angustifolia*, Bebb in Bot. California, ii, 84.—Rothrock in Wheeler's Rep. vi, 374.

A form with short, densely-flowered aments is—

var. *congesta*, Bebb in Bot. California, ii, 84.

Wood light, soft, not strong, brittle, close-grained, compact; medullary rays numerous, very thin; color, light brown tinged with red; specific gravity, 0.4872; ash, 0.58.

309.—*Salix lasiandra*, Bentham,

Pl. Hartweg. 336.—Torrey in Pacific R. R. Rep. iv, 138.—Newberry in Pacific R. R. Rep. vi, 89.—Bebb in Bot. California, ii, 84.

S. Hoffmanniana, Hooker & Arnott, Bot. Beechey, 159.

S. speciosa, Nuttall, Sylva, i, 58, t. 17; 2 ed. i, 74, t. 17 [not Hooker & Arnott].—Newberry in Pacific R. R. Rep. vi, 89.—Cooper in Pacific R. R. Rep. xii², 29.

S. lucida, var. *angustifolia*, forma *lasiandra*, Andersson in Ofv. af. Vet. Akad. Forh. 1858, 115 (Proc. Am. Acad. iv, 54).

S. arguta, var. *lasiandra*, Andersson in Kongl. Sven. Akad. Handl. vi, 33; De Candolle, Prodr. xvi², 206.

WILLOW.

British Columbia, shores of lake Kamloop (*Macoun*), southward to the valley of the Sacramento river, California; Rocky mountains, Utah, and through Colorado to New Mexico (var. *Fendleriana*).

A tree 12 to 18 meters in height, with a trunk sometimes 0.60 meter in diameter; banks of streams; very common; varying in the shape of the leaves and character of the aments.

The best marked forms are—

var. *lancifolia*, Bebb in Bot. California, ii, 84.

S. lancifolia, Andersson in Kongl. Sven. Akad. Handl. vi, 34, f. 23.—Gray in Proc. Am. Acad. vii, 402.—Hall in Coulter's Bot. Gazette, ii, 91.

S. lucida, var. *macrophylla*, Andersson in De Candolle, Prodr. xvi², 205.

The common form of British Columbia and western Washington territory and Oregon.

var. *Fendleriana*, Bebb in Bot. California, ii, 84.

S. pentandra, var. *caudata*, Nuttall, Sylva, i, 61, t. 18; 2 ed. i, 77, t. 18.

S. Fendleriana, Andersson in Ofv. af. Vet. Akad. Forh. 1858, 115 (Proc. Am. Acad. iv, 54).—Walpers, Ann. v, 745.

S. arguta, Andersson in Kongl. Sven. Akad. Handl. vi, 32; De Candolle, Prodr. xvi², 205, in part.

Wood light, soft, not strong, brittle, close-grained, compact; medullary rays numerous, very obscure; color, light brown, the sap-wood lighter or often nearly white; specific gravity, 0.4756; ash, 0.60. Var. *lancifolia*, specific gravity, 0.4547; ash, 0.79. Var. *Fendleriana*, the heart-wood brown, sap-wood light brown; specific gravity, 0.4598; ash, 0.56.

310.—*Salix longifolia*, Muhlenberg,

Neue Schriften Gesell. Nat. Fr. Berlin, iv, 238, t. 6, f. 6 (Ann. Bot. ii, 66, t. 5, f. 6).—Willdenow, Spec. iv, 670.—Persson, Syn. ii, 600.—Pursh, Fl. Am. Sept. ii, 613.—Nuttall, Genera, ii, 231.—Torrey in Ann. Lyc. N. York, ii, 248; Fl. N. York, ii, 209; Nicotlet's Rep. 160; Fremont's Rep. 97; Emory's Rep. 412; Sitgreaves' Rep. 172; Bot. Mex. Boundary Survey, 204.—Barratt, Sal. Am. No. 23.—Beck, Bot. 320.—Eaton, Manual, 6 ed. 319.—Eaton & Wright, Bot. 408.—Hooker, Fl. Bor.-Am. ii, 149.—Dietrich, Syn. v, 420.—Parry in Owen's Rep. 618.—Richardson, Arctic Exped. 439, 440.—Cooper in Smithsonian Rep. 1858, 261.—Andersson in Ofv. af. Vet. Akad. Forh. 1858, 116 (Proc. Am. Acad. iv, 56); Kongl. Sven. Akad. Handl. vi, 54, f. 35; De Candolle, Prodr. xvi³, 214.—Walpers, Ann. v, 745.—Lesquereux in Owen's 2d Rep. Arkansas, 389.—Wood, Cl. Book, 653; Bot. & Fl. 310.—Engelmann in Proc. Am. Phil. Soc. new ser. xii, 209.—Gray, Manual N. States, 5 ed. 465.—Watson in King's Rep. v, 324; Wheeler's Rep. 1872, 493.—Gray in Proc. Am. Acad. vii, 402.—Macoun in Geological Rep. Canada, 1875-76, 210.—Vasey, Cat. Forest Trees, 29.—Hall in Coulter's Bot. Gazette, ii, 91.—Bebb in Wheeler's Rep. vi, 240; Bot. California, ii, 84.—Ward in Bull. U. S. Nat. Mus. No. 22, 116.

S. fluviatilis, Nuttall, Sylva, i, 73; 2 ed. i, 89.

? *S. rubra*, Richardson, Arctic Exped. Appx. 37.

S. longifolia, var. *pedicellata*, Andersson in Kongl. Sven. Akad. Handl. vi, 55, f. 35; De Candolle, Prodr. xvi³, 214.—Macoun in Geological Rep. Canada, 1875-76, 210.

SAND-BAR WILLOW.

Valley of the Connecticut river (Sunderland, Massachusetts, *N. G. Jesup*) and of the Potomac river at Washington (*Ward*); west and northwest through the region of the great lakes to the valley of the Mackenzie river, in latitude 66° N. (*Richardson*), through the Mississippi basin, Texas, the Rocky Mountain region, and the Pacific Coast states.

A small tree, 6 to 9 meters in height, with a trunk rarely exceeding 0.30 meter in diameter; borders of streams and river sand-bars, in low, wet sandy soil, often forming low, dense clumps; rare east of the Alleghany mountains; very common throughout the Mississippi River basin, and reaching its greatest development in the valleys of Oregon and northern California.

Well-marked forms, varying from the type in the form of the leaves, aments, and nature of pubescens, etc., are—

var. *exigua*, Bebb in Bot. California, ii, 85.

S. exigua, Nuttall, Sylva, i, 75; 2 ed. i, 90.

S. longifolia, var. *angustissima*, Andersson in Ofv. af. Vet. Akad. Forh. 1858, 116 (Proc. Am. Acad. iv, 56).

Western Texas to California and Oregon.

var. *argyrophylla*, Andersson in Kongl. Sven. Akad. Handl. vi, 55; De Candolle, Prodr. xvi³, 214.—Watson in King's Rep. v, 324.—Porter in Hayden's Rep. 1872, 493.—Rothrock, Pl. Wheeler, 50.—Porter & Coulter, Fl. Colorado; Hayden's Surv. Misc. Pub. No. 4, 128.—Macoun in Geological Rep. Canada, 1875-76, 210.—Bebb in Bot. California, ii, 85.

S. argyrophylla, Nuttall, Sylva, i, 71, t. 20; 2 ed. i, 87, t. 20.

? *S. brachycarpa*, Nuttall, Sylva, i, 69; 2 ed. i, 85.

S. longifolia, var. *opaca*, Andersson in Kongl. Sven. Akad. Handl. vi, 55.

S. longifolia, var. *argyrophylla angustissima*, Andersson in Kongl. Sven. Akad. Handl. vi, 55; De Candolle, Prodr. xvi³, 214.

S. longifolia, var. *argyrophylla opaca*, Andersson in De Candolle, Prodr. xvi³, 214.

Western Texas to Oregon.

Wood light, soft, very close-grained, compact; medullary rays numerous, very obscure; color, brown tinged with red, the sap-wood brown; specific gravity, 0.4930; ash, 0.48. Var. *exigua*, heavier, the heart- and sap-wood darker colored; specific gravity, 0.5342; ash, 1.06.

311.—*Salix sessilifolia*, Nuttall,

Sylva, i, 68; 2 ed. i, 84.—Andersson in Ofv. af. Vet. Akad. Forh. 1858, 116 (Proc. Am. Acad. iv, 56); Kongl. Sven. Akad. Handl. vi, 55, f. 36; De Candolle, Prodr. xvi³, 214.—Walpers, Ann. v, 746.—Bebb in Bot. California, ii, 85.

S. sessilifolia, var. *villosa*, Andersson in De Candolle, Prodr. xvi³, 215.

Puget sound southward to northern California, near the coast.

A small tree, 9 to 12 meters in height, with a trunk rarely exceeding 0.30 to 0.45 meter in diameter; borders of streams, in low, wet ground.

A form with narrower entire leaves, of the Sacramento valley and the California Coast ranges, is—

var. *Hindsiana*, Andersson in Ofv. af. Vet. Akad. Forh. 1858, 117 (Proc. Am. Acad. iv, 56).—Bebb in Bot. California, ii, 85.

S. Hindsiana, Benthams, Fl. Hartweg. 335.—Newberry in Pacific R. R. Rep. vi, 89.—Torrey in Pacific R. R. Rep. iv, 138.—
*Andersson in Kongl. Sven. Akad. Handl. vi, 56, f. 37; De Candolle, Prodr. xvi², 215.—Walpers, Ann. v, 746.

S. Hindsiana, var. *tenuifolia*, Andersson in Kongl. Sven. Akad. Handl. vi, 56; De Candolle, Prodr. xvi², 215.

Wood light, soft, close-grained, compact; medullary rays thin; color, light red, the sap-wood nearly white; specific gravity, 0.4397; ash, 0.50.

312.—*Salix discolor*, Muhlenberg,

Neue Schriften Gesell. Nat. Fr. Berlin, iv, 234, t. 5, f. 1 (Ann. Bot. ii, 62, t. 5, f. 1).—Willdenow, Spec. iv, 665.—Persoon, Syn. ii, 599.—Pursh, Fl. Am. Sept. ii, 613.—Poiret, Suppl. v, 56.—Nuttall, Genera, ii, 231.—Elliott, Sk. ii, 669.—Torrey, Compend. Fl. N. States, 369; Fl. N. York, ii, 206.—Sprengel, Syst. i, 104.—Forbes, Sal. Woburn. 279.—Eaton, Manual, 6 ed. 319.—Smith in Rees' Cycl. No. 25.—Darlington, Fl. Cestrica, 3 ed. 257.—Eaton & Wright, Bot. 408.—London, Arboretum, iii, 1530, f. 1317, 1630, f. 147.—Bigelow, Fl. Boston. 3 ed. 392.—Hooker, Fl. Bor.-Am. ii, 147.—Barratt, Sal. Am. No. 3.—Emerson, Trees Massachusetts, 258; 2 ed. i, 296 & t.—Dietrich, Syn. v, 419.—Richardson, Arctic Exped. 312.—Darby, Bot. S. States, 506.—Andersson in Ofv. af. Vet. Akad. Forh. 1858, 114 (Proc. Am. Acad. iv, 63); Kongl. Sven. Akad. Handl. vi, 83, f. 49; De Candolle, Prodr. xvi², 225.—Walpers, Ann. v, 750.—Chapman, Fl. S. States, 430.—Gray, Manual N. States, 5 ed. 462.—Koch, Dendrologie, ii, 570.—Macoun in Geological Rep. Canada, 1874-'75, 210.—Ridgway in Proc. U. S. Nat. Mus. 1882, 86.

S. sensitiva, Barratt, Sal. Am. No. 8.

GLAUCOUS WILLOW.

Labrador, west to the valleys of the Peace and Athabasca rivers, southward through the Atlantic region to Delaware and southern Missouri.

A small tree, rarely exceeding 6 meters in height, with a trunk sometimes 0.30 meter in diameter, or more often a tall, straggling shrub 3 to 6 meters in height; along streams and borders of swamps in low, wet soil; varying greatly in the form of leaves, aments, and nature of pubescence.

The best marked forms are—

var. *eriocephala*, Andersson in Kongl. Sven. Akad. Handl. vi, 85; De Candolle, Prodr. xvi², 225.—Gray, Manual N. States, 5 ed. 463.

S. eriocephala, Michaux, Fl. Bor.-Am. ii, 225.—Lamarck, Dict. vi, 661.—Bigelow, Fl. Boston. 3 ed. 391.—Eaton, Manual, 6 ed. 321.—Eaton & Wright, Bot. 409.—Emerson, Trees Massachusetts, 1 ed. 259; 2 ed. i, 196 & t.—Carey in Gray's Manual N. States, 1 ed. 426.—Andersson in Ofv. af. Vet. Akad. Forh. 1858, 117 (Proc. Am. Acad. iv, 57).—Walpers, Ann. v, 746.

S. crassa, Barratt, Sal. Am. No. 7.

var. *prinoides*, Andersson in Kongl. Sven. Akad. Handl. vi, 86; De Candolle, Prodr. xvi², 225.—Emerson, Trees Massachusetts, 2 ed. i, 297.

S. prinoides, Pursh, Fl. Am. Sept. ii, 613.—Nuttall, Genera, ii, 231.—Sprengel, Syst. i, 102.—Poiret, Suppl. iv, 67.—Torrey, Compend. Fl. N. States, 366.—Smith in Rees' Cycl. No. 26.—Forbes, Sal. Woburn. 79, t. 40.—Eaton, Manual, 6 ed. 319.—Beck, Bot. 319.—Eaton & Wright, Bot. 407.—W. Koch, Comment. 46.—London, Arboretum, iii, 1530, f. 1317, 1612, f. 40.—Hooker, Fl. Bor.-Am. ii, 150.—Emerson, Trees Massachusetts, 1, ed. 259.—Dietrich, Syn. v, 419.

Wood light, soft, close-grained, compact, containing many evenly-distributed, small, open ducts; medullary rays and layers of annual growth not obscure; color, brown streaked with orange, the sap-wood light brown; specific gravity, 0.4261; ash, 0.43.

313.—*Salix flavescens*, Nuttall,

Sylva, i, 65; 2 ed. i, 81.—Bebb in Bot. California, ii, 86, in part.

Rocky mountains of Idaho and Montana southward to the Mogollon range, New Mexico (*E. L. Greene*); on the Cascade mountains, Oregon, and the Sierra Nevada, California.

A small tree, sometimes 6 to 9 meters in height, with a trunk rarely 0.30 meter in diameter; borders of streams, reaching its greatest development in the southern Rocky Mountain region.

Wood light, soft, not strong, close-grained, compact; medullary rays numerous, obscure; color, brown tinged with red, the sap-wood nearly white; specific gravity, 0.4969; ash, 0.61.

Var. *Scouleriana*, Bebb;

Coulter's Bot. Gazette, vii, 129.

S. brachystachys, Bentham, Pl. Hartweg. 336.—Andersson in Ofv. af. Vet. Akad. Forh. 1858, 121 (Proc. Am. Acad. iv, 61); Kongl. Sven. Akad. Handl. vi, 82, f. 48; De Candolle, Prodr. xvi³, 224.

S. Scouleriana, Barratt in Hooker, Fl. Bor.-Am. ii, 145, in part.—Cooper in Pacific R. R. Rep. xii³, 29.

S. brachystachys, var. *Scouleriana*, Andersson in De Candolle, Prodr. xvi³, 224.

S. flavescens, Bebb in Bot. California, ii, 86, in part.

BLACK WILLOW.

Kadiak island, Alaska (*Kellogg*), southward through British Columbia, western Washington territory, and Oregon to Santa Barbara, California.

A small tree, 8 to 9 meters in height, with a trunk rarely 0.60 meter in diameter; uplands, near springs or streams, or often in quite dry soil; common and reaching its greatest development near the shores of Puget sound.

Wood light, hard, strong, tough, close-grained, compact; medullary rays numerous, very obscure; color, light red, the sap-wood brown; specific gravity, 0.5412; ash, 0.39.

314.—*Salix Hookeriana*, Barratt;

Hooker, Fl. Bor.-Am. ii, 145, t. 180.—Nuttall, Sylva, i, 64; 2 ed. i, 80.—Andersson in Ofv. af. Vet. Akad. Forh. 1858, 119 (Proc. Am. Acad. iv, 59); De Candolle, Prodr. xvi³, 274.—Walpers, Ann. v, 747.—Macoun in Geological Rep. Canada, 1875-'76, 210.

Grand rapids of the Saskatchewan (*Douglas*); coast of Washington territory and Oregon.

A small tree, 8 to 9 meters in height, with a trunk rarely 0.30 meter in diameter, or more often a low, straggling shrub with many prostrate stems; on the coast generally along the edge of sea-beaches, or in low, rather moist, sandy soil.

Wood light, soft, close-grained, compact, containing many minute open ducts; medullary rays thin, very obscure; color, light brown tinged with red, the sap-wood nearly white; specific gravity, 0.5350; ash, 0.32.

315.—*Salix cordata*, var. *vestita*, Andersson,

Kongl. Sven. Akad. Handl. vi, 159; De Candolle, Prodr. xvi³, 252.

DIAMOND WILLOW.

Valley of the Missouri river and its tributaries, Fort Osage, Missouri (*Prince Newwied*), Iowa, Nebraska, and westward to about the one hundred and tenth degree of longitude.

A small tree, rarely 8 meters in height, with a trunk 0.15 to 0.20 meter in diameter, or more often a straggling shrub not exceeding 1.80 to 3 meters in height; low bottom lands, in wet, sandy soil.

Wood light, soft, close-grained, compact, the annual layers of growth clearly defined; medullary rays very obscure; color, brown or often tinged red, the sap-wood nearly white; specific gravity, 0.6069; ash, 0.59; heavier than that of other species examined, and largely used for fence posts, being said to equal, when thoroughly seasoned, red cedar in durability in contact with the soil.

NOTE.—The typical *Salix cordata*, Muhlenberg, of wide distribution through the Atlantic region, rarely, if ever, attains arborescent size or habit.

316.—*Salix lasiolepis*, Bentham,

Pl. Hartweg. 335.—Cooper in Smithsonian Rep. 1858, 261.—Andersson in Ofv. af. Vet. Akad. Forh. 1858, 118 (Proc. Am. Acad. iv, 58); De Candolle, Prodr. xvi³, 264.—Walpers, Ann. v, 747.—Vasey, Cat. Forest Trees, 29.—Bebb in Bot. California, ii, 86.

S. lasiolepis, var. *Bigelovii*, Bebb in Bot. California, ii, 86 (a vernal state, *teste* Bebb in lit.).

S. Bigelovii, Torrey in Pacific R. R. Rep. iv, 139.—Andersson in Ofv. af. Vet. Akad. Forh. 1858, 118 (Proc. Am. Acad. iv, 58); Kongl. Sven. Akad. Handl. vi, 163, f. 94; De Candolle, Prodr. xvi³, 255.—Walpers, Ann. v, 747.

S. Bigelovii, var. *fuscior*, Andersson in Kongl. Sven. Akad. Handl. vi, 163; De Candolle, Prodr. xvi³, 255.

S. ———, ? Watson in King's Rep. v, 325.

S. lasiolepis, var. *fallax*, Bebb in Bot. California, ii, 86.

WILLOW.

California, valley of the Klamath river, southward through the western portions of the state, reaching in the Sierra Nevadas an elevation of 3,500 to 4,000 feet above the sea.

A small tree, sometimes 12 to 18 meters in height, with a trunk 0.45 to 0.50 meter in diameter, or northward and at high elevations reduced to a low shrub; leaves varying greatly in shape and breadth (vars. *angustifolia* and *latifolia*, Andersson in *De Candolle Prodr.* xvi², 255), or toward its southern limit often persistent until spring (*S. Hartwegi*, Benth in *Pl. Hartwegi*, 52; *S. humilis*, var. *Hartwegi*, Andersson, l. c. 236).

Wood light, soft, not strong, close-grained, compact; medullary rays numerous, thin; color, light brown, the sap-wood nearly white; specific gravity, 0.5587; ash, 0.98; somewhat used as fuel, especially in the southern part of the state.

317.—*Salix Sitchensis*, Sanson;

Bongard in *Mem. Acad. St. Petersburg*, 6 ser. ii, 162.—Ledebour, *Fl. Rossica*, iii, 609.—Richardson, *Arctic Exped.* 439.—Andersson in *Ofv. af. Vet. Akad. Forh.* 1858, 126 (*Proc. Am. Acad.* iv, 66); *Kongl. Sven. Akad. Handl.* vi, 106, f. 59; *De Candolle, Prodr.* xvi², 233.—Walpers, *Ann.* v, 752.—Gray in *Proc. Am. Acad.* vii, 402.—Hall in *Coulter's Bot. Gazette*, ii, 93.—Bebb in *Bot. California*, ii, 87; *Coulter's Bot. Gazette*, vii, 25.

S. cuneata, Nuttall, *Sylva*, i, 66; 2 ed. i, 82.

SILKY WILLOW.

Alaska, southward near the coast to Santa Barbara, California.

A low, much-branched tree, rarely exceeding 8 meters in height, with a trunk 0.30 to 0.45 meter in diameter, or more often a straggling shrub; low, wet soil, borders of streams and ponds.

A form with narrow oblanceolate leaves is—

var. *angustifolia*, Bebb in *Bot. California*, ii, 87.

S. chlorophylla, var. *pellita*, Andersson in *Kongl. Sven. Akad. Handl.* 139, f. 72; *De Candolle, Prodr.* xvi², 244.

Wood light, soft, close-grained, compact; medullary rays numerous, thin; color, light red, the sap-wood nearly white; specific gravity, 0.5072; ash, 0.59.

318.—*Populus tremuloides*, Michaux.

Fl. Bor.-Am. ii, 243.—Nouveau Duhamel, ii, 184, t. 53.—Persoon, *Syn.* ii, 623.—Desfontaines, *Hist. Arb.* ii, 465.—Michaux f. *Hist. Arb.-Am.* iii, 285, t. 8, f. 1; *N. American Sylva*, 3 ed. ii, 175, t. 99, f. 1.—Poiret, *Suppl.* iv, 377.—Willdenow, *Enum. Suppl.* 67.—Torrey, *Ann. Lyc. N. York*, ii, 249; *Compend. Fl. N. States*, 375; *Fremont's Rep.* 97; *Fl. N. York*, ii, 214; *Sitgreaves' Rep.* 172; *Ives' Rep.* 27; *Bot. Wilkes Exped.* 468.—Beck, *Bot.* 323.—Darlington, *Fl. Cestrica*, 3 ed. 281.—Eaton, *Manual*, 117; 6 ed. 277.—Lindley, *Fl. Med.* 320.—Hooker, *Fl. Bor.-Am.* ii, 154.—Eaton & Wright, *Bot.* 370.—Bigelow, *Fl. Boston*, 3 ed. 397.—Spach in *Ann. Sci. Nat.* 2 ser. xv, 30; *Hist. Veg.* x, 384.—Nuttall, *Sylva*, i, 55; 2 ed. i, 70.—Seringe, *Fl. des Jard.* ii, 56.—Parry in *Owen's Rep.* 618.—Newberry in *Pacific R. R. Rep.* vi, 25, 89.—Cooper in *Smithsonian Rep.* 1858, 257; *Pacific R. R. Rep.* xii², 29, 68; *Am. Nat.* iii, 409.—Hooker f. in *Trans. Linnæan Soc.* xxiii², 301.—Wood, *Cl. Book*, 655; *Bot. & Fl.* 311.—Engelmann in *Trans. Am. Phil. Soc.* new ser. xii, 209.—Gray, *Manual N. States*, 5 ed. 466.—Wesmæl in *De Candolle, Prodr.* xvi², 325.—London Gard. *Chronicle*, 1871, 683.—Watson in *King's Rep.* v, 327; *Pl. Wheeler*, 17; *Am. Jour. Sci.* 3 ser. xv, 135; *Bot. California*, ii, 91.—Porter in *Hayden's Rep.* 1871, 494.—Porter & Coulter, *Fl. Colorado*; *Hayden's Surv. Misc. Pub. No. 4*, 128.—Hayden in *Warren's Rep. Nebraska & Dakota*, 2 ed. 121.—Vasey, *Cat. Forest Trees*, 29.—Hall in *Coulter's Bot. Gazette*, ii, 91.—Macoun in *Geological Rep. Canada*, 1875-'76, 210.—Rothrock in *Wheeler's Rep.* vi, 51.—Beal in *Am. Nat.* xv, 32, f. 1.—Trelease in *Coulter's Bot. Gazette*, vi, 284, f. 6.—Scars in *Bull. Essex Inst.* xiii, 183.—G. M. Dawson in *Canadian Nat. new ser.* ix, 231.—Ridgway in *Proc. U. S. Nat. Mus.* 1882, 87.

P. trepida, Willdenow, *Spec.* iv, 803.—Aiton, *Hort. Kew.* 2 ed. 395.—Pursh, *Fl. Am. Sept.* ii, 618.—Eaton, *Manual*, 117.—Nuttall, *Genera*, ii, 239.—Sprengel, *Syst.* ii, 244.—London, *Arboretum*, iii, 1649, f. 1510.

P. tremuliformis, Emerson, *Trees Massachusetts*, 243; 2 ed. i, 279 & t.

P. Atheniensis, Hort.—Koch, *Dendrologie*, ii, 486 (excl. syn.).

ASPEN. QUAKING ASP.

Northern Newfoundland and Labrador to the southern shores of Hudson bay, northwest to the Great Bear lake, the mouth of the Mackenzie river, and the valley of the Yukon river, Alaska; south in the Atlantic region to the mountains of Pennsylvania, the valley of the lower Wabash river, and northern Kentucky; in the Pacific region south to the valley of the Sacramento river, California, and along the Rocky mountains and interior ranges to southern New Mexico, Arizona, and central Nevada.

A small tree, 15 to 18 meters in height, with a trunk rarely exceeding 0.60 meter in diameter; very common through British America and spreading over enormous areas bared by fire of the coniferous forest; in the Pacific region very common upon moist mountain slopes and bottoms at an elevation of 6,000 to 10,000 feet; the most widely-distributed North American tree.

Wood light, soft, not strong, close-grained, compact, not durable, containing, as does that of the whole genus, numerous minute, scattered, open ducts; medullary rays very thin, hardly distinguishable; color, light brown, the thick sap-wood nearly white; specific gravity, 0.4032; ash, 0.55; largely manufactured into wood-pulp, a substitute for rags in the manufacture of paper; in the Pacific region sometimes used for fuel, flooring, in turnery, etc.

A bitter principle in the bark causes its occasional use as a tonic in the treatment of intermittent fevers and cases of debility (*U. S. Dispensatory*, 14 ed. 1763).

319.—*Populus grandidentata*, Michaux,

Fl. Bor.-Am. ii, 243.—Persoon, Syn. ii, 624.—Desfontaines, Hist. Arb. ii, 466.—Michaux f. Hist. Arb. Am. iii, 287, t. 8, f. 2; N. American Sylva, 3 ed. ii, 176, t. 99, f. 2.—Pursh, Fl. Am. Sept. ii, 619.—Poiret, Suppl. iv, 377.—Barton, Compend. Fl. Philadelph. ii, 197.—Nuttall, Genera, ii, 239.—Hayne, Dend. Fl. 200.—Willdenow, Enum. Suppl. 67.—Elliott, Sk. ii, 710.—Sprengel, Syst. ii, 244.—Torrey, Compend. Fl. N. States, 375; Fl. N. York, ii, 214.—Beck, Bot. 323.—Eaton, Manual, 6 ed. 277.—Hooker, Fl. Bor.-Am. ii, 154.—Eaton & Wright, Bot. 370.—London, Arboretum, iii, 1650, f. 1511.—Bigelow, Fl. Boston. 3 ed. 397.—Spach in Ann. Sci. Nat. xv, 2 ser. 33; Hist. Veg. x, 384.—Emerson, Trees Massachusetts, 242; 2 ed. i, 278 & t.—Seringe in Fl. des Jard. ii, 56.—Parry in Owen's Rep. 618.—Darlington, Fl. Cestrica, 3 ed. 281.—Darby, Bot. S. States, 507.—Cooper in Smithsonian Rep. 1858, 257.—Chapman, Fl. S. States, 431.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 73.—Wood, Cl. Book, 656; Bot. & Fl. 311.—Gray, Manual N. States, 5 ed. 466.—Koch, Dendrologie, ii, 487.—Wesmæl in De Candolle, Prodr. xvi², 327.—Vasey, Cat. Forest Trees, 29.—Watson in Am. Jour. Sci. 3 ser. xv, 135.—Beal in Am. Nat. xv, 34, f. 2.—Sears in Bull. Essex Inst. xiii, 182.—Trelease in Coulter's Bot. Gazette, vi, 285.—Bell in Geological Rep. Canada, 1879-'80, 56^c.

P. grandidentata, var. *pendula*, Torrey, Compend. Fl. N. States, 375.—Nuttall, Genera, ii, 239.

POPLAR.

Nova Scotia, New Brunswick, and west through Ontario to northern Minnesota, south through the northern states and along the Alleghany mountains to North Carolina, extending west to middle Kentucky and Tennessee.

A tree 21 to 24 meters in height, with a trunk 0.50 to 0.75 meter in diameter; rich woods and borders of streams and swamps.

Wood light, soft, not strong, close-grained, compact; medullary rays thin, obscure; color, light brown, the sap-wood nearly white; specific gravity, 0.4632; ash, 0.45; largely manufactured into wood-pulp and occasionally used in turnery, for woodenware, etc.

320.—*Populus heterophylla*, Linnæus,

Spec. 1 ed. 1034.—Marshall, Arbustum, 107.—Wangenheim, Amer. 85.—Walter, Fl. Caroliniana, 248.—Aiton, Hort. Kew. iii, 407; 2 ed. v, 397.—Nouveau Duhamel, ii, 181, t. 51.—Michaux, Fl. Bor.-Am. ii, 244.—Willdenow, Spec. iv, 806; Enum. 1017; Berl. Baumz. 293.—Desfontaines, Hist. Arb. ii, 466.—Pursh, Fl. Am. Sept. ii, 619.—Nuttall, Genera, ii, 239.—Hayne, Depd. Fl. 203.—Elliott, Sk. ii, 712.—Sprengel, Syst. ii, 244.—Torrey, Compend. Fl. N. States, 375; Fl. N. York, ii, 215.—Beck, Bot. 323.—Eaton, Manual, 6 ed. 278.—Darlington, Fl. Cestrica, 3 ed. 281.—London, Arboretum, iii, 1672, f. 1534.—Eaton & Wright, Bot. 371.—Spach in Ann. Sci. Nat. 2 ser. xv, 30; Hist. Veg. x, 386.—Seringe in Fl. des Jard. ii, 61.—Darby, Bot. S. States, 507.—Cooper in Smithsonian Rep. 1858, 257.—Chapman, Fl. S. States, 431.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 73.—Wood, Cl. Book, 656; Bot. & Fl. 311.—Gray, Manual N. States, 5 ed. 467.—Koch, Dendrologie, ii, 488.—Wesmæl in De Candolle, Prodr. xvi², 326.—Vasey, Cat. Forest Trees, 29.—Watson in Am. Jour. Sci. 3 ser. xv, 135.—Trelease in Coulter's Bot. Gazette, vi, 285.—Ridgway in Proc. U. S. Nat. Mus. 1881, 86.

P. cordifolia, Burgsdorf, Anleit. Erz. Holzart. 3 ed. 152.

P. argentea, Michaux f. Hist. Arb. Am. iii, 390, t. 9; N. American Sylva, 3 ed. ii, 170, t. 97.

P. heterophylla, var. *argentea*, Wesmæl in De Candolle, Prodr. xvi², 376.

RIVER COTTONWOOD. SWAMP COTTONWOOD.

Guilford, Connecticut (*W. R. Dudley*), Northport, Long island, south, generally near the coast, to southern Georgia, through the Gulf states to western Louisiana, and through Arkansas to central Tennessee and Kentucky, southern Illinois and Indiana.

A tree 24 to 27 meters in height, with a trunk 0.60 to 0.75 meter in diameter; borders of river swamps; most common and reaching its greatest development in the basin of the lower Ohio river; rare and local.

Wood light, soft, not strong, close-grained, compact; medullary rays thin, very obscure; color, dull brown, the thick sap-wood lighter brown; specific gravity, 0.4089; ash, 0.81.

321.—*Populus balsamifera*, Linnæus,

Spec. 1 ed. 1034.—Du Roi, Harbk. 82.—Marshall, Arbustum, 107.—Wangenheim, Amer. 85, t. 28, f. 59.—Aiton, Hort. Kew. iii, 406; 2 ed. v, 397.—Mœnch, Meth. 338.—B. S. Barton, Coll. i, 16.—Nouveau Duhamel, ii, 179, t. 50.—Michaux, Fl. Bor.-Am. ii, 244.—Willdenow, Spec. iv, 805; Enum. 1017; Berl. Baumz. 290.—Persoon, Syn. ii, 624.—Desfontaines, Hist. Arb. ii, 466.—Michaux f. Hist. Arb. Am. iii, 306, t. 13, f. 1; N. American Sylva, 3 ed. ii, 172, t. 98, f. 1.—Pursh, Fl. Am. Sept. ii, 618.—Eaton, Manual, 117; 6 ed. 278.—Nuttall, Genera, ii, 239; Sylva, i, 55; 2 ed. i, 70.—Hayne, Dend. Fl. 202.—Sprengel, Syst. ii, 244.—Beck, Bot. 322.—Lindley, Fl. Med. 320.—London, Arboretum, iii, 1637, f. 1535, 1536 & t.—Hooker, Fl. Bor.-Am. ii, 153.—Eaton & Wright, Bot. 370.—Hooker & Arnott, Bot. Beechey, 159.—Spach in Ann. Sci. Nat. 2 ser. xv, 33; Hist. Veg. x, 393.—Lindley, Bot. Reg. xxix, Misc. 20.—Seringe in Fl. des Jard. ii, 65.—Torrey, Fl. N. York, ii, 216; Bot. Wilkes Exped. 469.—Cooper in Smithsonian Rep. 1858, 257; Am. Nat. iii, 408.—Hooker f. in Trans. Linnæan Soc. xxiii², 301.—Wood, Cl. Book, 656; Bot. & Fl. 311.—Gray, Manual N. States, 5 ed. 467.—Koch, Dendrologie, ii, 495.—Vasey, Cat. Forest Trees, 29.—Macoun in Geological Rep. Canada, 1875-'76, 211.—Watson in Am. Jour. Sci. xv, 135.—Beal in Am. Nat. xv, 34, f. 4.—Trelease in Coulter's Bot. Gazette, vi, 285.—Sears in Bull. Essex Inst. xiii, 181.—Bell in Geological Rep. Canada, 1879-'80, 45^c.

P. Tacamahaca, Miller, Dict.

P. viminea, Bon Jard. 1845, 565.

P. balsamifera, var. *genuina*, Wesmæl in De Candolle, Prodr. xvi², 329.

BALSAM. TACAMAHAC. BALM OF GILEAD.

Straits of Belle Isle to Richmond gulf and cape Churchill, Hudson bay, northwest to the shores of the Great Bear lake and the valley of the Yukon river, Alaska, south to northern New England, central Michigan and Minnesota, the Rocky mountains and interior ranges of Montana and Idaho, Washington territory, and British Columbia.

A large tree, 18 to 24 meters in height, with a trunk 1.50 to 2.10 meters in diameter; very common on all islands and shores of the northern rivers; in British Columbia generally confounded with the allied *P. trichocarpa*, the range of the two species here still uncertain.

Wood very light, soft, not strong, close-grained, compact; medullary rays numerous, very obscure; color, brown, the thick sap-wood nearly white; specific gravity, 0.3635; ash, 0.66.

The buds, as well as those of several other species, covered with a resinous exudation, and occasionally used medicinally as a substitute for turpentine and other balms.

Var. *candicans*, Gray,

Manual N. States, 2 ed. 419; 5 ed. 467.—Cooper in Smithsonian Rep. 1858, 257.—Porter & Coulter, Fl. Colorado; Hayden's Surv. Misc. Pub. No. 4, 129.—Watson in Am. Jour. Sci. 3 ser. xv, 135.—Bull. Torrey Bot. Club, vii, 57.—Trelease in Coulter's Bot. Gazette, vi, 285.

P. balsamifera lanceolata, Marshall, Arbustum, 108.

P. candicans, Aiton, Hort. Kew. iii, 406; 2 ed. v, 397.—Nouveau Duhamel, ii, 179.—Willdenow, Spec. iv, 806; Enum. 1017; Berl. Baumz. 291.—Persoon, Syn. ii, 624.—Michaux f. Hist. Arb. Am. iii, 308, t. 13, f. 2; N. American Sylva, 3 ed. ii, 173, t. 98, f. 2.—Pursh, Fl. Am. Sept. ii, 618.—Barton, Prodr. Fl. Philadelph. 96.—Poiret, Suppl. iv, 378.—Nuttall, Genera, ii, 239.—Hayne, Dend. Fl. 202.—Sprengel, Syst. ii, 244.—Torrey, Compend. Fl. N. States, 375; Fl. N. York, ii, 217.—Audubon, Birds, t. 59.—Beck, Bot. 332.—Eaton, Manual, 6 ed. 278.—London, Arboretum, ii, 1676, f. 1537.—Hooker, Fl. Bor.-Am. ii, 154.—Eaton & Wright, Bot. 370.—Bigelow, Fl. Boston. 3 ed. 398.—Spach in Ann. Sci. Nat. 2 ser. xv, 33; Hist. Veg. x, 392.—Lindley, Bot. Reg. xxix, Misc. 22.—Emerson, Trees Massachusetts, 245; 2 ed. i, 281.—Seringe in Fl. des Jard. ii, 63.—Gray, Manual N. States, 1 ed. 431.—Wood, Cl. Book, 656; Bot. & Fl. 311.—Wesmæl in De Candolle, Prodr. xvi², 330.

P. Canadensis, Mœnch, Weiss. 81 [not Michaux f.].

P. latifolia, Mœnch, Meth. 338.

P. Ontariensis, Hort.—Loddiges, Cat. 1836.

P. macrophylla, Lindley in London, Encyc. Pl. 840.

P. aeladesca and *P. heterophylla*, Hort. (ex. Koch, Wachen. 1865, 238).

A large tree, rare or unknown in a wild state; very common in cultivation.

The wood heavier than that of the species; specific gravity, 0.4161; ash, 0.46.

322.—*Populus angustifolia*, James,

Long's Exped. i, 497.—Torrey in Ann. Lyc. N. York, ii, 249; Fremont's Rep. 97; Sitgreaves' Rep. 172; Ives' Rep. 27; Bot. Wilkes Exped. 469.—Nuttall, Sylva, i, 52, t. 16; 2 ed. i, 68, t. 16.—Cooper in Smithsonian Rep. 1858, 261; Am. Nat. iii, 408.—Hayden in Warren's Rep. Nebraska & Dakota, 2 ed. 121.—Vasey, Cat. Forest Trees, 29.—Watson in Am. Jour. Sci. 3 ser. xv, 136; Bot. California, ii, 91.

P. Canadensis, var. *angustifolia*, Wesmæl in De Candolle, Prodr. xvi², 329.

P. balsamifera, var. *angustifolia*, Watson in King's Rep. v, 327; Pl. Wheeler, 17.—Porter in Hayden's Rep. 1871, 494.—Porter & Coulter, Fl. Colorado; Hayden's Surv. Misc. Pub. No. 4, 128.—Macoun in Geological Rep. Canada, 1875-'76, 211.—Rusby in Bull. Torrey Bot. Club, ix, 106.

BLACK COTTONWOOD.

Black hills of Dakota (*R. Douglas*), Swimming Horse creek, and the Snowy Mountain region, Montana, Red Rock creek, southwestern Montana (*Watson*), east Humboldt and Shoshone mountains, Nevada, Rocky mountains of Colorado, and the ranges of southwestern New Mexico and eastern Arizona.

A small tree, 15 to 18 meters in height, with a trunk rarely exceeding 0.60 meter in diameter; borders of streams, between 6,000 and 10,000 feet elevation.

Wood light, soft, weak, close-grained, compact; medullary rays numerous, obscure; color, brown, the sap-wood nearly white; specific gravity, 0.3912; ash, 0.79.

323.—*Populus trichocarpa*, Torrey & Gray;

Hooker, Icon. v, 878.—Walpers, Ann. v, 767.—Cooper in Smithsonian Rep. 1858, 266.—Wesmæl in De Candolle, Prodr. xvi², 330.—Watson in King's Rep. v, 328; Am. Jour. Sci. 3 ser. xv, 136; Bot. California, ii, 91.—Torrey, Bot. Wilkes Exped. 469.—Macoun in Geological Rep. Canada, 1875-'76, 211.—Trelease in Coulter's Bot. Gazette, vi, 285, f. 5.—G. M. Dawson in Canadian Nat. new ser. ix, 331.

P. balsamifera, var. Hooker, Fl. Bor.-Am. ii, 154.

P. angustifolia, Newberry in Pacific R. R. Rep. vi, 89 [not James].—Cooper in Pacific R. R. Rep. xii², 29, 68.

P. balsamifera, Lyall in Jour. Linnæan Soc. vii, 134 [not Linnæus].—Hall in Coulter's Bot. Gazette, ii, 91.

P. trichocarpa, var. *cupulata*, Watson in Am. Jour. Sci. 3 ser. xv, 136; Bot. California, ii, 91.

P. balsamifera, var. ? *Californica*, Watson in Am. Jour. Sci. 3 ser. xv, 136.

BLACK COTTONWOOD. BALSAM COTTONWOOD.

Valley of the Fraser river, British Columbia, and probably much farther north, east to the eastern base of the Bitter Root mountains, Montana (*Watson*), south through Washington territory, western Oregon and California to the southern borders of the state.

A large tree, 24 to 60 meters in height, with a trunk 1.20 to 2.10 meters in diameter; banks of streams and bottom lands below 6,000 feet elevation; very common and reaching its greatest development in the valleys of the lower Columbia river and the streams flowing into Puget sound, here the largest deciduous tree of the forest.

Wood very light, soft, not strong, rather close-grained, compact; medullary rays thin, hardly distinguishable; color, light dull brown, the sap-wood lighter, nearly white; specific gravity, 0.3814; ash, 1.27; in Oregon and Washington territory largely manufactured into staves of sugar barrels, woodenware, etc.

324.—*Populus monilifera*, Aiton,

Hort. Kew. iii, 406; 2 ed. v, 396.—Abbot, Insects Georgia, ii, 71.—Nouveau Duhamel, ii, 186.—Willdenow, Spec. iv, 805; Enum. 1017; Berl. Baumz. 292.—Persoon, Syn. ii, 623.—Desfontaines, Hist. Arb. ii, 465.—Michaux f. Hist. Arb. Am. iii, 295, t. 10, f. 2; N. American Sylva, 3 ed. ii, 168, t. 96, f. 2.—Pursh, Fl. Am. Sept. ii, 618.—Nuttall, Genera, ii, 239; Trans. Am. Phil. Soc. 2 ser. v, 167.—Hayne, Dend. Fl. 202.—Sprengel, Syst. ii, 244.—Watson, Dend. Brit. ii, t. 102.—Beck, Bot. 323.—Eaton, Manual, 6 ed. 278.—London, Arboretum, iii, 1657, f. 1517 & t.—Eaton & Wright, Bot. 371.—Spach in Ann. Sci. Nat. 2 ser. xv, 32; Hist. Veg. x, 389.—Torrey in Fremont's Rep. 97; Fl. N. York, ii, 215; Pacific R. R. Rep. v, 365.—Emerson, Trees Massachusetts, 249; 2 ed. i, 287.—Seringe in Fl. des Jard. ii, 63.—Cooper in Smithsonian Rep. 1858, 257.—Gray in Pacific R. R. Rep. xii², 47; Manual N. States, 5 ed. 467.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 72.—Lesquereux in Owen's 2d Rep. Arkansas, 389.—Wood, Cl. Book, 655.—Engelmann in Trans. Am. Phil. Soc. xii, 209.—Watson in King's Rep. v, 327; Am. Jour. Sci. 3 ser. xv, 136.—Hayden in Warren's Rep. Nebraska & Dakota, 2 ed. 121.—Macoun in Geological Rep. Canada, 1875-'76, 211.—Trelease in Coulter's Bot. Gazette, vi, 285, f. 3, 4.—Ward in Bull. U. S. Nat. Mus. No. 22, 116.—Beal in Am. Nat. xv, 34, f. 3.—Bell in Geological Rep. Canada, 1879-'80, 56c.—Ridgway in Proc. U. S. Nat. Mus. 1882, 87.—Chapman, Fl. S. States, Suppl. 649.

? *P. deltoide*, Marshall, Arbustum, 106.

P. angulata, Aiton, Hort. Kew. iii, 406; 2 ed. v, 396.—Nouveau Duhamel, ii, 186.—Desfontaines, Hist. Arb. ii, 466.—Willdenow, Spec. iv, 805; Enum. 1017; Berl. Baumz. 294.—Michaux f. Hist. Arb. Am. iii, 302, t. 12; N. American Sylva, 3 ed. ii, 161, t. 94.—Pursh, Fl. Am. Sept. ii, 619.—Eaton, Manual, 117; 6 ed. 277.—Nuttall, Genera, ii, 239.—James in Long's Exped. ii, 164.—Torrey in Ann. Lyc. N. York, ii, 249.—Elliott, Sk. ii, 711.—Sprengel, Syst. ii, 244.—London, Arboretum, iii, 1670, 1533 & t.—Eaton & Wright, Bot. 370.—Spach in Ann. Sci. Nat. 2 ser. xv, 321; Hist. Veg. x, 391.—Seringe in Fl. des Jard. ii, 64.—Scheele in Roemer, Texas, 446.—Darby, Bot. S. States, 507.—Cooper in Smithsonian Rep. 1858, 257.—Chapman, Fl. S. States, 431.—Lesquereux in Owen's 2d Rep. Arkansas, 389.—Wood, Cl. Book, 655; Bot. & Fl. 311.—Gray, Manual N. States, 5 ed. 467.—Wesmæl in De Candolle, Prodr. xvi², 328.—Koch, Dendrologie, ii, 494.—Young, Bot. Texas, 514.—Porter & Coulter, Fl. Colorado; Hayden's Surv. Misc. Pub. No. 4, 129.—Vasey, Cat. Forest Trees, 29.—Sears in Bull. Essex Inst. xiii, 182.

P. lævigata, Aiton, Hort. Kew. iii, 406; 2 ed. v, 395.—Willdenow, Spec. iv, 803.—Pursh, Fl. Am. Sept. ii, 619.—Poiret, Suppl. iv, 378.—Nuttall, Genera, ii, 239; Sylva, i, 54; 2 ed. i, 70.—Sprengel, Syst. ii, 244.—Beck, Bot. 323.—Eaton, Manual, 6 ed. 278.—Loddiges, Cat. ed. 1836.—Eaton & Wright, Bot. 370.—Emerson, Trees Massachusetts, 246; 2 ed. i, 283.

P. glandulosa, Mönch, Meth. 339.

P. angulosa, Michaux, Fl. Bor.-Am. ii, 243.

P. Canadensis, Michaux f. Hist. Arb. Am. iii, 302, t. 12; N. American Sylva, 3 ed. ii, 164, t. 95.—Spach in Ann. Sci. Nat. 2 ser. xv, 32; Hist. Veg. x, 390.—Seringe in Fl. des Jard. ii, 65.—Fescali, Forst. Pfl. 122, t. 8, f. 10-14.—Wood, Bot. & Fl. 311.—Wesmæl in De Candolle, Prodr. xvi², 329.—Koch, Dendrologie, ii, 491.

P. Virginiana, Du Mont, Cours. Bot. Cult. vi, 400.

P. Marylandica, Bosc in Nouv. Dict. xi, 409.—Poiret, Suppl. iv, 378.—Sprengel, Syst. ii, 244.

P. macrophylla, Loddiges, Cat. ed. 1836.

P. Lindleyana, *P. neglecta*, and *P. lævigata*, Hort.

COTTONWOOD. NECKLACE POPLAR. CAROLINA POPLAR. BIG COTTONWOOD.

Shores of lake Champlain, Vermont, south through western New England to the Chattahoochee region of western Florida, west along the northern shores of lake Ontario to the eastern base of the ranges of the Rocky mountains of Montana, Colorado, and New Mexico.

A large tree, 24 to 51 meters in height, with a trunk 1.20 to 2.40 meters in diameter; low, moist soil; the common cottonwood of Texas and the western plains, bordering all streams flowing east from the Rocky mountains.

Wood very light, soft, not strong, close-grained, compact, liable to warp in drying, difficult to season; medullary rays numerous, obscure; color, dark brown, the thick sap-wood nearly white; specific gravity, 0.3889; ash, 0.96; largely used in the manufacture of paper-pulp, for light packing-cases, fence boards, and fuel.

325.—*Populus Fremontii*, Watson,

Proc. Am. Acad. x, 350; Am. Jour. Sci. 3 ser. xv, 136; Bot. California, ii, 92.

P. monilifera, Newberry in Pacific R. R. Rep. vi, 327 [not Aiton].—Watson in King's Rep. v, 327; Pl. Wheeler, 17.—Torrey, Bot. Wilkes Exped. 469.

COTTONWOOD.

California, valley of the upper Sacramento river, south to San Bernardino county (Colton, *Parry*), and eastward in Nevada and Utah.

A large tree, 24 to 30 meters in height, with a trunk 1.20 to 1.80 meter in diameter; borders of streams; the common cottonwood of the valleys of central California.

Wood light, soft, not strong, close-grained, compact, liable to warp in drying, difficult to season; medullary rays thin, very obscure; color, light brown, the sap-wood nearly white; specific gravity, 0.4914; ash, 0.77.

Var. *Wislizeni*, Watson,

Am. Jour. Sci. 3 ser. xv, 137; Bot. California, ii, 92; Proc. Am. Acad. xviii, 157.—Rusby in Bull. Torrey Bot. Club, ix, 79.

P. monilifera, Torrey in Sitgreaves' Rep. 172; Bot. Mex. Boundary Survey, 204; Ives' Rep. 27 [not Aiton].—Bigelow in Pacific R. R. Rep. iv, 21.

COTTONWOOD. WHITE COTTONWOOD.

San Diego county, California, through Arizona and New Mexico to western Texas and southern Colorado.

A large tree, 24 to 30 meters in height, with a trunk 1.20 to 1.80 meter in diameter; borders of streams; the prevalent cottonwood of the arid southwestern region, there largely planted as a shade tree and for fuel.

Wood light, soft, not strong, compact; specific gravity, 0.4621; ash, 1.13; furnishing the ordinary domestic fuel of the region.

CONIFERÆ.

326.—*Libocedrus decurrens*, Torrey,

Smithsonian Contrib., vi, 7, t. 3; Pacific R. R. Rep. iv, 140; Bot. Mex. Boundary Survey, 211; Bot. Wilkes Exped. t. 16.—Bentham, Pl. Hartweg, 338.—Lindley in London Gard. Chronicle, 1853, 695.—Newberry in Pacific R. R. Rep. vi, 63.—Cooper in Smithsonian Rep. 1858, 262.—Walpers, Ann. v, 795.—Bolander in Proc. California Acad. iii, 228.—Parlatore in De Candolle, Prodr. xvi², 456.—R. Brown Campst. in Trans. Edinburgh Bot. Soc. ix, 373.—Hoopes, Evergreens, 309, f. 40.—Watson in King's Rep. v, 335; Bot. California, ii, 116.—A. Murray in London Garden, ii, 542.—Gordon, Pinetum, 2 ed. 402.—Veitch, Manual Conif. 267.

Thuja Craigana, Murray in Rep. Oregon Exped. 2, t. 5.

Thuja gigantea, Carrière in Rev. Hort. 1854, 224, f. 12-14, in part; Fl. des Serres, ix, 199, f. 3-5, in part; Trait. Conif. 106, in part; 2 ed. 112, in part.—Gordon, Pinetum, 321, in part; Suppl. 102, in part.—Henkel & Hochstetter, Nadelhölz. 280, in part.

Heyderia decurrens, Koch, Dendrologie, ii², 179.

WHITE CEDAR. BASTARD CEDAR. POST CEDAR. INCENSE CEDAR.

North fork of the Santian river, Oregon, south along the western slopes of the Cascade and Sierra Nevada mountains between 3,000 and 8,500 feet elevation, and through the California Coast ranges to the San Bernardino and Cayumaca mountains.

A large tree, 30 to 45 meters in height, with a trunk 1.20 to 2.10 meters in diameter; slopes and valleys; common.

Wood light, soft, not strong, brittle, close-grained, compact, very durable in contact with the soil; bands of small summer cells thin, dark colored, conspicuous; medullary rays numerous, obscure; the thin sap-wood nearly white; specific gravity, 0.4017; ash, 0.08; largely used for fencing and in the construction of water-flumes, and for interior finish, furniture, laths, shingles, etc.; often injured by a species of dry rot (*Dædalia vorax*, Harkness in *Pacific Rural Press*, Jan. 25, 1879, f. 1, 2), rendering it unfit for lumber.

327.—*Thuja occidentalis*, Linnæus,

Spec. 1 ed. 1002.—Kalm, Travels, English ed. iii, 170.—Marshall, Arbustum, 152.—Wangenheim, Amer. 7, t. 2, f. 3.—Walter, Fl. Caroliniana, 238.—Aiton, Hort. Kew. iii, 371; 2 ed. v, 321.—Gærtner, Fruct. ii, 62, t. 91, f. 2.—Michaux, Fl. Bor.-Am. ii, 209.—Willdenow, Spec. iv, 508; Enum. 990; Berl. Baumz. 504.—Nouveau Duhamel, iii, 12, t. 4.—Poirot in Lamarck, Dict. vii, 369; Ill. iii, 369.—Schkuhr, Handb. iii, 287, t. 309.—Persoon, Syn. ii, 580.—Desfontaines, Hist. Arb. ii, 575.—Titford, Hort. Bot. Am. 98.—Michaux f. Hist. Arb. Am. iii, 29, t. 3; N. American Sylva, 3 ed. iii, 177, t. 156.—Pursh, Fl. Am. Sept. ii, 647.—Barton, Prodr. Fl. Philadelph. 93.—Eaton, Manual, 111; 6 ed. 364.—Nuttall, Genera, ii, 224.—Hayne, Dend. Fl. 177.—Elliott, Sk. ii, 641.—Watson, Dend. Brit. ii, 150.—Sprengel, Syst. iii, 888.—Richard, Conif. 43, t. 71, f. 1.—Torrey, Compend. Fl. N. States, 361; Fl. N. York, ii, 234.—Rafinesque, Med. Bot. ii, 268.—Beck, Bot. 338.—London, Arboretum, iv, 2454, f. 2312-2314 & t.—Forbes, Pinetum Woburn. 193.—Hooker, Fl. Bor.-Am. ii, 165.—Eaton & Wright, Bot. 451.—Bigelow, Fl. Boston. 3 ed. 388.—Spach, Hist. Veg. xi, 339.—Penn. Cycl. xxiv, 409.—Reid in London Gard. Chronicle, 1844, 276.—Emerson, Trees Massachusetts, 96; 2 ed. i, 112.—Endlicher, Syn. Conif. 51.—Lindley & Gordon in Jour. Hort. Soc. London, v, 206.—Parry in Owen's Rep. 618.—Darlington, Fl. Cestrica, 3 ed. 294.—Knight, Syn. Conif. 16.—Carrière in Rev. Hort. 1854, 224, f. 15; Trait. Conif. 103; 2 ed. 109.—Darby, Bot. S. States, 516.—Cooper in Smithsonian Rep. 1858, 257.—Gordon, Pinetum, 323; 2 ed. 403.—Chapman, Fl. S. States, 436.—Wood, Cl. Book, 662; Bot. & Fl. 315.—Porcher, Resources S. Forests, 507.—Henkel & Hochstetter, Nadelhölz. 278.—Nelson, Pinaceæ, 68.—R. Brown Campst. in Trans. Edinburgh Bot. Soc. ix, 363.—Gray, Manual N. States, 5 ed. 472.—Hoopes, Evergreens, 317.—Parlatore in De Candolle, Prodr. xvi², 458.—Schnizlein, Icon. t. 76, f. 2.—Koch, Dendrologie, ii², 173.—Vasey, Cat. Forest Trees, 36.—Macoun in Geological Rep. Canada, 1875-'76, 211.—Sears in Bull. Essex Inst. xiii, 183.—Veitch, Manual Conif. 261.—Bell in Geological Rep. Canada, 1879-'80, 47^c.

T. odorata, Marshall, Arbustum, 152.

T. obtusa, Moench, Meth. 691.

Cupressus Arbor-vitæ, Targione-Tozzetti, Obs. Bot. ii, 51.

T. Wareana and *T. Sibirica*, Hort.

WHITE CEDAR. ARBOR-VITÆ.

New Brunswick to Anticosti island, through the valley of the Saint Lawrence river to the southern shores of James' bay and southeast to the eastern extremity of lake Winnipeg, south through the northern states to central New York, northern Pennsylvania, central Michigan, northern Illinois, central Minnesota, and along the Alleghany mountains to the high peaks of North Carolina.

A tree 12 to 18 meters in height, with a trunk sometimes 1.20 to 1.50 meter in diameter; cold, wet swamps and along the rocky banks of streams; very common at the north, spreading over great areas of swamp; extensively cultivated as a hedge and ornamental plant, and producing innumerable seminal varieties of more or less horticultural value.

Wood very light, soft, not strong, brittle, rather coarse-grained, compact, very durable in contact with the soil; the bands of small summer cells very thin, dark colored; medullary rays numerous, indistinct; color, light brown, turning darker with exposure, the thin sap-wood nearly white; specific gravity, 0.3164; ash, 0.37; largely used for posts, fencing, railway ties, and shingles.

The distilled oil and a tincture of the leaves of *Thuya* have been found useful in the treatment of pulmonary and uterine complaints (*U. S. Dispensatory*, 14 ed. 1775.—*Nat. Dispensatory*, 2 ed. 1428).

328.—*Thuya gigantea*, Nuttall,

Jour. Philadelphia Acad. vii, 52; Sylva, iii, 102, t. iii; 2 ed. ii, 162, t. 111.—Loddiges, Cat. ed. 1836.—London, Arboretum, iv, 2458.—Hooker, Fl. Bor.-Am. ii, 165.—Spach, Hist. Veg. xi, 342.—Endlicher, Syn. Conif. 52.—Lindley & Gordon in Jour. Hort. Soc. London, v, 206.—Newberry in Pacific R. R. Rep. vi, 56, f. 22.—Carrière, Trait. Conif. 102; 2 ed. 112, in part.—Cooper in Smithsonian Rep. 1858, 262; Am. Nat. iii, 413.—Gordon, Pinetum, 321, in part; Suppl. 102; 2 ed. 181.—Torrey, Bot. Mex. Boundary Survey, 211.—Lyll in Jour. Linnæan Soc. vii, 133, 144.—Henkel & Hochstetter, Nadelhölz. 280, in part.—Nelson, Pinaceæ, 67.—Rothrock in Smithsonian Rep. 1867, 434.—Parlatore in De Candolle, Prodr. xvi², 457.—R. Brown Campst. in Trans. Edinburgh Bot. Soc. ix, 367.—Hoopes, Evergreens, 315.—London Gard. Chronicle, 1871, 683.—Gray in Proc. Am. Acad. vii, 402.—Fowler in London Gard. Chronicle, 1872, 1527.—Koch, Dendrologie, ii², 176.—Vasey, Cat. Forest Trees, 36.—E. Hall in Coulter's Bot. Gazette, ii, 91.—Watson, Bot. California, ii, 115.—G. M. Dawson in Canadian Nat. new ser. ix, 324.—T. Howell in Coulter's Bot. Gazette, vi, 267.—Veitch, Manual Conif. 256.

T. plicata, Don, Hort. Cantab. 6 ed. 249.—Lambert, Pinus, 1 ed. ii, 19; 2 ed. 114, in part.—Nuttall, Sylva, iii, 103; 2 ed. ii, 164.—Spach, Hist. Veg. xi, 342.—Endlicher, Syn. Conif. 51 (excl. syn. *Wareana & odorata*).—Lindley & Gordon in Jour. Hort. Soc. London, v, 205.—Knight, Syn. Conif. 16.—Carrière, Trait. Conif. 102 (excl. syn. *Wareana & odorata*); 2 ed. 106 (excl. syn. *Wareana*).—Cooper in Smithsonian Rep. 1858, 262; Pacific R. R. Rep. xii², 27.—Henkel & Hochstetter, Nadelhölz. 277 (excl. syn. *odorata*).—Nelson, Pinaceæ, 68.—Gordon, Pinetum, 2 ed. 406.—A. De Candolle, Prodr. xvi², 457, in part.—Vasey, Cat. Forest Trees, 36.—Veitch, Manual Conif. 263.

T. Menziesii, Douglas, Mss.—Carrière, Trait. Conif. 106; 2 ed. 107.—Gordon, Pinetum, 323.—Nelson, Pinaceæ, 67.—Henkel & Hochstetter, Nadelhölz. 281.

T. Lobbii, Hort.

T. occidentalis, var. *plicata*, Hort.—Hoopes, Evergreens, 321.

RED CEDAR. CANOE CEDAR.

Alaska, south along the Coast ranges and islands of British Columbia, through western Washington territory and Oregon and the Coast ranges of northern California to Mendocino county, extending east along the mountains of Washington territory to the Cœur d'Alène, Bitter Root, and Salmon River mountains of Idaho and the western slopes of the Rocky mountains of northern Montana (*Canby & Sargent*).

A large tree, 30 to 45 meters in height, with a trunk 0.90 to 3.60 meters in diameter; low, rich woods and swamps, less commonly on dry ridges and slopes below 5,200 feet elevation; common and reaching its greatest development in western Washington territory and Oregon; the large specimens generally hollow.

Wood very light, soft, not strong, brittle, rather coarse-grained, compact, easily worked, very durable in contact with the soil; bands of small summer cells thin, dark colored, distinct; medullary rays numerous, obscure; color, dull brown tinged with red, the thin sap-wood nearly white; specific gravity, 0.3796; ash, 0.17; largely used for interior finish, fencing, shingles, in cabinet-making and cooperage, and exclusively by the Indians of the northwest coast in the manufacture of their canoes.

329.—*Chamæcyparis sphæroidea*, Spach,

Hist. Veg. xi, 331.—Endlicher, Syn. Conif. 61.—Lindley & Gordon in Jour. Hort. Soc. London, v, 209.—Knight, Syn. Conif. 20.—Carrière, Trait. Conif. 133; 2 ed. 122.—Gordon, Pinetum, 49; 2 ed. 71.—Henkel & Hochstetter, Nadelhölz. 248.—Nelson, Pinaceæ, 69.—Parlatore in De Candolle, Prodr. xvi², 464.—Ridgway in Proc. U. S. Nat. Mus. 1882, 87.

Cupressus thyoides, Linnaeus, Spec. 1 ed. 1003.—Kalm, Travels, English ed. ii, 174.—Du Roi, Harbk. ii, 198.—Marshall, Arbustum, 39.—Wangenheim, Amer. 8, t. 2, f. 4.—Aiton, Hort. Kew. iii, 372; 2 ed. v, 323.—Bartram, Travels, 2 ed. 409.—Michaux, Fl. Bor.-Am. ii, 208.—Willdenow, Spec. iv, 512; Enum. 991; Berl. Baumz. 111.—Nouveau Duhamel, iii, 6.—Persoon, Syn. ii, 590.—Desfontaines, Ilust. Arb. ii, 567.—Schkuhr, Handb. iii, 286, t. 310.—Michaux f. Hist. Arb. Am. iii, 20, t. 2; N. American Sylva, 3 ed. iii, 162, t. 152.—Pursh, Fl. Am. Sept. ii, 646.—Eaton, Manual, 111; 6 ed. 115.—Nuttall, Genera, ii, 224.—Hayne, Dend. Fl. 178.—Elliott, Sk. ii, 644.—Watson, Dend. Brit. ii, 156.—Torrey, Compend. Fl. N. States, 361; Fl. N. York, ii, 233.—Beck, Bot. 338.—London, Arboretum, iv, 2475, f. 2327.—Forbes, Pinetum Woburn. 183, t. 61.—Hooker, Fl. Bor.-Am. ii, 165.—Eaton & Wright, Bot. 215.—Bigelow, Fl. Boston. 3 ed. 387.—Emerson, Trees Massachusetts, 98; 2 ed. i, 114.—Richardson, Arctic Exped. 442.—Darby, Bot. S. States, 516.—Cooper in Smithsonian Rep. 1858, 257.—Chapman, Fl. S. States, 435.—Curtis in Rep. Geological Surv. N. Carolina, 1660, iii, 23.—Wood, Cl. Book, 663; Bot. & Fl. 315.—Porcher, Resources S. Forests, 509.—Gray, Manual N. States, 5 ed. 473.—Hoopes, Evergreens, 346.—Koch, Dendrologie, ii², 162.—Vasey, Cat. Forest Trees, 36.—Veitch, Manual Conif. 232.

Thuya sphæroidea, Sprengel, Syst. iii, 889.

Thuya sphæroidalis, Richard, Conif. 45, t. 8, f. 2.

WHITE CEDAR.

Southern Maine, south near the coast to northern Florida, and along the Gulf coast to the valley of the Pearl river, Mississippi.

A tree 24 to 27 meters in height, with a trunk 0.60 to 1.20 meter in diameter; in deep, cold swamps; rare in the Gulf states, west of the bay of Mobile.

Wood very light and soft, not strong, close-grained, compact, easily worked, very durable in contact with the soil; bands of small summer cells thin, dark colored, conspicuous; medullary rays numerous, obscure; color, light brown tinged with red, growing darker with exposure, the sap-wood lighter; specific gravity, 0.3322; ash, 0.33; largely used in boat-building; for woodenware, cooperage, shingles, interior finish, telegraph and fence posts, railway ties, etc.

Along the Atlantic coast from New Jersey southward lumber is manufactured from buried trunks of this species dug from peat swamps.

330.—*Chamæcyparis Nutkaensis*, Spach,

Hist. Veg. xi, 333.—Nuttall, Sylva, iii, 105; 2 ed. ii, 165.—Endlicher, Syn. Conif. 62.—Ledebour, Fl. Rossica, iii, 680.—Lindley & Gordon in Jour. Hort. Soc. London, v, 209.—Carrière, Trait. Conif. 134; 2 ed. 127.—Walpers, Ann. v, 796.—Henkel & Hochstetter, Nadelhölz, 250.—Parlatore in De Candolle, Prodr. xvi², 465.—Hall in Coulter's Bot. Gazette, ii, 91.—G. M. Dawson in Canadian Nat. 2 ser. ix, 329.

Cupressus Nootkatensis, Lambert, Pinus, 1 ed. ii, 18; 2 ed. ii, No. 60.—London, Arboretum, iv, 2480.

Cupressus Nutkaensis, Hooker, Fl. Bor.-Am. ii, 165.—Newberry in Pacific R. R. Rep. vi, 63, f. 28.—Gordon, Pinetum, 66; 2 ed. 94.—Cooper in Smithsonian Rep. 1858, 263.—Nelson, Pinaceæ, 74.—Hoopes, Evergreens, 345.—Lawson, Pinetum Brit. ii, 199, t. 34, f. 1-12.—Koch, Dendrologie, ii², 165.—Vasey, Cat. Forest Trees, 36.—Macoun in Geological Rep. Canada, 1876-'77, 211.—Veitch, Manual Conif. 235.

Thuja excelsa, Bongard in Mem. Acad. St. Petersburg, 6 ser. ii, 164.

Cupressus Americana, Trautvetter, Imag. Pl. Fl. Rossica, 12, t. 7.

C. Nutkaensis, var. *glauca*, Walpers, Ann. v, 769.

Thuyopsis borealis, Hort.—Carrière, Trait. Conif. 1 ed. 113.

Thuyopsis cupressoides, Carrière, Man. des Pl. iv, 324.

C. excelsa, Fischer in herb. Sitka.

Thuyopsis Tchugatskoy and *T. Tchugatskoyæ*, Hort.

YELLOW CYPRESS. SITKA CYPRESS.

Sitka, south along the islands and Coast ranges of British Columbia and the Cascade mountains of Washington territory and Oregon to the valley of the Santian river, Oregon ("Lucky Camp mountain", *Cusick*).

A large tree of great economic value, 30 to 38 meters in height, with a trunk 1.20 to 1.80 meter in diameter, or toward its southern limits and at high elevations much smaller; common along the coast at the sea-level to about latitude 49° 30' N., then less common and only at higher elevations; south of British Columbia hardly below 5,000 feet elevation and very rare and local; the most valuable timber tree of Alaska.

Wood light, hard, not strong, brittle, very close-grained, compact, very durable in contact with the soil, easily worked, satiny, susceptible of a beautiful polish, possessing an agreeable, resinous odor; bands of small summer cells thin, not conspicuous; medullary rays thin, numerous, hardly distinguishable; color, bright, light clear yellow, the thin sap-wood nearly white; specific gravity, 0.4782; ash, 0.34; somewhat used in boat and ship-building, for furniture, interior finish, etc., probably unsurpassed in beauty as a cabinet wood by that of any North American tree.

331.—*Chamæcyparis Lawsoniana*, Parlatore,

Stud. Organ. Conif. 23, 29, t. 3, f. 22-25; De Candolle, Prodr. xvi², 464.—Gordon, Pinetum, 2 ed. 85.—Watson, Bot. California, ii, 155.—Sargent in London Gard. Chronicle, 1881, 8.

Cupressus Lawsoniana, Murray in Edinburgh New Phil. Jour. new ser. i, 292, t. 9.—Bot. Mag. t. 5551.—Nelson, Pinaceæ, 72.—Cooper in Smithsonian Rep. 1858, 263.—Lawson, Pinetum Brit. ii, 191, t. 31, f. 1-13.—Hoopes, Evergreens 342, f. 53.—Henkel & Hochstetter, Nadelhölz. 246.—Fowler in London Gard. Chronicle, 1872, 285.—London Garden, vii 508 & t.—Vasey, Cat. Forest Trees, 36.—Veitch, Manual Conif. 231.—Eichler in Monatsb. Acad. Berl. 1881, f. 29, 30.

Cupressus fragrans, Kellogg in Proc. California Acad. i, 103.

?*Cupressus attenuata*, Gordon, Pinetum, 1 ed. 57; 2 ed. 79.

C. Boursierii, Carrière, Trait. Conif. 2 ed. 125 [not Decaisne].

C. Nutkanus, Torrey, Bot. Wilkes Exped. t. 16.

PORT ORFORD CEDAR. OREGON CEDAR. WHITE CEDAR. LAWSON'S CYPRESS. GINGER PINE.

Oregon, Coos bay, south to the valley of the Rogue river, not extending more than thirty miles from the coast; California, valley of the upper Sacramento river (shores of Castle and Soda lakes, Shasta county).

A large tree of the first economic value, 45 to 61 meters in height, with a trunk 1.80 to 4 meters in diameter; rich woods, in low, moist soil, interspersed with the yellow fir and hemlock; most common and reaching its greatest development along the Oregon coast; local; in California very rare and local.

Wood light, hard, strong, very close-grained, compact, easily worked, very durable in contact with the ground, abounding in odoriferous resin, satiny, susceptible of a beautiful polish; layers of small summer cells thin, not conspicuous; medullary rays numerous, very obscure; color, light yellow or almost white, the thin sap-wood hardly distinguishable; specific gravity, 0.4621; ash, 0.10; largely manufactured into lumber and used for interior finish, flooring, railway ties, fence posts, matches, and in ship- and boat-building; the resin strongly diuretic and a powerful insecticide.

332.—*Cupressus macrocarpa*, Hartweg,

Jour. Hort. Soc. London, ii, 187.—Bentham, Pl. Hartweg. 337.—Gordon in Jour. Hort. Soc. London, iv, 296 & t.; Pinetum, 65; 2 ed. 91.—Lindley & Gordon in Jour. Hort. Soc. London, v, 206.—Knight, Syn. Conif. 20.—Torrey, Bot. Mex. Boundary Survey, 211.—Cooper in Smithsonian Rep. 1853, 263; Proc. California Acad. iii, 290.—Carrière, Trait. Conif. 1 ed. 124, in part.—Bolander in Proc. California Acad. iii, 228.—Henkel & Hochstetter, Nadelhölz. 239.—Nelson, Pinaceæ, 73.—Hoopes, Evergreens, 353.—Parlatore in De Candolle, Prodr. xvi³, 473.—Fowler in London Gard. Chronicle, 1872, 285.—Koch, Dendrologie, ii², 148.—Vasey, Cat. Forest Trees, 36.—Watson, Bot. California, ii, 113.—Veitch, Manual Conif. 234.—Lawson Pinetum Brit. ii, 195, t. 32.

C. Lambertiana, Carrière in Rev. Hort. 1855, 232; Trait. Conif. 124; 2 ed. 166.

C. Hartwegii, Carrière in Rev. Hort. 1855, 232; Trait. Conif. 2 ed. 168.

? *C. macrocarpa*, var. *fastigiata*, Knight, Conif. 20.—Parlatore in De Candolle, Prodr. xvi³, 473.—Veitch, Manual Conif. 234.

? *C. Hartwegii*, var. *fastigiata*, Carrière, Trait. Conif. 2 ed. 169.

MONTEREY CYPRESS.

California, Monterey (Cypress point, Pescadero ranch, and Carmelo point).

A tree 15 to 21 meters in height, with a trunk 1.20 to 1.80 meter in diameter; on granite rocks immediately upon the sea-coast; very local.

Wood heavy, hard, strong, rather brittle, very close-grained, compact, easily worked, very durable in contact with the soil, satiny, susceptible of a beautiful polish, odorous; bands of small summer cells thin, dark colored, conspicuous; medullary rays numerous, hardly distinguishable; color, clear bright brown streaked with red and yellow, the thin sap-wood light yellow; specific gravity, 0.6261; ash, 0.57; very beautiful and of undoubted value as a cabinet wood.

333.—*Cupressus Goveniana*, Gordon,

Jour. Hort. Soc. London, iv, 296 & f.; Pinetum, 60; 2 ed. 83.—Bentham, Pl. Hartweg. 337.—Lindley & Gordon in Jour. Hort. Soc. London, v, 206.—Carrière, Trait. Conif. 125; 2 ed. 170.—Torrey, Mex. Boundary Survey, 211.—Cooper in Smithsonian Rep. 1853, 263.—Henkel & Hochstetter, Nadelhölz. 240.—Hoopes, Evergreens, 252.—Parlatore in De Candolle, Prodr. xvi³, 472.—Fowler in London Gard. Chronicle, 1872, 285.—Watson, Bot. California, ii, 114.—Veitch, Manual Conif. 230.

? *C. Californica*, Carrière, Trait. Conif. 127; 2 ed. 164.

C. Californica gracilis, Nelson, Pinaceæ, 70, in part

? *C. cornuta*, Carrière in Rev. Hort. 1866, 251 & f.

? *Juniperus aromatica*, Hort.

Humboldt county, California, south along the coast and through the Coast ranges into Lower California.

A small tree, sometimes 12 to 15 meters in height, with a trunk 0.60 to 0.90 meter in diameter; borders of streams and mountain slopes, in rather rich soil, or often a low shrub, fruiting when 0.30 to 1 meter in height, and occupying extensive tracts of sandy barrens 1 to 5 miles inland from the coast, or thin, rocky soil (*Pringle*); widely but not generally distributed.

Wood light, soft, not strong, brittle, close-grained, compact; bands of small summer cells broad, dark colored, conspicuous; medullary rays thin, obscure; color, light brown, the thick sap-wood nearly white; specific gravity, 0.4689; ash, 0.45.

334.—*Cupressus Macnabiana*, Murray,

Edinburgh, New Phil. Jour. new ser. i, 293, t. 10.—Gordon, Pinetum, 64; 2 ed. 90.—Carrière, Trait. Conif. 2 ed. 165.—Hoopes, Evergreens, 353.—Parlatore in De Candolle, Prodr. xvi², 473.—Koch, Dendrologie, ii², 150.—Vasey, Cat. Forest Trees, 36.—Watson, Bot. California, ii, 114.—Veitch, Manual Conif. 233.

C. glandulosa, Hooker, (ex. Henkel & Hochstetter, Nadelhölz. 241).

C. Californici gracilis, Nelson, Pinaceæ, 70, in part.

California, mountains south of Clear lake, Lake county (*Torrey, Bolander, Pringle, Miller*).

A small tree, sometimes 9 meters in height, with a trunk 0.30 to 0.45 meter in diameter, or more often a tall shrub branching from the ground; very rare and local; not rediscovered in the original station reported by Jeffrey, the Mount Shasta region.

Wood not collected.

335.—*Cupressus Guadalupensis*, Watson,

Proc. Am. Acad. xiv, 300; Bot. California, ii, 114.

C. macrocarpa, ? Watson in Proc. Am. Acad. xi, 119 [not Hartweg].

C. Arizonica, E. L. Greene in Bull. Torrey Bot. Club, ix, 64.—Rusby in Bull. Torrey Bot. Club, ix, 79.—Watson in Proc. Am. Acad. xviii, 157.

San Francisco mountains of New Mexico and eastern Arizona (*Greene, Rusby*), Santa Catalina and Santa Rita mountains, Arizona (*Pringle, Lemmon*); on the Sierra Madre, near Saltillo, and Gaudalupe island, Mexico (*Palmer*).

A tree 18 to 21 meters in height, with a trunk 0.60 to 0.90 meter in diameter; rocky cañons and ridges; on the New Mexico and Arizona mountains, forming extensive forests between 5,000 and 8,000 feet elevation, generally on northern slopes; local.

Wood light, soft, very close-grained, compact, easily worked, susceptible of a good polish; bands of small summer cells, broad, conspicuous; medullary rays numerous, very obscure; color, gray, often faintly streaked with yellow, the thick sap-wood light yellow; specific gravity, 0.4843; ash, 0.44.

336.—*Juniperus Californica*, Carrière,

Rev. Hort. iii, 353 & f.; Trait. Conif. 58; 2 ed. 41.—Gordon, Pinetum, 121.—Vasey, Cat. Forest Trees, 37.—Engelmann in Trans. St. Louis Acad. iii, 588; Wheeler's Rep. vi, 375.—Palmer in Am. Nat. xii, 593.—Watson, Bot. California, ii, 113.

J. tetragona, var. *osteosperma*, Torrey in Pacific R. R. Rep. iv, 141; Bot. Mex. Boundary Survey, 210; Ives' Rep. 28.

J. tetragona, Cooper in Smithsonian Rep. 1858, 263 [not Schlechtendal].

J. Uerrosianus, Kellogg in Proc. California Acad. ii, 37.

J. occidentalis, Gordon, Pinetum, Suppl. 38; Pinetum, 2 ed. 162, in part.—Henkel & Hochstetter, Nadelhölz. 245, in part.—Hoopes, Evergreens, 299, in part.—Parlatore in De Candolle, Prodr. xvi², 469, in part.

J. Californica, var. *osteosperma*, Engelmann; Watson in Proc. Am. Acad. xi, 119.

JUNIPER.

California, San Francisco bay, south through the Coast ranges to Lower California.

A small tree, rarely 6 to 9 meters in height, with a trunk 0.30 to 0.60 meter in diameter, or more often a tall shrub, sending up many stems from the ground; sandy barrens and dry, rocky soil.

Wood light, soft, very close-grained, compact, very durable in contact with the soil; bands of small summer cells thin, dark colored, not conspicuous; medullary rays numerous, very obscure; color, light brown slightly tinged with red, the sap-wood nearly white; specific gravity, 0.6282; ash, 0.75; in southern California largely used for fencing and fuel.

Var. *Utahensis*, Engelmann,

Trans. St. Louis Acad. iii, 588; Wheeler's Rep. vi, 264.—Vasey, Cat. Forest Trees, 37.—Sargent in Am. Jour. Sci. 3 ser. xvii, 418.—Palmer in Am. Nat. xii, 594.—Watson, Bot. California, ii, 113.

J. occidentalis, Watson in King's Rep. v, 336, in part; Pl. Wheeler, 18 [not Hooker].

J. occidentalis, var. *Utahensis*, Veitch, Manual Conif. 289.

JUNIPER.

Western base of the Wahsatch mountains, Utah, to eastern California, south through the Great Basin to southeastern California (*Pringle*) and the San Francisco mountains, eastern Arizona (*Greene*).

A small, contorted tree, 6 to 9 meters in height, with a trunk 0.60 to 0.90 meter in diameter, or more often a tall, much-branched shrub; very common through the elevated valleys and along the lower slopes of all the ranges of central and southern Utah and Nevada, and the most generally-distributed arborescent species of the region.

Wood light, soft, close-grained, compact, very durable in contact with the soil; color, light brown, the thick sap-wood nearly white; specific gravity, 0.5522; ash, 0.49; the common fuel and fencing material of the region.

337.—*Juniperus pachyphloea*, Torrey,

Pacific R. R. Rep. iv, 142; Bot. Mex. Boundary Survey, 210; Ives' Rep. 28.—Cooper in Smithsonian Rep. 1858, 263.—Henkel & Hochstetter, Nadelhölz. 247.—Carrière, Trait. Conif. 2 ed. 56.—Parlatore in De Candolle, Prodr. xvi², 490.—Gordon, Pinetum, 2 ed. 164.—Engelmann in Trans. St. Louis Acad. iii, 589; Wheeler's Rep. vi, 264.—Palmer in Am. Nat. xii, 593.—Veitch, Manual Conif. 289.—Rusby in Bull. Torrey Bot. Club, ix, 79.—Hemsley, Bot. Am.-Cent. iii, 184.

J. plochyderma, Torrey in Sitgreaves' Rep. 173, t. 16.

J. Sabina pachyphloea, Antoine, Kupress. 39.

JUNIPER.

Eagle and Limpia mountains (*Havard*), west along the ranges of western Texas, southern New Mexico and Arizona south of latitude 34°; southward into Mexico.

A tree 9 to 15 meters in height, with a trunk 0.60 to 1.20 meter in diameter; dry, stony slopes and ridges, generally between 2,000 and 3,000 feet elevation; the prevailing and largest juniper of the mountains of western Texas.

Wood light, soft, not strong, brittle, very close-grained, compact, susceptible of a fine polish; bands of small summer cells very thin, dark colored, not conspicuous; medullary rays numerous, obscure; color, clear light red, often streaked with yellow, the thin sap-wood nearly white; specific gravity, 0.5829; ash, 0.11.

338.—*Juniperus occidentalis*, Hooker,

Fl. Bor.-Am. ii, 166.—Endlicher, Syn. Conif. 26.—Lindley & Gordon in Jour. Hort. Soc. London, v, 202.—Carrière, Conif. 42, in part; 2 ed. 40, in part.—Torrey in Pacific R. R. Rep. iv, 142.—Cooper in Smithsonian Rep. 1858, 263.—Gordon, Pinetum, 117 (excl. syn.); Suppl. 38 (excl. syn.); 2 ed. 162 (excl. syn.).—Henkel & Hochstetter, Nadelhölz. 345, in part.—Nelson, Pinaceæ, 142.—Hoopes, Evergreens, 299 (excl. syn. *Californica*).—Parlatore in De Candolle, Prodr. xvi², 489, in part.—Vasey, Cat. Forest Trees, 37.—Macoun in Geological Rep. Canada, 1875-76, 211.—Palmer in Am. Nat. xii, 594.—Watson, Bot. California, ii, 113.—Veitch, Manual Conif. 289.

J. excelsa, Pursh, Fl. Am. Sept. ii, 647.—Nuttall, Genera, ii, 245.

J. Andina, Nuttall, Sylva, iii, 95, t. 110; 2 ed. ii, 157, t. 110.—Carrière, Trait. Conif. 2 ed. 55.

Chamæcyparis Boursierii, Decaisne in Bull. Soc. Bot. France, i, 70.

J. Hermannii, Koch, Dendrologie, ii², 141 [not Sprengel].

J. occidentalis, var. *pleiosperma*, Engelmann in Trans. St. Louis Acad. ii, 590.

J. pyramidalis, Hort.

JUNIPER.

Blue mountains and high prairies of eastern Washington territory and Oregon, Cascade mountains of Oregon, valley of the Klamath river, California, and south along the high ridges of the Sierra Nevada, between 7,000 and 10,000 feet elevation, to the San Bernardino mountains (*Parish Bros.*).

A tree 9 to 15 meters in height, with a trunk 1.20 to 2.10 meters in diameter, or often a low, much-branched shrub; dry, rocky ridges and prairies, reaching its greatest development in the California sierras.

Wood light, soft, very close-grained, compact, very durable in contact with the soil; bands of small summer cells thin, not conspicuous; medullary rays numerous, very obscure; color, light red or brown, the sap-wood nearly white; specific gravity, 0.5765; ash, 0.12; largely used for fencing and fuel.

Var. *monosperma*, Engelmann,

Trans. St. Louis Acad. iii, 590; Wheeler's Rep. vi, 263.—Veitch, Manual Conif. 289.—Rusby in Bull. Torrey Bot. Club, ix, 79.

JUNIPER.

Eastern base of Pike's peak, Colorado, to the mountains of western Texas, and through New Mexico and southern Arizona to southern California.

A small, stunted tree, 6 to 9 meters in height, with a trunk sometimes 0.60 meter in diameter, or often branching from the ground with many stout, contorted stems; dry, gravelly slopes between 3,500 and 7,000 feet elevation.

Wood heavier than that of the type, the layers of annual growth often eccentric; specific gravity, 0.7119; ash, 0.78; largely used for fuel and fencing.

Var. *conjugens*, Engelmann,

Trans. St. Louis Acad. iii, 590.—Veitch, Manual Conif. 289.—Watson in Proc. Am. Acad. xviii, 158.

JUNIPER.

Western Texas, valley of the Colorado river (Austin), west and north.

A tree 11 to 15 meters in height, with a trunk sometimes 0.30 meter in diameter, covering with extensive forests the limestone hills of western Texas; its range not yet satisfactorily determined.

Wood light, hard, not strong, very close-grained, compact, very durable in contact with the soil; bands of small summer cells thin, dark colored, conspicuous; medullary rays numerous, very obscure; color, brown, often streaked with red, the thin sap-wood nearly white; specific gravity, 0.6907; ash, 0.46; largely used for fencing, fuel, telegraph poles, railway ties, etc.

339.—*Juniperus Virginiana*, Linnæus,

Spec. 1 ed. 1039.—Kalm, Travels, English ed. ii, 180.—Marshall, Arbustum, 70.—Wangenheim, Amer. 9, t. 2, f. 5.—Walter, Fl. Caroliniana, 248.—Aiton, Hort. Kew. iii, 414; 2 ed. v, 414.—Lamarck, Dict. iv, 627.—Willdenow, Spec. iv, 853; Enum. 1025; Berl. Baumz. 198.—Persoon, Syn. ii, 632.—Desfontaines, Hist. Arb. ii, 539.—Michaux f. Hist. Arb. Am. iii, 42, t. 5; N. American Sylva, 3 ed. 173, t. 155.—Pursh, Fl. Am. Sept. 647.—Nouveau Duhamel, vi, 49, t. 16.—Barton, Prodr. Fl. Philadelph. 96; Compend. Fl. Philadelph. ii, 200.—Eaton, Manual, 118; 2 ed. 194.—Nuttall, Genera, ii, 245; Sylva, iii, 97; 2 ed. ii, 159.—Bigelow, Med. Bot. iii, 49, t. 45; Fl. Boston. 3 ed. 398.—Hayne, Dend. Fl. 205.—Elliott, Sk. ii, 717.—Torrey in Nicolle's Rep. 167; Compend. Fl. N. States, 377; Fl. N. York, ii, 235; Marcy's Rep. 284; Pacific R. R. Rep. iv, 142; Bot. Mex. Boundary Survey, 211; Ives' Rep. 28.—Sprengel, Syst. iii, 908.—Richard, Conif. 37, t. 6, f. 2.—Audubon, Birds, t. 43.—Rafinesque, Med. Bot. ii, 13.—Beck, Bot. 337.—Lindley, Fl. Med. 556.—London, Arboretum, iv, 2495, f. 2357.—Forbes, Pinetum Woburn. 199.—Penn. Cycl. xiii, 147.—Eaton & Wright, Bot. 288.—Emerson, Trees Massachusetts, 102; 2 ed. i, 118.—Endlicher, Syn. Conif. 27, in part.—Scheele in Rømer, Texas, Appx. 447.—Lindley & Gordon in Jour. Hort. Soc. London, v, 202.—Parry in Owen's Rep. 618.—Darlington, Fl. Cestrica, 3 ed. 295.—Knight, Syn. Conif. 12.—Darby, Bot. S. States, 515.—Durand in Jour. Philadelphia Acad. 1355, 101.—Torrey & Gray in Pacific R. R. Rep. ii, 130, 175.—Carrière, Trait. Conif. 43; 2 ed. 44.—Bigelow in Pacific R. R. Rep. 20.—Gordon, Pinetum, 112; 2 ed. 154.—Cooper in Smithsonian Rep. 1858, 257; Am. Nat. iii, 413.—Chapman, Fl. S. States, 435.—Gray in Pacific R. R. Rep. xii, 48; Manual N. States, 5 ed. 474; Hall's Pl. Texas, 21.—Hooker f. in Trans. Linnæan Soc. xxiii, 302.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 71.—Lesquereux in Owen's 2d Rep. Arkansas, 389.—Wood, Cl. Book, 663; Bot. & Fl. 314.—Porcher, Resources S. Forests, 510.—Engelmann in Trans. Am. Phil. Soc. new ser. xii, 209; Trans. St. Louis Acad. iii, 591; Wheeler's Rep. vi, 263.—Lyll in Jour. Linnæan Soc. vii, 144.—Henkel & Hochstetter, Nadelhölz. 335.—Nelson, Pinaceæ, 153.—Hoopes, Evergreens, 291.—Parlatore in De Candolle, Prodr. xvi, 488.—Young, Bot. Texas, 517.—Koch, Dendrologie, ii, 138.—Watson in King's Rep. v, 335.—Rothrock in Pl. Wheeler, 28, 50; Wheeler's Rep. vi, 10.—Porter & Coulter, Fl. Colorado; Hayden's Surv. Misc. Pub. No. 4, 132.—Hayden in Warren's Rep. Nebraska & Dakota, 2 ed. 122.—Vasey, Cat. Forest Trees, 37.—Guibourt, Hist. Drogues, 7 ed. ii, 242.—Broadhead in Coulter's Bot. Gazette, iii, 60.—G. M. Dawson in Canadian Nat. new ser. ix, 329.—Sears in Bull. Essex Inst. xiii, 183.—Veitch, Manual Conif. 282.—Bell in Geological Rep. Canada, 1879-'80, 52.—Ridgway in Proc. U. S. Nat. Mus. 1882, 87.—Hemsley, Bot. Am.-Cent. iii, 184.

J. Caroliniana, Marshall, Arbustum, 71.—Du Roi, Harbk. 2 ed. 497.

J. arborescens, Mönch, Meth. 699.

J. Barbadosensis, Michaux, Fl. Bor.-Am. ii, 246 [not Linnæus].—Pursh, Fl. Am. Sept. ii, 647.—Nuttall, Genera, ii, 245; Sylva, iii, 96; 2 ed. ii, 158.

J. Virginiana, var. *Caroliniana*, Willdenow, Berl. Baumz. 198.—Hayne, Dend. Fl. 205.—London, Arboretum, iv, 2495.

J. Virginiana, var. *Hermannii*, Persoon, Syn. ii, 632.

J. Hermannii, Sprengel, Syst. iii, 908.

J. foetida, var. *Virginiana*, Spach in Ann. Sci. Nat. 2 ser. xvi, 298; Hist. Veg. xi, 316.

J. Virginiana vulgaris, Endlicher, Syn. Conif.

J. Sabina, var. *Virginiana*, Antoine, Kupress. t. 83, 84.

RED CEDAR. SAVIN.

Southern New Brunswick to the northern shores of Georgian bay, northern Michigan, Wisconsin and Minnesota, south to cape Malabar and Tampa bay, Florida, and the valley of the Colorado river, Texas, west to eastern Nebraska, Kansas, and the Indian territory to about the one hundredth parallel of west longitude; in the Pacific region, Rocky mountains of Colorado to Vancouver's island, British Columbia; not extending to western Texas, California, or Oregon; in Utah, Nevada, and Arizona rare and local.

The most widely distributed of North American Coniferae, a tree 24 to 30 meters in height, with a trunk 0.60 to 1.35-meter in diameter, or toward its northern and western limits much smaller, often reduced to a low shrub; dry, gravelly ridges, and limestone hills, or in the Gulf states, especially near the coast, in deep swamps; in northern Montana, borders of streams and lakes; common; and reaching its greatest development in the valley of the Red river, Texas.

Wood light, soft, not strong, brittle, very close- and straight-grained, compact, easily worked, very durable in contact with the soil; odorous; bands of small summer cells rather broad, conspicuous; medullary rays numerous, very obscure; color, dull red, the thin sap-wood nearly white; specific gravity, 0.4926; ash, 0.13; largely used for posts, sills, railway ties, interior finish, cabinet-making, and almost exclusively for lead-pencils.

A decoction of the leaves is occasionally used as a substitute for savine cerete, and an infusion of the berries as a diuretic (*U. S. Dispensatory*, 14 ed. 529.—*Nat. Dispensatory*, 2 ed. 795).

340.—*Taxodium distichum*, Richard,

Ann. Mus. xvi, 298; Conif. 52, t. 10.—Nouveau Duhamel, iii, 8.—Robin, Voyages, iii, 525.—Lambert, Pinus, 2 ed. 25 & t.—Torrey, Compend. Fl. N. States, 361; Bot. Mex. Boundary Survey, 210.—Brongniart in Ann. Sci. Nat. 1 ser. xxx, 182.—Loudon, Arboretum, iv, 2481, f. 2335-2339.—Forbes, Pinetum Woburn. 177, t. 60.—Endlicher, Syn. Conif. 68, in part.—Engelmann & Gray in Jour. Boston Soc. Nat. Hist. v, 234.—Scheele in Rømer, Texas, Appx. 447.—Lindley & Gordon in Jour. Hort. Soc. London, v, 269.—Knight, Syn. Conif. 20.—Darlington, Fl. Cestrica, 3 ed. 295.—Carrière, Trait. Conif. 143; 2 ed. 180; Rev. Hort. viii, 62 & f.—Morren in Belg. Hort. vi, 74 & t.—Gordon, Pinetum, 305; 2 ed. 382.—London Gard. Chronicle, 1857, 549.—Cooper in Smithsonian Rep. 1858, 257.—Chapman, Fl. S. States, 435.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 29.—Lesquereux in Owen's 2d Rep. Arkansas, 389.—Wood, Cl. Book, 663; Bot. & Fl. 315.—Henkel & Hochstetter, Nadelhölz. 258.—Gray, Manual N. States, 5 ed. 473.—Hoopes, Evergreens, 364, f. 58.—Parlatore in De Candolle, Prodr. xvi², 440.—Lawson, Pinetum Brit. ii, 305, f. 1-9.—Fowler in London Gard. Chronicle, 1872, 1526.—Young, Bot. Texas, 518.—Koch, Dendrologie, ii², 195.—Bertrand in Bull. Soc. Bot. France, xviii, 127.—Vasey, Cat. Forest Trees, 36.—Broadhead in Coulter's Bot. Gazette, iii, 60.—Veitch, Manual Conif. 214.—Ridgway in Proc. U. S. Nat. Mus. 87.—Watson in Proc. Am. Acad. xviii, 158.

Cupressus disticha, Linnæus, Spec. 1 ed. 1003.—Du Roi, Harbk. i, 201.—Marshall, Arbustum, 39.—Lamarek, Dict. ii, 244.—Wangenheim, Amer. 43.—Walter, Fl. Caroliniana, 238.—Aiton, Hort. Kew. iii, 372; 2 ed. v, 323.—Bartram, Travels, 2 ed. 88.—Michaux, Fl. Bor.-Am. ii, 208.—Desfontaines, Hist. Arb. ii, 567.—Willdenow, Spec. iv, 512; Enum. 991; Berl. Baumz. 111.—Schkuhr, Handb. iii, 288.—Michaux f. Hist. Arb. Am. iii, 4, t. 1; N. American Sylva, 3 ed. iii, 154, t. 151.—Pursh, Fl. Am. Sept. ii, 645.—Barton, Prodr. Fl. Philadelph. 93.—Rafinesque, Fl. Ludoviciana, 151.—Nuttall, Genera, ii, 224.—Hayne, Dend. Fl. 178.—James in Long's Exped. ii, 317, 318.—Elliott, Sk. ii, 642.—Beck, Bot. 238.—Eaton, Manual, 6 ed. 116.—Eaton & Wright, Bot. 215.—De Chambray, Trait. Arb. Res. Conif. 349.—Dickson & Brown in Am. Jour. Sci. 2 ser. v, 15.—Porcher, Resources S. Forests, 508.

Cupressus disticha, var. *patens* and var. *nutans*, Aiton, Hort. Kew. 2 ed. v, 323.

Cupressus disticha, var. *imbricaria*, Nuttall, Genera, ii, 224; Trans. Am. Phil. Soc. 2 ser. v, 163.—Croom in Ann. Jour. Sci. 1 ser. xxviii, 166.

Schubertia disticha, Mirbel in Mem. Mus. xiii, 75.—Sprengel, Syst. iii, 890.—Spach, Hist. Veg. xi, 349.

T. microphyllum, Brongniart in Ann. Sci. Nat. 1 ser. xxx, 182.—Endlicher, Syn. Conif. 68.—Lindley & Gordon in Jour. Hort. Soc. London, v, 207.—Carrière, Trait. Conif. 148.

T. adscendens, Brongniart in Ann. Sci. Nat. 1 ser. xxx, 182.—Endlicher, Syn. Conif. 69.—Lindley & Gordon in Jour. Hort. Soc. London, v, 207.—Carrière, Trait. Conif. 148.

T. distichum, var. *patens* and var. *nutans*, Endlicher, Syn. Conif. 68.—Loudon, Arboretum, iv, 2481.

T. distichum fustigiatum, Knight, Syn. Conif. 21.—Carrière, Trait. Conif. 145; 2 ed. 181.—Gordon, Pinetum, 307; 2 ed. 383.—Henkel & Hochstetter, Nadelhölz. 260.—Hoopes, Evergreens, 367.

T. distichum, var. *microphyllum*, Henkel & Hochstetter, Nadelhölz. 261.—Parlatore in De Candolle, Prodr. xvi², 441 (*T. Sinense pendulum*, Forbes, Pinetum Woburn. 180.—*Glyptostrobus pendulus*, Endlicher, Conif. 71.—Bot. Mag. t. 5603.—Carrière, trait. Conif. 152.—*T. Sinense*, Gordon, Pinetum, 309.—*Cupressus Sinense*, Hort.).

Cupressinnata disticha, Nelson, Pinaceæ, 61.

BALD CYPRESS. BLACK CYPRESS. RED CYPRESS. WHITE CYPRESS. DECIDUOUS CYPRESS.

Sussex county, Delaware, south near the coast to Mosquito inlet and cape Romano, Florida, west through the Gulf states near the coast to the valley of the Nueces river, Texas, and through Arkansas to western Tennessee, western and northern Kentucky, southeastern Missouri, and southern Illinois and Indiana.

A large tree of great economic value, 24 to 46 meters in height, with a trunk 1.80 to 4 meters in diameter; deep, submerged swamps, river-bottom lands, and pine-barren ponds; common and forming extensive forests in the south Atlantic and Gulf states.

Wood light, soft, close, straight-grained, not strong, compact, easily worked, very durable in contact with the soil; bands of small summer cells broad, resinous, conspicuous; medullary rays numerous, very obscure; color, light or dark brown, the sap-wood nearly white; specific gravity, 0.4543; ash, 0.42; largely manufactured into lumber and used for construction, cooperage, railway ties, posts, fencing, etc., often injured, especially west of the Mississippi river, by a species of *Dædalia*, not yet determined, rendering it unfit for lumber.

Two varieties of cypress, black and white, are recognized by lumbermen, the wood of the former heavier than water when green, rather harder and considered more durable than the other; the unseasoned wood of the latter lighter than water and rather lighter colored than black cypress.

341.—*Sequoia gigantea*, Decaisne,

Bull. Bot. Soc. France, i, 70; Rev. Hort. 1855, 9, t. 10, f. 1.—Gray in Proc. Am. Acad. iii, 94; Am. Jour. Sci. 2 ser. xvii. 440; xviii, 150, 286.—Torrey in Pacific R. R. Rep. iv, 140.—Kellogg in Proc. California Acad. i, 42.—Blake in Pacific R. R. Rep. v, 257, t. 13.—Carrière, Trait. Conif. 166.—Newberry in Pacific R. R. Rep. vi, 90.—Cooper in Smithsonian Rep. 1858, 263.—Wood, Bot. & Fl. 315.—Bloomer in Proc. California Acad. iii, 397.—Hoopes, Evergreens, 239, f. 29.—Parlatore in De Candolle Prodr. xvi², 437.—Koch, Dendrologie, ii², 194.—Bertrand in Ann. Sci. Nat. 5 ser. xx, 114.—Vasey, Cat. Forest Trees, 36.—Muir in Proc. Am. Assoc. xxv, 242.—Watson, Bot. California, ii, 117.

Wellingtonia gigantea, Lindley in London Gard. Chronicle, 1853, 819, 823; Bot. Mag. t. 4777, 4778.—C. Lemaire in Ill. Hort. 1854, 14 & t.—Naudin in Rev. Hort. 1854, 116.—Fl. des Serres, ix, 93 & t. 903 & t.—Flor. Cabinet, 1854, 121 & t.—Bigelow in Pacific R. R. Rep. iv, 22.—Gordon, Pinetum, 330; Suppl. 106; 2 ed. 415.—Murray in Edinburgh New Phil. Jour. new ser. xi, 205, t. 3-9 (Trans. Bot. Soc. Edinburgh, vi, 330, t. 6, f. 8, 9).—Henkel & Hochstetter, Nadelhölz. 222.—Carrière, Trait. Conif. 2 ed. 217.—Veitch, Manual Conif. 415.

Wellingtonia Californica, Winslow in California Farmer, September, 1854.—Hooker, Jour. Bot. & Kew Misc. vii, 26.

Taxodium Washingtonianum, Winslow in California Farmer, September, 1854.

Taxodium giganteum, Kellogg & Behr in Proc. California Acad. i, 51.

S. Wellingtonia, Seemann in Bonplandia, ii, 238; iii, 27; vi, 343; Ann. & Mag. Nat. Hist. 3 ser. March, 1859, 161.—Lawson, Pinetum Brit. iii, 299, t. 37, 51, 53, f. 1-37.

Gigantabies Wellingtonia, Nelson, Pinaceæ, 79.

BIG TREE.

California, western slopes of the Sierra Nevadas from Placer county (Calaveras Grove) south to Deer creek on the southern borders of Tulare county.

The largest tree of the American forest, 76 to 119 meters in height, with a trunk 6 to 11 meters in diameter; valleys and moist swales or hollows between 4,000 and 6,000 feet elevation, growing in small, isolated groves, except toward its southern limits, here mixed with the sugar pine and red and white firs, covering large tracts, often several hundred acres in extent.

Wood very light, soft, weak, brittle, rather coarse-grained, compact, remarkably durable in contact with the soil; bands of small summer cells thin, dark colored, conspicuous; medullary rays numerous, thin; color, bright clear red, turning much darker with exposure, the thin sap-wood white; specific gravity, 0.2882; ash, 0.50; in Fresno county formerly somewhat manufactured into lumber and locally used for fencing, shingles, construction, etc.

342.—*Sequoia sempervirens*, Endlicher,

Syn. Conif. 198.—Decaisne in Rev. Hort. 1855, 9, t. 11, f. 2.—Carrière, Trait. Conif. 164; 2 ed. 210.—Bigelow in Pacific R. R. Rep. iv, 23.—Newberry in Pacific R. R. Rep. vi, 57, 90, f. 23.—Torrey in Pacific R. R. Rep. iv, 140; Bot. Mex. Boundary Survey, 210; Ives' Rep. 28.—Gordon, Pinetum, 303; Suppl. 97; 2 ed. 379.—Cooper in Smithsonian Rep. 1858, 263.—Murray in Edinburgh New Phil. Jour. new ser. xi, 221 (Trans. Bot. Soc. Edinburgh, vi, 346).—Seemann in Ann. & Mag. Nat. Hist. 3 ser. March, 1859, 165.—Wood, Bot. & Fl. 315.—Bolander in Proc. California Acad. iii, 231.—Hoopes, Evergreens, 244.—Parlatore in De Candolle Prodr. xvi², 436.—Koch, Dendrologie, ii², 193.—Vasey, Cat. Forest Trees, 36.—Stearns in Am. Nat. x, 110.—Watson, Bot. California, ii, 116.—Veitch, Manual Conif. 212.—Lawson, Pinetum Brit. iii, t. 52 & figs.

Taxodium sempervirens, Lambert, Pinus, 114; 2 ed. ii, 107, t. 52.—London, Arboretum, iv, 2487, f. 2340, 2341.—Hooker, Fl. Bor.-Am. ii, 164; Icon. iv, t. 379.—Hooker & Arnott, Bot. Beechey, 1841.—Fremont, Geographical Mem. California, 36, 37.—Henkel & Hochstetter, Nadelhölz. 262.

Taxodi species, Douglas in Companion Bot. Mag. ii, 150.

Sequoia gigantea, Endlicher, Syn. Conif. 190, in part.—Bentham, Pl. Hartweg. 338.

Abies religiosa, Hooker & Arnott, Bot. Beechey, 160.

Schubertia sempervirens, Spach, Hist. Veg. xi, 353.

S. religiosa, Presl, Epimel. Bot. 357.—Walpers, Ann. iii, 448.

Gigantabies taxifolia, Nelson, Pinaceæ, 78.

REDWOOD.

California, from the northern boundary of the state, south through the Coast ranges to "Veers creek" near the southern border of Monterey county.

A large tree of great economic value, 61 to 92 meters in height, with a trunk 2.40 to 7 meters in diameter, sending up from the stump when cut many vigorous shoots; sides of cañons and gulches in low, wet situations, borders of streams, etc., not appearing on dry hillsides; generally confined to the western slopes of the Coast ranges, and nowhere extending far from the coast; most generally multiplied and reaching its greatest average density north of cape Mendocino.

Wood light, soft, not strong, very brittle, rather coarse-grained, compact, susceptible of a good polish, easily split and worked, very durable in contact with the soil; bands of small summer cells thin, dark colored, conspicuous; medullary rays numerous, very obscure; color, clear light red, the thin sap-wood nearly white; specific gravity, 0.4208; ash, 0.14; largely sawed into lumber; the prevailing and most valuable building material of the Pacific coast, and in California almost exclusively used for shingles, fence posts, telegraph poles, railway ties, wine-butts, tanning- and water-tanks, coffins, etc.; forms with curled or contorted grain are highly ornamental.

343.—*Taxus brevifolia*, Nuttall,

Sylva, iii, 86, t. 108; 2 ed. ii, 149, t. 108 (*T. occidentalis* on plate).—Torrey in Pacific R. R. Rep. iv, 140.—Newberry in Pacific R. R. Rep. vi, 60, 90, f. 26.—Cooper in Smithsonian Rep. 1858, 263; Pacific R. R. Rep. xii³, 26, 69; Am. Nat. iii, 414.—Wood, Bot. & Fl. 316.—Bolander in Proc. California Acad. iii, 229.—Carrière, Trait. Conif. 2 ed. 742.—Hoopes, Evergreens, 383.—Parlatore in De Candolle, Prodr. xvi³, 501.—Gray in Proc. Am. Acad. vii, 402.—Koch, Dendrologie, ii³, 95.—Gordon, Pinetum, 2 ed. 392.—Vasey, Cat. Forest Trees, 35.—Macoun in Geological Rep. Canada, 1875-'76, 211.—Hall in Coulter's Bot. Gazette, ii, 91.—Watson, Bot. California, ii, 110.—G. M. Dawson in Canadian Nat. new ser. ix, 329.—Veitch, Manual Conif. 305.

T. baccata, var. *Canadensis*, Bentham, Pl. Hartweg. 338.

T. baccata, Hooker, Fl. Bor.-Am. ii, 167, in part.

T. Boursierii, Carrière in Rev. Hort. 1854, 228 & t.; Trait. Conif. 523; 2 ed. 739.

T. Lindleyana, Murray in Edinburgh New Phil. Jour. new ser. i, 294; Trans. Bot. Soc. Edinburgh, vi, 1860.—Lawson, Cat. 1855, 15.—Gordon, Pinetum, 316; Suppl. 99.—Henkel & Hochstetter, Nadelhölz. 360.—Nelson, Pinaceæ, 174.

T. Canadensis, Bigelow in Pacific R. R. Rep. iv, 25 [not Willdenow].

YEW

Queen Charlotte islands and the valley of the Skeena river, south through the Coast ranges of British Columbia, through western and the mountain ranges of eastern Washington territory and Oregon to the western slopes of the Rocky mountains of northern Montana (*Canby & Sargent*), through the California Coast ranges to the bay of Monterey and along the western slopes of the Sierra Nevadas to about latitude 37° N.

A tree 18 to 24 meters in height, with a trunk 0.60 to 0.90 meter in diameter, or toward its eastern limits in Idaho and Montana much smaller, often reduced to a low shrub; rare; low, rich woods and borders of streams, reaching its greatest development in western Oregon, Washington territory, and British Columbia.

Wood heavy, hard, strong, brittle, very close-grained, compact, susceptible of a beautiful polish, very durable in contact with the soil; bands of small summer cells thin, dark colored, conspicuous; medullary rays thin, numerous, very obscure; color, light bright red, the thin sap-wood light yellow; specific gravity, 0.6391; ash, 0.22; used for fence posts and by the Indians of the northwest coast for paddles, spear handles, bows, fish-hooks, etc.

344.—*Taxus Floridana*, Nuttall,

Sylva, iii, 92; 2 ed. ii, 155.—Croom in Am. Jour. Sci. 1 ser. xxvi, 334.—Chapman, Fl. S. States, 436.—Carrière, Trait. Conif. 2 ed. 741.—Hoopes, Evergreens, 384.—Vasey, Cat. Forest Trees, 36.

T. montana, Nuttall, Sylva, iii, 92; 2 ed. ii, 155.

YEW.

Western Florida, banks of the Apalachicola river from Bristol to Aspalaga, Gadsden county, and Watson's Landing? (*Curtiss*).

A small tree, 3 to 6 meters in height, with a trunk 0.15 to 0.25 meter in diameter; rare and very local.

Wood heavy, hard, very close-grained, compact; bands of small summer cells very thin, dark colored, not conspicuous; medullary rays numerous, obscure; color, dark brown tinged with red, the thin sap-wood nearly white; specific gravity, 0.6340; ash, 0.21.

345.—*Torreya taxifolia*, Arnott,

Ann. Nat. Hist. i, 134; Hooker, Icon. iii, t. 232, 233.—Eaton & Wright, Bot. 454.—Nuttall, Sylva, iii, 91, t. 109; 2 ed. ii, 153, t. 109.—Spach, Hist. Veg. xi, 298.—Endlicher, Syn. Conif. 241.—Lindley & Gordon in Jour. Hort. Soc. London, v, 226.—Darby, Bot. S. States, 516.—Carrière, Trait. Conif. 514; 2 ed. 726.—Gordon, Pinetum, 329; 2 ed. 412.—Cooper in Smithsonian Rep. 1858, 259.—Chapman, Fl. S. States, 436.—Wood, Cl. Book, 664; Bot. & Fl. 316.—Hoopes, Evergreens, 387, f. 62.—Parlatore in De Candolle, Prodr. xvi², 505.—Koch, Dendrologie, ii², 100.—Vasey, Cat. Forest Trees, 35.—Veitch, Manual Conif. 311.

Caryotaxus taxifolia, Henkel & Hochstetter, Nadelhölz. 367.

Fetataxus montana, Nelson, Pinaceæ, 167.

STINKING CEDAR. SAVIN.

Western Florida, eastern bank of the Apalachicola river from Chattahoochee to the neighborhood of Bristol, Gadsden county; doubtfully reported from the shores of a small lake west of Ocheesee and at Wakulla Springs, Wakulla county (*Curtiss*).

A tree 12 to 18 meters in height, with a trunk 0.60 to 0.90 meter in diameter, sending up when cut many vigorous shoots from the stem and roots; borders of swamps on calcareous soil; very rare and local.

Wood light, rather hard, strong, brittle, very close-grained, compact, susceptible of a beautiful polish, very durable in contact with the soil; bands of small summer cells very thin, not conspicuous; medullary rays numerous, obscure; color, clear bright yellow, the thin sap-wood much lighter; specific gravity, 0.5145; ash, 0.73; largely used locally for fence posts, etc.

346.—*Torreya Californica*, Torrey,

N. York Jour. Pharm. iii, 49; Pacific R. R. Rep. iv, 140.—Bigelow in Pacific R. R. Rep. iv, 24.—Kellogg in Proc. California Acad. 1, 35.—Newberry in Pacific R. R. Rep. vi, 61, 90, f. 27.—Cooper in Smithsonian Rep. 1858, 263.—Bolander in Proc. California Acad. iii, 229.—Hoopes, Evergreens, 385.—Parlatore in De Candolle, Prodr. xvi², 506.—Koch, Dendrologie, ii², 101.—Gordon, Pinetum, 2 ed. 410.—Vasey, Cat. Forest Trees, 35.—Watson, Bot. California, ii, 110.

T. Myristica, Hooker f. in Bot. Mag. t. 4780.—Van Houtte in Fl. des Serres, ix, 175 & t.—Carrière, Conif. 315; 2 ed. 727.—Gordon, Pinetum, 1 ed. 327.—Murray in Edinburgh New Phil. Jour. new ser. x, 7, t. 3.—Veitch, Manual Conif. 311.

Caryotaxus Myristica, Henkel & Hochstetter, Nadelhölz. 368.

Fetataxus Myristica, Nelson, Pinaceæ, 168.

CALIFORNIA NUTMEG. STINKING CEDAR.

California, Mendocino county, and along the western slope of the Sierra Nevadas to Tulare county, between 3,000 and 5,000 feet elevation.

A tree 15 to 22 meters in height, with a trunk 0.30 to 0.90 meter in diameter, sending up from the stump when cut many vigorous shoots; borders of streams, in moist soil; rare.

Wood light, soft, not strong, very close-grained, compact, susceptible of a fine polish, very durable in contact with the soil; bands of small summer cells broad, not conspicuous; medullary rays numerous, obscure; color, clear light yellow, the thin sap-wood nearly white; specific gravity, 0.4760; ash, 1.34.

347.—*Pinus Strobis*, Linnæus,

Spec. 1 ed. 1001; Du Roi, Harbk. ii, 57.—Wangenheim, Amer. i, t. 1, f. 1.—Aiton, Hort. Kew. iii, 369; 2 ed. v, 318.—Swartz, Obs. 363.—Mönch, Meth. 364.—Michaux, Fl. Bor.-Am. ii, 205.—Poiret in Lamarck, Dict. v, 341; Ill. iii, 369, t. 786, f. 2.—Lambert, Pinus, 1 ed. t. 22; 2 ed. i, 27, t. 35; 3 ed. i, 51, t. 32.—Willdenow, Spec. iv, 501; Enum. 939; Berl Baumz. 213.—Persoon, Syn. ii, 579.—Desfontaines, Hist. Arb. ii, 612.—Michaux f. Hist. Arb. Am. i, 104, t. 10; N. American Sylva, 3 ed. iii, 126, t. 145.—Nouveau Duhamel, v, 249, t. 76.—Smith in Roes' Cycl. xxviii, No. 17.—Pursh, Fl. Am. Sept. ii, 644.—Eaton, Manual, 110; 6 ed. 265.—Nuttall, Genera, ii, 223; Sylva, iii, 118; 2 ed. ii, 176 (excl. syn. var. *monticola*).—Hayne, Dend. Fl. 175.—Elliott, Sk. ii, 638.—Sprengel, Syst. ii, 887.—Torrey, Compend. Fl. N. States, 360; Fl. N. York, ii, 229.—Richard, Conif. 60, t. 12, f. 2.—Audubon, Birds, t. 39.—Beck, Bot. 339.—Loudon, Arboretum, iv, 2280, f. 2193–2196.—Forbes, Pinetum Woburn. 83.—Hooker, Fl. Bor.-Am. ii, 161.—Eaton & Wright, Bot. 359.—Bigelow, Fl. Boston. 3 ed. 385.—Antoine, Conif. 43, t. 20, f. 3.—Lindley in Penn. Cycl. xvii, 173.—Link in Linnæa, xv, 514.—Spach, Hist. Veg. xi, 394.—De Chambray, Trait. Arb. Res. Conif. 262, t. 4, 5, f. 8.—Emerson, Trees Massachusetts, 60; 2 ed. i, 73 & t.—Endlicher, Syn. Conif. 147.—Gihoul, Arb. Resin. 35, t. 5.—Knight, Syn. Conif. 34.—Lindley & Gordon in Jour. Hort. Soc. London. v, 215.—Carrière, Trait. Conif. 302; 2 ed. 398.—Buckley in Am. Jour. Sci. 2 ser. xiii, 398.—Darlington, Fl. Cestrica, 3 ed. 290.—Darby, Bot. S. States, 515.—Gordon, Pinetum, 239; 2 ed. 322.—Cooper in Smithsonian Rep. 1858, 257.—Fescali, Forst. Pf. 56, t. 11, f. 7–13.—Chapman, Fl. S. States, 434.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 25.—Wood, Cl. Book, 660; Bot. & Fl. 312.—Porcher, Resources S. Forests, 505.—Henkel & Hochstetter, Nadelhölz. 92.—Nelson, Pinaceæ, 130.—Hoopes, Evergreens, 136, f. 19.—Gray, Manual N. States, 5 ed. 470.—Parlatore in De Candolle, Prodr. xvi², 405.—Schnizlein, Icon. t. 77, f. 10.—Koch, Dendrologie, ii², 319.—Vasey, Cat. Forest Trees, 32.—Macoun in Geological Rep. Canada, 1875–'76, 211.—Sears in Bull. Essex Inst. xiii, 187.—Veitch, Manual Conif. 183.—Bell in Geological Rep. Canada, 1879–'80, 49^c.

P. Strobis, var. *alba*, var. *brevifolia*, var. *compressa*, Loudon, Arboretum, iv, 2280.—Lindley & Gordon in Jour. Hort. Soc. London, v, 215.

P. Strobis, var. *nivea*, Hort.

WHITE PINE. WEYMOUTH PINE.

Newfoundland, northern shores of the gulf of Saint Lawrence to lake Nipigon and the valley of the Winnipeg river, south through the northern states to Pennsylvania, the southern shores of lake Michigan, "Starving rock," near La Salle, Illinois, near Davenport, Iowa (*Parry*), and along the Alleghany mountains to northern Georgia.

A large tree of the first economic value, 24 to 52 meters in height, with a trunk 1.20 to 3.50 meters in diameter; sandy loam upon drift formations, forming extensive forests, or in the region of the great lakes often in small bodies scattered through the hard-wood forests, here reaching its greatest development; north of latitude 47° N. and south of Pennsylvania, central Michigan, and Minnesota much smaller, less common and valuable.

Wood light, soft, not strong, very close, straight-grained, compact, easily worked, susceptible of a beautiful polish; bands of small summer cells thin, not conspicuous, resin passages small, not numerous nor conspicuous; medullary rays numerous, thin; color, light brown, often slightly tinged with red; the sap-wood nearly white; specific gravity, 0.3854; ash, 0.19; more largely manufactured into lumber, shingles, laths, etc., than that of any other North American tree; the common and most valuable building material of the northern states; largely used in cabinet-making, for interior finish, and in the manufacture of matches, woodenware, and for many domestic purposes.

Coniferin, a glucoside principle, has been discovered in the cambium layer of this and several other species of *Coniferæ* (*Jour. für Prakt. Chem.* xcvi, 243.—*Am. Jour. Pharm.* 1867, 261.—*U. S. Dispensatory*, 14 ed. 901).

348.—*Pinus monticola*, Douglas;

Lambert, Pinus, 1 ed. iii, 27, t. 35.—Loudon, Arboretum, iv, 2291, f. 2208, 2209.—Forbes, Pinetum Woburn. 81, t. 31.—Antoine, Conif. 40, t. 18, f. 3.—Hooker & Arnott, Bot. Beechey, 394.—Endlicher, Syn. Conif. 148.—Lindley & Gordon in Jour. Hort. Soc. London, v, 215.—Carrière, Trait. Conif. 305; 2 ed. 401.—Gordon, Pinetum, 233; 2 ed. 314.—Cooper in Smithsonian Rep. 1858, 262; Pacific R. R. Rep. xii², 27; Am. Nat. iii, 410.—Lyll in Jour. Linnæan Soc. vii, 141.—Henkel & Hochstetter, Nadelhölz. 94.—Nelson, Pinaceæ, 120.—Hoopes, Evergreens, 135.—Bolander in Proc. California Acad. iii, 318.—Parlatore in De Candolle, Prodr. xvi², 405.—Gray in Proc. Am. Acad. vii, 402.—Fowler in London Gard. Chronicle, 1872, 1071.—Koch, Dendrologie, ii², 322.—Vasey, Cat. Forest Trees, 32.—Macoun in Geological Rep. Canada, 1875–'76, 211.—Hall in Coulter's Bot. Gazette, ii, 91.—Engelmann in Bot. California, ii, 123.—G. M. Dawson in Canadian Nat. new ser. ix, 328.—Veitch, Manual Conif. 181, f. 41.—Lawson, Pinetum Brit. i, 69, f. 1–6.

P. Strobis, var. *monticola*, Nuttall, Sylva, iii, 118; 2 ed. ii, 176.

P. Grozelieri, Carrière in Rev. Hort. 1869, 126.

P. porphyrocarpa, Lawson, Pinetum Brit. i, 83, f. 1–8.

WHITE PINE.

Vancouver's island, Coast and Gold ranges of southern British Columbia, through the Cœur d'Alène and Bitter Root mountains of Idaho to the valley of the Flathead river, northern Montana (*Canby & Sargent*), south along the Cascade mountains of Washington territory and Oregon and the California sierras to Calaveras county.

A large tree, 30 to 46 meters in height, with a trunk 0.90 to 1.50 meter in diameter; most common and reaching its greatest development in the Pend d'Oreille and Clark's Fork regions of Idaho, here a valuable and important timber tree; in British Columbia generally below 3,000 feet, and in California between 7,000 and 10,000 feet elevation; not common.

Wood very light, soft, not strong, close, straight-grained, compact; bands of small summer cells thin, resinous, not conspicuous, resin passages numerous, not large, conspicuous; medullary rays numerous, obscure; color, light brown or red, the sap-wood nearly white; specific gravity, 0.3908; ash, 0.23; inferior in quality, although resembling that of the eastern white pine (*P. Strobus*); in Idaho and Montana somewhat manufactured into lumber.

349.—*Pinus Lambertiana*, Douglas,

Companion Bot. Mag. ii, 92, 106, 107, 130, 152; Trans. Linnæan Soc. xv, 500.—Lambert, *Pinus*, 1 ed. iii, 157, t. 68, 69.—Loudon, *Arboretum*, iv, 2288, f. 2203.—Forbes, *Pinetum Woburn*, 77, t. 30.—Hooker, *Fl. Bor.-Am.* ii, 161.—Antoine, *Conif.* 41, t. 19.—Lindley in *Penn. Cyl.* xvii, 173.—Hooker & Arnott, *Bot. Beechey*, 394.—Spach, *Hist. Veg.* xi, 397.—Nuttall, *Sylva*, iii, 122, t. 114; 2 ed. ii, 180, t. 114.—De Chambray, *Trait. Arb. Res. Conif.* 346.—Endlicher, *Syn. Conif.* 150.—Lindley & Gordon in *Jour. Hort. Soc. London*, v, 215.—Carrière, *Trait. Conif.* 307; 2 ed. 403.—Bigelow in *Pacific R. R. Rep.* iv, 21.—Torrey in *Pacific R. R. Rep.* iv, 141; *Bot. Mex. Boundary Survey*, 210; Ives' *Rep.* 28.—Newberry in *Pacific R. R. Rep.* vi, 42, 90, f. 14.—Gordon, *Pinetum*, 228; 2 ed. 307.—Cooper in *Smithsonian Rep.* 1858, 262.—Murray in *Trans. Bot. Soc. Edinburgh*, vi, 369.—Lawson, *Pinetum Brit.* i, 47, t. 7, f. 1-7.—Bolander in *Proc. California Acad.* iii, 226, 317.—Henkel & Hochstetter, *Nadelhölz.* 95.—Nelson, *Pinaceæ*, 115.—Hoopes, *Evergreens*, 134.—Parlatore in *De Candolle, Prodr.* xvi², 402.—Fowler in *London Gard. Chronicle*, 1872, 1071.—Koch, *Dendrologie*, ii², 323.—Vasey, *Cat. Forest Trees*, 32.—Veitch, *Manual Conif.* 179.

SUGAR PINE.

Oregon, Cascade and Coast ranges, from the head of the Mackenzie river and the valley of the Rogue river south along the western flank of the California sierras, through the Coast ranges to the Santa Lucia mountains, and in the San Bernardino and Cuyamaca mountains.

A large tree, 46 to 92 meters in height, with a trunk 3 to 7 meters in diameter; most common and reaching its greatest development upon the sierras of central and northern California between 4,000 and 8,000 feet elevation; in the Oregon Coast ranges descending to 1,000 feet above the sea-level.

Wood very light, soft, coarse, straight-grained, compact, satiny, easily worked; bands of small summer cells thin, resinous, conspicuous, resin passages numerous, very large and conspicuous; medullary rays numerous, obscure; color, light brown, the sap-wood nearly white; specific gravity, 0.3684; ash, 0.22; now largely manufactured into lumber and used for interior finish, door-blinds, sashes, etc., and for cooperage and woodenware; less valuable and less easily worked than that of the eastern white pine (*Pinus Strobus*); its quality injured by the larger and more numerous resin passages.

A saccharine exudation from the stumps of cut or partially-burned trees sometimes used as a substitute for sugar.

350.—*Pinus flexilis*, James,

Long's Exped. ii, 27, 34.—Torrey in *Ann. Lye. N. York*, ii, 249; *Pacific R. R. Rep.* iv, 141.—Eaton, *Manual*, 6 ed. 265.—Eaton & Wright, *Bot.* 359.—Nuttall, *Sylva*, iii, 107, t. 112; 2 ed. ii, 167, t. 107.—Lindley & Gordon in *Jour. Hort. Soc. London*, v, 220.—Carrière in *Fl. des Serres*, ix, 200; *Rev. Hort.* 1854, 228; *Trait. Conif.* 310; 2 ed. 392.—Bigelow in *Pacific R. R. Rep.* iv, 6, 20.—Gordon, *Pinetum*, 224; 2 ed. 302.—Cooper in *Smithsonian Rep.* 1858, 262.—Parry in *Trans. St. Louis Acad.* ii, 121.—Engelmann in *Am. Jour. Sci.* 2 ser. xxxiv, 331; *Trans. St. Louis Acad.* ii, 208; *Wheeler's Rep.* vi, 257; *Bot. California*, ii, 124.—Henkel & Hochstetter, *Nadelhölz.* 126.—Nelson, *Pinaceæ*, 112.—Bolander in *Proc. California Acad.* iii, 318.—Hoopes, *Evergreens*, 131, f. 18.—Parlatore in *De Candolle, Prodr.* xvi², 403.—Porter in *Hayden's Rep.* 1871, 494.—Watson in *King's Rep.* v, xxviii, 332; *Pl. Wheeler*, 17.—Rothrock, *Pl. Wheeler*, 27, 50; *Wheeler's Rep.* vi, 9.—Porter & Coulter, *Fl. Colorado*; *Hayden, Surv. Misc. Pub. No. 4*, 130.—Murray in *London Gard. Chronicle*, 1875, 106.—Vasey, *Cat. Forest Trees*, 32.—Sargent in *Am. Jour. Sci.* 3 ser. xvii, 420.—Lawson, *Pinetum Brit.* i, 35, f. 1.

P. Lambertiana, var. Hooker, *Fl. Bor.-Am.* ii, 161.

P. Lambertiana, var. *brevifolia*, Endlicher, *Syn. Conif.* 150.—Lindley & Gordon in *Jour. Hort. Soc. London*, v, 215.—Carrière, *Trait. Conif.* 2 ed. 404.

P. flexilis, var. *serrulata*, Engelmann in *Wheeler's Rep.* vi, 258.

P. flexilis, var. *macrocarpa*, Engelmann in *Wheeler's Rep.* vi, 258.

WHITE PINE.

Eastern slopes of the Rocky mountains, Montana, and probably much farther north, south to New Mexico, on the Guadalupe and Limpia mountains, western Texas (*Havard*), on the high mountain ranges of Utah, Nevada, and northern Arizona, Inyo mountains and mount Silliman, California.

A tree 15 to 18 meters in height, with a trunk 0.60 to 1.20 meter in diameter; dry, gravelly slopes and ridges between 4,000 and 10,000 feet elevation; common along the eastern slopes of the Rocky mountains of northern Montana, forming open, scattered forests, here low, round-topped, and the prevailing forest tree; in central Nevada the most valuable lumber tree of the region.

Wood light, soft, close-grained, compact; bands of small summer cells narrow, not conspicuous, resin passages numerous, large; medullary rays numerous, conspicuous; color, light clear yellow, turning red with exposure, the sap-wood nearly white; specific gravity, 0.4358; ash, 0.28; in northern Montana, Nevada, and Utah sometimes sawed into inferior lumber and used in construction and for various domestic purposes.

351.—*Pinus albicaulis*, Engelmann,

Trans. St. Louis Acad. ii, 209; Coulter's Bot. Gazette, vii, 4.—Gray in Proc. Am. Acad. vii, 402.—Vasey, Cat. Forest Trees, 32.—Hall in Coulter's Bot. Gazette, ii, 91.—Lawson, Pinetum Brit. i, 1, f. 1-4.

P. flexilis, Murray, Rep. Oregon Exped. i, t. 2, f. 1 [not James].—Lyal in Jour. Linnæan Soc. vii, 142.—Parlatore in De Candolle, Prodr. xvi², 403, in part.

P. cembroides, Newberry in Pacific R. R. Rep. vi, 44, 90, f. 15 [not Zuccarini].

P. Shasta, Carrière, Trait. Conif. 2 ed. 390.

P. flexilis, var. *albicaulis*, Engelmann in Bot. California, ii, 124.—G. M. Dawson in Canadian Nat. new. ser. ix, 328.

Coast ranges of British Columbia, from the valley of the Lltasyouco river (*G. M. Dawson*) south along the Cascade and Blue mountains of Washington territory and Oregon, extending east along the high ranges of northern Washington territory to the eastern slope of the Rocky mountains of northern Montana (Old Marias pass, *Canby & Sargent*); California, Scott's mountains, mount Shasta, and on the high peaks of the Sierra Nevadas to mount San Bernardino.

A small alpine tree, 6 to 12 meters in height, with a trunk rarely 0.60 meter in diameter, or at its highest elevation reduced to a low, prostrate shrub; dry, gravelly ridges at the extreme limit of tree growth, reaching in the San Bernardino mountains an elevation of 10,500 feet.

Wood light, soft, not strong, brittle, close-grained, compact; bands of small summer cells thin, not conspicuous, resin passages numerous, not large; medullary rays numerous, obscure; color, light brown, the sap-wood nearly white; specific gravity, 0.4165; ash, 0.27.

352.—*Pinus reflexa*, Engelmann,

Coulter's Bot. Gazette, vii, 4.—Rusby in Bull. Torrey Bot. Club, ix, 80.

P. flexilis, var. *reflexa*, Engelmann in Wheeler's Rep. vi, 258.

WHITE PINE.

High mountains of southwestern New Mexico (*Greene, Rusby*) to the Santa Rita mountains (*Rothrock, Engelmann & Sargent*) and Santa Catalina mountains (*Lemmon, Pringle*), Arizona.

A tree 24 to 30 meters in height, with a trunk sometimes exceeding 0.60 meter in diameter; rocky ridges and slopes of almost inaccessible cañons between 6,000 to 8,000 feet elevation.

Wood light, hard, not strong, close-grained, compact; bands of small summer cells thin, resinous, not conspicuous, resin passages large, not numerous; medullary rays numerous, obscure; color, light red, the sap-wood nearly white; specific gravity, 0.4877; ash, 0.26.

353.—*Pinus Parryana*, Engelmann,

Am. Jour. Sci. 2 ser. xxiv, 332, note; Bot. California, ii, 124.—Parlatore in De Candolle, Prodr. xvi², 402.—Vasey, Cat. Forest Trees, 30.

P. Llaveana, Torrey, Bot. Mex. Boundary Survey, 208, t. 55 [not Schiede & Deppe].—Cooper in Smithsonian Rep. 1858, 262.—Bolander in Proc. California Acad. iii, 318.

PIÑON. NUT PINE.

California, Larkin's station, 20 miles southeast of Campo, San Diego county (*Vasey*), and southward into Lower California.

A small tree, 6 to 9 meters in height, with a trunk 0.30 to 0.45 meter in diameter; very rare within the limits of the United States; south of the boundary forming extensive open forests upon the high *mesas* and slopes of Lower California (*Pringle*).

Wood light, soft, close-grained, compact; bands of small summer cells thin, not conspicuous, resin passages very numerous, large, conspicuous; medullary rays numerous, obscure; color, light brown or yellow, the sap-wood much lighter, nearly white; specific gravity, 0.5675; ash, 0.54.

The large seeds edible.

354.—*Pinus cembroides*, Zuccarini,

Flora, ii, 93.—Endlicher, Syn. Conif. 182.—Fl. des Serres, iv, 3446, t. 97.—Nelson, Pinaceæ, 107.—Parlatore in De Candolle, Prodr. xvi², 397.—Engelmann in Trans. St. Louis Acad. iv, 176.—Watson in Proc. Am. Acad. xviii, 158.

P. Llaveana, Schiede & Deppe in Linnæa, xii, 468.—Forbes, Pinetum Woburn. 49, t. 17.—Antoine, Conif. 36, t. 16, f. 1.—Spach, Hist. Veg. xi, 401.—Lindley & Gordon in Jour. Hort. Soc. London, v, 216.—Carrière, Trait. Conif. 405; 2 ed. 461.—Gordon, Pinetum, 199; 2 ed. 274 (excl. syn. *edulis*).—Henkel & Hochstetter, Nadelhölz. 64 (excl. syn. *edulis*).—Hoopes, Evergreens, 143.

P. osteosperma, Engelmann in Wislizenus' Rep. No. 3.—Lindley & Gordon in Jour. Hort. Soc. London, v, 216.—Carrière in Fl. des Serres, ix, 200; Rev. Hort. 1854, 227.

NUT PINE.

Santa Catalina mountains, Arizona (*Pringle*); through northern Mexico.

A small tree, in Arizona 6 to 7 meters in height, with a trunk hardly exceeding 0.30 meter in diameter; dry ridges and slopes at 3,500 feet elevation.

Wood light, soft, very close-grained, compact; bands of small summer cells thin, not conspicuous, resin passages few, small; medullary rays numerous, obscure; color, light clear yellow, the sap-wood nearly white; specific gravity, 0.6512; ash, 0.90.

The seeds edible.

355.—*Pinus edulis*, Engelmann,

Wislizenus' Rep. No. 4; Wheeler's Rep. vi, 260.—Lindley & Gordon in Jour. Hort. Soc. London, v, 216.—Carrière, Fl. des Serres, ix, 201; Rev. Hort. 1854, 227; Trait. Conif. 408.—Torrey in Sitgreaves' Rep. 173, t. 20; Pacific R. R. Rep. iv, 140; Ives' Rep. 28.—Bigelow in Pacific R. R. Rep. iv, 3, 19.—Cooper in Smithsonian Rep. 1853, 261.—Hoopes, Evergreens, 142.—Parlatore in De Candolle, Prodr. xvi², 398.—Watson in Pl. Wheeler, 17.—Porter & Coulter, Fl. Colorado; Hayden's Surv. Misc. Pub. No. 4, 130.—Vasey, Cat. Forest Trees, 30.—Rothrock in Wheeler's Rep. vi, 9.—Rusby in Bull. Torrey Bot. Club, ix, 106.—Veitch, Manual Conif. 172.

P. cembroides, Gordon in Jour. Hort. Soc. London, v, 236 & f.; Pinetum, 192; 2 ed. 265 [not Zuccarini].—Fl. des Serres, iv, 324^b, 325^b, t. 331, f. 97.—Lindley & Gordon in Jour. Hort. Soc. London, v, 216.—Carrière, Trait. Conif. 404; 2 ed. 460.

P. futilis, Roehl in herb. *Adel* Gordon, Pinetum, Suppl. 76; 2 ed. 265.

PIÑON. NUT PINE.

Eastern base of Pike's peak, Colorado, south through New Mexico to the mountains of western Texas.

A small tree, 6 to 9 meters in height, with a trunk 0.30 to 0.90 meter in diameter; dry *mesas* and slopes, generally on lime or sandstone, reaching in Colorado an elevation of 9,000 feet.

Wood light, soft, not strong, brittle, close-grained, compact, durable in contact with the soil; bands of small summer cells thin, not conspicuous, resin passages few, small; medullary rays numerous, obscure; color, light brown, the sap-wood nearly white; specific gravity, 0.6388; ash, 0.62; largely used for fuel, charcoal, fencing, etc., and in western Texas occasionally manufactured into inferior lumber.

The large edible nuts supply the Indians with a valuable article of food.

356.—*Pinus monophylla*, Torrey & Fremont,

Fremont's Rep. 319, t. 4.—Cooper in Smithsonian Rep. 1858, 261.—Bolander in Proc. California Acad. iii, 318.—Hoopes, Evergreens, 142.—Parlatore in De Candolle, Prodr. xvi², 378.—Lawson, Pinetum Brit. i, 65, t. 9, f. 1-12 (*P. Fremontiana* on plate).—Watson in King's Rep. v, 330; Pl. Wheeler, 17.—Koch, Dendrologie, ii², 271.—Bertrand in Bull. Soc. Bot. France, xviii, 81, t. 5, f. 81.—Rothrock in Pl. Wheeler, 28, 50.—Vasey, Cat. Forest Trees, 30.—Palmer in Am. Nat. xii, 594.—Engelmann in Wheeler's Rep. vi, 259, 374; Trans. St. Louis Acad. iv, 178; Bot. California, ii, 124.—Sargent in Am. Jour. Sci. 3 ser. xvii, 419.—Masters in London Gard. Chronicle, 1883, p. 48, f. 8.

P. Fremontiana, Endlicher, Syn. Conif. 1831, in part.—Gordon in Jour. Hort. Soc. London, iv, 293 & f.; Pinetum, 194; 2 ed. 235.—Knight, Syn. Conif. 28.—Lindley & Gordon in Jour. Hort. Soc. London, v, 216.—Carrière, Trait. Conif. 194; 2 ed. 462.—Henkel & Hochstetter, Nadelhölz. 62.

PIÑON. NUT PINE.

Near Utah lake, Utah, to the eastern foot-hills of the California sierras, south along the mountain ranges of the Great Basin to the San Francisco mountains of eastern Arizona.

A small, bushy tree, 4 to 6 meters in height, with a trunk sometimes 1 meter in diameter; dry, gravelly slopes and *mesas* between 3,000 and 6,000 feet elevation.

Wood light, soft, weak, brittle, close-grained, compact; bands of small summer cells thin, not conspicuous, resin passages few, not large; medullary rays numerous, obscure; color, yellow or light brown, the sap-wood nearly white; specific gravity, 0.5658; ash, 0.68; largely used for fuel and charcoal.

The large edible seeds furnish the principal food of the Indians of the Great Basin.

357.—*Pinus Balfouriana*, Murray,

Rep. Oregon Exped. i, t. 3, f. 1.—Gordon, Pinetum, 217; 2 ed. 293.—Henkel & Hochstetter, Nadelhölz. 109.—Bolander in Proc. California Acad. iii, 318.—Carrière, Trait. Conif. 2 ed. 425.—Nelson, Pinaceæ, 104.—Hoopes, Evergreens, 149.—Fowler in London Gard. Chronicle, 1872, 973.—Vasey, Cat. Forest Trees, 32.—Engelmann in Trans. St. Louis Acad. iv, 179; Bot. California, ii, 125.—Veitch, Manual Conif. 175.—Lawson, Pinetum Brit. i, 11, f. 1-5.

California, Scott's mountain, Siskiyou county (*Jeffrey, Lemmon*), mount Whitney, and about the headwaters of King and Kern rivers.

A small tree, 15 to 19 meters in height, with a trunk 0.60 to 0.90 meter in diameter; dry, gravelly slopes and ridges, forming upon Scott's mountain a broad belt of open forest growth between 5,000 and 8,000 feet elevation.

Wood light, soft, weak, brittle, very close-grained, compact, satiny, susceptible of a good polish; bands of small summer cells very narrow, dark colored, resin passages few, not conspicuous; medullary rays numerous obscure; specific gravity, 0.5434; ash, 0.41.

Var. *aristata*, Engelmann,

Wheeler's Rep. vi, 375.—Bot. California, ii, 125.—Veitch, Manual Conif. 175.

P. aristata, Engelmann in Am. Jour. Sci. 2 ser. xxxiv, 331; Trans. St. Louis Acad. ii, 205, t. 5, 6; iv, 179; Bot. California, ii, 125.—Parry in Trans. St. Louis Acad. ii, 123.—Wood, Bot. & Fl. 313.—Regel, Gartenflora, 1863, iii, 91.—Henkel & Hochstetter, Nadelhölz. 417.—Nelson, Pinaceæ, 103.—Carrière, Trait. Conif. 2 ed. 424.—Parlatore in De Candolle, Prodr. xvi³, 400.—Porter & Coulter, Fl. Colorado; Hayden's Surv. Misc. Pub. No. 4, 130.—Murray in London Gard. Chronicle, 1875, 106.—Gordon, Pinetum, 2 ed. 291.—Vasey, Cat. Forest Trees, 32.—Brandegge in Coulter's Bot. Gazette, 32.—Lawson, Pinetum Brit. i, 5, f. 1.

P. Balfouriana, Watson in King's Rep. v, 331; Pl. Wheeler, 17 [not Murray].—Rothrock in Pl. Wheeler, 28, 50.—Sargent in Am. Jour. Sci. 3 ser. xvii, 419.

FOXTAIL PINE. HICKORY PINE.

Mountains of southeastern California, Nevada, northern Arizona, and southern Utah to Colorado, above 7,500 feet, or in Colorado reaching 12,000 feet elevation.

A tree 15 to 30 meters in height, with a trunk 0.60 to 2.40 meters in diameter; dry, gravelly ridges; not common.

Wood light, soft, not strong, very close-grained, compact; bands of small summer cells thin, dark colored, not conspicuous, resin passages few, not prominent; medullary rays numerous, obscure; color, red, the thin sap-wood nearly white; specific gravity, 0.5572; ash, 0.30; in central Nevada largely used for the timbering of mines, and now nearly exterminated.

358.—*Pinus resinosa*, Aiton,

Hort. Kew. iii, 337; 2 ed. v, 316.—Lambert, Pinus, 1 ed. t. 14; 2 ed. i, 20, t. 14; 3 ed. i, 17, t. 13.—Willdenow, Spec. iv, 496; Enum. 988; Berl. Baumz. 267.—Poiret in Lamarck, Dict. v, 339.—Persoon, Syn. ii, 578.—Desfontaines, Hist. Arb. ii, 612.—Smith in Rees' Cycl. xxviii, No. 3.—Pursh, Fl. Am. Sept. ii, 642.—Eaton, Manual, 110; 6 ed. 264.—Nuttall, Genera, ii, 223.—Hayne, Dend. Fl. 173.—Sprengel, Syst. ii, 886.—Torrey, Compend. Fl. N. States, 360; Fl. N. York, ii, 227.—Beck, Bot. 339.—London, Arboretum, iv, 2210, f. 2094-2097.—Forbes, Pinetum Woburn. 19, t. 6.—Hooker, Fl. Bor.-Am. ii, 161, in part.—Eaton & Wright, Bot. 358.—Bigelow, Fl. Boston. 3 ed. 334.—Lindley in Penn. Cycl. xvii, 170.—Antoine, Conif. 7, t. 4, f. 1.—Link in Linnæa, xv, 501.—Endlicher, Syn. Conif. 178.—Knight, Syn. Conif. 27.—Lindley & Gordon in Jour. Hort. Soc. London, v, 219.—Parry in Owen's Rep. 618.—Carrière, Trait. Conif. 401.—Gordon, Pinetum, 183 (excl. syn. *Loiseleuriana*); 2 ed. 256.—Richardson Arctic Exped. 441.—Cooper in Smithsonian Rep. 1858, 257.—Wood, Cl. Book, 631; Bot. & Fl. 313.—Henkel & Hochstetter, Nadelhölz. 45 (excl. syn. *Loiseleuriana*).—Hoopes, Evergreens, 102.—Gray, Manual N. States, 5 ed. 470.—Parlatore in De Candolle, Prodr. xvi³, 388.—Koch, Dendrologie, ii³, 286.—Vasey, Cat. Forest Trees, 30.—Macoun in Geological Rep. Canada, 1875-'76, 211.—Engelmann in Trans. St. Louis Acad. iv, 179.—Sears in Bull. Essex Inst. xiii, 185.—Bell in Geological Rep. Canada, 1879-'80, 56c.—Veitch, Manual Conif. 159.

P. rubra, Michaux f. Hist. Arb. Am. i, 46, t. 1; N. American Sylva, 3 ed. iii, 91, t. 134 [not Lambert].—De Chambray, Trait. Arb. Res. 344.—Gihoul, Arb. Resin. 27.—Carrière, Trait. Conif. 2 ed. 496.

P. Laricio, var. *resinosa*, Spach, Hist. Veg. 385.

RED PINE. NORWAY PINE.

Newfoundland, northern shores of the gulf of Saint Lawrence and lake Nipigon to the valley of the Winnipeg river, south through the northern states to Chestnut Hill, Middlesex county, Massachusetts, the mountains of northern Pennsylvania, Isabella county, Michigan, and central Minnesota.

A large tree, 24 to 46 meters in height, with a trunk 0.60 to 1.37 meter in diameter; light sandy loam or dry, rocky ridges, forming scattered groves rarely exceeding a few hundred acres in extent; common and reaching its greatest development through northern Wisconsin and Minnesota; rare in the eastern States, except in the extreme northern portions of New England.

Wood light, not strong, hard, rather coarse-grained, compact; bands of small summer cells broad, dark colored, very resinous, resin passages few, small, not conspicuous; medullary rays numerous, thin; color, light red, the sap-wood yellow or often almost white; specific gravity, 0.4854; ash, 0.27; largely manufactured into lumber and used for all purposes of construction, flooring, piles, etc.

359.—*Pinus Torreyana*, Parry,

Bot. Mex. Boundary Survey, 210, t. 58, 59; Proc. San Diego Nat. Hist. Soc. Nov. 1883.—Carrière, Trait. Conif. 326; 2 ed. 423.—Gordon, Pinetum, 241.—Cooper in Smithsonian Rep. 1860, 442.—Henkel & Hochstetter, Nadelhölz. 117.—Bolander in Proc. California Acad. iii, 318.—Hoopes, Evergreens, 150.—Vasey, Cat. Forest Trees, 31.—Palmer in Am. Nat. xii, 594.—Engelmann in Trans. St. Louis Acad. iv, 181; Bot. California, ii, 125.—Veitch, Manual Conif. 173.

P. lophosperma, Lindley in London Gard. Chronicle, 1860, 46.—Gordon, Pinetum, Suppl. 69; 2 ed. 310.—Henkel & Hochstetter, Nadelhölz. 112.—Nelson, Pinaceæ, 117.—Parlatore in De Candolle, Prodr. xvi², 391.

California, mouth of the Soledad river, San Diego county; doubtfully reported from one of the islands off Santa Barbara and from Lower California.

A low, short-lived, gnarled, crooked tree, 6 to 8 meters in height, with a trunk 0.23 to 0.33 meter in diameter; crests of sandy bluffs immediately upon the sea-coast; very local and fast disappearing.

Wood light, soft, not strong, brittle, rather close-grained, compact; bands of small summer cells broad, resinous, conspicuous, resin passages small, few; medullary rays numerous, obscure; color, light red, the sap-wood yellow or nearly white; specific gravity, 0.4879; ash, 0.35; locally used for fuel.

360.—*Pinus Arizona*, Engelmann,

Wheeler's Rep. vi, 260; Trans. St. Louis Acad. iv, 181; Coulter's Bot. Gazette, vii, 4.

YELLOW PINE.

Santa Rita mountains (*Rothrock, Engelmann & Sargent*), Santa Catalina mountains (*Lemmon, Pringle*), and probably upon other ranges of southern Arizona.

A tree 24 to 30 meters in height, with a trunk 0.60 to 0.90 meter in diameter; high rocky ridges between 6,000 and 8,000 feet elevation; the prevailing forest tree over large areas near the summits of the Santa Catalina mountains (*Lemmon*).

Wood light, soft, not strong, rather brittle, close grained, compact; bands of small summer cells broad, very resinous, conspicuous, resin passages numerous, large; medullary rays thin, obscure; color, light red or often yellow, the sap-wood lighter yellow or white; specific gravity, 0.5038; ash, 0.20; sometimes sawed into inferior lumber.

361.—*Pinus ponderosa*, Douglas,

Companion Bot. Mag. ii, 111.—London, Arboretum, iv, 2243, f. 2132–2136.—Forbes, Pinetum Woburn. 44, t. 15.—Antoine, Conif. 28, t. 8, f. 1.—Lindley in Penn. Cycl. xvii, 172.—Link in Linnæa, xv, 306.—Nuttall, Sylva, iii, 114; 2 ed. ii, 173.—Spach, Hist. Veg. xi, 389.—Endlicher, Syn. Conif. 163.—Knight, Syn. Conif. 30.—Lindley & Gordon in Jour. Hort. Soc. London, v, 217.—Carrière, Trait. Conif. 340; 2 ed. 445.—Gordon, Pinetum, 205; Suppl. 67; 2 ed. 281.—Newberry in Pacific R. R. Rep. vi, 36, 90, t. 4, f. 12.—Cooper in Smithsonian Rep. 1858, 261; Pacific R. R. Rep. xii², 27, 68; Am. Nat. iii, 409.—Torrey, Bot. Mex. Boundary Survey, 209; Ives' Rep. 28.—Engelmann in Am. Jour. Sci. 2 ser. xxxiv, 332; Proc. Am. Phil. Soc. 2 ser. xii, 209; Wheeler's Rep. vi, 261; Trans. St. Louis Acad. iv, 181; Bot. California, ii, 125.—Lyll in Jour. Linnæan Soc. vii, 142.—Bolander in Proc. California Acad. iii, 226, 317.—Henkel & Hochstetter, Nadelhölz. 71.—Nelson, Pinaceæ, 125.—Hoopes, Evergreens, 117.—Parlatore in De Candolle, Prodr. xvi², 395 (excl. syn. *Sinclairii*).—Watson in King's Rep. v, 331; Pl. Wheeler, 17.—Gray in Proc. Am. Acad. vii, 402.—Fowler in London Gard. Chronicle, 1872, 1326.—Koch, Dendrologie, ii², 310.—Rothrock in Pl. Wheeler, 28, 50; Wheeler's Rep. vi, 9.—Porter & Coulter, Fl. Colorado; Hayden's Surv. Misc. Pub. No. 4, 129.—Hayden in Warren's Rep. Nebraska & Dakota, 2 ed. 121.—Vasey, Cat. Forest Trees, 30.—Hall in Coulter's Bot. Gazette, ii, 91.—Macoun in Geological Rep. Canada, 1875-'76, 211.—Brandegge in Coulter's Bot. Gazette, iii, 32.—G. M. Dawson in Canadian Nat. new ser. ix, 326.—Rusby in Bull. Torrey Bot. Club, ix, 106.

- P. Benthamiana*, Hartweg in Jour. Hort. Soc. London, ii, 189; iii, 223.—Gordon in Jour. Hort. Soc. London, iv, 212 & t.; (Fl. des Serres, vi, 85 & f.); Pinetum, 188; 2 ed. 261 (excl. syn. *Sinclairii*).—Knight, Syn. Conif. 30.—Lindley & Gordon in Jour. Hort. Soc. London, v, 216.—Carrière, Trait. Conif. 350; 2 ed. 452.—Murray in Edinburgh New Phil. Jour. new ser. i, 287, t. 8.—Henkel & Hochstetter, Nadelhölz. 84.—Nelson, Pinaceæ, 104.—Fowler in London Gard. Chronicle, 1872, 973.
- P. resinosa*, Torrey in Ann. Lyc. N. York, ii, 249 [not Aiton].—Douglas, Companion Bot. Mag. ii, 126.—Hooker, Fl. Bor.-Am. ii, 161, in part.—Winchell in Ludlow's Rep. Black Hills, 68.
- P. brachyptera*, Engelmann in Wislizenus' Rep. No. 4.—Lindley & Gordon in Jour. Hort. Soc. London, v, 216.—Carrière in Fl. des Serres, ix, 201; Rev. Hort. 1854, 227; Trait. Conif. 356; 2 ed. 454.—Bigelow in Pacific R. R. Rep. iv 18.—Gordon, Pinetum, 190; 2 ed. 263.—Henkel & Hochstetter, Nadelhölz. 85.—Nelson, Pinaceæ, 454.
- P. Beardsleyi*, Murray in Edinburgh New Phil. Jour. new ser. i, 286, t. 6.—Carrière, Trait. Conif. 359.
- P. Craigana*, Murray in Edinburgh New Phil. Jour. new ser. i, 288, t. 7.
- P. macrophylla*, ? Torrey in Sitgreaves' Rep. 173 [not Engelmann].
- P. Engelmanni*, Torrey in Pacific R. R. Rep. iv, 141 [not Carrière].
- P. Parryana*, Gordon, Pinetum, 202; 2 ed. 277 [not Engelmann].—Henkel & Hochstetter, Nadelhölz. 88.—Carrière, Trait. Conif. 2 ed. 446.
- P. ponderosa*, var. *Benthamiana*, Vasey, Cat. Forest Trees, 30.
- P. ponderosa*, var. *scopulorum*, Engelmann in Bot. California, ii, 126.

* YELLOW PINE. BULL PINE.

Interior of British Columbia, south of latitude 51°, south and east along the mountain ranges of the Pacific region to Mexico, the Black hills of Dakota, Colorado, and western Texas; not detected in central or southern Nevada.

A large tree, 61 to 91 meters in height, with a trunk 3.60 to 4.57 meters in diameter, or throughout the Rocky Mountain region much smaller, rarely exceeding 30 meters in height (var. *scopulorum*); dry, rocky ridges and prairies, or in northern California rarely in cold, wet swamps, reaching its greatest development along the western slope of the sierras of northern and central California; in western Washington territory and Oregon rare and local; after *Pseudotsuga Douglasii* the most generally distributed and valuable timber tree of the Pacific forests, furnishing the principal lumber of eastern Washington territory and Oregon, western Montana, Idaho, the Black hills of Dakota, western Texas, New Mexico, and Arizona.

Wood, varying greatly in quality and value, heavy, hard, strong, brittle, not coarse-grained nor durable, compact; bands of small summer cells broad or narrow, very resinous, conspicuous, resin passages few, small; medullary rays numerous, obscure; color, light red, the very thick sap-wood almost white; specific gravity, 0.4715; ash, 0.35; largely manufactured into lumber, and used for railway ties, fuel, etc.

NOTE.—A form with purple cones and long glaucous foliage, approaching *P. Jeffreyi* in habit, is the prevailing tree of the valley of Flathead lake, Montana (*Cowby & Sargent*).

362.—*Pinus Jeffreyi*, Murray,

Rep. Oregon Exped. 2, t. 1; Edinburgh New Phil. Jour. new ser. xi, 224, t. 8, 9 (Trans. Bot. Soc. Edinburgh, vi, 350 & t.); Carrière, Trait. Conif. 368; 2 ed. 439.—Gordon, Pinetum, 198; 2 ed. 272.—Henkel & Hochstetter, Nadelhölz. 87.—Nelson, Pinaceæ, 115.—Hoopes, Evergreens, 115.—Parlatore in De Candolle, Prodr. xvi², 393.—Lawson, Pinetum Brit. i, 45, t. 6, f. 1-4.—Koch, Dendrologie, ii², 314.—Engelmann in Coulter's Bot. Gazette, vii, 4.—Veitch, Manual Conif. 165.

P. deflexa, Torrey in Bot. Mex. Boundary Survey, 209, t. 56, in part.—Cooper in Smithsonian Rep. 1860, 442.—Henkel & Hochstetter, Nadelhölz. 416.—Carrière, Trait. Conif. 2 ed. 455.—Bolander in Proc. California Acad. iii, 318.—Parlatore in De Candolle, Prodr. xvi², 431.—Fowler in London Gard. Chronicle, 1872, 1070.—Murray in London Gard. Chronicle, 1875, 106.—Gordon, Pinetum, 2 ed. 289.

P. ponderosa, var. *Jeffreyi*, Vasey, Cat. Forest Trees, 31.—Engelmann in Trans. St. Louis Acad. iv, 181; Bot. California, ii, 126.

BULL PINE. BLACK PINE.

California, Scott's mountain, Siskiyou county, south along the Sierra Nevada to the San Bernardino and San Jacinto mountains.

A large tree, 30 to 31 meters in height, with a trunk 1.20 to 4 meters in diameter; dry, gravelly slopes between 6,000 and 8,000 feet elevation; most common and reaching its greatest development on the eastern slope of the Sierra Nevada, here generally replacing the allied *P. ponderosa*, from which it may be distinguished by its more deeply-cleft bark, glaucous branchlets and leaves, much larger cones, and by the strong, pungent odor of oil of orange of the freshly-cut branchlets.

Wood light, strong, hard, rather coarse-grained, compact; bands of small summer cells not broad, very resinous, conspicuous, resin passages few, not large; medullary rays numerous, obscure; color, light red, the sap-wood pale yellow or nearly white; specific gravity, 0.5206; ash, 0.26; largely manufactured into coarse lumber.

Abietine, a volatile carbo-hydrogen possessing powerful anæsthetic properties, is probably obtained by distilling the resinous exudation of this species, and not of *P. Sabiniana* (*Watt's Dict. Chemistry*, 2d Suppl. 1.—*Am. Jour. Pharm.* 1872, 97.—*U. S. Dispensatory*, 14 ed. 900).

363.—*Pinus Chihuahuana*, Engelmann,

Wislizenus' Rep. No. 26; Wheeler's Rep. vi, 262; Trans. St. Louis Acad. iv, 181; Coulter's Bot. Gazette, vii, 4.—Lindley & Gordon in Jour. Hort. Soc. London, v, 220.—Carrière in Fl. des Serres, ix, 200; Rev. Hort. 1854, 227; Trait. Conif. 357; 2 ed. 455.—Gordon, Pinetum, 193; 2 ed. 266.—Torrey, Bot. Mex. Boundary Survey, 209.—Cooper in Smithsonian Rep. 1860, 442.—Henkel & Hochstetter, Nadelhölz. 86.—Hoopes, Evergreens, 143.—Parlatore in De Candolle, Prodr. xvi³, 397.—Vasey, Cat. Forest Trees, 32.

Santa Rita mountains, Arizona (*Rothrock, Engelmann & Sargent*), San Francisco mountains of southwestern New Mexico and Arizona (*Greene*); in Chihuahua.

A small tree, 18 to 24 meters in height, with a trunk 0.45 to 0.60 meter in diameter; dry, rocky ridges and slopes between 5,000 and 7,000 feet elevation; not common.

Wood light, soft, strong, brittle, close-grained, compact; bands of small summer cells not broad, resinous, conspicuous, resin passages few, rather large, conspicuous; medullary rays numerous, thin; color, clear light orange, the thick sap-wood lighter; specific gravity, 0.5457; ash, 0.39.

364.—*Pinus contorta*, Douglas;

London, Arboretum, iv, 2292, f. 2210, 2211.—Nuttall, Sylva, iii, 117; 2 ed. ii, 176.—Endlicher, Syn. Conif. 168.—Carrière, Trait. Conif. 164; 2 ed. 474.—Torrey in Pacific R. R. Rep. iv, 141.—Gordon, Pinetum, 165; 2 ed. 232.—Cooper in Smithsonian Rep. 1858, 261.—Lyll in Jour. Linnæan Soc. vii, 133, 141, in part.—Henkel & Hochstetter, Nadelhölz. 24.—Rothrock in Smithsonian Rep. 1867, 433.—Hoopes, Evergreens, 81, in part.—Parlatore in De Candolle, Prodr. xvi³, 381, in part.—Watson in King's Rep. v, 330.—Fowler in London Gard. Chronicle, 1872, 1070.—Gray in Proc. Am. Acad. vii, 402.—Koch, Dendrologie, ii³, 301.—Vasey, Cat. Forest Trees, 29.—Hall in Coulter's Bot. Gazette, ii, 91.—Macoun in Geological Rep. Canada, 1875-'76, 211.—Engelmann in Trans. St. Louis Acad. iv, 182; Bot. California, ii, 126; London Gard. Chronicle, 1883, 351.—G. M. Dawson in Canadian Nat. 2 ser. ix, 327, in part.—Veitch, Manual Conif. 145.—Masters in London Gard. Chronicle, 1883, 45, f. 5.

P. inops, Bongard in Mem. Acad. St. Petersburg, 6 ser. ii, 163 [not Aiton].—Hooker, Fl. Bor.-Am. ii, 161, in part.—Ledebour, Fl. Rossica, iii, 676 [not Aiton].

P. Boursieri, Carrière in Rev. Hort. 1854, 233 & f.; Fl. des Serres, ix, 200 & f.; Trait. Conif. 398; 2 ed. 475.

P. Banksiana, Lindley & Gordon in Jour. Hort. Soc. London, v, 218, in part.

P. muricata, Bolander in Proc. California Acad. iii, 227, 317 [not Don].

P. Bolanderi, Parlatore in De Candolle, Prodr. xvi³, 379.

SCRUB PINE.

Alaska, south along the coast to Mendocino county, California, extending inland to the western slopes of the Coast ranges.

A small, stunted tree, 6 to 9 meters in height, with a trunk 0.30 to 0.50 meter in diameter; sandy dunes and exposed rocky points.

Wood light, hard, strong, brittle, coarse-grained; bands of small summer cells very broad, resinous, conspicuous, resin passages numerous, not large; medullary rays numerous, obscure; color, light brown tinged with red, the thick sap-wood nearly white; specific gravity, 0.5815; ash, 0.19.

365.—*Pinus Murrayana*, Balfour,

Rep. Oregon Exped. 2, t. 3, f. 2.—Murray in Edinburgh New Phil. Jour. new ser. xi, 226 (Trans. Bot. Soc. Edinburgh, vi, 351).

P. inops, Bentham, Pl. Hartweg. 337 [not Aiton].

P. contorta, Newberry in Pacific R. R. Rep. vi, 34, 90, t. 5, f. 11 [not Douglas].—Engelmann in Am. Jour. Sci. 2. ser. xxiv, 332.—Lyll in Jour. Linnæan Soc. vii, 141, in part.—Cooper in Am. Nat. iii, 409.—Parlatore in De Candolle, Prodr. xvi³, 381, in part.—Porter in Hayden's Rep. 1871, 494.—Gray in Proc. Am. Acad. vii, 402.—Rothrock in Pl. Wheeler, 27, 50.—Parry in Am. Nat. vii, 179.

P. contorta, var. *latifolia*, Engelmann in King's Rep. v, 331; Porter & Coulter, Fl. Colorado; Hayden's Surv. Misc. Pub. No. 4, 129; Wheeler's Rep. vi, 262.—Brandege in Coulter's Bot. Gazette, iii, 32.—G. M. Dawson in Canadian Nat. new ser. ix, 328.

P. contorta, var. *Bolanderi*, Vasey, Cat. Forest Trees, 29.

TAMARACK. BLACK PINE. LODGE-POLE PINE. SPRUCE PINE.

Valley of the Yukon river, Alaska (Fort Selkirk, *Dall*), south through the interior of British Columbia, along the mountain ranges of Washington territory and Oregon and the Sierra Nevadas of California to mount San Jacinto; on the high plateau east of the Rocky mountains in about latitude 56°, and south through the mountains of Idaho, Montana, Wyoming, Colorado, and Utah to New Mexico and northern Arizona.

A tree 18 to 24 meters in height, with a trunk 0.60 to 1.20 meter in diameter; reaching its greatest development in the California Sierras; in the interior regions in dry, gravelly soil, here the prevailing tree, covering immense areas, and generally replacing other species destroyed by fire; western Washington territory and southward only along the borders of moist alpine meadows between 6,000 and 9,000 feet elevation; generally confounded with the closely-allied *P. contorta* of the coast, from which it may be distinguished by its longer, broader leaves, very thin, scaly bark, thin sap-wood, and less resinous and finer-grained wood, resembling that of the white pines; the distribution of the two species in northern British Columbia and Alaska still undetermined.

Wood light, soft, not strong, close, straight-grained, easily worked, compact, not durable; bands of small summer cell a narrow, not conspicuous, resin passages few, not large; medullary rays numerous, obscure; color, light yellow or nearly white, the thin sap-wood lighter; specific gravity, 0.4096; ash, 0.32; occasionally manufactured into lumber, and used for fuel, railway ties, etc.

366.—*Pinus Sabiniana*, Douglas,

Companion Bot. Mag. ii, 150.—Lambert, *Pinus*, 1 ed. iii, 137, t. 58.—London, *Arboretum*, iv, 2246, f. 2138-2143.—Forbes, *Pinetum Woburn*, 63, t. 23, 24.—Hooker, *Fl. Bor.-Am.* ii, 162.—Lindley in *Penn. Cycl.* xvii, 172.—Antoine, *Conif.* 30, t. 11.—Hooker & Arnott, *Bot. Beechey*, 393.—Link in *Linnaea*, xv, 509.—Nuttall, *Sylva*, iii, 110, t. 113; 2 ed. ii, 169, t. 113.—Spach, *Hist. Veg.* xi, 390.—De Chambray, *Trait. Arb. Res.* 347.—Endlicher, *Syn. Conif.* 159.—Knight, *Syn. Conif.* 30.—Lindley & Gordon in *Jour. Hort. Soc. London*, v, 216.—*Fl. des Serres*, ix, 275, t. 964.—Carrière, *Trait. Conif.* 334; 2 ed. 435.—Torrey & Gray in *Pacific R. R. Rep.* ii, 130.—Bigelow in *Pacific R. R. Rep.* iv, 25.—Torrey in *Pacific R. R. Rep.* iv, 141; *Bot. Mex. Boundary Survey*, 210; t. 57; *Ives' Rep.* 28.—Newberry in *Pacific R. R. Rep.* vi, 39, 90, f. 13.—Gordon, *Pinetum*, 208; 2 ed. 284.—Cooper in *Smithsonian Rep.* 1858, 261.—Walpers, *Ann.* v, 799.—Bolander in *Proc. California Acad.* iii, 226, 318.—Henkel & Hochstetter, *Nadelhölz.* 75.—Lawson, *Pinetum Brit.* i, 85, t. 11, t. 1-3.—Nelson, *Pinaceae*, 129.—Hoopes, *Evergreens*, 121.—Parlatore in *De Candolle, Prodr.* xvi^a, 391.—Fowler in *London Gard. Chronicle*, 1872, 1326.—Koch, *Dendrologie*, ii^a, 312.—Vasey, *Cat. Forest Trees*, 31.—Engelmann in *Wheeler's Rep.* vi, 375; *Trans. St. Louis Acad.* iv, 182; *Bot. California*, ii, 127.—Veitch, *Manual Conif.* 169.

DIGGER PINE. BULL PINE.

California, Portuguese Flat, Shasta county, south along the foot-hills of the Coast ranges and the western slope of the Sierra Nevadas below 4,000 feet elevation.

A large tree, 24 to 30 meters in height, with a trunk 0.60 to 1.20 meter in diameter; very common through all the foot-hills region.

Wood light, soft, not strong, brittle, very coarse-grained, compact, not durable; bands of small summer cells broad, very resinous, conspicuous, resin passages few, large, prominent; medullary rays numerous, obscure; color, light brown or red, the thick sap-wood yellow or nearly white; specific gravity, 0.4840; ash, 0.40; largely used for fuel.

The large edible nuts furnish the Indians an important article of food.

367.—*Pinus Coulteri*, D. Don,

Trans. Linnæan Soc. xvii, 440.—London, *Arboretum*, iv, 2250, f. 2144-2146.—Forbes, *Pinetum Woburn*, 67, t. 25, 26.—Antoine, *Conif.* 31, t. 12, 13.—*Penn. Cycl.* xvii, 172.—Link in *Linnaea*, xv, 510.—Hooker & Arnott, *Bot. Beechey*, 393.—Nuttall, *Sylva*, iii, 112; 2 ed. ii, 171.—Endlicher, *Syn. Conif.* 160.—Carrière in *Fl. des Serres*, ix, 275 & t.; *Trait. Conif.* 334; 2 ed. 435.—Cooper in *Smithsonian Rep.* 1858, 261.—Torrey in *Ives' Rep.* 28.—Henkel & Hochstetter, *Nadelhölz.* 76.—Bolander in *Proc. California Acad.* iii, 318.—Parlatore in *De Candolle, Prodr.* xvi, 392.—Vasey, *Cat. Forest Trees*, 31.—Gordon, *Pinetum*, 2 ed. 266.—Engelmann in *Trans. St. Louis Acad.* iv, 182; *Bot. California*, ii, 127.—Lawson, *Pinetum Brit.* i, 23, f. 1-5.

P. macrocarpa, Lindley in *Bot. Reg.* xxvi, Misc. 61.—Knight, *Syn. Conif.* 30.—Lindley & Gordon in *Jour. Hort. Soc. London*, v, 216.—Gordon, *Pinetum*, 201.—Nelson, *Pinaceae*, 117.—Hoopes, *Evergreens*, 115.—Veitch, *Manual Conif.* 166.

P. Sabiniana Coulteri, London, *Encycl. Pl.* 985, f. 1839-1841.

P. Sabiniana macrocarpa, Hort.

California, Monte Diablo, south through the Coast ranges to the Cuyamaca mountains, and probably in Lower California.

A tree 24 to 46 meters in height, with a trunk 0.90 to 1.80 meter in diameter; dry ridges and slopes between 3,000 and 6,000 feet elevation; most common and reaching its greatest development in the San Jacinto mountains.

Wood light, soft, not strong, brittle, coarse-grained; bands of small summer cells broad, very resinous, conspicuous, resin passages few, large; medullary rays numerous, prominent; color, light red, the thick sap-wood nearly white; specific gravity, 0.4133; ash, 0.37.

368.—*Pinus insignis*, Douglas;

London, Arboretum, iv, 2243, f. 2132-2137.—Forbes, Pinetum Woburn, 51, t. 18.—Lindley in Penn. Cycl. xvii, 171.—Antoine, Conif. 27, t. 8, f. 1.—Hooker & Arnott, Bot. Beechey, 393.—Spach, Hist. Veg. xi, 389.—Nuttall, Sylva, iii, 115; 2 ed. ii, 174.—Bentham, Bot. Sulphur, 55.—Endlicher, Syn. Conif. 163.—Knight, Syn. Conif. 30.—Lindley & Gordon in Jour. Hort. Soc. London, v, 217.—Carrière, Trait. Conif. 339; 2 ed. 440.—Bigelow in Pacific R. R. Rep. iv, 25.—Torrey in Pacific R. R. Rep. iv, 141; Bot. Mex. Boundary Survey, 209, t. 55; Ives' Rep. 23.—Newberry in Pacific R. R. Rep. vi, 90.—Gordon, Pinetum, 197; 2 ed. 270.—Cooper in Smithsonian Rep. 1858, 261.—Murray in Edinburgh New Phil. Jour. new ser. xi, 222 (Trans. Bot. Soc. Edinburgh, vi, 347).—Henkel & Hochstetter, Nadelhölz. 69.—Bolander in Proc. California Acad. iii, 262, t. 317.—Nelson, Pinaceæ, 114.—Hoopes, Evergreens, 143.—Parlatore in De Candolle, Prodr. xvi², 395.—Lawson, Pinetum Brit. i, 37 t. 1, 5, f. 1-14.—Fowler in London Gard. Chronicle, 1872, 1070.—Vasey, Cat. Forest Trees, 31.—Engelmann in Trans. St. Louis Acad. iv, 182; Bot. California, ii, 128.—Veitch, Manual Conif. 163, f. 39.

?*P. Californica*, Loiseleur in Nouveau Duhamel, v, 243.—London, Arboretum, iv, 2268.—Endlicher, Syn. Conif. 162.—Hooker & Arnott, Bot. Beechey, 393.—Nuttall, Sylva, iii, 117; 2 ed. ii, 175.—Carrière, Trait. Conif. 1 ed. 253.

P. adunca, Bosc in Poiret, Suppl. iv, 418.

P. Sinclairii, Hooker & Arnott, Bot. Beechey, 392, 393, t. 93, in part.—Nuttall, Sylva, iii, 141; 2 ed. ii, 198.—Carrière, Trait. Conif. 2 ed. ii, 198.

P. radiata, D. Don in Trans. Linnæan Soc. xvii, 442; Lambert, Pinus, 1 ed. iii, 133, t. 86.—London, Arboretum, iv, 2270, f. 2182.—Antoine, Conif. 33, t. 14, f. 3.—Hooker & Arnott, Bot. Beechey, 392, 393, in part.—Nuttall, Sylva, iii, 116; 2 ed. ii, 175.—Endlicher, Syn. Conif. 161.—Hartweg in Jour. Hort. Soc. London, iii, 226.—Gordon in Jour. Hort. Soc. London, iv, 214 & f. (Fl. des Serres, vi, 434 & t.); Pinetum, 206; 2 ed. 282.—Knight, Syn. Conif. 37.—Lindley & Gordon in Jour. Hort. Soc. London, v, 216.—Carrière, Trait. Conif. 1 ed. 337.—Nelson, Pinaceæ, 127.—Hoopes, Evergreens, 118.—Koch, Dendrologie, ii², 307.—Vasey, Cat. Forest Trees, 31.

P. tuberculata, D. Don in Trans. Linnæan Soc. xvii, 441 [not Gordon].—Lambert, Pinus, 1 ed. iii, 131, t. 85.—London, Arboretum, iv, 2270, f. 2181.—Antoine, Conif. 33, t. 14, f. 2.—Hooker & Arnott, Bot. Beechey, 394.—Endlicher, syn. Conif. 162.—Carrière, Trait. Conif. 338; 2 ed. 441, in part.—Nelson, Pinaceæ, 137.—Hoopes, Evergreens, 123 (excl. syn. *Californica*).—Parlatore in De Candolle, Prodr. xvi², 394, in part.

P. rigida,? Hooker & Arnott, Bot. Beechey, 160 [not Miller].

P. insignis macrocarpa, Hartweg in Jour. Hort. Soc. London, iii, 226.—Carrière, Trait. Conif. 440.

MONTEREY PINE.

California, Pescadero to Monterey and San Simeon bay.

A tree 24 to 30 meters in height, with a trunk 0.60 to 0.90 meter in diameter; sandy soil, in immediate proximity to the sea-coast; rare and local; now widely cultivated on the Pacific coast for shelter and ornament. A form of Guadalupe island, off the coast of Lower California, with leaves in pairs, is var. *binata* (Engelmann in Proc. Am. Acad. xi, 119; Bot. California, ii, 128).

Wood light, soft, not strong, brittle, close-grained, compact; bands of small summer cells not broad, resinous, conspicuous; color, light brown, the very thick sap-wood nearly white; specific gravity, 0.4574; ash, 0.30; locally somewhat used for fuel.

369.—*Pinus tuberculata*, Gordon,

Jour. Hort. Soc. London, iv, 218 & f. (Fl. des Serres, v, 517^c & f.); Pinetum, 211; 2 ed. 288 [not Don].—Rep. Oregon Exped. 2, t. 2, f. 2.—Henkel & Hochstetter, Nadelhölz. 78, in part.—Bolander in Proc. California Acad. iii, 262, 317.—Lawson, Pinetum Brit. i, 93, t. 13, f. 1-9.—Carrière, Trait. Conif. 2 ed. 441, in part.—Parlatore in De Candolle, Prodr. xvi², 394 (excl. bib.).—Koch, Dendrologie, ii², 309.—Vasey, Cat. Forest Trees, 31.—Engelmann in Trans. St. Louis Acad. iv, 183; Bot. California, ii, 128.—Veitch, Manual Conif. 170.

P. Californica, Hartweg in Jour. Hort. Soc. London, ii, 189 [not Loiseleur].

KNOB-CONE PINE.

Valley of the Mackenzie river, Oregon, south along the western slope of the Cascade and Sierra Nevada mountains, and in the California Coast ranges from the Santa Cruz to the San Jacinto mountains.

A tree 18 to 22 meters in height, with a trunk 0.60 to 0.90 meter in diameter, or, rarely, reduced to a low shrub, fruiting when not more than 1 meter in height; dry, gravelly ridges and slopes from 2,500 (San Bernardino mountains) to 5,500 (mount Shasta) feet elevation; not common.

Wood light, soft, not strong, brittle, coarse-grained, compact; bands of small summer cells very broad, not conspicuous, resin passages numerous, large, prominent; medullary rays numerous, thin; color, light brown, the thick sap-wood nearly white or slightly tinged with red; specific gravity, 0.3499; ash, 0.33.

370.—*Pinus Tæda*, Linnæus,

Spec. 1 ed. 1000, in part.—Du Roi, Harbk. ii, 63.—Wangenheim, Amer. 41.—Aiton, Hort. Kew. iii, 368; 2 ed. v, 317.—Mönch, Meth. 365.—Michaux, Fl. Bor.-Am. ii, 205.—Lambert, Pinus, 1 ed. i, 23, t. 16, 17; 2 ed. i, 26, t. 17, 18; 3 ed. i, 30, t. 15.—Willdenow, Spec. iv, 498; Berl. Baumz. 269.—Persoon, Syn. ii, 578.—Desfontaines, Hist. Arb. ii, 612.—Michaux f. Hist. Arb. Am. i, 98, t. 9; N. American Sylva, 3 ed. iii, 123, t. 143.—Nouveau Duhamel, v, 245, t. 75, f. 2.—Smith in Rees' Cycl. xxviii, No. 13.—Pursh, Fl. Am. Sept. ii, 644.—Nuttall, Genera, ii, 223.—Hayne, Dend. Fl. 175.—Elliott, Sk. ii, 636.—Sprengel, Syst. ii, 887.—Eaton, Manual, 6 ed. 265.—Lawson, Ag. Manual, 351; Pinetum Brit. i, 89, t. 12.—London, Arboretum, iv, 2237, f. 2118-2122.—Forbes, Pinetum Woburn. 43, t. 14.—Antoine, Conif. 25, t. 7, f. 1.—Eaton & Wright, Bot. 359.—Link in Linnæa, xv, 503.—Spach, Hist. Veg. xi, 391.—Griffith, Med. Bot. 609.—Gihoul, Arb. Resin, 32.—Endlicher, Syn. Conif. 164.—Scheele in Roemer, Texas, Appx. 447.—Knight, Syn. Conif. 30.—Lindley & Gordon in Jour. Hort. Soc. London, v, 217.—Carrière, Trait. Conif. 344; 2 ed. 448.—Darby, Bot. S. States, 515.—Gordon, Pinetum, 210; 2 ed. 286.—Cooper in Smithsonian Rep. 1858, 257.—Chapman, Fl. S. States, 433.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 22.—Lesquereux in Owen's 2d Rep. Arkansas, 389.—Wood, Cl. Book, 660; Bot. & Fl. 313.—Porcher, Resources S. Forests, 506.—Henkel & Hochstetter, Nadelhölz. 65.—Nelson, Pinaceæ, 136.—Gray, Manual N. States, 5 ed. 469; Hall's Pl. Texas, 21.—Hoopes, Evergreens, 122.—Parlatore in De Candolle, Prodr. xvi², 393.—Young, Bot. Texas, 516.—Koch, Dendrologie, ii², 304.—Vasey, Cat. Forest Trees, 31.—Bentley & Trimen, Med. Pl. iv, 259, t. 259.—Engelmann in Trans. St. Louis Acad. iv, 183.—Veitch, Manual Conif. 172.

P. Tæda, var. *tenuifolia*, Aiton, Hort. Kew. iii, 368.

LOBLOLLY PINE. OLD-FIELD PINE. ROSEMARY PINE.

Southern Delaware, south to cape Malabar and Tampa bay, Florida, generally near the coast, through the Gulf states to the valley of the Colorado river, Texas, and north through southern Arkansas to the valley of the Arkansas river.

A tree 24 to 46 meters in height, with a trunk 0.90 to 1.50 meter in diameter; low, wet clay or dry sandy soil; springing up on all abandoned lands from Virginia southward, and now often replacing in the southern pine belt the original forests of *Pinus palustris*; in eastern North Carolina rarely on low, rich swamp ridges, here known as rosemary pine and attaining its greatest development and value.

Wood light, not strong, brittle, very coarse-grained, not durable; bands of small summer cells broad, very resinous, conspicuous, resin passages few, not prominent; medullary rays numerous, obscure; color, light brown, the very thick sap-wood orange, or often nearly white; wood of the rosemary pine close-grained, less resinous, lighter, with much thinner sap; specific gravity, 0.5441; ash, 0.26; largely used for fuel and manufactured into lumber of inferior quality.

Turpentine is occasionally manufactured from this species (*U. S. Dispensatory*, 14 ed. 901.—*Flückiger & Hanbury, Pharmacographia*, 545).

371.—*Pinus rigida*, Miller,

Diet. 7 ed. No. 10.—Du Roi, Harbk. ii, 60.—Marshall, Arbustum, 101.—Wangenheim, Amer. 41.—Lambert, Pinus, 1 ed. i, 25, t. 18, 19; 2 ed. i, 28; t. 18, 19; 3 ed. i, 32, t. 16, 17.—Willdenow, Spec. iv, 498; Enum. 988; Berl. Baumz. 268.—Persoon, Syn. ii, 578.—Desfontaines, Hist. Arb. ii, 612.—Michaux f. Hist. Arb. Am. i, 89, t. 8; N. American Sylva, 3 ed. iii, 118, t. 144.—Nouveau Duhamel, v, 244, t. 74.—Aiton, Hort. Kew. 2 ed. v, 317.—Smith in Rees' Cycl. xxviii, No. 14.—Pursh, Fl. Am. Sept. ii, 643.—Poiret, Suppl. iv, 417.—Eaton, Manual, 110; 6 ed. 265.—Barton, Compend. Fl. Philadelph. ii, 183.—Nuttall, Genera, ii, 223.—Hayne, Dend. Fl. 175.—Elliott, Sk. ii, 635.—Sprengel, Syst. ii, 887.—Torrey, Compend. Fl. N. States, 360; Fl. N. York, ii, 227.—Beck, Bot. 339.—London, Arboretum, iv, 2239, f. 2123-2126.—Forbes, Pinetum Woburn. 41, t. 13.—Eaton & Wright, Bot. 358.—Antoine, Conif. 26, t. 7, f. 2.—Bigelow, Fl. Boston. 3 ed. 385.—Lindley in Penn. Cycl. xvii, 172.—Link in Linnæa, xv, 503.—Spach, Hist. Veg. xi, 388.—Griffith, Med. Bot. 604.—Gihoul, Arb. Resin, 31.—Endlicher, Syn. Conif. 164.—Knight, Syn. Conif. 30.—Lindley & Gordon in Jour. Hort. Soc. London, v, 217.—Carrière, Trait. Conif. 342; 2 ed. 447.—Darlington, Fl. Cestrica, 3 ed. 290.—Darby, Bot. S. States, 514.—Gordon, Pinetum, 207; 2 ed. 283.—Cooper in Smithsonian Rep. 1858, 257.—Chapman, Fl. S. States, 433.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 21.—Wood, Cl. Book, 660; Bot. & Fl. 313.—Henkel & Hochstetter, Nadelhölz, 67.—Nelson, Pinaceæ, 128.—Gray, Manual N. States, 5 ed. 469.—Hoopes, Evergreens, 119.—Parlatore in De Candolle, Prodr. xvi², 394.—Koch, Dendrologie, ii², 307.—Vasey, Cat. Forest Trees, 31.—Engelmann in Trans. St. Louis Acad. iv, 183.—Sears in Bull. Essex Inst. xiii, 186.—Veitch, Manual Conif. 169.

P. Tæda, var. *rigida*, Aiton, Hort. Kew. iii, 368.

P. Tæda, var. *a.* Poiret in Lamarck, Dict. v, 340.

P. Fraseri, Loddiges, Cat. ed. 1836, 50 [not Pursh].

P. Loddigesii, London, Arboretum, iv, 2269.

PITCH PINE.

Valley of the Saint John's river, New Brunswick, to the northern shores of lake Ontario, south through the Atlantic states to northern Georgia, extending to the western slope of the Alleghany mountains in West Virginia and Kentucky (Pineville, Bell county, *De Friesse*).

A tree 12 to 24 meters in height, with a trunk 0.60 to 0.90 meter in diameter; dry, sandy, barren soil, or less commonly in deep, cold swamps; very common.

Wood light, soft, not strong, brittle, coarse-grained, compact; bands of small summer cells broad, very resinous, conspicuous, resin passages numerous, not large; medullary rays numerous, obscure; color, light brown or red, the thick sap-wood yellow or often nearly white; specific gravity, 0.5151; ash, 0.23; largely used for fuel, charcoal, and occasionally manufactured into coarse lumber.

NOTE.—Upon the island of Nantucket, Massachusetts, this species is now greatly injured by the attacks of the destructive caterpillar of the pine moth (*Retina frustrana*, Souder in *Pub. Massachusetts Ag. Soc.* 1883 & t).

372.—*Pinus serotina*, Michaux,

Fl. Bor.-Am. ii, 205.—Willdenow, Spec. iv, 499.—Persoon, Syn. ii, 578.—Michaux f. Hist. Arb. Am. i, 86, t. 7; N. American Sylva, 3 ed. iii, 117, t. 142.—Nouveau Duhamel, v, 246, t. 75, f. 1.—Pursh, Fl. Am. Sept. ii, 643.—Poiret, Suppl. iv, 417.—Nuttall, Genera, ii, 223.—Lambert, Pinus, 1 ed. iii, 35, t. 12.—Elliott, Sk. ii, 634.—Sprengel, Syst. ii, 887.—Torrey, Compend. Fl. N. States, 360.—Beck, Bot. 339.—Eaton, Manual, 6 ed. 265.—London, Arboretum, iv, 2242, f. 2127–2131.—Forbes, Pinetum Woburn. 47, t. 16.—Eaton & Wright, Bot. 359.—Antoine, Conif. 27, t. 8, f. 2.—Lindley in Penn. Cycl. xvii, 172.—Link in Linnæa, xv, 504.—Spach, Hist. Veg. xi, 389.—Giboul, Arb. Resin. 32.—Endlicher, Syn. Conif. 163.—Knight, Syn. Conif. 30.—Lindley & Gordon in Jour. Hort. Soc. London, v, 217.—Carrière, Trait. Conif. 341; 2 ed. 449.—Darby, Bot. S. States, 514.—Gordon, Pinetum, 209; 2 ed. 285.—Chapman, Fl. S. States, 433.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 21.—Henkel & Hochstetter, Nadelhölz. 70.—Nelson, Pinaceæ, 129.—Parlatore in De Candolle, Prodr. xvi², 394.—Koch, Dendrologie, ii², 305.—Vasey, Cat. Forest Trees, 31.

P. Tæda, var. *alopecuroides*, Aiton, Hort. Kew. 2 ed. v, 317.—London, Arboretum, iv, 2237.

P. rigida, var. *serotina*, London, Encycl. Pl. 979, f. 1824–1827.—Cooper in Smithsonian Rep. 1858, 257.—Hoopes, Evergreens, 120.—Engelmann in Trans. St. Louis Acad. iv, 183.

POND PINE.

North Carolina, south near the coast to the head of the Saint John's river, Florida.

A tree 12 to 24 meters in height, with a trunk 0.60 to 0.90 meter in diameter; inundated borders of streams and ponds in low, peaty soil; not common.

Wood heavy, soft, not strong, brittle, coarse-grained, compact; bands of small summer cells broad, forming fully one-half the annual growth, very resinous, dark colored, conspicuous, resin passages few, large; medullary rays numerous, obscure; color, dark orange, the thick sap-wood pale yellow; specific gravity 0.7942; ash, 0.17.

373.—*Pinus inops*, Aiton,

Hort. Kew. iii, 367; 2 ed. v, 316.—Michaux, Fl. Bor.-Am. ii, 204.—Lambert, Pinus, 1 ed. i, 18, t. 13; 2 ed. i, 21, t. 14; 3 ed. i, 25, t. 12.—Willdenow, Spec. iv, 496; Enum. 988; Berl. Baumz. 266.—Persoon, Syn. ii, 578.—Michaux f. Hist. Arb. Am. i, 58, t. 4; N. American Sylva, 3 ed. iii, 103, t. 139.—Nouveau Duhamel, v, 236, t. 69, f. 1.—Pursh, Fl. Am. Sept. ii, 641.—Smith in Rees' Cycl. xxviii, No. 10.—Barton, Prodr. Fl. Philadelph. 93.—Compend. Fl. Philadelph. ii, 183.—Nuttall, Genera, ii, 223.—Hayne, Dend. Fl. 173.—Elliott, Sk. ii, 633.—Sprengel, Syst. ii, 886.—Torrey, Compend. Fl. N. States, 359.—Audubon, Birds, t. 97.—Beck, Bot. 338.—Eaton, Manual, 6 ed. 265.—Bon Jard. 1837, 976.—London, Arboretum, iv, 2192, f. 2068–2071.—Forbes, Pinetum Woburn. 15, t. 4.—Hooker, Fl. Bor.-Am. ii, 161, in part.—Eaton & Wright, Bot. 358.—Antoine, Conif. 17, t. 5, f. 3.—Lindley in Penn. Cycl. xvii, 171.—Link in Linnæa, xv, 500.—Spach, Hist. Veg. xi, 386.—Endlicher, Syn. Conif. 167.—Knight, Syn. Conif. 26.—Lindley & Gordon in Jour. Hort. Soc. London, v, 217.—Carrière, Trait. Conif. 361; 2 ed. 471.—Darlington, Fl. Cestrica, 3 ed. 290.—Darby, Bot. S. States, 514.—Gordon, Pinetum, 167; 2 ed. 238.—Cooper in Smithsonian Rep. 1858, 257.—Chapman, Fl. S. States, 433.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 20.—Wood, Cl. Book, 661; Bot. & Fl. 313.—Henkel & Hochstetter, Nadelhölz. 22.—Nelson, Pinaceæ, 113.—Gray, Manual N. States, 5 ed. 470.—Hoopes, Evergreens, 84.—Parlatore in De Candolle, Prodr. xvi², 380 (excl. syn. *variabilis*).—Vasey, Cat. Forest Trees, 30.—Veitch, Manual Conif. 158.

P. Virginiana, Miller, Gard. Diet. 7 ed. No. 9.—Du Roi, Obs. Bot. 43; Harbk. 2 ed. ii, 35.—Marshall, Arbustum, 102.—Wangenheim, Amer. 74.—Koch, Dendrologie, ii², 299.

P. Tæda, var. *Virginiana*, Poiret in Lamarck, Diet. v, 340.

JERSEY PINE. SCRUB PINE.

Middle Island, Long island, Tottenville, and Clifton, Staten island, New York, south, generally near the coast, to the valley of the Savannah river (Aiken, South Carolina), and through eastern and middle Kentucky to "the knobs" of southeastern Indiana.

A tree 24 to 36 meters in height, with a trunk 0.60 to 0.90 meter in diameter, or in the Atlantic states generally much smaller; sandy, generally barren soil, reaching its greatest development west of the Alleghany mountains.

Wood light, soft, not strong, brittle, very close-grained, compact, durable; bands of small summer cells broad, very resinous, conspicuous, resin passages few, not prominent; medullary rays numerous, thin; color, light orange, the thick sap-wood nearly white; specific gravity, 0.5309; ash, 0.30; largely used for fuel, and in Kentucky and Indiana preferred for and largely manufactured into water-pipes and pump-logs.

374.—*Pinus clausa*, Vasey.

Cat. Forest Trees, 30.

P. inops, var. *clausa*, Engelmann in Trans. St. Louis Acad. iv, 183.—Chapman, Fl. S. States, Suppl. 650.

SAND PINE. SCRUB PINE. SPRUCE PINE.

Florida, shores of Pensacola bay, south, generally within 30 miles of the coast, to Pease creek, and occupying a narrow ridge along the east coast south of Saint Augustine.

A tree 21 to 24 meters in height, with a trunk 0.60 to 0.75 meter in diameter, or on the west coast rarely 6 to 9 meters in height; barren, sandy dunes and ridges; most common and reaching its greatest development about the head of Halifax bay.

Wood light, soft, not strong, brittle; bands of small summer cells broad, very resinous, conspicuous, resin passages numerous, prominent; medullary rays numerous, thin; color, light orange or yellow, the thick sap-wood nearly white; specific gravity, 0.5576; ash, 0.31; occasionally used for the masts of small vessels.

375.—*Pinus pungens*, Michaux f.

Hist. Arb. Am. i, 61, t. 5; N. American Sylva, 3 ed. iii, 105, t. 140.—Nouveau Duhamel, v. 236, t. 67, f. 4.—Aiton, Hort. Kew. 2 ed. v, 314.—Pursh, Fl. Am. Sept. ii, 643.—Poiret, Suppl. iv, 417.—Elliott, Sk. ii, 635.—Sprengel, Syst. ii, 886.—Eaton, Manual, 6 ed. 265.—Lambert, Pinus, 1 ed. iii, 34, t. 17.—London, Arboretum, iv, 2197, f. 2077-2080.—Forbes, Pinetum Woburn. 17, t. 5.—Eaton & Wright, Bot. 359.—Antoine, Conif. 18, t. 5, f. 4.—Lindley in Penn. Cycl. xvii, 171.—Nuttall, Sylva, iii, 125; 2 ed. ii, 184.—Spach, Hist. Veg. xi, 287.—Endlicher, Syn. Conif. 166.—Knight, Syn. Conif. 27.—Lindley & Gordon in Jour. Hort. Soc. London, v, 217.—Carrière, Trait. Conif. 359; 2 ed. 470.—Darby, Bot. S. States, 515.—Gordon, Pinetum, 181; 2 ed. 254.—Cooper in Smithsonian Rep. 1858, 257.—Chapman, Fl. S. States, 432.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 20.—Wood, Cl. Book, 660; Bot. & Fl. 313.—Henkel & Hochstetter, Nadelhölz, 21.—Nelson, Pinaceæ, 127.—Gray, Manual N. States, 5 ed. 469.—Hoopees, Evergreens, 98.—Parlatore in De Candolle, Prodr. xvi³, 379.—Koch, Dendrologie ii³, 304.—Vasey, Cat. Forest Trees, 30.—Meehan in Rep. Penn. Fruit Growers' Soc. 1877 & t.—Engelmann in Trans. St. Louis Acad. iv, 183.—Veitch, Manual Conif. 158.

TABLE-MOUNTAIN PINE. HICKORY PINE.

Alleghany mountains, Pennsylvania to Tennessee.

A tree 9 to 18 meters in height, with a trunk 0.60 to 1.05 meter in diameter; most common and reaching its greatest development upon the high mountains of East Tennessee, here often the prevailing species and forming extensive forests.

Wood light, soft, not strong, brittle, coarse-grained, compact; bands of small summer cells broad, resinous, conspicuous, resin passages numerous, large; medullary rays numerous, prominent; color, light brown, the thick sap-wood nearly white; specific gravity, 0.4935; ash, 0.27; in Pennsylvania largely manufactured into charcoal.

376.—*Pinus muricata*, D. Don,

Trans. Linnæan Soc. xvii, 441.—Lambert, Pinus, 1 ed. iii, t. 84.—London, Arboretum, iv, 2269, f. 2180.—Hooker & Arnott, Bot. Beechey, 393.—Antoine, Conif. 32, t. 14, f. 1.—Nuttall, Sylva, iii, 113; 2 ed. ii, 172.—Endlicher, Syn. Conif. 161.—Knight, Syn. Conif. 26.—Gordon in Jour. Hort. Soc. London, iv, 216 & f. (Fl. des Serres, v, 517^b & f.); Pinetum, 173; 2 ed. 246 (excl. syn. *Murrayana*).—Lindley & Gordon in Jour. Hort. Soc. London, v, 217.—Carrière, Trait. Conif. 359; 2 ed. 470.—Torrey, Bot. Mex. Boundary Survey, 209, t. 54 (*P. Edgariana* on plate).—Cooper in Smithsonian Rep. 1858, 261.—Henkel & Hochstetter, Nadelhölz, 60.—Nelson, Pinaceæ, 121.—Hoopees, Evergreens, 92.—Parlatore in De Candolle, Prodr. xvi³, 379.—Fowler in London Gard. Chronicle, 1872, 1164.—Koch, Dendrologie, ii³, 302.—Vasey, Cat. Forest Trees, 30.—Engelmann in Trans. St. Louis Acad. iv, 183; Bot. California, ii, 128.—Veitch, Manual Conif. 151.—London Gard. Chronicle, 1884, 49, f. 7-9.

P. inops, var. Bentham, Pl. Hartweg. 337.

P. Edgariana, Hartweg in Jour. Hort. Soc. London, iii, 217, 226.

P. contorta, Bolander in Proc. California Acad. iii, 227, 317 [not Douglas].

OBISPO PINE. BISHOP'S PINE.

California, Mendocino county south through the Coast ranges to San Luis Obispo county.

A tree 24 to 36 meters in height, with a trunk 0.30 to 0.90 meter in diameter, or more often not exceeding 15 meters in height; cold peat bogs or barren, sandy gravel; always exposed to the winds and fogs of the ocean, and not found above 2,000 feet elevation, reaching its greatest development in Mendocino county; rare and local.

Wood light, very strong and hard, rather coarse-grained, compact; bands of small summer cells broad, resinous, resin passages few, not prominent; medullary rays numerous, thin; color, light brown, the thick sap-wood nearly white; specific gravity, 0.4942; ash, 0.26.

377.—*Pinus mitis*, Michaux,

Fl. Bor.-Am. ii, 204.—Michaux f. Hist. Arb. Am. i, 52, t. 3; N. American Sylva, 3 ed. iii, 96, t. 137.—Barton, Prodr. Fl. Philadelph. 93.—Poiret, Suppl. iv, 417.—London, Arboretum, iv, 2195, f. 2072-2076.—Antoine, Conif. 16, t. 5, f. 1.—Lindley in Penn. Cycl. xvii, 171.—Spach, Hist. Veg. xi, 386.—Torrey, Fl. N. York, ii, 229.—Endlicher, Syn. Conif. 167.—Knight, Syn. Conif. 26.—Lindley & Gordon in Jour. Hort. Soc. London, v, 217.—Carrière, Trait. Conif. 361; 2 ed. 472.—Gordon, Pinetum, 170; 2 ed. 243 (excl. syn. Boylei).—Cooper in Smithsonian Rep. 1858, 275.—Chapman, Fl. S. States, 433.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 19.—Lesquereux in Owen's 2d Rep. Arkansas, 389.—Wood, Cl. Book, 660; Bot. & Fl. 313.—Henkel & Hochstetter, Nadelholz. 23.—Gray, Manual N. States, 5 ed. 470.—Hoopes, Evergreens, 88.—Parlatore in De Candolle, Prodr. xvi³, 380.—Young, Bot. Texas, 516.—Koch, Dendrologie, ii³, 300.—Vasey, Cat. Forest Trees, 30.—Broadhead in Coulter's Bot. Gazette, iii, 60.—Engelmann in Trans. St. Louis Acad. iv, 184.—Ridgway in Proc. U. S. Nat. Mus. 88.

P. echinata, Miller, Dict. 7 ed. No. 12.—Marshall, Arbustum, 180†.—Wangenheim, Amer. 74.

P. Virginiana, var. *echinata*, Du Roi, Harbk. ii, 38.

P. Teda, var. *variabilis*, Aiton, Hort. Kew. iii, 368.

P. variabilis, Lambert, Pinus, 1 ed. i, 22, t. 15; 2 ed. i, 25, t. 16; 3 ed. i, 29, t. 14.—Willdenow, Spec. iv, 498.—Persoon, Syn. ii, 578.—Nouveau Duhamel, v, 235, t. 69, f. 2.—Aiton, Hort. Kew. 2 ed. v, 316.—Pursh, Fl. Am. Sept. ii, 643.—Smith in Rees' Cycl. xxviii, No. 12.—Barton, Compend. Fl. Philadelph. ii, 183.—Nuttall, Genera, ii, 223.—Elliott, Sk. ii, 633.—Sprengel, Syst. ii, 886.—Torrey, Compend. Fl. N. States, 360.—Beck, Bot. 339.—Eaton, Manual, 6 ed. 265.—Forbes, Pinetum Woburn. 35, t. 11.—Eaton & Wright, Bot. 358.—Antoine, Conif. 15, t. 5, f. 2.—Link in Linnæa, xv, 502.—Endlicher, Syn. Conif. 168 (excl. syn.).—Darby, Bot. S. States, 514.

P. rigida, Porcher, Resources S. States, 504 [not Miller].

YELLOW PINE. SHORT-LEAVED PINE. SPRUCE PINE. BULL PINE.

Staten island, New York, south to the Chattahoochee region of western Florida, through the Gulf states to Tennessee and eastern Texas, and through Arkansas to the Indian territory, southeastern Kansas, southern Missouri, and in Union county, Illinois.

A tree 24 to 30 meters in height, with a trunk 0.60 to 1.35 meter in diameter; light sandy soil or, less commonly, along the low borders of swamps; forming west of the Mississippi river, mixed with oaks and other deciduous trees, extensive forests; the only species of northern Arkansas, Kansas, and Missouri, reaching its greatest development in western Louisiana, southern Arkansas, and eastern Texas.

Wood, varying greatly in quality and amount of sap, heavy, hard, strong, generally coarse-grained, compact; bands of small summer cells broad, often occupying half the width of the annual growth; very resinous, resin passages numerous, large; medullary rays numerous, conspicuous; color, orange, the sap-wood nearly white; specific gravity, 0.6104; ash, 0.29; largely manufactured into lumber, especially in the states west of the Mississippi river, and among yellow pines only inferior in value to that of *P. palustris*.

378.—*Pinus glabra*, Walter,

Fl. Caroliniana, 237.—Poiret in Lamarck, Dict. v, 342.—Ravenel in Proc. Elliott Soc. i, 52.—Chapman, Fl. S. States, 433.—Porcher, Resources S. Forests, 506.—Hoopes, Evergreens, 82.—Vasey, Cat. Forest Trees, 30.—Engelmann in Trans. St. Louis Acad. iv, 184.

?*P. mitis*, var. *paupera*, Wood, Cl. Book, 660.

CEDAR PINE. SPRUCE PINE. WHITE PINE.

South Carolina, south to the Ohattahoochee region of western Florida, generally near the coast, and through the Gulf states south of latitude 32° 30' to the valley of the Pearl river, Louisiana.

A tree 24 to 30 meters in height, with a trunk 0.60 to 1.20 meter in diameter; rich bottom lands and hummocks in dense forests of hard-wood trees, reaching its greatest development in Alabama and Mississippi; not common and local.

Wood light, soft, not strong, brittle, very coarse-grained, not durable; bands of small summer cells broad, not resinous, resin passages few, not large; medullary rays numerous, obscure; color, light brown, the sap-wood nearly white; specific gravity, 0.3931; ash, 0.45.

379.—*Pinus Banksiana*, Lambert,

Pinus, 1 ed. i, 7, t. 3; 2 ed. i, 7, t. 3; 3 ed. i, 9, t. 3.—Persoon, Syn. ii, 578.—Desfontaines, Hist. Arb. ii, 611.—Nouveau Duhamel, v, 234, t. 67, f. 3.—Aiton, Hort. Kew. 2 ed. v, 315.—Pursh, Fl. Am. Sept. ii, 642.—Smith in Rees' Cycl. xxviii, No. 4.—Nuttall, Genera, ii, 223; Sylva, iii, 124; 2 ed. ii, 182.—Sprengel, Syst. ii, 886.—Torrey, Compend. Fl. N. States, 360.—Beck, Bot. 339.—Eaton, Manual, 6 ed. 265.—London, Arboretum, iv, 2190, f. 2064–2067.—Forbes, Pinetum Woburn. 13, t. 3.—Hooker, Fl. Bor.-Am. ii, 161.—Eaton & Wright, Bot. 358.—Antoine, Conif. 8, t. 4, f. 2.—Lindley in Penn. Cycl. xvii, 171.—Link in Linnæa, xv, 491.—Spach, Hist. Veg. xi, 379.—Endlicher, Syn. Conif. 177.—Knight, Syn. Conif. 26.—Lindley & Gordon in Jour. Hort. Soc. London, v, 218 (excl. syn. *contorta*).—Parry in Owen's Rep. 618.—Carrière, Trait. Conif. 381; 2 ed. 485.—Gordon, Pinetum, 163; 2 ed. 230.—Richardson, Arctic Exped. 441.—Cooper in Smithsonian Rep. 1858, 257.—Hooker f. in Trans. Linnæan Soc. xxiii, 301.—Wood, Cl. Book, 661.—Henkel & Hochstetter, Nadelhölz. 44.—Nelson, Pinaceæ, 104.—Gray, Manual N. States, 5 ed. 470.—Hoopes, Evergreens, 78.—Vasey, Cat. Forest Trees, 29.—Macoun in Geological Rep. Canada, 1875–76, 211.—Engelmann in Trans. St. Louis Acad. iv, 184.—Sears in Bull. Essex Inst. xiii, 186.—Bell in Geological Rep. Canada, 1879–80, 46.—Veitch, Manual Conif. 158.

P. sylvestris, var. *divaricata*, Aiton, Hort. Kew. iii, 366.

P. Hudsonica, Poir. in Lamarek, Dict. v, 339.—Parlatore in De Candolle, Prodr. xvi, 380.—Wood, Bot. & Fl. 313.—Koch, Dendrologie, ii, 298.

P. rupestris, Michaux f. Hist. Arb. Am. i, 49, t. 2; N. American Sylva, 3 ed. iii, 95, t. 136.

GRAY PINE. SCRUB PINE. PRINCE'S PINE.

Bay of Chaleur, New Brunswick, to the southern shores of Hudson bay, northwest to the Great Bear lake, the valley of the Mackenzie river, and the eastern slope of the Rocky mountains between the fifty-second and sixty-fifth degrees of north latitude; south to northern Maine, Ferrisburg, Vermont (*R. E. Robinson*), the southern shore of lake Michigan, and central Minnesota.

A small tree, 9 to 22 meters in height, with a trunk rarely exceeding 0.75 meter in diameter; barren, sandy soil or, less commonly, in rich loam; most common north of the boundary of the United States, and reaching its greatest development in the region north of lake Superior, here often forming considerable forests; toward its extreme western limits associated and often confounded with the closely allied *P. contorta* and *P. Murrayana* of the Pacific region.

Wood light, soft, not strong, rather close-grained, compact; bands of small summer cells not broad, very resinous, conspicuous, resin passages few, not large; medullary rays numerous, obscure; color, clear light brown or, rarely, orange, the thick sap-wood almost white; specific gravity, 0.4761; ash, 0.23; largely used for fuel, railway ties, etc.

380.—*Pinus palustris*, Miller,

Dict. 7 ed. No. 14.—Marshall, Arbustum, 100.—Wangenheim, Amer. 73.—Walter, Fl. Caroliniana, 237.—Aiton, Hort. Kew. iii, 368; 2 ed. v, 317.—Abbot, Insects Georgia, i, t. 42.—Du Roi, Harbk. 2 ed. ii, 66.—Michaux, Fl. Bor.-Am. ii, 204.—Lambert, *Pinus*, 1 ed. i, 27, t. 20; 2 ed. i, 30, t. 21; 3 ed. i, 41, t. 24, 25.—Willdenow, Spec. iv, 499.—Poir. in Lamarek, Dict. v, 341.—Persoon, Syn. ii, 578.—Desfontaines, Hist. Arb. ii, 612.—Pursh, Fl. Am. Sept. ii, 644.—Smith in Rees' Cycl. xxviii, No. 15.—Nuttall, Genera, ii, 223; Sylva, iii, 126; 2 ed. ii, 185.—Hayne, Dend. Fl. 174.—Elliott, Sk. ii, 637.—Sprengel, Syst. ii, 887.—Eaton, Manual, 6 ed. 266.—Forbes, Pinetum Woburn. 59, t. 22.—Eaton & Wright, Bot. 359.—Antoine, Conif. 23, t. 6, f. 2.—Link in Linnæa, xv, 206.—Griffith, Med. Bot. 604.—Darby, Bot. S. States, 515.—Cooper in Smithsonian Rep. 1858, 257.—Wood, Cl. Book, 660.—Porcher, Resources S. Forests, 495.—Michaux f. N. American Sylva, 3 ed. iii, 106, t. 141 (the plate as *P. australis*).

P. australis, Michaux f. Hist. Arb. Am. i, 64, t. 6.—Nouveau Duhamel, v, 246, t. 75, f. 3.—London, Arboretum, iv, 2255, f. 2156–2160.—Lindley in Penn. Cycl. xvii, 171.—Spach, Hist. Veg. xi, 392.—Endlicher, Syn. Conif. 165.—Carson, Med. Bot. ii, 43, t. 87.—Gihoul, Arb. Resin. 33.—Knight, Syn. Conif. 30.—Lindley & Gordon in Jour. Hort. Soc. London, v, 217.—Carrière, Trait. Conif. 345; 2 ed. 450.—Gordon, Pinetum, 187; Suppl. 63; 2 ed. 260.—Chapman, Fl. S. States, 434.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 24.—Wood, Bot. & Fl. 313.—Henkel & Hochstetter, Nadelhölz. 65.—Nelson, Pinaceæ, 103.—Hoopes, Evergreens, 109.—Parlatore in De Candolle, Prodr. xvi, 392.—Young, Bot. Texas, 517.—Vasey, Cat. Forest Trees, 31.—Bentley & Trimen, Med. Pl. iv, 258, t. 258.—Engelmann in Trans. St. Louis Acad. iv, 185.—Veitch, Manual Conif. 172.

LONG-LEAVED PINE. SOUTHERN PINE. GEORGIA PINE. YELLOW PINE. HARD PINE.

Southeastern Virginia, south to cape Canaveral and Tampa bay, Florida, and through the Gulf states to the valley of the Red river, Louisiana, and the Trinity river, Texas, rarely extending beyond 150 miles from the coast.

A tree of the first economic value, 18 to 29 meters in height, with a trunk 0.60 to 1.20 meter in diameter; dry, sandy loam of the maritime plain, generally of Tertiary formation, and forming, outside of the river bottoms, extensive forests almost to the exclusion of other species, or toward its extreme interior range, especially in the Gulf states, occupying rolling hills, here mixed with oaks and various deciduous trees; rarely along the borders of swamps in low, wet soil.

Wood heavy, exceedingly hard, very strong, tough, coarse-grained, compact, durable; bands of small summer cells broad, occupying fully half the width of the annual growth, very resinous, dark colored, resin passages few, not conspicuous; medullary rays numerous, conspicuous; color, light red or orange, the thin sap-wood nearly white; specific gravity, 0.6999; ash, 0.25; largely manufactured into lumber and used in construction of all sorts, for ship-building, fencing, railway ties, etc.

The turpentine, tar, pitch, rosin, and spirits of turpentine manufactured in the United States are almost exclusively produced by this species (*U. S. Dispensatory*, 14 ed. 709, 899.—*Nat. Dispensatory*, 2 ed. 1417.—*Flückiger & Hanbury, Pharmacographia*, 545).

381.—*Pinus Cubensis*, Grisebach,

Mem. Am. Acad. viii, 530; Cat. Pl. Cuba, 217.—Parlatore in De Candolle, Prodr. xvi², 396.

P. Tæda, var. *heterophylla*, Elliott, Sk. ii, 636.

P. Elliottii, Engelmann; Vasey, Cat. Forest Trees, 30; Trans. St. Louis Acad. iv, 186, t. 1, 2, 3.—Chapman, Fl. S. States, Suppl. 650.

P. Cubensis, var. *terthrocarpa*, Wright.—Grisebach, Cat. Pl. Cuba, 217.

SLASH PINE. SWAMP PINE. BASTARD PINE. MEADOW PINE.

South Carolina (Bluffton, *Mellichamp*), south near the coast to the southern keys of Florida, west along the Gulf coast to the valley of the Pearl river, Louisiana, not extending beyond 50 or 60 miles inland; in the West Indies.

A tree 24 to 30 meters in height, with a trunk 0.60 to 0.90 meter in diameter; light sandy soil along the dunes and marshes of the coast, or wet clay borders of ponds, abandoned fields, etc., and now rapidly taking possession of ground from which the forests of *P. palustris* have been removed; the only species of Florida south of cape Canaveral and bay Biscayne.

Wood heavy, exceedingly hard, very strong, tough, coarse-grained, compact, durable; bands of small summer cells very broad, occupying fully half the width of the annual growth, very resinous, conspicuous, resin passages few, not large; medullary rays numerous, rather prominent; color, rich dark orange, the sap-wood lighter, often nearly white; specific gravity, 0.7504; ash, 0.26; hardly inferior in value to that of *P. palustris*, although rarely manufactured into lumber.

Turpentine is occasionally manufactured in southern Florida from this species.

NOTE.—Specimens collected upon the southern keys of Florida by A. H. Curtiss connect the forms of South Carolina, Georgia, and northern Florida with the West Indian tree.

382.—*Picea nigra*, Link,

Linnæa, xv, 520.—Carrière, Trait. Conif. 241; 2 ed. 323.—Hooker f. in Trans. Linnæan Soc. xxiii², 301.—Brunet, Hist. Picea, 10 & t. f. B.—Peck in Trans. Albany Inst. viii, 283.—Engelmann in London Gard. Chronicle, 1879, 334.—Sears in Bull. Essex Inst. xiii, 185.

Abies Mariana, Miller, Dict.—Wangenheim, Amer. 75.

Pinus Mariana, Du Roi, Obs. Bot. 38; Harbk. ii, 107.—Ehrhart, Beitr. iii, 24.

Pinus Abies Canadensis, Marshall, Arbustum, 103.

Pinus Americana rubra, Wangenheim, Amer. 75.

Pinus nigra, Aiton, Hort. Kew. iii, 370; 2 ed. v, 319.—Lambert, Pinus, 1 ed. i, 41, t. 27; 2 ed. i, 45, t. 27; 3 ed. i, 64, t. 37.—Willdenow, Spec. iv, 506; Enum. 990; Berl. Baumz. 278.—Persoon, Syn. ii, 579.—Pursh, Fl. Am. Sept. ii, 640.—Smith in Rees' Cycl. xxviii, No. 20.—Barton, Compend. Fl. Philadelph. ii, 182.—Nuttall, Genera, ii, 223.—Hayne, Dend. Fl. 177.—Elliott, Sk. ii, 640.—Sprengel, Syst. ii, 885.—Torrey, Compend. Fl. N. States, 359; Fl. N. York, ii, 230.—Beck, Bot. 340.—Eaton, Manual, 6 ed. 264.—Hooker, Fl. Bor.-Am. ii, 163.—Eaton & Wright, Bot. 358.—Bigelow, Fl. Boston. 3 ed. 386.—Antoine, Conif. 88, t. 34, f. 3.—Endlicher, Syn. Conif. 115.—Darby, Bot. S. States, 515.—Porcher, Resources S. Forests, 505.—Parlatore in De Candolle, Prodr. xvi², 413.

Pinus Americana, Gärtner, Fruct. ii, 60, t. 91, f. 1.

Pinus rubra, Lambert, Pinus, 1 ed. i, 48, t. 28; 2 ed. i, 47, t. 30; 3 ed. i, 66, t. 38 [not Michaux f.].—Persoon, Syn. ii, 579.—Aiton, Hort. Kew. 2 ed. v, 319.—Pursh, Fl. Am. Sept. ii, 640.—Smith in Rees' Cycl. xxviii, No. 23.—Nuttall, Genera, ii, 223.—Sprengel, Syst. ii, 825.—Torrey, Compend. Fl. N. States, 359.—Beck, Bot. 340.—Eaton, Manual, 6 ed. 264.—Hooker, Fl. Bor.-Am. ii, 164.—Eaton & Wright, Bot. 358.—Antoine, Conif. 87, t. 34, f. 2.—Endlicher, Syn. Conif. 113.—Gihoul, Arb. Resin. 44.—Parlatore in De Candolle, Prodr. xvi², 413.

Abies denticulata, Michaux, Fl. Bor.-Am. ii, 206.—Poiret in Lamarck, Dict. vi, 520.

Abies nigra, Poiret in Lamarck, Dict. vi, 520.—Desfontaines, Hist. Arb. ii, 580.—Michaux f. Hist. Arb. Am. i, 124, t. 11; N. American Sylva, 3 ed. iii, 139, t. 147.—Nouveau Duhamel, v, 292, t. 81, f. 1.—Lindley in Penn. Cycl. i, 32.—London, Arboretum, iv, 2312, f. 2225-2227.—Spach, Hist. Veg. xi, 410, in part.—Emerson, Trees Massachusetts, 81; 2 ed. ii, 96.—Griffith, Med. Bot. 606.—Knight, Syn. Conif. 36.—Lindley & Gordon in Jour. Hort. Soc. London, v, 211.—Parry in Owen's Rep. 618.—Gordon, Pinetum, 11; 2 ed. 17.—Richardson, Arctic Exped. 442.—Cooper in Smithsonian Rep. 1858, 257.—Chapman, Fl. S. States, 434.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 27.—Wood, Cl. Book, 662; Bot. & Fl. 313.—Porcher, Resources S. Forests, 507.—Henkel & Hochstetter, Nadelhölz. 191.—Nelson, Pinaceæ, 50.—Gray, Manual N. States, 5 ed. 471.—Hoopes, Evergreens, 169.—Vasey, Cat. Forest Trees, 33.—Guibourt, Hist. Drogues, 7 ed. ii, 247.—Macoun in Geological Rep. Canada, 1875-'76, 211.—Bell in Geological Rep. Canada, 1879-'80, 44^c.—Veitch, Manual Conif. 74.

Abies rubra, Poiret in Lamarck, Dict. vi, 520.—Desfontaines, Hist. Arb. ii, 580.—London, Arboretum, iv, 2316, f. 2228.—Forbes, Pinetum Woburn. 101, t. 35.—Knight, Syn. Conif. 37.—Lindley & Gordon in Jour. Hort. Soc. London, v, 211.—Gordon, Pinetum, 11; 2 ed. 17.—Henkel & Hochstetter, Nadelhölz. 189.—Nelson, Pinaceæ, 51.

P. rubra, Link in Linnæa, xv, 521.—Carrière, Trait. Conif. 240; 2 ed. 322.

Abies nigra, var. *rubra*, Michaux f. Hist. Arb. Am. i, 123; N. American Sylva, 3 ed. iii, 141.—Spach, Hist. Veg. xi, 411.—Hoopes, Evergreens, 170.

? *Abies rubra*, var. *arctica*, Lindley & Gordon in Jour. Hort. Soc. London, v, 211.

Abies alba, Chapman, Fl. S. States, 435 [not Poiret].

Abies Americana, Koch, Dendrologie, ii², 241.

P. nigra, var. *rubra*, Engelmann in London Gard. Chronicle, 1879, 334.

Abies arctica, Hort.

Abies Marylandica, Hort.

BLACK SPRUCE.

Newfoundland, northern Labrador to Ungava bay, Nastapokee sound, cape Churchill, Hudson bay, and northwest to the mouth of the Mackenzie river and the eastern slope of the Rocky mountains; south through the northern states to Pennsylvania, central Michigan, Wisconsin, and Minnesota, and along the Alleghany mountains to the high peaks of North Carolina.

A tree 15 to 21 meters in height, with a trunk 0.60 to 0.90 meter in diameter; light, dry, rocky soil, forming, especially north of the fiftieth degree of latitude, extensive forests on the water-sheds of the principal streams or in cold, wet swamps; then small, stunted, and of little value (*P. rubra*).

Wood light, soft, not strong, close, straight-grained, compact, satiny; bands of small summer cells thin, resinous, resin passages few, minute; medullary rays few, conspicuous; color, light red or often nearly white, the sap-wood lighter; specific gravity, 0.4584; ash, 0.27; largely manufactured into lumber, used in construction, for ship-building, piles, posts, railway ties, etc.

Essence of spruce, prepared by boiling the young branches of this species, is used in the manufacture of spruce beer, a popular beverage (*U. S. Dispensatory*, 14 ed. 901).

383.—*Picea alba*, Link,

Linnæa, xv, 519.—Carrière, Trait. Conif. 238; 2 ed. 319.—Fl. des Serres, xxi, 157, t. 2251.—Brunet, Hist. Picea, 4 & t. f. A.—Engelmann in London Gard. Chronicle, 1879, 334.—Sears in Bull. Essex Inst. xiii, 184.

Abies Canadensis, Miller, Diet. No. 1.

Pinus Canadensis, Du Roi, Obs. Bot. 38; Harbk. ii, 124 [not Linnæus].—Wangenheim, Amer. 5, t. 1, f. 2.

P. laxa, Ehrhart, Beitr. iii, 24.

P. glauca, Mönch, Weiss. 73.

Pinus alba, Aiton, Hort. Kew. iii, 371; 2 ed. v, 318.—Lambert, Pinus, 1 ed. i, 39 t. 26; 2 ed. i, 43, t. 28; 3 ed. i, 61, t. 35.—Willdenow, Spec. iv, 507; Enum. 990; Berl. Baumz. 280.—Persoon, Syn. ii, 579.—Pursh, Fl. Am. Sept. ii, 641.—Smith in Rees' Cycl. xxviii, No. 21.—Eaton, Manual, 6 ed. 264.—Nuttall, Genera, ii, 223.—Hayne, Dend. Fl. 177.—Elliott, Sk. ii, 640.—Sprengel, Syst. ii, 835.—Torrey, Compend. Fl. N. States, 359; Fl. N. York, ii, 231.—Meyer, Pl. Labrador, 30.—Beck, Bot. 340.—Hooker, Fl. Bor.-Am. ii, 163.—Eaton & Wright, Bot. 358.—Bigelow, Fl. Boston. 3 ed. 386.—Antoine, Conif. 86, t. 34, f. 1.—Endlicher, Syn. Conif. 112.—Darby, Bot. S. States, 515.—Tuinbouw Flora, 1855, 1, t. 14, 15.—Walpers, Ann. v, 799.—Parlatore in De Candolle, Prodr. xvi², 414.

Pinus tetragona, Mönch, Meth. 364.

Abies alba, Poiret in Lamarck, Diet. vi, 521.—Michaux, Fl. Bor.-Am. ii, 207.—Desfontaines, Hist. Arb. ii, 560.—Michaux f. Hist. Arb. Am. i, 133, t. 12; N. American Sylva, 3 ed. iii, 144, t. 148.—Nouveau Duhamel, v, 291, t. 81, f. 2.—London, Arboretum, iv, 2310, f. 2224.—Forbes, Pinetum Woburn. 95, t. 33.—Nuttall, Sylva, iii, 129; 2 ed. ii, 189.—Spach, Hist. Veg. xi, 412.—Emerson, Trees Massachusetts, 84; 2 ed. i, 99.—Gihoul, Arb. Resin. 43.—Knight, Syn. Conif. 36.—Lindley & Gordon in Jour. Hort. Soc. London, v, 211.—Parry in Owen's Rep. 618.—Gordon, Pinetum, 2; 2 ed. 3.—Richardson, Arctic Exped. 442.—Cooper in Smithsonian Rep. 1858, 257.—Hooker f. in Trans. Linnæan Soc. xxiii², 301.—Engelmann in Am. Jour. Sci. 2 ser. xxxiv, 330.—Wood, Cl. Book, 661; Bot. & Fl. 313.—Porcher, Resources S. Forests, 507.—Henkel & Hochstetter, Nadelholz. 188.—Nelson, Pinaceæ, 47.—Gray, Manual N. States, 5 ed. 471.—Murray in Seemann, Jour. Bot. v, 253, t. 69, f. 2-7.—Hoopes, Evergreens, 157, f. 20.—Vasey, Cat. Forest Trees, 32.—Guibourt, Hist. Drogues, 7 ed. ii, 247.—Macoun in Geological Rep. Canada, 1875-76, 211.—Bell in Geological Rep. Canada, 1879-80, 44^c.

Abies rubra, var. *cærulea*, London, Arboretum, iv, 2316.—Lindley & Gordon in Jour. Hort. Soc. London, v, 211.

Abies cærulea, Forbes, Pinetum Woburn. 99.

P. cærulea, Link in Linnæa, xv, 522.

Pinus rubra, var. *violacea*, Endlicher, Syn. Conif. 114.

P. nigra, var. *glauca*, Carrière, Trait. Conif. 1 ed. 242.

Abies arctica, Murray in Seemann, Jour. Bot. v, 253, t. 69, f. 1, 8-13.

Abies laxa, Koch, Dendrologie, ii², 243.

Abies alba, var. *cærulea*, Carrière, Trait. Conif. 2 ed. 320.

Abies alba, var. *arctica*, Parlatore in De Candolle, Prodr. xvi², 414.

WHITE SPRUCE.

Newfoundland, northern shore of Labrador to Ungava bay, cape Churchill, and northwestward to the mouth of the Mackenzie river and the valley of the Yukon river, Alaska; south to the coast of Maine, northeastern Vermont (West Burke and Elmwood, *Pringle*), northern Michigan, Minnesota to Moose lake and the White Earth Indian reservation, the Black hills of Dakota (*R. Douglas*), along the Rocky mountains of northern Montana to the valley of the Blackfoot river (*Canby & Sargent*), Sitka, and British Columbia.

A tree 15 to 50 meters in height, with a trunk 0.60 to 0.90 meter in diameter; low, rather wet soil, borders of ponds and swamps; most common north of the boundary of the United States, and reaching its greatest development along the streams and lakes of the Flathead region of northern Montana at an elevation of 2,500 to 3,500 feet; the most important timber tree of the American subarctic forests north of the sixtieth degree of latitude, here more generally multiplied and of larger size than the allied *P. nigra*, with which it is associated; its distribution southward in British Columbia not yet satisfactorily determined.

Wood light, soft, not strong, close, straight-grained, compact, satiny; bands of small summer cells thin, not conspicuous, resin passages few, minute; medullary rays numerous, prominent; color, light yellow, the sap-wood hardly distinguishable; specific gravity, 0.4051; ash, 0.32; largely manufactured into lumber, although not distinguished in commerce from that of the black spruce (*P. nigra*).

384.—*Picea Engelmanni*, Engelmann,

Trans. St. Louis Acad. ii, 212; Wheeler's Rep. vi, 256; London Gard. Chronicle, 1879, 334; 1882, 145.—Carrière, Trait. Conif. 2 ed. 348.—G. M. Dawson in Canadian Nat. new ser. ix, 325.—Rusby in Bull. Torrey Bot. Club, ix, 80.

Abies alba, ? Torrey in Fremont's Rep. 97.

Abies nigra, Engelmann in Am. Jour. Sci. 2 ser. xxxiii, 330 [not Poirét].

Abies Engelmanni, Parry in Trans. St. Louis Acad. ii, 122; London Gard. Chronicle, 1863, 1035; Am. Nat. viii, 179; Proc. Davenport Acad. i, 149.—Regel, Gartenflora, 1864, 244.—Henkel & Hochstetter, Nadelhölz. 418.—Hoopes, Evergreens, 177, f. 22.—Watson in King's Rep. v, 332; Pl. Wheeler, 17.—Porter in Hayden's Rep. 1871, 494.—Porter & Coulter, Fl. Colorado; Hayden's Surv. Misc. Pub. No. 4, 130.—Vasey, Cat. Forest Trees, 33.—Koch, Dendrologie, ii², 242.—Hall in Coulter's Bot. Gazette, ii, 91.—Sargent in London Gard. Chronicle, 1877, 631.—Macoun in Geological Rep. Canada, 1875-'76, 211.—Brandeggee in Coulter's Bot. Gazette, iii, 32.—Bell in Geological Rep. Canada, 1879-'80, 56c.—Veitch, Manual Conif. 68.

Pinus Engelmanni, Engelmann in Proc. Am. Phil. Soc. new ser. xii, 209.

Pinus commutata, Parlatore in De Candolle, Prodr. xvi², 417.—Gordon, Pinetum, 2 ed. 5.

WHITE SPRUCE.

Peace River plateau, in latitude 55° 46' N. (*G. M. Dawson*), through the interior of British Columbia and along the Cascade mountains of Washington territory and Oregon to the valley of the Mackenzie river; along the principal ranges of the Rocky and Wahsatch mountains to the San Francisco mountains, Sierra Blanco, and mount Graham, Arizona.

A large tree, 24 to 46 meters in height, with a trunk 0.90 to 1.20 meter in diameter, or at its extreme elevation reduced to a low, prostrate shrub; dry, gravelly slopes and ridges between 5,000 and 11,500 feet elevation; the most valuable timber tree of the central Rocky Mountain region, here forming extensive forests, generally above 8,500 feet elevation; rare and of small size in the mountains of Washington territory, Oregon, and Montana.

Wood very light, soft, not strong, very close, straight-grained, compact, satiny; bands of small summer cells narrow, not conspicuous, resin passages few, minute; medullary rays numerous, conspicuous; color, pale yellow tinged with red, the sap-wood hardly distinguishable; specific gravity, 0.3449; ash, 0.32; in Colorado manufactured into lumber and largely used for fuel, charcoal, etc.

The bark rich in tannin, and in Utah sometimes used in tanning leather.

NOTE.—Forms of northern Montana too closely connect this species with the allied *P. alba*. The two species occur here, however, only at different elevations, in different soils, and never mingle.

385.—*Picea pungens*, Engelmann,

London Gard. Chronicle, 1879, 334; 1882, 145.—Masters in London Gard. Chronicle, 1883, 725, f. 130.

P. Menziesii, Engelmann in Trans. St. Louis Acad. ii, 214 [not Carrière].

Abies Menziesii, Engelmann in Am. Jour. Sci. 2 ser. xxxiii, 330 [not Lindley].—Gray in Proc. Philadelphia Acad. 1863, 76.—Watson in King's Rep. v, 333, in part.—Parry in Am. Nat. viii, 179 [not Lindley].—Porter in Hayden's Rep. 1871, 494.—Hoopes, Evergreens, 166, in part.—Rothrock in Pl. Wheeler, 28; Wheeler's Rep. vi, 10 [not Lindley].—Porter & Coulter, Fl. Colorado; Hayden's Surv. Misc. Pub. No. 4, 131 [not Lindley].—Vasey, Cat. Forest Trees, 33, in part.—Brandeggee in Coulter's Bot. Gazette, iii, 32.

Abies Menziesii Parryana, André in Ill. Hort. xxiii, 198; xxiv, 53, 119.—Roezl in Ill. Hort. xxiv, 86.

Abies Engelmanni glauca, Veitch, Manual Conif. 69.

. WHITE SPRUCE. BLUE SPRUCE.

Valley of the Wind river, south through the mountain ranges of Wyoming, Colorado, and Utah.

A tree 30 to 46 meters in height, with a trunk 0.60 to 0.90 meter in diameter; borders of streams, in damp or wet soil, generally between 6,000 and 9,000 feet elevation, never forming forests or reaching as high elevations as the allied *P. Engelmanni*; rare and local.

Wood very light, soft, weak, close-grained, compact, satiny; bands of small summer cells narrow, not conspicuous, resin passages few, small; medullary rays numerous, prominent; color, very light brown or often nearly white, the sap-wood hardly distinguishable; specific gravity, 0.3740; ash, 0.38.

386.—*Picea Sitchensis*, Carrière,

Trait. Conif. 1 ed. 260; Engelmann in London Gard. Chronicle, 1879, 344; Bot. California, ii, 122.

Pinus Sitchensis, Bongard in Mem. Acad. St. Petersburg, 6 ser. ii, 104.—Hooker, Fl. Bor.-Am. ii, 164.—Endlicher, Syn. Conif. 123.

Abies Menziesii, Lindley in Penn. Cycl. 1, 32.—London, Arboretum, iv, 2321, f. 2232.—Forbes, Pinetum Woburn. 93, t. 32.—Nuttall, Sylva, iii, 131, t. 116; 2 ed. ii, 189, t. 116.—Knight, Syn. Conif. 37.—Lindley & Gordon in Jour. Hort. Soc. London, v, 211.—Newberry in Pacific R. R. Rep. vi, 56, 90, t. 9, f. 21.—Gordon, Pinetum, 6; 2 ed. 12.—Cooper in Smithsonian Rep. 1858, 262; Pacific R. R. Rep. xii², 25, 69, in part.—Wood, Bot. & Fl. 314.—Lyll in Jour. Linnæan Soc. vii, 131, 133, 144.—Henkel & Hochstetter, Nadelhölz. 187.—Nelson, Pinaceæ, 148.—Rothrock in Smithsonian Rep. 1867, 433.—Hoopes, Evergreens, 166, in part.—Watson in King's Rep. v, 333, in part.—Veitch, Manual Conif. 73.

Pinus Menziesii, Douglas in Lambert, Pinus, 1 ed. iii, 161, t. 71.—Hooker, Fl. Bor.-Am. ii, 162.—Antoine, Conif. 85, t. 33, f. 1, 2.—Hooker & Arnott, Bot. Beechey, 394.—Endlicher, Syn. Conif. 112.—Parlatore in De Candolle, Prodr. xvi², 418.

? *Abies trigona*, Rafinesque, Atlant. Jour. 119.—Endlicher, Syn. Conif. 124.—Carrière, Trait. Conif. 1 ed. 264.

? *Abies falcata*, Rafinesque, Atlant. Jour. 119.—Endlicher, Syn. Conif. 124.—Lindley & Gordon in Jour. Hort. Soc. London, v, 213.—Carrière, Trait. Conif. 268; 2 ed. 314.

Pinus Menziesii, var. *crispa*, Antoine, Conif. 85, t. 35, f. 2.

Abies Sitchensis, Lindley & Gordon in Jour. Hort. Soc. London, v, 212.—Koch, Dendrologie, ii², 247.

P. Menziesii, Carrière, Man. des Pl. iv, 339; Trait. Conif. 237; 2 ed. 318.

? *Sequoia Rafinesquei*, Carrière, Trait. Conif. 2 ed. 213.

TIDE-LAND SPRUCE.

Alaska, south to Mendocino county, California, not extending more than 50 miles inland from the coast.

A large tree of great economic value, 46 to 61 meters in height, with a trunk 2.40 to 5.19 meters in diameter; gravelly ridges and swamps, reaching its greatest development in Washington territory and Oregon near the mouth of the Columbia river, here forming a belt of nearly continuous forest growth 50 or, farther north and south, rarely more than 10 or 15 miles in width.

Wood light, soft, not strong, close, straight-grained, compact, satiny; bands of small summer cells narrow, not conspicuous, resin passages few, obscure; medullary rays numerous, rather prominent; color, light brown tinged with red, the sap-wood nearly white; specific gravity, 0.4287; ash, 0.17; largely manufactured into lumber and used for construction, interior finish, fencing, boat-building, the dunnage of vessels, cooperage, woodenware, etc.

387.—*Tsuga Canadensis*, Carrière,

Trait. Conif. 189; 2 ed. 248.—Sears in Bull. Essex Inst. xiii, 184.—Engelmann in Coulter's Bot. Gazette, vi, 224.

Pinus Canadensis, Linnæus, Spec. 2 ed. 1421.—Wangenheim, Amer. 39, t. 15, f. 36.—Ehrhart, Beitr. iii, 23.—Aiton, Hort. Kew. iii, 370; 2 ed. v, 320.—Michaux, Fl. Bor.-Am. ii, 206.—Lambert, Pinus, 1 ed. 50, t. 32; 2 ed. i, 56, t. 35; 3 ed. ii, 79, t. 45.—Willdenow, Spec. iv, 505; Enum. 989; Berl. Baumz. 277.—Poirot in Lamarek, Dict. vi, 521.—Persoon, Syn. ii, 579.—Pursh, Fl. Am. Sept. ii, 640.—Smith in Rees' Cycl. xxviii, No. 29.—Barton, Compend. Fl. Philadelph. ii, 182.—Nuttall, Genera, ii, 223.—Hayne, Dend. Fl. 176.—Elliott, Sk. ii, 639.—Sprengel, Syst. ii, 885.—Torrey, Compend. Fl. N. States, 359; Fl. New York, ii, 230.—Beck, Bot. 340.—Eaton, Manual, 6 ed. 264.—Darlington, Fl. Cestrica, 2 ed. 548.—Hooker, Fl. Bor.-Am. ii, 164, in part.—Eaton & Wright, Bot. 358.—Bigelow, Fl. Boston. 3 ed. 386.—Antoine, Conif. 80, t. 32, f. 3.—Endlicher, Syn. Conif. 86.—Gihoul, Arb. Resin. 46.—Darby, Bot. S. States, 515.—Parlatore in De Candolle, Prodr. xvi², 428.—McNab in Proc. Royal Irish Acad. 2 ser. ii, 211, 212, t. 23, f. 3.—Bentley & Trimen, Med. Pl. iv, 264, t. 264.

Pinus Americana, Miller, Dict. 7 ed. No. 6.—Du Roi, Obs. Bot. 41; Harbk. 2 ed. ii, 151.

Pinus Abies Americana, Marshall, Arbustum, 103.

Abies Canadensis, Desfontaines, Hist. Arb. ii, 590.—Michaux f. Hist. Arb. Am. i, 138, t. 13; N. American Sylva, 3 ed. iii, 146, t. 140.—Nouveau Duhamel, v, 293, t. 83, f. 1.—Eaton, Manual, 111.—Richard, Conif. 77, t. 17, f. 2.—Audubon, Birds, t. 197.—London, Arboretum, iv, 2322 & t.—Forbes, Pinetum Woburn. 129.—Nuttall, Sylva, iii, 133; 2 ed. ii, 190.—Spach, Hist. Veg. xi, 424.—Emerson, Trees Massachusetts, 77; 2 ed. i, 92 & t.—Griffith, Med. Bot. 606.—Knight, Syn. Conif. 37.—Lindley & Gordon in Jour. Hort. Soc. London, v, 209.—Parry in Owen's Rep. 618.—Darlington, Fl. Cestrica, 3 ed. 291.—Gordon, Pinetum, 14; 2 ed. 22.—Cooper in Smithsonian Rep. 1858, 257.—Chapman, Fl. S. States, 434.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 27.—Wood, Cl. Book, 661; Bot. & Fl. 313.—Porcher, Resources S. Forests, 506.—Henkel & Hochstetter, Nadelhölz. 153 (excl. syn. *aromatica*).—Nelson, Pinaceæ, 30.—Gray, Manual N. States, 5 ed. 471.—Hoopes, Evergreens, 184, f. 23.—Koch, Dendrologie, ii², 249.—Vasey, Cat. Forest Trees, 23.—Fl. des Serres, xxii, 206.—Guibourt, Hist. Drogues, ii, 247.—Bell in Geological Rep. Canada, 1879-'80, 51^c.—Veitch, Manual Conif. 114, f. 29.

Picea Canadensis, Link in Linnæa, xv, 524.

HEMLOCK.

Nova Scotia, southern New Brunswick, valley of the Saint Lawrence river to the shores of lake Temiscaming, and southwest to the western borders of northern Wisconsin; south through the northern states to New Castle county, Delaware, southeastern Michigan, central Wisconsin, and along the Alleghany mountains to Clear Creek falls, Winston county, Alabama (*Mohr*).

A tree 21 to 33 meters in height, with a trunk 0.90 to 1.15 meter in diameter; dry, rocky ridges, generally facing the north and often forming extensive forests almost to the exclusion of other species, or, less commonly, borders of swamps in deep, rich soil; most common at the north, although reaching its greatest individual development in the high mountains of North Carolina and Tennessee.

Wood light, soft, not strong, brittle, coarse, crooked-grained, difficult to work, liable to wind-shake and splinter, not durable; bands of small summer cells rather broad, conspicuous; medullary rays numerous, thin; color, light brown tinged with red or often nearly white, the sap-wood somewhat darker; specific gravity, 0.4239; ash, 0.46; largely manufactured into coarse lumber and used in construction for outside finish, railway ties, etc.; two varieties, red and white, produced apparently under precisely similar conditions of growth, are recognized by lumbermen.

The bark, rich in tannin, is the principal material used in the northern states in tanning leather, and yields a fluid extract sometimes used medicinally as a powerful astringent.

Canada or hemlock pitch, prepared from the resinous secretion of this species, is used in the preparation of stimulating plasters, etc. (*U. S. Dispensatory*, 14 ed. 709, 903.—*Nat. Dispensatory*, 2 ed. 1109.—*Flückiger & Hanbury, Pharmacographia*, 552).

388.—*Tsuga Caroliniana*, Engelm.

Coulter's Bot. Gazette, vi, 223.

Abies species, Gibbs in Proc. Elliott Soc. i, 286.

Abies Caroliniana, Chapman, Fl. S. States, Suppl. 650.

HEMLOCK.

Southern Alleghany region, Bluff mountain, North Carolina (*A. Gray*), "Saluda mountain," South Carolina (*L. S. Gibbs*), Pinnacle mountain, North Carolina (*Curtiss*), New river, North Carolina, and Cæsar's head, South Carolina (*Canby*), Whitesides mountain and Devil's Court-House peak, Jackson county, North Carolina (*J. Donnell Smith*).

A small tree, 12 to 15 meters in height, with a trunk 0.60 to 0.75 meter in diameter; dry, rocky ridges between 4,000 and 5,000 feet elevation; rare and local; long confounded with the closely allied *T. Canadensis*, from which it may be distinguished by its larger, glossier, blunter leaves, and larger cones with wide-spreading scales.

Wood light, soft, not strong, brittle, coarse-grained; bands of small summer cells narrow, not conspicuous; medullary rays numerous, thin; color, light brown tinged with red, the sap-wood nearly white; specific gravity, 0.4275; ash, 0.40.

389.—*Tsuga Mertensiana*, Carrière,

Trait. Conif. 2 ed. 250.—Engelmann in Bot. California, ii, 121; Coulter's Bot. Gazette, vi, 224.—G. M. Dawson in Canadian Nat. new ser. ix, 324.

?*Abies heterophylla*, Rafinesque, Atlant. Jour. 119.—Endlicher, Syn. Conif. 124.—Carrière, Trait. Conif. 1 ed. 265.

Pinus Mertensiana, Bongard in Mem. Acad. St. Petersburg, 6 ser. iii, 163.—Hooker, Fl. Bor.-Am. ii, 164.—Endlicher, Syn. Conif. 111.—Ledebour, Fl. Rossica, iii, 668.—Parlatore in De Candolle, Prodr. xvi², 428.—McNab in Proc. Royal Irish Acad. 2 ser. ii, 211, 212, t. 23, f. 4.

Pinus Canadensis, Bongard in Mem. Acad. St. Petersburg, 6 ser. iii, 163 [not Linnæus].—Douglas in Companion Bot. Mag. ii, 127.—Hooker, Fl. Bor.-Am. ii, 164, in part.—Ledebour, Fl. Rossica, iii, 668.

Abies Mertensiana, Lindley & Gordon in Jour. Hort. Soc. London, v, 211.—Carrière, Trait. Conif. 1 ed. 232.—Gordon, Pinetum, 18; Suppl. 12; 2 ed. 29.—Lyll in Jour. Linnæan Soc. vii, 133, 144.—Henkel & Hochstetter, Nadelhölz. 152.—Rothrock in Smithsonian Rep. 1867, 433.—Cooper in Am. Nat. iii, 412.—Gray in Proc. Am. Acad. vii, 402.—Heopes, Evergreens, 192.—Koch, Dendrologie, ii², 250.—Vasey, Cat. Forest Trees, 33.—Macoun in Geological Rep. Canada 1875-'76, 211.—Hall in Coulter's Bot. Gazette, ii, 91.

Abies Canadensis,? Cooper in Smithsonian Rep. 1858, 262; Pacific R. R. Rep. xii², 69 [not Desfontaines].

Abies Bridgesii, Kellogg in Proc. California Acad. ii, 37.

Abies Albertiana, Murray in Proc. Hort. Soc. London, iii, 149 & f.—Lawson, Pinetum Brit. ii, 111, t. 16, f. 1-18.—Nelson, Pinaceæ, 31.—Fowler in London Gard. Chronicle, 1872, 75.

Abies taxifolia, Hartweg, *ined.* (*Ades* Murray in Proc. Hort. Soc. London, iii, 148).

Pinus Pattoniana, McNab in Proc. Royal Irish Acad. 2 ser. ii, 211, 212, t. 23, f. 2 [not Parlatores] (*Ades* Engelmann in London Gard. Chronicle, 1882, 145).

Abies Pattonii, McNab in Jour. Linnæan Soc. xix, 308.

HEMLOCK.

Alaska, south along the islands and coast of British Columbia, and through the Selkirk, Gold, and other interior ranges to the Bitter Root mountains of Idaho, and the western slopes of the Rocky mountains of Montana (valley of the Flathead river, *Canby & Sargent*), extending south along the Cascade mountains to southern Oregon and in the Coast ranges to Marin county, California, between 1,000 and 4,000 feet elevation.

A large tree, 30 to 61 meters in height, with a trunk 1.20 to 3 meters in diameter; low, moist bottoms or rocky ridges; very common and reaching its greatest development in western Oregon and Washington territory, often forming extensive forests, especially along the western base of the Cascade mountains.

Wood light, hard, not strong, rather close-grained; bands of small summer cells thin, not conspicuous; medullary rays numerous, prominent; color, light brown tinged with yellow, the sap-wood nearly white; specific gravity, 0.5182; ash, 0.42; occasionally manufactured into coarse lumber.

The bark, rich in tannin, is the principal material used on the northwest coast in tanning leather.

390.—*Tsuga Pattoniana*, Engelmann,

Bot. California, ii, 121; London Gard. Chronicle, 145.

Abies Pattoniana, Jeffrey in Rep. Oregon Exped. i, t. 4, f. 2.—Murray in Edinburgh New Phil. Jour. new ser. i, 291, t. 9, f. 1-7.—Lawson, Pinetum Brit. ii, 157, t. 22.—Gray in Proc. Am. Acad. vii, 402.—Koch, Dendrologie, ii*, 252.—Hoopes, Evergreens, 172.—Carrière, Trait. Conif. 2 ed. 30.—Hall in Coulter's Bot. Gazette, ii, 91.—Veitch, Manual Conif. 116, f. 31, 32.

? *Picea Californica*, Carrière, Trait. Conif. 261; 2 ed. 346.

Abies Hookeriana, Murray in Edinburgh New Phil. Jour. new ser. i, 289, t. 9, f. 11-17.—Lawson, Pinetum Brit. ii, 153, t. 21, 22, f. 1-22.—Nelson, Pinaceæ, 31.—McNab in Proc. Royal Irish Acad. 2 ser. ii, 211, 212, t. 23, f. 1.—Veitch, Manual Conif. 115, t. 32.

Abies Williamsonii, Newberry in Pacific R. R. Rep. vi, 53, 90, t. 7, f. 19.—Wood, Bot. & Fl. 313.—Cooper in Am. Nat. iii, 412.—Vasey, Cat. Forest Trees, 33.

Pinus Pattoniana, Parlatores in De Candolle, Prodr. xvi*, 429.

Abies Pattonii, Gordon, Pinetum, 1 ed. 10 (excl. syn. *trigona*).

Abies Pattoni, Gordon, Pinetum, Suppl. 12.—Henkel & Hochstetter, Nadelhölz. 151 (excl. syn. *trigona*).

Valley of the Fraser river, British Columbia, and probably much farther north, south along the Cascade mountains and the California Sierras to the headwaters of the San Joaquin river, extending east along the high mountains of northern Washington territory to the western slopes and summits of the Cœur d'Alêne and Bitter Root mountains of Idaho (Lolo trail, *Watson*), and the divide between Thompson and Little Bitter Root creeks, northern Montana (*H. B. Ayres*).

An alpine tree, rarely 30 meters in height, with a trunk 1.50 to 2.10 meters in diameter; dry slopes and ridges near the limits of tree growth, ranging from an elevation of 2,700 feet in British Columbia to 10,000 feet in the Sierras of central California.

Wood light, soft, not strong, close-grained, satiny, susceptible of a good polish; bands of small summer cells thin, not conspicuous; medullary rays numerous, obscure; color, light brown or red, the sap-wood nearly white; specific gravity, 0.4454; ash, 0.44.

391.—*Pseudotsuga Douglasii*, Carrière,

Trait. Conif. 2 ed. 256.—Engelmann in *Wheeler's Rep.* vi, 257; *Bot. California*, ii, 120.—G. M. Dawson in *Canadian Nat. new ser.* ix, 323.—Eichler in *Monatsb. Acad. Berl.* 1881, f. 18-22.—Rusby in *Bull. Torrey Bot. Club*, ix, 79.

Pinus taxifolia, Lambert, *Pinus*, 1 ed. i, 51, t. 33; 2 ed. i, 58, t. 36; 3 ed. ii, 82, t. 47.—Pursh, *Fl. Am. Sept.* ii, 640.—Smith in *Rees' Cycl.* xxviii, No. 28.—Sprengel, *Syst.* ii, 885.—Eaton, *Manual*, 6 ed. 264.—Eaton & Wright, *Bot.* 358.

Abies taxifolia, Poiret in *Lamarek, Dict.* vi, 523.—Nouveau Duhamel, v, 293.—Torrey & Gray in *Pacific R. R. Rep.* ii, 130.—Cooper in *Smithsonian Rep.* 1858, 262; *Pacific R. R. Rep.* xii³, 69.

Abies Douglasii, Lindley in *Penn. Cycl.* i, 32.—London, *Arboretum*, iv, 2319, f. 2230.—Forbes, *Pinetum Woburn.* 127, t. 45.—Bentham, *Pl. Hartweg.* 57.—Nuttall, *Sylva*, iii, 129, t. 115; 2 ed. ii, 187, t. 115.—Spach, *Hist. Veg.* xi, 423.—Knight, *Syn. Conif.* 37.—Lindley & Gordon in *Jour. Hort. Soc. London*, v, 209.—London *Gard. Chronicle*, 1854, 163.—Bigelow in *Pacific R. R. Rep.* iv, 17.—Torrey in *Pacific R. R. Rep.* iv, 141; *Bot. Mex. Boundary Survey*, 210; *Ives' Rep.* 28.—Newberry in *Pacific R. R. Rep.* vi, 54, 90, t. 8, f. 20.—Gordon, *Pinetum*, 15; *Suppl.* 10; 2 ed. 24.—Cooper in *Smithsonian Rep.* 1858, 262; *Pacific R. R. Rep.* xii³, 24, 69; *Am. Nat.* iii, 411.—Wood, *Bot. & Fl.* 313.—Engelmann in *Am. Jour. Sci.* 2 ser. xxxiv, 330; *Proc. Am. Phil. Soc. new ser.* xii, 209.—Lyll in *Jour. Linnæan Soc.* vii, 131, 133, 143.—Henkel & Hochstetter, *Nadelhölz.* 155.—Nelson, *Pinaceæ*, 32.—Rothrock in *Smithsonian Rep.* 1867, 433; *Pl. Wheeler*, 28, 50; *Wheeler's Rep.* vi, 9.—Hoopes, *Evergreens*, 189.—Lawson, *Pinetum Brit.* ii, 115, t. 17, 18, f. 1-23.—Porter in *Hayden's Rep.* 1871, 494.—Watson in *King's Rep.* v, 334; *Pl. Wheeler*, 17.—Fowler in *London Gard. Chronicle*, 1872, 75.—Gray in *Proc. Am. Acad.* vii, 402.—Koch, *Dendrologie*, ii³, 255.—Porter & Coulter, *Fl. Colorado*; *Hayden's Surv. Misc. Pub. No. 4*, 131.—Murray in *London Gard. Chronicle*, 1872, 106.—Vasey, *Cat. Forest Trees*, 33.—Hayden in *Warren's Rep. Nebraska & Dakota*, 2 ed. 122.—Macoun in *Geological Rep. Canada*, 1875-76, 211.—Hall in *Coulter's Bot. Gazette*, ii, 91.—Brandegge in *Coulter's Bot. Gazette*, iii, 32.—Veitch, *Manual Conif.* 119, f. 35.

Abies mucronata, Rafinesque, *Jour. Atlant.* 119.—Endlicher, *Syn. Conif.* 126.—Lindley & Gordon in *Jour. Hort. Soc. London*, v, 213.—Carrière, *Trait. Conif.* 268; 2 ed. 312.

?*Abies mucronata palustris*, Rafinesque, *Jour. Atlant.* 129.—Carrière, *Trait. Conif.* 268; 2 ed. 313.

Pinus Douglasii, Lambert, *Pinus*, 1 ed. iii, 163, t. 21.—Hooker, *Fl. Bor.-Am.* ii, 162, t. 183.—Antoine, *Conif.* 84, t. 33, f. 3.—Hooker & Arnott, *Bot. Beechey*, 394.—Endlicher, *Syn. Conif.* 87.—Torrey in *Sitgreaves' Rep.* 173.—Parlatore in *De Candolle, Prodr.* xvi³, 430.—McNab in *Proc. Royal Irish Acad.* 2 ser. ii, 703, t. 49, f. 32, 32^a, 32^b.

Abies Douglasii, var. *taxifolia*, London, *Arboretum*, iv, 2319, f. 2231.—Gordon, *Pinetum*, 16; 2 ed. 25.—Henkel & Hochstetter, *Nadelhölz.* 156.

Pinus Douglasii, var. *brevibracteata*, Antoine, *Conif.* 84, t. 33, f. 4.

Picea Douglasii, Link in *Linnæa*, xv, 524.

Tsuga Douglasii, Carrière, *Trait. Conif.* 192.—Bolander in *Proc. California Acad.* iii, 232.

Tsuga Lindleyana, Roehl, *Cat. Grain Mex.* 8.

RED FIR. YELLOW FIR. OREGON PINE. DOUGLAS FIR.

Coast ranges and interior plateau of British Columbia south of latitude 55° N. (not reaching the coast archipelago north of Vancouver's island), east to the eastern slope of the Rocky mountains in latitude 51° N. (Bow River pass, *Macoun*); south along the mountain ranges of Washington territory, Oregon, the California Coast ranges, and the western slope of the Sierra Nevadas, through the mountain ranges east to Montana, Wyoming, Colorado, and the Guadalupe mountains of Texas; in the Wahsatch and Uintah mountains, the ranges of northern and eastern Arizona, and southward into Mexico; not detected in the interior region between the Sierra Nevada and the Wahsatch mountains, south of the Blue mountains of Oregon, and north of Arizona.

A large tree, 61 to 92 meters in height, with a trunk 0.83 to 3.66 meters in diameter, or in the Rocky mountains much smaller, here rarely 30 meters in height; the most generally-distributed and valuable timber tree of the Pacific region, growing from the sea-level to an elevation in Colorado of nearly 10,000 feet; often forming extensive forests, almost to the exclusion of other species, and reaching in western Oregon and Washington territory its greatest development and value.

Wood hard, strong, varying greatly with age and conditions of growth in density, quality, and amount of sap; difficult to work, durable; bands of small summer cells broad, occupying fully half the width of the annual growth, dark colored, conspicuous, soon becoming flinty and difficult to cut; medullary rays numerous, obscure; color, varying from light red to yellow, the sap-wood nearly white; specific gravity, 0.5157; ash, 0.08; largely manufactured into lumber and used for all kinds of construction, railway ties, piles, fuel, etc.; two varieties, red and yellow fir, are distinguished by lumbermen, dependent probably upon the age of the tree; the former coarse-grained, darker colored, and considered less valuable than yellow fir.

The bark is found valuable in tanning leather.

Var. *macrocarpa*, Eng. lmann,

Bot. California, ii, 120.

Abies Douglasii, var. *macrocarpa*, Torrey in Ives' Rep. 23.—Vasey, Cat. Forest Trees, 33.

Abies macrocarpa, Vasey in Gard. Monthly, Jan. 1876.

HEMLOCK.

California Coast ranges; San Bernardino mountains to the Cuyamaca mountains.

A tree 30 to 54 meters in height, with a trunk 1.20 to 1.80 meter in diameter; dry ridges and cañons between 2,500 and 4,000 feet elevation.

Wood heavy, hard, strong, cross-grained, very durable, difficult to work; color, rather darker red than that of the species; specific gravity, 0.4563; ash, 0.08; somewhat manufactured into coarse lumber and largely used for fuel.

392.—*Abies Fraseri*, Lindley,

Penn. Cycl. i, 30.—Forbes, Pinetum Woburn. iii, t. 38.—Link in Linnæa, xv, 531.—Nuttall, Sylva, iii, 139, t. 119; 2 ed. ii, 196, t. 119.—Lindley & Gordon in Jour. Hort. Soc. London, v, 209.—Carrière, Trait. Conif. 200; 2 ed. 270.—Cooper in Smithsonian Rep. 1858, 257.—Chapman, Fl. S. States, 434.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 26.—Wood, Cl. Book, 661; Bot. & Fl. 314.—Henkel & Hochstetter, Nadelhölz. 169.—Gray, Manual N. States, 5 ed. 472, in part.—Hoopes, Evergreens, 202.—Bertrand in Bull. Soc. Bot. France, xviii, 379.—Koch, Dendrologie, ii³, 216.—Vasey, Cat. Forest Trees, 35.—Engelmann in Trans. St. Louis Acad. iii, 596; London Gard. Chronicle, 1877, 147.—Veitch, Manual Conif. 96.

Pinus Fraseri, Pursh, Fl. Am. Sept. ii, 639.—Smith in Rees' Cycl. xxviii, No. 27.—Poiret, Suppl. v, 35.—Sprengel, Syst. ii, 884.—Beck, Bot. 340.—Eaton, Manual, 6 ed. 264.—Lambert, Pinus, 1 ed. iii, 74, t. 42.—Eaton & Wright, Bot. 358.—Antoine, Conif. 76, t. 29, f. 1.—Endlicher, Syn. Conif. 91.—Parlatore in De Candolle, Prodr. xvi³, 419.—McNab in Proc. Royal Irish Acad. 2 ser. ii, 684, t. 47, f. 10.

A. balsamea, var. *Fraseri*, Nuttall, Genera, ii, 223.—Spach, Hist. Veg. xi, 422.

Pinus balsamea, var. *Fraseri*, Torrey, Compend. Fl. N. States, 359.

Picea Fraseri, Loudon, Arboretum, iv, 2340, f. 2243, 2244.—Knight, Syn. Conif. 39.—Gordon, Pinetum, 148; 2 ed. 205.

BALSAM. SHE BALSAM.

High mountains of North Carolina and Tennessee.

A tree 18 to 24 meters in height, with a trunk sometimes 0.60 meter in diameter; moist slopes between 5,000 and 6,500 feet elevation, often forming considerable forests.

Wood very light, soft, not strong, coarse-grained, compact; bands of small summer cells rather broad, light colored, not conspicuous; medullary rays numerous, thin; color, light brown, the sap-wood lighter, nearly white; specific gravity, 0.3565; ash, 0.54.

393.—*Abies balsamea*, Miller,

Dict. No. 5.—Desfontaines, Hist. Arb. ii, 579.—Nouveau Duhamel, v, 295, t. 83, f. 2.—Richard, Conif. 74, t. 16.—Lindley, Penn. Cycl. i, 30; Fl. Med. 554.—Forbes, Pinetum Woburn. 109, t. 37.—Link in Linnæa, xv, 530.—Spach, Hist. Veg. xi, 421.—Griffith, Med. Bot. 605, f. 268.—Lindley & Gordon in Jour. Hort. Soc. London, v, 210.—Carrière, Trait. Conif. 217; 2 ed. 292.—Richardson, Arctic Exped. 441.—Darlington, Fl. Cestrica, 3 ed. 291.—Cooper in Smithsonian Rep. 1858, 257.—Wood, Cl. Book, 661; Bot. & Fl. 314.—Porcher, Resources S. Forests, 506.—Henkel & Hochstetter, Nadelhölz. 176.—Gray, Manual N. States, 5 ed. 471.—Hoopes, Evergreens, 197.—Bertrand in Bull. Soc. Bot. France, xviii, 379.—Koch, Dendrologie, ii³, 214.—Vasey, Cat. Forest Trees, 34.—Guibourt, Hist. Drogues, 7 ed. ii, 246.—Engelmann in Trans. St. Louis Acad. iii, 597.—Macoun in Geological Rep. Canada, 1875-76, 211.—Sears in Bull. Essex Inst. xiii, 184.—Bell in Geological Rep. Canada, 1879-80, 46c.—Veitch, Manual Conif. 88.

Pinus balsamea, Linnæus, Spec. 1 ed. 1002.—Wangenheim, Amer. 40.—Aiton, Hort. Kew. iii, 370; 2 ed. v, 319.—Mönch, Meth. 364.—Du Roi, Harbk. 2 ed. 144.—Lambert, Pinus, 1 ed. i, 48, t. 31; 2 ed. i, 52, t. 33; 3 ed. i, 72, t. 41.—Willdenow, Spec. iv, 504; Enum. 989; Berl. Baumz. 276.—Persoon, Syn. ii, 579.—Pursh, Fl. Am. Sept. ii, 639.—Eaton, Manual, 111; 6 ed. 264.—Nuttall, Genera, ii, 223.—Hayne, Dend. Fl. 176.—Elliott, Sk. ii, 639.—Sprengel, Syst. ii, 884.—Torrey, Compend. Fl. N. States, 359; Fl. N. York, ii, 229.—Descourtiz, Fl. Med. Antilles, iv, 59, t. 246.—Woodville, Med. Bot. 3 ed. v, 1, t. 1.—Beck, Bot. 340.—Hooker, Fl. Bor.-Am. ii, 163.—Eaton & Wright, Bot. 358.—Bigelow, Fl. Boston. 3 ed. 385.—Antoine, Conif. 66, t. 26, f. 3.—Endlicher, Syn. Conif. 103.—Gihoul, Arb. Resin. 45.—Darby, Bot. S. States, 515.—Parlatore in De Candolle, Prodr. xvi³, 423.—McNab in Proc. Royal Irish Acad. 2 ser. ii, 684, t. 47, f. 11.—Bentley & Trimen, Med. Fl. iv, 263, t. 263.

Pinus Abies Balsamea, Marshall, Arbustum, 102.

A. balsamifera, Michaux, Fl. Bor.-Am. ii, 207, in part.—Michaux f. Hist. Arb. Am. i, 145, t. 14; N. American Sylva, 3 ed. iii, 150, t. 150, in part.

Picea balsamea, Loudon, Arboretum, iv, 2339, f. 2240, 2241.—Knight, Syn. Conif. 39.—Gordon, Pinetum, 143; 2 ed. 200.—Henkel & Hochstetter, Nadelhölz. 176.—Emerson, Trees Massachusetts, 85; 2 ed. i, 101.—Nelson, Pinaceae, 37.

Picea balsamea, var. *longifolia*, Hort.—London, Arboretum, iv, 2839.

Picea Fraseri, Emerson, Trees Massachusetts, 88; 2 ed. i, 104 [not London].

BALSAM FIR. BALM OF GILEAD FIR.

Northern Newfoundland and Labrador to the southern shores of Hudson bay, northwest to the Great Bear lake and the eastern base of the Rocky mountains; south through the northern states to Pennsylvania, central Michigan and Minnesota, and along the Alleghany mountains to the high peaks of Virginia.

A tree 21 to 27 meters in height, with a trunk rarely exceeding 0.60 meter in diameter, or at high elevations reduced to a low, prostrate shrub (*A. Hudsonica*, Hort.); damp woods and mountain swamps.

Wood very light, soft, not strong, coarse-grained, compact, not durable; bands of small summer cells not broad, resinous, conspicuous; medullary rays numerous, obscure; color, light brown, often streaked with yellow, the sap-wood lighter; specific gravity, 0.3819; ash, 0.45.

Canadian balsam or balm of fir, an aromatic liquid oleo-resin obtained from this and other species of *Abies* by puncturing the vesicles formed under the bark of the stem and branches, is used medicinally, chiefly in the treatment of chronic catarrhal affections, and in the arts (*U. S. Dispensatory*, 14 ed. 898, 900.—*Nat. Dispensatory*, 2 ed. 1417.—*Flückiger & Hanbury, Pharmacographia*, 552).

394.—*Abies subalpina*, Engelmann,

Am. Nat. x, 554; Trans. St. Louis Acad. iii, 597; Wheeler's Rep. vi, 255.—Vasey, Cat. Forest Trees, 34.—Hall in Coulter's Bot. Gazette, ii, 91.—Brandegge in Coulter's Bot. Gazette, iii, 32.—G. M. Dawson in Canadian Nat. newser. ix, 326.—Masters in London Gard. Chronicle, 1881, 236, f. 43, 44, 45.

?*Pinus lasiocarpa*, Hooker, Fl. Bor.-Am. ii, 163 [not Hort.].—Endlicher, Syn. Conif. 105.—McNab in Proc. Royal Irish Acad. 2 ser. ii, 682, t. 46, f. 7, 7^a; t. 47, 48, 49 (excl. syn.).

?*A. lasiocarpa*, Nuttall, Sylva, iii, 138; 2 ed. ii, 195.—Lindley & Gordon in Jour. Hort. Soc. London, v, 210.—Carrière, Trait. Conif. 1 ed. 221.—Cooper in Smithsonian Rep. 1858, 262.—Murray in Proc. Hort. Soc. London, iii, 313, f. 27-31.—Henkel & Hochstetter, Nadelhölz. 161 (excl. syn.).

?*Pinus* species, Torrey in Fremont's Rep. 97.

Picea amabilis, Gordon, Pinetum, 154, in part; 2 ed. 213, in part.

A. bifolia, Murray in Proc. Hort. Soc. London, iii, 320, f. 51-56; London Gard. Chronicle, 1875, 465, f. 96, 97.—Regel, Gartenflora, xiii, 119.—Henkel & Hochstetter, Nadelhölz. 420.

A. grandis, Engelmann in Am. Jour. Sci. 2 ser. xxxiv, 310 [not Lindley].—Carrière, Trait. Conif. 2 ed. 296, in part.—Watson in King's Rep. v, 334, in part.—Gray in Proc. Am. Acad. vii, 402 [not Lindley].—Porter & Coulter, Fl. Colorado; Hayden's Surv. Misc. Pub. No. 4, 131 [not Lindley].

Pinus amabilis, Parlatores in De Candolle, Prodr. xvi², 426, in part.

Picea bifolia, Murray in London Gard. Chronicle, 1875, 105.

A. subalpina, var. *fallax*, Engelmann in Trans. St. Louis Acad. iii, 597.

BALSAM.

Valley of the Stakhin river, Alaska, in latitude 60° N. (*Muir*), south through British Columbia and along the Cascade mountains to northern Oregon (*Collier*), through the Blue mountains of Oregon and the ranges of Idaho, Montana, Wyoming, Utah, and Colorado.

A tree 24 to 40 meters in height, with a trunk rarely exceeding 0.60 meter in diameter; mountain slopes and cañons between 4,000 (British Columbia) and 12,000 (Colorado) feet elevation; generally scattered and rarely forming the prevailing forest growth.

Wood very light, soft, not strong, rather close-grained, compact; bands of small summer cells very narrow, not conspicuous; medullary rays numerous, obscure; color, light brown or nearly white, the sap-wood lighter; specific gravity, 0.3476; ash, 0.44.

395.—*Abies grandis*, Lindley,

Penn. Cycl. i, 30.—Forbes, *Pinetum Woburn*, 123, t. 43.—Spach, *Hist. Veg.* xi, 422.—Nuttall, *Sylva*, iii, 134; 2 ed. ii, 192.—Lindley & Gordon in *Jour. Hort. Soc. London*, v, 210.—Carrière, *Trait. Conif.* 220; 2 ed. 296 (excl. syn.).—Cooper in *Smithsonian Rep.* 1858, 262; *Pacific R. R. Rep.* xii^a, 25, 69; *Am. Nat.* iii, 410.—Wood, *Bot. & Fl.* 314.—Lyll in *Jour. Linnean Soc.* vii, 143.—Bolander in *Proc. California Acad.* iii, 232.—Henkel & Hochstetter, *Nadelhölz.* 160.—Nelson, *Pinaceæ*, 38.—Hoopes, *Evergreens*, 211.—Bertrand in *Bull. Soc. Bot. France*, xviii, 378.—Vasey, *Cat. Forest Trees*, 34.—Hall in *Coulter's Bot. Gazette*, ii, 91.—Macoun in *Geological Rep. Canada*, 1875-76, 211.—Engelmann in *Trans. St. Louis Acad.* iii, 593; *London Gard. Chronicle*, 1879, 684; 1880, 660, f. 119; *Bot. California*, ii, 118.—G. M. Dawson in *Canadian Nat. new ser.* ix, 326.—Masters in *London Gard. Chronicle*, 1881, 179, f. 33-36.—Veitch, *Manual Conif.* 97, f. 23, 24.

Pinus grandis, Douglas in *Companion Bot. Mag.* ii, 147.—Hooker, *Fl. Bor.-Am.* ii, 163.—Antoine, *Conif.* 63, t. 25, f. 1.—Hooker & Arnott, *Bot. Beechey*, 394.—Endlicher, *Syn. Conif.* 105.—Parlatore in *De Candolle, Prodr.* xvi^a, 427 (excl. syn.).—McNab in *Proc. Royal Irish Acad.* 2 ser. ii, 678, t. 46, f. 4, 4^a.

?*A. aromatica*, Rafinesque, *Atlant. Jour.* 119.—Endlicher, *Syn. Conif.* 125.—Lindley & Gordon in *Jour. Hort. Soc. London*, v, 213.—Carrière, *Trait. Conif.* 266; 2 ed. 310.

Picea grandis, London, *Arboretum*, iv, 2341, f. 2245, 2246, in part.—Knight, *Syn. Conif.* 39.—Gordon, *Pinetum*, 155; *Suppl.* 5 (excl. syn. *Parsonsii*); 2 ed. 216.—Newberry in *Pacific R. R. Rep.* vi, 46, 90, f. 16, t. 6, in part.—Murray in *London Gard. Chronicle*, 1875, 135, f. 23.

A. Gordoniana, Carrière, *Trait. Conif.* 2 ed. 298 (excl. syn. *Parsonsii*).—Bertrand in *Bull. Soc. Bot. France*, xviii, 379.

A. amabilis, Murray in *Proc. Hort. Soc. London*, iii, 310, f. 22-24 [not Forbes].

WHITE FIR.

Vancouver's island, south to Mendocino county, California, near the coast; interior valleys of western Washington territory and Oregon south to the Umpqua river, Cascade mountains below 4,000 feet elevation, through the Blue mountains of Oregon (*Cusick*) to the eastern slope of the Cœur d'Alène mountains (*Cooper*), the Bitter Root mountains, Idaho (*Watson*), and the western slopes of the Rocky mountains of northern Montana (Flathead region, *Canby & Sargent*).

A large tree, 61 to 92 meters in height, with a trunk 0.90 to 1.50 meter in diameter; most common and reaching its greatest development in the bottom lands of western Washington territory and Oregon in rich, moist soil; or moist mountain slopes, then much smaller, rarely exceeding 30 meters in height.

Wood very light, soft, not strong, coarse-grained, compact; bands of small summer cells broader than in other American species, dark colored, resinous, conspicuous; medullary rays numerous, obscure; color, light brown, the sap-wood rather lighter; specific gravity, 0.3545; ash, 0.49; in western Oregon manufactured into lumber and used for interior finish, packing-cases, cooperage, etc.

396.—*Abies concolor*, Lindley & Gordon,

Jour. Hort. Soc. London, v, 210.—Parry in *Am. Nat.* ix, 204.—Vasey, *Cat. Forest Trees*, 34.—Engelmann in *Trans. St. Louis Acad.* iii, 600; *Wheeler's Rep.* vi, 255; *London Gard. Chronicle*, 1879, 684, f. 114, 115; *Bot. California*, ii, 118.—Brandegge in *Coulter's Bot. Gazette*, iii, 32.—Masters in *London Gard. Chronicle*, 1879, 684, f. 114, 115.—Veitch, *Manual Conif.* 93.

Pinus concolor, Engelmann in *herb.*; Parlatore in *De Candolle, Prodr.* xvi^a, 426.—McNab in *Proc. Royal Irish Acad.* 2 ser. ii, 681, t. 46, f. 6.

Picea concolor, Gordon, *Pinetum*, 155; 2 ed. 216.—Murray in *London Gard. Chronicle*, 1875, 135, f. 26.

Pinus lasiocarpa, Balfour in *Rep. Oregon Exped.* i, t. 4, f. 1 [not Hooker].—Murray in *Proc. Hort. Soc. London*, iii, 314, f. 25.—Henkel & Hochstetter, *Nadelhölz.* 429.

?*A. balsamea*, Bigelow in *Pacific R. R. Rep.* iv, 18 [not Miller].—Torrey in *Pacific R. R. Rep.* iv, 141.

Picea grandis, Newberry in *Pacific R. R. Rep.* vi, 46, in part.

Abies grandis, Carrière, *Trait. Conif.*; 2 ed. 296, in part.—Watson in *Pl. Wheeler*, 17 [not Lindley].

Picea Lowiana, Gordon, *Pinetum*, *Suppl.* 53; 2 ed. 218.—Henkel & Hochstetter, *Nadelhölz.* 419.

A. Lowiana, Murray in *Proc. Hort. Soc. London*, iii, 317, f. 38-41.

A. amabilis, Watson in *King's Rep.* v, 333 [not Forbes].

A. grandis, var. *Lowiana*, Hoopes, *Evergreens*, 212.

Pinus grandis, Parlatore in *De Candolle, Prodr.* xvi^a, 427, in part.

Picea concolor, var. *violacea*, Murray in *London Gard. Chronicle*, 1875, 464, f. 94, 95.

Pinus Lowiana, McNab in *Proc. Royal Irish Acad.* 2 ser. ii, 680, t. 46, f. 5.

A. lasiocarpa, *Hort.* [not Nuttall].

A. Parsonsii, *Hort.*

WHITE FIR. BALSAM FIR.

Northern slopes of the Siskiyou mountains, Oregon, and perhaps farther north in the Cascade mountains, south along the western slope of the Sierra Nevadas to the San Bernardino and San Jacinto mountains, California; along the high mountains of northern Arizona to the Mogollon mountains, New Mexico, northward to the Pike's Peak region of Colorado, and in the Wahsatch mountains of Utah.

A large tree, 30 to 40 meters in height, with a trunk 1.20 to 1.50 meter in diameter; moist slopes and cañons between 3,000 and 9,000 feet elevation, reaching its greatest development in the California sierras, varying greatly in the color and length of leaves, habit, etc., and perhaps merely a southern form of the too nearly allied *A. grandis*, from which it cannot be always readily distinguished.

Wood very light, soft, not strong, coarse-grained, compact; bands of small summer cells narrow, resinous, not conspicuous; medullary rays, numerous, obscure; color, very light brown or nearly white, the sap-wood somewhat darker; specific gravity, 0.3638; ash, 0.85; occasionally manufactured into lumber and used for packing-cases, butter-tubs, and other domestic purposes.

397.—*Abies bracteata*, Nuttall,

Sylva, iii, 137, t. 118; 2 ed. ii, t. 118.—Hartweg in Jour. Hort. Soc. London, iii, 225.—Lindley & Gordon in Jour. Hort. Soc. London, v, 209.—Carrière, Trait. Conif. 193; 2 ed. 265.—London Gard. Chronicle, 1853, 435; 1854, 459; 1859, 928.—Bot. Mag. t. 4740.—Lemaire in Ill. Hort. i, 14, t. 5.—Fl. des Serres, ix, 109 & t.—Naudin in Rev. Hort. 1854, 31.—Cooper in Smithsonian Rep. 1858, 262.—Murray in Edinburgh New Phil. Jour. new ser. x, 1, t. 1, 2 (Trans. Bot. Soc. Edinburgh, vi, 211, t. 1, 2).—Henkel & Hochstetter, Nadelhölz. 167.—Hoopes, Evergreens, 199.—Bertrand in Bull. Soc. Bot. France, xviii, 379.—Vasey, Cat. Forest Trees, 35.—Engelmann in Trans. St. Louis Acad. iii, 601; London Gard. Chronicle, 1879, 684; Bot. California, ii, 118.—Veitch, Manual Conif. 89, f. 14, 15.

Pinus venusta, Douglas in Companion Bot. Mag. ii, 152.

Pinus bracteata, D. Don in Trans. Linnæan Soc. xvii, 443.—Lambert, Pinus, 1 ed. iii, 169, t. 91.—Antoine, Conif. 77, t. 30.—Hooker & Arnott, Bot. Beechey, 394.—Hooker, Icon. t. 379.—Endlicher, Syn. Conif. 89.—Walpers, Ann. v, 798.—Parlatore in De Candolle, Prodr. xvi³, 419.—McNab in Proc. Royal Irish Acad. 2 ser. ii, 674, t. 46, f. 1.

Picea bracteata, London, Arboretum, iv, 2348, f. 2256.—Gordon, Pinetum, 145; 2 ed. 202.—Lawson, Pinetum Brit. ii, 171, t. 25, 26, f. 1-7.—Nelson, Pinaceæ, 37.—Fowler in London Gard. Chronicle, 1872, 286.

A. venusta, Koch, Dendrologie, ii³, 210.

Santa Lucia mountains, California, from the northern boundary of San Luis Obispo county about 40 miles northward.

A tree 46 to 61 meters in height, with a trunk 0.90 to 1.20 meter in diameter; moist, cold soil, occupying 4 or 5 cañons between 3,000 and 6,000 feet elevation, generally west of the summit of the range (*G. R. Vasey*).

Wood heavy, not hard, coarse-grained, compact; bands of small summer cells broad, resinous, conspicuous; medullary rays numerous, obscure; color, light brown tinged with yellow, the sap-wood not seen; specific gravity, 0.6783; ash, 2.04; probably more valuable than the wood of the other North American *Abies*.

398.—*Abies amabilis*, Forbes,

Pinetum Woburn. 125, t. 44.—Lindley & Gordon in Jour. Hort. Soc. London, v, 210.—Carrière, Trait. Conif. 219; 2 ed. 296.—Cooper in Smithsonian Rep. 1858, 262.—Lyll in Jour. Hort. Soc. London, vii, 143.—Henkel & Hochstetter, Nadelhölz. 159.—Nelson, Pinaceæ, 36.—Hoopes, Evergreens, 209 (excl. syn. *lasiocarpa*).—Fowler in London Gard. Chronicle, 1872, 285.—Koch, Dendrologie, ii³, 211 (excl. syn. *lasiocarpa*).—Macoun in Geological Rep. Canada, 1875-76, 211.—Engelmann in London Gard. Chronicle, 1880, 720, f. 136-141; Coulter's Bot. Gazette, vii, 4.—Veitch, Manual Conif. 86.

Pinus amabilis, Douglas in Companion Bot. Mag. ii, 93.—Antoine, Conif. 63, t. 25, f. 2.—Hooker & Arnott, Bot. Beechey, 394.—Endlicher, Syn. Conif. 104.—Parlatore in De Candolle, Prodr. xvi³, 426, in part.

Pinus grandis, Lambert, Pinus, 1 ed. iii, t. 26 [not Douglas].

Picea amabilis, London, Arboretum, iv, 2342, f. 2247, 2248.—Knight, Syn. Conif. 39.—Gordon, Pinetum, 154; 2 ed. 213 (excl. syn.).—Newberry in Pacific R. R. Rep. vi, 51, 90, f. 18.

A. grandis, Murray in Proc. Hort. Soc. London, iii, 308, f. 18-21 [not Lindley].

A. grandis, var. *densiflora*, Engelmann in Trans. St. Louis Acad. iv, 599.

Valley of the Fraser river, British Columbia (*Engelmann & Sargent*), and probably farther north, south along the Cascade mountains of Washington territory and Oregon.

A tree 30 to 45 meters in height, with a trunk sometimes 1.20 meter in diameter, forming extensive forests on the mountains of British Columbia, between 3,500 and 5,000 feet, and upon the mountains south of the Columbia river between 3,000 and 4,000 feet elevation, here reaching its greatest development; its northern range not yet determined.

Wood light, not strong, close-grained, compact; bands of small summer cells broad, resinous, dark colored, conspicuous; medullary rays numerous, thin; color, light brown, the sap-wood nearly white; specific gravity, 0.4228; ash, 0.23.

399.—*Abies nobilis*, Lindley.

Penn. Cycl. i, 30.—Forbes, Pinetum Woburn. 115, t. 40.—Link in Linnæa, xv, 532.—Spach, Hist. Veg. xi, 419.—Nuttall, Sylva, iii, 136, t. 117; 2 ed. ii, 193, t. 117.—Lindley & Gordon in Jour. Hort. Soc. London, v, 209.—Carrière, Trait. Conif. 198; 2 ed. 268.—Jour. Bot. & Kew Gard. Misc. ix, 85.—Cooper in Smithsonian Rep. 1858, 262.—Henkel & Hochstetter, Nadelhölz. 168.—Hoopes, Evergreens, 203.—Koch, Dendrologie, ii^a, 209.—Vasey, Cat. Forest Trees, 34.—Engelmann in Trans. St. Louis Acad. iii, 601, in part; London Gard. Chronicle, 1879, 885; Bot. California, ii, 119, in part; Coulter's Bot. Gazette, vii, 4.—Veitch, Manual Conif. 101.

Pinus nobilis, Douglas in Companion Bot. Mag. ii, 147.—Lambert, Pinus, 1 ed. iii, 167, t. 74.—Hooker, Fl. Bor.-Am. ii, 162.—Antoine, Conif. 77, t. 29, f. 2.—Hooker & Arnott, Bot. Beechey, 394.—Endlicher, Syn. Conif. 90.

Picea nobilis, London, Arboretum, iv, 2342, f. 2249, 2250.—Knight, Syn. Conif. 39.—Lindley & Gordon in Jour. Hort. Soc. London, v, 209.—Gordon, Pinetum, 149; Suppl. 48; 2 ed. 207.—Newberry in Pacific R. R. Rep. vi, 49, 90, f. 17.—Lawson, Pinetum, Brit. ii, 181, t. 28, 29, f. 1-18.—Nelson, Pinaceæ, 39.

Pseudotsuga nobilis, Bertrand in Bull. Soc. Bot. France, xviii, 86.—McNab in Proc. Royal Irish Acad. 2 ser. ii, 699, t. 49, f. 29, 29^a.

A. magnifica, Engelmann in Bot. California, ii, 119, in part.

RED FIR.

Oregon, Cascade mountains from the Columbia river south to the valley of the upper Rogue river, and along the summits of the Coast Range from the Columbia to the Nestucca river (*Collier*).

A large tree, 61 to 92 meters in height, with a trunk 2.40 to 3 meters in diameter, forming, with *A. amabilis*, extensive forests along the slopes of the Cascade Range, between 3,000 and 4,000 feet elevation; less multiplied in the coast ranges, here reaching its greatest individual development.

Wood light, hard, strong, rather close grained, compact; bands of small summer cells broad, resinous, dark colored, conspicuous; medullary rays thin, hardly distinguishable; color, light brown streaked with red, the sap-wood a little darker; specific gravity, 0.4561; ash, 0.34.

400.—*Abies magnifica*, Murray,

Proc. Hort. Soc. London, iii, 318, f. 42-50; London Gard. Chronicle, 1875, 134.—Regel, Gartenflora, xiii, 119.—Henkel & Hochstetter, Nadelhölz. 419.—Koch, Dendrologie, ii^a, 213.—Engelmann in Trans. St. Louis Acad. iii, 601; London Gard. Chronicle, 1879, 885, f. 116; Bot. California, ii, 119; Coulter's Bot. Gazette, vii, 4.—Veitch, Manual Conif. 99.

A. campylocarpa, Murray in Trans. Bot. Soc. Edinburgh, vi, 370.

A. nobilis robusta, Hort.—Carrière, Trait. Conif. 2 ed. 269.

Picea magnifica, Gordon, Pinetum, 2 ed. 219.—Murray in London Gard. Chronicle, 1875, 105.

Pinus amabilis, Parlatores in De Candolle, Prodr. xvi^a, 426, in part.—McNab in Proc. Royal Irish Acad. 2 ser. ii, 677, t. 46, f. 3, 3^a?

A. amabilis, Vasey, Cat. Forest Trees, 34 [not Forbes].

Pseudotsuga magnifica, McNab in Proc. Royal Irish Acad. 2 ser. ii, 700, t. 49, f. 30, 30^a.

A. nobilis, Engelmann, Bot. California, ii, 119, in part.

RED FIR.

California, mount Shasta, south along the western slope of the Sierra Nevadas to Kern county.

A large tree, 61 to 76 meters in height, with a trunk 2.40 to 3 meters in diameter, forming about the base of mount Shasta extensive forests between 4,900 and 8,000 feet elevation; farther south less common and reaching an extreme elevation of 10,000 feet.

Wood light, soft, not strong, rather close-grained, compact, satiny, durable in contact with the soil, liable to twist and warp in seasoning; bands of small summer cells broad, resinous, dark colored, conspicuous; medullary rays numerous, thin; color, light red, the sap-wood somewhat darker; specific gravity, 0.4701; ash, 0.30; largely used for fuel and occasionally manufactured into coarse lumber.

401.—*Larix Americana*, Michaux,

Fl. Bor.-Am. ii, 203.—Michaux f. Hist. Arb. Am. iii, 37, t. 4; N. American Sylva, 3 ed. iii, 167, t. 153.—Audubon, Birds, t. 4.—London, Arboretum, iv, 2399.—Emerson, Trees Massachusetts, 89; 2 ed. i, 105 & t.—Gihoul, Arb. Resin. 51.—Parry in Owen's Rep. 618.—Richardson, Arctic. Exped. 442.—Cooper in Smithsonian Rep. 1858, 257.—Hooker f. in Trans. Linnæan Soc. xxiii, 302.—Wood, Cl. Book, 662; Bot. & Fl. 314.—Nelson, Pinaceæ, 86.—Gray, Manual N. States, 5 ed. 442.—Hoopes, Evergreens, 247.—Regel, Gartenflora, xx, 105, t. 684, f. 7, 8 (Belg. Hort. xxii, 105, t. 10, f. 2, 3).—Bertrand in Ann. Sci. Nat. 5 ser. xx, 90.—Vasey, Cat. Forest Trees, 35.—Macoun in Geological Rep. Canada, 1875-76, 211.—Sears in Bull. Essex Inst. xiii, 185.

Pinus laricina, Du Roi, Obs. Bot. 49; Harbk. ii, 83.—Wangenheim, Amer. 42, t. 16, f. 37.—Mœnch, Meth. 364.

Pinus Larix rubra, alba and nigra, Marshall, Arbustum, 103, 104.

Pinus intermedia, Wangenheim, Amer. 42, t. 16, f. 37.—Du Roi, Harbk. 2 ed. ii, 114.

Pinus pendula, Aiton, Hort. Kew, iii, 369; 2 ed. v, 320.—Lambert, Pinus, 1 ed. i, 55, t. 36; 2 ed. ii, 63, t. 39; 3 ed. ii, 86, t. 49.—Willdenow, Spec. iv, 502.—Persoon, Syn. ii, 579.—Pursh, Fl. Am. Sept. ii, 645.—Smith in Rees' Cycl. xxviii, No. 32.—Eaton, Manual, 110; 6 ed. 365.—Nuttall, Genera, ii, 223.—Sprengel, Syst. ii, 887.—Audubon, Birds, t. 90, 180.—Beck, Bot. 339.—Hooker, Fl. Bor.-Am. ii, 164.—Eaton & Wright, Bot. 359.—Torrey, Fl. N. York, ii, 232.—Parlatore in De Candolle, Prodr. xvi*, 409.

Pinus microcarpa, Lambert, Pinus, 1 ed. i, 56, t. 37; 2 ed. ii, 65, t. 40; 3 ed. ii, 88, t. 50.—Willdenow, Spec. iv, 502; Enum. 989; Berl. Baumz. 273.—Persoon, Syn. ii, 579.—Aiton, Hort. Kew. 2 ed. v, 321.—Pursh, Fl. Am. Sept. ii, 645.—Smith in Rees' Cycl. xxviii, No. 33.—Eaton, Manual, 110; 6 ed. 365.—Nuttall, Genera, ii, 223.—Hayne, Dend. Fl. 175.—Sprengel, Syst. ii, 887.—Torrey, Compend. Fl. N. States, 360.—Meyer, Pl. Labrador, 30.—Beck, Bot. 340.—Hooker, Fl. Bor.-Am. ii, 164.—Eaton & Wright, Bot. 359.—Bigelow, Fl. Boston. 3 ed. 387.—Antoine, Conif. 54, t. 21, f. 1.—Endlicher, Syn. Conif. 132.

Abies pendula, Poiret in Lamarck, Dict. vi, 514.—Nouveau Duhamel, v, 288.—Lindley & Gordon in Jour. Hort. Soc. London, v, 213.

Abies microcarpa, Poiret in Lamarck, Dict. vi, 514.—Nouveau Duhamel, v, 289, t. 79, f. 2.—Lindley in Penn. Cycl. i, 33.—Lindley & Gordon in Jour. Hort. Soc. London, 213.

L. tenuifolia, Salisbury in Trans. Linnæan Soc. viii, 313.

L. pendula, Salisbury in Trans. Linnæan Soc. viii, 313.—Forbes, Pinetum Woburn. 137, t. 46.—Carrière, Trait. Conif. 1 ed. 272.—Gordon, Pinetum, 129; 2 ed. 177.—Hooker f. in Trans. Linnæan Soc. xxiii, 302.

L. microcarpa, Desfontaines, Hist. Arb. ii, 597.—Forbes, Pinetum Woburn. 139, t. 47.—Spach, Hist. Veg. xi, 436.—Link in Linnæa, xv, 536.—Carrière, Trait. Conif. 275; 2 ed. 355.—Gordon, Pinetum, 129; 2 ed. 175.—Henkel & Hochstetter, Nadelhölz. 137.—Hooker f. in Trans. Linnæan Soc. xxiii, 302, 341.—Veitch, Manual Conif. 180.

L. intermedia, Loddiges, Cat. ed. 1836, 50.—Forbes, Pinetum Woburn. 141.—Link in Linnæa, xv, 535.

L. Americana rubra, London, Arboretum, iv, 2400.—Knight, Syn. Conif. 40.

L. Americana, var. *pendula*, London, Arboretum, iv, 2400.—Carrière, Trait. Conif. 2 ed. 356.

L. Americana, var. *prolifera*, London, Arboretum, iv, 2401.—Carrière, Trait. Conif. 2 ed. 356.

L. decidua, var. *Americana*, Henkel & Hochstetter, Nadelhölz. 133.

LARCH. BLACK LARCH. TAMARACK. HACKMATAACK.

Northern Newfoundland and Labrador to the eastern shores of Hudson bay, cape Churchill and northwest to the northern shores of the Great Bear lake and the valley of the Mackenzie river within the Arctic circle; south through the northern states to northern Pennsylvania, northern Indiana and Illinois, and central Minnesota.

A tree 24 to 30 meters in height, with a trunk 0.60 to 0.90 meter in diameter; moist uplands and intervalle lands, or south of the boundary of the United States in cold, wet swamps, often covering extensive areas, here much smaller and less valuable.

Wood heavy, hard, very strong, rather coarse-grained, compact, durable in contact with the soil; bands of small summer cells broad, very resinous, dark colored, conspicuous, resin passages few, obscure; medullary rays numerous, hardly distinguishable, color, light brown, the sap-wood nearly white; specific gravity, 0.6236; ash, 0.33; preferred and largely used for the upper knees of vessels, for ship timbers, fence posts, telegraph poles, railway ties, etc.

The inner bark of the closely-allied European larch is recommended in the treatment of chronic catarrhal affections of the pulmonary and urinary passages; probably that of the American species would be equally efficacious.

402.—*Larix occidentalis*, Nuttall,

Sylva, iii, 143, t. 120; 2 ed. ii, 199, t. 120.—Newberry in Pacific R. R. Rep. vi, 59, f. 24, 25.—Cooper in Smithsonian Rep. 1858, 263; Am. Nat. iii, 412.—Lyll in Jour. Linnæan Soc. vii, 143.—Nelson, Pinacæ, 91.—Hoopes, Evergreens, 253.—Regel, Gartenflora, xx, 103, t. 685, f. 8-10 (Belg. Hort. xxii, 101, t. 8, f. 3-5).—Vasey, Cat. Forest Trees, 35.—Gordon, Pinetum, 2 ed. 176.—Macoun in Geological Rep. Canada, 1875-'76, 211.—G. M. Dawson in Canadian Nat. new ser. ix, 329.—Veitch, Manual Conif. 130.

Pinus Larix, Douglas in Companion Bot. Mag. ii, 109 [not Linnæus].

L. Americana, var. *brevifolia*, Carrière, Trait. Conif. 2 ed. 357.

Pinus Nuttallii, Parlatores in De Candolle, Prodr. xvi⁹, 412.

TAMARACK.

British Columbia, Selkirk and Gold ranges, south of latitude 53° N., extending west to the head of Okanagan lake (*G. M. Dawson*), south along the eastern slopes of the Cascade mountains to the Columbia river, through the mountain ranges of northern Washington territory to the western slopes of the Rocky mountains of Montana, and in the Blue mountains of Washington territory and Oregon.

A noble tree of great economic value, 30 to 45 meters in height, with a trunk 0.90 to 1.50 meter in diameter; moist mountain slopes and benches between 2,500 and 5,000 feet elevation; scattered among other trees and never exclusively forming forests; the thick bark long resisting the action of forest fires; very common, and perhaps reaching its greatest development in the region north of the Big Blackfoot river and in the valley of the Flathead river, Montana, here the largest and most valuable timber tree.

Wood heavy, exceedingly hard and strong, rather coarse-grained, compact, satiny, susceptible of a fine polish, very durable in contact with the soil; bands of small summer cells broad, occupying fully half the width of annual growth, very resinous, dark colored, conspicuous, resin passages few, obscure; medullary rays numerous, thin; color, light bright red, the thin sap-wood nearly white; specific gravity, 0.7407; ash, 0.09; occasionally manufactured into lumber, but principally used for fuel, posts, railway ties, etc.

403.—*Larix Lyallii*, Parlatores,

Enum. Sem. Hort. Reg. Mus. Flor. 1863; London Gard. Chronicle, 1863, 916 (Regel, Gartenflora, xiii, 244).—Lyll in Jour. Linnæan Soc. vii, 143.—Henkel & Hochstetter, Nadelhölz. 417.—Carrière, Trait. Conif. 2 ed. 361.—Hoopes, Evergreens, 256.—Regel, Gartenflora, xx, 103, t. 685, f. 11-13 (Belg. Hort. xxii, 102, t. 9, f. 1-3).—Bertrand in Ann. Sci. Nat. 5 ser. xx, 90.—Vasey, Cat. Forest Trees, 35.—Macoun in Geological Rep. Canada, 1875-'76, 211.—Veitch, Manual Conif. 130.

Pinus Lyallii, Parlatores in De Candolle, Prodr. xvi⁹, 412.

"Cascade mountains, 6,500 to 7,000 feet, forming an open belt of trees mingled with *P. flexilis* (*P. albicaulis*); on the Galton range at 6,000 feet and in the Rocky mountains at 7,000 feet, growing with *P. flexilis*" (*Lyall*); mount Stewart, Washington territory (*Brandege & Tweedy*, August, 1883); Grave Creek pass, northern Montana (*H. B. Ayres*, September, 1883).

A low, much-branched, straggling, alpine tree, rarely exceeding 15 meters in height, with a trunk sometimes 1.50 meter in diameter; dry, rocky soil, generally upon northern exposures, and associated with *Pinus albicaulis* and *Tsuga Pattoniana* along the upper limits of tree-growth between 5,500 and 7,000 feet elevation (*Brandeges*).

The wood not collected.

NOTE.—A well-marked species, distinguished from *L. occidentalis* by its alpine habit, the larger green or purple deciduous cones with ciliated scales, and by the dense tomentum covering the young shoots and leaf buds.

PALMACEÆ.

404.—*Sabal Palmetto*, Loddiges;

Römer & Schultes, Syst. vii, 1487.—Croom in Am. Jour. Sci. 1 ser. xxvi, 315.—Martius, Hist. Palm. iii, 247.—Kunth, Enum. iii, 247.—Spach, Hist. Veg. xii, 107.—Chapman, Fl. S. States, 438.—Curtis in Rep. Geological Surv. N. Carolina, 1860, iii, 64.—Wood, Cl. Book, 666; Bot. & Fl. 317.—Vasey, Cat. Forest Trees, 38.

Corypha Palmetto, Walter, Fl. Caroliniana, 119.

Chamærops Palmetto, Michaux, Fl. Bor.-Am. i, 206.—Michaux f. Hist. Arb.-Am. ii, 186, t. 10; N. American Sylva, 3 ed. iii, 5, t. 101.—Aiton, Hort. Kew. 2 ed. v, 490.—Nuttall, Genera, i, 231.—Elliott, Sk. i, 431.—Sprengel, Syst. ii, 137.—Eaton, Manual, 6 ed. 89.—Eaton & Wright, Bot. 191.—Darby, Bot. S. States, 546.—Cooper in Smithsonian Rep. 258.—Porcher, Resources, S. Forests, 526.

CABBAGE TREE. CABBAGE PALMETTO.

Smith island, off the mouth of Cape Fear river, North Carolina, south along the coast to Key Largo, Florida, and along the Gulf coast to the Apalachicola river.

A tree 7 to 12 meters in height, with a trunk 0.60 to 0.90 meter in diameter; sandy maritime shores; very common and reaching its greatest development upon the west coast of the Florida peninsula south of Cedar Keys.

Wood light, soft; fibro-vascular bundles hard, difficult to work, dark colored; color, light brown; specific gravity, 0.4404; ash, 7.66; impervious to the attacks of the teredo, and very durable under water; largely used for piles, wharves, etc.

405.—*Washingtonia filifera*, Wendland,

Bot. Zeit. xxxvii, 68.—Watson, Bot. California, ii, 211, 485.

Brahea dulcis,? Cooper in Smithsonian Rep. 1860, 442 [not Martius].

Pritchardia filamentosa, Wendland in Bot. Zeit. xxxiv, 807.—Vasey, Cat. Forest Trees, 38.—Fenzl in Bull. Soc. Toscana, i, 116 & f.—Palmer in Am. Nat. xii, 598.

Brahea filamentosa, Hort.—Williams in London Gard. Chronicle, 1876, 80.

FAN-LEAF PALM.

San Bernardino county, California, from the eastern base of the San Bernardino mountains to the valley of the Colorado river.

A tree 12 to 18 meters in height, with a trunk 0.60 to 1.05 meter in diameter, forming groves of 250 to 500 plants in the depressions of the desert, in moist alkaline soil, or solitary and scattered near the heads of small ravines formed by water-courses; often stunted and greatly injured by fire.

Wood light, soft; fibro-vascular bundles hard, difficult to cut, dark colored, conspicuous; specific gravity 0.5173; ash, 1.89.

406.—*Thrinax parviflora*, Swartz,

Prodr. 57; Fl. Ind. Occ. i, 614.—Aiton, Hort. Kew. iii, 614; 2 ed. ii, 307.—Willdenow, Spec. ii, 202.—Persoon, Syn. i, 383.—Poiret in Lamarek, Dict. vii, 633.—Titford, Hort. Bot. Am. 112.—Sprengel, Syst. ii, 20.—Römer & Schultes, Syst. vii, 1300.—Martius, Hist. Palm. iii, 255, t. 103.—Kunth, Enum. iii, 253.—Dietrich, Syn. ii, 1091.—Walpers, Ann. v, 818.—Grisebach, Fl. British West Indies, 515.—Vasey, Cat. Forest Trees, 38.—Chapman in Coulter's Bot. Gazette, iii, 12; Fl. S. States, Suppl. 651.

T. Garberi, Chapman in Coulter's Bot. Gazette, iii, 12; Fl. S. States, Suppl. 651.

SILK-TOP PALMETTO.

Semi-tropical Florida, southern keys from Bahia Honda to Long's Key; in the West Indies.

A small tree, 9 meters in height with a trunk rarely exceeding 0.10 meter in diameter, or in pine-barren soil often low and stemless (*T. Garberi*).

Wood light, soft; fibro-vascular bundles small, hard, not conspicuous; color, light brown; specific gravity, 0.5991; ash, 3.99; the trunk used in making sponge- and turtle-crawls.

407.—*Thrinax argentea*, Loddiges;

Deafontaines, Cat. 3 ed. 31.—Römer & Schultes, Syst. vii, 1300.—Martius, Hist. Palm. iii, 256, t. 103, f. 3, t. 163.—Kunth, Enum. iii, 253.—Dietrich, Syn. ii, 1091.—Walpers, Ann. v, 818.—Grisebach, Fl. British West Indies, 515.—Chapman, Fl. S. States, Suppl. 651.

Palma argentea, Jacquin, Fragm. 38, No. 125, t. 43, f. 1.—Märter in Bom. Physik. Arbeiten. ii, 76.

SILVER-TOP PALMETTO. BRICKLEY THATCH. BEITTLE THATCH.

Semi-tropical Florida, on a nameless key 10 miles west of Key West, Elliott's Key, Key Largo, Piney Key, Boca Chica Key, Key West, Gordon Key, and on the small keys south and west of Bahia Honda Key (*Ourtiss*); in the West Indies.

A small tree, 7 to 9 meters in height, with a trunk 0.15 to 0.20 meter in diameter.

Wood light, soft; fibro-vascular bundles small, very numerous; interior of the trunk spongy, much lighter than the exterior; specific gravity, 0.7172; ash, 3.01; used for piles, the foliage in the manufacture of ropes, for thatch, etc.

408.—*Oreodoxa regia*, HBK.

Nov. Genera & Spec. i, 305.—Martius, Hist. Palm. iii, 168, t. 156, f. 3-5.—Richard, Fl. Cuba, 348.—Kunth, Enum. iii, 182.—Spach, Hist. Veg. xii, 68.—Ill. Hort. ii, 28 & t.—Walpers, Ann. v, 807.—Grisebach, Fl. British West Indies, 327.—London Gard. Chronicle, 1875, 302, f. 66.—Chapman, Fl. S. States, Suppl. 651.

Enocarpus regia, Sprengel, Syst. ii, 140.

O. oleracea, ? Cooper in Smithsonian Rep. 1860, 440.

ROYAL PALM.

Semi-tropical Florida, "Little and Big Palm hummocks," 15 and 25 miles east of cape Romano (*Ourtiss*), near the mouth of Little river, and on Elliott's Key; in the West Indies.

A tree 18 to 30 meters in height, with a trunk 0.60 meter in diameter; rich hummocks, often forming extensive groves; in Florida rare and local.

Wood heavy, hard; fibro-vascular bundles large, very dark, conspicuous; interior of the trunk spongy, much lighter than the exterior; color, brown; specific gravity, exterior of the trunk, 0.7982, interior, 0.2128; ash, 2.54.

LILIACEÆ.

409.—*Yucca canaliculata*, Hooker,

• Bot. Mag. t. 5201.—Baker in London Gard. Chronicle, 1870, 1217.—Engelmann in Trans. St. Louis Acad. iii, 43.

Y. Treculiana, Carrière in Rev. Hort. vii, 280.—Baker in London Gard. Chronicle, 1870, 828.—Engelmann in Trans. St. Louis Acad. iii, 41.—Vasey, Cat. Forest Trees, 38.—London Garden, xii, 328, t. 94.

SPANISH BAYONET.

Southern Texas, Matagorda bay, and from the Brazos and Guadalupe rivers south into Mexico.

A small tree, 5 to 8 meters in height, with a trunk 0.30 to 0.75 meter in diameter; dry, gravelly, arid soil.

Wood, like that of the whole genus, showing distinct marks of concentric arrangement, fibrous, spongy, heavy, difficult to cut and work; color, light brown; specific gravity, 0.6677; ash, 6.27.

The bitter, sweetish fruit cooked and eaten by the Mexicans; the root stock, as in the whole genus, saponaceous and largely used by the Mexicans as a substitute for soap.

410.—*Yucca brevifolia*, Engelmann,

King's Rep. v, 496; Trans. St. Louis Acad. iii, 47.—Parry in Am. Nat. ix, 141, 351.—Vasey, Cat. Forest Trees, 38.—Watson, Bot. California, ii, 164.

Y. Draconis, ? var. *arborescens*, Torrey in Pacific R. R. Rep. iv, 147.

THE JOSHUA. JOSHUA TREE.

Southwestern Utah, northwestern Arizona to southern Nevada, and the valley of the Mohave river, California.

A tree 6 to 12 meters in height, with a trunk 0.60 to 0.90 meter in diameter; dry, gravelly soil, forming upon the Mohave desert at 2,500 feet elevation an open, straggling forest.

Wood light, soft, spongy, difficult to work; color, very light brown or nearly white; specific gravity, 0.3737; ash, 4.00; occasionally manufactured into paper-pulp.

411.—*Yucca elata*, Engelmann,

Coulter's Bot. Gazette, vii, 17.

Y. angustifolia, var. *radiosa*, Engelmann in King's Rep. v, 496.

Y. angustifolia, var. *elata*, Engelmann in Trans. St. Louis Acad. iii, 50; Wheeler's Rep. vi, 270.

SPANISH BAYONET.

Western Texas to southern Arizona and Utah; southward into Mexico.

A small tree, 3 to 5 meters in height, with a trunk 0.20 to 0.25 meter in diameter; dry, gravelly *mesas*.

Wood light, soft, spongy; color, light brown or yellow; specific gravity, 0.4470; ash, 9.28.

412.—*Yucca baccata*, Torrey,

Bot. Mex. Boundary Survey, 221; Ives' Rep. 29.—Cooper in Smithsonian Rep. 1858, 266.—Baker in London Gard. Chronicle, 1870, 923.—

André in Ill. Hort. 3 ser. xx, 23, t. 115.—Gray, Hall's Pl. Texas, 23.—Engelmann in Trans. St. Louis Acad. iii, 44; King's Rep. v, 496;

Wheeler's Rep. vi, 270.—Loew in Wheeler's Rep. iii, 609.—Rothrock in Wheeler's Rep. vi, 52.—Watson, Bot. California, ii, 164.

Y. filamentosa, ? Wood in Proc. Philadelphia Acad. 1868, 167 [not Torrey].

SPANISH BAYONET. MEXICAN BANANA.

Western Texas, south of latitude 32° N., west through New Mexico to southern Colorado and San Diego county, California; southward into northern Mexico.

A tree 7 to 12 meters in height, with a trunk 0.60 meter in diameter, or often much smaller, and toward the northern limits of its range stemless; forming upon the plains of Presidio county, Texas, extensive open forests (*Havard*).

Wood light, soft, spongy, difficult to work; color, light brown; specific gravity, 0.4470; ash, 9.28.

The large juicy fruit edible and an important article of food to Mexicans and Indians; a strong coarse fiber, prepared by macerating the leaves in water, is manufactured into rope by the Mexicans.

INDEX TO CATALOGUE.

[NOTE.—In this index the names of families are indicated by "SMALL CAPS", of species by "Roman" type, and of synonyms by "Italics".]

A.		Page.		Page.
<i>Abies alba</i> (Picea alba).....	204	<i>Abies Marylandica</i>	203	
<i>Abies alba</i> ? (Picea Engelmanni).....	205	<i>Abies Menziesii</i> (Picea pungens).....	205	
<i>Abies alba</i> (Picea nigra).....	203	<i>Abies Menziesii</i> (Picea Sitchensis).....	206	
<i>Abies alba</i> , var. <i>arctica</i>	204	<i>Abies Menziesii</i> Parryana.....	205	
<i>Abies alba</i> , var. <i>cærulea</i>	204	<i>Abies Mertensiana</i>	207	
<i>Abies Albertiana</i>	208	<i>Abies microcarpa</i>	215	
<i>Abies amabilis</i>	213	<i>Abies mucronata</i>	209	
<i>Abies amabilis</i> (Abies concolor).....	212	<i>Abies mucronata palustris</i>	209	
<i>Abies amabilis</i> (Abies grandis).....	212	<i>Abies nigra</i> (Picea Engelmanni).....	205	
<i>Abies amabilis</i> (Abies magnifica).....	214	<i>Abies nigra</i> (Picea nigra).....	203	
<i>Abies Americana</i>	203	<i>Abies nigra</i> , var. <i>rubra</i>	203	
<i>Abies arctica</i> (Picea alba).....	204	<i>Abies nobilis</i>	214	
<i>Abies arctica</i> (Picea nigra).....	203	<i>Abies nobilis</i> (Abies magnifica).....	214	
<i>Abies aromatica</i>	212	<i>Abies nobilis robusta</i>	214	
<i>Abies balsamea</i>	210	<i>Abies Parsonii</i>	213	
<i>Abies balsamea</i> (Abies concolor).....	212	<i>Abies Pattonii</i>	208	
<i>Abies balsamea</i> , var. <i>Fraseri</i>	210	<i>Abies Pattoniana</i>	208	
<i>Abies balsamifera</i>	211	<i>Abies Pattonii</i> (Tsuga Mertensiana).....	208	
<i>Abies bifolia</i>	211	<i>Abies Pattonii</i> (Tsuga Pattoniana).....	208	
<i>Abies bracteata</i>	213	<i>Abies pendula</i>	215	
<i>Abies Bridgesii</i>	207	<i>Abies religiosa</i>	185	
<i>Abies campylocarpa</i>	214	<i>Abies rubra</i>	203	
<i>Abies Canadensis</i> (Picea alba).....	204	<i>Abies rubra</i> , var. <i>arctica</i>	203	
<i>Abies Canadensis</i> (Tsuga Canadensis).....	206	<i>Abies rubra</i> , var. <i>cærulea</i>	204	
<i>Abies Canadensis</i> ? (Tsuga Morteusiana).....	207	<i>Abies Sitchensis</i>	206	
<i>Abies Caroliniana</i>	207	<i>Abies speciosa</i> (Tsuga Caroliniana).....	207	
<i>Abies cærulea</i>	204	<i>Abies subalpina</i>	211	
<i>Abies concolor</i>	212	<i>Abies subalpina</i> , var. <i>fallax</i>	211	
<i>Abies denticulata</i>	203	<i>Abies taxifolia</i> (Pseudotsuga Douglasii).....	209	
<i>Abies Douglasii</i>	209	<i>Abies taxifolia</i> (Tsuga Mertensiana).....	208	
<i>Abies Douglasii</i> , var. <i>macrocarpa</i>	210	<i>Abies trigona</i>	206	
<i>Abies Douglasii</i> , var. <i>taxifolia</i>	209	<i>Abies venusta</i>	213	
<i>Abies Engelmanni</i>	205	<i>Abies Williamsonii</i>	208	
<i>Abies Engelmanni glauca</i>	205	<i>Acacia Bahamensis</i>	64	
<i>Abies falcata</i>	206	<i>Acacia Berlandieri</i>	63	
<i>Abies Fraseri</i>	210	<i>Acacia biocope</i>	62	
<i>Abies Gordoniana</i>	212	<i>Acacia esculenta</i>	63	
<i>Abies grandis</i>	212	<i>Acacia frondosa</i>	62	
<i>Abies grandis</i> (Abies amabilis).....	213	<i>Acacia glauca</i>	62	
<i>Abies grandis</i> (Abies concolor).....	212	<i>Acacia Green-bark</i>	60	
<i>Abies grandis</i> (Abies subalpina).....	211	<i>Acacia Greggii</i>	63	
<i>Abies grandis</i> , var. <i>densiflora</i>	213	<i>Acacia latistipula</i>	64	
<i>Abies grandis</i> , var. <i>Louisiana</i>	212	<i>Acacia leucocephala</i>	62	
<i>Abies heterophylla</i>	207	<i>Acacia pulverulenta</i>	63	
<i>Abies Hookeriana</i>	208	<i>Acacia tricholoba</i>	63	
<i>Abies Hudsonica</i>	211	<i>Acacia, Three-thorned</i>	59	
<i>Abies lasiocarpa</i> (Abies concolor).....	212	<i>Acacia Wrightii</i>	63	
<i>Abies lasiocarpa</i> (Abies subalpina).....	211	<i>Acer barbatum</i> (Acer glabrum).....	48	
<i>Abies laxa</i>	204	<i>Acer barbatum</i> (Acer saccharinum).....	48	
<i>Abies Louisiana</i>	212	<i>Acer Californicum</i>	51	
<i>Abies macrocarpa</i>	210	<i>Acer Canadense</i>	48	
<i>Abies magnifica</i>	214	<i>Acer Caroliniana</i>	50	
<i>Abies magnifica</i> (Abies nobilis).....	214	<i>Acer circinatum</i>	47	
<i>Abies Mariana</i>	203	<i>Acer coccineum</i>	56	
		<i>Acer dasycarpum</i>	49	

	Page.		Page.
<i>Acer Douglasii</i>	48	<i>Alnus glauca</i>	164
<i>Acer Drummondii</i>	50	? <i>Alnus glutinosa</i> (<i>Alnus rubra</i>).....	163
<i>Acer eriocarpum</i>	49	<i>Alnus glutinosa</i> , var. <i>rugosa</i>	164
<i>Acer glabrum</i>	47	<i>Alnus glutinosa</i> , var. <i>serrulata</i> (<i>Alnus rhombifolia</i>).....	163
? <i>Acer glaucum</i>	50	<i>Alnus glutinosa</i> , var. <i>serrulata</i> (<i>Alnus serrulata</i>).....	164
<i>Acer grandidentatum</i>	48	<i>Alnus hybrida</i>	164
<i>Acer macrophyllum</i>	47	<i>Alnus incana</i>	164
<i>Acer montanum</i>	46	<i>Alnus incana</i> (Regel).....	164
<i>Acer Negundo</i>	51	<i>Alnus incana</i> , var. <i>glauca</i> (<i>Alnus incana</i>).....	164
<i>Acer nigrum</i>	49	<i>Alnus incana</i> , var. <i>glauca</i> (<i>Alnus incana</i> , var. <i>virescens</i>).....	165
<i>Acer palmatum</i>	47	<i>Alnus incana</i> , var. <i>rubra</i>	163
<i>Acer parviflorum</i>	46	<i>Alnus incana</i> , var. <i>virescens</i>	165
<i>Acer Pennsylvanicum</i>	46	<i>Alnus Japonica</i>	162
<i>Acer Pennsylvanicum</i> (<i>Acer spicatum</i>).....	46	<i>Alnus maritima</i>	162
<i>Acer rubrum</i>	50	<i>Alnus maritima typica</i>	162
<i>Acer rubrum</i> , var. <i>Drummondii</i>	50	<i>Alnus oblongata</i> (<i>Alnus maritima</i>).....	162
<i>Acer rubrum</i> , var. <i>pallidum</i>	49	<i>Alnus oblongata</i> (<i>Alnus serrulata</i>).....	164
<i>Acer saccharinum</i>	48	<i>Alnus oblongifolia</i>	163
<i>Acer saccharinum</i> (<i>Acer dasycarpum</i>).....	49	<i>Alnus obtusifolia</i>	164
<i>Acer saccharinum</i> (<i>Acer saccharinum</i> , var. <i>nigrum</i>).....	49	<i>Alnus Oregona</i>	163
<i>Acer saccharinum</i> , var. <i>nigrum</i>	49	<i>Alnus rhombifolia</i>	163
<i>Acer saccharum</i>	48	<i>Alnus rubra</i>	163
<i>Acer sanguineum</i>	50	<i>Alnus rubra</i> (<i>Alnus serrulata</i>).....	164
<i>Acer spicatum</i>	46	<i>Alnus rugosa</i>	164
<i>Acer striatum</i>	46	<i>Alnus serrulata</i>	164
<i>Acer tripartitum</i>	48	? <i>Alnus serrulata</i> (Regel).....	164
<i>Acer virgatum</i>	47	<i>Alnus serrulata genuina</i>	164
<i>Achras cuneifolia</i>	103	<i>Alnus serrulata</i> , var. <i>latifolia</i>	164
<i>Achras mammosa</i>	103	<i>Alnus serrulata</i> , var. <i>macrophylla</i>	164
<i>Achras pallida</i>	101	<i>Alnus serrulata</i> , var. <i>oblongata</i>	164
<i>Achras salicifolia</i>	101	<i>Alnus serrulata</i> , var. <i>oblongifolia</i>	163
? <i>Achras serrata</i>	69	<i>Alnus serrulata</i> , var. <i>rugosa</i> (<i>Alnus incana</i> , var. <i>virescens</i>).....	165
<i>Achras Zapotilla</i> , var. <i>parviflora</i>	103	<i>Alnus serrulata</i> , var. <i>rugosa</i> (<i>Alnus rhombifolia</i>).....	163
<i>Adelia acuminata</i>	112	<i>Alnus serrulata</i> , var. <i>rugosa</i> (<i>Alnus serrulata</i>).....	164
<i>Esculus Californica</i>	43	<i>Alnus serrulata</i> , var. <i>vulgaris</i>	164
? <i>Esculus carnea</i>	42	<i>Alnus undulata</i>	164
<i>Esculus discolor</i>	43	<i>Amelanchier alnifolia</i>	85
<i>Esculus echinata</i>	42	<i>Amelanchier Bartramiana</i>	84
<i>Esculus flava</i>	43	<i>Amelanchier Botryapium</i>	84
<i>Esculus flava</i> , var. <i>purpurascens</i>	43	<i>Amelanchier Canadensis</i>	84
<i>Esculus glabra</i>	42	<i>Amelanchier Canadensis</i> , var. <i>oblongifolia</i>	84
<i>Esculus Hippocastanum</i> , var. <i>glabra</i>	42	<i>Amelanchier Canadensis</i> , var. <i>oligocarpa</i>	84
<i>Esculus Hippocastanum</i> , var. <i>Ohioensis</i>	42	<i>Amelanchier Canadensis</i> , var. <i>rotundifolia</i>	84
<i>Esculus Hippocastanum</i> , var. <i>pallida</i>	42	<i>Amelanchier intermedia</i>	85
<i>Esculus hybrida</i>	43	<i>Amelanchier oblongifolia</i>	85
<i>Esculus lutea</i>	43	<i>Amelanchier ovalis</i> (<i>Amelanchier Canadensis</i>).....	84
<i>Esculus neglecta</i>	43	<i>Amelanchier ovalis</i> (<i>Amelanchier Canadensis</i> , var. <i>oblongifolia</i>).....	85
<i>Esculus octandra</i>	48	<i>Amelanchier spicata</i>	85
<i>Esculus Ohioensis</i>	42	<i>Amelanchier Wangenheimiana</i>	84
<i>Esculus pallida</i>	42	American Crab.....	73
<i>Esculus Pavia</i> , var. <i>discolor</i>	43	American Crab Apple.....	73
? <i>Esculus Watsoniana</i>	42	American Elm.....	123
<i>Agastianis secundiflora</i>	57	American Holly.....	35
<i>Ageria Cassena</i>	36	American Linden.....	27
<i>Ageria geminata</i>	36	<i>Amyris cymosa</i>	33
<i>Ageria heterophylla</i>	35	<i>Amyris dyatripa</i>	33
<i>Ageria obovata</i>	35	<i>Amyris clemifera</i>	31
<i>Ageria opaca</i>	35	<i>Amyris Floridana</i>	33
<i>Ageria palustris</i>	35	<i>Amyris maritima</i>	33
Alder (<i>Alnus oblongifolia</i>).....	163	<i>Amyris Plumieri</i>	33
Alder (<i>Alnus rhombifolia</i>).....	163	<i>Amyris sylvatica</i>	33
Alder (<i>Alnus rubra</i>).....	163	ANACARDIACE.....	52-54
Alder, Black (<i>Alnus incana</i>).....	165	<i>Anamomis punctata</i>	88
Alder, Black (<i>Alnus serrulata</i>).....	164	Annua.....	114
Alder, Hoary.....	165	<i>Andromeda arborea</i>	98
Alder, Seaside.....	162	<i>Andromeda arborescens</i>	98
Alder, Smooth.....	164	<i>Andromeda ferruginea</i>	96
Alder, Speckled.....	165	<i>Andromeda ferruginea</i> , var. <i>arborescens</i>	96
Algaroba.....	62	<i>Andromeda ferruginea</i> , var. <i>fruticosa</i>	96
<i>Algarobia glandulosa</i>	61	<i>Andromeda plumata</i>	37
<i>Alnus Americana</i> (<i>Alnus incana</i>).....	164	<i>Andromeda rhomboidalis</i>	96
<i>Alnus Americana</i> (<i>Alnus serrulata</i>).....	164	<i>Andromeda rigida</i>	96
<i>Alnus arguta</i>	162	<i>Anona glabra</i>	23
<i>Alnus Canadensis</i>	164	<i>Anona laurifolia</i>	23
<i>Alnus genuina</i>	164	<i>Anona pendula</i>	23

	Page.		Page.
<i>Anona</i> species (<i>Anona laurifolia</i>)	28	Bartram's Oak	188
<i>Anona triloba</i>	28	Basket Oak	141
ANONACEÆ	28	Basswood	27
<i>Anonymos aquatica</i>	124	Basswood, White	28
<i>Anthomeles æstivalis</i>	82	Bastard Cedar	176
<i>Anthomeles Douglasii</i>	75	Bastard Pine	202
<i>Anthomeles flava</i>	82	<i>Batodendron arboresum</i>	96
<i>Anthomeles glandulosa</i>	82	Bayonet, Spanish (<i>Yucca baccata</i>)	219
<i>Anthomeles rotundifolia</i>	77	Bayonet, Spanish (<i>Yucca canaliculata</i>)	218
<i>Anthomeles turbinata</i>	82	Bayonet, Spanish (<i>Yucca elata</i>)	219
Ant's Wood	103	Bay, Bull	19
Apple, American Crab	73	Bay, Lobloily	25
Apple, Custard	23	Bay, Red	118
Apple Haw	82	Bay, Rose	90
Apple, Oregon Crab	73	Bay, Sweet	20
Apple, Pond	23	Bay, Tan	25
Apple, Seven-year	95	Bay, White	20
Apple, Southern Crab	72	Bayberry	136
Arbol de Hierro	56	Bay Tree, California	120
Arbor-vitæ	176	Bean Tree	115
<i>Arbutus laurifolia</i>	97	Bearberry	41
<i>Arbutus macrophylla</i>	97	Bearwood	41
<i>Arbutus Menziesii</i>	97	Beaver Tree	20
<i>Arbutus Menziesii</i> (<i>Arbutus Texana</i>)	97	Beech	158
<i>Arbutus Menziesii</i> (<i>Arbutus Xalapensis</i>)	97	Beech, Blue	159
<i>Arbutus procera</i>	97	Beech, Water (<i>Carpinus Caroliniana</i>)	159
<i>Arbutus Texana</i>	97	Beech, Water (<i>Platanus occidentalis</i>)	129
<i>Arbutus varians</i>	97	Beefwood	117
<i>Arbutus Xalapensis</i>	97	Bee Tree	27
<i>Arbutus Xalapensis</i> (<i>Arbutus Texana</i>)	97	<i>Benthamedia florida</i>	90
Ardisia Pickeringia	100	<i>Betula acuminata</i>	159
<i>Aronia arborea</i>	84	<i>Betula alba</i> , subspecies <i>commutata</i>	160
<i>Aronia arbutifolia</i>	83	<i>Betula alba</i> , subspecies <i>occidentalis typica</i>	160
<i>Aronia Botryopium</i>	84	<i>Betula alba</i> , subspecies <i>papyrifera</i>	160
<i>Aronia cordata</i>	84	<i>Betula alba</i> , subspecies <i>papyrifera</i> , var. <i>communis</i>	160
<i>Aronia ovalis</i>	85	<i>Betula alba</i> , subspecies <i>papyrifera</i> , var. <i>cordifolia</i>	160
Arrowwood	38	<i>Betula alba</i> , subspecies <i>populifolia</i>	159
<i>Asagracea spinosa</i>	55	<i>Betula alba</i> , var. <i>papyrifera</i>	160
Ash	107	<i>Betula alba</i> , var. <i>populifolia</i>	159
Ash, Black	111	<i>Betula alba</i> , var. <i>populifolia</i> (<i>Betula papyrifera</i>)	160
Ash, Blue	111	<i>Betula angulata</i>	161
Ash, Green	109	<i>Betula Canadensis</i>	160
Ash, Ground	111	<i>Betula carpinifolia</i>	162
Ash, Hoop	111	<i>Betula cordifolia</i>	160
Ash, Mountain (<i>Pyrus Americana</i>)	78	<i>Betula excelsa</i> (<i>Betula lenta</i>)	162
Ash, Mountain (<i>Pyrus sambucifolia</i>)	74	<i>Betula excelsa</i> (<i>Betula lutea</i>)	161
Ash, Oregon	111	<i>Betula grandis</i>	160
Ash, Prickly (<i>Xanthoxylum Americanum</i>)	29	<i>Betula inoana</i>	164
Ash, Prickly (<i>Xanthoxylum Clava-Herculis</i>)	80	<i>Betula lanulosa</i>	161
Ash, Red	109	<i>Betula lenta</i>	162
Ash, Sea	80	<i>Betula lenta</i> (<i>Betula alba</i> , var. <i>populifolia</i>)	159
Ash, Wafer	81	<i>Betula lenta</i> (<i>Betula lutea</i>)	161
Ash, Water	110	<i>Betula lutea</i>	161
Ash, White	107	<i>Betula nigra</i>	161
Ash, Yellow	57	<i>Betula nigra</i> (<i>Betula lenta</i>)	162
Ash-leaved Maple	51	<i>Betula nigra</i> (<i>Betula papyrifera</i>)	160
<i>Asimina campaniflora</i>	23	<i>Betula occidentalis</i>	160
<i>Asimina triloba</i>	23	<i>Betula occidentalis</i> (<i>Betula papyrifera</i>)	160
Aspen	171	<i>Betula papyracea</i>	159
Asp, Quaking	171	<i>Betula papyrifera</i>	159
<i>Avicennia nitida</i>	117	<i>Betula populifolia</i>	159
<i>Avicennia oblongifolia</i>	117	<i>Betula rubra</i> (<i>Betula nigra</i>)	161
<i>Avicennia tomentosa</i>	117	<i>Betula rubra</i> (<i>Betula papyrifera</i>)	160
		<i>Betula rugosa</i>	164
Bald Cypress	184	<i>Betula serrulata</i>	164
Balm of Gilead	178	? <i>Betula-Ainus glauca</i>	164
Balm of Gilead Fir	211	<i>Betula-Ainus maritima</i>	162
Balsam (<i>Abies Fraseri</i>)	210	? <i>Betula-Ainus rubra</i>	164
Balsam (<i>Abies subalpina</i>)	211	<i>Betula-Ainus</i> , var. <i>β inoana</i>	164
Balsam (<i>Populus balsamifera</i>)	173	BETULACEÆ	159-165
Balsam Cottonwood	174	Big-bud Hickory	124
Balsam Fir (<i>Abies balsamea</i>)	211	Big Cottonwood	175
Balsam Fir (<i>Abies concolor</i>)	213	Big Laurel	19
Balsam, She	210	Big Shell-bark	183
Banana, Mexican	219	Big Tree	184

223

	Page.		Page.
<i>Bignelovia acuminata</i>	112	<i>Bourreria Havanensis</i>	114
<i>Bignonia Catalpa</i>	115	<i>Bourreria Havanensis, var. radula</i>	114
<i>Bignonia linearis</i>	116	<i>Bourreria ovata</i>	114
BIGNONIACEÆ	115, 116	<i>Bourreria radula</i>	114
Bilsted	86	<i>Bourreria recurva</i>	114
Birch, Black (<i>Betula lenta</i>)	162	<i>Bourreria tomentosa</i>	114
Birch, Black (<i>Betula occidentalis</i>)	160	<i>Bourreria tomentosa, var. Havanensis</i>	114
Birch, Canoe	160	Box Elder (<i>Negundo aceroides</i>)	51
Birch, Cherry	162	Box Elder (<i>Negundo Californicum</i>)	51
Birch, Gray (<i>Betula alba, var. populifolia</i>)	159	Boxwood (<i>Cornus florida</i>)	91
Birch, Gray (<i>Betula lutea</i>)	161	Boxwood (<i>Schæfferia frutescens</i>)	39
Birch, Mahogany	162	<i>Brahea dulcis?</i>	217
Birch, Old-field	159	<i>Brahea filamentosa</i>	21
Birch, Paper	160	Brickley Thatch	218
Birch, Red	161	Brittle Thatch	218
Birch, River	161	Broad-leaved Maple	47
Birch, Sweet	162	<i>Broussonetia secundiflora</i>	57
Birch, West Indian	33	<i>Broussonetia tinctoria</i>	128
Birch, White (<i>Betula alba, var. populifolia</i>)	159	Brown Hickory	184
Birch, White (<i>Betula papyrifera</i>)	160	<i>Bucida Buceras</i>	87
Birch, Yellow	161	Buckeye, California	44
Bishop's Pine	200	Buckeye, Fetid	42
Bitter Nut	135	Buckeye, Ohio	42
Bitter Pecan	136	Buckeye, Spanish	44
Black Alder (<i>Alnus incana</i>)	165	Buckeye, Sweet	43
Black Alder (<i>Alnus serrulata</i>)	164	Buckthorn, Southern	103
Black Ash	111	Buckwheat Tree	38
Black Birch (<i>Betula lenta</i>)	162	Bull Bay	19
Black Birch (<i>Betula occidentalis</i>)	160	Bull Nut	134
Black Calabash Tree	116	Bull Pine (<i>Pinus Jeffreyi</i>)	193
Black Cherry, Wild	68	Bull Pine (<i>Pinus mitis</i>)	200
Black Cottonwood (<i>Populus angustifolia</i>)	174	Bull Pine (<i>Pinus ponderosa</i>)	193
Black Cottonwood (<i>Populus trichocarpa</i>)	174	Bull Pine (<i>Pinus Sabiniana</i>)	195
Black Cypress	184	<i>Bumelia angustifolia</i>	103
Black Gum	92	<i>Bumelia arborea</i>	102
Black Haw	94	<i>Bumelia chrysophylloides</i>	101
Black Hickory (<i>Carya porcina</i>)	134	<i>Bumelia cuneata</i>	103
Black Hickory (<i>Carya tomentosa</i>)	134	<i>Bumelia ferruginea</i>	102
Black Ironwood	39	<i>Bumelia fetidissima</i>	101
Black Jack (<i>Quercus Catesbeii</i>)	151	<i>Bumelia lanuginosa</i>	102
Black Jack (<i>Quercus nigra</i>)	150	<i>Bumelia lanuginosa, var. macrocarpa</i>	102
Black Jack, Forked-leaf	151	<i>Bumelia lycioides</i>	102
Black Larch	215	<i>Bumelia lycioides, var. reclinatam</i>	103
Black Locust (<i>Gleditschia triacanthos</i>)	59	<i>Bumelia macrocarpa</i>	102
Black Locust (<i>Robinia Pseudacacia</i>)	55	<i>Bumelia Mastichodendron</i>	101
Black Mangrove	117	<i>Bumelia myrsinifolia</i>	103
Black Oak (<i>Quercus Emoryi</i>)	146	? <i>Bumelia oblongifolia</i>	102
Black Oak (<i>Quercus Kelloggii</i>)	149	<i>Bumelia pallida</i>	101
Black Oak (<i>Quercus rubra</i>)	148	<i>Bumelia parvifolia</i>	103
Black Oak (<i>Quercus tinctoria</i>)	149	<i>Bumelia reclinata</i> (<i>Bumelia cuneata</i>)	103
Black Persimmon	105	<i>Bumelia reclinata</i> (<i>Bumelia lycioides</i>)	103
Black Pine (<i>Pinus Jeffreyi</i>)	193	? <i>Bumelia reclinata</i> (<i>Bumelia tenax</i>)	101
Black Pine (<i>Pinus Murrayana</i>)	195	<i>Bumelia salicifolia</i>	101
Black Sloe	67	? <i>Bumelia serrata</i>	69
Black Spruce	208	<i>Bumelia spinosa</i>	102
Black Sugar Maple	49	<i>Bumelia tenax</i>	101
Black Thorn	79	<i>Bumelia tomentosa</i>	102
Black Tree	117	Bumwood	54
Black Walnut	131	Burning Bush	38
Black Willow (<i>Salix flavescons, var. Scouleriana</i>)	170	Burr Oak	140
Black Willow (<i>Salix nigra</i>)	166	<i>Bursara acuminata</i>	32
Blackwood	117	<i>Bursara gummifera</i>	32
Blue Ash	111	BURSARACEÆ	32, 33
Blue Beech	159	Bustle	101</

	Page.		Page.
CACTACEÆ	89, 90	<i>Castanea vesca</i> (<i>Castanea vulgaris</i> , var. <i>Americana</i>)	157
<i>Cactus</i> , Giant	90	<i>Castanea vesca</i> , var. <i>Americana</i>	157
Cagiput	120	<i>Castanea vulgaris</i> , var. <i>Americana</i>	157
Calabash Tree, Black	116	<i>Castanopsis chrysophylla</i>	156
Calico Bush	98	<i>Castanopsis chrysophylla</i> , var. <i>minor</i>	156
Calicowood	106	<i>Castanopsis chrysophylla</i> , var. <i>pumila</i>	156
California Bay Tree	120	Catalpa	115
California Buckeye	44	<i>Catalpa bignonioides</i>	115
California Holly	84	<i>Catalpa bignonioides</i> (<i>Catalpa speciosa</i>)	115
California Laurel	120	<i>Catalpa communis</i>	115
California Nutmeg	186	<i>Catalpa cordifolia</i> (<i>Catalpa bignonioides</i>)	115
California Olive	120	<i>Catalpa cordifolia</i> (<i>Catalpa speciosa</i>)	115
<i>Calothyrsus Californica</i>	43	<i>Catalpa speciosa</i>	115
Calyptranthes Chytraculla	88	<i>Catalpa syringaefolia</i>	115
Canada Plum	65	Catalpa, Western	115
Canella alba	24	Catawba	115
<i>Canella laurifolia</i>	24	Cat's Claw (<i>Acacia Greggii</i>)	63
<i>Canella Winterana</i>	24	Cat's Claw (<i>Acacia Wrightii</i>)	63
CANELLACEÆ	24	Cat's Claw (<i>Pithecolobium Ungula-cati</i>)	64
Canoe Birch	160	<i>Ceanothus ferreus</i>	39
Canoe Cedar	177	<i>Ceanothus laevigatus</i>	39
<i>Canotia holocantha</i>	32	<i>Ceanothus reclinatus</i>	41
CAPPARIDACEÆ	24	<i>Ceanothus thyrsiflorus</i>	41
<i>Capparis Breynia</i>	24	Cedar, Bastard	176
<i>Capparis cynophyllophora</i>	24	Cedar, Canoe	177
<i>Capparis emarginata</i>	24	Cedar Elm	122
<i>Capparis Jamaicensis</i>	24	Cedar, Incense	176
<i>Capparis siliquosa</i>	24	Cedar, Oregon	179
<i>Capparis torulosa</i>	24	Cedar Pine	201
<i>Capparis uncinata</i>	24	Cedar, Port Orford	179
CAPRIFOLIACEÆ	93, 94	Cedar, Post	176
<i>Cardioplepis ob-ovata</i>	41	Cedar, Red (<i>Juniperus Virginiana</i>)	182
Carolina Poplar	175	Cedar, Red (<i>Thuja gigantea</i>)	177
<i>Carpinus Americana</i>	158	Cedar, Stinking (<i>Torreya Californica</i>)	186
<i>Carpinus Betulus Virginiana</i>	158	Cedar, Stinking (<i>Torreya taxifolia</i>)	186
<i>Carpinus Caroliniana</i>	158	Cedar, White (<i>Chamaecyparis Lawsoniana</i>)	179
<i>Carpinus Ostrya</i>	158	Cedar, White (<i>Chamaecyparis sphaeroides</i>)	178
<i>Carpinus Ostrya</i> , var. <i>Americana</i>	158	Cedar, White (<i>Libocedrus decurrens</i>)	176
<i>Carpinus triflora</i>	158	Cedar, White (<i>Thuja occidentalis</i>)	176
<i>Carpinus Virginiana</i>	158	<i>Cedrus Mahagoni</i>	33
<i>Carya alba</i>	132	CELASTRACEÆ	38, 39
<i>Carya alba</i> (<i>Carya tomentosa</i>)	134	<i>Celtis alba</i>	125
<i>Carya amara</i>	135	<i>Celtis Audubertiana</i>	126
<i>Carya amara</i> , var. <i>myristiciformis</i>	135	<i>Celtis Audubertiana</i> , var. <i>oblongata</i>	126
<i>Carya amara</i> , var. <i>porcina</i>	134	<i>Celtis Audubertiana</i> , var. <i>ovata</i>	126
<i>Carya angustifolia</i>	132	<i>Celtis Berlandieri</i>	126
<i>Carya aquatica</i>	135	<i>Celtis brevipes</i>	126
<i>Carya cathartica</i>	130	<i>Celtis canina</i>	125
<i>Carya cordiformis</i>	133	<i>Celtis cordata</i>	125
<i>Carya glabra</i>	134	<i>Celtis crassifolia</i>	125
<i>Carya Illinoensis</i>	132	<i>Celtis crassifolia</i> , var. <i>eucalyptifolia</i>	125
<i>Carya integrifolia</i>	135	<i>Celtis crassifolia</i> , var. <i>morifolia</i>	125
<i>Carya microcarpa</i>	133	<i>Celtis crassifolia</i> , var. <i>tiliaefolia</i>	125
<i>Carya myristiciformis</i>	135	<i>Celtis Douglasii</i>	126
<i>Carya oliviformis</i>	132	<i>Celtis Floridiana</i>	125
<i>Carya porcina</i>	134	<i>Celtis fuscata</i>	125
<i>Carya sulcata</i>	133	<i>Celtis heterophylla</i>	125
? <i>Carya tetraptera</i>	132	<i>Celtis integrifolia</i>	125
<i>Carya tomentosa</i>	133	<i>Celtis laevigata</i>	125
<i>Carya tomentosa</i> , var. <i>maxima</i>	134	<i>Celtis Lindheimeri</i>	125
<i>Caryotaxus Myristica</i>	186	<i>Celtis longifolia</i>	125
<i>Caryotaxus taxifolia</i>	186	<i>Celtis maritima</i> (in <i>Am. Monthly Mag. and Crit. Rev.</i>)	125
Cassida	101	<i>Celtis maritima</i> (in <i>New Fl. and Bot.</i>)	125
Cassena	36	<i>Celtis Mississippiensis</i>	125
<i>Cassine Caroliniana</i>	36	<i>Celtis morifolia</i>	125
<i>Cassine Peragua</i>	36	<i>Celtis obliqua</i>	125
<i>Cassine ramulosa</i>	36	<i>Celtis occidentalis</i>	125
<i>Castanea alnifolia</i>	156	<i>Celtis occidentalis</i> , var. <i>cordata</i>	125
<i>Castanea Americana</i>	157	<i>Celtis occidentalis</i> , var. <i>crassifolia</i>	126
<i>Castanea chrysophylla</i>	156	<i>Celtis occidentalis</i> , var. <i>grandidentata</i>	125
<i>Castanea chrysophylla</i> , var. <i>minor</i>	156	<i>Celtis occidentalis</i> , var. <i>integrifolia</i>	125
<i>Castanea nana</i>	156	<i>Celtis occidentalis</i> , var. <i>pumila</i> (<i>Celtis occidentalis</i>)	126
<i>Castanea pumila</i>	156	? <i>Celtis occidentalis</i> , var. <i>pumila</i> (<i>Celtis occidentalis</i> var. <i>reticulata</i>)	126
<i>Castanea sempervirens</i>	156	<i>Celtis occidentalis</i> , var. <i>reticulata</i>	126
<i>Castanea vesca</i> (<i>Castanea pumila</i>)	156	<i>Celtis occidentalis</i> , var. <i>scabriscula</i>	125

	Page.		Page.
<i>Celtis occidentalis</i> , var. <i>serrulata</i>	125	<i>Chilopsis linearis</i>	116
<i>Celtis occidentalis</i> , var. <i>tenuifolia</i>	125	<i>Chilopsis saligna</i>	116
<i>Celtis patula</i>	125	<i>Chimanthus amygdalinus</i>	69
<i>Celtis pumila</i>	125	China, Wild	44
<i>Celtis reticulata</i>	126	Chinquapin (<i>Castanea pumila</i>)	156
<i>Celtis salicifolia</i>	125	Chinquapin (<i>Castanopsis chrysophylla</i>)	156
<i>Celtis tenuifolia</i>	125	Chinquapin Oak	143
<i>Celtis Texana</i>	126	<i>Chionanthus angustifolia</i>	112
<i>Cerasus Americana</i>	65	<i>Chionanthus heterophylla</i>	112
<i>Cerasus borealis</i>	66	<i>Chionanthus longifolia</i>	112
<i>Cerasus Capollin</i>	68	<i>Chionanthus maritima</i>	112
<i>Cerasus Capuli</i>	68	<i>Chionanthus montana</i>	112
<i>Cerasus Caroliniana</i>	69	<i>Chionanthus trifida</i>	112
<i>Cerasus Chicasa</i>	66	<i>Chionanthus Virginica</i>	112
<i>Cerasus demissa</i>	69	<i>Chionanthus Virginica</i> , var. <i>angustifolia</i>	112
<i>Cerasus emarginata</i>	67	<i>Chionanthus Virginica</i> , var. <i>latifolia</i>	112
<i>Cerasus erecta</i>	67	<i>Chionanthus Virginica</i> , var. <i>maritima</i>	112
<i>Cerasus glandulosa</i>	67	<i>Chionanthus Virginica</i> , var. <i>montana</i>	112
<i>Cerasus hiemalis</i>	65	Chittamwood	52
<i>Cerasus ilicifolia</i>	70	<i>Chloromeles sempervirens</i>	72
<i>Cerasus mollis</i>	67	<i>Chrysobalanus Icaco</i>	64
<i>Cerasus nigra</i>	65	<i>Chrysobalanus Icaco</i> , var. <i>pellocarpa</i>	65
<i>Cerasus Pennsylvanica</i>	66	<i>Chrysobalanus pellocarpa</i>	65
? <i>Cerasus persicifolia</i>	66	<i>Chrysophyllum Barbascio</i>	100
<i>Cerasus serotina</i> (<i>Prunus demissa</i>)	69	<i>Chrysophyllum Caneto</i> , β.	100
<i>Cerasus serotina</i> (<i>Prunus serotina</i>)	68	<i>Chrysophyllum ferrugineum</i>	100
<i>Cerasus sphaerocarpa</i>	70	<i>Chrysophyllum microphyllum</i>	100
<i>Cerasus umbellata</i>	67	<i>Chrysophyllum monopyrenum</i>	100
<i>Cerasus Virginiana</i>	68	<i>Chrysophyllum oliviforme</i>	100
<i>Cercidium floridum</i>	60	Cigar Tree	115
<i>Cercis Canadensis</i>	61	<i>Cinchona Caribaea</i>	95
<i>Cercis Canadensis</i> , var. <i>pubescens</i>	61	<i>Cinchona Caroliniana</i>	95
<i>Cercis occidentalis</i>	61	<i>Cinchona Jamaicensis</i>	95
<i>Cercis occidentalis</i> , var.	61	Cinnamon Bark	24
<i>Cercis occidentalis</i> , var. <i>Texensis</i>	61	Cinnamon, Wild	24
<i>Cercis reniformis</i>	61	<i>Citharexylum villosum</i>	116
<i>Cercocarpus betulafolia</i>	71	<i>Cladrastis hutea</i>	57
<i>Cercocarpus betuloides</i>	71	<i>Cladrastis tinctoria</i>	57
<i>Cercocarpus brevifolius</i>	71	Clammy Locust	56
<i>Cercocarpus intricatus</i>	71	Cliff Elm	123
<i>Cercocarpus ledifolius</i>	71	<i>Cliftonia ligustrina</i>	38
<i>Cercocarpus ledifolius</i> , var. <i>intricatus</i>	71	<i>Clusia flava</i>	25
<i>Cercocarpus parvifolius</i>	71	<i>Clusia rosea</i>	25
<i>Cercocarpus parvifolius</i> , var. <i>glaber</i>	71	Coast Live Oak	147
<i>Cercocarpus parvifolius</i> , var. <i>paucidentatus</i>	71	<i>Coccoloba Floridana</i>	117
<i>Cereus giganteus</i>	89	<i>Coccoloba Laganensis</i>	118
Chapote	105	<i>Coccoloba parvifolia</i>	117
<i>Chamaecyparis Boursierii</i> (<i>Chamaecyparis Lawsoniana</i>)	178	<i>Coccoloba uvifera</i>	118
<i>Chamaecyparis Boursierii</i> (<i>Juniperus occidentalis</i>)	181	<i>Coccoloba uvifera</i> , var. <i>Laganensis</i>	118
<i>Chamaecyparis excoleta</i>	178	<i>Coccoloba uvifera</i> , var. <i>ovalifolia</i>	118
<i>Chamaecyparis Lawsoniana</i>	178	Cockspur Thorn	76
<i>Chamaecyparis Nutkaensis</i>	178	Cocoa Plum	65
<i>Chamaecyparis Nutkaensis</i> , var. <i>glauca</i>	178	Coffee Nut	58
<i>Chamaecyparis Nutkanus</i>	178	Coffee Tree, Kentucky	58
<i>Chamaecyparis sphaeroides</i>	177	<i>Colubrina reclinata</i>	41
<i>Chamaecyparis Palmetto</i>	217	COMBRETACEÆ	87
<i>Chetranthodendron Californicum</i>	26	<i>Condalia ferrea</i>	39
Cherry	100	<i>Condalia obovata</i>	46
Cherry Birch	162	CONIFERÆ	176-216
Cherry, Indian	40	<i>Conocarpus acutifolia</i>	87
Cherry, May	84	<i>Conocarpus erecta</i>	87
Cherry, Pigeon	66	<i>Conocarpus erecta</i> , var. <i>procumbens</i>	87
Cherry, Pin	66	<i>Conocarpus procumbens</i>	87
Cherry, Rum	68	<i>Conocarpus racemosa</i>	87
Cherry, Wild (<i>Prunus Capuli</i>)	69	Coral Sumach	54
Cherry, Wild (<i>Prunus demissa</i>)	69	<i>Cordia Boissieri</i>	114
Cherry, Wild Black	68	<i>Cordia Floridana</i>	114
Cherry, Wild Red	68	? <i>Cordia juglandifolia</i>	113
Chestnut	157	<i>Cordia Sebestena</i>	113
Chestnut Oak (<i>Quercus densiflora</i>)	155	<i>Cordia speciosa</i>	113
Chestnut Oak (<i>Quercus prinoides</i>)	143	Cork Elm	123
Chestnut Oak (<i>Quercus Prinus</i>)	142	Corkwood	117
Chestnut Oak, Rock	142	CORNACEÆ	90-93
Chickasaw Plum	66	<i>Cornus alternata</i>	90
<i>Chilopsis glutinosa</i>	116	<i>Cornus alternifolia</i>	90

	Page.		Page.
<i>Cornus florida</i>	90	<i>Oratagus linearis</i>	77
<i>Cornus florida</i> (<i>Cornus Nuttallii</i>).....	91	<i>Oratagus lobata</i> (<i>Cratægus flava</i>).....	83
<i>Cornus Nuttallii</i>	91	<i>Oratagus lobata</i> (<i>Cratægus tomentosa</i>).....	79
<i>Corypha Palmetto</i>	217	<i>Oratagus lucida</i> (<i>Cratægus æstivalis</i>).....	82
<i>Cotinus Americanus</i>	52	<i>Oratagus lucida</i> (<i>Cratægus Crus-galli</i>).....	76
<i>Cotinus coggygria</i>	52	<i>Oratagus macracantha</i>	77
Cotton Gum.....	98	<i>Oratagus Michauxii</i>	82
Cottonwood (<i>Populus Fremontii</i>).....	175	<i>Oratagus microcarpa</i>	81
Cottonwood (<i>Populus Fremontii</i> , var. <i>Wallseni</i>).....	175	<i>Oratagus mollis</i>	78
Cottonwood (<i>Populus monilifera</i>).....	175	<i>Oratagus obovatifolia</i>	80
Cottonwood, Balsam.....	174	<i>Oratagus opaca</i>	82
Cottonwood, Big.....	175	<i>Oratagus ovalifolia</i>	76
Cottonwood, Black (<i>Populus angustifolia</i>).....	174	<i>Oratagus ozyacantha</i>	81
Cottonwood, Black (<i>Populus trichocarpa</i>).....	174	<i>Oratagus ozyacantha</i> , var. <i>apifolia</i>	81
Cottonwood, River.....	172	<i>Cratægus parvifolia</i>	83
Cottonwood, Swamp.....	172	<i>Oratagus populifolia</i> (<i>Cratægus coccinea</i>).....	78
Cottonwood, White.....	175	<i>Oratagus populifolia</i> (<i>Cratægus cordata</i>).....	80
Cow Oak.....	141	<i>Oratagus prunellifolia</i>	76
Crab, American.....	72	<i>Oratagus prunifolia</i>	77
Crab Apple, American.....	72	<i>Oratagus punctata</i>	80
Crab Apple, Oregon.....	78	<i>Oratagus punctata</i> , var. <i>bracteopina</i>	75
Crab Apple, Southern.....	72	<i>Oratagus punctata</i> , var. <i>rubra</i> and <i>ovata</i>	80
Crab, Sweet-scented.....	72	<i>Oratagus punctata</i> , var. <i>zanthocarpa</i>	80
Crabwood.....	121	<i>Oratagus pyrifolia</i>	79
<i>Cratægus æstivalis</i>	82	<i>Oratagus racemosa</i>	84
<i>Cratægus apifolia</i>	81	<i>Cratægus rivularis</i>	74
<i>Cratægus arborescens</i>	75	<i>Oratagus rivularis</i> (<i>Cratægus Douglasii</i>).....	75
<i>Oratagus arbutifolia</i>	83	<i>Oratagus salicifolia</i>	76
<i>Cratægus berberifolia</i>	82	<i>Oratagus sanguinea</i>	75
<i>Oratagus Bosciiana</i>	77	<i>Oratagus sanguinea</i> , var. <i>Douglasii</i> (<i>Cratægus Douglasii</i>).....	75
<i>Cratægus brachyacantha</i>	75	<i>Oratagus sanguinea</i> , var. <i>Douglasii</i> (<i>Cratægus rivularis</i>).....	74
<i>Oratagus Caroliniana</i>	82	<i>Oratagus sanguinea</i> , var. <i>villosa</i>	78
<i>Cratægus coccinea</i>	77	<i>Cratægus spathulata</i>	81
<i>Oratagus coccinea</i> (<i>Cratægus tomentosa</i>).....	79	<i>Oratagus spathulata</i> (<i>Cratægus flava</i> , var. <i>pubescens</i>).....	82
<i>Oratagus coccinea</i> , var. <i>cordata</i>	79	<i>Oratagus spicata</i>	85
<i>Oratagus coccinea</i> , var. <i>mollis</i>	78	<i>Cratægus subvillosa</i>	78
<i>Cratægus coccinea</i> var. <i>oligandra</i>	78	<i>Oratagus Texana</i>	78
<i>Cratægus coccinea</i> , var. <i>populifolia</i>	78	<i>Cratægus tomentosa</i>	79
<i>Oratagus coccinea</i> , var. <i>typica</i>	78	<i>Oratagus tomentosa</i> (<i>Amelanchier Canadensis</i>).....	84
<i>Cratægus coccinea</i> , var. <i>viridis</i>	78	<i>Oratagus tomentosa</i> , var. <i>mollis</i>	78
? <i>Oratagus coccinea</i> , var. <i>viridis</i> (<i>Cratægus tomentosa</i>).....	79	<i>Oratagus tomentosa</i> , var. <i>plicata</i>	80
<i>Cratægus cordata</i>	80	<i>Cratægus tomentosa</i> , var. <i>punctata</i>	80
<i>Oratagus coronaria</i>	72	<i>Oratagus tomentosa</i> , var. <i>pyrifolia</i>	79
<i>Oratagus Courretiana</i>	76	<i>Oratagus turbinata</i>	82
<i>Cratægus Crus-galli</i>	76	<i>Oratagus Virginica</i>	83
<i>Oratagus Crus-galli</i> (<i>Cratægus coccinea</i>).....	77	<i>Oratagus viridis</i> (<i>Cratægus coccinea</i>).....	78
<i>Oratagus Crus-galli</i> (<i>Cratægus tomentosa</i> , var. <i>punctata</i>).....	80	<i>Oratagus viridis</i> (<i>Cratægus flava</i> , var. <i>pubescens</i>).....	83
<i>Cratægus Crus-galli</i> , var. <i>linearis</i>	76	<i>Oratagus Watsoniana</i>	76
<i>Cratægus Crus-galli</i> , var. <i>ovalifolia</i>	76	<i>Crescentia cucurbitina</i>	116
<i>Cratægus Crus-galli</i> , var. <i>prunifolia</i>	77	<i>Crescentia latifolia</i>	116
<i>Cratægus Crus-galli</i> , var. <i>pyracanthifolia</i>	76	<i>Crescentia lathifera</i>	116
<i>Oratagus Crus-galli</i> , var. <i>pyracanthifolia</i> (<i>Cratægus arborescens</i>).....	75	<i>Crescentia obovata</i>	116
<i>Oratagus Crus-galli</i> , var. <i>salicifolia</i>	76	<i>Crescentia ovata</i>	116
<i>Oratagus Crus-galli</i> , var. <i>splendens</i>	76	<i>Crescentia toxicaria</i>	116
<i>Oratagus cuneifolia</i>	80	Cucumber Tree (<i>Magnolia acuminata</i>).....	20
<i>Cratægus Douglasii</i>	75	Cucumber Tree (<i>Magnolia cordata</i>).....	21
<i>Oratagus elliptica</i> (<i>Cratægus æstivalis</i>).....	82	Cucumber Tree, Large-leaved.....	21
<i>Oratagus elliptica</i> (<i>Cratægus flava</i> , var. <i>pubescens</i>).....	83	Cucumber Tree, Long-leaved.....	22
<i>Cratægus flava</i>	82	<i>Cupressinata disticha</i>	183
<i>Oratagus flava</i> (<i>Cratægus flava</i> , var. <i>pubescens</i>).....	83	<i>Cupressus Americana</i>	178
<i>Oratagus flava</i> (<i>Cratægus tomentosa</i>).....	79	<i>Cupressus Arbor-vitæ</i>	176
<i>Oratagus flava</i> (<i>Cratægus tomentosa</i> , var. <i>punctata</i>).....	80	<i>Cupressus Arizonica</i>	180
<i>Oratagus flava</i> , var. <i>lobata</i>	82	? <i>Cupressus attenuata</i>	178
<i>Cratægus flava</i> , var. <i>pubescens</i>	83	? <i>Cupressus Californica</i>	179
? <i>Oratagus flexuosa</i>	80	<i>Cupressus Californica gracilis</i> (<i>Cupressus Goveniana</i>).....	179
<i>Oratagus glandulosa</i> (<i>Cratægus coccinea</i>).....	77	<i>Cupressus Californica gracilis</i> (<i>Cupressus Macnabiana</i>).....	180
? <i>Oratagus glandulosa</i> (<i>Cratægus Douglasii</i>).....	75	? <i>Cupressus cornuta</i>	179
<i>Oratagus glandulosa</i> (<i>Cratægus flava</i>).....	77	<i>Cupressus disticha</i>	183
<i>Oratagus glandulosa</i> (<i>Cratægus flava</i> , var. <i>pubescens</i>).....	83	<i>Cupressus disticha</i> , var. <i>imbricaria</i>	183
<i>Oratagus glandulosa</i> , var. <i>macracantha</i>	77	<i>Cupressus disticha</i> , var. <i>nulans</i>	183
<i>Oratagus glandulosa</i> , var. <i>rotundifolia</i>	78	<i>Cupressus disticha</i> , var. <i>patens</i>	183
<i>Cratægus latifolia</i> (<i>Cratægus tomentosa</i>).....	79	<i>Cupressus fragrans</i>	178
<i>Cratægus latifolia</i> (<i>Cratægus tomentosa</i> , var. <i>punctata</i>).....	80	<i>Cupressus glandulosa</i>	184
<i>Oratagus leucocephalus</i>	79	<i>Cupressus Goveniana</i>	179
<i>Cratægus leucophloeos</i>	79	<i>Cupressus Guadalupensis</i>	184

227

[illegible]

229

	Page.		Page.
<i>Gleditschia inermis</i> (<i>Gleditschia monosperma</i>).....	59	Haw, Black.....	84
<i>Gleditschia inermis</i> (<i>Gleditschia triacanthos</i> , var. <i>inermis</i>).....	59	Haw, Hogs'.....	75
<i>Gleditschia macrocarpa</i>	59	Haw, May.....	82
<i>Gleditschia Melloba</i>	59	Haw, Paralely.....	81
<i>Gleditschia monosperma</i>	59	Haw, Pear.....	79
<i>Gleditschia spinosa</i>	59	Haw, Purple.....	40
<i>Gleditschia triacantha</i>	59	Haw, Red (<i>Cratægus coccinea</i>).....	78
<i>Gleditschia triacanthos</i>	59	Haw, Red (<i>Cratægus flava</i> , var. <i>pubescens</i>).....	83
<i>Gleditschia triacanthos</i> , var. <i>brachycarpus</i>	59	Haw, Scarlet (<i>Cratægus coccinea</i>).....	78
<i>Gleditschia triacanthos</i> , var. <i>inermis</i>	59	Haw, Scarlet (<i>Cratægus subvillosa</i>).....	78
<i>Gleditschia triacanthos</i> , var. <i>monosperma</i>	59	Haw, Small-fruited.....	81
Goose-foot Maple.....	46	Haw, Summer (<i>Cratægus flava</i>).....	86
Gopher Plum.....	91	Haw, Summer (<i>Cratægus flava</i> , var. <i>pubescens</i>).....	86
Gopherwood.....	57	Haw, Yellow.....	83
<i>Gordonia Franklinii</i>	25	Hazel, Witch.....	85
<i>Gordonia Lasianthus</i>	25	Hemlock (<i>Pseudotsuga Douglasii</i> , var. <i>macrocarpa</i>).....	210
<i>Gordonia pubescens</i>	25	Hemlock (<i>Tsuga Canadensis</i>).....	207
<i>Gordonia pyramidalis</i>	25	Hemlock (<i>Tsuga Caroliniana</i>).....	207
Grape, Sea.....	118	Hemlock (<i>Tsuga Mertensiana</i>).....	208
Gray Birch (<i>Betula alba</i> , var. <i>populifolia</i>).....	159	Heteromeles arbutifolia.....	83
Gray Birch (<i>Betula lutea</i>).....	161	<i>Heteromeles Fremontiana</i>	83
Gray Pine.....	201	<i>Heyderia decurrens</i>	176
Great Laurel.....	99	<i>Heymassoli spinosa</i>	34
Green Ash.....	109	<i>Hickoria</i> , species.....	132
Green-bark Acacia.....	60	<i>Hickorius amara</i>	125
Ground Ash.....	111	Hickory, Big-bud.....	134
<i>Guaiacum angustifolium</i>	29	Hickory, Black (<i>Carya porcina</i>).....	134
<i>Guaiacum sanctum</i>	28	Hickory, Black (<i>Carya tomentosa</i>).....	134
<i>Guaiacum verticillate</i>	28	Hickory, Brown.....	134
<i>Guettarda elliptica</i>	96	Hickory Elm.....	123
<i>Guettarda Blodgettii</i>	96	Hickory, Nutmeg.....	135
Guiana Plum.....	121	Hickory Pine (<i>Pinus Balfouriana</i> , var. <i>aristata</i>).....	191
<i>Guilandina dioica</i>	58	Hickory Pine (<i>Pinus pungens</i>).....	199
Gum, Black.....	92	Hickory, Shag-bark.....	133
Gum, Cotton.....	93	Hickory, Shell-bark.....	133
Gum, Doctor.....	54	Hickory, Swamp (<i>Carya amara</i>).....	135
Gum Elastic.....	102	Hickory, Swamp (<i>Carya aquatica</i>).....	126
Gum Elemi.....	33	Hickory, Switch-bud.....	134
Gum, Red.....	86	Hickory, Water.....	126
Gum, Sour.....	92	Hickory, White-heart.....	134
Gum, Star-leaved.....	86	<i>Hicorius integrifolia</i>	125
Gum, Sweet.....	86	<i>Hierophyllum Cassini</i>	86
Gum, Tupelo.....	98	<i>Hippomane Mancinella</i>	121
Gumbo Limbo.....	83	Hoary Alder.....	185
Gurgeon Stopper.....	88	Hog Plum (<i>Prunus angustifolia</i>).....	66
GUTTIFERÆ.....	25	Hog Plum (<i>Rhus Metopium</i>).....	54
<i>Gymnanthes lucida</i>	121	Hog Plum (<i>Ximenia Americana</i>).....	84
<i>Gymnocladus Oatesiana</i>	119	Hogs' Haw.....	75
<i>Gymnocladus Canadensis</i>	58	Holly, American.....	35
<i>Gymnocladus dioica</i>	58	Holly, California.....	84
		Holly, Dahoon.....	35
III.		Honey Locust (<i>Gleditschia triacanthos</i>).....	59
Hackberry (<i>Celtis occidentalis</i>).....	126	Honey Locust (<i>Prosopis juliflora</i>).....	62
Hackberry (<i>Celtis occidentalis</i> , var. <i>reticulata</i>).....	126	Honey Pod.....	62
Hackmatack.....	215	Honey Shucks.....	59
<i>Halesia diptera</i>	105	Hoop Ash.....	111
<i>Halesia parviflora</i>	106	<i>Hopea tinctoria</i>	105
<i>Halesia reticulata</i>	105	Hop Hornbeam.....	158
<i>Halesia tetraptera</i>	106	Hop Tree.....	31
<i>Halmia cornifolia</i>	80	Hornbeam.....	159
<i>Halmia flabellata</i>	77	Hornbeam, Hop.....	158
<i>Halmia lobata</i>	79	Horse Plum.....	65
<i>Halmia punctata</i>	80	Horse Sugar.....	105
<i>Halmia tomentosa</i>	79	<i>Hypelate oblongifolia</i>	45
HAMAMELACEÆ.....	85, 86	<i>Hypelate paniculata</i>	45
<i>Hamamelis androgyna</i>	85	<i>Hypelate trifoliata</i>	45
<i>Hamamelis corylifolia</i>	85	<i>Hyperanthera dioica</i>	58
<i>Hamamelis dioica</i>	85	<i>Hypericum Lasianthus</i>	26
<i>Hamamelis macrophylla</i>	85		
<i>Hamamelis parvifolia</i>	85	I.	
<i>Hamamelis Virginiana</i> , var. <i>parvifolia</i>	85	<i>Ilex astivalis</i>	37
<i>Hamamelis Virginica</i>	85	<i>Ilex ambigua</i>	37
Hard Maple.....	43	<i>Ilex angustifolia</i>	35
Hard Pine.....	202	<i>Ilex aquifolium</i>	35
Haw, Apple.....	82	<i>Ilex Canadensis</i>	35

	Page.		Page.
King Nut.....	124	Loblolly Pine.....	197
Knackaway	114	Locust (Robinia Neo-Mexicana)	56
Knob-cone Pine.....	196	Locust (Robinia Pseudacacia)	55
L.			
<i>Lacathea florida</i>	25	Locust, Black (Gleditschia triacanthos)	59
<i>Laguncularia glabrifolia</i>	87	Locust, Black (Robinia Pseudacacia)	55
<i>Laguncularia racemosa</i>	87	Locust, Clammy	56
Lancewood	119	Locust, Honey (Gleditschia triacanthos)	59
Larch	215	Locust, Honey (Prosopis juliflora)	62
Larch, Black	215	Locust, Sweet	59
Large-leaved Cucumber Tree.....	21	Locust, Water	60
Large Tupelo.....	93	Locust, Yellow	55
<i>Larix Americana</i>	215	Lodge-pole Pine	196
<i>Larix Americana rubra</i>	215	Logwood.....	40
<i>Larix Americana</i> , var. <i>brevipolia</i>	216	Long-leaved Cucumber Tree.....	22
<i>Larix Americana</i> , var. <i>pendula</i>	215	Long-leaved Pine	202
<i>Larix Americana</i> , var. <i>prolifera</i>	215	<i>Lyonia arborea</i>	96
<i>Larix decidua</i> , var. <i>Americana</i>	215	<i>Lyonia ferruginea</i>	96
<i>Larix intermedia</i>	215	<i>Lyonia rigida</i>	96
<i>Larix Lyallii</i>	216	<i>Lysiloma Bahamensis</i>	64
<i>Larix microcarpa</i>	215	<i>Lysiloma latifolia</i>	64
<i>Larix occidentalis</i>	216	M.	
<i>Larix pendula</i>	215	<i>Maclura aurantiaca</i>	128
<i>Larix tenuifolia</i>	215	Madeira	34
LAURACEÆ	118-120	Madroña.....	97
Laurel	98	<i>Magnolia acuminata</i>	20
Laurel, Rig	19	<i>Magnolia auricularis</i>	22
Laurel, California.....	120	<i>Magnolia auriculata</i>	22
Laurel, Great	99	<i>Magnolia cordata</i>	20
Laurel, Mountain.....	120	<i>Magnolia De Candollii</i>	20
Laurel Oak (<i>Quercus imbricaria</i>)	154	<i>Magnolia fragrans</i>	20
Laurel Oak (<i>Quercus laurifolia</i>)	158	<i>Magnolia Fraseri</i>	22
Laurel, Swamp	20	<i>Magnolia glauca</i>	19
Laurel, White	20	<i>Magnolia glauca</i> , var. <i>latifolia</i>	20
<i>Laurocerasus Caroliniana</i>	69	<i>Magnolia glauca</i> , var. <i>longifolia</i>	20
<i>Laurocerasus ilicifolia</i>	70	<i>Magnolia grandiflora</i>	19
<i>Laurus Borbonia</i>	118	<i>Magnolia grandiflora</i> , var. <i>elliptica</i> and <i>obovata</i>	19
<i>Laurus Caroliniana</i>	118	<i>Magnolia grandiflora</i> , var. <i>lanceolata</i>	19
<i>Laurus Carolinensis</i>	118	<i>Magnolia longifolia</i>	20
<i>Laurus Carolinensis</i> , var. <i>glabra</i>	118	<i>Magnolia macrophylla</i>	21
<i>Laurus Carolinensis</i> , var. <i>obtus.</i>	118	<i>Magnolia, Mountain</i>	20
<i>Laurus Carolinensis</i> , var. <i>pubescens</i>	119	<i>Magnolia pyramidalis</i>	22
<i>Laurus Catesbaei</i>	119	<i>Magnolia tripetala</i>	21
<i>Laurus Catesbyana</i>	119	<i>Magnolia Umbrella</i>	21
<i>Laurus regia</i>	120	<i>Magnolia Virginiana</i> , var. <i>a. glauca</i>	20
<i>Laurus sanguinea</i>	119	<i>Magnolia Virginiana</i> , var. <i>β. fastida</i>	19
<i>Laurus Sassafras</i>	119	<i>Magnolia Virginiana</i> , var. <i>c.</i>	20
Lawson's Cypress	179	<i>Magnolia Virginiana</i> , var. <i>tripetala</i>	21
Leaf, Sweet	105	MALVACEÆ	19-22
LEGUMINOSÆ	55-64	Mahogany.....	34
<i>Leptocarpa Caroliniana</i>	69	Mahogany Birch.....	162
<i>Leucena glauca</i>	62	Mahogany, Mountain (<i>Cercocarpus ledifolius</i>)	71
<i>Leucena pulverulenta</i>	63	Mahogany, Mountain (<i>Cercocarpus parvifolius</i>)	71
Leverwood	158	MALPIGHIACEÆ	28
<i>Libocedrus decurrens</i>	176	<i>Malpighia lucida</i>	28
<i>Lignum-vitæ</i>	28	<i>Malus angustifolia</i>	72
LILIACEÆ	218, 219	<i>Malus coronaria</i>	73
Lime, Ogeechee	91	<i>Malus discolorifolia</i>	73
Lime Tree	27	<i>Malus microcarpa coronaria</i>	73
Lime, Wild (<i>Xanthoxylum Pterota</i>)	81	<i>Malus rivularis</i>	73
Lime, Wild (<i>Ximenia Americana</i>)	84	<i>Malus sempervirens</i>	73
Lín	27	<i>Malus subcordata</i>	73
Linden, American	27	Manchineel	121
<i>Liquidambar macrophylla</i>	86	Manchineel, Mountain.....	54
<i>Liquidambar Styraciflua</i>	86	<i>Mancinella venenata</i>	121
<i>Liquidambar Styraciflua</i> , var. <i>Mexicana</i>	86	Mangrove	87
<i>Liquidamber</i>	86	Mangrove, Black	117
<i>Liriodendron procera</i>	22	Mangrove, White.....	87
<i>Liriodendron Tulipifera</i>	22	Maple, Ash-leaved.....	51
Live Oak (<i>Quercus chrysolepis</i>)	146	Maple, Black Sugar	49
Live Oak (<i>Quercus virens</i>)	145	Maple, Broad-leaved.....	47
Live Oak (<i>Quercus Wlailzenii</i>)	147	Maple, Dwarf	46
Live Oak, Coast	147	Maple, Goose-foot	46
Loblolly Bay	25	Maple, Hard.....	46
		Maple, Mountain.....	46

	Page.		Page.
Maple, Red	50	<i>Mespilus Watsoniana</i>	76
Maple, Rock	48	? <i>Mespilus Wendlandii</i>	77
Maple, Silver	49	Mesquit	62
Maple, Soft (<i>Acer dasycarpum</i>)	49	Mesquit, Screw-pod	62
Maple, Soft (<i>Acer rubrum</i>)	50	<i>Melospium Linnæi</i>	54
Maple, Striped	48	Mexican Banana	219
Maple, Sugar	48	Mexican Mulberry	126
Maple, Swamp	50	Mexican Persimmon	105
Maple, Vine	47	<i>Michauxia sessilis</i>	25
Maple, Water	50	<i>Mimosa biceps</i>	62
Maple, White	49	<i>Mimosa frondosa</i>	62
Marlberry	100	<i>Mimosa glauca</i>	62
Mastic	101	<i>Mimosa Guadalupeensis</i>	64
Manul Oak	146	<i>Mimosa latifolia</i>	64
May Cherry	84	<i>Mimosa leucocephala</i>	62
May Haw	82	<i>Mimosa rosea</i>	64
Meadow Pine	202	<i>Mimosa Unguis-cati</i>	64
MELIACEÆ	83, 84	<i>Mimusops dissecta</i>	103
<i>Melicocca paniculata</i>	45	<i>Mimusops Sieberi</i>	108
<i>Melilobus heterophylla</i>	59	Mocker Nut	124
<i>Mespilus acerifolia</i>	80	Mock Orange	70
<i>Mespilus acuticulis</i>	82	Monterey Cypress	179
<i>Mespilus apifolia</i>	81	Monterey Pine	196
<i>Mespilus arborea</i>	84	Moose Elm	123
<i>Mespilus arbutifolia</i>	83	Moosewood	46
<i>Mespilus Azarolus</i>	81	<i>Morus Canadensis</i> (Lamarck and Rafinesque)	127
<i>Mespilus berberifolia</i>	82	<i>Morus microphylla</i>	128
<i>Mespilus Bosciiana</i>	77	<i>Morus Missouriensis</i>	127
<i>Mespilus Calpodendron</i>	79	<i>Morus parvifolia</i> (<i>Morus microphylla</i>)	128
<i>Mespilus Canadensis</i>	84	<i>Morus parvifolia</i> (<i>Morus rubra</i>)	127
<i>Mespilus Canadensis</i> , var. <i>cordata</i>	84	<i>Morus reticulata</i>	127
<i>Mespilus Canadensis</i> , var. <i>obovata</i>	85	<i>Morus riparia</i>	127
<i>Mespilus Caroliniana</i>	82	<i>Morus rubra</i>	127
<i>Mespilus coccinea</i>	77	<i>Morus rubra</i> , var. <i>Canadensis</i>	127
<i>Mespilus cordata</i>	80	<i>Morus rubra</i> , var. <i>incisa</i>	127
<i>Mespilus cornifolia</i>	80	<i>Morus rubra</i> , var. <i>tomentosa</i>	127
<i>Mespilus Crus-galli</i>	76	<i>Morus scabra</i>	127
<i>Mespilus Crus-galli</i> , var. <i>pyracanthifolia</i>	76	<i>Morus tomentosa</i>	127
<i>Mespilus Crus-galli</i> , var. <i>salicifolia</i>	76	Mossy-cup Oak	146
<i>Mespilus cuneifolia</i> (<i>Cratægus Crus-galli</i>)	76	Mountain Ash (<i>Pyrus Americana</i>)	78
<i>Mespilus cuneifolia</i> (<i>Cratægus tomentosa</i> var. <i>punctata</i>)	80	Mountain Ash (<i>Pyrus sambucifolia</i>)	79
? <i>Mespilus cuneiformis</i>	76	Mountain Laurel	120
<i>Mespilus elliptica</i> (<i>Cratægus Crus-galli</i>)	76	Mountain Magnolia	20
<i>Mespilus elliptica</i> (<i>Cratægus flava</i> , var. <i>pubescens</i>)	83	Mountain Mahogany (<i>Cercocarpus ledifolius</i>)	71
<i>Mespilus flabellata</i>	77	Mountain Mahogany (<i>Cercocarpus parvifolius</i>)	71
<i>Mespilus flava</i>	82	Mountain Manchinel	54
<i>Mespilus flexispina</i>	82	Mountain Maple	46
<i>Mespilus glandulosa</i>	77	Mountain Plum	84
<i>Mespilus hiemalis</i>	83	Mountain White Oak	143
<i>Mespilus latifolia</i>	79	Mulberry, Mexican	128
<i>Mespilus linearis</i>	77	Mulberry, Red	128
<i>Mespilus lobata</i>	79	<i>Myginda pallens</i>	88
<i>Mespilus lucida</i>	76	<i>Mylocaryum ligustrinum</i>	88
<i>Mespilus lucida</i> , var. <i>angustifolia</i>	77	<i>Myrica Californica</i>	127
<i>Mespilus Michauxii</i>	83	<i>Myrica Carolinensis</i>	126
<i>Mespilus monogyna</i> , var. <i>apifolia</i>	81	<i>Myrica cerifera</i>	126
<i>Mespilus nivea</i>	84	<i>Myrica cerifera humilis</i>	126
<i>Mespilus odorata</i>	79	<i>Myrica cerifera sempervirens</i>	126
<i>Mespilus ovalifolia</i>	76	<i>Myrica cerifera</i> , var. <i>angustifolia</i>	126
<i>Mespilus Phænopyrum</i>	80	<i>Myrica cerifera</i> , var. <i>arborescens</i>	126
<i>Mespilus populiifolia</i>	78	<i>Myrica cerifera</i> , var. <i>latifolia</i>	126
<i>Mespilus pruinosa</i>	79	<i>Myrica cerifera</i> , var. <i>media</i>	126
<i>Mespilus prunellifolia</i>	76	<i>Myrica cerifera</i> , var. <i>pumila</i>	126
<i>Mespilus prunifolia</i> ?	77	<i>Myrica Pennsylvanica</i>	126
<i>Mespilus pubescens</i>	77	? <i>Myrica Xalapensis</i>	127
<i>Mespilus punctata</i>	80	MYRICACEÆ	126, 127
<i>Mespilus pyrifolia</i> (<i>Cratægus tomentosa</i>)	79	MYRSINACEÆ	99, 100
<i>Mespilus pyrifolia</i> (<i>Cratægus tomentosa</i> , var. <i>punctata</i>)	80	<i>Myrsine floribunda</i>	99
<i>Mespilus rotundifolia</i> (<i>Cratægus coccinea</i>)	77	<i>Myrsine Florida</i>	99
<i>Mespilus rotundifolia</i> (<i>Cratægus Crus-galli</i> , var. <i>prunifolia</i>)	77	<i>Myrsine Rapanæ</i>	99
<i>Mespilus salicifolia</i>	76	MYRTACEÆ	83, 89
<i>Mespilus spathulata</i>	81	Myrtle, Blue	41
<i>Mespilus tiliaefolia</i>	78	Myrtle, Wax	126
<i>Mespilus turbinata</i>	82	<i>Myrtus asilaris</i>	88
<i>Mespilus viridis</i>	78	<i>Myrtus buxifolia</i>	88

	Page.		Page.
<i>Myrtus Oxytrachelis</i>	88	Oak, Chestnut (<i>Quercus Prinns</i>)	142
<i>Myrtus dichotoma</i>	88	Oak, Chinquapin	143
<i>Myrtus monticola</i>	89	Oak, Coast Live	147
<i>Myrtus Poireri</i>	88	Oak, Cow	141
<i>Myrtus procera</i>	89	Oak, Duck	152
		Oak, Iron	189
N.		Oak, Jack	150
Naked Wood (<i>Colubrina reclinata</i>)	42	Oak, Laurel (<i>Quercus imbricaria</i>)	154
Naked Wood (<i>Eugenia dichotoma</i>)	88	Oak, Laurel (<i>Quercus laurifolia</i>)	153
Nannyberry	94	Oak, Live (<i>Quercus chrysolepis</i>)	146
Necklace Poplar	175	Oak, Live (<i>Quercus virens</i>)	145
<i>Nectandra Bredemeyeri</i>	119	Oak, Live (<i>Quercus Wailsoni</i>)	147
<i>Nectandra Willdenoviana</i>	119	Oak, Maul	146
<i>Negundium fraxinifolium</i>	51	Oak, Mosey-cup	140
<i>Negundo aceroides</i>	50	Oak, Mountain White	143
<i>Negundo aceroides</i> (<i>Negundo Californicum</i>)	51	Oak, Over-cup (<i>Quercus lyrata</i>)	140
<i>Negundo Californicum</i>	51	Oak, Over-cup (<i>Quercus macrocarpa</i>)	140
<i>Negundo Californicum</i> (<i>Negundo aceroides</i>)	51	Oak, Peach (<i>Quercus densiflora</i>)	155
<i>Negundo fraxinifolium</i>	51	Oak, Peach (<i>Quercus Phellos</i>)	154
<i>Negundo lobatum</i>	51	Oak, Pin	153
<i>Negundo Mezianum</i>	51	Oak, Possum	153
<i>Negundo trifoliatum</i>	51	Oak, Post	139
Newcastle Thorn	76	Oak, Punk	153
Norway Pine	192	Oak, Quercitron	149
Nut, Bitter	135	Oak, Red (<i>Quercus falcata</i>)	151
Nut, Bull	134	Oak, Red (<i>Quercus rubra</i>)	148
Nut, Coffee	58	Oak, Red (<i>Quercus rubra</i> , var. <i>Texana</i>)	148
Nut, Illinois	182	Oak, Rock Chestnut	142
Nut, King	184	Oak, Scarlet	148
Nut, Mocker	184	Oak, Scrub (<i>Quercus Catesbei</i>)	151
Nut, Pig	184	Oak, Scrub (<i>Quercus undulata</i> , var. <i>Gambelli</i>)	139
Nut Pine (<i>Pinus cembroides</i>)	190	Oak, Shingle	151
Nut Pine (<i>Pinus edulis</i>)	190	Oak, Spanish	151
Nut Pine (<i>Pinus monophylla</i>)	190	Oak, Swamp Post	140
Nut Pine (<i>Pinus Parryana</i>)	189	Oak, Swamp Spanish	153
Nut, Tallow	84	Oak, Swamp White	141
Nutmeg, California	186	Oak, Tanbark	155
Nutmeg, Hickory	185	Oak, Turkey	151
NYCTAGINACEÆ	117	Oak, Upland Willow	153
<i>Nyssa aquatica</i> (<i>Nyssa sylvatica</i>)	92	Oak, Valparaiso	146
<i>Nyssa aquatica</i> (<i>Nyssa uniflora</i>)	92	Oak, Water (<i>Quercus aquatica</i>)	152
<i>Nyssa angustiana</i>	98	Oak, Water (<i>Quercus palustris</i>)	152
<i>Nyssa angulosa</i>	98	Oak, Water White	140
<i>Nyssa biflora</i>	92	Oak, Weeping	188
<i>Nyssa Canadensis</i>	92	Oak, White (<i>Quercus alba</i>)	137
<i>Nyssa candicans</i>	91	Oak, White (<i>Quercus Garryana</i>)	138
<i>Nyssa capitata</i>	91	Oak, White (<i>Quercus grisea</i>)	144
<i>Nyssa capitata</i> , var. <i>grandidentata</i>	93	Oak, White (<i>Quercus lobata</i>)	138
<i>Nyssa Caroliniana</i>	92	Oak, White (<i>Quercus oblongifolia</i>)	144
<i>Nyssa coccinea</i>	91	Oak, Willow	154
<i>Nyssa denticulata</i>	92	Oak, Yellow (<i>Quercus prinoides</i>)	148
<i>Nyssa grandidentata</i>	98	Oak, Yellow (<i>Quercus tinctoria</i>)	149
<i>Nyssa integrifolia</i>	92	Oak, Yellow-bark	149
<i>Nyssa montana</i>	91	Obispo Pine	200
<i>Nyssa multiflora</i>	92	<i>Enocarpus regia</i>	218
<i>Nyssa multiflora</i> , var. <i>sylvatica</i>	92	Ogechee Lime	91
<i>Nyssa Ogeche</i>	91	Ohio Buckeye	42
<i>Nyssa palustris</i>	98	OLACINÆ	34
<i>Nyssa sylvatica</i>	92	Old-field Birch	159
<i>Nyssa tomentosa</i> (<i>Nyssa capitata</i>)	91	Old-field Pine	197
<i>Nyssa tomentosa</i> (<i>Nyssa uniflora</i>)	93	Old Man's Beard	113
<i>Nyssa uniflora</i>	92	<i>Olea Americana</i>	113
<i>Nyssa villosa</i>	92	OLEACEÆ	106-113
		Olive, California	120
		Olneya Tesota	56
Oak, Bartram's	153	Orange, Mock	70
Oak, Basket	141	Orange, Osage	128
Oak, Black (<i>Quercus Emoryi</i>)	146	Orange, Wild (<i>Prunus Caroliniana</i>)	70
Oak, Black (<i>Quercus Kelloggii</i>)	149	Orange, Wild (<i>Xanthoxylum Clava-Herculis</i>)	80
Oak, Black (<i>Quercus rubra</i>)	148	<i>Orchidocarpum aristatum</i>	23
Oak, Black (<i>Quercus tinctoria</i>)	149	Oregon Ash	111
Oak, Blue	143	Oregon Cedar	179
Oak, Burr	140	Oregon Crab Apple	73
Oak, Chestnut (<i>Quercus densiflora</i>)	155	Oregon Pine	209
Oak, Chestnut (<i>Quercus prinoides</i>)	143	<i>Oreodaphne Californica</i>	120

	Page.		Page.
<i>Oreodoxa oleracea?</i>	218	<i>Photinia arbutifolia</i>	83
<i>Oreodoxa regia</i>	218	<i>Photinia salicifolia</i>	83
<i>Osage Orange</i>	128	<i>Picea alba</i>	204
<i>Osmanthus Americanus</i>	113	<i>Picea amabilis</i> (<i>Abies amabilis</i>)	213
<i>Ostrya Virginiana</i>	158	<i>Picea amabilis</i> (<i>Abies subalpina</i>)	211
<i>Ostrya Virginica</i>	158	<i>Picea balsamea</i>	211
<i>Ostrya Virginica</i> , var. <i>eglandulosa</i>	158	<i>Picea balsamea</i> , var. <i>longifolia</i>	211
<i>Ostrya Virginica</i> , var. <i>glandulosa</i>	158	<i>Picea bifolia</i>	211
Over-cup Oak (<i>Quercus lyrata</i>)	140	<i>Picea bracteata</i>	213
Over-cup Oak (<i>Quercus macrocarpa</i>)	140	<i>Picea Californica</i>	208
<i>Oxydendrum arboreum</i>	98	<i>Picea Canadensis</i>	206
P.			
<i>Padus cartilaginea</i>	68	<i>Picea canadensis</i>	204
<i>Padus demissa</i>	69	<i>Picea concolor</i>	212
<i>Padus serotina</i>	68	<i>Picea concolor</i> , var. <i>violacea</i>	212
<i>Padus Virginiana</i>	68	<i>Picea Douglasii</i>	209
Palm, Fan-leaf	217	<i>Picea Engelmanni</i>	205
Palm, Royal	218	<i>Picea Fraseri</i> (<i>Abies balsamea</i>)	211
<i>Palma argentea</i>	218	<i>Picea Fraseri</i> (<i>Abies Fraseri</i>)	210
PALMACEAE	217, 218	<i>Picea glauca</i>	204
Palmetto, Cabbage	217	<i>Picea grandis</i> (<i>Abies concolor</i>)	213
Palmetto, Silk-top	217	<i>Picea grandis</i> (<i>Abies grandis</i>)	212
Palmetto, Silver-top	218	<i>Picea laza</i>	204
Palo Blanco	126	<i>Picea Lowiana</i>	212
Palo Verde	60	<i>Picea magnifica</i>	214
Pupaw	23	<i>Picea Menziesii</i> (<i>Picea pungens</i>)	205
Paper Birch	100	<i>Picea Menziesii</i> (<i>Picea Sitchenensis</i>)	206
Paradise Tree	32	<i>Picea nigra</i>	202
<i>Parkinsonia aculeata</i>	60	<i>Picea nigra</i> , var. <i>glauca</i>	204
<i>Parkinsonia microphylla</i>	60	<i>Picea nigra</i> , var. <i>rubra</i>	203
<i>Parkinsonia Torreyana</i>	60	<i>Picea nobilis</i>	214
Parsley Haw	81	<i>Picea pungens</i>	205
<i>Pasania densiflora</i>	155	<i>Picea rubra</i>	203
<i>Pavia Californica</i>	43	<i>Picea Sitchenensis</i>	206
<i>Pavia carnea</i>	42	<i>Pickeringia paniculata</i>	100
<i>Pavia discolor</i>	43	Pigeon Cherry	66
<i>Pavia flava</i>	43	Pigeon Plum	117
<i>Pavia glabra</i>	42	Pigeonwood	117
<i>Pavia hybrida</i>	43	Pig Nut	124
<i>Pavia lutea</i>	43	<i>Pinocereus Engelmanni</i>	80
<i>Pavia neglecta</i>	43	Pin Cherry	66
<i>Pavia pallida</i>	42	Pin Oak	152
<i>Pavia Watsoniana</i>	42	<i>Pinckneya pubens</i>	95
Peach Oak (<i>Quercus densiflora</i>)	155	<i>Pinckneya pubescens</i>	95
Peach Oak (<i>Quercus Phellos</i>)	154	Pine, Bastard	202
Peach, Wild	70	Pine, Bishop's	200
Pear Haw	79	Pine, Black (<i>Pinus Jeffreyi</i>)	193
Pecan	132	Pine, Black (<i>Pinus Murrayana</i>)	195
Pecan, Bitter	136	Pine, Bull (<i>Pinus Jeffreyi</i>)	198
Pepperidge	92	Pine, Bull (<i>Pinus mitis</i>)	200
Pepperwood	30	Pine, Bull (<i>Pinus ponderosa</i>)	193
<i>Persea Borbonia</i>	118	Pine, Bull (<i>Pinus Sabiniana</i>)	195
<i>Persea Carolinensis</i>	118	Pine, Cedar	201
<i>Persea Carolinensis</i> , var. <i>glabrescens</i>	118	Pine, Digger	195
<i>Persea Carolinensis</i> , var. <i>palustris</i>	119	Pine, Foxtail	191
<i>Persea Carolinensis</i> , var. <i>pubescens</i>	119	Pine, Georgia	202
<i>Persea Catesbyana</i>	119	Pine, Ginger	179
<i>Persea Sassafras</i>	119	Pine, Gray	201
Persimmon	104	Pine, Hard	202
Persimmon, Black	105	Pine, Hickory (<i>Pinus Balfouriana</i> , var. <i>aristata</i>)	191
Persimmon, Mexican	105	Pine, Hickory (<i>Pinus pungens</i>)	199
<i>Phænopyrum acerifolium</i>	80	Pine, Jersey	199
<i>Phænopyrum arborescens</i>	75	Pine, Knob-cone	196
<i>Phænopyrum Carolinianum</i>	82	Pine, Loblolly	197
<i>Phænopyrum coccineum</i>	77	Pine, Lodge-pole	195
<i>Phænopyrum cordatum</i>	80	Pine, Long-leaved	202
<i>Phænopyrum ellipticum</i>	82	Pine, Meadow	202
<i>Phænopyrum populifolium</i>	78	Pine, Monterey	196
<i>Phænopyrum pruinosum</i>	79	Pine, Norway	192
<i>Phænopyrum spathulatum</i>	81	Pine, Nut (<i>Pinus oemboideae</i>)	190
<i>Phænopyrum subvillosum</i>	78	Pine, Nut (<i>Pinus edulis</i>)	190
<i>Phænopyrum Virginicum</i>	82	Pine, Nut (<i>Pinus monophylla</i>)	190
<i>Phænopyrum viride</i>	78	Pine, Nut (<i>Pinus Parryana</i>)	189
<i>Phænopyrum Wendlandii</i>	77	Pine, Obispo	200
		Pine, Old-field	197
		Pine, Oregon	200

	Page.		Page.
Pine, Pitch	198	<i>Pinus contorta</i> (Pinus Murrayana)	194
Pine, Pond	198	<i>Pinus contorta</i> , var. <i>Bolanderi</i>	194
Pine, Prince's	201	<i>Pinus contorta</i> , var. <i>latifolia</i>	194
Pine, Red	192	<i>Pinus Coulteri</i>	195
Pine, Rosemary	197	<i>Pinus Oraigana</i>	193
Pine, Sand	199	<i>Pinus Cubensis</i>	202
Pine, Scrub (Pinus Banksiana)	201	<i>Pinus Oubensis</i> , var. <i>terthrocarpa</i>	202
Pine, Scrub (Pinus clausa)	199	<i>Pinus deflexa</i>	193
Pine, Scrub (Pinus contorta)	194	<i>Pinus Douglasii</i>	209
Pine, Scrub (Pinus inops)	199	<i>Pinus Douglasii</i> , var. <i>brevibracteata</i>	209
Pine, Short-leaved	200	<i>Pinus echinata</i>	200
Pine, Slash	202	<i>Pinus Edgariana</i>	199
Pine, Southern	202	<i>Pinus edulis</i>	199
Pine, Spruce (Pinus clausa)	199	<i>Pinus Elliottii</i>	202
Pine, Spruce (Pinus glabra)	201	<i>Pinus Engelmanni</i> (Picea Engelmanni)	205
Pine, Spruce (Pinus mitis)	200	<i>Pinus Engelmanni</i> (Pinus ponderosa)	193
Pine, Spruce (Pinus Murrayana)	195	<i>Pinus flexilis</i>	188
Pine, Sugar	188	<i>Pinus flexilis</i> (Pinus albicaulis)	189
Pine, Swamp	202	<i>Pinus flexilis</i> , var. <i>albicaulis</i>	189
Pine, Table-mountain	199	<i>Pinus flexilis</i> , var. <i>macrocarpa</i>	188
Pine, Weymouth	187	<i>Pinus flexilis</i> , var. <i>reflexa</i>	189
Pine, White (Pinus flexilis)	188	<i>Pinus flexilis</i> , var. <i>serrulata</i>	188
Pine, White (Pinus glabra)	201	<i>Pinus Fraseri</i> (Abies Fraseri)	210
Pine, White (Pinus monticola)	187	<i>Pinus Fraseri</i> (Pinus rigida)	197
Pine, White (Pinus reflexa)	189	<i>Pinus Fremontiana</i>	190
Pine, White (Pinus Strobus)	187	<i>Pinus futilis</i>	190
Pine, Yellow (Pinus Arizonica)	192	<i>Pinus glabra</i>	200
Pine, Yellow (Pinus mitis)	209	<i>Pinus grandis</i> (Abies amabilis)	213
Pine, Yellow (Pinus palustris)	202	<i>Pinus grandis</i> (Abies concolor)	212
Pine, Yellow (Pinus ponderosa)	193	<i>Pinus grandis</i> (Abies grandis)	212
Piñon (Pinus edulis)	190	<i>Pinus Grozelleri</i>	187
Piñon (Pinus monophylla)	190	<i>Pinus Hudsonica</i>	201
Piñon (Pinus Parryana)	189	<i>Pinus inops</i>	198
<i>Pinus Abies Americana</i>	206	<i>Pinus inops</i> (Pinus contorta)	194
<i>Pinus Abies Balsamea</i>	211	<i>Pinus inops</i> (Pinus Murrayana)	194
<i>Pinus Abies Canadensis</i>	202	<i>Pinus inops</i> , var. (Pinus muricata)	199
<i>Pinus adunca</i>	196	<i>Pinus inops</i> , var. <i>clausa</i>	199
<i>Pinus alba</i>	204	<i>Pinus insignis</i>	196
<i>Pinus albicaulis</i>	189	<i>Pinus insignis macrocarpa</i>	196
<i>Pinus amabilis</i> (Abies amabilis)	213	<i>Pinus insignis</i> , var. <i>binata</i>	196
<i>Pinus amabilis</i> (Abies magnifica)	214	<i>Pinus intermedia</i>	215
<i>Pinus amabilis</i> (Abies subalpina)	211	<i>Pinus Jeffreyi</i>	193
<i>Pinus Americana</i> (Picea nigra)	203	<i>Pinus Lambertiana</i>	188
<i>Pinus Americana</i> (Tauga Canadensis)	206	<i>Pinus Lambertiana</i> , var. (Pinus flexilis)	188
<i>Pinus Americana rubra</i>	202	<i>Pinus Lambertiana</i> , var. <i>brevifolia</i>	188
<i>Pinus aristata</i>	191	<i>Pinus laricina</i>	215
<i>Pinus Arizonica</i>	192	<i>Pinus Laricio</i> , var. <i>resinosa</i>	191
<i>Pinus australis</i>	201	<i>Pinus Larix</i>	216
<i>Pinus Balfouriana</i>	191	<i>Pinus Larix alba</i>	215
<i>Pinus Balfouriana</i> (Pinus Balfouriana, var. <i>aristata</i>)	191	<i>Pinus Larix nigra</i>	215
<i>Pinus Balfouriana</i> , var. <i>aristata</i>	191	<i>Pinus Larix rubra</i>	215
<i>Pinus balsamea</i>	210	<i>Pinus lasiocarpa</i> (Abies concolor)	212
<i>Pinus balsamea</i> , var. <i>Fraseri</i>	210	<i>Pinus lasiocarpa</i> (Abies subalpina)	211
<i>Pinus Banksiana</i>	201	<i>Pinus Llaveana</i> (Pinus cembroides)	190
<i>Pinus Banksiana</i> (Pinus contorta)	194	<i>Pinus Llaveana</i> (Pinus Parryana)	189
<i>Pinus Beardsleyi</i>	193	<i>Pinus Loddigesii</i>	197
<i>Pinus Benthamiana</i>	193	<i>Pinus lophosperma</i>	192
<i>Pinus Bolanderi</i>	194	<i>Pinus Louiana</i>	212
<i>Pinus Boursieri</i>	194	<i>Pinus Lyallii</i>	216
<i>Pinus brachyptera</i>	193	<i>Pinus Mariana</i>	202
<i>Pinus bracteata</i>	213	<i>Pinus macrocarpa</i>	195
<i>Pinus Californica</i> (Pinus insignis)	196	<i>Pinus macrophylla</i> ?	193
<i>Pinus Californica</i> (Pinus tuberculata)	196	<i>Pinus Menziesii</i>	206
<i>Pinus Canadensis</i> (Picea alba)	204	<i>Pinus Menziesii</i> , var. <i>crispa</i>	206
<i>Pinus Canadensis</i> (Tauga Canadensis)	206	<i>Pinus Mertensiana</i>	207
<i>Pinus Canadensis</i> (Tauga Mertensiana)	207	<i>Pinus microcarpa</i>	215
<i>Pinus cembroides</i>	190	<i>Pinus mitis</i>	200
<i>Pinus cembroides</i> (Pinus albicaulis)	189	<i>Pinus mitis</i> , var. <i>paupers</i>	200
<i>Pinus cembroides</i> (Pinus edulis)	190	<i>Pinus monophylla</i>	190
<i>Pinus Chihuahuana</i>	194	<i>Pinus monticola</i>	187
<i>Pinus clausa</i>	199	<i>Pinus muricata</i>	199
<i>Pinus commutata</i>	205	<i>Pinus muricata</i> (Pinus contorta)	194
<i>Pinus concolor</i>	212	<i>Pinus Murrayana</i>	194
<i>Pinus contorta</i>	194	<i>Pinus nigra</i>	202
<i>Pinus contorta</i> (Pinus muricata)	199	<i>Pinus nobilis</i>	214

	Page.		Page.
<i>Pinus Nuttallii</i>	216	<i>Platanus lobata</i>	129
<i>Pinus obovata</i>	190	<i>Platanus Mexicana</i> (<i>Platanus racemosa</i>)	129
<i>Pinus palustris</i>	201	<i>Platanus Mexicana</i> (<i>Platanus Wrightii</i>)	129
<i>Pinus Parryana</i>	189	<i>Platanus occidentalis</i>	129
<i>Pinus Parryana</i> (<i>Pinus ponderosa</i>)	193	<i>Platanus occidentalis</i> (<i>Platanus racemosa</i>)	129
<i>Pinus Pattoniana</i> (<i>Tunga Mertensiana</i>)	208	<i>Platanus racemosa</i>	129
<i>Pinus Pattoniana</i> (<i>Tunga Pattoniana</i>)	208	<i>Platanus racemosa</i> (<i>Platanus Wrightii</i>)	129
<i>Pinus pendula</i>	215	<i>Platanus vulgaris</i> , var. <i>angulosa</i>	129
<i>Pinus ponderosa</i>	192	<i>Platanus Wrightii</i>	129
<i>Pinus ponderosa</i> , var. <i>Benthiana</i>	193	<i>Plum, Canada</i>	65
<i>Pinus ponderosa</i> , var. <i>Jeffreyi</i>	193	<i>Plum, Chickasaw</i>	65
<i>Pinus ponderosa</i> , var. <i>scopulorum</i>	193	<i>Plum, Cocoa</i>	65
<i>Pinus porphyrocarpa</i>	187	<i>Plum, Darling</i>	39
<i>Pinus pungens</i>	199	<i>Plum, Downward</i>	103
<i>Pinus radiata</i>	196	<i>Plum, Gopher</i>	91
<i>Pinus reflexa</i>	189	<i>Plum, Guiana</i>	121
<i>Pinus resinosa</i>	191	<i>Plum, Hog</i> (<i>Prunus angustifolia</i>)	65
<i>Pinus resinosa</i> (<i>Pinus ponderosa</i>)	193	<i>Plum, Hog</i> (<i>Rhus Metopium</i>)	54
<i>Pinus rigida</i>	197	<i>Plum, Hog</i> (<i>Ximenia Americana</i>)	34
<i>Pinus rigida</i> ? (<i>Pinus insignis</i>)	196	<i>Plum, Horse</i>	65
<i>Pinus rigida</i> (<i>Pinus mitis</i>)	200	<i>Plum, Mountain</i>	34
<i>Pinus rigida</i> , var. <i>serotina</i>	198	<i>Plum, Pigeon</i>	117
<i>Pinus rubra</i> (<i>Picea nigra</i>)	203	<i>Plum, Saffron</i>	103
<i>Pinus rubra</i> (<i>Pinus resinosa</i>)	191	<i>Plum, Wild</i>	65
<i>Pinus rubra</i> , var. <i>violacea</i>	204	<i>Poison Elder</i>	54
<i>Pinus rupestris</i>	201	<i>Poison Samach</i>	54
<i>Pinus Sabinalana</i>	195	<i>Poisonwood</i> (<i>Rhus Metopium</i>)	54
<i>Pinus Sabintana Coultteri</i>	195	<i>Poisonwood</i> (<i>Sebastiania lucida</i>)	121
<i>Pinus Sabintana macrocarpa</i>	195	POLYGONACEÆ	117, 118
<i>Pinus serotina</i>	198	<i>Polygonum uvifera</i>	118
<i>Pinus Shasta</i>	189	<i>Pond Apple</i>	23
<i>Pinus Sinclairii</i>	196	<i>Pond Pine</i>	198
<i>Pinus Sitchensis</i>	206	<i>Poplar</i>	172
<i>Pinus species</i> (<i>Abies subalpina</i>)	211	<i>Poplar, Carolina</i>	175
<i>Pinus Strobus</i>	187	<i>Poplar, Neoklaco</i>	175
<i>Pinus Strobus</i> , var. <i>alba</i>	187	<i>Poplar, Yellow</i>	22
<i>Pinus Strobus</i> , var. <i>brevifolia</i>	187	<i>Populus acledesca</i>	173
<i>Pinus Strobus</i> , var. <i>compressa</i>	187	<i>Populus angulata</i>	175
<i>Pinus Strobus</i> , var. <i>monticola</i>	187	<i>Populus angulosa</i>	175
<i>Pinus Strobus</i> , var. <i>nivea</i>	187	<i>Populus angustifolia</i>	174
<i>Pinus sylvestris</i> , var. <i>disaricata</i>	201	<i>Populus angustifolia</i> (<i>Populus trichocarpa</i>)	174
<i>Pinus Tæda</i>	197	<i>Populus argentea</i>	172
<i>Pinus Tæda</i> , var. <i>a</i>	197	<i>Populus atheniensis</i>	171
<i>Pinus Tæda</i> , var. <i>alopeuroidea</i>	198	<i>Populus balsamifera</i>	173
<i>Pinus Tæda</i> , var. <i>heterophylla</i>	202	<i>Populus balsamifera</i> (<i>Populus trichocarpa</i>)	174
<i>Pinus Tæda</i> , var. <i>rigida</i>	197	<i>Populus balsamifera lanceolata</i>	173
<i>Pinus Tæda</i> , var. <i>tenuifolia</i>	197	<i>Populus balsamifera</i> , var.	174
<i>Pinus Tæda</i> , var. <i>variabilis</i>	200	<i>Populus balsamifera</i> , var. <i>angustifolia</i>	174
<i>Pinus Tæda</i> , var. <i>Virginiana</i>	198	<i>Populus balsamifera</i> , var. <i>! Californica</i>	174
<i>Pinus taxifolia</i>	219	<i>Populus balsamifera</i> , var. <i>candicans</i>	173
<i>Pinus tetragona</i>	201	<i>Populus balsamifera</i> , var. <i>genuina</i>	173
<i>Pinus Torreyana</i>	192	<i>Populus Canadensis</i> (<i>Populus balsamifera</i> , var. <i>candicans</i>)	173
<i>Pinus tuberculata</i>	190	<i>Populus Canadensis</i> (<i>Populus monilifera</i>)	175
<i>Pinus tuberculata</i> (<i>Pinus insignis</i>)	196	<i>Populus Canadensis</i> , var. <i>angustifolia</i>	174
<i>Pinus variabilis</i>	200	<i>Populus candicans</i>	173
<i>Pinus venusta</i>	213	<i>Populus cordifolia</i>	172
<i>Pinus Virginiana</i>	198	<i>! Populus deltoide</i>	174
<i>Pinus Virginiana</i> , var. <i>echinata</i>	200	<i>Populus Fremontii</i>	175
<i>Piscidia Carthagenensis</i>	57	<i>Populus Fremontii</i> , var. <i>Wializeni</i>	175
<i>Piscidia Erythrina</i>	57	<i>Populus glandulosa</i>	175
<i>Pisonia aculeata</i>	117	<i>Populus grandidentata</i>	173
<i>Pisonia obtusata</i>	117	<i>Populus grandidentata</i> , var. <i>pendula</i>	173
<i>Pistacia Mexicana</i>	54	<i>Populus heterophylla</i>	173
<i>Pitch Pine</i>	198	<i>Populus heterophylla</i> (<i>Populus balsamifera</i> , var. <i>candicans</i>)	173
<i>Pithecolobium forbesi</i>	61	<i>Populus heterophylla</i> , var. <i>argentea</i>	172
<i>Pithecolobium Guadalupeensis</i>	64	<i>Populus laevigata</i> (<i>Populus monilifera</i> , <i>Aiton</i> , etc.)	175
<i>Pithecolobium microphyllum</i>	64	<i>Populus laevigata</i> (<i>Populus monilifera</i> , <i>Hort.</i>)	175
<i>Pithecolobium Unguis-cati</i>	64	<i>Populus latifolia</i>	173
<i>Planera aquatica</i>	124	<i>Populus Lindleyana</i>	175
<i>Planera Gmelini</i>	124	<i>Populus macrophylla</i> (<i>Populus balsamifera</i> , var. <i>candicans</i>)	173
<i>Planera Richardi</i>	124	<i>Populus macrophylla</i> (<i>Populus monilifera</i>)	175
<i>Planera ulmifolia</i>	124	<i>Populus Marylandica</i>	175
PLATANACEÆ	129, 130	<i>Populus monilifera</i>	174
<i>Platanus Californica</i>	129	<i>Populus monilifera</i> (<i>Populus Fremontii</i>)	175
<i>Platanus hybridæ</i>	129	<i>Populus monilifera</i> (<i>Populus Fremontii</i> , var. <i>Wializeni</i>)	175

	Page.		Page.
<i>Populus neglecta</i>	175	<i>Ptelea mollis</i>	81
<i>Populus Ontariensis</i>	173	<i>Ptelea trifoliata</i>	81
<i>Populus Tasamahaca</i>	173	<i>Ptelea trifoliata, var. mollis</i>	81
<i>Populus tremuliformis</i>	171	<i>Ptelea viticifolia</i>	81
<i>Populus tremuloides</i>	171	Punk Oak	152
<i>Populus tremula</i>	171	Purple Haw	40
<i>Populus trichocarpa</i>	174	Pyrus Americana	73
<i>Populus trichocarpa, var. cuspidata</i>	174	<i>Pyrus Americana (Pyrus sambucifolia)</i>	74
<i>Populus viminalis</i>	173	<i>Pyrus Americana, var. microcarpa</i>	74
<i>Populus Virginiana</i>	175	<i>Pyrus angustifolia</i>	72
<i>Porcelia triloba</i>	23	<i>Pyrus aucuparia (Pyrus Americana)</i>	73
Porkwood	117	<i>Pyrus aucuparia (Pyrus sambucifolia)</i>	74
Portiera angustifolia	29	<i>Pyrus Hartwegiana</i>	84
Port Orford Cedar	170	<i>Pyrus Botryospermum</i>	84
Possum Oak	132	<i>Pyrus coronaria</i>	72
Post Cedar	176	<i>Pyrus coronaria (Pyrus angustifolia)</i>	72
Post Oak	130	<i>Pyrus coronaria, var. angustifolia</i>	72
Post Oak, Swamp	140	<i>Pyrus diversifolia</i>	73
Prickly Ash (<i>Xanthoxylum Americanum</i>)	20	<i>Pyrus fusca</i>	73
Prickly Ash (<i>Xanthoxylum Clava-Herculis</i>)	30	<i>Pyrus glandulosa</i>	77
Prince's Pine	201	<i>Pyrus microcarpa</i>	74
<i>Prinos deciduus</i>	37	<i>Pyrus ovalis</i>	65
<i>Pritchardia filamentosa</i>	217	<i>Pyrus rivularis</i>	73
Privet	112	<i>Pyrus sambucifolia</i>	74
<i>Prosopis Emoryi</i>	62	<i>Pyrus subcordata</i>	73
<i>Prosopis glandulosa</i>	61	<i>Pyrus Wangenheimiana</i>	84
<i>Prosopis juliflora</i>	61		
<i>Prosopis odorata (Prosopis juliflora)</i>	61	Quaking Asp.	171
<i>Prosopis odorata (Prosopis pubescens)</i>	62	<i>Quassia dioica</i>	82
<i>Prosopis pubescens</i>	62	<i>Quassia Simaruba</i>	82
<i>Prunus Americana</i>	65	Quercitron Oak	149
<i>Prunus Americana, var. mollis</i>	65	<i>Quercus acutidens</i>	155
<i>Prunus angustifolia</i>	66	<i>Quercus acutiglans</i>	146
<i>Prunus borealis</i>	66	<i>Quercus agrifolia</i>	146
<i>Prunus Canadensis</i>	68	<i>Quercus agrifolia, var. frutescens</i>	147
<i>Prunus Capolin</i>	68	<i>Quercus alba</i>	137
<i>Prunus Capuli</i>	68	<i>Quercus alba minor</i>	139
<i>Prunus Carolina</i>	60	<i>Quercus alba palustris</i>	141
<i>Prunus Caroliniana</i>	69	<i>Quercus alba, var. ? Gunnisonii</i>	139
<i>Prunus cartilaginea</i>	68	<i>Quercus alba, var. microcarpa</i>	137
<i>Prunus Ohicosa</i>	66	<i>Quercus alba, var. pinnatifida</i>	137
<i>Prunus coccinea</i>	65	<i>Quercus alba, var. pinnatifido-sinuata</i>	137
<i>Prunus demissa</i>	69	<i>Quercus alba, var. repanda</i>	137
<i>Prunus emarginata</i>	67	<i>Quercus alba, var. sinuata</i>	137
<i>Prunus emarginata, var. mollis</i>	67	<i>Quercus ambigua</i>	147
<i>Prunus hiemalis (Prunus Americana)</i>	65	<i>Quercus annulata</i>	145
<i>Prunus hiemalis (Prunus Americana, var. mollis)</i>	65	<i>Quercus aquatica</i>	152
<i>Prunus ilicifolia</i>	70	<i>! Quercus aquatica (Quercus nigra)</i>	150
<i>Prunus inistitia</i>	66	<i>Quercus aquatica, var. attenuata</i>	152
<i>! Prunus lanceolata</i>	66	<i>Quercus aquatica, var. cuneata</i>	152
<i>Prunus Lusitanica</i>	69	<i>Quercus aquatica, var. elongata</i>	152
<i>Prunus Mississippi</i>	65	<i>Quercus aquatica, var. heterophylla</i>	153
<i>Prunus mollis (Prunus Americana, var. mollis)</i>	65	<i>Quercus aquatica, var. hybrida</i>	152
<i>Prunus mollis (Prunus emarginata, var. mollis)</i>	67	<i>Quercus aquatica, var. indivisa</i>	152
<i>Prunus nigra</i>	65	<i>Quercus aquatica, var. laurifolia</i>	152
<i>Prunus Pennsylvanica</i>	60	<i>Quercus aquatica, var. myrtifolia</i>	155
<i>! Prunus persicifolia</i>	66	<i>Quercus Banisteri</i>	155
<i>Prunus pumila</i>	67	<i>Quercus berberidifolia</i>	155
<i>Prunus sempervirens</i>	69	<i>Quercus bicolor</i>	141
<i>Prunus serotina</i>	68	<i>Quercus bicolor, var. Michauxii</i>	141
<i>Prunus serratifolia</i>	69	<i>Quercus bicolor, var. mollis</i>	141
<i>Prunus sphaerocarpa</i>	70	<i>! Quercus bicolor, var. platanioides</i>	141
<i>Prunus spinosa</i>	65	<i>Quercus Breweri</i>	155
<i>Prunus umbellata</i>	67	<i>Quercus Californica</i>	149
<i>Prunus Virginiana</i>	68	<i>Quercus Castanea (Quercus prinoides)</i>	142
<i>Prunus Virginiana (Prunus serotina)</i>	68	<i>Quercus Castanea (Quercus Prinus)</i>	142
<i>Prunus Virginiana, var. demissa</i>	69	<i>Quercus Catesbaei</i>	151
<i>Pseudocedra odorata</i>	55	<i>Quercus Chinquapin</i>	143
<i>Pseudopetalon glandulosum</i>	30	<i>Quercus chrysolpis</i>	146
<i>Pseudopetalon tricarpum</i>	30	<i>Quercus chrysopsis, var. racemifolia</i>	146
<i>Pseudotsuga Douglasii</i>	209	<i>Quercus cinerea</i>	153
<i>Pseudotsuga Douglasii, var. macrocarpa</i>	210	<i>Quercus cinerea, var. pumila</i>	155
<i>Pseudotsuga magnifica</i>	214	<i>Quercus cinerea, var. sericea</i>	155
<i>Pseudotsuga nobilis</i>	214	<i>Quercus coccinea</i>	148
<i>Paidium Gualva</i>	80	<i>Quercus coccinea, var. ambigua</i>	147

	Page.		Page.
<i>Quercus coccinea</i> , var. <i>microcarpa</i>	148	<i>Quercus oblongifolia</i> , var. <i>breviloba</i>	143
<i>Quercus coccinea</i> , var. <i>rubra</i>	147	<i>Quercus obtusa</i>	152
<i>Quercus coccinea</i> , var. <i>tinctoria</i>	149	<i>Quercus obtusifolia</i> , var. <i>? breviloba</i>	145
<i>Quercus confertifolia</i>	154	<i>Quercus obtusiloba</i>	138
<i>Quercus crassipetala</i>	146	<i>Quercus obtusiloba</i> , var. <i>depressa</i>	140
<i>Quercus cuneata</i>	150	<i>Quercus obtusiloba</i> , var. <i>parvifolia</i>	139
<i>Quercus decipiens</i>	144	<i>Quercus Cretediana</i>	138
<i>Quercus densiflora</i>	154	<i>Quercus oleoides</i>	145
<i>Quercus discolor</i>	149	<i>Quercus oliviformis</i>	140
<i>Quercus discolor</i> , var. <i>triloba</i>	151	<i>Quercus oxyadenia</i>	146
<i>Quercus Douglassii</i>	143	<i>Quercus palustris</i>	151
<i>Quercus Douglasii</i> , var. <i>Gambelii</i>	139	<i>Quercus palustris</i> (<i>Quercus rubra</i> , var. <i>Texana</i>)	148
<i>Quercus Douglasii</i> , var. <i>? Neesii</i>	138	<i>Quercus parvifolia</i>	130
<i>? Quercus Drummondii</i>	139	<i>Quercus Phellos</i>	154
<i>Quercus dumosa</i>	155	<i>Quercus Phellos angustifolia</i>	154
<i>Quercus dumosa</i> , var. <i>bullata</i>	155	<i>Quercus Phellos</i> × <i>coccinea</i>	158
<i>Quercus Durandii</i>	145	<i>Quercus Phellos latifolia</i>	154
<i>Quercus echinacea</i>	155	<i>Quercus Phellos pumila</i>	155
<i>Quercus echinoides</i>	155	<i>Quercus Phellos</i> × <i>tinctoria</i>	158
<i>Quercus elongata</i>	150	<i>Quercus Phellos</i> , var. (<i>Quercus heterophylla</i>)	153
<i>Quercus Emoryi</i>	146	<i>Quercus Phellos</i> , var. <i>arenaria</i>	155
<i>? Quercus Emoryi</i> (<i>Quercus undulata</i> , var. <i>Gambelii</i>)	139	<i>Quercus Phellos</i> , var. <i>cinerea</i>	153
<i>Quercus falcata</i>	150	<i>Quercus Phellos</i> , var. <i>humilis</i>	154
<i>Quercus falcata</i> , var. <i>Ludoviciana</i>	151	<i>Quercus Phellos</i> , var. <i>imbricaria</i>	154
<i>Quercus falcata</i> , var. <i>pagodafolia</i>	151	<i>Quercus Phellos</i> , var. <i>laurifolia</i>	152
<i>Quercus falcata</i> , var. <i>triloba</i>	151	<i>Quercus Phellos</i> , var. <i>sempervirens</i>	145
<i>Quercus fulvescens</i>	146	<i>Quercus Phellos</i> , var. <i>sericea</i>	155
<i>Quercus Gambelii</i>	139	<i>Quercus Phellos</i> , var. <i>viridis</i>	154
<i>Quercus Garryana</i>	138	<i>Quercus prinoides</i>	142
<i>Quercus Georgiana</i>	155	<i>Quercus Prinus</i>	142
<i>Quercus grisea</i>	144	<i>Quercus Prinus</i> β. (<i>Quercus cinerea</i>)	153
<i>Quercus hastata</i>	146	<i>Quercus Prinus</i> (<i>Quercus Michauxii</i>)	141
<i>Quercus hemisphaerica</i>	152	<i>Quercus Prinus Olinquapin</i>	143
<i>Quercus hemisphaerica</i> , var. <i>nana</i>	152	<i>Quercus Prinus humilis</i>	142
<i>Quercus heterophylla</i>	153	<i>Quercus Prinus palustris</i>	141
<i>Quercus Hindsii</i>	138	<i>? Quercus Prinus platanoidea</i>	141
<i>Quercus humilis</i>	153	<i>Quercus Prinus pumila</i>	142
<i>Quercus hypoleuca</i>	154	<i>Quercus Prinus tomentosa</i>	141
<i>Quercus ilicifolia</i>	155	<i>Quercus Prinus</i> , var. <i>acuminata</i>	142
<i>Quercus imbricaria</i>	154	<i>Quercus Prinus</i> , var. <i>bicolor</i>	141
<i>Quercus Jacobi</i>	138	<i>Quercus Prinus</i> , var. <i>discolor</i>	141
<i>Quercus Kelloggii</i>	149	<i>Quercus Prinus</i> , var. <i>lata</i>	142
<i>? Quercus laevis</i>	151	<i>Quercus Prinus</i> , var. <i>Michauxii</i>	141
<i>Quercus laurifolia</i>	152	<i>Quercus Prinus</i> , var. <i>monticola</i>	142
<i>Quercus laurifolia hybrida</i>	152	<i>Quercus Prinus</i> , var. <i>oblongata</i>	143
<i>Quercus laurifolia</i> , var. <i>acuta</i>	152	<i>Quercus Prinus</i> , var. <i>prinoides</i>	143
<i>Quercus laurifolia</i> , var. <i>obtus</i>	152	<i>Quercus pumila</i>	155
<i>Quercus lobata</i>	138	<i>Quercus pungens</i>	144
<i>Quercus lobata</i> , var. <i>fruticosa</i>	155	<i>Quercus Ransomii</i>	138
<i>Quercus lobulata</i>	139	<i>Quercus reticulata</i>	144
<i>Quercus longiglанда</i>	138	<i>? Quercus reticulata</i> , var. <i>Greggii</i>	144
<i>Quercus lyrata</i>	140	<i>Quercus retusa</i>	145
<i>Quercus macrocarpa</i>	140	<i>Quercus rubra</i>	147
<i>Quercus macrocarpa</i> , var. <i>abbreviata</i>	140	<i>Quercus rubra</i> β. (<i>Quercus coccinea</i>)	148
<i>Quercus macrocarpa</i> , var. <i>minor</i>	140	<i>Quercus rubra</i> (<i>Quercus Kelloggii</i>)	149
<i>Quercus macrocarpa</i> , var. <i>oleaformis</i>	140	<i>Quercus rubra</i> (<i>Quercus tinctoria</i>)	149
<i>Quercus Marylandica</i>	150	<i>Quercus rubra maxima</i>	147
<i>Quercus Michauxii</i>	141	<i>Quercus rubra montana</i>	150
<i>Quercus montana</i>	142	<i>Quercus rubra ramosissima</i>	151
<i>Quercus Morehus</i>	147	<i>Quercus rubra</i> , var. <i>dissecta</i>	151
<i>Quercus Muhlenbergii</i>	143	<i>Quercus rubra</i> var. <i>latifolia</i>	147
<i>Quercus myrtifolia</i>	155	<i>Quercus rubra</i> , var. <i>montana</i>	147
<i>Quercus nana</i>	152	<i>Quercus rubra</i> , var. <i>runcinata</i>	147
<i>Quercus Neesii</i>	138	<i>Quercus rubra</i> , var. <i>Texana</i>	148
<i>Quercus nigra</i>	150	<i>Quercus San-Sabeana</i>	145
<i>Quercus nigra</i> (<i>Quercus aquatica</i>)	152	<i>Quercus sempervirens</i>	145
<i>Quercus nigra</i> (<i>Quercus tinctoria</i>)	149	<i>Quercus sericea</i>	155
<i>Quercus nigra aquatica</i>	152	<i>? Quercus Shumardii</i>	149
<i>Quercus nigra digitata</i>	150	<i>? Quercus sinuata</i>	137
<i>Quercus nigra integrifolia</i>	150	<i>Quercus Sonomensis</i>	140
<i>Quercus nigra trifida</i>	152	<i>Quercus epicata</i>	144
<i>Quercus nigra</i> , var. (<i>Quercus heterophylla</i>)	153	<i>Quercus stellata</i>	139
<i>Quercus nigra</i> , var. <i>latifolia</i>	150	<i>Quercus stellata</i> , var. <i>depressa</i>	140
<i>Quercus oblongifolia</i>	143	<i>Quercus stellata</i> , var. <i>Floridana</i>	139
<i>? Quercus oblongifolia</i>	146	<i>Quercus stellata</i> , var. <i>Utahensis</i>	139

	Page.		Page.
<i>Quercus Texana</i>	148	<i>Rhus Canadense</i>	52
<i>Quercus tinctoria</i>	149	<i>Rhus copallina</i>	53
<i>Quercus tinctoria</i> , var. <i>angulosa</i>	149	<i>Rhus copallina</i> , var. <i>angustialata</i>	53
<i>Quercus tinctoria</i> , var. <i>Californica</i>	149	<i>Rhus copallina</i> , var. <i>angustifolia</i>	53
<i>Quercus tinctoria</i> , var. <i>sinuosa</i>	149	<i>Rhus copallina</i> , var. <i>integrifolia</i>	53
<i>Quercus triloba</i>	150	<i>Rhus copallina</i> , var. <i>lanceolata</i>	53
<i>Quercus uliginosa</i>	152	<i>Rhus copallina</i> , var. <i>latialata</i>	53
<i>Quercus undulata</i>	155	<i>Rhus copallina</i> , var. <i>latifolia</i>	53
<i>Quercus undulata</i> (<i>Quercus Durandii</i>).....	145	<i>Rhus copallina</i> , var. <i>leucantha</i>	53
<i>Quercus undulata</i> , var. <i>Gambelii</i>	139	<i>Rhus copallina</i> , var. <i>serrata</i>	53
<i>Quercus undulata</i> , var. <i>grisea</i>	144	<i>Rhus cotinoides</i>	52
<i>Quercus undulata</i> , var. <i>oblongata</i>	143	<i>Rhus cotinus</i>	52
<i>Quercus undulata</i> , var. <i>pungens</i>	144	<i>Rhus glabra</i>	53
<i>Quercus undulata</i> , var. <i>Wrightii</i>	144	<i>Rhus hypselodendron</i>	52
<i>Quercus vaccinifolia</i>	146	<i>Rhus leucantha</i>	53
<i>Quercus velutina</i>	149	<i>Rhus Metopium</i>	54
<i>Quercus villosa</i>	139	<i>Rhus Toxicodendron</i>	54
<i>Quercus virens</i>	145	<i>Rhus typhina</i>	52
<i>Quercus Virginiana</i>	145	<i>Rhus typhina</i> , var. <i>laciniata</i>	52
<i>Quercus Wislizeni</i>	147	<i>Rhus typhina</i> , var. <i>viridiflora</i>	52
R.		<i>Rhus venenata</i>	54
<i>Randia chusafolia</i>	95	<i>Rhus vernix</i>	54
<i>Rapanea Guyanensis</i>	99	<i>Rhus viridiflora</i>	52
Rattle-box.....	106	River Birch.....	161
Red Ash.....	109	River Cottonwood.....	172
Red Bay.....	118	<i>Robinia fragilis</i>	55
Red Birch.....	161	<i>Robinia glutinosa</i>	56
Red Cedar (<i>Juniperus Virginiana</i>).....	183	<i>Robinia Neo-Mexicana</i>	56
Red Cedar (<i>Thuja gigantea</i>).....	177	<i>Robinia Pseudacacia</i>	55
Red Cherry, Wild.....	66	<i>Robinia viscosa</i>	56
Red Cypress.....	184	Rock Chestnut Oak.....	142
Red Elm.....	122	Rock Elm.....	123
Red Fir (<i>Abies magnifica</i>).....	214	Rock Maple.....	48
Red Fir (<i>Abies nobilis</i>).....	214	ROSACEAE.....	64-85
Red Fir (<i>Pseudotsuga Douglasii</i>).....	209	Rose Bay.....	90
Red Gum.....	86	Rosemary Pine.....	197
Red Haw (<i>Crataegus coccinea</i>).....	78	Royal Palm.....	218
Red Haw (<i>Crataegus flava</i> , var. <i>pubescens</i>).....	88	RUBIACEAE.....	95, 96
Red Ironwood.....	89	Rum Cherry.....	68
Red Maple.....	50	RUTACEAE.....	20-32
Red Mulberry.....	128	S.	
Red Oak (<i>Quercus falcata</i>).....	151	Sabal Palmetto.....	217
Red Oak (<i>Quercus rubra</i>).....	148	Saffron Plum.....	103
Red Oak (<i>Quercus rubra</i> , var. <i>Texana</i>).....	148	Saguaro.....	90
Red Pine.....	102	SALICACEAE.....	165-175
Red Stopper.....	89	<i>Salix</i>	170
Redbud (<i>Cercis Canadensis</i>).....	61	<i>Salix ambigua</i>	165
Redbud (<i>Cercis reniformis</i>).....	61	<i>Salix amygdaloides</i>	166
Redwood.....	185	<i>Salix argophylla</i>	168
Reynoldsia latifolia.....	39	<i>Salix arguta</i>	167
<i>Rhamnidium revolutum</i>	39	<i>Salix arguta</i> , var. <i>lasiantha</i>	167
RHAMNACEAE.....	39-42	<i>Salix Bigelovii</i>	170
<i>Rhamnus alnifolia</i>	41	<i>Salix Bigelovii</i> , var. <i>fuscol.</i>	170
<i>Rhamnus Californica</i>	40	<i>Salix brachycarpa</i>	168
<i>Rhamnus Californica</i> , var. <i>tomentella</i>	41	<i>Salix brachystachys</i>	170
<i>Rhamnus Caroliniana</i>	40	<i>Salix brachystachys</i> , var. <i>Scouleriana</i>	170
<i>Rhamnus ellipticus</i>	41	<i>Salix Caroliniana</i>	165
<i>Rhamnus ferrea</i>	39	<i>Salix chlorophylla</i> , var. <i>pellita</i>	171
<i>Rhamnus laevigatus</i>	39	<i>Salix cordata</i>	170
<i>Rhamnus laurifolius</i>	40	<i>Salix cordata</i> , var. <i>falcata</i>	165
<i>Rhamnus oleifolius</i>	40	<i>Salix cordata</i> , var. <i>vestita</i>	170
<i>Rhamnus Purshiana</i>	41	<i>Salix crassa</i>	169
<i>Rhamnus tomentellus</i>	41	<i>Salix cuneata</i>	171
<i>Rhizophora Americana</i>	86	<i>Salix discolor</i>	169
<i>Rhizophora Mangle</i>	86	<i>Salix discolor</i> , var. <i>eriocephala</i>	169
<i>Rhizophora racemosa</i>	86	<i>Salix discolor</i> , var. <i>prinoides</i>	169
RHIZOPHORACEAE.....	86, 87	<i>Salix eriocephala</i>	169
<i>Rhododendron maximum</i>	90	<i>Salix exigua</i>	168
<i>Rhododendron maximum</i> , var. <i>album</i>	90	<i>Salix falcata</i>	165
<i>Rhododendron maximum</i> , var. <i>purpureum</i>	90	<i>Salix Fendleriana</i>	167
<i>Rhododendron maximum</i> , var. <i>roseum</i>	90	<i>Salix flavescens</i>	169
<i>Rhododendron procerum</i>	90	<i>Salix flavescens</i> (<i>Salix flavescens</i> , var. <i>Scouleriana</i>).....	170
<i>Rhododendron purpureum</i>	90	<i>Salix flavescens</i> , var. <i>Scouleriana</i>	170
<i>Rhododendron Purshii</i>	90	<i>Salix flavo-virens</i>	165
<i>Rhus arborecens</i>	33	<i>Salix fluviatilis</i>	168

	Page.		Page.
<i>Salix Hartwegi</i>	171	<i>Sapindus Drummondii</i>	44
<i>Salix Hindiana</i>	169	<i>Sapindus falcatus</i>	44
<i>Salix Hindiana</i> , var. <i>tenusifolia</i>	169	<i>Sapindus inaequalis</i>	44
<i>Salix Hoffmanniana</i>	167	<i>Sapindus marginatus</i>	44
<i>Salix Hookeriana</i>	170	<i>Sapindus Saponaria</i>	45
<i>Salix Houstoniana</i>	165	<i>Sapindus Saponaria</i> (<i>Sapindus marginatus</i>)	44
<i>Salix humilis</i> , var. <i>Hartwegi</i>	171	SAPOTACEAE	100-108
<i>Salix laevigata</i>	167	<i>Sarcophallus Carolinianus</i>	40
<i>Salix laevigata</i> , var. <i>angustifolia</i>	167	Sassafras	120
<i>Salix laevigata</i> , var. <i>congesta</i>	167	<i>Sassafras officinale</i>	119
<i>Salix lanceifolia</i>	167	Satinwood	31
<i>Salix lasiandra</i>	167	Savin (<i>Juniperus Virginiana</i>)	183
<i>Salix lasiandra</i> , var. <i>Fendleriana</i>	167	Savin (<i>Torreya taxifolia</i>)	186
<i>Salix lasiandra</i> , var. <i>lanceifolia</i>	167	Scarlet Haw (<i>Crataegus coccinea</i>)	78
<i>Salix lasiolepis</i>	170	Scarlet Haw (<i>Crataegus subvillosa</i>)	78
<i>Salix lasiolepis</i> , var. <i>angustifolia</i>	171	Scarlet Oak	143
<i>Salix lasiolepis</i> , var. <i>Bigelovii</i>	170	<i>Schaefferia buxifolia</i>	30
<i>Salix lasiolepis</i> , var. <i>fallax</i>	170	<i>Schaefferia completa</i>	30
<i>Salix lasiolepis</i> , var. <i>latifolia</i>	171	<i>Schaefferia frutescens</i>	30
<i>Salix ligustrina</i>	165	<i>Schaefferia lateriflora</i>	120
<i>Salix longifolia</i>	168	<i>Schoubaea commutata</i>	87
<i>Salix longifolia</i> , var. <i>angustissima</i>	168	<i>Schubertia disticha</i>	183
<i>Salix longifolia</i> , var. <i>argyrophylla</i>	168	<i>Schubertia sempervirens</i>	185
<i>Salix longifolia</i> , var. <i>argyrophylla angustissima</i>	168	Screw Bean	62
<i>Salix longifolia</i> , var. <i>argyrophylla opaca</i>	168	Screw-pod Mesquit	62
<i>Salix longifolia</i> , var. <i>exigua</i>	168	Scrub Oak (<i>Quercus Catesbaei</i>)	151
<i>Salix longifolia</i> , var. <i>opaca</i>	168	Scrub Oak (<i>Quercus undulata</i> , var. <i>Gambellii</i>)	139
<i>Salix longifolia</i> , var. <i>pedicellata</i>	168	Scrub Pine (<i>Pinus Banksiana</i>)	201
<i>Salix longipes</i>	166	Scrub Pine (<i>Pinus clausa</i>)	190
<i>Salix longipes</i> , var. <i>pubescens</i>	166	Scrub Pine (<i>Pinus contorta</i>)	194
<i>Salix lucida</i> , var. <i>angustifolia</i> , forma <i>lasiandra</i>	167	Scrub Pine (<i>Pinus inops</i>)	199
<i>Salix lucida</i> , var. <i>macrophylla</i>	167	<i>Scutia ferrea</i> (<i>Condalia ferrea</i>)	39
<i>Salix marginata</i>	166	<i>Scutia ferrea</i> (<i>Reynoldsia latifolia</i>)	39
<i>Salix melanopsis</i>	166	Sea Ash	30
<i>Salix nigra</i>	165	Sea Grape	118
<i>Salix nigra</i> , var. <i>amygdaloides</i>	166	Seaside Alder	162
<i>Salix nigra</i> , var. <i>angustifolia</i>	166	<i>Sebastiania lucida</i>	121
<i>Salix nigra</i> , var. <i>falcata</i>	165	<i>Sebestena scabra</i>	113
<i>Salix nigra</i> , var. <i>latifolia</i>	166	<i>Sequoia gigantea</i>	184
<i>Salix nigra</i> , var. <i>longifolia</i>	166	<i>Sequoia gigantea</i> (<i>Sequoia sempervirens</i>)	185
<i>Salix nigra</i> , var. <i>longipes</i>	166	<i>Sequoia Rafinesquii</i>	206
<i>Salix nigra</i> , var. <i>longipes</i> , subvar. <i>gongylocarpa</i>	166	<i>Sequoia religiosa</i>	185
<i>Salix nigra</i> , var. <i>longipes</i> , subvar. <i>venulosa</i>	166	<i>Sequoia sempervirens</i>	184
<i>Salix nigra</i> , var. <i>marginata</i>	166	<i>Sequoia Wellingtonia</i>	184
<i>Salix nigra</i> , var. <i>Wardii</i>	166	Service Tree	84
<i>Salix occidentalis</i>	166	Seven-year Apple	95
<i>Salix pentandra</i>	165	Shad Bush	84
<i>Salix pentandra</i> , var. <i>caudata</i>	167	Shag-bark Hickory	133
<i>Salix prinoides</i>	169	She Balsam	210
<i>Salix Purshiana</i>	165	Sheepberry	94
<i>Salix rubra</i>	168	Shell-bark, Big	133
<i>Salix Scouleriana</i>	170	Shell-bark, Bottom	133
<i>Salix sensitiva</i>	169	Shell-bark Hickory	133
<i>Salix sessilifolia</i>	168	Shingle Oak	154
<i>Salix sessilifolia</i> , var. <i>Hindiana</i>	169	Shittimwood (<i>Bumelia lanuginosa</i>)	102
<i>Salix sessilifolia</i> , var. <i>villosa</i>	168	Shittimwood (<i>Rhamnus Purshiana</i>)	41
<i>Salix Sitchenensis</i>	171	Short-leaved Pine	200
<i>Salix Sitchenensis</i> , var. <i>angustifolia</i>	171	Shrubby Trefoll	31
<i>Salix speciosa</i>	167	<i>Sideroxylon Carolinense</i>	101
<i>Salix subvillosa</i>	166	<i>Sideroxylon chrysophyllodes</i>	101
<i>Salix Wrightii</i>	166	<i>Sideroxylon cuneatum</i>	108
<i>Samara floribunda</i>	99	<i>Sideroxylon decandrum</i>	103
<i>Samara pentandra</i>	99	<i>Sideroxylon lanuginosum</i>	102
<i>Sambucus Californica</i>	93	<i>Sideroxylon laeve</i>	108
<i>Sambucus glauca</i>	93	<i>Sideroxylon lycioides</i>	103
<i>Sambucus glauca</i> (<i>Sambucus Mexicana</i>)	93	<i>Sideroxylon Mastichodendron</i>	101
<i>Sambucus Mexicana</i>	93	<i>Sideroxylon pallidum</i>	101
<i>Sambucus Mexicana</i> (<i>Sambucus glauca</i>)	93	<i>Sideroxylon reclinatum</i>	103
<i>Sambucus velutina</i>	93	<i>Sideroxylon salicifolium</i>	101
Sand Jack	153	<i>Sideroxylon sericeum</i>	101
Sand Pine	199	<i>Sideroxylon tenax</i> (<i>Bumelia lanuginosa</i>)	102
Sand-bar Willow	168	<i>Sideroxylon tenax</i> (<i>Bumelia tenax</i>)	101
SAPINDACEAE	42-51	<i>Silquastrum cordatum</i>	61
<i>Sapindus acuminata</i>	44	Silk-top Palmetto	217
		Silky Willow	171

241

16 FOR

	Page.		Page.
TERNSTROMIACEÆ	25, 26	<i>Tsuga Caroliniana</i>	207
<i>Tetranthera Californica</i>	120	<i>Tsuga Douglasii</i>	209
Thatch, Brickley.....	218	<i>Tsuga Lindleyana</i>	209
Thatch, Brittle.....	218	<i>Tsuga Mertensiana</i>	207
The Joshua.....	219	<i>Tsuga Pattoniana</i>	208
Thorn, Black.....	79	<i>Tulipastrum Americanum</i>	20
Thorn, Cockspur.....	76	<i>Tulipastrum Americanum</i> , var. <i>subcordatum</i>	20
Thorn, Newcastle.....	76	<i>Tulipifera Liriodendron</i>	22
Thorn, Washington.....	81	Tulip Tree.....	22
Thorn, White.....	78	Tupelo.....	92
Three-thorned Acacia.....	59	Tupelo Gum.....	93
<i>Thrinax argentea</i>	218	Tupelo, Large.....	93
<i>Thrinax Garberi</i>	217	Tupelo, Sour.....	91
<i>Thrinax parviflora</i>	217	Turkey Oak.....	151
<i>Thuja Craigiana</i>	176		
<i>Thuja excelsa</i>	178	U.	
<i>Thuja gigantea</i>	177	<i>Ulmus alata</i>	124
<i>Thuja gigantea</i> (Libocedrus decurrens).....	176	<i>Ulmus alba</i>	123
<i>Thuja Lobbii</i>	177	<i>Ulmus Americana</i>	123
<i>Thuja Menziesii</i>	177	<i>Ulmus Americana</i> (Ulmus racemosa).....	123
<i>Thuja obtusa</i>	176	<i>Ulmus Americana</i> , var. <i>alata</i>	124
<i>Thuja occidentalis</i>	176	<i>Ulmus Americana</i> , var. <i>alba</i>	123
<i>Thuja occidentalis</i> , var. <i>placata</i>	177	<i>Ulmus Americana</i> , var. <i>aspera</i>	123
<i>Thuja odorata</i>	176	<i>Ulmus Americana</i> , var. <i>Bartramii</i>	123
<i>Thuja plicata</i>	177	<i>Ulmus Americana</i> , var. <i>pendula</i>	123
<i>Thuja Sibirica</i>	176	<i>Ulmus Americana</i> , var. <i>rubra</i>	122
<i>Thuja sphaeroidalis</i>	177	<i>Ulmus Americana</i> , var. <i>scabra</i>	123
<i>Thuja sphaeroides</i>	177	<i>Ulmus aquatica</i>	124
<i>Thuja Wareana</i>	176	<i>Ulmus crassifolia</i>	122
<i>Thuyopsis borealis</i>	178	? <i>Ulmus crispus</i>	122
<i>Thuyopsis cupressoides</i>	178	<i>Ulmus Floridana</i>	123
<i>Thuyopsis Tchugatskoy</i>	178	<i>Ulmus fulva</i>	122
<i>Thuyopsis Tchugatskoyae</i>	178	<i>Ulmus mollifolia</i>	123
<i>Thylax fraxineum</i>	29	? <i>Ulmus nemoralis</i>	124
Tide-land Spruce.....	206	<i>Ulmus opaca</i>	122
<i>Tilia alba</i>	27	<i>Ulmus pendula</i>	123
<i>Tilia Americana</i>	26	<i>Ulmus pubescens</i>	122
<i>Tilia Americana</i> (Tilia Americana, var. <i>pubescens</i>).....	27	<i>Ulmus pumila</i>	124
<i>Tilia Americana</i> , var. <i>heterophylla</i>	27	<i>Ulmus racemosa</i>	123
<i>Tilia Americana</i> , var. <i>pubescens</i>	27	<i>Ulmus rubra</i>	122
<i>Tilia Americana</i> , var. <i>Walleri</i>	27	<i>Umbellularia Californica</i>	120
<i>Tilia Canadensis</i>	26	<i>Umbrella Tree</i>	21
<i>Tilia Caroliniana</i>	27	<i>Ungnadia heptaphylla</i>	44
<i>Tilia glabra</i>	26	<i>Ungnadia heterophylla</i>	44
<i>Tilia grata</i>	27	<i>Ungnadia speciosa</i>	44
<i>Tilia heterophylla</i>	27	Upland Willow Oak.....	153
<i>Tilia heterophylla</i> , var. <i>alba</i>	27	<i>Urostigma pedunculatum</i>	127
<i>Tilia latifolia</i>	26	URTICACEÆ	122-128
<i>Tilia laxiflora</i> (Tilia Americana, var. <i>pubescens</i>).....	27	<i>Urtica triloba</i>	23
<i>Tilia laxiflora</i> (Tilia heterophylla).....	27		
<i>Tilia neglecta</i>	26	V.	
<i>Tilia nigra</i>	26	<i>Vaccinium arboreum</i>	96
<i>Tilia pubescens</i>	27	<i>Vaccinium difusum</i>	96
<i>Tilia pubescens</i> , var. <i>leptophylla</i>	27	<i>Vaccinium mucronatum</i>	96
? <i>Tilia stenopetala</i>	27	Valparaiso Oak.....	146
<i>Tilia truncata</i>	27	<i>Vauquelinia corymbosa</i>	70
TILIACEÆ	26-28	<i>Vauquelinia Torreyi</i>	70
<i>Tili.</i>	88	VERBENACEÆ	116, 117
<i>Tollon</i>	84	<i>Viburnum Lentago</i>	94
Toothache Tree (Xanthoxylum Americanum).....	29	<i>Viburnum prunifolium</i>	94
Toothache Tree (Xanthoxylum Clava-Herculis).....	30	<i>Viburnum prunifolium</i> , var. <i>ferrugineum</i>	94
Torchwood.....	33	<i>Viburnum pyrifolium</i>	94
Tornilla.....	62	Vine Maple.....	47
<i>Torreya Californica</i>	186	<i>Virgilia lutea</i>	57
<i>Torreya Myristica</i>	186	<i>Virgilia secundiflora</i>	57
<i>Torreya taxifolia</i>	186		
<i>Toxicodendron arborescens</i>	33	W.	
<i>Toxylon MacLurea</i>	128	<i>Wafer Ash</i>	31
<i>Toyon</i>	84	<i>Wahoo</i> (Euonymus atropurpureus).....	38
Trefoll, Shrubby.....	31	<i>Wahoo</i> (Tilia heterophylla).....	28
<i>Trilopus dentata</i>	85	<i>Wahoo</i> (Ulmus alata).....	124
<i>Trilopus nigra</i>	85	<i>Wallia cinerea</i>	130
<i>Trilopus parvifolia</i>	85	<i>Wallia nigra</i>	131
<i>Trilopus rotundifolia</i>	85	Walnut.....	131
<i>Trilopus Virginiana</i>	85	Walnut, Black.....	131
<i>Tsuga Canadensis</i>	206	Walnut, White.....	136
		<i>Washingtonia filifera</i>	217
		<i>Washington Thorn</i>	31

243

	Page.		Page.
Water Ash	110	Willow (<i>Salix amygdaloides</i>)	168
Water Beech (<i>Carpinus Caroliniana</i>)	159	Willow (<i>Salix laevigata</i>)	167
Water Beech (<i>Platanus occidentalis</i>)	129	Willow (<i>Salix lasiandra</i>)	167
Water Elm	123	Willow (<i>Salix lasiolepis</i>)	171
Water Hickory	136	Willow, Black (<i>Salix flaveoens</i> , var. <i>Scouleriana</i>)	170
Water Locust	60	Willow, Black (<i>Salix nigra</i>)	166
Water Maple	50	Willow, Desert	116
Water Oak (<i>Quercus aquatica</i>)	153	Willow, Diamond	170
Water Oak (<i>Quercus palustris</i>)	152	Willow, Glaucons	169
Water White Oak	140	Willow Oak	154
Wax Myrtle	136	Willow Oak, Upland	153
Weeping Oak	138	Willow, Sand-bar	168
<i>Wellingtonia Californica</i>	184	Willow, Silky	171
<i>Wellingtonia gigantea</i>	184	Winged Elm	124
Western Catalpa	115	<i>Winters Canella</i>	24
West-Indian Birch	93	Witch Hazel	85
Weymouth Pine	187		
Whistlewood	46	Xanthoxylum Americanum	29
White Ash	107	Xanthoxylum aromaticum	30
White Basswood	28	Xanthoxylum Caribaeum	30
White Bay	20	Xanthoxylum Carolinianum	30
White Birch (<i>Betula alba</i> , var. <i>populifolia</i>)	159	Xanthoxylum Oatesbianum	30
White Birch (<i>Betula papyrifera</i>)	160	Xanthoxylum Clava-Herculis	30
White Buttonwood	87	Xanthoxylum Olava-Herculis (<i>Xanthoxylum Americanum</i>)	29
White Cedar (<i>Chamaecyparis Lawsoniana</i>)	179	Xanthoxylum Olava-Herculis (<i>Xanthoxylum Caribaeum</i>)	30
White Cedar (<i>Chamaecyparis sphaeroides</i>)	178	Xanthoxylum Olava-Herculis, var.	30
White Cedar (<i>Libocedrus decurrens</i>)	17	Xanthoxylum Clava-Herculis, var. <i>fruticosum</i>	30
White Cedar (<i>Thuja occidentalis</i>)	176	Xanthoxylum Floridanum	30
White Cottonwood	175	Xanthoxylum frazineum	29
White Cypress	184	Xanthoxylum fraziniifolium (<i>Xanthoxylum Americanum</i>)	29
White Elm (<i>Ulmus Americana</i>)	123	Xanthoxylum fraziniifolium (<i>Xanthoxylum Clava-Herculis</i>)	30
White Elm (<i>Ulmus racemosa</i>)	123	Xanthoxylum hirsutum	30
White Fir (<i>Abies concolor</i>)	213	Xanthoxylum lanceolatum	30
White Fir (<i>Abies grandis</i>)	212	Xanthoxylum macrophyllum	30
White Ironwood	45	Xanthoxylum mite	29
White Laurel	20	Xanthoxylum Pterota	31
White Mangrove	87	Xanthoxylum ramiflorum	29
White Maple	49	Xanthoxylum tricarpum (<i>Xanthoxylum Americanum</i>)	29
White Oak (<i>Quercus alba</i>)	137	Xanthoxylum tricarpum (<i>Xanthoxylum Clava-Herculis</i>)	30
White Oak (<i>Quercus Garryana</i>)	138	Ximenia Americana	34
White Oak (<i>Quercus grisea</i>)	144	Ximenia montana	34
White Oak (<i>Quercus lobata</i>)	138	Ximenia multiflora	34
White Oak (<i>Quercus oblongifolia</i>)	144		
White Oak, Mountain	143	Yaupon	36
White Oak, Swamp	141	Yellow Ash	57
White Oak, Water	140	Yellow-bark Oak	149
White Pine (<i>Pinus flexilis</i>)	188	Yellow Birch	161
White Pine (<i>Pinus glabra</i>)	201	Yellow Cypress	178
White Pine (<i>Pinus monticola</i>)	187	Yellow Fir	209
White Pine (<i>Pinus reflexa</i>)	189	Yellow Haw	83
White Pine (<i>Pinus Strobus</i>)	187	Yellow Locust	55
White Spruce (<i>Picea alba</i>)	204	Yellow Oak (<i>Quercus prinoides</i>)	143
White Spruce (<i>Picea Engelmanni</i>)	205	Yellow Oak (<i>Quercus tinctoria</i>)	149
White Spruce (<i>Picea pungens</i>)	205	Yellow Pine (<i>Pinus Arizona</i>)	192
White Stopper	89	Yellow Pine (<i>Pinus mitis</i>)	200
White Thorn	78	Yellow Pine (<i>Pinus palustris</i>)	202
White Walnut	130	Yellow Pine (<i>Pinus ponderosa</i>)	193
White-heart Hickory	184	Yellow Poplar	22
Whitewood (<i>Canella alba</i>)	24	Yellowwood (<i>Cladrastis tinctoria</i>)	57
Whitewood (<i>Drypetes crocea</i>)	121	Yellowwood (<i>Schaefferia frutescens</i>)	39
Whitewood (<i>Liriodendron Tulipifera</i>)	22	Yew (<i>Taxus brevifolia</i>)	185
Wild Black Cherry	68	Yew (<i>Taxus Florida</i>)	186
Wild Cherry (<i>Prunus Capuli</i>)	69	Yopon	36
Wild Cherry (<i>Prunus demissa</i>)	69	<i>Yucca angustifolia</i> , var. <i>elata</i>	219
Wild China	44	<i>Yucca angustifolia</i> , var. <i>radiosa</i>	219
Wild Cinnamon	24	<i>Yucca baccata</i>	219
Wild Dilly	103	<i>Yucca brevifolia</i>	218
Wild Fig	127	<i>Yucca canaliculata</i>	218
Wild Lime (<i>Xanthoxylum Pterota</i>)	31	<i>Yucca Draconis</i> , ? var. <i>arborescens</i>	218
Wild Lime (<i>Ximenia Americana</i>)	34	<i>Yucca elata</i>	219
Wild Orange (<i>Prunus Caroliniana</i>)	70	<i>Yucca filamentosa</i> ?	219
Wild Orange (<i>Xanthoxylum Clava-Herculis</i>)	30	<i>Yucca Treculiana</i>	218
Wild Peach	70		
Wild Plum	65	Zizyphus Dominicensis	41
Wild Red Cherry	66	Zizyphus emarginatus	39
Wild Tamarind	64	ZYGOPHYLLACEAE	28, 29

PART II.

THE WOODS OF THE UNITED STATES.

THE WOODS OF THE UNITED STATES.

A critical examination of the wood produced by the indigenous trees of North America, exclusive of Mexico, has been made in connection with the investigation of the forest wealth of the United States.

Mr. S. P. Sharples, special agent in charge of this department of the investigation, has had general direction of such experiments, and suggested the methods adopted for their execution.

The object of this examination has been to determine, first, the fuel value of the woods of the United States; second, the value as material for construction of the wood of the principal timber trees of the country. The results thus obtained are highly suggestive; they must not, however, be considered conclusive, but rather valuable as indicating what lines of research should be followed in a more thorough study of this subject.

The fuel value has been obtained by a determination of the specific gravity and the ash of the absolutely dry wood, supplemented by a determination of the actual chemical composition of the wood of some of the most important trees; the value of our woods for construction has been obtained by experiments made with the United States testing-machine at the Watertown arsenal. Each specimen as received was at once numbered, and this number, designated in the following tables as "Office number", was carefully repeated on every fragment cut from the original tree, and always refers to the same specimen. In a few cases in the early part of the work a sub-number was used to designate a specimen from another tree of the same species received from the same collector. In most cases the specimens were taken from the butt-cut of the tree, and unless it is otherwise mentioned in the remarks, were free from sap and knots; they may be regarded as representing the best wood that could be obtained from the tree.

The specimens used in the different series of experiments are deposited in the National Museum at Washington and in the museum of the Arboretum of Harvard College. It was found necessary, in order to secure proper material upon which to carry out the various experiments, to obtain a much larger amount of wood of the different species than was actually consumed in the experiments. This surplus material has been worked into 12,961 museum specimens, of convenient size, showing as far as possible the bark, sap-, and heart-wood of each species. These have been made into sixty sets, more or less complete, and distributed to the following educational institutions in the United States and Europe:

Institute of Technology, Boston, Massachusetts.
United States Military Academy, West Point, New York.
Academy of Natural Science, Philadelphia, Pennsylvania.
United States Naval Academy, Annapolis, Maryland.
Sheffield Scientific School, New Haven, Connecticut.
School of Mines, Columbia College, New York, New York.
National School of Forestry, Nancy, France.
Museum of Science and Art, Edinburgh, Scotland.
Agricultural Museum, Rome, Italy.
Brown University, Providence, Rhode Island.
Rensselaer Polytechnic Institute, Troy, New York.
Lawrence Scientific School, Cambridge, Massachusetts.
Iowa Agricultural College, Ames, Iowa.
Administration of National Forests, Lisbon, Portugal.
National Forest Administration, Paris, France.
McGill University, Montreal, Canada.
Royal Botanic Gardens, Sydney, New South Wales.
State Agricultural College, Lansing, Michigan.

Peabody Academy of Science, Salem, Massachusetts.
 Arkansas Industrial University, Fayetteville, Arkansas.
 Imperial Botanic Gardens, St. Petersburg, Russia.
 American Society of Civil Engineers, New York, New York.
 Portland Society of Natural History, Portland, Maine.
 New Jersey Agricultural College, New Brunswick, New Jersey.
 State Agricultural College, Burlington, Vermont.
 State Agricultural Collège, College Station, Maryland.
 Union College Engineering School, Schenectady, New York.
 Cornell University, Ithaca, New York.
 Hampton Agricultural and Normal Institute, Hampton, Virginia.
 Pennsylvania State College, State College, Pennsylvania.
 Ohio State University, Columbus, Ohio.
 Agricultural College of Missouri, Columbia, Missouri.
 University of Wisconsin, Madison, Wisconsin.
 State Agricultural and Mechanical College, Auburn, Alabama.
 University of Minnesota, Minneapolis, Minnesota.
 North Carolina Agricultural College, Chapel Hill, North Carolina.
 West Virginia University, Morgantown, West Virginia.
 State Agricultural College, Orono, Maine.
 Georgia Agricultural College, Athens, Georgia.
 Massachusetts Agricultural College, Amherst, Massachusetts.
 Tennessee Agricultural College, Knoxville, Tennessee.
 New Hampshire College of Agriculture, Hanover, New Hampshire.
 Illinois Industrial University, Champaign, Illinois.
 State Agricultural College, Corvallis, Oregon.
 State Agricultural College, Manhattan, Kansas.
 Agricultural College of Mississippi, Starkville, Mississippi.
 Kentucky Agricultural College, Lexington, Kentucky.
 Claflin University, Orangeville, South Carolina.
 Purdue University, Lafayette, Indiana.
 Botanic Garden, Königsberg, Germany.
 Engineer's office, Water-works, Boston, Massachusetts.
 Franklin Society, Providence, Rhode Island.
 Madison University, Hamilton, New York.
 Rochester University, Rochester, New York.
 Colby Academy, New London, New Hampshire.

SPECIFIC GRAVITY AND ASH.

The specific gravity and the ash of every tree of the United States have been determined (Table I) by Mr. Sharples, with the exception of the following: *Olusia flava*, once detected upon the keys of southern Florida, but not rediscovered; *Gordonia pubescens*, a rare and local species discovered in the last century upon the banks of the Altamaha river of Georgia and never rediscovered; *Pistacia Mexicana* and *Acacia Berlandieri*, economically unimportant species of the valley of the lower Rio Grande; *Cratægus berberifolia*, a little known species of the Red River valley; *Cupressus Macnabiana*, a rare and local species of California of little economic importance, and *Larix Lyallii*, a rare and local species of the northern Rocky mountains.

At least two determinations of specific gravity have been made for each species studied, and, in the case of woods of commercial importance, specimens were taken from many trees growing in widely different parts of the country, and under different conditions of soil and climate.

The specimens used for specific gravity determinations were made 100 millimeters long and about 35 millimeters square, and were dried at 100° centigrade until they ceased to lose weight. The specific gravity was then obtained by measurement with micrometer calipers and calculation from the weights of the blocks.

Two determinations of ash were made from each specimen studied by burning small, dried blocks in a muffle furnace at a low temperature.

An average of the specific gravity and of the ash of all the specimens taken from the same tree was made, and the average of these averages is given as the final result for the species; equal weight is thus given to each tree in the calculations without regard to the number of specimens representing it.

In the following table the trees of the United States are arranged in the order of the weight of the dry wood:

Catalogue number.	Species.	Relative specific gravity.	Catalogue number.	Species.	Relative specific gravity.	Catalogue number.	Species.	Relative specific gravity.
43	<i>Condalia ferrea</i>	1.8020	163	<i>Gnetta elliptica</i>	0.8337	402	<i>Larix occidentalis</i>	0.7407
44	<i>Condalia obovata</i>	1.1999	159	<i>Viburnum prunifolium</i>	0.8332	273	<i>Quercus coccinea</i>	0.7405
140	<i>Rhizophora Mangle</i>	1.1017	26	<i>Ptelea trifoliata</i>	0.8319	87	<i>Gleditsia monosperma</i>	0.7343
20	<i>Gnailacum sanctum</i>	1.1432	119	<i>Pyrus rivularis</i>	0.8310	77	<i>Robinia Pseudacacia</i>	0.7333
114	<i>Vauquelinia Torreyi</i>	1.1374	257	<i>Quercus lyrata</i>	0.8313	276	<i>Quercus nigra</i>	0.7324
147	<i>Eugenia longipes</i>	1.1235	292	<i>Ostrya Virginica</i>	0.8284	158	<i>Viburnum Lentago</i>	0.7303
21	<i>Porlieria angustifolia</i>	1.1101	270	<i>Quercus agrifolia</i>	0.8263	278	<i>Quercus Catesbei</i>	0.7294
220	<i>Sebastiania lucida</i>	1.0905	244	<i>Carya tomentosa</i>	0.8218	178	<i>Bumelia tenax</i>	0.7293
183	<i>Mimusops Sieberi</i>	1.0838	245	<i>Carya porcin</i>	0.8217	130	<i>Crataegus cordata</i>	0.7293
115	<i>Cercocarpus ledifolius</i>	1.0731	49	<i>Colubrina reclinata</i>	0.8208	228	<i>Celtis occidentalis</i>	0.7287
42	<i>Reynoldsia latifolia</i>	1.0715	106	<i>Prunus umbellata</i>	0.8202	293	<i>Carpinus Caroliniana</i>	0.7286
80	<i>Olneya Tesota</i>	1.0602	151	<i>Cornus florida</i>	0.8153	31	<i>Swietenia Mahagoni</i>	0.7283
80	<i>Amyris sylvatica</i>	1.0459	54	<i>Sapindus marginatus</i>	0.8126	228	<i>Celtis occidentalis, var. reticulata</i>	0.7275
162	<i>Genipa clusifolia</i>	1.0316	201	<i>Osmanthus Americanus</i>	0.8111	35	<i>Ilex Cassine</i>	0.7270
176	<i>Sideroxylon Mastichodendron</i>	1.0109	243	<i>Carya sulcata</i>	0.8108	225	<i>Ulmus racemosa</i>	0.7263
264	<i>Quercus grisea</i>	1.0092	78	<i>Robinia viscosa</i>	0.8094	222	<i>Ulmus crassifolia</i>	0.7245
141	<i>Conocarpus erecta</i>	0.9900	204	<i>Bourreria Havanensis</i>	0.8073	280	<i>Quercus aquatica</i>	0.7244
12	<i>Canella alba</i>	0.9893	259	<i>Quercus Michauxii</i>	0.8039	103	<i>Prunus Americana</i>	0.7215
83	<i>Sophora secundiflora</i>	0.9642	79	<i>Robinia Neo-Mexicana</i>	0.8034	126	<i>Crataegus Crus-galli</i>	0.7194
213	<i>Coccoloba Floridana</i>	0.9835	247	<i>Carya myristiciformis</i>	0.8016	196	<i>Fraxinus quadrangulata</i>	0.7184
113	<i>Prunus ilicifolia</i>	0.9803	284	<i>Quercus hypoleuca</i>	0.8009	241	<i>Carya oliviformis</i>	0.7180
214	<i>Coccoloba uvifera</i>	0.9635	182	<i>Bumelia cuneata</i>	0.7959	407	<i>Thrinax argentea</i>	0.7172
56	<i>Hypelate paniculata</i>	0.9533	128	<i>Crataegus subvillosa</i>	0.7953	170	<i>Kalmia latifolia</i>	0.7160
266	<i>Quercus Durandii</i>	0.9507	372	<i>Pinus serotina</i>	0.7942	132	<i>Crataegus spathulata</i>	0.7159
267	<i>Quercus virens</i>	0.9501	73	<i>Rhus Metopium</i>	0.7917	16	<i>Fremontia Californica</i>	0.7143
265	<i>Quercus reticulata</i>	0.9479	184	<i>Diospyros Virginiana</i>	0.7906	142	<i>Laguncularia racemosa</i>	0.7137
148	<i>Eugenia procera</i>	0.9453	189	<i>Fraxinus Greggii</i>	0.7904	338	<i>Juniperus occidentalis, var. monosperma</i>	0.7118
263	<i>Quercus oblongifolia</i>	0.9441	109	<i>Prunus Capuli</i>	0.7879	194	<i>Fraxinus viridis</i>	0.7117
97	<i>Acacia Wrightii</i>	0.9392	271	<i>Quercus Wislizeni</i>	0.7855	202	<i>Cordia Sebestena</i>	0.7108
116	<i>Cercocarpus parvifolius</i>	0.9366	137	<i>Amelanchier Canadensis</i>	0.7838	167	<i>Arbutus Xalapensis</i>	0.7099
144	<i>Eugenia buxifolia</i>	0.9360	135	<i>Crataegus flava</i>	0.7809	166	<i>Arbutus Menziesii</i>	0.7052
175	<i>Chrysophyllum oliviforme</i>	0.9360	41	<i>Schæfferia frutescens</i>	0.7745	117	<i>Pyrus coronaria</i>	0.7048
219	<i>Drypetes crocea, var. latifolia</i>	0.9346	234	<i>Maclura aurantiaca</i>	0.7736	274	<i>Quercus tinctoria</i>	0.7045
136	<i>Heteromeles arbutifolia</i>	0.9326	233	<i>Morus microphylla</i>	0.7715	390	<i>Pinus palustris</i>	0.6999
177	<i>Dipholia salicifolia</i>	0.9316	102	<i>Chrysobalanus Icaco</i>	0.7709	11	<i>Capparis Jamaicensis</i>	0.6971
160	<i>Exostemma Caribæum</i>	0.9310	122	<i>Crataegus rivularis</i>	0.7703	223	<i>Ulmus fulva</i>	0.6956
269	<i>Quercus Emoryi</i>	0.9263	216	<i>Neotandra Willdenoviana</i>	0.7693	110	<i>Prunus demissa</i>	0.6951
95	<i>Leucana glauca</i>	0.9235	135	<i>Crataegus flava, var. pubescens</i>	0.7683	123	<i>Crataegus Douglasii</i>	0.6950
219	<i>Drypetes crocea</i>	0.9200	281	<i>Quercus laurifolia</i>	0.7673	174	<i>Jacquinia armillaris</i>	0.6948
32	<i>Ximenia Americana</i>	0.9196	258	<i>Quercus bicolor</i>	0.7662	279	<i>Quercus palustris</i>	0.6923
146	<i>Eugenia monticola</i>	0.9156	93	<i>Prosopis juliflora</i>	0.7652	85	<i>Gymnocladus Canadensis</i>	0.6924
211	<i>A vicennia nitida</i>	0.9138	192	<i>Fraxinus Americana, var. Texensis</i>	0.7636	277	<i>Quercus falcata</i>	0.6923
57	<i>Hypelate trifoliata</i>	0.9102	129	<i>Crataegus tomentosa</i>	0.7633	64	<i>Acer saccharinum, var. nigrum</i>	0.6915
272	<i>Quercus rubra, var. Texana</i>	0.9080	299	<i>Betula lenta</i>	0.7617	64	<i>Acer saccharinum</i>	0.6912
161	<i>Pithecolobium Unguis-cati</i>	0.9049	164	<i>Vaccinium arboreum</i>	0.7610	338	<i>Juniperus occidentalis, var. conjugens</i>	0.6907
40	<i>Myginda pallens</i>	0.9048	94	<i>Prosopis pubescens</i>	0.7609	63	<i>Acer grandidentatum</i>	0.6902
24	<i>Xanthoxylum Caribæum</i>	0.9002	246	<i>Carya amara</i>	0.7552	118	<i>Pyrus angustifolia</i>	0.6895
112	<i>Prunus sphaerocarpa</i>	0.8998	285	<i>Quercus imbricaria</i>	0.7529	27	<i>Canotia holocantha</i>	0.6885
143	<i>Calyptranthes Chytraculia</i>	0.8992	92	<i>Cercia reniformis</i>	0.7513	104	<i>Prunus angustifolia</i>	0.6884
145	<i>Eugenia dichotoma</i>	0.8983	381	<i>Pinus Cubensis</i>	0.7504	291	<i>Fagus ferruginea</i>	0.6883
262	<i>Quercus Douglasii</i>	0.8928	168	<i>Arbutus Texana</i>	0.7500	138	<i>Hamamelis Virginica</i>	0.6856
75	<i>Eysenhardtia orthocarpa</i>	0.8740	165	<i>Andromeda ferruginea</i>	0.7500	282	<i>Quercus heterophylla</i>	0.6834
81	<i>Piscidia Erythrina</i>	0.8734	260	<i>Quercus Prinus</i>	0.7499	287	<i>Quercus densiflora</i>	0.6827
210	<i>Citharexylum villosum</i>	0.8710	226	<i>Ulmus alata</i>	0.7491	191	<i>Fraxinus pistaciifolia</i>	0.6810
111	<i>Prunus Caroliniana</i>	0.8688	152	<i>Cornus Nuttallii</i>	0.7481	124	<i>Crataegus brachysantha</i>	0.6798
127	<i>Crataegus coccinea</i>	0.8618	286	<i>Quercus Phellos</i>	0.7472	203	<i>Cordia Bolanderi</i>	0.6790
261	<i>Quercus prinoides</i>	0.8605	251	<i>Quercus alba</i>	0.7470	37	<i>Cyrilla racemiflora</i>	0.6784
173	<i>Ardisia Pickeringia</i>	0.8602	181	<i>Bumelia lycioides</i>	0.7467	397	<i>Abies bracteata</i>	0.6783
98	<i>Acacia Greggii</i>	0.8550	169	<i>Oxydendrum arboreum</i>	0.7438	86	<i>Gleditsia triacanthos</i>	0.6740
84	<i>Sophora affinis</i>	0.8509	131	<i>Crataegus apifolia</i>	0.7453	96	<i>Leucaena pulverulenta</i>	0.6733
268	<i>Quercus chrysoplepis</i>	0.8493	253	<i>Quercus Garryana</i>	0.7449	250	<i>Myrica Californica</i>	0.6703
185	<i>Diospyros Texana</i>	0.8460	256	<i>Quercus macrocarpa</i>	0.7453	150	<i>Cornus alternifolia</i>	0.6696
235	<i>Quercus undulata, var. Gambelii</i>	0.8407	89	<i>Parkinsonia microphylla</i>	0.7453	409	<i>Yucca canaliculata</i>	0.6677
242	<i>Carya alba</i>	0.8372	25	<i>Xanthoxylum Pterota</i>	0.7444	61	<i>Acer circinatum</i>	0.6660
55	<i>Sapindus Saponaria</i>	0.8367	36	<i>Ilex decidua</i>	0.7420	180	<i>Bumelia spinosa</i>	0.6603
254	<i>Quercus obtusiloba</i>	0.8367	252	<i>Quercus lobata</i>	0.7409	190	<i>Fraxinus anomala</i>	0.6597
172	<i>Myrsine Rapanca</i>	0.8341	248	<i>Carya aquatica</i>	0.7407	39	<i>Euonymus atropurpureus</i>	0.6592

Catalogue number.	Species.	Relative specific gravity.	Catalogue number.	Species.	Relative specific gravity.	Catalogue number.	Species.	Relative specific gravity.
134	<i>Crataegus metivialis</i>	0.6564	187	<i>Halesia diptera</i>	0.5705	379	<i>Pinus Banksiana</i>	0.4761
240	<i>Juglans rupestris</i>	0.6554	235	<i>Platanus occidentalis</i>	0.5678	346	<i>Torreya Californica</i>	0.4780
297	<i>Betula lutea</i>	0.6553	353	<i>Pinus Parryana</i>	0.5675	309	<i>Salix lasiandra</i>	0.4754
179	<i>Bumelia lanuginosa</i>	0.6544	47	<i>Rhamnus Purshiana</i>	0.5672	231	<i>Ficus pedunculata</i>	0.4739
192	<i>Fraxinus Americana</i>	0.6543	356	<i>Pinus monophylla</i>	0.5658	237	<i>Platanus Wrightii</i>	0.4736
272	<i>Quercus rubra</i>	0.6540	22	<i>Xanthoxylum Americanum</i>	0.5654	14	<i>Gordonia Lasianthus</i>	0.4728
88	<i>Parkinsonia Torreyana</i>	0.6531	249	<i>Myrica cerifera</i>	0.5637	361	<i>Pinus ponderosa</i>	0.4715
212	<i>Pisonia obtusata</i>	0.6529	188	<i>Halesia tetraptera</i>	0.5628	400	<i>Abies magnifica</i>	0.4701
218	<i>Umbellularia Californica</i>	0.6517	316	<i>Salix lasiolepis</i>	0.5587	3	<i>Magnolia acuminata</i>	0.4690
354	<i>Pinus cembroides</i>	0.6512	374	<i>Pinus clausa</i>	0.5576	333	<i>Cupressus Goveniana</i>	0.4689
224	<i>Ulmus Americana</i>	0.6506	288	<i>Castanopsis chrysophylla</i>	0.5574	304	<i>Alnus serrulata</i>	0.4686
125	<i>Crataegus arborescens</i>	0.6491	357	<i>Pinus Balfouriana</i> , var. <i>aristata</i>	0.5573	319	<i>Populus grandidentata</i>	0.4632
205	<i>Ehretia elliptica</i>	0.6440	76	<i>Dalea spinosa</i>	0.5536	325	<i>Populus Fremontii</i> , var. <i>Wialisii</i>	0.4621
275	<i>Quercus Kelloggii</i>	0.6435	336	<i>Juniperus Californica</i> , var. <i>Utahensis</i>	0.5522	331	<i>Chamaecyparis Lawsoniana</i>	0.4621
215	<i>Persea Carolinensis</i>	0.6429	45	<i>Rhamnus Caroliniana</i>	0.5462	157	<i>Sambucus Mexicana</i>	0.4614
69	<i>Rhus cotinoides</i>	0.6425	66	<i>Acer rubrum</i> , var. <i>Drummondii</i>	0.5459	153	<i>Nyssa capitata</i>	0.4613
283	<i>Quercus cinerea</i>	0.6420	363	<i>Pinus Chihuahuana</i>	0.5457	305	<i>Alnus incana</i>	0.4607
100	<i>Lysiloma latisiliqua</i>	0.6418	120	<i>Pyrus Americana</i>	0.5451	309	<i>Salix lasiandra</i> , var. <i>Fendleriana</i>	0.4598
230	<i>Ficus brevifolia</i>	0.6398	370	<i>Pinus Taeda</i>	0.5441	382	<i>Picea nigra</i>	0.4584
215	<i>Persea Carolinensis</i> , var. <i>palustris</i>	0.6396	357	<i>Pinus Balfouriana</i>	0.5434	368	<i>Pinus insignis</i>	0.4574
343	<i>Taxus brevifolia</i>	0.6391	313	<i>Salix flavescens</i> , var. <i>Sconleriana</i>	0.5412	391	<i>Pseudotsuga Douglasii</i> , var. <i>macrocarpa</i>	0.4563
355	<i>Pinus edulis</i>	0.6388	101	<i>Pinckneya pubens</i>	0.5350			
200	<i>Chionanthus Virginica</i>	0.6372	314	<i>Salix Hookeriana</i>	0.5350	399	<i>Abies nobilis</i>	0.4561
91	<i>Cercis Canadensis</i>	0.6363	310	<i>Salix longifolia</i> , var. <i>exigua</i>	0.5342	309	<i>Salix lasiandra</i> , var. <i>lanceifolia</i>	0.4547
1	<i>Magnolia grandiflora</i>	0.6360	59	<i>Acer spicatum</i>	0.5330	340	<i>Taxodium distichum</i>	0.4543
154	<i>Nyssa sylvatica</i>	0.6356	186	<i>Symlocos tinctoria</i>	0.5325	50	<i>Aesculus glabra</i>	0.4542
199	<i>Forestiera acuminata</i>	0.6345	5	<i>Magnolia macrophylla</i>	0.5309	17	<i>Tilia Americana</i>	0.4525
344	<i>Taxus Floridana</i>	0.6340	373	<i>Pinus inops</i>	0.5309	290	<i>Castanea vulgaris</i> , var. <i>Americana</i>	0.4504
53	<i>Ungnadia speciosa</i>	0.6332	58	<i>Acer Pennsylvanicum</i>	0.5299	107	<i>Prunus emarginata</i>	0.4502
209	<i>Crescentia cucurbitina</i>	0.6319	227	<i>Planera aquatica</i>	0.5294	307	<i>Salix amygdaloides</i>	0.4502
198	<i>Fraxinus sambucifolia</i>	0.6318	71	<i>Rhus copallina</i>	0.5273	6	<i>Magnolia Umbrella</i>	0.4487
171	<i>Rhododendron maximum</i>	0.6303	65	<i>Acer dasycarpum</i>	0.5269	208	<i>Catalpa bignonioides</i>	0.4474
336	<i>Juniperus Californica</i>	0.6282	362	<i>Pinus Jeffreyi</i>	0.5206	411	<i>Yucca elata</i>	0.4470
82	<i>Cladonia tinctoria</i>	0.6278	155	<i>Nyssa uniflora</i>	0.5194	308	<i>Salix nigra</i>	0.4456
332	<i>Cupressus macrocarpa</i>	0.6261	71	<i>Rhus copallina</i> , var. <i>lanceolata</i>	0.5184	390	<i>Tsuga Pattoniana</i>	0.4454
193	<i>Fraxinus pubescens</i>	0.6251	389	<i>Tsuga Mertensiana</i>	0.5182	404	<i>Sabal Palmetto</i>	0.4404
38	<i>Cliftonia ligustrina</i>	0.6249	405	<i>Washingtonia filifera</i>	0.5173	311	<i>Salix sessilifolia</i>	0.4397
401	<i>Larix Americana</i>	0.6236	391	<i>Pseudotsuga Douglasii</i>	0.5157	72	<i>Rhus venenata</i>	0.4382
66	<i>Acer rubrum</i>	0.6178	371	<i>Pinus rigida</i>	0.5151	350	<i>Pinus flexilis</i>	0.4358
90	<i>Parkinsonia aculeata</i>	0.6116	345	<i>Torreya taxifolia</i>	0.5145	70	<i>Rhus typhina</i>	0.4357
239	<i>Juglans nigra</i>	0.6115	156	<i>Sambucus glauca</i>	0.5087	67	<i>Negundo aceroides</i>	0.4328
377	<i>Pinus mitis</i>	0.6104	317	<i>Salixitchensis</i>	0.5072	386	<i>Piceaitchensis</i>	0.4287
315	<i>Salix cordata</i> , var. <i>vestita</i>	0.6069	23	<i>Xanthoxylum Clava-Herculia</i>	0.5056	388	<i>Tsuga Caroliniana</i>	0.4275
408	<i>Oreodoxa regia</i>	0.6034	10	<i>Anona laurifolia</i>	0.5053	51	<i>Aesculus flava</i>	0.4274
296	<i>Betula occidentalis</i>	0.6030	217	<i>Sassafras officinale</i>	0.5042	312	<i>Salix discolor</i>	0.4261
62	<i>Acer glabrum</i>	0.6028	360	<i>Pinus Arizona</i>	0.5038	18	<i>Tilia heterophylla</i>	0.4253
46	<i>Rhamnus Californica</i>	0.6000	2	<i>Magnolia glauca</i>	0.5035	387	<i>Tsuga Canadensis</i>	0.4239
406	<i>Thrinax parviflora</i>	0.5991	105	<i>Prunus Pennsylvanica</i>	0.5023	8	<i>Liriodendron Tulipifera</i>	0.4230
23	<i>Xanthoxylum Clava-Herculia</i> , var. <i>fruticosum</i>	0.5967	7	<i>Magnolia Fraseri</i>	0.5003	398	<i>Abies amabilis</i>	0.4228
			300	<i>Alnus maritima</i>	0.4996	342	<i>Sequoia sempervirens</i>	0.4208
295	<i>Betula papyrifera</i>	0.5955	52	<i>Aesculus Californica</i>	0.4980	207	<i>Catalpa speciosa</i>	0.4165
121	<i>Pyrus sambucifolia</i>	0.5928	313	<i>Salix flavescens</i>	0.4969	351	<i>Pinus albicaulis</i>	0.4165
139	<i>Liquidambar Styraciflua</i>	0.5909	376	<i>Pinus muricata</i>	0.4942	321	<i>Populus balsamifera</i> , var. <i>candicans</i>	0.4161
208	<i>Chilopsis saligna</i>	0.5902	375	<i>Pinus pungens</i>	0.4935	4	<i>Magnolia cordata</i>	0.4139
22	<i>Morus rubra</i>	0.5898	310	<i>Salix longifolia</i>	0.4930	28	<i>Simaruba glauca</i>	0.4136
19	<i>Byrsonima lucida</i>	0.5888	339	<i>Juniperus Virginiana</i>	0.4926	367	<i>Pinus Coulteri</i>	0.4133
289	<i>Castanea pumila</i>	0.5887	325	<i>Populus Fremontii</i>	0.4914	302	<i>Alnus rhombifolia</i>	0.4127
34	<i>Ilex Dahoon</i> , var. <i>myrtifolia</i>	0.5873	60	<i>Acer macrophyllum</i>	0.4909	305	<i>Pinus Murrayana</i>	0.4096
337	<i>Juniperus pachyphloea</i>	0.5829	236	<i>Platanus racemosa</i>	0.4880	320	<i>Populus heterophylla</i>	0.4089
108	<i>Prunus serotina</i>	0.5822	359	<i>Pinus Torreyana</i>	0.4879	238	<i>Juglans cinerea</i>	0.4066
33	<i>Ilex opaca</i>	0.5818	352	<i>Pinus reflexa</i>	0.4877	17	<i>Tilia Americana</i> , var. <i>pubescens</i>	0.4074
364	<i>Pinus contorta</i>	0.5815	308	<i>Salix levigata</i>	0.4872	383	<i>Picea alba</i>	0.4051
194	<i>Fraxinus viridis</i> , var. <i>Berlandieriana</i>	0.5780	358	<i>Pinus resinosa</i>	0.4854	318	<i>Populus tremuloides</i>	0.4032
11	<i>Hippomane Mancinella</i>	0.5772	325	<i>Cupressus Guadalupensis</i>	0.4843	326	<i>Libocedrus decurrens</i>	0.4017
338	<i>Juniperus occidentalis</i>	0.5765	366	<i>Pinus Sabiniana</i>	0.4840	303	<i>Alnus oblongifolia</i>	0.3981
298	<i>Betula nigra</i>	0.5762	68	<i>Negundo Californicum</i>	0.4821	9	<i>Asimina triloba</i>	0.3969
294	<i>Betula alba</i> , var. <i>populifolia</i>	0.5760	301	<i>Alnus rubra</i>	0.4813	378	<i>Pinus glabra</i>	0.3931
48	<i>Ceanothus thyrsiflorus</i>	0.5750	34	<i>Ilex Dahoon</i>	0.4806	322	<i>Populus angustifolia</i>	0.3912
197	<i>Fraxinus Oregana</i>	0.5731	330	<i>Chamaecyparis Nutkaensis</i>	0.4782	348	<i>Pinus monticola</i>	0.3908

Catalogue number.	Species.	Relative specific gravity.	Catalogue number.	Species.	Relative specific gravity.	Catalogue number.	Species.	Relative specific gravity.
324	<i>Populus monillifera</i>	0.3839	396	<i>Abies concolor</i>	0.3638	329	<i>Chamaecyparis sphaeroides</i>	0.3823
347	<i>Pinus Strobus</i>	0.3854	321	<i>Populus balsamifera</i>	0.3635	149	<i>Cereus giganteus</i>	0.3188
398	<i>Abies balsamea</i>	0.3819	392	<i>Abies Fraseri</i>	0.3565	327	<i>Thuja occidentalis</i>	0.3164
323	<i>Populus trichocarpa</i>	0.3814	395	<i>Abies grandis</i>	0.3545	29	<i>Bursera gummifera</i>	0.3008
328	<i>Thuja gigantea</i>	0.3796	195	<i>Fraxinus platycarpa</i>	0.3541	341	<i>Sequoia gigantea</i>	0.2983
385	<i>Picea pungens</i>	0.3740	369	<i>Pinus tuberculata</i>	0.3499	412	<i>Yucca baccata</i>	0.3724
410	<i>Yucca brevifolia</i>	0.3737	394	<i>Abies subalpina</i>	0.3476	229	<i>Ficus aurea</i>	0.3616
349	<i>Pinus Lambertiana</i>	0.3684	384	<i>Picea Engelmanni</i>	0.3449			

It will be noticed that all species in which the wood is heavier than water belong to the semi-tropical region of Florida or to the arid Mexican and interior Pacific regions. There seems to be a certain, but by no means constant relation, as shown in this table, between aridity of climate and the weight of the wood produced by closely allied species or by individuals of the same species. The wood of the form of *Quercus rubra* peculiar to western Texas is nearly 39 per cent. heavier than the average of all the specimens of the typical species grown in the northern states. Among the white oaks the wood of species belonging to regions of little rainfall, *Quercus grisea*, *oblongifolia*, *Durandii*, and *Douglasii*, is heavier than that of allied species peculiar to regions more favorable for the growth of trees. The average of two specimens of *Quercus prinoides* grown in western Texas is 19 per cent. heavier than the average of all the other specimens of this species grown in other parts of the country. In *Fraxinus*, the wood of *F. Greggii* of the Rio Grande valley is heavier than that of any other species; it only just surpasses in weight, however, the wood of the western Texas form of *F. Americana*, which is 20 per cent. heavier than the average of all specimens of the typical species grown north of Texas. On the other hand, the wood of Texas forms of *Fraxinus viridis* is constantly lighter than that of northern specimens, and the wood of *Celtis* grown in Arizona is lighter than that of the average of all the other specimens of this species. In *Juglans*, the heaviest wood is that of *J. rupestris*, a species belonging to a region of little rainfall, and a specimen of *J. nigra* from western Texas is 33 per cent. heavier than the average of all specimens grown in the Mississippi basin. In the case of *Platanus*, the heaviest wood is that of the Atlantic species, but wood of the species peculiar to the comparatively moist climate of southwestern Arizona is, however, considerably lighter than that of the drier climate of southern California.

FUEL VALUE.

The relative fuel values are obtained by deducting the percentage of ash from the specific gravity, and are based on the hypothesis that the real value of the combustible material in all woods is the same.

A number of analyses was also made of the wood of several of the principal trees of the United States (Table II) and their absolute fuel value calculated. Mr. Sharples describes the methods adopted by him to obtain these results, as follows:

The carbon and hydrogen determinations were made by the ordinary processes of organic analysis, by burning the wood in a current of oxygen. The moisture was determined by drying the wood at 100° centigrade until its weight became sensibly constant. The calculations were then made on the dry wood. The results contain a slight constant error, arising from the fact that the nitrogen in the wood was not determined. This error is, however, very slight, the nitrogen, which is included in the percentage of oxygen, rarely amounting, in any wood, to one per cent. The column headed "Hydrogen combined with oxygen", is found by dividing the amount in the column headed "Oxygen" by eight, and represents the hydrogen that may be considered as already combined with oxygen in the form of water, and is therefore useless for fuel. The fuel value per kilogram is found by multiplying the percentage of carbon by 8,080, and that of excess of hydrogen by 34,462 (these being the values obtained by *Favre* and *Silberman*), adding these together and deducting from the sum the product of the total hydrogen multiplied by 4,833, which represents the heat required to evaporate the water produced by burning the hydrogen. The constants used above represent the number of kilograms of water raised one degree centigrade, by burning one kilogram of carbon or hydrogen. The fuel value per cubic decimeter is found by multiplying the value per kilogram by the specific gravity. It need hardly be said that this fuel value is rarely attained in practice, and that it is never utilized. There are too many sources of loss; the calculation supposes that the combustion is perfect, that no smoke is given off, and that the heat of the products of combustion, with the exception of that necessary to convert the water into vapor, is all utilized.

It appears from Mr. Sharples' experiments that resinous woods give upward of 12 per cent. more heat from equal weights burned than non-resinous woods; the heat produced by burning a kilogram of dry non-resinous wood being about 4,000 units, while the heat produced by burning a kilogram of dry resinous wood is about 4,500 units, a unit being the quantity of heat required to raise 1 kilogram of water 1 degree centigrade.

Count Rumford first propounded the theory that the value of equal weights of wood for fuel was the same without reference to specific distinctions; that is, that a pound of wood, whatever the variety, would always produce the same amount of heat (*Count Rumford's Works*, Boston, 1873, vol. ii). Marcus Bull, experimenting in 1826 upon the fuel value of different woods (*Trans. Am. Phil. Soc.*, new ser., iii, 1), found a variation of only 11 per cent. between the different species tested. Rumford's theory must be regarded as nearly correct, if woods are

separated into resinous and non-resinous classes. The specific gravity gives a direct means of comparing heat values of equal volumes of wood of different resinous and non-resinous species. In burning wood, however, various circumstances affect its value; few fire-places are constructed to fully utilize the fuel value of resinous wood, and carbon escapes unconsumed in the form of smoke. Pine, therefore, which, although capable of yielding more heat than oak or hickory, may in practice yield considerably less, the pine losing both carbon and hydrogen in the form of smoke, while hickory or oak, burning with a smokeless flame, is practically entirely consumed. The ash in a wood, being non-combustible, influences its fuel value in proportion to its amount. The state of dryness of wood also has much influence upon its fuel value, though to a less degree than is generally supposed. The water in green wood prevents its rapid combustion, evaporation reducing the temperature below the point of ignition. Green wood may often contain as much as 50 per cent. of water, and this water must evaporate during combustion; but as half a kilogram of ordinary wood will give 2,000 units of heat, while half a kilogram of water requires only 268.5 units to evaporate it, 1731.5 units remain available for generating heat in wood containing even a maximum amount of water. In cases where the pressure was perpendicular to the grain of the wood it was applied on the side of the specimen nearest to the heart of the tree.

A factor in the general value of wood as fuel is the ease with which it can be seasoned; beech, for example, a very dense wood of high fuel value when dried, is generally considered of little value as fuel, on account of the rapidity with which it decays when cut and the consequent loss of carbon by decomposition.

THE STRENGTH OF WOOD.

The specimens tested for the purpose of determining the strength of the wood produced by the different trees of the United States were cut, with few exceptions, before March, 1881, and were slowly and carefully seasoned.

Those used in determining the resistance to transverse strain were made 4 centimeters square and long enough to give the necessary bearing upon the supports. These were shod with flat iron plates, slightly rounded on the edges and were set exactly 1 meter apart; they remained perfectly rigid under the pressure applied. Each specimen was weighed, measured, and its specific gravity calculated before it was tested. The result thus obtained represents the specific gravity of the air-dried wood.

To eliminate the action of their weight the specimens were placed upright, and hydraulic pressure was applied by means of an iron rod 12 millimeters in radius, acting midway between the supports, the deflections being read at this point.

The direction of the grain of the wood is shown by diagrams in the table (Table III), the pressure acting upon it horizontally from the left.

The pressure was applied slowly and uniformly, a reading of the deflections being taken for every 50 kilograms. When a load of 200 kilograms had been applied it was removed and the set read. Pressure was again applied in the same way, and the readings of deflections were resumed when 200 kilograms was again reached.

The formula used in calculating the coefficient of elasticity was $E = \frac{P l}{4 \Delta b d^3}$; l, b, d , being taken in millimeters; that of the modulus of rupture, $R = \frac{3 P l}{2 b d^2}$, l, b, d being in centimeters, P , in both formulas, in kilograms.

A few experiments were also made in the same manner, for purposes of comparison, to determine the transverse strength of specimens 1 meter long between the bearings and 8 centimeters square (Table IV).

The specimens tested by longitudinal compression were 4 centimeters square and 32 centimeters (8 diameters) long. They were placed between the platforms of the machine, and pressure was gradually applied until they failed. The figures given represent the number of kilograms required to cause failure.

The specimens tested under pressure applied perpendicularly to the fibers were 4 centimeters square and 16 centimeters long. They were placed upon the platform of the machine and indented with an iron punch 4 centimeters square on its face, covering the entire width of the specimen and one-quarter of its length at the center. In this series of experiments the direction of the annual rings was noted, horizontal pressure being also applied from the left. Readings were taken of the pressure necessary to produce each successive indentation of 0.254 up to 2.54 millimeters, and in the case of specimens which did not fail with this pressure a further test was made of the weight required to produce indentations of 3.81 and 5.08. The remarks (Table V) upon the behavior of the wood of the different species under compression were furnished by Mr. James E. Howard, in charge of the testing machine.

COMPARATIVE VALUES.

In the following table the number standing opposite each species represents its relative value in the column in which it appears.

This table is purely an arbitrary one, since the introduction of one or more species would of course change the value of all species standing lower in value, or results based on an examination of a larger number of specimens of any species may change the relative numbers in regard to it very considerably. In other words, any twenty or thirty species bearing consecutive numbers may change places with each other. This arises partly from the want of uniformity of the wood of any species, and partly from the fact that where so many determinations fall between comparatively narrow limits the mere order of sequence must be largely accidental.

TABLE OF RELATIVE VALUES.

Catalogue number.	Species.	Approximate fuel value.	Elasticity.	Ultimate transverse strength.	Ultimate resistance to longitudinal crushing.	Resistance to indentation to 1.27 millimeters.	Catalogue number.	Species.	Approximate fuel value.	Elasticity.	Ultimate transverse strength.	Ultimate resistance to longitudinal crushing.	Resistance to indentation to 1.27 millimeters.
MAGNOLIACEÆ.							ANACARDIACEÆ.						
1	Magnolia grandiflora	145	183	135	115	124	71	Rhus copallina	198	210	205	227	216
2	Magnolia glauca	205	127	163	173	234	78	Rhus Metopium	59	74	208	73	100
3	Magnolia acuminata	230	122	201	189	236	LEGUMINOSÆ.						
4	Magnolia cordata	265	119	236	191	248	77	Robinia Pseudacacia	87	19	8	12	71
5	Magnolia macrophylla	189	41	184	104	247	79	Robinia Neo-Mexicana	51	48	81	16	52
6	Magnolia Umbrella	249	207	245	234	257	80	Olneya Tesota	7	148	153	235	2
7	Magnolia Fraseri	207	114	179	182	202	81	Piscidia Erythrina	35	154	151	34	24
8	Liriodendron Tulipifera	259	124	208	229	262	82	Cladrastris tinctoria	150	95	88	71	141
ANONACEÆ.							84	Sophora affinis	34	102	121	48	27
9	Asimina triloba	277	281	293	296	285	85	Gymnocladus Canadensis	114	77	144	200	163
10	Anona laurifolia	220	278	234	277	198	86	Gleditschia triacanthos	124	64	77	93	156
CANELACEÆ.							87	Gleditschia monosperma	88	40	49	41	53
12	Canella alba	11	56	50	6	5	88	Parkinsonia Torreyana	134	264	259	184	90
TERNSTROMIACEÆ.							91	Cercis Canadensis	146	228	165	123	142
14	Gordonia Lasianthus	228	187	201	214	240	93	Prosopis juliflora	69	259	273	40	26
TILIACEÆ.							94	Prosopis pubescens	67	172	91	18	28
17	Tilia Americana	246	161	241	240	290	ROSACEÆ.						
17	Tilia Americana, var. pubescens	272	183	256	195	298	103	Prunus Americana	97	161	103	39	104
18	Tilia heterophylla	257	158	248	202	263	104	Prunus angustifolia	119	253	276	197	190
MALPIGHIACEÆ.							107	Prunus emarginata	248	149	194	124	266
19	Byrsonima lucida	169	273	288	206	100	108	Prunus serotina	164	153	115	61	114
ZYGOPHYLLACEÆ.							110	Prunus demissa	112	199	186	84	76
20	Gnaisson sanctum	8	147	187	10	1	111	Prunus Caroliniana	29	120	76	50	83
RUTACEÆ.							113	Prunus ilicifolia	10	212	188	63	89
23	Xanthoxylum Clava-Herculis	204	217	219	148	165	117	Pyrus coronaria	109	245	273	181	74
24	Xanthoxylum Caribæum	27	143	150	15	18	121	Pyrus sambucifolia	165	248	284	220	225
SIMARUBEÆ.							125	Crataegus arborescens	135	189	228	95	140
26	Simaruba glauca	268	121	262	171	251	126	Crataegus Crus-galli	100	240	210	169	107
BURSERACEÆ.							128	Crataegus subvillosa	52	135	162	68	63
29	Bursaria gummiifera	298	292	300	300	300	129	Crataegus tomentosa	65	212	177	153	80
30	Amyris sylvatica	5	65	2	9	6	132	Crataegus spatulata	102	237	268	189	90
MELIACEÆ.							134	Crataegus aestivalis	128	256	176	152	94
31	Swietenia Mahogoni	96	71	56	20	36	135	Crataegus flava	61	225	167	76	81
ILICINEÆ.							137	Amelanchier Canadensis	57	84	24	19	48
33	Ilex opaca	167	244	188	180	149	HAMAMELACEÆ.						
34	Ilex Dahoon	224	246	250	252	214	139	Liquidambar Styraciflua	162	167	214	128	192
CYRILLACEÆ.							RHIZOPHORACEÆ.						
38	Cliftonia ligustrina	152	191	262	281	177	140	Rhizophora Mangle	2	2	10	2	8
RHAMNACEÆ.							COMBRETACEÆ.						
42	Reynosa latifolia	6	74	116	3	4	141	Conocarpus erecta	9	89	72	32	19
43	Condalia ferrea	1	45	85	5	3	142	Laguncularia racemosa	108	219	264	147	174
45	Rhamnus Caroliniana	184	208	251	154	187	MYRTACEÆ.						
47	Rhamnus Purshiana	175	129	152	28	131	144	Eugenia buxifolia	18	6	38	1	20
SAPINDACEÆ.							146	Eugenia monticola	24	65	15	55	12
50	Æsculus glabra	245	243	271	273	282	148	Eugenia procera	19	38	14	17	9
52	Æsculus Californica	208	232	222	245	220	CORNACEÆ.						
54	Sapindus marginatus	47	165	112	122	58	151	Cornus florida	44	176	55	69	40
60	Acer macrophyllum	213	194	190	223	159	152	Cornus Nuttallii	74	85	60	22	77
61	Acer circinatum	126	221	147	186	119	153	Nyssa capitata	235	234	194	167	169
64	Acer saccharinum	117	9	18	30	73	154	Nyssa sylvatica	141	178	112	125	127
64	Acer saccharinum, var. nigrum	118	88	68	57	66	155	Nyssa uniflora	195	275	210	237	161
65	Acer dasycarpum	192	59	53	113	145	CAPRIFOLIACEÆ.						
66	Acer rubrum	154	117	121	183	151	156	Sambucus glauca	206	299	295	287	184
67	Negundo aceroides	256	290	263	270	215	159	Viburnum prunifolium	38	131	69	37	35
68	Negundo Californicum	221	113	132	155	221	RUBIACEÆ.						
							160	Exostemma Caribæum	15	36	55	8	7
							161	Pinckneya pubens	187	231	291	269	230

TABLE OF RELATIVE VALUES—Continued.

Catalogue number.	Species.	Approximate fuel value.	Elasticity.	Ultimate transverse strength.	Ultimate resistance to longitudinal crushing.	Resistance to indentation to 1.27 millimeters.	Catalogue number.	Species.	Approximate fuel value.	Elasticity.	Ultimate transverse strength.	Ultimate resistance to longitudinal crushing.	Resistance to indentation to 1.27 millimeters.
ERICACEÆ.													
165	Andromeda ferruginea.....	72	179	194	107	92	224	Ulmus Americana.....	136	205	110	146	138
166	Arbutus Menziesii.....	107	164	81	89	110	225	Ulmus racemosa.....	93	62	36	38	112
167	Arbutus Xalapensis.....	104	251	230	198	75	226	Ulmus alata.....	80	273	167	158	67
169	Oxydendrum arboreum.....	78	137	164	90	95	227	Planera aquatica.....	191	266	228	203	179
170	Kalmia latifolia.....	101	258	219	168	65	228	Celtis occidentalis.....	94	229	135	178	101
171	Rhododendron maximum.....	148	242	205	156	133	228	Celtis occidentalis, var. reticulata.....	98	143	127	160	56
SAPOTACEÆ.							229	Ficus aurea.....	300	300	298	299	296
175	Chrysophyllum oliviforme.....	17	54	107	33	15	231	Ficus pedunculata.....	244	292	290	284	207
176	Sideroxylon Mastichodendron.....	12	61	67	23	25	232	Morus rubra.....	163	173	141	179	148
177	Dipholia salicifolia.....	16	16	18	11	49	234	Maclura aurantiaca.....	63	114	25	4	22
178	Bumelia tenax.....	90	204	200	143	146	PLATANACEÆ.						
179	Bumelia lanuginosa.....	133	280	294	239	162	235	Platanus occidentalis.....	173	146	222	144	158
181	Bumelia lycioides.....	81	193	254	106	97	236	Platanus racemosa.....	218	249	254	269	245
182	Bumelia cuneata.....	56	252	265	117	47	237	Platanus Wrightii.....	231	286	285	267	211
183	Mimusops Sieberi.....	4	95	79	135	16	JUGLANDACEÆ.						
EBENACEÆ.							238	Juglans cinerea.....	270	181	238	205	244
184	Diospyros Virginiana.....	55	192	98	88	29	239	Juglans nigra.....	155	63	109	46	126
STYRACACEÆ.							240	Juglans rupestris.....	132	218	236	159	143
186	Symplocos tinctoria.....	190	250	230	218	139	241	Carya oliviformis.....	103	239	246	164	84
187	Halesia diptera.....	172	229	107	166	123	242	Carya alba.....	36	12	11	27	60
OLEACEÆ.							243	Carya sulcata.....	46	78	33	52	45
191	Fraxinus pistaciifolia.....	122	254	226	217	108	244	Carya tomentosa.....	43	42	20	36	51
192	Fraxinus Americana.....	130	91	106	121	153	245	Carya porcina.....	42	92	31	43	41
192	Fraxinus Americana, var. Texensis.....	66	69	27	65	121	246	Carya amara.....	71	86	30	78	78
193	Fraxinus pubescens.....	151	182	101	162	113	247	Carya myristiciformis.....	49	8	1	25	34
194	Fraxinus viridis.....	105	138	91	114	98	248	Carya aquatica.....	86	93	96	110	55
195	Fraxinus platycarpa.....	291	283	261	296	185	MYRICACEÆ.						
196	Fraxinus quadrangulata.....	73	196	121	94	111	249	Myrica cerifera.....	177	137	119	151	182
197	Fraxinus Oregana.....	171	156	203	80	157	250	Myrica Californica.....	125	99	46	74	135
198	Fraxinus sambucifolia.....	149	142	125	175	130	CUPULIFERÆ.						
199	Forestiera acuminata.....	147	227	175	199	154	251	Quercus alba.....	76	104	85	82	105
201	Osmanthus Americanus.....	45	28	40	51	64	252	Quercus lobata.....	85	222	103	172	136
BORRAGINACEÆ.							253	Quercus Garryana.....	79	183	99	97	79
204	Bourreria Havanensis.....	53	97	71	45	43	254	Quercus obtusiloba.....	37	169	100	108	54
205	Ehretia elliptica.....	142	206	169	215	87	255	Quercus undulata, var. Gambellii.....	82	262	194	186	69
BIGNONIACEÆ.							256	Quercus macrocarpa.....	82	122	65	103	82
206	Catalpa bignonioides.....	250	283	241	238	273	257	Quercus lyrata.....	89	17	50	100	72
207	Catalpa speciosa.....	263	175	224	194	254	258	Quercus bicolor.....	60	131	81	102	96
208	Chilopsis saligna.....	161	267	247	278	181	259	Quercus Michauxii.....	48	106	28	112	83
VERBENACEÆ.							260	Quercus Prinus.....	75	27	48	67	86
210	Citharexylum villosum.....	28	26	73	13	37	261	Quercus prinoides.....	31	53	6	44	61
NYCTAGINACEÆ.							262	Quercus Douglassii.....	26	197	57	53	17
212	Pisonia obtusata.....	157	284	296	274	218	263	Quercus oblongifolia.....	20	150	175	165	10
POLYGONACEÆ.							264	Quercus grisea.....	8	209	73	116	21
213	Coccoloba Florida.....	14	49	78	7	13	266	Quercus Durandii.....	21	165	57	70	38
LAURACEÆ.							267	Quercus virens.....	13	49	54	59	30
215	Persea Carolinensis.....	138	162	88	47	128	268	Quercus chrysolepis.....	33	33	4	62	33
215	Persea Carolinensis, var. palustris.....	140	162	116	232	132	269	Quercus Emoryi.....	22	247	180	176	11
217	Sassafras officinale.....	202	273	235	221	188	270	Quercus agrifolia.....	41	110	75	131	81
218	Umbellularia Californica.....	131	70	127	49	120	271	Quercus Wislizeni.....	58	148	118	72	57
EUPHORBIACEÆ.							272	Quercus rubra.....	129	48	63	83	150
219	Drypetes crocea.....	80	79	132	24	23	273	Quercus rubra, var. Texana.....	23	83	52	42	44
219	Drypetes crocea, var. latifolia.....	25	168	178	79	13	273	Quercus coccinea.....	84	65	38	87	116
URTICACEÆ.							274	Quercus tinctoria.....	106	81	44	92	115
222	Ulmus crassifolia.....	99	226	142	142	68	275	Quercus Kelloggii.....	137	206	145	149	152
223	Ulmus fulva.....	111	110	101	66	172	276	Quercus nigra.....	89	102	43	98	46
							277	Quercus falcata.....	113	11	12	35	117
							278	Quercus Catesbaei.....	91	80	42	137	88
							279	Quercus palustris.....	115	55	32	101	134
							280	Quercus aquatica.....	95	29	40	91	122
							281	Quercus laurifolia.....	62	25	13	77	70

TABLE OF RELATIVE VALUES—Continued.

Catalogue number.	Species.	Approximate fuel value.	Elasticity.	Ultimate transverse strength.	Ultimate resistance to longitudinal crushing.	Resistance to indentation to 1.27 millimeters.	Catalogue number.	Species.	Approximate fuel value.	Elasticity.	Ultimate transverse strength.	Ultimate resistance to longitudinal crushing.	Resistance to indentation to 1.27 millimeters.
282	<i>Quercus heterophylla</i>	121	80	35	190	144	347	<i>Pinus Strobus</i>	283	154	225	212	278
283	<i>Quercus cinerea</i>	144	203	57	150	118	348	<i>Pinus monticola</i>	279	112	232	203	288
284	<i>Quercus hypoleuca</i>	54	114	29	279	59	349	<i>Pinus Lambertiana</i>	287	187	238	202	270
285	<i>Quercus imbricaria</i>	68	37	8	56	89	350	<i>Pinus flexilis</i>	253	235	227	251	219
286	<i>Quercus Phellos</i>	77	190	63	211	102	351	<i>Pinus albicaulis</i>	262	276	244	265	224
287	<i>Quercus densiflora</i>	123	106	70	119	93	352	<i>Pinus reflexa</i>	215	127	143	105	200
288	<i>Castanopsis chrysophylla</i>	180	94	158	163	206	353	<i>Pinus Parryana</i>	174	297	287	258	129
289	<i>Castanea pumila</i>	160	46	60	99	209	355	<i>Pinus edulis</i>	143	291	283	253	106
290	<i>Castanea vulgaris, var. Americana</i>	247	152	184	222	227	356	<i>Pinus monophylla</i>	176	289	297	288	155
291	<i>Fagus ferruginea</i>	120	32	18	120	125	357	<i>Pinus Balfouriana</i>	185	255	288	260	178
292	<i>Ostrya Virginica</i>	40	15	23	64	85	357	<i>Pinus Balfouriana, var. aristata</i>	179	223	210	268	189
293	<i>Carpinus Caroliniana</i>	92	43	18	96	103	358	<i>Pinus resinosa</i>	217	51	131	140	256
BETULACEÆ.							359	<i>Pinus Torreyana</i>	200	270	149	272	180
294	<i>Betula alba, var. populifolia</i>	168	214	140	254	196	360	<i>Pinus Arizona</i>	203	174	210	204	217
295	<i>Betula papyrifera</i>	159	18	37	109	195	361	<i>Pinus ponderosa</i>	226	140	171	226	222
296	<i>Betula occidentalis</i>	158	126	127	207	171	362	<i>Pinus Jeffreyi</i>	201	125	156	185	212
297	<i>Betula lutea</i>	127	3	5	29	160	363	<i>Pinus Chihuahuana</i>	182	216	114	259	170
298	<i>Betula nigra</i>	170	58	66	157	183	364	<i>Pinus contorta</i>	166	4	60	54	175
299	<i>Betula lenta</i>	64	10	9	81	91	365	<i>Pinus Murrayana</i>	269	198	253	264	253
301	<i>Alnus rubra</i>	222	72	121	188	210	366	<i>Pinus Sabiniana</i>	219	257	139	261	186
302	<i>Alnus rhombifolia</i>	267	158	190	244	268	367	<i>Pinus Coulteri</i>	266	46	148	233	246
303	<i>Alnus oblongifolia</i>	276	199	189	285	279	368	<i>Pinus insignis</i>	239	101	158	183	229
SALICACEÆ.							369	<i>Pinus tuberculata</i>	293	290	290	293	255
307	<i>Salix amygdaloides</i>	251	277	257	291	264	370	<i>Pinus Tæda</i>	183	52	95	170	223
308	<i>Salix lævigata</i>	216	280	217	272	208	371	<i>Pinus rigida</i>	197	261	158	245	191
309	<i>Salix lasiandra, var. lancifolia</i>	243	141	199	257	249	372	<i>Pinus serotina</i>	50	39	17	86	42
309	<i>Salix lasiandra, var. Fendleriana</i>	236	298	276	281	261	373	<i>Pinus inops</i>	188	269	207	241	168
313	<i>Salix flaveacens</i>	209	65	125	192	241	374	<i>Pinus clausa</i>	178	268	269	228	194
313	<i>Salix flaveacens, var. Scouleriana</i>	186	22	81	126	199	375	<i>Pinus pungens</i>	211	186	166	247	213
316	<i>Salix lasiolepis</i>	181	137	120	216	183	376	<i>Pinus muricata</i>	210	35	47	85	204
318	<i>Populus tremuloides</i>	275	180	198	266	265	377	<i>Pinus mitis</i>	156	13	45	118	197
319	<i>Populus grandidentata</i>	234	108	169	243	295	378	<i>Pinus glabra</i>	278	288	270	280	228
320	<i>Populus heterophylla</i>	271	220	218	283	250	379	<i>Pinus Banksiana</i>	225	118	214	201	237
321	<i>Populus balsamifera</i>	288	150	257	271	276	380	<i>Pinus palustris</i>	110	7	18	26	167
321	<i>Populus balsamifera, var. candicans</i>	264	215	232	286	291	381	<i>Pinus Cubensis</i>	70	5	15	21	137
322	<i>Populus angustifolia</i>	280	284	292	296	274	382	<i>Picea nigra</i>	237	60	156	193	272
323	<i>Populus trichocarpa</i>	285	57	204	209	293	383	<i>Picea alba</i>	273	90	155	256	283
324	<i>Populus monilifera</i>	281	98	145	249	258	384	<i>Picea Engelmanni</i>	295	185	249	292	275
325	<i>Populus Fremontii</i>	214	73	183	225	252	385	<i>Picea pungens</i>	289	265	281	295	267
325	<i>Populus Fremontii, var. Wislizeni</i>	238	160	186	230	239	386	<i>Picea Sitchensis</i>	254	100	216	248	281
CONIFERÆ.							387	<i>Tsuga Canadensis</i>	258	135	171	219	260
326	<i>Libocedrus decurrens</i>	274	157	190	130	242	388	<i>Tsuga Caroliniana</i>	255	224	279	196	211
327	<i>Thuja occidentalis</i>	297	271	267	275	297	389	<i>Tsuga Mertensiana</i>	194	13	80	60	235
328	<i>Thuja gigantea</i>	284	81	154	145	284	390	<i>Tsuga Pattoniana</i>	252	195	171	224	232
329	<i>Chamaecyparis sphaeroides</i>	296	294	281	294	287	391	<i>Pseudotsuga Douglasii</i>	196	20	96	81	238
330	<i>Chamaecyparis Nutkaensis</i>	223	87	130	138	236	392	<i>Pseudotsuga Douglasii, var. macrocarpa</i>	240	74	111	132	233
331	<i>Chamaecyparis Lawsoniana</i>	233	31	93	129	259	393	<i>Abies Fraseri</i>	290	104	219	255	289
333	<i>Cupressus Goveniana</i>	232	279	260	242	147	394	<i>Abies balsamea</i>	283	177	265	236	277
338	<i>Juniperus occidentalis, var. conjugens</i>	116	211	276	75	50	395	<i>Abies subalpina</i>	294	201	275	276	294
339	<i>Juniperus Virginiana</i>	212	238	158	177	176	396	<i>Abies grandis</i>	292	109	271	268	299
340	<i>Taxodium distichum</i>	242	84	190	174	280	397	<i>Abies concolor</i>	289	130	180	210	269
341	<i>Sequoia gigantea</i>	299	287	280	213	286	398	<i>Abies amabilis</i>	260	24	134	127	292
342	<i>Sequoia sempervirens</i>	261	236	238	187	271	399	<i>Abies nobilis</i>	241	21	105	141	205
343	<i>Taxus brevifolia</i>	189	202	84	111	62	400	<i>Abies magnifica</i>	229	241	182	161	243
345	<i>Torreya taxifolia</i>	198	170	94	134	166	401	<i>Larix Americana</i>	153	23	90	58	231
346	<i>Torreya Californica</i>	227	294	243	250	203	402	<i>Larix occidentalis</i>	83	1	7	14	173
PALMACEÆ.							405	<i>Washingtonia filifera</i>	199	263	285	297	164

The following table gives the figures from which the table of relative values was computed, and includes all species upon which complete tests have been made.

The coefficient of elasticity is derived from the second deflection, the measurements being taken in millimeters and the weight in kilograms.

The ultimate transverse strength is the force, applied at the middle of the stick, required to break a stick 4 centimeters square and 1 meter between the supports.

In the compression tests the surface exposed to pressure was 4 centimeters square. To give the pressure on a square centimeter these results must be divided by 16.

The indentation to 1.27 millimeters, or the fifth in the series, is the one selected for comparison.

TABLE OF AVERAGES.

Catalogue number.	Species.	Approximate relative fuel value.	Coefficient of elasticity, kilograms on millimeters.	Ultimate transverse strength, in kilograms.	Ultimate resistance to longitudinal crushing, in kilograms.	Resistance to indentation to 1.27 millimeters, in kilograms.	Catalogue number.	Species.	Approximate relative fuel value.	Coefficient of elasticity, kilograms on millimeters.	Ultimate transverse strength, in kilograms.	Ultimate resistance to longitudinal crushing, in kilograms.	Resistance to indentation to 1.27 millimeters, in kilograms.
MAGNOLIACEÆ.							SAPINDACEÆ.						
1	<i>Magnolia grandiflora</i>	63.26	903	338	7,705	3,156	50	<i>Æsculus glabra</i>	45.03	644	211	5,017	1,132
2	<i>Magnolia glauca</i>	50.11	914	313	6,790	1,627	52	<i>Æsculus Californica</i>	49.45	683	271	5,686	1,723
3	<i>Magnolia acuminata</i>	46.76	929	286	6,633	1,709	54	<i>Sapindus marginatus</i>	50.05	837	360	7,523	4,350
4	<i>Magnolia cordata</i>	41.26	941	256	6,552	1,427	60	<i>Acer macrophyllum</i>	48.83	780	292	6,100	2,597
5	<i>Magnolia macrophylla</i>	52.90	1,169	297	7,829	1,427	61	<i>Acer circinatum</i>	66.34	718	327	7,349	3,205
6	<i>Magnolia Umbrella</i>	44.78	744	249	5,861	1,343	64	<i>Acer saccharinum</i>	68.75	1,465	480	9,907	4,019
7	<i>Magnolia Fraseri</i>	49.89	944	302	6,691	1,966	64	<i>Acer saccharinum, var. nigrum</i> ..	68.66	1,027	410	8,803	4,149
8	<i>Liriodendron Tulipifera</i>	42.20	926	280	5,955	1,296	65	<i>Acer dasycarpum</i>	52.52	1,110	435	7,711	2,899
ANONACEÆ.							66	<i>Acer rubrum</i>	61.65	943	346	7,402	2,795
9	<i>Asimina triloba</i>	39.61	482	167	3,395	1,098	67	<i>Negundo aceroides</i>	42.82	582	226	5,151	1,781
10	<i>Anona laurifolia</i>	48.11	501	259	4,829	2,037	68	<i>Negundo Californicum</i>	47.95	945	340	7,073	1,719
CANELACEÆ.							ANACARDIACEÆ.						
12	<i>Canella alba</i>	97.20	1,117	438	12,519	9,163	71	<i>Rhus copallina</i>	52.42	786	223	6,033	1,744
TERNSTROMIACEÆ.							73	<i>Rhus Metopium</i>	77.28	1,050	280	8,523	3,348
14	<i>Gordonia Lasianthus</i>	46.92	794	296	6,195	1,591	LEGUMINOSÆ.						
TILIACEÆ.							77	<i>Robinia Pseudacacia</i>	72.96	1,301	543	11,272	4,038
17	<i>Tilia Americana</i>	45.00	840	252	5,768	1,044	79	<i>Robinia Neo-Mexicana</i>	79.86	1,149	388	10,931	4,427
17	<i>Tilia Americana, var. pubescens</i> ..	40.47	811	239	6,467	950	80	<i>Olneya Tesota</i>	103.59	868	320	5,851	10,478
18	<i>Tilia heterophylla</i>	42.27	846	246	6,307	1,296	81	<i>Placidia Erythrina</i>	84.39	851	321	9,548	5,698
MALPIGHIACEÆ.							82	<i>Cladrastis tinctoria</i>	62.61	1,002	385	8,550	2,937
19	<i>Byrsonima lucida</i>	57.43	525	181	6,260	3,475	84	<i>Sophora affinis</i>	84.46	977	346	9,129	5,343
ZYGOPHYLLACEÆ.							85	<i>Gymnocladus Canadensis</i>	68.88	1,048	329	6,406	2,560
20	<i>Guaicum sanctum</i>	113.88	863	336	11,789	12,689	86	<i>Gleditsia triacanthos</i>	66.86	1,086	394	8,001	2,697
RUTACEÆ.							87	<i>Gleditsia monosperma</i>	72.89	1,170	439	9,344	4,420
23	<i>Xanthoxylum Clava-Herculis</i>	50.15	726	273	7,189	2,548	88	<i>Parkinsonia Torreyana</i>	64.58	556	233	6,679	3,620
24	<i>Xanthoxylum Caribæum</i>	88.20	868	322	10,955	5,964	91	<i>Cercis Canadensis</i>	63.18	688	310	7,510	2,917
SIMARUBEÆ.							93	<i>Prosopis juliflora</i>	74.86	583	207	9,412	5,484
28	<i>Simaruba glauca</i>	40.98	932	241	6,816	1,383	94	<i>Prosopis pubescens</i>	75.87	824	382	10,732	5,267
BURSERACEÆ.							ROSACEÆ.						
29	<i>Bursera gummiifera</i>	29.41	417	63	2,473	749	103	<i>Prunus Americana</i>	72.02	827	369	9,419	3,405
30	<i>Amyris sylvatica</i>	103.97	1,085	557	11,975	8,795	104	<i>Prunus angustifolia</i>	68.65	603	200	6,441	2,132
MELIACEÆ.							107	<i>Prunus emarginata, var. mollis</i> ..	44.93	861	290	7,507	1,280
31	<i>Swietenia Mahogoni</i>	72.08	1062	428	10,660	4,951	108	<i>Prunus serotina</i>	58.14	832	354	8,746	3,269
ILICINEÆ.							110	<i>Prunus demissa</i>	69.16	769	295	8,165	3,937
33	<i>Ilex opaca</i>	57.74	643	293	6,709	2,826	111	<i>Prunus Caroliniana</i>	86.52	937	396	8,989	5,090
34	<i>Ilex Dahoon</i>	47.62	642	244	5,582	1,808	113	<i>Prunus ilicifolia</i>	97.27	732	334	8,709	4,888
CYRILLACEÆ.							117	<i>Pyrus coronaria</i>	70.11	642	207	6,706	3,999
38	<i>Cliftonia ligustrina</i>	62.23	783	225	5,938	2,356	121	<i>Pyrus sambucifolia</i>	58.08	626	190	6,123	1,715
RHAMNACEÆ.							125	<i>Cratægus arborescens</i>	64.55	788	265	7,969	2,951
42	<i>Reynosia latifolia</i>	103.72	1,050	350	13,426	9,753	126	<i>Cratægus Crus-galli</i>	71.54	664	279	6,884	3,368
43	<i>Condalia ferrea</i>	119.38	1,143	386	12,848	10,388	128	<i>Cratægus subvillosa</i>	78.98	901	315	8,612	4,207
45	<i>Rhamnus Caroliniana</i>	54.27	741	242	7,112	2,195	129	<i>Cratægus tomentosa</i>	75.96	732	303	7,117	3,844
47	<i>Rhamnus Purshiana</i>	56.24	913	320	9,984	3,075	132	<i>Cratægus spathulata</i>	71.12	673	216	7,280	3,454
							134	<i>Cratægus setivialis</i>	65.27	592	304	7,122	3,583
							135	<i>Cratægus flava, var. pubescens</i> ..	76.13	708	309	8,437	5,103
							137	<i>Amelanchier Canadensis</i>	77.95	1,197	483	10,712	4,483
							HAMAMELACEÆ.						
							139	<i>Liquidambar Styraciflua</i>	58.78	837	278	7,463	2,123

TABLE OF AVERAGES—Continued.

Catalogue number.	Species.	Approximate relative fuel value.	Coefficient of elasticity, kilograms on millimeters.	Ultimate transverse strength, in kilograms.	Ultimate resistance to longitudinal crushing, in kilograms.	Resistance to indentation to 1.27 millimeters, in kilograms.	Catalogue number.	Species.	Approximate relative fuel value.	Coefficient of elasticity, kilograms on millimeters.	Ultimate transverse strength, in kilograms.	Ultimate resistance to longitudinal crushing, in kilograms.	Resistance to indentation to 1.27 millimeters, in kilograms.
RHIZOPHORACEÆ.							BIGNONIACEÆ.						
140	Rhizophora Mangle.....	114.06	1,666	515	13,767	7,394	206	Catalpa bignonioides.....	44.57	683	253	5,981	1,236
COMBRETACEÆ.							207	Catalpa speciosa.....	41.48	822	270	6,521	1,877
141	Conocarpus erecta.....	98.68	1,025	402	9,568	5,926	208	Chilopsis saligna.....	58.79	544	247	4,753	2,304
142	Laguncularia racemosa.....	70.21	734	221	7,190	2,386	VERBENACEÆ.						
MYRTACEÆ.							210	Citharexylum villosum.....	86.75	1,257	400	11,034	4,937
144	Eugenia buxifolia.....	92.20	1,575	450	14,198	5,851	NYCTAGINACEÆ.						
146	Eugenia monticola.....	89.83	1,065	500	8,845	6,582	212	Pisonia obtusata.....	60.31	465	127	4,982	1,787
148	Eugenia procera.....	92.05	1,191	502	10,750	7,099	POLYGONACEÆ.						
CORNACEÆ.							213	Coccoloba floridana.....	98.40	1,136	392	12,337	6,810
151	Cornus florida.....	80.98	821	398	8,553	4,875	LAURACEÆ.						
152	Cornus Nuttallii.....	74.44	1,081	423	10,803	3,883	215	Persea Carolinensis.....	68.81	839	385	9,173	3,128
153	Nyssa capitata.....	45.97	681	290	6,895	2,484	216	Persea Carolinensis, var. palustris.....	68.78	849	350	5,874	3,073
154	Nyssa sylvatica.....	68.66	818	360	7,497	3,181	217	Sassafras officinale.....	50.38	519	267	6,110	2,144
155	Nyssa uniflora.....	51.58	518	279	5,848	2,575	218	Umbellularia Californica.....	64.92	1,068	344	9,065	3,196
CAPRIFOLIACEÆ.							EUPHORBIACEÆ.						
156	Sambucus glauca.....	50.07	305	158	4,400	2,218	219	Drypetes crocea.....	86.44	1,089	340	10,410	5,797
159	Viburnum prunifolium.....	82.89	907	406	9,474	5,009	219	Drypetes crocea, var. latifolia.....	86.65	886	302	8,324	6,510
RUBIACEÆ.							URTICACEÆ.						
160	Exostemma Caribæum.....	92.89	1,194	429	12,030	7,707	222	Ulmus crassifolia.....	71.59	704	330	7,248	4,060
161	Pinckneya pubens.....	53.28	683	173	4,365	1,678	223	Ulmus fulva.....	68.77	963	371	8,628	2,899
ERICACEÆ.							224	Ulmus Americana.....	64.54	747	264	7,191	2,970
166	Andromeda ferruginea.....	74.66	814	290	7,802	3,611	225	Ulmus racemosa.....	72.20	1,096	455	9,474	3,281
166	Arbutus Menziesii.....	70.24	838	387	8,084	3,322	226	Ulmus alata.....	74.17	523	309	7,001	4,085
167	Arbutus Kalapensis.....	70.81	616	264	6,419	3,947	227	Planera aquatica.....	52.71	552	265	6,805	2,384
169	Oxydendrum arboreum.....	74.30	839	311	8,025	3,550	228	Celtis occidentalis.....	72.08	685	337	6,739	3,472
170	Kalmia latifolia.....	71.31	565	278	6,890	4,196	228	Celtis occidentalis, var. reticulata.....	71.86	868	344	6,985	4,373
171	Rhododendron maximum.....	62.80	646	283	7,030	3,086	229	Ficus aurea.....	24.84	257	102	2,597	980
SAPOTACEÆ.							231	Ficus pedunculata.....	45.07	407	98	4,491	1,905
173	Chrysophyllum oliviforme.....	92.44	1,124	368	9,571	6,108	232	Morus rubra.....	58.56	824	331	6,721	2,806
176	Sideroxylon Mastichodendron.....	95.89	1,099	414	10,410	5,682	234	Madura aurantiaca.....	76.01	944	483	12,939	5,806
177	Dipholia salicifolia.....	92.86	1,336	490	11,680	4,480	PLATANACEÆ.						
178	Bumelia tenax.....	72.39	751	287	7,235	2,894	235	Platanus occidentalis.....	58.52	864	371	7,207	2,645
179	Bumelia lanuginosa.....	64.64	463	165	5,799	2,564	236	Platanus racemosa.....	48.26	624	240	5,190	1,486
181	Bumelia lycioides.....	74.07	751	240	7,825	3,529	237	Platanus Wrightii.....	46.72	457	183	5,228	1,867
182	Bumelia cuneata.....	78.06	608	220	7,643	4,581	JUGLANDACEÆ.						
182	Mimusops Sieber.....	105.55	1,002	390	7,390	6,001	238	Juglans cinerea.....	40.66	812	255	6,270	1,488
EBENACEÆ.							239	Juglans nigra.....	60.91	1,092	365	9,178	3,140
184	Diospyros Virginiana.....	78.32	782	375	8,045	5,192	240	Juglans rupestris.....	64.89	727	256	6,997	2,909
STYRACEÆ.							241	Carya oliviformis.....	70.99	666	247	6,961	3,714
186	Symplocos tinctoria.....	52.88	622	264	6,146	2,987	242	Carya alba.....	88.11	1,390	512	10,007	4,344
187	Halesia diptera.....	56.81	683	366	6,940	3,153	243	Carya sulcata.....	80.35	1,039	464	8,939	4,609
OLEACEÆ.							244	Carya tomentosa.....	81.29	1,159	482	9,485	4,429
191	Fraxinus pistaciifolia.....	67.68	601	266	6,158	3,368	245	Carya porcina.....	81.36	1,014	466	9,232	4,822
192	Fraxinus Americana.....	65.16	1,015	367	7,585	2,745	246	Carya amara.....	74.74	1,030	470	8,357	3,878
192	Fraxinus Americana, var. Texensis.....	75.83	1,062	480	8,664	3,177	247	Carya myristiciformis.....	79.31	1,465	595	10,206	5,042
193	Fraxinus pubescens.....	62.35	812	371	6,990	3,272	248	Carya aquatica.....	73.13	1,013	376	7,776	4,397
194	Fraxinus viridis.....	70.71	903	382	7,711	3,521	MYRICACEÆ.						
195	Fraxinus platycarpa.....	85.16	476	229	4,014	2,209	249	Myrica cerifera.....	56.08	888	348	7,122	2,304
196	Fraxinus quadrangulata.....	74.50	774	346	7,980	3,322	250	Myrica Californica.....	66.81	903	442	8,516	3,017
197	Fraxinus Oregana.....	57.12	848	284	8,320	2,653	CUPULIFERÆ.						
198	Fraxinus sambucifolia.....	62.72	872	345	6,766	3,106	251	Quercus alba.....	74.39	971	366	8,183	3,388
199	Forestiera acuminata.....	63.00	708	306	6,418	2,717	252	Quercus lobata.....	73.87	717	369	6,793	3,014
201	Oemanthus Americanus.....	80.74	1,231	449	8,966	4,206	253	Quercus Garryana.....	74.24	811	375	7,957	3,846
BORRAGINACEÆ.							254	Quercus obtusiloba.....	83.01	833	372	7,790	4,415
204	Bourreria Havanensis.....	78.48	996	403	9,197	4,702	255	Quercus undulata, var. Gambellii.....	85.38	571	290	6,668	4,072
205	Ehretia elliptica.....	63.56	397	308	6,192	3,663	256	Quercus macrocarpa.....	74.06	929	419	7,843	3,730
							257	Quercus lyrata.....	82.59	1,234	438	7,864	4,033
							258	Quercus bicolor.....	76.18	906	388	7,850	3,534

TABLE OF AVERAGES—Continued.

Catalogue number.	Species.	Approximate relative fuel value.	Coefficient of elasticity, kilograms on millimeters.	Ultimate transverse strength, in kilograms.	Ultimate resistance to longitudinal crushing, in kilograms.	Resistance to indentation to 1.27 millimeters, in kilograms.	Catalogue number.	Species.	Approximate relative fuel value.	Coefficient of elasticity, kilograms on millimeters.	Ultimate transverse strength, in kilograms.	Ultimate resistance to longitudinal crushing, in kilograms.	Resistance to indentation to 1.27 millimeters, in kilograms.
250	Quercus Michauxii	80.03	964	477	7,715	3,725	324	Populus monilifera	38.53	994	328	5,651	1,327
260	Quercus Prinus	74.42	1,255	440	8,615	3,698	325	Populus Fremontii	48.77	1,051	298	6,055	1,382
261	Quercus prinoides	86.09	1,125	528	9,204	4,224	326	Populus Fremontii, var. Wislizeni	45.69	843	295	5,950	1,607
262	Quercus Douglasii	88.53	771	424	8,913	5,988	CONIFERÆ.						
263	Quercus oblongifolia	97.60	857	307	6,941	7,081	326	Libocedrus decurrens	40.14	847	291	7,446	1,561
264	Quercus grisea	99.10	740	400	7,686	5,829	327	Thuja occidentalis	31.58	533	219	4,903	957
266	Quercus Durandii	91.00	837	424	8,550	4,922	328	Thuja gigantea	37.90	1,034	319	7,197	1,114
267	Quercus virens	93.93	1,186	434	8,748	5,185	329	Chamaecyparis sphaeroides	33.12	404	194	4,149	1,074
268	Quercus chrysolepis	84.43	1,198	541	8,721	5,079	330	Chamaecyparis Nutkaensis	47.06	1,029	342	7,281	1,618
269	Quercus Emoryi	90.44	638	300	6,759	6,646	331	Chamaecyparis Lawsoniana	46.16	1,217	379	7,454	1,317
270	Quercus agrifolia	81.47	953	399	7,416	3,770	332	Cupressus Goveniana	46.68	499	230	5,742	2,832
271	Quercus Wislizeni	77.75	861	349	8,527	4,362	333	Juniperus occidentalis, var. conferta	63.75	734	200	8,505	4,464
272	Quercus rubra	65.28	1,137	423	8,172	2,825	339	Juniperus Virginiana	49.11	670	316	6,750	2,376
273	Quercus rubra, var. Texana	90.08	1,033	437	9,310	4,685	340	Taxodium distichum	45.24	1,032	291	6,771	1,166
278	Quercus coccinea	73.91	1,055	450	8,074	3,224	341	Sequoia gigantea	28.67	451	196	6,210	1,091
274	Quercus tinctoria	70.10	1,034	444	8,012	3,243	342	Sequoia sempervirens	42.03	676	255	6,656	1,242
275	Quercus Kelloggii	64.18	745	328	7,184	2,783	343	Taxus brevifolia	63.78	761	400	7,734	4,223
276	Quercus nigra	72.39	977	445	7,954	4,582	345	Torreya taxifolia	51.08	821	378	7,364	2,523
277	Quercus falcata	69.11	1,402	509	9,532	3,223	346	Torreya Californica	46.06	401	249	5,625	1,962
278	Quercus Catesbeii	72.31	1,685	447	7,316	3,646	347	Pinus Strobus	33.47	851	267	6,219	1,194
279	Quercus palustris	68.82	1,123	465	7,862	3,040	348	Pinus monticola	33.99	950	260	5,349	1,071
280	Quercus aquatica	72.07	1,227	449	8,023	3,169	349	Pinus Lambertiana	36.76	794	255	5,382	1,244
281	Quercus laurifolia	76.10	1,259	504	8,424	4,056	350	Pinus flexilis	43.42	676	266	5,561	1,727
282	Quercus heterophylla	68.22	1,225	458	6,600	2,068	351	Pinus albicaulis	41.54	512	249	5,296	1,716
283	Quercus cinerea	63.47	751	424	7,167	3,221	352	Pinus reflexa	48.65	913	329	7,825	2,002
284	Quercus hypoleuca	78.41	944	475	4,695	4,348	353	Pinus Parryana	58.44	378	183	5,420	3,126
285	Quercus imbricaria	74.97	1,193	520	8,839	3,623	355	Pinus edulis	63.49	421	191	5,579	3,368
286	Quercus Phellos	74.35	784	423	6,236	3,452	356	Pinus monophylla	56.20	435	123	4,389	2,718
287	Quercus densiflora	67.25	964	404	7,609	3,598	357	Pinus Balfouriana	54.17	594	181	5,368	2,350
288	Castanopsis chrysophylla	55.55	1,012	316	6,959	1,912	358	Pinus Balfouriana, var. aristata	55.56	715	279	5,209	2,140
289	Castanea pumila	58.80	1,141	423	7,923	1,887	359	Pinus resinosa	48.41	1,132	341	7,274	3,353
290	Castanea vulgaris, var. Americana	44.95	856	297	6,106	1,068	360	Pinus Torreyana	50.62	542	323	4,548	2,309
291	Fagus ferruginea	68.48	1,210	490	7,550	3,145	361	Pinus Arizonica	50.28	824	279	6,292	1,740
292	Ostrya Virginica	82.42	1,373	484	8,669	3,696	362	Pinus ponderosa	46.99	887	307	6,037	1,719
293	Carpinus Caroliniana	72.26	1,149	490	7,969	3,405	363	Pinus Jeffreyi	50.53	925	318	6,679	1,850
BETULACEÆ.							364	Pinus Chihuahuana	54.37	726	355	5,396	2,470
294	Betula alba, var. populifolia	57.43	780	332	5,564	2,073	365	Pinus contorta	58.04	1,585	423	8,068	2,382
295	Betula papyrifera	59.40	1,306	454	7,781	2,083	366	Pinus Murrayana	40.83	771	241	5,328	1,379
296	Betula occidentalis	60.12	924	344	6,260	2,459	367	Pinus Sabiniana	48.18	585	333	5,367	2,202
297	Betula lutea	65.34	1,618	533	9,907	2,581	368	Pinus Coulteri	41.18	1,141	325	5,874	1,475
298	Betula nigra	57.42	1,113	415	7,007	2,117	369	Pinus insignis	45.60	979	316	6,680	1,687
299	Betula lenta	75.97	1,432	519	9,907	3,615	370	Pinus tuberculata	34.88	429	175	4,207	1,372
301	Alnus rubra	47.98	1,080	346	6,644	1,870	371	Pinus Taeda	54.27	1,128	377	6,834	1,719
302	Alnus rhombifolia	41.14	846	291	5,696	1,257	372	Pinus rigida	51.39	581	316	5,687	2,123
303	Alnus oblongifolia	39.65	769	298	4,452	1,189	373	Pinus serotina	79.29	1,170	497	8,079	4,740
SALICACEÆ.							374	Pinus inops	52.93	543	281	5,765	2,496
307	Salix amygdaloides	44.68	501	235	4,224	1,294	375	Pinus clausa	55.09	543	214	6,028	2,100
308	Salix laevigata	48.44	488	275	5,114	1,894	376	Pinus pungens	49.22	803	310	5,670	1,842
309	Salix lasiandra, var. lancifolia	45.73	305	200	4,581	1,311	377	Pinus muricata	49.29	1,194	441	8,142	1,950
309	Salix lasiandra, var. Fendleriana	45.12	879	288	5,457	1,400	378	Pinus mitis	60.96	1,375	443	7,628	2,064
313	Salix flavescens	53.91	1,263	388	7,484	2,019	379	Pinus glabra	39.13	448	212	4,604	1,694
313	Salix flavescens, var. Scouleriana	49.39	1,085	345	6,582	1,581	380	Pinus Banksiana	47.50	942	278	6,329	1,609
316	Salix lasiolepis	55.32	888	347	6,169	2,241	381	Pinus palustris	69.82	1,488	490	10,074	2,508
318	Populus tremuloides	40.10	814	289	5,285	1,281	382	Pinus Cubensis	74.83	1,577	500	10,626	2,985
319	Populus grandidentata	46.11	963	308	5,727	994	383	Picea nigra	45.71	1,100	318	6,520	1,340
320	Populus heterophylla	40.57	723	274	4,527	1,384	384	Picea alba	40.38	1,023	319	5,489	1,117
321	Populus balsamifera	36.11	857	235	5,126	1,202	385	Picea Engelmanni	34.88	808	245	4,271	1,217
321	Populus balsamifera, var. candicans	41.42	730	260	4,418	1,030	386	Picea pungens	37.26	553	194	4,128	1,267
322	Populus angustifolia	38.81	458	171	4,382	1,225	387	Picea Sitchensis	42.60	990	277	5,663	1,160
323	Populus trichocarpa	37.66	1,117	284	6,248	1,018	388	Teuga Canadensis	42.20	900	307	6,142	1,314
							389	Teuga Caroliniana	42.58	713	197	6,450	1,996
								Teuga Morteniana	51.61	1,375	388	8,747	1,622

TABLE OF AVERAGES—Continued.

Catalogue number.	Species.	Approximate relative fuel value.	Coefficient of elasticity, kilograms on millimeters.	Ultimate transverse strength in kilograms.	Ultimate resistance to longitudinal crushing, in kilograms.	Resistance to indentation to 1.27 millimeters in kilograms.	Catalogue number.	Species.	Approximate relative fuel value.	Coefficient of elasticity, kilograms on millimeters.	Ultimate transverse strength in kilograms.	Ultimate resistance to longitudinal crushing in kilograms.	Resistance to indentation to 1.27 millimeters in kilograms.
390	<i>Tsuga Pattoniana</i>	44.35	775	307	6,074	1,664	398	<i>Abies amabilis</i>	42.18	1260	338	7,480	1,029
391	<i>Pseudotsuga Douglasii</i>	51.58	1283	376	8,289	1,608	399	<i>Abies nobilis</i>	45.46	1277	368	7,256	1,017
391	<i>Pseudotsuga Douglasii</i> , var. <i>macrocarpa</i>	45.59	1050	361	7,405	1,642	400	<i>Abies magnifica</i>	46.87	662	299	6,963	1,545
							401	<i>Larix Americana</i>	62.16	1261	384	8,763	1,675
392	<i>Abies Fraseri</i>	35.46	972	273	5,557	1,048	402	<i>Larix occidentalis</i>	74.00	1658	524	11,023	2,395
393	<i>Abies balsamea</i>	33.02	819	220	5,851	1,202							
394	<i>Abies subalpina</i>	34.61	762	202	4,829	1,015							
395	<i>Abies grandis</i>	35.08	958	211	6,255	810							
396	<i>Abies concolor</i>	36.07	909	300	6,237	1,248	405	<i>Washingtonia filifera</i>	50.75	563	183	3,633	2,550

The following table illustrates the relation between the specific gravity and the transverse strength of the wood of species upon which a sufficient number of tests has been made to render such a comparison valuable. The determinations of the specific gravity and transverse strength were, in every case, made upon the same specimen, at the same time. The table is arranged according to the specific gravity of the specimens.

It will be noticed that the strength of the different specimens closely but not invariably follows their specific gravity. An examination of Table III will show, however, that in nearly every case where any wide difference occurs it is due to imperfections in the stick disproportionately affecting its strength. Moreover, in the case of species where the specific gravity and strength of different specimens are nearly identical, their order of arrangement becomes largely accidental. A slight difference in the time occupied in the strength tests, or slight variations in the direction of the grain of the wood, may considerably affect the sequence in such a table:

TABLE ILLUSTRATING THE RELATION BETWEEN TRANSVERSE STRENGTH AND SPECIFIC GRAVITY IN THE WOOD OF CERTAIN SPECIES.

Catalogue number.	Species.	Office number.	Relative specific gravity.	Relative transverse strength.	Catalogue number.	Species.	Office number.	Relative specific gravity.	Relative transverse strength.
8	<i>Magnolia acuminata</i>	534	1	1	66	<i>Acer rubrum</i>	20	1	8
		534	2	2			878	2	3
		246	3	3			878	3	1
		261	4	5			1048	4	2
		246	5	4			1048	5	4
		261	6	6			580	6	5
8	<i>Liriodendron Tulipifera</i>	818	1	1			580	7	6
		818	2	5			743	8	9
		1231	3	2			743	9	7
		1236	4	6	77	<i>Robinia Pseudacacia</i>	1248	1	1
		1236	5	4			400	2	2
		1232	6	3			1248	3	5
		1232	7	8			1247	4	3
		395	8	7			1247	5	4
60	<i>Acer macrophyllum</i>	982	1	1			815	6	6
		982	2	2			815	7	7
		1023	3	4	108	<i>Prunus serotina</i>	15	1	1
		1023	4	8			15	2	2
64	<i>Acer saccharinum</i>	290	1	2			127	3	4
		1233	2	1			1053	4	5
		1235	3	7			1053	5	6
		1234	4	6			127	6	3
		1235	5	5			768	7	8
		1234	6	4			768	8	7
		1233	7	3			368	9	9
		876	8	8			317	10	10
		409	9	9			317	11	11
64	<i>Acer saccharinum</i> , var. <i>nigrum</i>	274	1	1			1154	12	13
		213	2	2			408	13	12
		757	3	3	117	<i>Pyrus coronaria</i>	1068	1	1
		757	4	4			7108	2	3

FOREST TREES OF NORTH AMERICA.

RELATION BETWEEN TRANSVERSE STRENGTH AND SPECIFIC GRAVITY, ETC.—Continued.

Catalogue number.	Species.	Office number.	Relative specific gravity.	Relative transverse strength.	Catalogue number.	Species.	Office number.	Relative specific gravity.	Relative transverse strength.
117	<i>Pyrus coronaria</i> —continued	1088	3	2	194	<i>Fraxinus viridis</i> —continued	948	2	4
		1087	4	4			957	3	1
129	<i>Liquidambar Styraciflua</i>	1178	1	2			438	4	3
		1182	2	9			57	5	7
		1182	3	5			937	6	5
		1183	4	8			308	7	6
		1173	5	1			308	8	3
		1095	6	3	196	<i>Fraxinus quadrangulata</i>	96	1	3
		1181	7	11			96	2	2
		1181	8	10			296 ¹	3	1
		546	9	6			518	4	5
		1095	10	4			296 ²	5	6
		546	11	12			291	6	4
		1183	12	7			125	7	8
151	<i>Cornus florida</i>	1077	3	1	217	<i>Sassafras officinale</i>	125	8	7
		1077	4	3			814	1	6
		1092	5	8			814	2	4
		812	6	4			71	3	1
		812	7	5			854	4	7
		761	8	7			854	5	8
		97	9	6			446	6	5
		97	10	4			387	7	3
154	<i>Nyssa sylvatica</i>	750	1	2			71	8	2
		835	2	9			387	9	9
		750	3	1	223	<i>Ulmus fulva</i>	134	1	1
		833	4	6			134	2	2
		833	5	4	264	<i>Ulmus Americana</i>	583	1	6
		834	6	7			583	2	4
		834	7	3			1049	3	2
		813	8	5			19	4	1
		813	9	8			19	5	3
155	<i>Nyssa uniflora</i>	128	1	2			1036	6	11
		128	2	1			1036	7	10
		604	3	6			958	8	7
		604	4	4			261	9	8
		550	5	5			281	10	9
		550	6	3			958	11	5
184	<i>Diospyros Virginiana</i>	425	1	1	225	<i>Ulmus racemosa</i>	116	1	1
		1084	2	6			314	2	2
		1102	3	4			314	3	5
		811	4	3			116 ²	4	3
		1084	5	2			423	5	7
		811	6	5			116 ³	6	4
		61	7	8			116	7	6
		61	8	7	226	<i>Celtis occidentalis</i>	873	1	5
192	<i>Fraxinus Americana</i>	1045	1	1			873	2	4
		1045	2	4			1111	3	1
		114 ³	3	3			306	4	6
		987	4	19			306	5	7
		227 ²	5	2			1111	6	3
		130	6	7			75	7	3
		431	7	6			75	8	3
		114 ⁴	8	5	232	<i>Morus rubra</i>	132	1	1
		227 ¹	9	10			1255	2	4
		39 ²	10	8			132	3	2
		212	11	11			1255	4	6
		212	12	9			1244	5	3
		747	13	21			1245	6	5
		551	14	14			1246	7	7
		227	15	13	238	<i>Juglans cinerea</i>	1057	1	1
		207	16	12			76 ²	2	2
		747	17	17			16	3	7
		551	18	15			16	4	4
		114 ⁵	19	16			76	5	5
		114	20	18			76	6	3
		114	21	20			123	7	6
194	<i>Fraxinus viridis</i>	949	1	2			303	8	3

Catalogue number.	Species.	Office number.	Relative specific gravity.	Relative transverse strength.	Catalogue number.	Species.	Office number.	Relative specific gravity.	Relative transverse strength.
239	Juglans nigra.....	951	1	4	251	Quercus alba—continued	49 ¹	19	15
		951	2	3			259 ²	20	21
		706	3	1			49 ³	21	17
		706	4	6			49 ⁴	23	24
		318	5	9			8	23	25
		325	6	8			403	24	28
		407	7	5			805	25	29
		117	8	2			251	26	4
		117	9	7			118 ³	27	31
242	Carya alba.....	29 ⁴	1	4			403	28	33
		539	2	7			251	29	26
		531	3	2			32 ⁵	30	20
		531	4	3			118 ⁶	31	34
		1056	5	12			49	32	22
		1056	6	13			443	33	35
		3	7	6			82 ⁷	34	23
		1697	8	1	253	Quercus Garryana.....	118	35	31
		249	9	11			985	1	1
		118	10	14			1027	2	6
		249	11	10			985	3	3
		118	12	16			988	4	2
		3	13	5			1027	5	7
		816	14	9			1029	6	4
		539	15	8			988	7	5
		816	16	15	254	Quercus obtusiloba.....	1029	8	8
243	Carya sulcata.....	338	1	3			771	1	5
		1083	2	5			771	2	7
		391	3	1			256	3	3
		1166	4	6			151	4	1
		391	5	4			151	5	2
		1083	6	7			351	6	4
		1164	7	2	256	Quercus macrocarpa.....	351	7	4
		1170	8	8			137	1	1
		1165	9	9			310	2	2
245	Carya porcina.....	88	1	1			1071	3	5
		88	2	2			310	4	10
		1168	3	7			143	5	13
		1168	4	4			983	6	12
		442	5	9			1073	7	6
		538	6	3			1073	8	7
		6	7	5			983	9	11
		6	8	6			79	10	3
		121	9	8			432	11	9
246	Carya aquatica.....	740	1	1	257	Quercus lyrata.....	831	12	4
		740	2	5			79	13	2
		363	3	3			545	1	1
		363	4	4			762	2	2
		129	5	2			762	3	3
		917	6	6			545	4	4
251	Quercus alba.....	1257	1	18	258	Quercus bicolor.....	54	1	1
		749	2	27			846	2	5
		547	3	1			846	3	4
		547	4	11			54 ³	4	2
		1257	5	14	259	Quercus Michauxii.....	54	5	3
		8	6	19			755	1	3
		1050	7	10			755	2	1
		749	8	8			240	3	5
		259	9	9			240	4	6
		238	10	5			524	5	4
		743	11	3	260	Quercus Prinus.....	524	6	2

FOREST TREES OF NORTH AMERICA.

RELATION BETWEEN TRANSVERSE STRENGTH AND SPECIFIC GRAVITY, ETC.—Continued.

Catalogue number.	Species.	Office number.	Relative specific gravity.	Relative transverse strength.	Catalogue number.	Species.	Office number.	Relative specific gravity.	Relative transverse strength.
261	<i>Quercus prinoides</i> —continued.....	514	3	3	291	<i>Fagus ferruginea</i> —continued.....	765	7	9
		34 ³	4	1			765	8	7
		514	5	6			443	9	10
		34	6	5			44 ³	10	8
267	<i>Quercus virens</i>	404	1	3	295	<i>Betula papyrifera</i>	886	1	1
		954	2	7			1065	2	2
		954	3	6			1065	3	4
		799	4	2			1067	4	3
		919	5	4			723	5	5
		799	6	1			990	6	9
		919	7	5			1066	7	6
273	<i>Quercus rubra</i>	1043	1	1			1066	8	7
		146	2	10			890	9	11
		1043	3	4			1067	10	10
		217	4	6			723	11	8
		215	5	7	297	<i>Betula lutea</i>	843	1	3
		218	6	3			843	2	8
		7	7	2			1068	4	7
		7	8	5			1069	5	6
		149	9	8			1070	3	4
		215	10	11			1068	6	5
		92	11	13			1069	7	1
		45 ²	12	9			1070	8	3
		141	13	15	298	<i>Betula nigra</i>	843	1	5
		920	14	6			841	2	4
		920	15	7			841	3	2
		45 ¹	16	12			843	4	6
		45 ²	17	14			136	5	1
274	<i>Quercus tinctoria</i>	74	1	4			136	6	3
		36 ²	2	9	301	<i>Alnus rubra</i>	991	1	4
		36 ³	3	2			991	2	3
		86	4	6			1025	3	1
		36	5	8			997	4	5
		247	6	1			997	5	6
		17	7	5			1025	6	3
		247	8	3	324	<i>Populus monilifera</i>	809	1	3
		437	9	10			754	2	2
		17	10	7			754	3	1
		244	11	11			809	1	3
		444	12	12			804	5	5
277	<i>Quercus falcata</i>	548	1	3			804	6	6
		265	2	1	325	<i>Populus Fremontii</i> , var. <i>Wicksoni</i>	912	1	5
		548	3	2			646	2	2
		181	4	5			646	3	4
		181	5	7			909	4	1
		265	6	4			912	5	6
		245	7	6			909	6	3
		245	8	8	327	<i>Thuja occidentalis</i>	1099	1	2
280	<i>Quercus aquatica</i>	743	1	1			874	2	1
		511	2	8			874	3	5
		743	3	9			783	4	7
		511	4	2			1099	5	3
		349	5	4			783	6	8
		349	6	3			783	7	10
		264 ¹	7	6			790	8	4
		264 ²	8	5			790	9	6
		264 ³	9	7			379	10	9
280	<i>Castanea vulgaris</i> , var. <i>Americana</i>	18	1	2			379	11	11
		258	2	1			793	12	12
		516	3	3			793	13	13
		18	4	4	329	<i>Chamaecyparis sphaeroides</i>	850	1	7
291	<i>Fagus ferruginea</i>	119	1	2			850	2	3
		9	2	1			850	3	1
		853	3	4			851	4	4
		9	4	3			851	5	2
		853	5	6			853	6	5
		119	6	5			850	7	6

RELATION BETWEEN TRANSVERSE STRENGTH AND SPECIFIC GRAVITY, ETC.—Continued.

Catalogue number.	Species.	Office number.	Relative specific gravity.	Relative transverse strength.	Catalogue number.	Species.	Office number.	Relative specific gravity.	Relative transverse strength.
230	<i>Chamaecyparis Nutkaensis</i>	989	1	5	378	<i>Pinus glabra</i> —continued.....	764	2	2
		983	2	1			142	2	4
		1000	3	8			142	4	2
		983	4	3			544	5	6
		1000	5	6			544	6	5
		994	6	4	379	<i>Pinus Banksiana</i>	780	1	1
		994	7	2			879	2	2
		989	8	7			894	3	3
239	<i>Juniperus Virginiana</i>	1249	1	5			894	4	4
		1250	2	7			879	5	6
		784	3	1			780	6	5
		800	4	4	380	<i>Pinus palustris</i>	81	1	1
		800	5	3			358	2	7
		827	6	2			559	3	2
		827	7	6			358	4	3
242	<i>Sequoia sempervirens</i>	711	1	5			357	5	1
		710	2	1			81	6	10
		711	3	3			359	7	11
		713	4	6			360	8	4
		713	5	2			361	9	3
		712	6	4			361	10	5
		712	7	7			360	11	12
247	<i>Pinus Strobus</i>	222	1	2			243	12	9
		1044	2	7			243	13	6
		797	3	1			385	14	13
		1	4	4			390	15	15
		788	5	3			387	16	12
		788	6	5			85	17	19
		797	7	6			384	18	16
		1044	8	11			390	19	17
		789	9	8			85	20	20
		789	10	9			384	21	21
		777	11	10	381	<i>Pinus Oubensis</i>	498	1	6
258	<i>Pinus resinosa</i>	315	1	1			498	2	4
		315	2	2			356	3	2
		1076	3	3			356	4	1
		1076	4	5			84	5	3
		785	5	4			84	6	5
		1074	6	6	382	<i>Picea nigra</i>	231	1	1
		1075	7	7			231	2	2
		785	8	8			776	3	3
261	<i>Pinus ponderosa</i>	632	1	3			880	4	4
		910	2	10			776	5	9
		907	3	11			373	6	7
		699	4	1			880	7	3
		619	5	4			794	8	5
		630	6	9			794	9	6
		731	7	6	383	<i>Picea alba</i>	513	1	6
		626	8	2			773	2	3
		718	9	5			773	3	1
		630	10	12			784	4	5
		718	11	8			513	5	4
		636	12	7			784	6	8
270	<i>Pinus Taeda</i>	82	1	1			791	7	2
		355	2	4	385	<i>Picea Sitchenensis</i>	970	1	3
		355	3	6			970	2	2
		388	4	2			1015	3	1
		389	5	3			977	4	4
		388	6	7			1026	5	8
		389	7	5			977	6	5
273	<i>Pinus inops</i>	1172	1	4			1026	7	9
		1172	2	2			1019	8	6
		1169	3	6			1019	9	7
		1169	4	1	387	<i>Tsuga Canadensis</i>	790	1	6
		622	5	5			793	2	2
		621	6	3			772	3	1
378	<i>Pinus glabra</i>	764	1	1			772	4	3

RELATION BETWEEN TRANSVERSE STRENGTH AND SPECIFIC GRAVITY, ETC.—Continued.

Catalogue number.	Species.	Office number.	Relative specific gravity.	Relative transverse strength.	Catalogue number.	Species.	Office number.	Relative specific gravity.	Relative transverse strength.
387	<i>Tsuga Canadensis</i> —continued.....	1040	5	4	391	<i>Pseudotsuga Douglasii</i> —continued.....	1022	5	26
		1040	6	5			1008	20	23
		775	7	18			271 ¹	27	24
		787	8	9			1020	26	1
		787	9	8			1022	29	6
		5	10	10			986	30	13
		219	11	7			1020	31	4
		775	12	15			702	32	14
		1042	13	11			709	33	23
		1042	14	12			709	34	27
		5	15	17	394	<i>Abies subalpina</i>	449	1	3
		817	16	14			449 ²	2	6
		219	17	16			449 ¹	3	5
		817	18	18			449 ¹	4	1
		778	19	20			820	5	3
		778	20	19			820	6	4
391	<i>Pseudotsuga Douglasii</i>	708	1	1	396	<i>Abies concolor</i>	733	1	1
		708	2	2			639	2	3
		704	3	3			733	3	2
		1018	4	8			639	4	4
		1018	5	10			529	5	6
		1016	6	33			529	6	5
		989	7	7	401	<i>Larix Americana</i>	226 ²	1	1
		1016	8	18			226	2	3
		705	9	14			774	3	10
		627	10	5			840	4	8
		881	11	9			796	5	2
		881	12	13			795	6	4
		1008	13	19			774	7	5
		706	14	11			795	8	7
		720	15	15			840	9	9
		1011	16	17			781	10	12
		627	17	12			786	11	6
		720	18	30			781	12	11
		1011	19	32	402	<i>Larix occidentalis</i>	1006	1	3
		974	20	16			1006	2	4
		732	21	21			984	3	1
		732	22	29			984	4	2
		973	23	25			719	5	5
		973	24	20			719	6	6

GENERAL REMARKS.

An examination of the results obtained from the various tests made upon the woods of North America indicate at least the important fact that within the limits of any species the weight and strength of any specimen of wood depends upon the actual proportion of the space occupied in the layers of annual growth with open ducts to the space occupied with compact, woody tissue, and to the size of these ducts; or in the case of the wood of Coniferae, the proportion of space occupied with cells formed early in the season to that occupied with the smaller cells of the summer growth. The proportion between these two kinds of growth varies not only in every individual tree, but in different parts of the same tree. The causes which thus affect the growth of wood are not very apparent. It is not soil, nor age, nor general climatic conditions, it appears, which produce the different proportion between the solid and the light portions of the annual growth in any species, because in the same individual this proportion is found to vary from year to year. It varies very irregularly; nor does the rapidity of growth, as has been supposed, greatly affect the strength of wood, because the proportion of open to compact growth is little affected by rapid or slow increase of the tree's diameter. How far annual climatic variations affect the nature of the annual layers of growth has not been demonstrated, although it is not impossible that in years in which conditions favorable to rapid growth are extended late into the season, the proportion of the annual layer occupied by open, weak growth to the growth of the whole year would be greater than that formed in a year during which the season favorable for rapid growth was less extended.

It follows that while such experiments as those conducted by Mr. Sharples are necessary to establish maximum and relative values for any species, these being established, actual values of any given specimen of

wood may be determined by microscopic examination of its structure; that is, two specimens of the wood of any species to which the census tests have been applied being given, their relative values can be determined by an examination of their structure as well as or better than by any elaborate experiments.

TANNIN VALUES.

The amount of tannin contained in the bark of various trees of the United States has been determined.

These determinations give the proportion of tannin. They do not indicate the real value of the bark of the species for tanning, which can only be obtained by actual experiments made on a large scale, other properties in the bark, beside the percentage of tannin, affecting the value of the leather prepared with it.

These determinations must therefore be regarded as approximations, which will serve, in some cases, to indicate species not now in general use for this purpose, which may be looked to as possible sources of tannin supply.

The methods adopted by Mr. Sharples in making these determinations are described by him as follows:

The tannin in each case was determined in the roased bark; that is, bark deprived of the main part of the outside coating. The method employed was that devised by Lowenthal, which may be thus briefly described: A standard decoction of the bark is titrated with permanganate of potash, a quantity of indigo being first added to it. In a second portion the tannin is precipitated by means of gelatine, and the gallic acid in the liquid again determined by permanganate and indigo. The difference between these two readings gives the amount of tannin in the bark, the value of the permanganate having previously been determined by pure tannic acid, or by oxalic acid and calculation.

The bark of the following species has been examined:

Catalogue number.	Botanical name.	Common name.	Per cent. of tannin.	Per cent. of ash.	Catalogue number.	Botanical name.	Common name.	Per cent. of tannin.	Per cent. of ash.
14	<i>Gordonia Lasianthus</i>	Loblolly Bay. Tan Bay	13.14	2.85	275	<i>Quercus Kelloggii</i>	Black Oak	6.76	2.64
93	<i>Prosopis juliflora</i>	Mesquit. Algaroba. Honey Locust. Honey Pod.	4.04	8.71	276	<i>Quercus nigra</i>	Black Jack. Jack Oak	4.26	6.28
140	<i>Rhizophora Mangle</i>	Mangrove	21.04	6.70	277	<i>Quercus falcata</i>	Spanish Oak. Red Oak	8.59	4.32
160	<i>Exostemma Caribaeum</i>		5.81	7.16	287	<i>Quercus densiflora</i>	Tanbark Oak. Chestnut Oak. Peach Oak.	16.46	2.84
251	<i>Quercus alba</i>	White Oak	5.99	6.11	290	<i>Castanea vulgaris</i> , var. Ameri- cana.	Chestnut	6.25	2.00
256	<i>Quercus macrocarpa</i>	Burr Oak. Mossy-cup Oak. Over-cup Oak.	4.59	8.05	382	<i>Picea nigra</i>	Black Spruce	7.20	2.84
260	<i>Quercus prinus</i>	Chestnut Oak. Rock Chest- nut Oak.	6.25	8.33	384	<i>Picea Engelmanni</i>	White Spruce	20.56	2.75
261	<i>Quercus prinoides</i> (old tree) ...	Yellow Oak. Chestnut Oak. Chinquapin Oak.	4.33	8.38	384	<i>Picea Engelmanni</i>	do	17.01	2.32
261	<i>Quercus prinoides</i> (young tree) ..	do	10.33	6.23	384	<i>Picea Engelmanni</i>	do	12.60	0.75
267	<i>Quercus virens</i>	Live Oak	10.46	8.89	387	<i>Tsuga Canadensis</i>	Hemlock	13.11	1.31
269	<i>Quercus Emoryi</i>	Black Oak	2.76	15.09	389	<i>Tsuga Mertensiana</i>	do	14.42	1.44
272	<i>Quercus rubra</i>	Red Oak. Black Oak	4.56	4.43	389	<i>Tsuga Mertensiana</i>	do	15.87	1.49
274	<i>Quercus tinctoria</i>	Black Oak. Yellow-bark Oak. Quercitron Oak. Yellow Oak.	5.90	5.78	390	<i>Tsuga Pattoniana</i>	do	15.73	2.48
					391	<i>Pseudotsuga Douglasii</i>	Red Fir. Yellow Fir. Ore- gon Pine. Douglas Fir.	12.79	1.56

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
MAGNOLIACEÆ.								
1. <i>Magnolia grandiflora</i> <i>Big Laurel. Bull Bay.</i>	346	Alabama	Cottage Hill	C. Mohr	Rich loam	0.372	116	
2. <i>Magnolia glauca</i> <i>Sweet Bay. White Bay. Beaver Tree. White Laurel. Swamp Laurel.</i>	354	do	do	do	Swampy	0.268	91	20
3. <i>Magnolia acuminata</i> <i>Ocumber Tree. Mountain Magnolia.</i>	246	Virginia	Wytheville	H. Shriver	Clay limestone			
	261 ¹	do	Fancy Gap	do	Rich, light			
	261 ²	do	do	do	do			
	261 ³	do	do	do	do			
	534	Mississippi	Salvers' mill	C. Mohr	do	0.198	28	14
4. <i>Magnolia cordata</i> <i>Ocumber Tree.</i>	1178	Alabama	Winston county	C. Mohr				
5. <i>Magnolia macrophylla</i> <i>Large-leaved Ocumber Tree.</i>	22	North Carolina	Statesville	M. E. Hyams	Rich	0.050	18	
	532	Mississippi	Quitman	C. Mohr	Rich, low			
6. <i>Magnolia Umbrella</i> <i>Umbrella Tree. Elk Wood.</i>	266 ¹	Virginia	Wytheville	H. Shriver		0.072	15	8
	266 ²	do	do	do				
	266 ³	do	do	do				
7. <i>Magnolia Fraseri</i> <i>Long-leaved Ocumber Tree.</i>	260 ¹	do	Fancy Gap	do	Damp	0.065	20	
	260 ²	do	do	do	do			
	260 ³	do	do	do	do			
8. <i>Liriodendron Tulipifera</i> <i>Tulip Tree. Yellow Poplar. White Wood.</i>	138	Michigan	Danville	W. J. Beal	Sandy			
	165	Ohio	D. E. McSherry & Co.	E. E. Barney				
	174	Tennessee	Woodsum Machine Company.	do				
	177	Ohio	Barney & Smith Manufacturing Co.	do				
	178	do	D. E. McSherry & Co.	do				
	187	do	J. W. Stoddard & Co.	do				
	188	do	Barney & Smith Manufacturing Co.	do				
	395	Michigan	Lansing	W. J. Beal				
	818	West Virginia	Grafton	C. G. Pringle				
	1231	Pennsylvania	Chester county	P. P. Sharples				
	1232	do	do	do				
ANONACEÆ.								
9. <i>Asimina triloba</i> <i>Papaw. Oustard Apple.</i>	211	Missouri	Meramec river, Jefferson county.	G. W. Letterman	Alluvial	0.169		
	332	Tennessee	Cumberland river	A. Gittinger	do	0.086		
10. <i>Anona laurifolia</i> <i>Pond Apple.</i>	479	Florida	Bay Biscayne	A. H. Curtiss	Swampy	0.240	47	

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.6040	0.6690		0.6360	0.40	0.65	0.53	39.63	Second sp. gr. determination made on sap-wood	346
0.5034	0.5067		0.5035	0.43	0.53	0.47	31.33		354
0.5012	0.5412		0.5212	0.29	0.26	0.27			246
0.4399			0.4399	0.25		0.25			261 ¹
0.4562			0.4562	0.30		0.30			261 ²
0.4215			0.4215	0.30		0.30			261 ³
0.5065	0.5058		0.5061	0.34	0.33	0.34			534
.....			0.4690			0.29	29.23		
0.4095	0.4124	0.4101	0.4130	0.25	0.39	0.32	25.79	Third sp. gr. determination made on sap-wood; fourth sp. gr. determination, 0.4175.	1178
0.5375	0.4859		0.5117	0.32		0.32		All sap-wood	22
0.5498	0.5584		0.5501	0.45	0.34	0.39			532
.....			0.5309			0.35	33.09		
0.3787			0.3787	0.19		0.19		Growth rapid	260 ¹
0.5067			0.5067	0.18		0.18		Growth rapid; 0.5 sap-wood	260 ²
0.4806			0.4806	0.24		0.24		Growth rapid; all sap-wood	260 ³
.....			0.4487			0.30	27.96		
0.5430			0.5430	0.25		0.25		All sap-wood	260 ¹
0.4976			0.4976	0.27		0.27			260 ²
0.4602			0.4602	0.33		0.33		All sap-wood	260 ³
.....			0.5003			0.28	31.18		
0.3843			0.3843	0.25		0.25			138
0.3831	0.3783		0.3807	0.27	0.27	0.27		Yellow poplar	165
0.3798	0.3787		0.3792	0.20	0.22	0.21		Yellow poplar (soft)	174
0.4475	0.4361		0.4418	0.17	0.15	0.16		Hard poplar	177
0.4512	0.4442		0.4477	0.19	0.19	0.19		Hard poplar	178
0.4362	0.4150		0.4256	0.18	0.21	0.19		Yellow poplar	187
0.4436	0.4551		0.4493	0.25	0.30	0.27			188
0.3774			0.3774	0.33	0.31	0.32			395
0.4768	0.4822		0.4793	0.26	0.29	0.28			818
0.4444	0.4409		0.4427	0.20	0.22	0.21		White poplar	1231
0.4199	0.4712		0.4455	0.19	0.16	0.18		Yellow poplar	1232
.....			0.4230			0.23	26.36		
0.3549	0.3810		0.3679	0.16	0.14	0.15			211
0.4259			0.4259	0.24	0.30	0.27			332
.....			0.3969			0.31	24.74		
0.4812	0.5199	0.5048	0.5053	4.94	4.79	4.86	31.49		479

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
CAPPARIDACEÆ.								
11. <i>Capparis Jamaicensis</i>	477	Florida	Upper Metacombe Key.	A. H. Curtiss	Coral	0.086	24	
	1186	do	do	do	do			
CANELLACEÆ.								
12. <i>Canella alba</i>	490	do	Umbrella Key	do	do	0.020	17	
White Wood. Cinnamon Bark. Wild Cinnamon.	1181	do	Elliott's Key	do	do	0.120	5	14
TERNSTROMIACEÆ.								
14. <i>Gordonia Lasianthus</i>	236	South Carolina....	Bonneau's Depot ...	H. W. Ravenel	Wet pine-barren			
Loblolly Bay. Ten Bay.	414	do	Aiken	do	Swampy	0.330	22	25
STERCULIACEÆ.								
16. <i>Fremontia Californica</i>	1230	California	San Bernardino mountains.	C. G. Pringle				
Slippery Elm.								
TILIACEÆ.								
17. <i>Tilia Americana</i>	2	Massachusetts	Arnold Arboretum..	C. S. Sargent	Drift	0.470	9	26
Lime Trees. Bass Wood. American Linden. Lin. Bee Tree.	124	Michigan	Big Rapids	W. J. Beal	Gravelly			
	262	Missouri	Allenton	G. W. Letterman	Alluvial	0.270	3	73
	816	Michigan	Hersey	W. J. Beal	Rich loam			
	1089	Massachusetts	Danvers	J. Robinson	Moist gravel	0.330	2	35
17. <i>Tilia Americana</i> , var. <i>pubescens</i>	745	Georgia	Bainbridge	A. H. Curtiss	Low	0.300	23	
18. <i>Tilia heterophylla</i>	285 ¹	Kentucky	Cliffs Kentucky river	W. M. Linney	Limestone			
White Bass Wood. Wahoo.	285 ²	do	Mercer county	do	do			
	285 ³	do	do	do	do			
	320	Tennessee	Cumberland river ..	A. Gittinger	Alluvial			
MALPIGHIACEÆ.								
19. <i>Byrsonima lucida</i>	510	Florida	Boca Chica Key	A. H. Curtiss	Coral	0.080	21	
Yellowberry. Glanberry.	1118	do	No-Name Key	do	do			
	1117	do	Boca Chica Key	do	do	0.074	19	12
	1190	do	No-Name Key	do	do			
ZYGOPHYLLACEÆ.								
20. <i>Guaicum sanctum</i>	476	do	Upper Metacombe Key.	do	do	0.178	18	55
<i>Lignum-vita</i> .	898	do		Department of Agriculture.				
	1183	do	Elliott's Key	A. H. Curtiss	Coral			
21. <i>Porliera angustifolia</i>	947	Texas	San Antonio	C. Mohr	Limestone	0.056	31	
RUTACEÆ.								
22. <i>Xanthoxylum Americanum</i>	80	Missouri	Allenton	G. W. Letterman	Alluvial	0.087	5	9
Prickly Ash. Toothache Trees.	392	Michigan	Lansing	W. J. Beal	do	0.086	4	28

THE WOODS OF THE UNITED STATES.

269

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.6690	0.6609		0.6649	5.60	5.77	5.68			477
0.7272	0.7315		0.7293	3.82	3.88	3.85			1186
			0.6971			4.76	43.44		
0.9570	0.9508		0.9539	0.75	0.95	0.85		All sap-wood	439
1.0890	1.0017		1.0203	2.66	2.65	2.66		0.5 sap-wood	1121
			0.9893			1.75	61.65		
0.3852	0.4659		0.4255	0.49	0.44	0.47			236
0.5146	0.5255		0.5201	1.02	1.08	1.05			414
			0.4728			0.76	29.47		
0.6995	0.7288		0.7142	1.51	1.86	1.69	44.51		1230
0.4626	0.4722		0.4674	0.30	0.33	0.31			3
0.3815			0.3815	0.32	0.38	0.35			124
0.5035	0.4378		0.4706	0.97	1.08	1.02			252
0.5166			0.5166	0.55	0.64	0.60		All sap-wood	316
0.4272	0.4163	0.4362	0.4266	0.42	0.48	0.45			1039
			0.4525			0.55	28.20		
0.4103	0.3885		0.4074	0.68	0.67	0.65	25.39	Growth very rapid	745
0.4087	0.4220			0.62	0.64				
0.4927			0.4927	0.45		0.45		Second growth	285
0.3791			0.3791	0.67		0.67			285
0.3674			0.3674	0.86		0.86			285
0.4695	0.4545		0.4620	0.48	0.53	0.50			320
			0.4253			0.62	26.51		
0.5257	0.5300		0.5308	2.84	2.85	2.85			510
0.5909	0.6783	0.6743	0.6479	2.40	1.94	2.17			1113
0.5681	0.5791	0.5875	0.5783	2.75	2.56	2.65			1119
0.6009	0.5964		0.5987	1.94	2.40	2.17			1190
			0.5888			2.46	36.69		
1.1845	1.2180	0.9563	1.1196	0.87	0.92	0.90		0.5 sap-wood	476
1.2736	1.1700		1.2218	0.51	0.81	0.66		Second sp. gr. determination made on 0.5 sap-wood	896
1.0997	1.0996	1.0652	1.0882	0.86	0.94	0.90		0.66 sap-wood	1123
			1.1432			0.82	71.24		
1.1230	1.0843	1.1230	1.1101	0.53	0.49	0.51	69.18	Sap-wood	947
0.6122	0.5969		0.6045	0.60	0.58	0.59			80
0.5235	0.5290		0.5262	0.54	0.55	0.54			302
			0.5654			0.57	35.23		

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
23. <i>Xanthoxylum Clava-Herculis</i> <i>Toothache Tree. Prickly Ash. Sea Ash. Pepper Wood. Wild Orange.</i>	735	Florida	Chattahoochee	A. H. Curtiss	Dry, sandy			
	807	Georgia	Cumberland island	do				
	1086	Texas	Palestine	C. Mohr	Damp, sandy	0.852	20	10
23. <i>Xanthoxylum Clava-Herculis</i> , var. <i>fruticosum</i>	938	do	Austin	do	Dry, calcareous	0.096	33	
24. <i>Xanthoxylum Caribaeum</i> <i>Satin Wood.</i>	1109	Florida	Bahia Honda Key	A. H. Curtiss	Coral			
	1140	do	do	do	do	0.136	4	54
25. <i>Xanthoxylum Pterota</i> <i>Wild Lime.</i>	481	do	Bay Biscayne	A. H. Curtiss	Coral			
	950	Texas	Matagorda bay	C. Mohr	Calcareous			
	1128	Florida	Bay Biscayne	A. H. Curtiss	Coral	0.84	13	31
26. <i>Ptelia trifoliata</i> <i>Hop Tree. Shrubby Trefoil. Wafer Ash.</i>	768	do	Aspalaga	do	Calcareous	0.94	23	
27. <i>Canotia holocantha</i>	1228	Arizona	Wickenburg	C. G. Pringle				
SIMARUBEÆ								
28. <i>Simaruba glauca</i> <i>Paradise Tree.</i>	487	Florida	Bay Biscayne	A. H. Curtiss	Coral			
BURSERACEÆ								
29. <i>Bursera gummifera</i> <i>Gum Elemi. Gumbo Limbo. West Indian Birch.</i>	462	do	Upper Metacombe Key	do	do			
	908	do		Department of Agriculture.				
30. <i>Amyris sylvatica</i> <i>Torah Wood.</i>	475	do	Upper Metacombe Key	A. H. Curtiss	Coral	0.128	61	
MELIACEÆ								
31. <i>Swietenia Mahagoni</i> <i>Mahogany. Madeira.</i>	452	do	do	do	do	0.228	16	81
OLACINÆÆ								
32. <i>Ximenia Americana</i> <i>Wild Lime. Tallow Nut. Hog Plum. Mountain Plum.</i>	472	do	do	do	do	0.112	3	43
	1134	do	Umbrella Key	do	do			
ILICINÆÆ								
33. <i>Ilex opaca</i> <i>American Holly.</i>	280	South Carolina	Waverly Mills	W. St. J. Mayek	Sandy loam	0.144	35	
	902	do		Department of Agriculture.				
34. <i>Ilex Dahoon</i> <i>Dahoon. Dahoon Holly.</i>	484	Florida	Bay Biscayne	A. H. Curtiss	Low, damp	0.128	29	
34. <i>Ilex Dahoon</i> , var. <i>myrtifolia</i>	802	do	Jacksonville	do	do	0.148	28	

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.5139	0.5002		0.5070	0.92	0.91	0.91			7
0.4880	0.4772		0.4826	0.68	0.65	0.67			8
0.5392	0.5153		0.5272	0.90	0.84	0.87			10
			0.5056			0.82	31.51		
0.6129	0.5723	0.6080	0.5967	0.76	0.76	0.76	37.19		8
0.8598	0.8965		0.8782	1.59	2.01	1.80			11
0.9090	0.9275	0.9302	0.9222	2.34	2.15	2.25			11
			0.9002			2.02	56.10		
0.7480	0.7666		0.7573	0.90	1.00	0.95			4
0.5807	0.5775		0.5791	0.68	0.59	0.61			9
0.9000	0.8949	0.8955	0.8968	0.69	0.89	0.79			11
			0.7444			0.78	46.39		
0.8238	0.8400		0.8319	0.84	0.27	0.86	51.84		7
0.6464	0.7305		0.6885	6.13	4.54	5.33	42.91		12
0.4100	0.4172		0.4136	0.93	0.92	0.93	25.78		4
0.2677	0.2488	0.2587	0.2584	2.05	1.91	1.98			4
0.2823	0.4022		0.3428	2.10	2.07	2.09			9
			0.3008			2.04	18.71		
1.0469	1.0450		1.0459	0.66	0.51	0.59	65.18		4
0.7335	0.7464	0.7047	0.7282	1.06	1.11	1.09	45.38		4
0.8799	0.8925		0.8862	0.55	0.69	0.62			4
0.9638	0.9505	0.9249	0.9531	0.84	0.82	0.83		0.5 asp-wood	11
			0.9196			0.73	57.31		
0.6046	0.6324		0.6335	0.71	0.55	0.63			3
0.5524	0.5078		0.5801	1.04	0.71	0.88			9
			0.5818			0.76	36.26		
0.4791	0.4820		0.4806	0.94	0.86	0.91	29.95		4
0.5896	0.5990		0.5878	0.95	0.84	0.90	36.60		8

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
35. <i>Ilex Cassine</i> <i>Cassina. Yaupon. Yupon.</i>	845	Alabama	Cottage Hill	C. Mohr	Sandy	0.076	17	
	804	Florida	Saint John's river ..	A. H. Curtiss	Rich, sandy			
	952	Texas	Matagorda bay	C. Mohr	Light			
36. <i>Ilex decidua</i>	56	Missouri	Allenton	G. W. Letterman	Low, damp			
	835	Texas	Dallas	J. Reverchon	Upland			
	753	Florida	Chattahoochee river ..	A. H. Curtiss	Clay	0.088	23	
	945	Texas	New Braunfels	C. Mohr	Alluvial			
CYRILLACEÆ.								
37. <i>Cyrilla racemiflora</i>	841	Alabama	Chunchula	..do	Damp, sandy	0.195		
	615	Georgia	Ogeechee river	A. H. Curtiss	Low			
38. <i>Cliftonia ligustrina</i>	338	Alabama	Cottage Hill	C. Mohr	Wet	0.194	47	
CELASTRACEÆ.								
39. <i>Eunonymus atropurpureus</i>	63	Missouri	Allenton	G. W. Letterman	Alluvial			
	1078	..do	..do	..do	..do	0.092	22	
40. <i>Myginda pallens</i>	1188	Florida	Umbrella Key	A. H. Curtiss	Calcareous			
41. <i>Schsefferia frutescens</i>	478	..do	Upper Metacombe	..do	Coral	0.110	71	
	1201	..do	Key	..do	..do			
RHAMNACEÆ.								
42. <i>Reynosa latifolia</i>	454	..do	..do	..do	..do	0.112	7	23
43. <i>Condalia ferrea</i>	460	..do	..do	..do	..do	0.105	22	56
44. <i>Condalia obovata</i>	941	Texas	New Braunfels	C. Mohr	Dry, calcareous	0.100	5	36
45. <i>Rhamnus Caroliniana</i>	43	Missouri	Allenton	G. W. Letterman	Limestone			
	521	Tennessee	Nashville	A. Gattinger	..do			
	803	Florida	Saint John's river ..	A. H. Curtiss	Rich hummock	0.108	19	10
	1004	Arkansas	Jonesboro'	T. B. Kitchens		0.059	14	
46. <i>Rhamnus Californica</i>	1256	California	Santa Cruz mount- ains.	C. G. Pringle				
47. <i>Rhamnus Purshiana</i>	993	Oregon	Portland	G. Engelmann and C. S. Sargent.	Rich, alluvial			
48. <i>Ceanothus thyrsiflorus</i>	1101	California	Santa Cruz	C. L. Anderson		0.090	11	

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
1.7257	0.7185		0.7221	0.74	0.70	0.72			845
0.6890	0.7101		0.7041	1.04	1.11	1.08			804
0.7475	0.7619		0.7547	6.89	0.72	0.81			953
			0.7270			0.87	45.31		
0.7076			0.7076	0.44	0.50	0.47			56
0.7888			0.7888	0.83	0.84	0.84			335
0.5910	0.5861		0.5885	0.87	0.79	0.83			753
0.7080	0.7894	0.7089	0.7931	0.67	0.67	0.67			945
			0.7420			0.70	46.35		
0.7147	0.6563		0.6855	0.41	0.43	0.42			341
0.6818	0.6808		0.6713	0.43	0.40	0.41			615
			0.6784			0.42	42.28		
0.6350	0.6147		0.6249	0.35	0.48	0.42	38.95		338
									
0.6240			0.6240	0.70	0.58	0.64			63
0.6868	0.7018		0.6943	0.51	0.53	0.52			1078
			0.6592			0.58	41.08		
0.9057	0.9039		0.9048	3.88	2.95	3.42	50.89		1188
0.7487	0.7728		0.7608	2.79	2.91	2.85			478
0.7912	0.7850		0.7881	2.15	2.28	2.22			1201
			0.7745			2.54	48.27		
1.0805	1.0825		1.0715	3.24	3.15	3.20	66.78		454
1.3020	1.3020		1.3020	8.00	8.61	8.31	81.14		460
1.2049	1.1949		1.1999	7.03	7.02	7.03	74.78	0.125 sap-wood.....	941
0.5954			0.5954	0.19	0.18	0.19			48
0.5940	0.5962		0.5951	0.76	0.96	0.87			521
0.4971	0.5801		0.5186	0.47	0.52	0.50		All sap-wood.....	808
0.4894	0.4720	}	0.4807	1.15	0.81	0.98		0.5 sap-wood.....	1094
0.4789	0.4825								
			0.5462			0.64	34.04		
0.6000			0.6000	0.63	0.53	0.58	37.39		1256
0.5685	0.5678		0.5672	0.74	0.50	0.67	35.35	0.25 sap-wood.....	998
0.5805	0.5804	}	0.5750	0.70	0.68	0.69	35.83	Growth rapid.....	1101
0.5865	0.5625								

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
409. <i>Colubrina reclinata</i>	502	Florida	Umbrella Key	A. H. Curtiss	Coral	0.128	6	45
Naked Wood.	1189	do	do	do	do	0.100	13	24
SAPINDACEÆ.								
30. <i>Æsculus glabra</i>	297	Missouri	Allenton	G. W. Letterman	Rich, moist			
Ohio Buckeye. Fetid Buckeye.	386	do	do	do	Alluvial			
	427	Tennessee	Nashville	A. Gettinger	Rich, moist			
31. <i>Æsculus sava</i>	445	do	do	do	Rich upland			
Sweet Buckeye.								
32. <i>Æsculus Californica</i>	684	California	Marin county	G. R. Vasey	do	0.250		
California Buckeye.								
33. <i>Ungnadia speciosa</i>	944	Texas	New Braunfels	C. Mohr	Limestone	0.124	26	
Spanish Buckeye.								
34. <i>Sapindus marginatus</i>	307	do	Dallas	J. Reverchon	Rich, damp			
Wild China. Soapberry.	589	do	do	do	do			
	824	New Mexico	Rio Gila cañon	E. L. Greene	do			
	928	Texas	Austin	C. Mohr	Limestone	0.222	24	36
35. <i>Sapindus Saponaria</i>	568	Florida	Cape Sable	A. H. Curtiss	Rich, sandy, damp	0.124	41	
Soapberry.	1123	do	Key Largo	do	Coral			
36. <i>Hypelate paniculata</i>	463	do	Upper Metacombe Key	do	do	0.262	15	80
Ink Wood. Iron Wood.								
37. <i>Hypelate trifoliata</i>	464	do	do	do	do	0.224	13	72
White Iron Wood.								
38. <i>Acer Pennsylvanicum</i>	90	Vermont	Huntingdon	C. G. Pringle	Gravelly			
Striped Maple. Moose Wood. Striped Dogwood. Goose-foot Maple. Whistle Wood.	372	do	do	do	do			
39. <i>Acer spicatum</i>	98	do	do	do	do			
Mountain Maple.	371	do	do	do	do			
40. <i>Acer macrophyllum</i>	982	Oregon	Portland	G. Englemann and C. S. Sargent	Rich, alluvial	0.229	30	40
Broad-leaved Maple.	1023	do	Portland Furniture Company	do				
41. <i>Acer circinnatum</i>	982	do	Portland	do	Moist, alluvial			
Vine Maple.	1013	do	do	do	do			
	1014	Washington territory	Wilkeson	do	do			
42. <i>Acer glabrum</i>	526	Colorado	Englemann's cañon	R. Douglas	dry, gravelly	0.048	28	
Dwarf Maple.								

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.8100	0.8516		0.8308	1.29	1.14	1.21			503
0.8046	0.8166	0.8112	0.8108	2.31	2.26	2.29			1129
			0.8208			1.75	51.15		
0.4270	0.4358		0.4314	1.01	1.00	1.01			207
0.4787	0.4625		0.4706	0.83	0.85	0.84			368
0.4606			0.4606	0.77	0.68	0.78			427
			0.4542			0.86	28.81		
0.4241	0.4307		0.4274	0.99	1.00	1.00	27.24		446
0.4921	0.5038		0.4980	0.66	0.73	0.70	81.04		681
0.6392	0.6272		0.6332	1.15	1.19	1.17	89.46		944
0.7520	0.7980		0.7750	1.42	1.27	1.35			307
0.8324	0.8616		0.8470	1.86	1.52	1.69			589
0.7978	0.8345		0.8162	1.44		1.44			824
0.7550	0.8522	0.8296	0.8128	1.54	1.52	1.53			926
			0.8126			1.50	50.64		
0.8595	0.8768		0.8679	3.82	3.30	3.56			563
0.7940	0.8024	0.8204	0.8056	5.00	5.25	5.13			1123
			0.8367			4.84	52.14		
0.9800	0.9405		0.9633	1.23	1.27	1.25	59.41		466
0.9656	0.8648		0.9102	1.35	1.41	1.38	56.72		464
0.5111	0.5119		0.5115	0.32	0.35	0.34			96
0.5564	0.5100	0.5780	0.5484	0.44	0.30	0.37			372
			0.5299			0.36	33.02		
0.5198	0.5205		0.5202	0.42	0.37	0.40		All sap-wood	96
0.5490	0.5528	0.5353	0.5457	0.48	0.41	0.45			371
			0.5330			0.43	33.22		
0.5057	0.5029	0.5254	0.5113	0.59	0.57	0.58			962
0.4323	0.5087		0.4705	0.49	0.49	0.49			1028
			0.4909			0.54	30.59		
0.6824	0.6720		0.6772	0.39	0.39	0.39			968
0.6587	0.6600		0.6624	0.35	0.37	0.36			1013
0.6370	0.6800		0.6585	0.41	0.44	0.43			1014
			0.6680			0.39	41.51		
0.6032	0.6023		0.6028	0.81	0.29	0.30	37.57		526

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
63. <i>Acer grandidentatum</i>	886	Utah	City Creek cañon ..	M. E. Jones	Moist, gravelly ..	0.050	13	
64. <i>Acer saccharinum</i>	105	Vermont		C. G. Pringle	Gravelly			
	203	Ohio	Woodsum Machine Company.	E. E. Barney				
	298	Missouri	Allenton	G. W. Letterman	Rich uplands ..			
	299	do	do	do	do			
	376	Vermont	Charlotte	C. G. Pringle	Gravelly			
	409	New England	Charlestown Navy-yard.	S. H. Pook				
	1233	Vermont	Charlotte	F. H. Horsford				
	1234	do	do	do				
	1235	do	do	do				
64. <i>Acer saccharinum</i> , var. <i>nigrum</i>	213	do	do	C. G. Pringle	Clay			
	274 ¹	Missouri	Allenton	G. W. Letterman	Low, alluvial ..			
	274 ²	do	do	do	Rich, alluvial ..			
	399	do	do	do	Low, alluvial ..			
	440	Tennessee	Nashville	A. Gattinger	Rich			
	757	Florida	Chattahoochee	A. H. Curtiss	Clay	0.268	51	
	1167	Missouri	Allenton	G. W. Letterman	Rich, alluvial ..			
65. <i>Acer dasycarpum</i>	103	Vermont	Charlotte	C. G. Pringle	Swampy			
	367	do	do	do	do			
	448	Missouri	Allenton	G. W. Letterman				
	1052	Massachusetts	Topsfield	J. Robinson	Low meadow ..			
66. <i>Acer rubrum</i>	20	do	Arnold Arboretum ..	C. S. Sargent	Drift	0.285	18	36
	530	Mississippi	Kemper's mill	C. Mohr	Rich, swampy ..	0.222	41	
	743	Georgia	Bainbridge	A. H. Curtiss	Low	0.252	22	54
	878	Massachusetts	Danvers	J. Robinson	do	0.168	30	3
	1048	do	North Reading	do	do	0.204	60	6
66. <i>Acer rubrum</i> , var. <i>Drummondii</i>	1239	Missouri	Poplar Bluff	G. W. Letterman ..				
	1240	do	do	do				
67. <i>Negundo aceroides</i>	290	Missouri	Allenton	G. W. Letterman	Rich bottom ..			
	811	Texas	Dallas	J. Reverchon	do			
68. <i>Negundo Californicum</i>	645	California	Contra Costa county	G. R. Vasey	Rich, moist	0.240	14	
ANACARDIACEÆ.								
69. <i>Rhus cotinoides</i>	1176	Alabama	Huntsville	C. Mohr	Rich loam			

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.7012	0.6791		0.6902	0.66	0.61	0.64	43.01		898
0.6139	0.6003		0.6071	0.44	0.40	0.42			106
0.6363	0.6300		0.6332	0.51	0.45	0.48			203
0.7697	0.8098		0.7898	0.54	0.59	0.57			298
0.7828	0.7680		0.7750	0.60	0.57	0.59			299
0.6538	0.6783		0.6661	0.93	1.10	1.02			376
0.6242			0.6242	0.86	0.44	0.40			400
0.7106			0.7106	0.91	0.33	0.62			1233
0.6893	0.6899		0.6896	0.33	0.29	0.31			1234
0.7239			0.7239	0.48	0.35	0.42			1235
			0.6912			0.54	43.08		
0.7319	0.6935	0.6639	0.6964	1.48	1.01	1.25			213
0.7249	0.6980		0.7115	0.56		0.56			274 ¹
0.7214	0.7002		0.7108	0.52		0.52			274 ²
0.7117	0.6799		0.6958	0.70	0.56	0.63			399
0.6410	0.6429		0.6419	1.31	1.02	1.17			440
0.6868	0.7008	0.7165	0.7014	0.55	0.33	0.44			757
0.6814	0.6840		0.6827	0.42	0.36	0.39			1167
			0.6915			0.71	43.09		
0.4889	0.5254		0.5072	0.31	0.32	0.32			103
0.4890	0.4828		0.4844	0.30	0.31	0.31			367
0.4760	0.5062		0.4911	0.40	0.42	0.41			443
0.6247	0.6163	0.6344	0.6251	0.29	0.27	0.28			1053
			0.5269			0.33	32.84		
0.6761	0.6845		0.6803	0.24	0.25	0.25		All sap-wood	20
0.5770	0.6064		0.5917	0.43	0.42	0.43			530
0.5288	0.5510	0.5601	0.5466	0.49	0.49	0.49			743
0.6397	0.6496	0.6406	0.6433	0.32	0.33	0.33		Second and third sp. gr. determinations made on sap-wood	873
0.6374	0.6185	0.6260	0.6273	0.38	0.32	0.35			1048
			0.6178			0.37	38.50		
0.5563			0.5563	0.31	0.31	0.31			1239
0.5355			0.5355	0.36	0.36	0.36			1240
			0.5459			0.34	34.02		
0.4332	0.4288		0.4310	0.76	1.22	0.99			290
0.4217	0.4474		0.4346	1.01	1.30	1.10			311
			0.4328			1.07	26.97		
0.4786	0.4856		0.4821	0.51	0.57	0.54	30.04		645
0.6196	0.6393	0.6686	0.6425	0.48	0.51	0.50	40.04		1176

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
70. <i>Rhus typhina</i> <i>Staghorn Sumach.</i>	158	Vermont.....	Hinesburgh.....	C. G. Pringle.....	Gravelly.....			
	1060	Massachusetts.....	Danvers.....	J. Robinson.....	do.....	0.120	2	34
71. <i>Rhus copallina</i> <i>Dwarf Sumach.</i>	70	Missouri.....	Allenton.....	G. W. Letterman.....	Moist limestone.....			
	610	Georgia.....	Lower Altamaha river.....	A. H. Curtiss.....	Dry, clay.....			
	736	Florida.....	Chattahoochee river.....	do.....	do.....	0.175	5	19
71. <i>Rhus copallina</i> , var. <i>lanceolata</i>	330	Texas.....	Dallas.....	J. Beverchon.....	Dry, gravelly.....			
72. <i>Rhus venonata</i> <i>Poison Sumach. Poison Elder.</i>	876	Massachusetts.....	Danvers.....	J. Robinson.....	Wet, swampy.....	0.070	1	20
	1037	do.....	do.....	do.....	do.....	0.085	4	16
	1041	do.....	do.....	do.....	do.....	0.067	4	27
73. <i>Rhus Metopium</i> <i>Poison Wood. Coral Sumach. Mount-ain Manchmet. Bum Wood. Hog Plum. Doctor Gum.</i>	467	Florida.....	Upper Metacombe Key.....	A. H. Curtiss.....	Coral.....	0.222	39	39
LEGUMINOSÆ.								
75. <i>Eysenhardtia orthocarpa</i>	1147	Arizona.....	Santa Rita mount-ains.....	C. G. Pringle.....	Dry, rocky.....			
76. <i>Dalea spinosa</i>	1079	California.....	Agua Caliente.....	Parish Brothers.....	Dry, sandy.....			
77. <i>Robinia Pseudacacia</i> <i>Locust. Black Locust. Yellow Locust.</i>	405		Charlestown Navy-yard.....	S. H. Pook.....				
	441	Tennessee.....	Nashville.....	A. Gattinger.....	Limestone.....			
	815	West Virginia.....	Grafton.....	C. G. Pringle.....				
	845	Massachusetts.....	Danvers.....	J. Robinson.....	Gravelly.....	0.184	4	35
78. <i>Robinia viscosa</i> <i>Clammy Locust.</i>	1061	do.....	do.....	do.....	Loam.....	0.060	4	8
79. <i>Robinia Neo-Mexicana</i> <i>Locust.</i>	1031	Colorado.....	Trinidad.....	W. B. Strong.....	Low, moist.....			
80. <i>Olneya Tesota</i> <i>Iron Wood. Arbol de Hierro.</i>	650	California.....	Lower Colorado valley.....	G. Engelmann and C. S. Sargent.....	Dry, gravelly.....			
81. <i>Piscidia Erythrina</i> <i>Jamaica Dogwood.</i>	564	Florida.....	Upper Metacombe Key.....	A. H. Curtiss.....	Coral.....			
82. <i>Cladrastis tinctoria</i> <i>Yellow Wood. Yellow Ash. Gopher Wood.</i>	33	Kentucky.....	Mercer county.....	W. M. Linney.....	Limestone.....			
	480	Tennessee.....	Nashville.....	A. Gattinger.....	Alluvial.....			
83. <i>Sophora secundiflora</i> <i>Fraxinella.</i>	940	Texas.....	New Braunfels.....	C. Mohr.....	Limestone.....			

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.4476	0.4593		0.4535	0.48	0.54	0.49			158
0.4079	0.4855		0.4170	0.52	0.52	0.52			1080
0.3912	0.3969		0.4357			0.50	27.15		
0.4987	0.5080		0.5089	0.62	0.62	0.62			70
0.5478	0.5579		0.5529	0.63	0.56	0.60			610
0.5472	0.4910	0.5367	0.5250	0.55	0.60	0.57			736
			0.5273			0.60	32.86		
0.5121	0.5236		0.5184	0.89	0.80	0.85	32.31		330
0.4883	0.4363		0.4326	0.90	0.88	0.89			876
0.4192	0.4368								
0.4365	0.4322		0.4418	0.49	0.49	0.49			1837
0.4535	0.4449								
0.4259	0.4511	0.4440	0.4403	0.50	0.60	0.55			1043
			0.4382			0.64	27.31		
0.8106	0.7728		0.7917	2.31	2.47	2.39	49.34		462
0.8691	0.8884	0.8646	0.8740	1.29	1.28	1.28	54.47	0.125 sap-wood	1147
0.5714	0.5904		0.5536	3.41	4.68	4.04	34.50	First, second, and third sp. gr. determinations made on sap-wood; fourth sp. gr. determination made on 0.1 sap-wood.	1079
0.5265	0.5241								
0.7550	0.7904		0.7727	0.22	0.26	0.24			405
0.7908			0.7908	1.26	1.12	1.19			443
0.6490	0.6387		0.6430	0.35		0.35			815
0.7410	0.7279	0.7087	0.7259	0.22	0.25	0.24		Third sp. gr. determination made on sap-wood. Cultivated	845
			0.7333			0.51	45.70		
0.8130	0.8017		0.8094	0.22	0.19	0.20	50.44	0.125 sap-wood. Cultivated	1063
0.7943	0.8207								
0.7968	0.7855	0.8258	0.8034	0.62	0.58	0.60	50.07		1031
0.8953	0.9000	1.1374	1.0902	1.85	2.15				650
1.1542	1.1542			2.75	2.48	2.29	60.07		
0.8779	0.8689		0.8734	3.42	3.34	3.38	51.43		504
0.6072	0.6485		0.6278	0.21	0.17	0.19			33
0.6277			0.6277	0.39	0.36	0.38			439
			0.6278			0.28	39.12		
1.0810	0.9690	0.9325	0.9942	1.44	1.74	1.59	61.34	First and second sp. gr. determinations made on sap-wood	910

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
84. <i>Sophora affinis</i>	329	Texas	Dallas	J. Reverchon	Dry, calcareous.....			
	932	do	Austin	C. Mohr	do	0.084	10	16
85. <i>Gymnocladus Canadensis</i>	206	Missouri.....	Allenton.....	G. W. Letterman	Low, rich			
<i>Kentucky Coffee Tree. Coffee Nut.</i>	402	do	do	do	Alluvial			
	519	Tennessee	Nashville	A. Gattinger	Limestone			
86. <i>Gleditsia triacanthos</i>	531	Missouri.....	Allenton.....	G. W. Letterman	Low, rich			
<i>Honey Locust. Black Locust. Thre-</i>	532	do	do	do	do			
<i>horned Acacia. Sweet Locust.</i>	444	Tennessee	Nashville	A. Gattinger	Dry, sandy barren			
<i>Honey Shucks.</i>								
87. <i>Gleditsia monosperma</i>	760	Florida	Chattahoochee river.	A. H. Curtiss	Alluvial	0.294	21	26
<i>Water Locust.</i>								
88. <i>Parkinsonia Torreyana</i>	678	Arizona	Lower Colorado river.	G. Engelmann and C. S. Sargent.	Sandy			
<i>Green-bark Acacia. Palo Verde.</i>								
89. <i>Parkinsonia microphylla</i>	1258	do	Valley of the Gila river.	C. G. Pringle				
90. <i>Parkinsonia aculeata</i>	829	Florida		Department of Agriculture.				
	1208	Texas	Austin	S. B. Buckley				
91. <i>Cercis Canadensis</i>	59	Missouri.....	Allenton.....	G. W. Letterman	Low, rich	0.200	5	35
<i>Redbud. Judas Tree.</i>	436	Tennessee	Nashville	A. Gattinger	Limestone			
	1089	Missouri.....	Allenton.....	G. W. Letterman	Rich			
	1090	do	do	do	do			
	1091	do	do	do	do			
92. <i>Cercis reniformis</i>	1142	Texas	Austin	S. B. Buckley	Limestone			
<i>Redbud.</i>								
93. <i>Prosopis juliflora</i>	561	do	Fort Stockton	B. L. Baldrige	Light, sandy			
<i>Mesquit. Algaroba. Honey Locust.</i>	590	Arizona	Tucson	C. S. Sargent				
<i>Honey Pod.</i>	680	do	do	do				
	927	Texas	Austin	C. Mohr	Rich, calcareous			
94. <i>Prosopis pubescens</i>	600	California.....	Fort Yuma	G. Engelmann and C. S. Sargent.				
<i>Screw Bean. Screw-pod Mesquit. Tor-</i>	658	do	do	do	Sandy			
<i>nilla.</i>	998	do	do	do		0.084	8	
95. <i>Leucaena glauca</i>	1224	Mexico	Lampasas mountains	S. B. Buckley				
96. <i>Leucaena pulverulenta</i>	1222	Texas	Brownsville	do	Sandy			

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.8019	0.8161		0.8090	0.42	0.84	0.38		0.25 sap-wood	329
0.8540	0.9316		0.8928	1.04	1.13	1.09		0.5 sap-wood	932
			0.8509			0.73	53.03		
0.6700	0.6786		0.6743	0.70	0.57	0.64			236
0.6663			0.6663	0.45	0.50	0.48			402
0.7395			0.7395	0.98	0.82	0.90			519
			0.6934			0.67	43.21		
0.7330	0.7359		0.7345	0.55	0.61	0.58			531
0.6418	0.6275		0.6346	0.69		0.69			532
0.6328			0.6528	1.29	0.98	1.14			444
			0.6740			0.80	42.00		
0.7509	0.7245	0.7272	0.7342	0.66	0.79	0.73	45.76	First sp. gr. determination made on sap-wood	760
0.6330	0.6732		0.6531	1.25	1.00	1.12	40.70	Second sp. gr. determination made on sap wood	678
0.7176	0.7722		0.7449	4.20	3.08	3.64	49.75		1258
0.6325			0.6325	2.32	2.26	2.29			829
0.5848	0.5968		0.5908	2.17	2.51	2.34			1208
			0.6116			2.32	38.11		
0.6104	0.6070	0.6157	0.6110	0.59	0.56	0.58			59
0.7098			0.7098	0.67	0.67	0.67			436
0.6254	0.5959		0.6107	0.80	0.77	0.79			1069
0.6493	0.6084		0.6289	0.82	0.74	0.78			1090
0.6009	0.6417		0.6213	0.67	0.84	0.76			1091
			0.6363			0.72	39.65		
0.7392	0.7645	0.7502	0.7513	0.76	0.77	0.77	46.82	First and second sp. gr. determinations made on half sap wood	1142
0.8322	0.8064		0.8498	2.25	2.69	3.02		Root	561
0.7137	0.6997		0.7067	1.67	1.71	1.69			500
0.7247	0.7726		0.7457	1.45	2.49	1.97			680
0.7478	0.7620	0.7587	0.7562	2.09	2.00	2.05		0.05 sap-wood	927
			0.7652			2.18	47.13		
0.7772	0.7910		0.7841	1.02	0.98	1.00		Dead tree; 0.75 sap-wood	600
0.7527	0.7930		0.7729	0.91	0.90	0.91		All sap-wood	658
0.7237	0.7278		0.7258	0.95		0.95		All sap-wood	996
			0.7609			0.95	47.42		
0.9265	0.9205		0.9235	3.27	3.31	3.29	57.55		1224
0.6955	0.6809		0.6732	0.90	1.13	1.01	41.95		1222

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
97. <i>Acacia Wrightii</i> <i>Cat's Claw.</i>	1209	Texas	Austin	S. B. Buckley				
98. <i>Acacia Greggii</i> <i>Cat's Claw.</i>	598	Arizona	Santa Rita mount- ains.	G. Engelmann and C. S. Sargent.	Dry, gravelly			
	697	do	do	do	do			
	893	do	Clifton	E. L. Greene				
100. <i>Lyalloma latifolia</i> <i>Wild Tamarind.</i>	509	Florida	Boca Chica Key	A. H. Curtiss	Coral	0.202	14	29
	1112	do	Key Largo	do	do			
101. <i>Pithecolobium Unguis-cati</i> <i>Cat's Claw.</i>	405	do	Upper Metacombe Key.	do	do			
	1108	do	Bahai Honda Key	do	do			
ROSACEÆ.								
102. <i>Chrysobalanus Icaco</i> <i>Ocoa Plum.</i>	480	do	Bay Biscayne	A. H. Curtiss	Swampy	0.100	47	
103. <i>Prunus Americana</i> <i>Wild Plum. Canada Plum. Horse Plum.</i>	68	Missouri	Allenton	G. W. Letterman	Rich uplands			
	220	Vermont	Charlotte	C. G. Pringle	Gravelly			
	334	Texas	Dallas	J. Reverchon	Rich			
104. <i>Prunus angustifolia</i> <i>Chickasaw Plum. Hog Plum.</i>	495	Tennessee	Nashville	A. Gittinger	River bluff			
105. <i>Prunus Pennsylvanica</i> <i>Wild Red Cherry. Pin Cherry. Pigeon Cherry.</i>	233	Vermont	Charlotte	C. G. Pringle	Cold, gravelly			
106. <i>Prunus umbellata</i> <i>Sloe. Black Sloe.</i>	606	Georgia	Altamaha river	A. H. Curtiss	Clay	0.216	21	24
107. <i>Prunus emarginata, var. mollis</i>	968	Washington ter- ritory.	Wilkeson	G. Engelmann and C. S. Sargent.	Low, rich			
108. <i>Prunus serotina</i> <i>Wild Black Cherry. Rum Cherry.</i>	15	Massachusetts	Roxbury	C. S. Sargent	Gravelly	0.205	9	31
	106	Vermont	Charlotte	C. G. Pringle	do			
	115	Michigan	Danville	W. J. Beal	do			
	127	Missouri	Allenton	G. W. Letterman	Rich loam			
	148	Illinois	Waukegan	H. Douglas	Gravelly			
	198	Ohio	Barney & Smith Manufacturing Co.	E. E. Barney				
	317	Michigan	Hersey	W. J. Beal	Rich			
	368	Vermont	Charlotte	C. G. Pringle	Gravelly			
	406	Virginia or Mid- dle states.	Charlestown Navy- yard.	S. H. Pook				
	725	Pennsylvania	Williamsport	C. G. Pringle				
	763	Florida	Chattahoochee river.	A. H. Curtiss	Clay	0.218	15	48
	1063	Massachusetts	Topsfield	J. Robinson	Gravelly	0.320	11	38
109. <i>Prunus Capuli</i> <i>Wild Cherry.</i>	294	New Mexico	Pinos Altos mount- ains.	E. L. Greene	Alluvial	0.238	7	12
	418	do	do	do	do			

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.9380	0.9404		0.9392	0.68	0.57	0.63	58.53		1209
0.8703			0.8703	0.68	0.89	0.79		0.05 sap-wood	596
0.8162	0.8616		0.8389	0.95		0.95		0.05 sap-wood	607
0.8558			0.8558	1.00		1.00			893
			0.8550			0.91	53.28		
0.6033	0.5603		0.5848	2.00	2.44	2.23			509
0.6901	0.7159	0.6001	0.6987	1.98	2.07	2.03		Rough bark	1112
			0.6418			2.12	40.00		
0.8829	0.8349		0.8589	2.48	2.25	2.37			465
0.8885	1.0135		0.9510	3.00	2.09	2.55			1108
			0.9049			2.46	56.39		
0.7290	0.8129		0.7709	0.87	0.87	0.87	48.04		480
0.6750	0.6745		0.6748	0.11	0.09	0.10			68
0.7035	0.6800		0.6918	0.17	0.21	0.19			220
0.7876	0.8079		0.7978	0.33	0.21	0.27			334
			0.7215			0.18	44.96		
0.6675	0.7092		0.6884	0.28	0.28	0.28	42.90		485
0.5014	0.5031		0.5023	0.36	0.43	0.40	31.30		223
0.8162	0.8242		0.8202	0.14	0.10	0.12	51.11		606
0.4310	0.4604		0.4502	0.18	0.23	0.21	28.06		968
0.7047	0.7079		0.7063	0.13	0.14	0.14		0.5 sap-wood	15
0.5525	0.5539		0.5532	0.13	0.11	0.12			106
0.5809	0.4832		0.5321	0.12	0.15	0.14			115
0.6736	0.6235		0.6486	0.19	0.17	0.18			127
0.6473	0.6508		0.6491	0.11	0.13	0.12			148
0.5675	0.5659		0.5667	0.13	0.11	0.12			198
0.5373	0.5186		0.5280	0.10	0.17	0.18			317
0.5131	0.5351		0.5241	0.14	0.17	0.16			368
0.4680			0.4680	0.06	0.09	0.08			406
0.5734	0.5777		0.5756	0.09	0.10	0.10			726
0.5755	0.5221		0.5488	0.17	0.34	0.25			703
0.6833	0.6857	0.6880	0.6857	0.15	0.17	0.16			1058
			0.5822			0.15	36.28		
0.7282	0.6940		0.7111	0.16	0.12	0.14			294
0.8459	0.8837		0.8648	0.28	0.26	0.27		0.33 sap-wood	418
			0.7879			0.20	49.10		

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
110. <i>Prunus demissa</i> <i>Wild Cherry.</i>	637	California.....	Strawberry valley ..	G. Engelmann and C. S. Sargent.	Low, rich			
111. <i>Prunus Caroliniana</i> <i>Wild Orange. Mock Orange. Wild Peach.</i>	549	Alabama	Mobile.....	C. Mohr	Sandy			
	806	Florida	Jacksonville.....	A. H. Curtiss	do	0.171	8	61
	916	do	Chattahoochee.....	C. Mohr	Alluvial	0.202	12	50
	1032	do	Jacksonville.....	A. H. Curtiss.....	Sandy			
	1062	Texas	Victoria	C. Mohr	Rich, moist	0.180	26	34
112. <i>Prunus sphaerocarpa</i>	1217	Florida	Edge of Everglades.	A. H. Curtiss	Rocky			
113. <i>Prunus ilicifolia</i>	1158	California	Santa Cruz	C. L. Anderson		0.206	80	
114. <i>Vauquelinia Torreyi</i>	1152	Arizona	Santa Rita mount- ains.	C. G. Pringle.....				
115. <i>Cercocarpus ledifolius</i> <i>Mountain Mahogany.</i>	883	Utah	City Creek cañon...	M. E. Jones	Rocky			
	904			Department of Ag- riculture.				
116. <i>Cercocarpus parvifolius</i>	825	California.....	Siakiyou county...	G. Engelmann and C. S. Sargent.		0.062	3	27
117. <i>Pyrus coronaria</i>	808	Delaware	Kiamensi	W. M. Canby	Clay			
<i>American Crab. Sweet-scented Crab.</i>	1087	Pennsylvania.....	Nazareth	J. Henry	Moist			
	1088	do	do	do	do			
118. <i>Pyrus angustifolia</i>	313	South Carolina...	Aiken	H. W. Ravenel.....	Rich, damp	0.130	25	
<i>American Crab Apple. Southern Crab Apple.</i>								
119. <i>Pyrus rivularis</i>	1002	Oregon	Portland.....	G. Engelmann and C. S. Sargent.	Rich hills.....	0.166		
<i>Oregon Crab Apple.</i>								
120. <i>Pyrus Americana</i>	214	Vermont.....	Charlotte	C. G. Pringle.....	Gravelly			
<i>Mountain Ash.</i>	365	do	Huntingdon	do	do			
121. <i>Pyrus sambucifolia</i>	410	do	Mount Mansfield ..	do	do	0.106	17	
<i>Mountain Ash.</i>								
122. <i>Crataegus rivularis</i>	885	Utah	City Creek cañon ..	M. E. Jones	do	0.064	14	13
123. <i>Crataegus Douglasii</i>	999	Oregon	Cascades of the Co- lumbia river.	G. Engelmann and C. S. Sargent.	Rich	0.200	29	
124. <i>Crataegus brachyacantha</i>	926	Louisiana.....	Webster parish.....	C. Mohr	Clay			
<i>Hoge's Hawth.</i>								
125. <i>Crataegus arborecens</i>	363 ¹	South Carolina ..	Aiken	H. W. Ravenel.....	Rich			
	363 ²	do	do	do	do			
	607	Georgia.....	Ogeechee	A. H. Curtiss	Low	0.168	46	

OF DRY SPECIMENS OF WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.7025	0.6937	0.6800	0.6951	0.51	0.48	0.50	43.32		637
0.9515	0.9855		0.9685	0.32	0.33	0.33			549
0.8874	0.8227	0.8314	0.8472	0.39	0.37	0.38			806
0.7706	0.7797		0.7782	0.44	0.44	0.44			916
0.8650	0.8721		0.8681	0.46	0.49	0.48			1032
0.9080	0.8610	0.8790	0.8810	0.43	0.43	0.43			1002
			0.8088			0.41	54.14		
0.9062	0.8934		0.8998	0.97	0.76	0.87	56.07		1217
0.9685	0.9720		0.9803	0.80	0.76	0.78	61.09		1158
1.1840	1.1408		1.1374	1.44	1.46	1.45	70.88		1152
1.0608	1.0332	1.0400	1.0447	1.27	1.12	1.19		0.125 sap-wood	883
1.0950	1.1080		1.1015	1.09	0.69	0.89			904
			1.0731			1.04	66.87		
0.9365			0.9365	0.45		0.45	58.36		825
0.6973	0.7228	0.7320	0.7174	0.52	0.48	0.50			808
0.6640	0.6762	0.6788	0.6730	0.50	0.38	0.44			1087
0.7402	0.7260	0.7057	0.7240	0.49	0.75	0.62			1088
			0.7048			0.52	43.92		
0.6945	0.6845		0.6895	0.33	0.33	0.33	42.97		313
0.8266	0.8366		0.8316	0.41	0.41	0.41	51.82		1002
0.5345	0.5429	0.5539	0.5438	0.98	1.23	1.11			214
0.5512	0.5416		0.5464	0.50	0.56	0.56		0.25 sap-wood	365
			0.5451			0.83	33.97		
0.5893	0.5963		0.5928	0.39	0.31	0.35	36.94	Sap-wood	410
0.7672	0.7734		0.7703	0.33	0.37	0.35	48.00		885
0.7025	0.6928	0.6898	0.6950	0.29	0.36	0.33	43.31		999
0.6750	0.6835		0.6793	0.37	0.46	0.42	42.33		926
0.6306	0.6230		0.6228	0.63	0.70	0.67			363 ¹
0.6083	0.6225								
0.6263	0.6390		0.6127	0.38		0.38			363 ²
0.6934	0.6899		0.6817	0.62	0.68	0.65			607
			0.6491			0.57	40.45		

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
126. <i>Crataegus Crus-galli</i> <i>Cockspur Thorn. Newcastle Thorn.</i>	828	Massachusetts....	Brookline.....	J. Robinson.....	Loam.....	0.180		
	1093	Missouri.....	Allenton.....	G. W. Letterman.....	Low, wet.....			
127. <i>Crataegus coccinea</i> <i>Scarlet Haw. Red Haw. White Thorn.</i>	100	Vermont.....	Monkton.....	C. G. Pringle.....	Gravelly.....			
128. <i>Crataegus subvillosa</i> <i>Scarlet Haw.</i>	949	Texas.....	Victoria.....	C. Mohr.....	Alluvial.....	0.180	61	
	1081	Missouri.....	Saint Louis.....	H. Eggert.....		0.106	28	
129. <i>Crataegus tomentosa</i> <i>Black Thorn. Pear Haw.</i>	154	Vermont.....	Charlotte.....	C. G. Pringle.....	Clay.....			
	426	Tennessee.....	Nashville.....	A. Gittinger.....	Limestone.....			
130. <i>Crataegus cordata</i> <i>Washington Thorn.</i>	447	do.....	do.....	do.....	Low.....			
131. <i>Crataegus apifolia</i> <i>Parsley Haw.</i>	759	Florida.....	Chattahoochee.....	A. H. Curtiss.....	Alluvial.....	0.080	27	
132. <i>Crataegus spatulata</i> <i>Small-fruited Haw.</i>	800	South Carolina.....	Aiken.....	H. W. Ravenel.....	Rich, damp.....			
	614	Georgia.....	Ogeechee river.....	A. H. Curtiss.....	Low.....	0.880	38	
134. <i>Crataegus setivialis</i> <i>May Haw. Apple Haw.</i>	239	South Carolina.....	Bonneau's Depot.....	H. W. Ravenel.....	Damp, rich.....			
135. <i>Crataegus flava</i> <i>Summer Haw. Yellow Haw.</i>	301	do.....	Aiken.....	do.....	Dry, fertile.....	0.224	58	
	569	Florida.....	Tampa.....	A. H. Curtiss.....	Sandy.....			
135. <i>Crataegus flava</i> , var. <i>pubescens</i> <i>Summer Haw. Red Haw.</i>	767	do.....	Aspalaga.....	do.....	Dry clay.....	0.142	44	
136. <i>Heteromeles arbutifolia</i> <i>Toyon. Tollyn. California Holly.</i>	1180	California.....	Santa Cruz.....	C. L. Anderson.....		0.184	21	34
137. <i>Amelanchier Canadensis</i> <i>Juneberry. Shad Bush. Service Tree. May Cherry.</i>	156	Vermont.....	Charlotte.....	C. G. Pringle.....	Gravelly.....			
	241	Kentucky.....	Brunfield Station.....	W. M. Linney.....	Waverly shale.....	0.140	34	69
	849	Massachusetts.....	Danvers.....	J. Robinson.....	Loam.....	0.190	43	21
HAMAMELACEÆ.								
138. <i>Hamamelis Virginica</i> <i>Witch Hazel.</i>	867	do.....	Beverly.....	do.....	Rocky.....			
	875	do.....	Danvers.....	do.....	do.....			
	1259	Tennessee.....	Montvale.....	A. H. Curtiss.....				
139. <i>Liquidambar Styraciflua</i> <i>Sweet Gum. Star-leaved Gum. Liquid-amber. Red Gum. Bilsted.</i>	546	Alabama.....	Kemper's mill.....	C. Mohr.....	Rich, alluvial.....			
	1095	Arkansas.....	Little Rock.....	G. W. Letterman.....				
	1173	New Jersey.....	Mount Holly.....	S. P. Sharples.....	Clay.....	0.208	39	17
	1181	Mississippi.....	Yazoo River bottom.....	R. Abbey.....	Alluvial.....			

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.6590	0.6705	0.6632	0.6642	0.57	0.55	0.56		Cultivated	323
0.7781	0.7667	0.7791	0.7746	0.49	0.60	0.55			1063
			0.7194			0.56	44.83		
0.8654	0.8582		0.8618	0.36	0.40	0.38	53.71		100
0.7909	0.7144		0.7527	0.77	0.78	0.78			949
0.8376	0.8382		0.8379	0.54	0.68	0.61			1061
			0.7953			0.69	49.56		
0.7649	0.7712		0.7681	0.50	0.44	0.47			154
0.8059	0.7110		0.7585	0.53	0.51	0.52			426
			0.7683			0.50	47.57		
0.7105	0.7461		0.7293	0.48	0.44	0.46	45.45		447
0.7506	0.7400		0.7453	0.72	1.22	0.97	46.45		759
0.7524	0.7696		0.7611	0.71	0.67	0.69			809
0.6716	0.6700		0.6708	0.60	0.63	0.62			614
			0.7159			0.66	44.61		
0.6625	0.6503		0.6564	0.57	0.56	0.57	40.91		339
0.7724	0.7620		0.7672	0.82	0.94	0.88			801
0.8172	0.7718		0.7945	0.65	0.78	0.69			599
			0.7809			0.79	48.67		
0.7770	0.7596		0.7683	0.91	0.91	0.91	47.88		767
0.9610	0.9042		0.9326	0.62	0.46	0.54	58.18		1166
0.6074	0.7642		0.6858	0.61	0.64	0.63		First sp. gr. determination made on 0.75 sap-wood; second sp. gr. determination made on 0.5 sap-wood.	156
0.7904	0.7944		0.7924	0.42	0.53	0.48		0.25 sap-wood	241
0.8600	0.9028	0.8570	0.8733	0.65	0.45	0.55		First and third sp. gr. determinations made on 0.9 sap-wood; second sp. gr. determination made on sap-wood.	849
			0.7888			0.55	48.85		
0.7067	0.7398		0.7342	0.43	0.44	0.44			867
0.7031	0.7826								
0.6662	0.7828		0.7126	0.27	0.33	0.30			875
0.7028	0.6985								
0.6099			0.6099	0.32	0.39	0.36			1269
			0.6836			0.37	42.72		
0.6415	0.6176		0.6295	0.47	0.49	0.48			546
0.5773	0.5665		0.5719	0.48	0.73	0.61			1093
0.5841	0.5991		0.5916	0.29	0.34	0.32		All sap-wood	1173
0.5605	0.5549	0.5692	0.5615	0.67	0.66	0.67			1181

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
139. Liquidambar styraciflua—continued.	1182	Mississippi	Yazoo River bottom	R. Abbey	Alluvial			
	1183	do	do	do	do			
RHIZOPHORACEÆ.								
140. Rhizophora Mangle. Mangrove.	485	Florida	Bay Biscayne.	A. H. Curtiss	Salt-marsh	0.250	31	50
COMBRETACEÆ.								
141. Conocarpus erecta. Button Wood.	489	do	do	do	do	0.164	15	24
142. Laguncularia racemosa. White Button Wood. White Mangrove.	507	do	Sugar-loaf Sound	do	do			
MYRTACEÆ.								
143. Calyptranthes Chytraculia	1205	do	Key Largo	do	Coral			
144. Eugenia buxifolia. Gurgon Stopper. Spanish Stopper.	456	do	Upper Metacombe Key.	do	do	0.084	43	
	1118	do	Lost Man's river	do	Humus and coral			
	1120	do	Elliott's Key	do	Coral			
	1198	do	Upper Metacombe Key.	do	do			
145. Eugenia dichotoma. Naked Wood.	506	do	Caximbas pass	do	Sandy	0.084	5	53
	1200	do	Palm Hummock	do	Coral			
146. Eugenia monticola. Stopper. White Stopper.	1115	do	Umbrella Key	do	do	0.150	15	
	1185	do	do	do	do			
	1189	do	do	do	do			
147. Eugenia longipes. Stopper.	1197	do	No-Name Key	do	do			
148. Eugenia procera. Red Stopper.	1127	do	Miami	do	do	0.141	87	
CACTACEÆ.								
149. Cereus giganteus. Suearrow. Saguaro. Giant Cactus.	603	Arizona	Tucson	G. Engelmann and C. S. Sargent.	Dry, gravelly			
CORNACEÆ.								
150. Cornus alternifolia. Dogwood.	860	Massachusetts	Danvers	J. Robinson	Lean			
	861	do	do	do	do			
151. Cornus florida. Flowering Dogwood. Boz Wood.	67	Missouri	Allenton	G. W. Letterman	Uplands	0.143	47	
	761	Florida	Chattahoochee	A. H. Curtiss	Calcareous	0.128	72	
	812	West Virginia	Grafton	C. G. Pringle	Dry			
	1077	Missouri	Allenton	G. W. Letterman	Gravelly	0.122	44	7
	1092	do	do	do	Flinty			
152. Cornus Nuttallii. Flowering Dogwood.	960	Oregon	Portland	G. Engelmann and C. S. Sargent.				

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.6125	0.5891	0.6256	0.6091	0.72	0.77	0.75			1182
0.5825	0.5765	0.5878	0.5823	0.85	0.77	0.81			1183
			0.5910			0.61	36.83		
1.1291	1.1842		1.1617	1.80	1.83	1.82	72.40		485
0.9939	0.9860		0.9900	0.32	0.32	0.32	61.70		489
0.7100	0.7174		0.7137	1.69	1.54	1.62	44.48		507
0.8930	0.9282	0.8765	0.8992	3.04	3.59	3.32	56.04		1205
0.8753			0.8753	1.11	1.26	1.19			456
1.0625	1.0400		1.0513	1.19	1.05	1.12			1118
0.8645	0.8540	1.0200	0.9128	2.23	2.15	2.19			1120
0.8857	0.9232		0.9045	1.66	1.88	1.52			1198
			0.8960			1.50	58.23		
0.8526	0.8392		0.8459	0.85	0.91	0.88			566
0.9542	0.9472		0.9507	0.58	0.60	0.59			1200
			0.8983			0.74	55.98		
0.8910	0.9110		0.9010	2.05	1.95	2.00			1115
0.9180	0.9340	0.9202	0.9241	1.60	1.95	1.77			1125
0.9140	0.9295		0.9217	1.83	1.99	1.91			1189
			0.9156			1.89	57.06		
1.1450	1.1020		1.1235	3.36	3.60	3.48	70.02		1197
0.9425	0.9482		0.9453	2.44	2.79	2.62	58.91		1127
0.3259	0.3116		0.3188	3.89	3.01	3.45	19.87		693
0.6572	0.6748	0.6048	0.6456	0.43	0.41	0.42		All sap-wood	860
0.6967	0.6932	0.6993	0.6937	0.39	0.43	0.41		All sap-wood	861
			0.6696			0.41	41.73		
0.8264	0.8264		0.8264	0.51	0.60	0.56			67
0.7599	0.7710		0.7655	0.75	0.84	0.80			761
0.7892	0.7863		0.7878	0.73	0.83	0.78			812
0.8690	0.8916		0.8803	0.66	0.58	0.62			1077
0.8114	0.8215		0.8165	0.62	0.60	0.61			1092
			0.8153			0.67	50.81		
0.7487	0.7474		0.7481	0.48	0.52	0.50	46.62		960

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
153. <i>Nyssa capitata</i> <i>Ogeechee Lime. Sour Tupelo. Gopher Plum.</i>	605	Georgia.....	Ogeechee river.....	A. H. Curtiss.....	Swampy.....	0.220	27	
154. <i>Nyssa sylvatica</i> <i>Tupelo. Sour Gum. Pepperidge. Black Gum.</i>	235	South Carolina....	Bonneau's Depot...	H. W. Ravenel....	Muck.....			
	517	Tennessee.....	Cumberland river...	A. Gattlinger.....				
	608	Georgia.....	Ogeechee river.....	A. H. Curtiss.....	Muck.....	0.220	49	
	750	Florida.....	Chattahoochee.....	do.....	Clay.....	0.214	49	
	813	West Virginia....	Grafton.....	C. G. Pringle.....				
	833	Massachusetts....	West Newbury.....	J. Robinson.....	Rich.....	0.256	66	23
	834	do.....	do.....	do.....	do.....			
	835	do.....	Chebaeco pond.....	do.....	Low, rich.....			
155. <i>Nyssa uniflora</i> <i>Large Tupelo. Cotton Gum. Tupelo Gum.</i>	123	South Carolina....	Bonneau's Depot...	H. W. Ravenel.....	Swampy.....			
	235	do.....	do.....	do.....		0.188	30	
	550	Alabama.....	Stockton.....	C. Mohr.....	Alluvial.....	0.320	67	19
	604	Georgia.....	Ogeechee river.....	A. H. Curtiss.....	Swampy.....			
CAPRIFOLIACEÆ.								
156. <i>Sambucus glauca</i> <i>Elder.</i>	681	California.....	Contra Costa county.	G. R. Vasey.....	Gravelly.....	0.320	6	28
157. <i>Sambucus Mexicana</i> <i>Elder.</i>	1220	Arizona.....	Santa Catalina mountains.	C. G. Pringle.....				
158. <i>Viburnum Lentago</i> <i>Sheepberry. Nannyberry.</i>	108	Vermont.....	Charlotte.....	do.....	Muck.....			
	370	do.....	Hinesburg.....	do.....	Swampy.....			
159. <i>Viburnum prunifolium</i> <i>Black Haw. Stag Bush.</i>	42	Kentucky.....	Mercer county.....	W. M. Linney.....	Shale.....			
	1104	do.....	do.....	do.....	Trenton limestone.			
	739	Georgia.....	Bainbridge.....	A. H. Curtiss.....	Clay.....	0.076	36	
RUBIACEÆ.								
160. <i>Rxostemma Caribæum</i>	466	Florida.....	Upper Metacombe Key.	do.....	Coral.....	0.076	28	25
161. <i>Pinckneya pubens</i> <i>Georgia Bark.</i>	257	South Carolina....	Bluffton.....	J. H. Mellichamp...	Sandy, swamp.....	0.108	5	24
	281	do.....	do.....	do.....	do.....			
162. <i>Genipa olusifolia</i> <i>Seven-year Apple.</i>	457	Florida.....	Upper Metacombe Key.	A. H. Curtiss.....	Sandy.....			
	1132	do.....	Elliott's Key.....	do.....	Coral.....			
	1195	do.....	Upper Metacombe Key.	do.....	do.....			
163. <i>Guettarda elliptica</i>	471	do.....	do.....	do.....	do.....	0.095	34	
	1129	do.....	Umbrella Key.....	do.....	do.....			
	1194	do.....	do.....	do.....	do.....			

THE WOODS OF THE UNITED STATES.

291

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.4680	0.4546		0.4613	0.85	0.83	0.84	28.75		606
0.5656	0.5649		0.5652	0.66	0.81	0.74			235
0.5650			0.5650	0.89	0.79	0.84			517
0.6079	0.6104		0.6092	0.49	0.43	0.46		N. Caroliniana	608
0.6436	0.6682		0.6559	0.49	0.49	0.49			750
0.6198	0.5965		0.6031	0.52	0.53	0.53			813
0.6649	0.6787	0.6881	0.6539	0.41	0.40	0.41			833
0.7467	0.6748	0.6708	0.6974	0.88	0.89	0.88			834
0.7429	0.6626	0.7022	0.7026	0.82	0.84	0.83			835
			0.6358			0.52	39.59		
0.5589	0.5700		0.5645	0.72	0.76	0.74			128
0.5656	0.5649		0.5653	0.66	0.81	0.74			236
0.5002	0.5525		0.5264	0.58	0.59	0.59			550
0.4424	0.4002		0.4213	0.76	0.66	0.71			604
			0.5194			0.70	82.87		
0.5076	0.5098		0.5087	1.60	1.55	1.57	81.70		681
0.4588	0.4608	0.4586	0.4614	1.88	2.17	2.00	28.75		1220
0.7285	0.7075		0.7180	0.27	0.28	0.28		0.1 sap-wood	108
0.7519	0.7832		0.7426	0.29	0.30	0.30			870
			0.7303			0.29	45.51		
0.8106			0.8106	0.46	0.49	0.48		0.75 sap-wood	42
0.9140			0.9140	0.47		0.47		All sap-wood	110
0.7749			0.7749	0.60		0.60		All sap-wood	739
			0.8332			0.52	51.92		
0.9200	0.9419		0.9310	0.24	0.23	0.23	58.02		406
0.5528	0.5613		0.5571	0.28	0.27	0.28			267
0.5126	0.5181		0.5129	0.49	0.58	0.54			381
			0.5850			0.41	33.41		
1.0219			1.0219	0.71	0.60	0.66		All sap-wood	457
1.0425	1.0279	0.9175	0.9957	1.00	1.02	1.01		Second and third sp. gr. determinations made on 0.3 sap-wood	1123
1.0765	1.0840		1.0772	1.48	1.55	1.52			1195
			1.0816			1.06	64.29		
0.8375	0.8983		0.9154	0.88	0.79	0.84			471
0.8409	0.8508		0.8504	0.94	1.03	0.99			1129
0.7499	0.7217		0.7353	1.83	1.82	1.82			1194
			0.8337			1.05	51.96		

FOREST TREES OF NORTH AMERICA.

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
ERICACEÆ.								
164. <i>Vaccinium arboreum</i> <i>Falseberry.</i>	243	Alabama	Citronelle.....	C. Mohr	Sandy	0.127	65	
	612	Georgia.....	Altamaha river.....	A. H. Curtis	Swampy			
165. <i>Andromeda ferruginea</i>	1083	Florida	Jacksonville.....	do	Hummock			
	1084	do	do	do		0.216	5	29
166. <i>Arbutus Menziesii</i> <i>Madroña.</i>	643	California	Contra Costa county	G. R. Vasey.....	Gravelly.....			
	679	do	Marin county.....	do	do			
167. <i>Arbutus Kalapensis</i>	596	Arizona	Santa Rita mount- ains.	G. Engelmann and C. S. Sargent.	Rocky			
168. <i>Arbutus Texana</i>	1085	Texas	Hays county.....	S. B. Buckley	Limestone	0.086	22	26
169. <i>Oxydendrum arboreum</i> <i>Sorrel Tree. Sour Wood.</i>	853	Alabama	Cottage Hill.....	C. Mohr	Light, rich	0.130	28	
	515	Tennessee	Nashville.....	A. Gattinger.....	Sandy, rocky			
170. <i>Kalmia latifolia</i> <i>Laurel. Calico Bush. Spoon Wood.</i> <i>Ivy.</i>	262 ¹	Virginia	Fancy Gap	H. Shriver	Moist.....	0.082	74	
	262 ²	do	do	do	do			
	262 ³	do	do	do	do			
171. <i>Rhododendron maximum</i> <i>Great Laurel. Rose Bay.</i>	263 ¹	do	do	do	do	0.078	52	
	263 ²	do	do	do	do			
MYRSINACEÆ.								
172. <i>Myrsine Rapaia</i>	1123	Florida	Bay Biscayne.....	A. H. Curtis	Coral			
173. <i>Ardisia Pickeringia</i> <i>Marlberry. Cherry.</i>	494	do	do	do	do	0.080	15	18
	1186	do	do	do	do			
	1192	do	Palm creek.....	do	do			
174. <i>Jacquinia armillaris</i> <i>Joe Wood.</i>	498	do	Key Largo.....	do	do			
	1130	do	Elliott's Key	do	do			
	1199	do	Umbrella Key	do	do			
SAPOTACEÆ.								
175. <i>Chrysophyllum oliviforme</i>	492	do	Bay Biscayne.....	do	do	0.105	18	
176. <i>Sideroxylon Mastichodendron</i> <i>Mastic.</i>	461	do	Upper Metacombe Key.	do	do	0.206	29	49

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.7814	0.7479	-----	0.7647	0.42	0.45	0.44	-----	-----	348
0.7740	0.7405	-----	0.7573	0.35	0.32	0.34	-----	-----	612
			0.7610			0.39	47.42	-----	
0.7766	0.8021	-----	0.7894	0.45	0.37	0.41	-----	-----	1083
0.7037	0.7174	-----	0.7106	0.56	0.45	0.51	-----	-----	1034
			0.7500			0.46	46.74	-----	
0.6452	0.6989	-----	0.6696	0.42	0.38	0.38	-----	-----	643
0.7482	0.7332	-----	0.7407	0.42	0.39	0.41	-----	-----	679
			0.7052			0.40	43.95	-----	
{ 0.7202	{ 0.6955	-----	0.7099	{ 0.21	{ 0.24	0.25	-----	-----	596
{ 0.6842	{ 0.7397			{ -----	{ 0.30				
0.7585	0.7674	0.7292	0.7500	0.54	0.48	0.51	46.74	0.5 sap-wood.....	1085
0.7920	0.7574	0.7916	0.7803	0.42	0.49	0.46	-----	-----	358
0.7114			0.7114	0.27	0.31	0.29	-----	-----	515
			0.7458			0.37	46.48	-----	
0.7586			0.7536	0.61		0.61	-----	-----	262 ¹
0.7214			0.7214	0.31		0.31	-----	-----	262 ²
0.6730			0.6730	0.32		0.32	-----	-----	262 ³
			0.7160			0.41	44.62	-----	
0.6266			0.6266	0.48		0.48	-----	-----	263 ¹
0.6341			0.6341	0.29		0.29	-----	-----	263 ²
			0.6303			0.36	39.28	-----	
0.8491	0.8541	0.7990	0.8341	0.74	0.88	0.81	51.96	-----	1123
0.8154	0.8412	-----	0.8283	1.81	1.84	1.83	-----	-----	494
0.8800	0.8532	-----	0.8666	1.74	1.95	1.85	-----	-----	1136
0.8770	0.8942	-----	0.8856	1.75	1.95	1.85	-----	-----	1192
			0.8602			1.85	53.61	-----	
0.5693	0.6500	0.6582	0.6258	4.09	3.82	3.96	-----	-----	496
0.6437	0.6475	-----	0.6456	3.47	3.74	3.61	-----	-----	1130
0.8300	0.7965	-----	0.8132	2.85	2.73	2.79	-----	-----	1199
			0.6948			3.45	43.30	-----	
0.8986	0.9433	0.9690	0.9360	1.36	1.11	1.24	58.33	-----	492
1.0125	1.0149	1.0052	1.0109	4.90	5.37	5.14	63.00	-----	461

FOREST TREES OF NORTH AMERICA.

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
177. <i>Dipholis salicifolia</i> <i>Busia. Cassada.</i>	488	Florida	Bay Biscayne	A. H. Curtiss	Coral	0.170	28	14
	500	do	Umbrella Key	do	do	0.150	7	25
	1191	do	No-Name Key	do	do			
178. <i>Bumelia tenax</i>	746	Georgia	Bainbridge	do	Low	0.128	29	
179. <i>Bumelia lanuginosa</i> <i>Gum Elastic. Shittim Wood.</i>	60	Missouri	Allenton	G. W. Letterman	Limestone	0.286	57	
	930	Texas	Austin	C. Mohr	do	0.237	65	
	1083	Missouri	Allenton	G. W. Letterman	do			
180. <i>Bumelia spinosa</i>	1146	Arizona	Santa Catalina mountains.	C. G. Pringle	Rocky			
	1151	do	do	do	do			
181. <i>Bumelia lycioides</i> <i>Iron Wood. Southern Buckthorn.</i>	333	Tennessee	Nashville	A. Gittinger	Alluvial			
182. <i>Bumelia cuneata</i> <i>Ants' Wood. Downward Plum. Saffron Plum.</i>	503	Florida	Long Key	A. H. Curtiss	Coral			
	1124	do	Boca Chica Key	do	do			
183. <i>Mimusops Sieberi</i> <i>Wild Dill.</i>	458	do	Upper Metacombe Key.	do	do	0.170	12	40
EBENACEÆ.								
184. <i>Diospyros Virginiana</i> <i>Persimmon.</i>	61	Missouri	Allenton	G. W. Letterman	Rich upland	0.158	56	
	425	Tennessee	Nashville	A. Gittinger	Rich loam			
	811	West Virginia	Grafton	C. G. Pringle				
	1084	Missouri	Allenton	G. W. Letterman	Rich upland			
	1162	do	do	do	do			
185. <i>Diospyros Texana</i> <i>Black Persimmon. Mexican Persimmon. Chapote.</i>	936	Texas	Austin	C. Mohr	Calcareous	0.098	37	
STYRACACEÆ.								
186. <i>Symplocos tinctoria</i> <i>Horse Sugar. Sweet Leaf.</i>	347	Alabama	Cottage Hill	do	Sandy	0.134	26	
	560	Arkansas	Texarkana	G. W. Letterman	do	0.123	47	
187. <i>Halesia diptera</i> <i>Snow-drop Tree. Silver-bell Tree.</i>	788	Georgia	Bainbridge	A. H. Curtiss	Low			
	922	Florida	Apalachicola	C. Mohr	Alluvial			

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.9613	0.8887		0.9350	0.84	0.83	0.83			488
0.9737	0.9233		0.9485	0.86	0.81	0.29			500
0.9312	0.8912		0.9112	0.87	0.83	0.85			1194
			0.9316			0.32	58.06		
0.7106	0.7479		0.7293	0.71	0.84	0.78	45.45		746
0.7541	0.7299		0.7420	1.10	1.02	1.06			60
0.5947	0.5997	0.5777	0.5907	1.44	1.28	1.36			930
0.6221	0.6386		0.6304	1.27	1.27	1.27			1083
			0.6544			1.23	40.78		
0.6580	0.6991		0.6786	1.19	1.04	1.11			1146
0.6392	0.6063	{	0.6419	1.32	1.43	1.37			1151
0.6565	0.6658								
			0.6903			1.24	41.15		
0.7487			0.7487	0.81	0.81	0.81	46.53		333
0.7587	0.7465		0.7516	2.29	2.31	2.30			503
0.8420	0.8384		0.8402	1.60	1.43	1.51			1124
			0.7959			1.90	49.60		
1.0525	1.0999	1.1000	1.0838	2.69	2.32	2.61	67.54		456
0.7639	0.7465		0.7552	1.03	1.03	1.03			61
0.8552			0.8552	0.95	1.12	1.04		Female	425
0.7962	0.7864		0.7913	0.83	0.70	0.77			811
0.7790	0.7920		0.7855	0.86	1.02	0.94			1084
0.7710	0.7722	0.7870	0.7687	0.97	1.05	1.01		Wood with black streaks	1162
			0.7908			0.96	49.28		
0.8325	0.8391	0.8964	0.8460	3.34	3.32	3.33	52.72		966
0.5265	0.5598		0.5429	0.72	0.81	0.77			347
0.5079	0.5360		0.5220	0.62	0.57	0.60			509
			0.5325			0.68	32.18		
0.5481	0.5381		0.5431	0.37	0.42	0.50			736
0.5999	0.5957		0.5978	0.35	0.35	0.35			923
			0.5705			0.42	35.55		

FOREST TREES OF NORTH AMERICA.

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
188. <i>Halesia tetraptera</i>	275	Virginia.....	Carter's ferry.....	H. Shriver.....	Rocky.....			
<i>Rattlebox. Snow-drop Tree. Silver-bell Tree. Oatlico Wood.</i>	613	Georgia.....	Altamaha river.....	A. H. Curtiss.....	Clay.....	0.088	17	
OLEACEÆ.								
189. <i>Fraxinus Greggii</i>	1221	Mexico.....	Lampasas mount-ains.	S. B. Buckley.....				
190. <i>Fraxinus anomala</i>	1105	Utah.....	Kane county.....	A. L. Siler.....	Sandy.....			
191. <i>Fraxinus pistaciæfolia</i>	591	Arizona.....	Santa Rita mount-ains.	G. Engelmann and C. S. Sargent.	Low.....			
<i>Ash.</i>	892	Eastern Arizona.	San Francisco mount-ains.	E. L. Greene.....				
192. <i>Fraxinus Americana</i>	25	Massachusetts.....	Acushnet.....	E. A. Dana.....				
<i>White Ash.</i>	391	Missouri.....	Allenton.....	G. W. Letterman.....	Siliceous hills.....			
	392	do.....	do.....	do.....	Alluvial.....			
	393	do.....	do.....	do.....	do.....			
	1141	Michigan.....	Dansville.....	W. J. Beal.....	Clay.....			
	1142	do.....	Hudson.....	do.....				
	1144	do.....	Lansing.....	do.....				
	1145	do.....	Dansville.....	do.....	Clay.....			
	180	South Carolina.....	Bonneau's Depot.....	H. W. Ravenel.....	Wet.....			
	144	Illinois.....	Waukegan.....	Robert Douglas.....	Clay.....			
	170	Ohio.....	Pineo, Daniels & Co.	E. E. Barney.....				
	173	do.....	Farmers' Friend Manufacturing Co.	do.....				
	175	do.....	J. W. Stoddard & Co.	do.....				
	190	do.....	Woodsum Machine Co.	do.....				
	191	do.....	B. E. McSherry & Co.	do.....				
	212	Virginia.....	Wytheville.....	H. Shriver.....				
	2271	Vermont.....	Charlotte.....	C. G. Pringle.....	Gravelly.....			
	2272	do.....	do.....	do.....	Swampy.....			
	2671	Virginia.....	Wytheville.....	H. Shriver.....				
	2672	do.....	do.....	do.....				
	2673	do.....	do.....	do.....				
	431	Tennessee.....	Nashville.....	A. Gattinger.....	Limestone.....			
	551	Alabama.....	Kemper's mill.....	C. Mohr.....	Alluvial.....	0.268	79	
	728	Pennsylvania.....	Williamsport.....	C. G. Pringle.....				
	747	Georgia.....	Bainbridge.....	A. H. Curtiss.....	River-bottom.....	0.260	95	
	1045	Massachusetts.....	Reading.....	J. Robinson.....		0.215	64	
192. <i>Fraxinus Americana, var. Texensis</i>	364	Texas.....	Dallas.....	J. Beverchon.....	Dry, calcareous.....	0.384	69	
	937	do.....	Anstin.....	C. Mohr.....	Rich, calcareous.....			
193. <i>Fraxinus pubescens</i>	139	Michigan.....	Lansing.....	W. J. Beal.....	Poor.....			
<i>Red Ash.</i>	2291	Vermont.....	Charlotte.....	C. G. Pringle.....	Clay.....			
	2292	do.....	do.....	do.....	do.....			

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.5617	0.5480		0.5589	0.38	0.36	0.37			275
0.5569	0.5864		0.5717	0.48	0.38	0.43			613
			0.5628			0.40	35.07		
0.7778	0.8080		0.7904	0.97	0.80	0.93	49.26		1221
0.6708	0.6485		0.6597	0.81	0.88	0.85	41.11	Growth slow.....	1105
0.6336	0.6243	}	0.6625	0.84	0.88	0.71			591
0.7287	0.6656			0.44	0.60				
0.6868	0.7120		0.6994	0.51	0.54	0.53			892
			0.6810			0.62	42.43		
0.5911	0.6683		0.6297	0.29	0.32	0.31			25
0.6253	0.6481		0.6367	0.33	0.34	0.34		Wood from hills not considered valuable.....	39 ¹
0.6223			0.6223	0.51		0.51		All sap-wood.....	39 ²
0.6771			0.6771						39 ³
0.5285	0.4711		0.4988	0.53		0.53			114 ¹
0.7225			0.7225	0.60		0.60			114 ²
0.6506			0.6506	0.38		0.38			114 ⁴
0.6086			0.6086	0.32		0.32			114 ⁵
0.6744	0.7005		0.6875	0.36	0.39	0.38		All sap-wood.....	130
0.7631	0.7600		0.7616	0.45	0.45	0.45		All sap-wood; second growth. Growth rapid.....	144
0.7449			0.7449	0.35	0.37	0.36			170
0.7179	0.7116		0.7148	0.39	0.39	0.39			178
0.6470	0.6473		0.6472	0.81	0.29	0.30			175
0.6383	0.6387		0.6385						190
0.6116			0.6116	0.24	0.24	0.24			191
0.6157	0.6213		0.6185	0.45	0.42	0.44		0.5 sap-wood.....	212
0.6253	0.6326		0.6290	0.30		0.30			227 ¹
0.6489	0.6687		0.6738	0.26		0.26			227 ²
0.6097			0.6097	0.43		0.43		0.5 sap-wood.....	267 ¹
0.6329			0.6329	0.35		0.35		0.5 sap-wood.....	267 ²
0.6763			0.6763	0.44		0.44		All sap-wood.....	267 ³
0.6449			0.6449	0.74	0.69	0.72			431
0.6176	0.6481		0.6329	0.55	0.48	0.52			551
0.6423	0.5944		0.6184	0.37	0.28	0.33			728
0.5599	0.5972		0.5786	0.68	0.71	0.70			747
0.9131	0.8141	0.8016	0.8429	0.47	0.39	0.43			1045
			0.6543			0.42	40.77		
0.8252	0.8530		0.8391	0.56	0.49	0.53			864
0.7095	0.6667		0.6881	0.86	0.87	0.87			987
			0.7636			0.70	47.59		
0.5066	0.6529	0.6779	0.6125	0.26		0.26		Poor quality.....	189
0.6619			0.6619	0.21		0.21			239 ¹
0.6455			0.6455	0.22		0.22			239 ²

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
193. <i>Fraxinus pubescens</i> —continued	229 ^a	Vermont	Charlotte	C. G. Pringle	Clay			
	1059	Massachusetts	Topsfield	J. Robinson	River-bottom			
194. <i>Fraxinus viridis</i> <i>Green Ash.</i>	57	Missouri.....	Allenton.....	G. W. Letterman.....	Rich, wet.....			
	308	Texas	Dallas	J. Reverchon	do	0.512	45	42
	488	Tennessee	Nashville.....	A. Gattinger.....	Rich upland			
	948	Texas	Victoria	C. Mohr.....	Rich, wet.....	0.144	31	
	957	do	Matagorda bay	do	do	0.184	34	
194. <i>Fraxinus viridis</i> , var. <i>Berlandieriana</i> <i>Ash.</i>	1211	do	Austin	S. B. Buckley	do			
195. <i>Fraxinus platycarpa</i> <i>Water Ash.</i>	536	Alabama.....	Stockton.....	C. Mohr	Rich, alluvial	0.090	27	
196. <i>Fraxinus quadrangulata</i> <i>Blue Ash.</i>	66	Missouri.....	Allenton.....	G. W. Letterman	Dry upland	0.138	35	
	125	Michigan	Lansing	W. J. Beal	Rich loam			
	286 ¹	Kentucky	Mercer county.....	W. M. Linney.....	Limestone			
	286 ²	do	do	do	do			
	286 ³	do	do	do	do			
	291	Missouri.....	Allenton.....	G. W. Letterman.....	Sandy loam			
	423	do	do	do	Rich upland			
	518	Tennessee	Nashville.....	A. Gattinger.....	Rich limestone			
197. <i>Fraxinus Oregana</i> <i>Oregon Ash.</i>	964	Oregon	Portland	G. Engelmann and C. S. Sargent.	Low, wet	0.355	46	47
	1001	do	Weidler's saw-mill	do				
	1024	do	Portland Furniture Company.	do				
	1080	do	do	do				
198. <i>Fraxinus sambucifolia</i> <i>Black Ash. Hoop Ash. Ground Ash.</i>	102	Vermont.....	Charlotte	C. G. Pringle.....	Wet, peaty			
	122	Michigan	Dansville.....	W. J. Beal	do			
	147	Illinois	Waukegan	Robert Douglas.....	Low, wet			
	376	Vermont.....	Charlotte	C. G. Pringle.....	Wet, peaty			
	839	Massachusetts.....	Danvers	J. Robinson.....	Rich, loamy	0.236	55	
199. <i>Forestiera acuminata</i> <i>Priest.</i>	616	Georgia.....	Ogeechee river	A. H. Curtiss	Rich, moist	0.075	31	
	787	do	Bainbridge	do		0.150	55	
	955	Texas	Matagorda bay	C. Mohr	Alluvial			
200. <i>Chionanthus Virginica</i> <i>Fringe Tree. Old Man's Beard.</i>	276	Virginia	Carter's ferry.....	H. Shriver.....	Rich, moist			
	751	Florida	Chattahoochee.....	A. H. Curtiss	Clay upland	0.141	47	

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.6459			0.6459	0.20		0.20			229 ^a
0.5957	0.5260		0.5598	0.42	0.41	0.42		All sap-wood.	1039
0.5685	0.5489								
			0.6251			0.26	38.96		
0.6640	0.6701		0.6671	0.47	0.62	0.55		First sp. gr. determination made on 0.5 sap-wood.	57
0.5947	0.6277		0.6112	0.54	0.57	0.56			308
0.7350			0.7350	0.82	0.80	0.81		All sap-wood.	488
0.7728	0.8217	0.7902	0.7949	0.63	0.56	0.60		Second and third sp. gr. determinations made on sap-wood.	948
0.7198	0.7807		0.7503	0.82	0.76	0.79		Second sp. gr. determination made on sap-wood.	987
			0.7117			0.65	44.35		
0.5774	0.5786		0.5780	0.51	0.56	0.54	36.02		1211
0.3567	0.3515		0.3541	0.69	0.76	0.73	22.07		536
0.8246	0.8440		0.8343	0.63	0.67	0.65			66
0.4929			0.4929	0.96		0.96			125
0.7789			0.7789	0.74		0.74			266 ¹
0.8131			0.8131	0.81		0.81		Second growth.	286 ^a
0.7469			0.7469	0.82		0.82		Second growth.	286 ^a
0.5967	0.6086		0.6027	0.88	0.90	0.89			291
0.6897	0.6746		0.6822	0.78	0.74	0.76			423
0.7960			0.7960	0.60	0.61	0.61			518
			0.7184			0.78	46.78		
0.6432	0.5947		0.6189	0.11	0.43	0.27		Second sp. gr. determination made on sap-wood.	964
0.5314	0.6123		0.5719	0.73		0.73			1001
0.4553	0.4786		0.4645	0.12	0.15	0.14		Branch.	1024
0.6317	0.6425		0.6371	0.21	0.24	0.23		Tough.	1030
			0.5731			0.34	35.72		
0.6209	0.6338		0.6274	0.62	0.57	0.60			162
0.5084			0.5084	0.89		0.89			122
0.6859	0.6812		0.6836	0.84	0.88	0.86		Second sp. gr. determination made on 0.125 sap-wood.	147
0.6485	0.5928		0.6207	0.77	0.83	0.80			378
0.7020	0.7465		0.7243	0.46	0.48	0.47		All sap-wood.	839
			0.6318			0.72	39.37		
0.7067	0.7500		0.7284	0.75	0.72	0.74			616
0.5629	0.5681		0.5755	0.65	0.72	0.69			737
0.5988	0.6002		0.5995	0.88	0.60	0.74			955
			0.6345			0.72	39.54		
0.6476	0.4628		0.5552	0.53		0.53			276
0.7150	0.7165	0.7259	0.7191	0.39	0.56	0.48			751
			0.6372			0.51	39.71		

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
201. <i>Oemanthus Americanus</i> <i>Devil Wood.</i>	283	Louisiana.....	Amite.....	C. Mohr.....	Rich, alluvial.....			
	344	Alabama.....	Cottage Hill.....	do.....	Low, rich.....			
	584	Florida.....	Saint John's river.....	A. H. Curtiss.....	Sandy loam.....	0.190	45	19
BORRAGINACEÆ.								
202. <i>Cordia Sebestena</i> <i>Geiger Tree.</i>	1202	do.....	Key West.....	do.....	Coral.....			
	1218	do.....	do.....	do.....	do.....			
203. <i>Cordia Boissieri</i>	1223	Texas.....	Brownsville.....	S. B. Buckley.....	Limestone.....			
204. <i>Bourreria Havanensis</i> <i>Strong Bark.</i>	455	Florida.....	Upper Metacombe Key.....	A. H. Curtiss.....	Coral.....	0.092	45	
	1187	do.....	Key Largo.....	do.....	do.....			
205. <i>Ehretia elliptica</i> <i>Knackaway. Anagua.</i>	827	Texas.....		Department of Agriculture.....				
	942	do.....	New Braunfels.....	C. Mohr.....	Rich, alluvial.....	0.170	25	
BIGNONIACEÆ.								
206. <i>Catalpa bignonioides</i> <i>Catalpa. Catawba. Bean Tree. Ogar Tree. Indian Bean.</i>	167	Ohio.....	Alexandersville.....	S. H. Binkley and E. E. Barney.....	Clayey loam.....			
	540	Alabama.....	Stockton.....	C. Mohr.....	Low, wet.....			
	744	Georgia.....	Bainbridge.....	A. H. Curtiss.....	Clay.....	0.238	8	17
207. <i>Catalpa speciosa</i> <i>Western Catalpa.</i>	38	Missouri.....	Charleston.....	C. S. Sargent.....	Wet clay.....	0.288	8	51
	160	Ohio.....	Dayton.....	E. E. Barney.....	Clay.....			
	166	Illinois.....	Cairo.....	D. Axtell.....	do.....			
	171	Indiana.....	Wabash river.....	E. E. Barney.....	Low, alluvial bottom.....			
	180	Tennessee.....	Obion river.....	E. P. Hynds and E. E. Barney.....	do.....			
	181	Missouri.....	New Madrid.....	E. E. Barney.....	do.....			
	182	Illinois.....	Ullin.....	do.....	do.....			
	183	Missouri.....	New Madrid.....	do.....	do.....			
	184	do.....	do.....	do.....	do.....			
	210	Indiana.....	Vincennes.....	do.....	do.....			
208. <i>Chilopsis saligna</i> <i>Desert Willow.</i>	556	New Mexico.....	Valley of the Upper Gila river.....	E. L. Greene.....	Alluvial.....			
	595	Arizona.....	Tucson.....	G. Engelmann and C. S. Sargent.....	Moist, gravelly.....			
	682	do.....	do.....	do.....	do.....			
209. <i>Crescentia cucurbitina</i> <i>Black Calabash Tree.</i>	1216	Florida.....	Bay Blacayne.....	A. H. Curtiss.....	Coral.....			
VERBENACEÆ.								
210. <i>Citharexylum villosum</i> <i>Fiddle Wood.</i>	483	do.....	do.....	do.....	do.....			

THE WOODS OF THE UNITED STATES.

301

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.8156	0.8652		0.8404	0.68	0.65	0.67		0.5 sap-wood	283
0.8606			0.8606	0.26	0.29	0.28		All sap-wood	344
0.7100	0.7544		0.7322	0.43	0.43	0.43		First sp. gr. determination made on sap-wood; second sp. gr. determination made on 0.75 sap-wood.	584
			0.8111			0.46	50.55		
0.7740			0.7740		4.09	4.09			1202
0.6008	0.6942		0.6475	4.43	4.28	4.35			1218
			0.7108			4.22	44.30		
0.6710	0.6870		0.6790	3.67	3.39	3.53	42.31		1222
0.8280	0.8248		0.8264	3.38	3.20	3.29			455
0.7708	0.7960	0.7960	0.7883	2.28	2.27	2.28			1137
			0.8073			2.79	50.31		
0.6636	0.6548		0.6592	1.23	1.27	1.25			827
0.6274	0.6302		0.6288	1.33	1.41	1.37			942
			0.6440			1.31	40.13		
0.4293	0.4270		0.4282	0.40	0.41	0.41		Cultivated	167
0.4694	0.4979		0.4837	0.26	0.34	0.30			540
0.4426	0.4177		0.4302	0.47	0.37	0.42			744
			0.4474			0.38	27.88		
0.4601	0.4446		0.4524	0.30	0.41	0.36			28
0.4586	0.4467		0.4527	0.34	0.39	0.37		Cultivated	160
0.3897	0.3829		0.3863	0.36	0.34	0.35			166
0.3850	0.3878		0.3864	0.32	0.30	0.31			171
0.4156	0.3783	0.4260	0.4066	0.45	0.50	0.48			180
0.3900			0.3900	0.38	0.42	0.40		From an old fence-rail	181
0.3968			0.3968	0.47	0.43	0.45		From a post 47 years in the ground	182
0.4493			0.4493	0.48	0.37	0.43		From a standing tree killed by earthquake in 1812	183
0.4535			0.4535	0.40	0.34	0.37			184
0.3911	0.3907		0.3909	0.38	0.39	0.39			210
			0.4165			0.39	25.96		
0.6059			0.6059	0.37	0.37	0.37			556
0.5631			0.5631	0.43	0.41	0.42			595
0.6003	0.6026		0.6015	0.32	0.32	0.32			682
			0.5802			0.37	36.78		
0.6270	0.6368		0.6319	1.47	1.23	1.35	30.38		1216
0.8775	0.8644		0.8710	0.49	0.54	0.52	54.28		483

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
211. <i>Avicennia nitida</i> <i>Black Mangrove. Black Tree. Black Wood.</i>	490 886 828	Florida do do	Bay Biscayne..... do do	A. H. Curtiss..... Department of Agriculture. do	Coral..... do do	0.158	39	18
NYCTAGINACEÆ.								
212. <i>Pisonia obtusata</i> <i>Pigeon Wood. Beef Wood. Cork Wood. Pork Wood.</i>	474	do	Upper Metacombe Key.	A. H. Curtiss.....	Coral.....			
POLYGONACEÆ.								
213. <i>Coccoloba floridana</i> <i>Pigeon Plum.</i>	473	do	do	do	do	0.156	48	27
214. <i>Coccoloba uvifera</i> <i>Sea Grape.</i>	453	do	do	do	Sandy	0.214	17	29
LAURACEÆ.								
215. <i>Persea carolinensis</i> <i>Red Bay.</i>	585	do	Saint John's river	do	Sandy loam	0.188	12	45
215. <i>Persea carolinensis</i> , var. <i>palustris</i>	340	Alabama	Mobile county	C. Mohr	Damp, sandy	0.372	119	
216. <i>Nectandra Willdenoviana</i> <i>Lance Wood.</i>	470 1138 1196	Florida do do	Upper Metacombe Key. Bay Biscayne..... Upper Metacombe Key.	A. H. Curtiss..... do do	Coral..... do do			
217. <i>Sassafras officinale</i> <i>Sassafras.</i>	71 387 446 814 854 1163	Missouri do Tennessee West Virginia Massachusetts Missouri	Allenton..... do Nashville..... Grafton..... Danvers..... Allenton.....	G. W. Letterman..... do A. Gittinger..... C. G. Pringle..... J. Robinson..... G. W. Letterman	Low, rich..... Alluvial..... Rich..... Rich loam..... Low, alluvial			
218. <i>Umbellularia Californica</i> <i>Mountain Laurel. California Laurel. Spice Tree. Copiput. California Olive. California Bay Tree.</i>	703 897	Oregon California	Coco bay.....	G. Engelmann and C. S. Sargent. Department of Agriculture.				
EUPHORBIACEÆ.								
219. <i>Drypetes crocea</i> <i>Guiana Plum. White Wood.</i>	468 1185 1196	Florida do do	Upper Metacombe Key. No-Name Key..... Upper Metacombe Key.	A. H. Curtiss..... do do	Coral..... do do			
219. <i>Drypetes crocea</i> , var. <i>latifolia</i>	459 1187	do do	do do	do do	do do	0.232	112	52

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
1.0919	1.0485		1.0702	1.99	1.12	1.56		0.5 sap-wood	490
0.9066	1.0622		0.9794	1.71	1.72	1.71		First sp. gr. determination made on sap-wood; second sp. gr. determination made on 0.5 sap-wood.	826
0.7074	0.9792		0.6918	4.54	4.01	4.27		All sap-wood	828
			0.9138			2.51	59.95		
0.6475	0.6582		0.6529	7.44	7.79	7.62	40.60		474
0.9629	1.0040		0.9835	5.25	4.81	5.03	61.29	0.66 sap-wood	478
0.9149	0.9998	0.9759	0.9635	1.11	1.63	1.37	60.04	First sp. gr. determination made on 0.5 sap-wood; third sp. gr. determination made on 0.25 sap-wood.	453
0.6373	0.6485		0.6429	0.66	0.85	0.76	40.07		586
0.5977	0.6815		0.6396	0.46	0.27	0.37	39.86		340
0.8206	0.7650		0.7928	0.82	0.68	0.75		0.9 sap-wood	470
0.7980	0.8482	}	0.8147	0.52	0.59	0.56		All sap-wood	1128
0.8148	0.7980		0.7005	0.48	0.47	0.48			1196
0.7222	0.6787		0.7003			0.60	47.94		
0.5090	0.5210		0.5120	0.11	0.09	0.10			71
0.4900	0.4828		0.4864	0.08	0.09	0.09			387
0.4542			0.4542	0.04	0.05	0.05			446
0.5296	0.5863		0.5315	0.06	0.07	0.07			814
0.5765		0.5773	0.5769	0.06	0.23	0.15			854
0.4558	0.4319	0.5055	0.4644	0.12	0.15	0.13			1168
			0.5042			0.10	31.42		
0.6326	0.6302		0.6264	0.36	0.27	0.32			708
0.6897	0.6840		0.6769	0.55	0.36	0.46			897
			0.6517			0.39	40.61		
0.9195	0.8448		0.8821	3.58	3.16	3.37		All sap-wood	468
0.8918	0.9119		0.9019	3.19	3.26	3.23			1185
0.9696	0.9682		0.9786	7.07	6.58	6.83			1193
			0.9209			6.14	57.89		
1.0009	0.9890		0.9780	3.60	3.09	3.39			450
0.9048	0.8878		0.8963	7.70	7.70	7.70			1187
			0.9346			3.29	58.24		

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
220. <i>Sebastiania lucida</i> Crab Wood. Poison Wood.	409	Florida	Upper Metacombe Key.	A. H. Curtiss.	Coral	0.116	30	40
	1126	do	do	do	do			
	1206	do	Key Largo	do	do			
221. <i>Hippomane Mancinella</i> Manchineel.	1110	do	Key West	do	do	0.129	35	
URTICACEÆ.								
222. <i>Ulmus crassifolia</i> Cedar Elm.	324	Texas	Dallas	J. Reverchon	Rich loam	0.484	52	76
	929	do	Austin	C. Mohr	do			
223. <i>Ulmus fulva</i> Red Elm. Shippery Elm. Mooses Elm.	30 ¹	Kentucky	Mercer county	W. M. Linney	Limestone			
	30 ²	do	do	do	do			
	30 ⁴	do	do	do	do			
	101	Vermont	Hinesburg	C. G. Pringle	Gravelly loam			
	120	Michigan	Dansville	W. J. Beal	Gravelly			
	184	Missouri	Allenton	G. W. Letterman	Rich, alluvial			
	366	Vermont	Charlotte	C. G. Pringle	Gravelly			
	369	do	do	do	do			
	429	Tennessee	Nashville	A. Gattinger	Clay			
	899	Massachusetts	Boxford	J. Robinson	Rich loam			
224. <i>Ulmus Americana</i> White Elm. American Elm. Water Elm.	19	do	Arnold Arboretum	C. S. Sargent	Drift	0.561	19	28
	202	Ohio	S. M. Brown & Co.	E. E. Barney	Limestone			
	281 ¹	Missouri	Allenton	G. W. Letterman	Alluvial			
	281 ²	do	do	do	do			
	968	Texas	Colorado river	C. Mohr	do	0.230	17	5
	1086	Massachusetts	Danvers	J. Robinson	Gravelly	0.160	7	17
	1049	do	North Reading	do	do	0.215	21	10
225. <i>Ulmus racemosa</i> Rock Elm. Cork Elm. Hickory Elm. White Elm. Olive Elm.	116 ¹	Michigan	Dansville	W. J. Beal	do			
	116 ²	do	Big Rapids	do	Low, gravelly			
	116 ³	do	do	do	do			
	116 ⁴	do	Hudson	do	Alluvial			
	814	do	Hersey	do	Rich loam			
	428	Tennessee	Nashville	A. Gattinger	do			
226. <i>Ulmus alata</i> Wahoo. Winged Elm.	133	South Carolina	Bonneau's Depot	H. W. Ravenel	do			
	380	Tennessee	Davidson county	A. Gattinger	Loam			
	583	Mississippi	Kemper's mill	C. Mohr	Alluvial	0.244	83	38

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
1.1195	1.0855		1.1025	4.96	4.48	4.72		0.25 sap-wood	469
1.0900	1.0765	1.0640	1.0768	1.82	1.96	1.89		0.5 sap-wood	1126
1.0832	1.1010		1.0921	1.65	1.82	1.74			1206
			1.0905			2.78	67.96		
0.5045	0.5600		0.5772	5.20	5.11	5.16	35.97		1110
0.6286	0.7185		0.6736	1.05	1.03	1.04		Ash of a knot, 1.85	224
0.7897	0.7640		0.7754	1.41	1.30	1.36			929
			0.7245			1.20	45.15		
0.8363	0.8358		0.8361	1.80	1.00	1.15			30 ¹
0.7082			0.7082	0.79		0.79			80 ²
0.6499			0.6499	0.60		0.60			80 ⁴
0.5699	0.5538		0.5619	0.86	0.81	0.84			101
0.6769			0.6769	0.82		0.82			120
0.6841	0.6841		0.6841	0.68	0.79	0.74			134
0.6540	0.7589		0.7065	1.24	0.84	1.04			360
0.7330	0.6908		0.6817	0.12	0.11	0.12			869
0.6706			0.6706	1.22	1.31	1.27			429
0.7613	0.7622		0.7623	1.07	0.70	0.88			869
0.7978	0.8180								
			0.6956			0.83	43.35		
0.5462	0.5309		0.5386	0.86	0.82	0.84		Specific gravity of wet sap-wood, 1.026	19
0.7459	0.7287		0.7373	0.40	0.56	0.48		Second sp. gr. determination made on sap-wood	202
0.6221	0.6054	0.6346	0.6207	0.93		0.93			281 ¹
0.6299			0.6299	0.86		0.86			281 ²
0.6495	0.6480	0.6265	0.6413	1.04	0.99	1.02		First sp. gr. determination made on 0.5 sap-wood; second sp. gr. determination made on 0.75 sap-wood; third sp. gr. determination made on sap-wood.	958
0.6382	0.7102	0.6742	0.6742	0.75	1.01	0.88		Third sp. gr. determination made on sap-wood	1036
0.7173	0.7085	0.7116	0.7124	0.75	0.45	0.60		First sp. gr. determination made on 0.25 sap-wood; second and third sp. gr. determinations made on sap-wood.	1049
			0.6506			0.80	40.54		
0.8144			0.8144	0.42		0.42			116 ¹
0.6846			0.6846						116 ²
0.6803	0.6006		0.6400	0.34		0.34			116 ³
0.7345			0.7345	0.79		0.79			116 ⁴
0.7290	0.7570		0.7430	0.59	0.69	0.64			314
0.7414			0.7414	0.90	0.72	0.81			428
			0.7263			0.60	45.26		
0.8710	0.8544		0.8627	1.02	1.12	1.07		All sap-wood	133
0.5979	0.6220		0.6100	1.14	1.17	1.16		0.5 sap-wood	380
0.7632	0.7880		0.7746	0.76	0.72	0.74			533
			0.7491			0.99	46.69		

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
227. <i>Planera aquatica</i>	758	Florida	Chattahoochee	A. H. Curtiss	Rich, alluvial	0.246	38	
	918	do	do	C. Mohr	do			
228. <i>Celtis occidentalis</i>	69	Missouri	Allenton	G. W. Letterman	Limestone	0.108	50	
<i>Sugarberry. Hackberry.</i>	75 ¹	do	do	do	Low, rich			
	75 ²	do	do	do	Alluvial			
	806	Texas	Dallas	J. Reverchon	do			
	375	Tennessee	Davidson county	A. Gattinger	Loam			
	864	Massachusetts	Plum Island	J. Robinson	Sandy		4	30
	878	do	Salem	do	Loam			
	1111	Missouri	Saint Louis	Henry Eggert	Moist loam	0.182	23	
228. <i>Celtis occidentalis</i> , var. <i>reticulata</i>	652	Arizona	Santa Rita mount- ains.	G. Engelmann and C. S. Sargent.	Dry			
<i>Hackberry. Palo Blanco.</i>								
229. <i>Ficus aurea</i>	486	Florida	Bay Biscayne	A. H. Curtiss	Coral			
230. <i>Ficus brevifolia</i>	1204	do	Key Largo	do	do			
231. <i>Ficus pedunculata</i>	508	do	Boca Chica Key	do	do			
<i>Wild Fig. India-rubber Tree.</i>								
232. <i>Morus rubra</i>	132	Missouri	Allenton	G. W. Letterman	Rich loam			
<i>Red Mulberry.</i>	438	Tennessee	Nashville	A. Gattinger	do			
233. <i>Morus microphylla</i>	450	New Mexico	Silver City	E. L. Greene	Moist clay	0.078	7	25
<i>Mexican Mulberry.</i>	1106	Texas	Austin	S. B. Buckley	Limestone	0.068	4	25
234. <i>Maclura aurantiaca</i>	253	do	Dallas	J. Reverchon	Bottom			
<i>Osage Orange. Bois d'Arc.</i>	421	Pennsylvania	West Chester	S. P. Sharples	Clay loam	0.710	8	44
PLATANACEÆ.								
235. <i>Platanus occidentalis</i>	21	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	0.680	35	53
<i>Sycamore. Button Wood. Button-ball Tree. Water Beech.</i>	126	Missouri	Allenton	G. W. Letterman	Rich, alluvial			
	195	Ohio	Miami valley	E. E. Barney	do			
236. <i>Platanus racemosa</i>	686	California	Carmel river	G. R. Vasey	Clay	0.280	20	
<i>Sycamore. Button Wood.</i>								
237. <i>Platanus Wrightii</i>	648	Arizona	Santa Rita mount- ains.	G. Engelmann and C. S. Sargent.	Rich, gravelly			
<i>Sycamore.</i>								
JUGLANDACEÆ.								
238. <i>Juglans cinerea</i>	16	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	0.560	5	50
<i>Butternut. White Walnut.</i>	76 ¹	Missouri	Allenton	G. W. Letterman	Rich loam			
	70 ²	do	do	do	Alluvial			
	123	Michigan	Dansville	W. J. Peck	Gravelly clay			

THE WOODS OF THE UNITED STATES.

307

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.5735	0.5631		0.5683	0.48	0.48	0.48		All sap-wood	758
0.4674	0.5137		0.4906	0.42	0.42	0.42		All sap-wood	918
			0.5294			0.45	32.99		
0.7802	0.7419		0.7611	1.89	1.89	1.89		C. pumila.....	69
0.6858			0.6858	0.74		0.74			751
0.6592			0.6592	0.68		0.68			752
0.6910	0.7637		0.7274	1.70	1.60	1.65			306
0.8384	0.7956		0.8170	0.97	0.90	0.94		C. Mississippiana	375
0.7784	0.7186	0.7385	0.7453	1.05	0.75	0.90			864
0.7099	0.7908	0.7604	0.7837	0.89	1.07	0.98			873
0.6570	0.6547	0.6396	0.6504	0.89	0.94	0.92			1111
			0.7287			1.09	45.41		
0.7400	0.7150		0.7275	1.32	1.12	1.22	45.34		653
0.2537	0.2695		0.2616	5.17	4.88	5.03	16.30		486
0.6321	0.6475		0.6398	4.55	4.18	4.36	39.87		1204
0.4680	0.4798		0.4739	4.83	5.00	4.92	29.53		508
0.6242	0.5993		0.6118	0.77	0.60	0.68			133
0.5679			0.5679	0.69	0.78	0.74			433
			0.5898			0.71	36.76		
0.7169	0.7157		0.7163	0.57	0.54	0.56			450
0.8270	0.8263		0.8266	0.79	0.82	0.81			1106
			0.7715			0.68	48.08		
0.7841	0.7495		0.7668	0.87	0.88	0.88	47.78		258
0.8551	0.7054		0.7803	0.38	0.59	0.49	48.63	Cultivated. First sp. gr. determination made on limb-wood; second sp. gr. determination made on root-wood.	421
			0.7736			0.68	48.21		
0.6060	0.6111		0.6086	0.35	0.32	0.33			21
0.6078	0.5748		0.5911	0.47	0.67	0.57			126
0.4986	0.5091		0.5038	0.40	0.57	0.48			195
			0.5678			0.46	35.38		
0.5151	0.4609		0.4880	1.17	1.05	1.11	30.41		686
0.4783	0.4688		0.4736	1.34	1.36	1.35	29.51	Grown at 4,000 feet altitude.....	648
0.4183	0.4343		0.4263	0.35	0.30	0.33			16
0.4014	0.4005		0.4010	0.40	0.43	0.40			76
0.4633	0.4355		0.4494	0.69		0.69			762
0.3638			0.3638	0.70		0.79			123

FOREST TREES OF NORTH AMERICA.

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
238. <i>Juglans cinerea</i> —continued	145	Illinois	Waukegan	Robert Douglas	Alluvial			
	176	Ohio	Barney & Smith Manufacturing Co.	E. E. Barney				
	393	Michigan	Lansing	W. J. Beal	Gravelly loam			29
	1057	Massachusetts	Topsfield	J. Robinson	Drift	0.152	8	37
239. <i>Juglans nigra</i> <i>Black Walnut.</i>	112	Missouri	Allenton	G. W. Letterman	Alluvial			
	117	Michigan	Dansville	W. J. Beal	Gravelly			
	149	Illinois	Waukegan	Robert Douglas	Loam			
	209	Ohio	Barney & Smith Manufacturing Co.	E. E. Barney				
	318	Michigan	Lansing	W. J. Beal	Loam			
	325	Texas	Dallas	J. Reverchon	Alluvial	0.272	24	84
	407		Charlestown Navy-yard.	S. H. Pook				
	430	Tennessee	Nashville	A. Gattinger	Limestone			
	766	Florida	Aspalaga	A. H. Curtiss	Clay	0.296	8	28
	934	Texas	Austin	C. Mohr	Moist, calcareous			
	951	do	New Braunfels	do	do			
240. <i>Juglans rupestris</i> <i>Walnut.</i>	415	New Mexico	Pinos Altos mountains.	E. L. Greene	Alluvial	0.207	15	32
	672	California	Contra Costa county.	G. R. Vasey	do	0.313	11	18
	1227	Arizona	Santa Catalina mountains.	C. G. Pringle				
241. <i>Carya oliviformis</i> <i>Pecan. Illinois Nut.</i>	323 ¹	Mississippi	Greenville	C. Mohr	Alluvial	0.260	33	48
	323 ²	do	do	do	do			
	326	Texas	Dallas	J. Reverchon	do			
242. <i>Carya alba</i> <i>Shell-bark Hickory. Shag-bark Hickory.</i>	3	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	0.305	25	37
	29 ¹	Kentucky	Danville	W. M. Linney	Shale			
	29 ²	do	do	do	do			
	29 ³	do	do	do	do			
	43	Missouri	Allenton	G. W. Letterman	Upland			
	118 ¹	Michigan	Hudson	W. J. Beal	Clay			
	118 ²	do	do	do	do			
	118 ³	do	Lansing	do	do			
	153	Missouri	Allenton	G. W. Letterman	Rich upland			
	249	Virginia	Wytheville	H. Shriver	Clay			
	422	Missouri	Allenton	G. W. Letterman	Alluvial			
	531	Mississippi	Kemper's mill	C. Mohr	do	0.288	28	81
	539	do	do	do	do	0.368	31	102
	816	West Virginia	Grafton	C. G. Pringle	do			
	1056	Massachusetts	Topsfield	J. Robinson	Rich loam	0.250	53	
	1097	Missouri	Allenton	G. W. Letterman	Alluvial			

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.4394	0.4388		0.4391	0.51		0.51			145
0.3941	0.4023		0.3982	0.44	0.44	0.44			176
0.3379			0.3379	0.43		0.43			398
0.4538	0.4355	0.4098	0.4530	0.41	0.53	0.47			1057
			0.4086			0.51	25.46		
0.5778	0.5881		0.5830	0.63		0.63			112
0.5807			0.5807	0.48		0.48			117
0.5579	0.5659		0.5569	0.12	0.12	0.12			149
0.5025	0.5265		0.5145	0.63	0.49	0.56			209
0.6313	0.6191		0.6252	0.62		0.62			318
0.5608	0.5833		0.5721	1.92	1.99	1.96			325
0.5750			0.5750	0.60	0.69	0.65			407
0.6376			0.6376	0.93	1.05	0.99			430
0.6415	0.6363	}	0.6403	0.78	0.65	0.68		First sp. gr. determination made on sap-wood	766
0.6435	0.6400			0.51	0.77				
0.7830	0.8002		0.7916	1.22	1.64	1.43		First sp. gr. determination made on 0.5 sap-wood	934
0.6609	0.6868		0.6499	0.71	0.57	0.64			951
			0.6115			0.79	38.11		
0.6789	0.6469	0.6753	0.6670	0.57	1.09	0.83		All sap-wood	415
0.6266	0.5611		0.5939	0.85	0.97	0.91			672
0.7258	0.6948		0.7058	1.15	1.42	1.28			1227
			0.6554			1.01	40.85		
0.7890	0.6925		0.7158	0.96	1.18	1.07		Second sp. gr. determination made on 0.5 sap-wood	322 ^a
0.7020	0.6982		0.7001	0.93		0.93		Second sp. gr. determination made on 0.5 sap-wood	322 ^b
0.7314	0.7445		0.7380	1.33	1.46	1.40			826
			0.7180			1.13	44.75		
0.7618	0.7814		0.7716	0.61	0.58	0.60		All sap-wood	8
0.9680			0.9680	0.73		0.73		All sap-wood	29 ^a
0.9205			0.9205	0.86		0.86		All sap-wood	29 ^b
0.9859	1.0080		0.9970	0.65		0.65		All sap-wood	29 ^c
0.8249	0.8260		0.8255	0.87	0.43	0.40			48
0.8028			0.8028	0.28		0.28			118 ^a
0.7974			0.7974	0.66		0.66			118 ^b
0.8019			0.8019	1.14		1.14			118 ^c
0.8084	0.7677		0.7871	0.87	0.66	0.77			152
0.8470	0.7477		0.7974	0.63	0.98	0.78		All sap-wood	249
0.7289	0.7756		0.7523	0.92	0.90	0.91			422
0.8986	0.9370		0.9088	0.79	0.78	0.79			531
0.8729	0.8692		0.8711	0.78	0.88	0.83			539
0.7470	0.7505		0.7523	0.76	0.66	0.71			816
0.8800	0.9035	0.9245	0.9027	0.54	0.69	0.62			1056
0.6835	0.8051		0.7443	0.90	0.90	0.90			1097
			0.8372			0.73	52.17		

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
243. <i>Carya sulcata</i> <i>Big Shell-bark. Bottom Shell-bark.</i>	27	Kentucky	Mercer county	W. M. Linney	Limestone			
	91 ^a	do	do	do	Alluvial			
	91 ^b	do	do	do	do			
	336	Tennessee	Nashville	A. Gattinger	do			
	383	Missouri	Allenton	G. W. Letterman	do			
	391	do	do	do	do			
	1082	do	do	do	do			
	1164	do	do	do	do			
	1165	do	do	do	do			
	1166	do	do	do	do			
	1170	do	do	do	do			
244. <i>Carya tomentosa</i> <i>Mocker Nut. Black Hickory. Bull Nut. Big-bud Hickory. White-heart Hickory. King Nut.</i>	52	do	do	do	Rich upland			
	72	Kentucky	Perryville	W. M. Linney	Utica shale			
	254	Missouri	Allenton	G. W. Letterman	Rich upland			
	289	do	do	do	do			
	348	Alabama	Citronelle	C. Mohr	Sandy	0.200	42	31
245. <i>Carya porcina</i> <i>Pig Nut. Brown Hickory. Black Hickory. Switch-bud Hickory.</i>	6	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	0.317	29	55
	51	Missouri	Allenton	G. W. Letterman	Flinty			
	88	do	do	do	Rich loam	0.159	41	6
	121	Michigan	Danville	W. J. Beal	Gravelly clay			
	288	Missouri	Allenton	G. W. Letterman	Rich loam	0.100	47	13
	442	Tennessee	Nashville	A. Gattinger	Upland			
	538	Alabama	Kemper's mill	C. Mohr	Alluvial	0.354	63	50
	1051	Massachusetts	North Reading	J. Robinson	Drift	0.228	31	42
	1098	Missouri	Allenton	G. W. Letterman	Flinty			
246. <i>Carya amara</i> <i>Bitter Nut. Swamp Hickory.</i>	153	do	do	do	Rich upland			
	401	do	do	do	do			
	838	Massachusetts	Danvers	J. Robinson	Rich loam			
	871	do	do	do	do	0.115	18	19
247. <i>Carya myriaticiformis</i> <i>Nutmeg Hickory.</i>	237	South Carolina	Bonneau's Depot	H. W. Ravenel	Rich, swampy	0.402	7	107
248. <i>Carya aquatica</i> <i>Water Hickory. Swamp Hickory. Bitter Pecan.</i>	129	do	do	do	Swampy			
	862	Mississippi	Vicksburg	C. Mohr		0.277	28	18
	740	Georgia	Bainbridge	A. H. Curtiss	Alluvial			
	917	Florida	Chattahoochee	C. Mohr	do	0.262	28	28

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.8791	0.8767		0.8779	0.72	0.70	0.71			27
0.7527	0.6859		0.7193	1.09		1.09			91 ^a
0.7345			0.7345	1.04		1.04			91 ^b
0.7034			0.7054						336
0.9020	0.8610		0.8815	0.99	0.92	0.96		First sp. gr. determination made on 0.5 sap-wood.	333
0.8512	0.8424		0.8466	0.91	1.24	1.07			391
0.8911	0.9061	0.8831	0.8934	0.83	0.90	0.87		All sap-wood	1083
0.8827	0.8854		0.8840	0.81	0.84	0.83			1164
0.7054	0.6898	0.7414	0.7122	0.73	0.72	0.73			1165
0.8508	0.8432		0.8470	0.90	0.95	0.93		0.5 sap-wood	1106
0.8024	0.7275	0.7369	0.7563	0.84	0.71	0.78			1170
.....			0.8108			0.90	50.53		
0.8524	0.8163		0.8346	0.97	0.84	0.91			52
0.8610	0.8709		0.8660	1.02	1.06	1.04		0.5 sap-wood	72
0.8334	0.8500		0.8417	0.72	1.05	0.89		Second sp. gr. determination made on sap-wood; second growth	254
.....	0.7279		0.7279	1.29	1.65	1.47			289
0.8316	0.8462		0.8389	0.99	1.08	1.01		First sp. gr. determination made on 0.5 sap-wood; second sp. gr. determination made on sap-wood.	348
.....			0.8218			1.06	51.21		
0.8157	0.8139		0.8148	0.60	0.60	0.60			6
0.8827			0.8827	1.25	1.40	1.33			51
0.9189	0.9290		0.9240	0.85	0.85	0.85		Second sp. gr. determination made on sap-wood; second growth	89
0.6803			0.6803	1.58		1.58			121
0.8584	0.8530		0.8542	0.88	0.71	0.77		First sp. gr. determination made on sap-wood; second sp. gr. determination made on 0.9 sap-wood.	268
0.7926			0.7926	0.68	0.70	0.69		All sap-wood	443
0.8530	0.7152		0.7841	0.82	0.86	0.84		First sp. gr. determination made on 0.8 sap-wood; second sp. gr. determination made on sap-wood.	538
0.8842	0.8481	0.8537	0.8620	0.74	0.75	0.75		Third sp. gr. determination made on sap-wood	1061
0.8990	0.7640		0.8315	1.27	1.29	1.28		Second sp. gr. determination made on sap-wood	1098
0.7470	0.7355	0.8913	0.7913	1.15	1.19	1.17		All sap-wood	1168
.....			0.8217			0.99	51.21		
0.7814	0.7830		0.7822	1.00	0.97	0.99			153
0.5927	0.5754		0.5841	0.97	0.93	0.95			401
0.7580	0.7142		0.7336	0.94	1.03	0.99			838
0.9208			0.9208	1.42	0.92	1.17		0.5 sap-wood	871
.....			0.7552			1.03	47.06		
0.7919	0.8112		0.8016	1.07	1.05	1.06	40.96	All sap-wood	237
0.7639	0.7332		0.7486	1.31	2.03	1.67		All sap-wood	129
0.7719	0.7700		0.7710	1.05	1.32	1.19		0.5 sap-wood	363
0.8248	0.8244		0.8246	1.32	1.05	1.19		0.5 sap-wood	746
0.6422	0.6313	0.5821	0.6185	0.91	1.17	1.04		0.5 sap-wood	917
.....			0.7407			1.27	44.16		

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
MYRICACEÆ.								
248. <i>Myrica cerifera</i> Bayberry. Wax Myrtle.	586	Florida	Saint John's river	A. H. Curtiss	Sandy loam	0.198	18	22
250. <i>Myrica Californica</i>	665	California	Santa Cruz	G. Engelmann and C. S. Sargent.	Rich loam			
CUPULIFERÆ.								
251. <i>Quercus alba</i>	8	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	0.500	12	88
White Oak.	26	do	Acushnet	E. A. Dana				
	32 ¹	Kentucky	Mercer county	W. M. Linney	Limestone			
	32 ²	do	Boyle county	do	Slate			
	32 ³	do	do	do	Shale			
	49 ¹	Missouri	Allenton	G. W. Lettorman	Rich upland			
	49 ²	do	do	do	do			
	49 ³	do	do	do	do			
	113 ¹	Michigan	Big Rapids	W. J. Beal	Gravelly	0.374	34	18
	113 ²	do	Dansville	do	Sandy			
	113 ³	do	Hudson	do				
	150	Illinois	Waukegan	Robert Douglas	Gravelly clay			
	160	Ohio	Pineo, Daniels & Co.	E. E. Barney				
	196	do	Barney & Smith Manufacturing Co.	do				
	228 ¹	Vermont	Champlain valley	C. G. Pringle	Clay			
	228 ²	do	Charlotte	do	Gravelly loam			
	228 ³	do	do	do	do			
	238	South Carolina	Bonneau's Depot	H. W. Ravenel	Rich, damp loam			
	250	Virginia	Wytheville	H. Shriver	Clay			
	251	do	do	do		0.225	56	
	250 ¹	do	do	do	Gravelly			
	250 ²	do	do	do	do			
	250 ³	do	do	do	do			
	403	Maryland	Charlestown Navy-yard.	S. H. Pook				
	443	Tennessee	Nashville	A. Gattinger	Rich bottom			
	547	Alabama	Kemper's mill	C. Mohr	Alluvial			
	748	Florida	Chattahoochee	A. H. Curtiss	Clay	0.236	8	48
	749	do	do	do	do	0.248	18	74
	895	Massachusetts		M. C. Beedle				
	1050	do	North Reading	J. Robinson		0.208	7	11
	1257		Charlestown Navy-yard.	S. H. Pook				
252. <i>Quercus lobata</i>	670	California	Redding	G. R. Vasey	Gravelly loam	0.356	15	17
White Oak. Weeping Oak.								
253. <i>Quercus Garryana</i>	985	Oregon	Weidler's saw-mill.	G. Engelmann and C. S. Sargent.				
White Oak.	988	do	Portland	do	Rich loam	0.393	20	82
	1027	do	Portland Furniture Company.	do				
	1029	do	do	do				
254. <i>Quercus obtusiloba</i>	371	Kentucky	Harrodsburg	W. M. Linney	Shale			
Post Oak. Iron Oak.	372	do	do	do	do			

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
{ 0.5689	0.5592 }	0.5711	0.5637	{ 0.54 }	0.51	0.51	35.13		538
0.5728	0.5519 }			{ 0.47 }					
0.6793	0.6613		0.6703	0.33	0.33	0.33	41.77		665
0.7165	0.7068		0.7129	0.37	0.31	0.34			8
0.7970	0.8012		0.7991	0.30	0.25	0.28			26
0.8480	0.8451	0.7965	0.8299	0.45	0.47	0.46			32 ^b
0.6328			0.6328						32 ^a
0.6305			0.6305						36 ^a
0.6945	0.6526		0.6746	0.37	0.43	0.40			49 ^b
0.7826			0.7826	0.45		0.45			49 ^a
0.6829	0.7385		0.7107	0.30		0.30			49 ^a
0.6006			0.6006	0.37		0.37			113 ^b
0.6852	0.7012		0.6932	0.41		0.41			113 ^a
0.7017			0.7017	0.24		0.24			113 ^a
0.7892	0.8735		0.8314	0.47	0.42	0.45			150
0.7672			0.7672	0.21	0.23	0.22			169
0.6203	0.6305		0.6254	0.57	0.65	0.61			196
0.8304			0.8304	0.39		0.39			228 ^b
0.8670			0.8670	0.33		0.33			228 ^a
0.8091			0.8091	0.37		0.37			228 ^a
0.7830	0.7391		0.7611	0.43	0.42	0.43			238
0.7874	0.7419		0.7647	0.35	0.33	0.34			250
0.6948	0.6984		0.6916	0.21	0.25	0.23			251
0.7556			0.7556						259 ^b
0.7069			0.7069	0.30		0.30			259 ^a
0.8056			0.8056	0.37		0.37			259 ^a
0.7390	0.7437		0.7414	0.49	0.44	0.47			403
0.6549			0.6549	1.37	1.64	1.51			443
0.8566	0.7784		0.8180	0.45	0.29	0.37			547
0.7732	0.7562		0.7647	0.47	0.50	0.49		Braeh: rough bark	743
0.8066	0.8228		0.8147	0.34	0.31	0.33		Tough: smooth bark	749
0.7672	0.7734		0.7703	0.33	0.37	0.35		Taken from beam in old court-house at Cambridge, built in 1757.	895
0.8065	0.7584	0.7358	0.7669	0.26	0.22	0.24			1050
			0.8439						1257
			0.7470			0.41	46.35		
0.7849	0.7672	0.6706	0.7409	0.25	0.34	0.30	46.17	Third sp. gr. determination made on sap-wood	670
0.7814	0.8425		0.8120	0.33	0.25	0.29			985
0.6549	0.7800		0.7075	0.36	0.23	0.33			988
0.7478	0.7574	0.7704	0.7585	0.37	0.41	0.39		Braeh	1027
0.6998	0.7369		0.7034	0.62	0.49	0.56		Tough	1029
			0.7453			0.39	46.45		
0.8200	0.8008		0.8104	0.63		0.62		Second sp. gr. determination made on 0.5 sap-wood	37 ^b
0.8765			0.8765	0.74		0.74		0.5 sap-wood	37 ^a

FOREST TREES OF NORTH AMERICA.

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
254. <i>Quercus obtusiloba</i> —continued	373	Kentucky	Harrodsburg	W. M. Linney	Shale			
	151	South Carolina	Bonneau's Depot	H. W. Ravenel	Rich upland			
	256	Missouri	Allenton	G. W. Letterman	Clay			
	351	Alabama	Citronelle	C. Mohr	do	0.244	38	88
	771	Florida	Aspalaga	A. H. Curtiss	Gravelly barrens	0.204	32	115
255. <i>Quercus undulata</i> , var. <i>Gambellii</i>	205	New Mexico	Pinos Altos mount-ains.	E. L. Greene		0.196	18	67
	417	do	do	do				
	525	Colorado	Engelmann's cañon	Robert Douglas	Rocky			
	1150	Arizona	Santa Rita mount-ains.	C. G. Pringle				
256. <i>Quercus macrocarpa</i>	701	Kentucky	Mercer county	W. M. Linney	Alluvial			
	792	do	do	do	do			
	793	do	do	do	do			
	794	do	do	do	do			
	137	Missouri	Allenton	G. W. Letterman	Moist upland			
	143	Illinois	Waukegan	Robert Douglas	Rich			
	168	Ohio	Woodsum Machine Company.	E. E. Barney				
	204	do	Barney & Smith Manufacturing Co.	do				
	310	Texas	Dallas	J. Reverchon	Rich, moist	0.528	13	120
	400	Missouri	Allenton	G. W. Letterman	Alluvial			
	412	Vermont	Charlotte	C. G. Pringle	Clay			
	432	Tennessee	Nashville	A. Gattinger	Alluvial			
	583	Illinois	Waukegan	Robert Douglas				
	831	do	Winnabago county	M. S. Bebb	Loam			
	832	do	do	do	do			
	933	Texas	Austin	C. Mohr	Alluvial			
	1071	Vermont	Charlotte	C. G. Pringle				
	1072	do	do	do				
	1073	do	do	do				
257. <i>Quercus lyrata</i>	424	Tennessee	Nashville	A. Gattinger	Low			
	545	Mississippi	Kemper's mill	C. Mohr	Alluvial	0.339	19	73
	762	Florida	Chattahoochee	A. H. Curtiss	do	0.266	20	70
	953	Texas	Matagorda bay	C. Mohr	Loam			
258. <i>Quercus bicolor</i>	12	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	0.305	8	60
	541	Missouri	Allenton	G. W. Letterman	Alluvial			
	542	do	do	do	do			
	543	do	do	do	do			
	846	Massachusetts	West Newbury	J. Robinson	Low, swampy	0.206	5	30

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.8459			0.8459	1.56		1.56			37 ²
0.8920	0.8882	0.9170	0.8991	0.79	0.79	0.79			151
0.8522	0.8196		0.8359	0.46	0.52	0.49			256
0.7888	0.7516		0.7702	0.52	0.53	0.53			351
0.8548	0.7942	0.8072	0.8187	0.63	0.88	0.76			771
			0.8367			0.79	52.14		
0.8382	0.8562		0.8472						295
0.8680	0.7748	0.7862	0.8097	0.73	0.79	0.79		First sp. gr. determination made on 0.75 sap-wood.	417
0.7684	0.7894		0.7789	1.03	1.37	1.23			525
0.9487	0.9052		0.9270	1.10	0.80	0.95			1150
			0.8407			0.99	52.39		
0.7070			0.7070	0.96		0.96			79 ¹
0.7205			0.7205	0.66		0.66			79 ²
0.7275	0.7275		0.7275	0.70		0.70			79 ³
0.6426	0.6300		0.6363	1.18		1.18			79 ⁴
0.8469			0.8469	0.41	0.37	0.39		Second growth.	137
0.8280	0.8276		0.8278	0.60	0.93	0.76			143
0.6440	0.6388		0.6416	0.31	0.32	0.32			168
0.6458	0.6177		0.6318	0.37	0.48	0.40			204
0.7722	0.8166		0.7944	1.08	1.03	1.06			310
0.6183	0.6411		0.6297	0.60	0.60	0.60			400
0.8630	0.8493		0.8563	0.26	0.27	0.27			412
0.7271			0.7271	0.74	0.75	0.75			432
0.9229	0.9543	0.9390	0.9387	1.18	1.46	1.32			583
0.6842	0.7112		0.6977	0.78	0.70	0.74		From fence-post 6 years in the ground.	831
0.6107	0.7075		0.6591	0.96	0.93	0.96		Old fence-post.	832
0.7605	0.8402	0.8047	0.8018	1.11	1.18	1.15			923
0.7740	0.8087	0.7360	0.7729	0.45	0.46	0.46			1071
0.7845	0.7924	0.8018	0.7929	0.32	0.45	0.39			1073
0.7598	0.7630	0.7317	0.7515	0.44	0.35	0.40			1078
			0.7453			0.71	46.45		
0.8060	0.7863		0.7962	0.59	0.56	0.58			424
0.7802	0.7502		0.7652	0.22	0.23	0.23			545
0.8050	0.7649		0.7850	0.23	0.18	0.21			762
0.9790			0.9790	1.72	1.44	1.58			953
			0.8313			0.65	51.61		
0.7562	0.7315		0.7434	0.26	0.28	0.27			12
0.8616	0.8628		0.8622	0.39		0.39			54 ¹
0.6833			0.6833	0.96		0.96			54 ²
0.7114			0.7114	0.96		0.96			54 ³
0.8494	0.8196	0.8330	0.8307	0.27	0.31	0.29			846
			0.7662			0.58	47.75		

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
259. <i>Quercus Michauxii</i> Basket Oak. Cow Oak.	240	South Carolina	Bonneau's Depot....	H. W. Ravenel	Alluvial			
	524	Alabama	Kemper's mill	C. Mohr	do	0.322	22	69
	755	Florida	Chattahoochee	A. H. Curtiss	do	0.260	12	32
260. <i>Quercus Prinus</i> Chestnut Oak. Rock Chestnut Oak.	311	Kentucky	Boyle county	W. M. Linney	Shale			
	312	do	do	do	do			
	313	do	do	do	do			
	35	do	do	do	Limestone			
	484	Tennessee	Nashville	A. Gattinger	Rocky upland			
	925	Alabama	Cullman	C. Mohr	Dry, rocky	0.436	34	84
261. <i>Quercus prinoides</i> Yellow Oak. Chestnut Oak. Chinquapin Oak.	28	Kentucky	Harrodsburg	W. M. Linney				
	341	do	Mercer county	do	Limestone			
	342	do	Boyle county	do	Waverly shale			
	343	do	Mercer county	do	Utica shale			
	58	Missouri	Allenton	G. W. Letterman	Poor, hilly			
	273	do	do	do	Limestone			
	287	do	do	do	Flinty			
	323	Texas	Dallas	J. Reverchon	Calcareous	0.226	24	35
	514	Tennessee	Nashville	A. Gattinger	Alluvial			
	588	Texas	Dallas	J. Reverchon	do			
	856	Massachusetts	Boxford	J. Robinson	Damp	0.364	10	123
262. <i>Quercus Douglasii</i> Mountain White Oak. Blue Oak.	688	California	Contra Costa county	G. R. Vasey	Clay	0.338	59	
263. <i>Quercus oblongifolia</i> White Oak.	601	do	San Diego county	do	Dry, gravelly			
	700	do	San Gabriel	G. Engelmann	do	0.202	16	5
264. <i>Quercus grisea</i> White Oak.	419	New Mexico	Silver City	E. L. Greene	Dry, rocky	0.212	38	40
	696	Arizona	Santa Rita mountains.	G. Engelmann and C. S. Sargent	do			
	1145	do	do	C. G. Pringle	do			
265. <i>Quercus reticulata</i>	1148	do	do	do	do			
266. <i>Quercus Durandii</i>	985	Texas	Austin	C. Mohr	Damp, calcareous	0.164	24	39
	1103	do	do	S. B. Buckley	do			
267. <i>Quercus virens</i> Live Oak.	404	Florida	Charlestown Navy-yard.	S. H. Pook				
	799	do	Saint John's river	A. H. Curtiss	Sandy	0.238	39	15

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.7674	0.8312		0.7993	0.33	0.48	0.40			240
0.8253	0.7704		0.7979	0.57	0.57	0.57			524
0.8200	0.8090		0.8145	0.38	0.35	0.37			755
			0.8039			0.45	50.10		
0.7126	0.7102		0.7114	0.36		0.36			81 ¹
0.6549	0.6659		0.6604	0.33		0.33			81 ¹
0.7726			0.7726						81 ¹
0.8308			0.8308	0.70	0.87	0.79			85
0.6870			0.6870	1.94	1.93	1.94			434
0.8550	0.8185		0.8368	0.40	0.49	0.45			925
			0.7409			0.77	46.73		
0.8401	0.8419		0.8410	1.50	1.40	1.45			28
0.7951			0.7951	1.49		1.49			84 ¹
0.8712			0.8712	0.39	0.49	0.44			34 ¹
0.9643			0.9643	1.62		1.62			34 ¹
0.6566	0.6525		0.6546	1.39	1.25	1.32			58
0.9237	0.8548		0.8893	1.16	1.31	1.24			273
0.8248	0.8644		0.8446	0.64	0.57	0.61			287
1.0240	1.0728		1.0484	1.25	1.09	1.17			823
0.8592	0.8750		0.8671	1.43	1.49	1.46			514
0.9183	0.9405		0.9294	1.38	1.23	1.30			563
0.7652	0.7458	0.7706	0.7605	0.43	0.43	0.43			856
			0.8605			1.14	53.63		
0.9058	0.7898		0.8928	0.80	0.88	0.84	55.64	Second sp. gr. determination made on sap-wood	688
1.0078	1.0790	1.0169	1.0486	4.02	4.69	4.07		First and second sp. gr. determinations made on sap-wood	601
0.9495	1.1500	1.0882		2.28	4.68				
0.8420	0.8374		0.8397	0.99	1.31	1.15		All sap-wood	700
			0.9441			2.61	58.84		
0.9171	0.9835	0.8763	0.9236	1.22	1.26	1.24		0.5 sap-wood	419
0.9897			0.9897	2.57	3.33	2.95		0.5 sap-wood	698
1.1340	1.0945		1.1143	1.32	1.19	1.26			1145
			1.0092			1.82	62.89		
0.9430	0.9528		0.9479	0.51	0.53	0.52	59.07		1148
0.9762	0.8000		0.8775	1.71		1.75		0.5 sap-wood	935
0.8648	0.8690			1.80					
0.9640	1.0835		1.0238	1.82		1.82			1102
			0.9507			1.78	59.25		
1.0350			1.0350						404
0.9005	0.8949		0.8977	1.23	1.21	1.22			794

FOREST TREES OF NORTH AMERICA.

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
267. <i>Quercus virens</i> —continued	919	Alabama.....	Mobile county	C. Mohr	Rich, sandy			
	954	Texas.....	Matagorda bay	do	Sandy loam			
268. <i>Quercus chrysolepis</i>	574	Oregon	Saw-mill, Ashland	G. Engelmann and				
	649	California	San Bernardino	C. S. Sargent. W. G. Wright				
	653	do	Marin county	G. R. Vasey	Gravelly.....			
269. <i>Quercus Emoryi</i>	594	Arizona	Santa Rita mount- ains.....	G. Engelmann and C. S. Sargent.....	Dry, rocky.....			
270. <i>Quercus agrifolia</i>	663	California.....	Marin county.....	G. R. Vasey	Loam	0.247		
271. <i>Quercus Wislizeni</i>	677	do	Auburn.....	G. Engelmann				
272. <i>Quercus rubra</i> Red Oak. Black Oak.	7	Massachusetts.....	Arnold Arboretum	C. S. Sargent	Drift	0.534	8	73
	45 ¹	Kentucky	Mercer county.....	W. M. Linney	Shale			
	45 ²	do	do	do	do			
	89	Missouri.....	Allenton	G. W. Letterman	Rich loam	0.149	5	40
	92 ¹	Kentucky	Mercer county.....	W. M. Linney	Alluvial			
	92 ²	do	do	do	Limestone			
	140	Michigan	Dansville	W. J. Beal.....	Sandy			
	141	do	do	do	do			
	146	Illinois	Waukegan	Robert Douglas	Gravelly.....			
	197	Ohio	Barney & Smith Manufacturing Co	E. E. Barney				
	215	Vermont.....	Charlotte	C. G. Pringle	Gravelly.....			
	216	do	do	do	do			
	217	do	do	do	do			
	218	do	do	do	do			
	553	Alabama	Kemper's mill	C. Mohr	Alluvial	0.368	38	97
	806	Massachusetts.....	Danvers	J. Robinson	Drift			
	920	Mississippi	Enterprise	C. Mohr	Alluvial			
	1043	Massachusetts	North Reading	J. Robinson	Drift	0.230	6	27
272. <i>Quercus rubra</i> , var. <i>Texana</i>	981	Texas	Austin	C. Mohr	Calcareous.....	0.118	13	20
273. <i>Quercus coccinea</i> Scarlet Oak.	23	Massachusetts.....	Hingham	T. T. Bouvè	Light, sandy			
	752	Florida	Aspalaga	A. H. Curtiss	Clay	0.230	15	
274. <i>Quercus tinctoria</i>	17	Massachusetts.....	Arnold Arboretum	C. S. Sargent	Drift	0.630	8	74
	36 ¹	Kentucky	Danville Junction	W. M. Linney	Shale			
	36 ²	do	do	do	do			

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.8225	0.9330		0.8778	1.25	1.80	1.27			919
1.0090	0.9709		0.9900	1.09	0.76	0.98		0.33 sap-wood.....	954
			0.9501			1.14	59.21		
0.7888	0.7592		0.7740	0.80	0.63	0.71			574
0.8830	0.8827		0.8829	0.88	0.84	0.84			649
0.8787	0.9083		0.8910	0.73	0.79	0.76			658
			0.8498			0.60	52.98		
{ 0.9975	{ 0.9019		0.9263	{ 1.86	{ 2.19	2.96	57.73	Second sp. gr. determination made on sap-wood.....	594
{ 0.9201	{ 0.8855			{ 2.61	{ 2.78				
0.8290	0.8216		0.8253	1.21	1.84	1.28	51.43		668
{ 0.7956	{ 0.7788		0.7855	{ 0.94	{ 0.98	1.02	48.95		677
{ 0.8018	{ 0.7657			{ 0.96	{ 1.18				
0.6881	0.6440		0.6411	0.14	0.10	0.12			7
0.5186	0.5910		0.5548	0.46		0.46			451
0.5169	0.6251		0.5710	0.48		0.48			459
0.7480			0.7480	0.27		0.27			89
0.5899			0.5899	0.47	0.47	0.47			921
0.7516	0.7512	0.7516	0.7515	0.15		0.15			924
0.6410			0.6410	0.20	0.21	0.21			140
0.5932			0.5952	0.23	0.24	0.24			141
0.7481	0.7514		0.7498	0.22	0.81	0.27			146
0.6516	0.6615		0.6566	0.16	0.17	0.17			197
0.6423	0.6997		0.6710	0.27	0.22	0.25			215
0.6897	0.7090		0.6994	0.20	0.24	0.22			216
0.6869	0.6867		0.6768	0.20	0.26	0.23		Second growth from stump.....	217
0.6765	0.7029		0.6897	0.38	0.83	0.33			218
0.6389	0.6589		0.6489	0.37	0.34	0.36			553
0.6636	0.6252	0.6578	0.6487	0.07	0.14	0.11			866
0.5244	0.5904		0.5424	0.25	0.21	0.23			920
0.7130	0.6806	0.6920	0.6952	0.14	0.19	0.17			1043
			0.6540			0.26	40.75		
0.9142	0.9018		0.9080	0.93	0.76	0.85	56.59	0.5 sap-wood.....	931
0.7111	0.7079		0.7095	0.11	0.16	0.14			28
0.7667	0.7762		0.7715	0.22	0.28	0.25			782
			0.7405			0.19	46.15		
0.7384	0.7305		0.7335	0.13	0.08	0.11			17
0.6863			0.6863	0.21		0.21			361
0.7215			0.7205	0.31		0.31			367

FOREST TREES OF NORTH AMERICA.

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
274. <i>Quercus tinctoria</i> —continued.....	363	Kentucky	Danville Junction ..	W. M. Linney	Slate			
	41	Missouri	Allenton	G. W. Letterman ..	Hilly			
	74	do	do	do	Rich upland	0.165	6	18
	88	do	do	do	do			
	244	Virginia	Wytheville	H. Shriver	Clay			
	247	do	do	do	do			
	437	Tennessee	Nashville	A. Gattinger	do			
	921	Alabama	Cullman	C. Mohr	Sandy			
275. <i>Quercus Kelloggii</i>	623	Oregon	Saw-mill, Ashland ..	G. Engelmann and C. S. Sargent.				
	963	do	Eugene City	G. H. Collier		0.224	17	78
276. <i>Quercus nigra</i>	268	Missouri	Allenton	G. W. Letterman ..	Clay			
	339	Alabama	Citronelle	C. Mohr	Sandy	0.123	13	46
277. <i>Quercus falcata</i>	131	South Carolina ..	Bonneau's Depot ..	H. W. Ravenel	Rich loam			
	245	Virginia	Wytheville	H. Shriver	Clay			
	265 ¹	do	Carroll county	do	do			
	265 ²	do	do	do	do			
	265 ³	do	do	do	do			
	548	Mississippi	Kemper's mill	C. Mohr	Rich loam			
278. <i>Quercus Catesbaei</i>	342	Alabama	Cottage Hill	do	Barren, sandy			
	770	Florida	Aspalaga	A. H. Curtiss	do	0.301	62	
279. <i>Quercus palustris</i>	47	Missouri	Allenton	G. W. Letterman ..	Rich, alluvial			
	282	do	do	do	do			
280. <i>Quercus aquatica</i>	349	Alabama	Cottage Hill	C. Mohr	Sandy loam	0.350	8	16
	511	Tennessee	Tallahoma	A. Gattinger	do			
	742	Georgia	Bainbridge	A. H. Curtiss	Alluvial	0.310	13	23
281. <i>Quercus laurifolia</i>	756	Florida	Saint John's river ..	do	Sandy loam			
	801	do	do	do	do	0.240	33	26
282. <i>Quercus heterophylla</i>	1171	New Jersey	Mount Holly	S. P. Sharples	Clay	0.329	6	19
283. <i>Quercus cinerea</i>	852	Alabama	Citronelle	C. Mohr	Pine-barren			

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.7749			0.7749	0.18		0.18			365
0.7596	0.7852		0.7474	0.17	0.14	0.16			41
0.8014			0.8014	0.22	0.19	0.21		0.33 sap-wood; second growth	74
0.7192			0.7192	0.28	0.18	0.23			86
0.5070	0.5175		0.5123	0.44	0.68	0.54			244
0.7164	0.6586		0.6875	0.14	0.15	0.15			247
0.6765			0.6765	0.85	0.87	0.86			487
0.6622	0.6972	0.7100	0.6898	0.59	0.70	0.65			921
			0.7045			0.28	43.90		
0.6565	0.6582		0.6573	0.18	0.21	0.20			628
0.6273	0.6322		0.6297	0.33	0.30	0.32			968
			0.6435			0.26	40.10		
0.7304	0.7492		0.7398	0.96	0.92	0.94			268
0.7309	0.7192		0.7251	1.41	1.32	1.37		All sap-wood	339
			0.7324			1.16	45.64		
0.6892	0.7003		0.6948	0.27	0.27	0.27			181
0.5701	0.6780		0.6241	0.27	0.33	0.30			245
0.6613			0.6613	0.15		0.15			2651
0.7334			0.7334	0.22		0.22			2652
0.6556			0.6556	0.25		0.25			2652
0.8052	0.7696		0.7874	0.29	0.28	0.29			548
			0.6928			0.25	43.17		
0.6679	0.6736		0.6708	0.85	0.80	0.83			342
0.7906	0.7854		0.7880	0.90	0.90	0.90			770
			0.7294			0.87	45.45		
0.6917	0.6465		0.6691	0.65	0.57	0.61			47
0.7480	0.6890		0.7185	0.92	1.07	1.00		Second growth	282
			0.6938			0.81	43.24		
0.7167	0.7149		0.7158	0.45	0.39	0.42			349
0.7084	0.7202		0.7143	0.82	0.84	0.83			511
0.7237	0.7624		0.7431	0.35	0.19	0.27			742
			0.7244			0.51	45.14		
0.7474	0.7075		0.7275	0.48	0.46	0.47		All sap-wood	756
0.8146	0.7995		0.8071	0.98	1.41	1.17			801
			0.7673			0.82	47.82		
0.6818	0.6834		0.6824	0.19	0.14	0.17	42.59		1171
0.6882	0.6802								
0.6502	0.6337		0.6420	1.27	1.15	1.21	40.00		352

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
284. <i>Quercus hypoleuca</i>	416	New Mexico.....	Pinos Altos mount- ains.	E. L. Greene	Dry, rocky.....	0.203	49	20
	590	Arizona	Santa Rita mount- ains.	G. Engelmann and C. S. Sargent.	do			
285. <i>Quercus imbricaria</i>	401	Kentucky	Harrodsburg	W. M. Linney.....	Utica shale			
	402	do	do	do	do			
	403	do	do	do	do			
	50	Missouri.....	Allenton.....	G. W. Letterman.....	Rich, moist			
	135	do	do	do	Rich loam			
286. <i>Quercus Phellos</i>	512	Tennessee	Tullahoma.....	A. Gattinger.....	Moist, siliceous...	0.184	15	
<i>Willow Oak. Peach Oak.</i>								
287. <i>Quercus densiflora</i>	687	California	Marin county.....	G. R. Vasey.....	Gravelly.....	0.400	25	85
<i>Tender Oak. Chestnut Oak. Peach Oak.</i>								
288. <i>Castanopsis chrysophylla</i>	729	do	Mendocino county	A. Kellogg				
<i>Chinquapin.</i>								
289. <i>Castanea pumila</i>	573	Arkansas	Hot Springs	G. W. Letterman.....	Sandy loam	0.615		
<i>Chinquapin.</i>								
290. <i>Castanea vulgaris</i> , var. <i>Americana</i>	18	Massachusetts....	Arnold Arboretum	C. S. Sargent.....	Drift	0.666	34	42
	2581	Virginia	Fancy Gap	H. Shriver	Moist.....			
	2582	do	do	do	do			
	2583	do	do	do	do			
	516	Tennessee	Nashville.....	A. Gattinger.....	Sandy			
	727	Pennsylvania.....	Williamsport.....	C. G. Fringle.....				
	863	Massachusetts....	Danvers	J. Robinson.....	Loam	0.110	4	21
291. <i>Fagus ferruginea</i>	9	do	Arnold Arboretum	C. S. Sargent.....	Drift	0.190	9	26
	442	Kentucky	Mercer county.....	W. M. Linney.....	Hudson River shale.			
	443	do	do	do	do			
	553	do	do	do	do			
	554	do	do	do	do			
	119	Michigan	Dansville	W. J. Beal	Gravelly.....			
	765	Florida	Chattahoochee.....	A. H. Curtiss	do	0.273	62	
	853	Massachusetts....	Hamilton	J. Robinson.....	do			
292. <i>Ostrya Virginica</i>	11	do	Arnold Arboretum	C. S. Sargent.....	Drift	0.285	26	44
	87	Missouri.....	Allenton	G. W. Letterman.....	Rich loam	0.085	35	
	870	Massachusetts....	Danvers	J. Robinson.....	Rocky			
	877	do	do	do	Rich loam			
	1047	do	North Reading	do		0.190	62	14
293. <i>Carpinus Caroliniana</i>	46	Missouri.....	Allenton	G. W. Letterman.....	Damp, alluvial...			
	731	Kentucky	Mercer county.....	W. M. Linney.....	Trenton limestone.			
	732	do	do	do	do			

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.7828	0.8304		0.8065	1.27	1.27	1.27		All sap-wood	416
{ 0.7774			0.7952	{ 1.21	1.35	1.41		0.5 sap-wood	509
{ 0.8288	0.7785	0.8089		{ 1.64	1.42				
.....			0.8009			1.34	49.91		
0.7402	0.7394		0.7398	0.23		0.23			40 ¹
0.7284			0.7284	0.65		0.65			40 ²
0.8332			0.8332	0.79		0.79			40 ³
0.7517	0.7380		0.7439	0.23	0.34	0.28			50
0.7200	0.7285		0.7243	0.23	0.18	0.21		Second growth	185
.....			0.7529			0.43	46.92		
0.7582	0.7412		0.7472	0.51	0.48	0.50	46.56		512
0.6850	0.7004		0.6827	1.55	1.41	1.49	42.55		687
0.5520	0.5627		0.5574	0.35		0.35	34.74	0.33 sap-wood	729
0.5698	0.6076		0.5887	0.12	0.11	0.12	30.69		573
0.3829	0.3827		0.3828	0.10	0.12	0.11			18
0.4720			0.4720	0.26		0.26			258 ¹
0.4716			0.4716	0.21		0.21			258 ²
0.4494			0.4494						258 ³
0.4693			0.4693	0.18	0.21	0.20			516
0.4693	0.4347		0.4455	0.16	0.17	0.17			727
0.4513	0.4594	0.4656	0.4621	0.14	0.12	0.13			868
.....			0.4504			0.18	28.07		
0.6068	0.6556		0.6610	0.34	0.23	0.34			9
0.6200			0.6200	0.75		0.75		Red beech	44 ²
0.6343			0.6343	0.70		0.70		Red beech	44 ³
0.7992			0.7992	0.57		0.57		White beech	58 ²
0.7150	0.7300		0.7175	0.54		0.54		White beech	58 ⁴
0.7619	0.6905		0.7112	0.31	0.28	0.25			119
0.6897	0.6590		0.6729	0.47		0.47			765
0.7017	0.6825	0.6870	0.6904	0.29	0.28	0.34		First and second sp. gr. determinations made on 0.5 sap-wood; third sp. gr. determination made on sap-wood.	853
.....			0.6883			0.51	42.89		
0.7608	0.7636		0.7622	0.34	0.37	0.36			11
0.8919			0.8919	0.49	0.67	0.58			87
0.8512	0.8402	0.8370	0.8595	0.53	0.66	0.60			870
0.8491	0.8296	0.8534	0.8440	0.55	0.51	0.53			877
0.7818	0.7940	0.7768	0.7842	0.51	0.39	0.45			1047
.....			0.8284			0.50	51.62		
0.7657	0.7711		0.7684	0.76	0.80	0.78			46
0.7264			0.7264	0.80	0.79	0.80			73 ¹
0.7526	0.7504		0.7515	1.34		1.34			73 ²

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
293. <i>Carpinus Caroliniana</i> —continued	77	Missouri.....	Allenton.....	G. W. Letterman.....	Rich, alluvial	0.052	22	
	872	Massachusetts.....	Danvers	J. Robinson.....	Low, rich			
	1038	do	do	do	Gravelly.....			
BETULACEÆ.								
294. <i>Betula alba</i> , var. <i>populifolia</i>	10	do	Arnold Arboretum..	C. S. Sargent.....	Drift	0.170	10	26
White Birch. Old-field Birch. Gray Birch.	848	do	Danvers	J. Robinson.....	Gravelly.....	0.196	16	33
295. <i>Betula papyrifera</i>	223	Vermont.....	Charlotte	C. G. Pringle	Gravelly.....			
Ounce Birch. White Birch. Paper Birch.	224	do	do	do	do			
	225	do	do	do	do			
	722	Montana.....	Missoula.....	Sereno Watson	Wet			
	836	Massachusetts.....	Townsend	J. Robinson.....		0.234	26	
	990	Alaska.....	Chilcoot inlet.....	Paul Schultze.....		0.188	57	
	1065	Vermont.....	Charlotte	C. G. Pringle.....				
	1066	do	do	do				
	1067	do	do	do				
296. <i>Betula occidentalis</i>	528	Colorado.....	Engelmann's cañon.	Robert Douglas.....	Wet, sandy			
Black Birch.	629	California	Strawberry valley ..	G. Engelmann and C. S. Sargent.	Wet, peaty			
297. <i>Betula lutea</i>	155	Vermont.....	Charlotte	C. G. Pringle.....	Swampy			
Yellow Birch. Gray Birch.	230 ¹	do	do	do	Clay			
	230 ²	do	do	do	Gravelly.....			
	843	Massachusetts.....	Danvers	J. Robinson.....	do	0.160	34	
	1068	Vermont.....	Charlotte	C. G. Pringle.....	do			
	1069	do	do	do	do			
	1070	do	do	do	do			
298. <i>Betula nigra</i>	136	Missouri.....	Allenton.....	G. W. Letterman ..	Moist loam			
Red Birch. River Birch.	398	do	do	do	Alluvial			
	841	Massachusetts.....	North Andover	J. Robinson.....	do	0.192	30	
	842	do	do	do	do	0.214	32	8
	1184	Missouri.....	Allenton.....	G. W. Letterman ..	Sandy loam			
299. <i>Betula lenta</i>	4	Massachusetts.....	Arnold Arboretum..	C. S. Sargent.....	Drift	0.362	12	61
Cherry Birch. Black Birch. Sweet Birch. Mahogany Birch.	221	Vermont.....	Charlotte	C. G. Pringle.....	Gravelly.....			
	844	Massachusetts.....	Danvers	J. Robinson.....	do	0.118	41	22

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.7591	0.6907		0.7249	0.60	0.52	0.56			77
0.7014	0.7085	0.7187	0.7079	0.88	0.93	0.91			872
0.7080	0.6804	0.6896	0.6927	0.55	0.65	0.60			1088
			0.7286			0.83	45.41		
0.5961			0.5961	0.82	0.27	0.30		Sap-wood.....	19
0.6330	0.6088	0.6061	0.6160	0.28	0.28	0.28		First and third sp. gr. determinations made on sap-wood.....	868
			0.5760			0.29	35.90		
2.4594	0.4757		0.4676	0.24	0.24	0.24		All sap-wood.....	222
0.6021	0.5979		0.6000	0.30	0.31	0.31		0.5 sap-wood.....	224
0.6886	0.6207		0.6297	0.25	0.22	0.24		First sp. gr. determination made on 0.5 sap-wood; second sp. gr. determination made on 0.75 sap-wood.	225
0.6240	0.6096		0.6168	0.25	0.22	0.24		All sap-wood.....	722
0.6290	0.6280	0.6396	0.6315	0.20	0.19	0.20		All sap-wood.....	826
0.6081	0.6060		0.6070	0.24	0.25	0.25		All sap-wood.....	899
0.6801	0.6227	0.6112	0.6380	0.28	0.26	0.27		All sap-wood.....	1085
0.6002	0.5965	0.5843	0.5887	0.25	0.24	0.25		All sap-wood.....	1086
0.5894	0.5908	0.5745	0.5849	0.26	0.22	0.24		All sap-wood.....	1087
			0.5965			0.25	37.11		
0.5880	0.6183		0.6032	0.18	0.18	0.18		All sap-wood.....	525
0.6543	0.5777	0.5767	0.6028	0.44	0.37	0.41			629
			0.6080			0.30	37.58		
0.5759	0.6237		0.5998	0.29	0.20	0.25		0.5 sap-wood.....	155
0.7047			0.7047	0.33		0.33		0.875 sap-wood.....	239
0.6416			0.6416	0.60		0.60			239
0.7032	0.7200	0.7080	0.7104	0.16	0.25	0.20		All sap-wood.....	842
0.6468	0.6293	0.6573	0.6445	0.28	0.25	0.27			1089
0.6380	0.6472	0.6256	0.6369	0.28	0.23	0.25			1090
0.6557	0.6543	0.6382	0.6494	0.27	0.27	0.27			1079
			0.6563			0.31	40.84		
0.5608	0.5721		0.5665	0.30	0.31	0.31		All sap-wood.....	128
0.5554	0.5508		0.5531	0.27	0.31	0.29		All sap-wood.....	898
0.5910	0.5759	0.5811	0.5827	0.38	0.38	0.38		From butt: Sap-wood.....	841
0.6049	0.5975	0.5965	0.5996	0.38	0.38	0.38		From top of trunk: Sap-wood.....	842
0.5556	0.5836	0.5983	0.5793	0.44	0.40	0.42			1184
			0.5762			0.35	35.91		
0.7555	0.7604		0.7579	0.16	0.20	0.18		All sap-wood.....	4
0.7844	0.6945		0.7145	0.25	0.32	0.29			221
0.8115	0.8160	0.8109	0.8128	0.26	0.33	0.30		All sap-wood.....	844
			0.7617			0.26	47.47		

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
300. <i>Alnus maritima</i> <i>Seaside Alder.</i>	809	Delaware	Adger's mill	W. M. Canby	Moist, sandy loam.	0.108	25	
	810	do	Pepper's mill	do	do			
301. <i>Alnus rubra</i> <i>Alder.</i>	907	Alaska	Sitka	Paul Schnitze				
	991	Washington ter- ritory.	Puyallup	G. Engelmann and C. S. Sargent.				
	1025	Oregon	Portland Furniture Company.	do				
302. <i>Alnus rhombifolia</i> <i>Alder.</i>	717	Montana	Missoula	Sereno Watson	Wet			
	979	Oregon	Drain	C. S. Sargent	Moist loam			
303. <i>Alnus oblongifolia</i> <i>Alder.</i>	694	California	San Bernardino	W. G. Wright				
304. <i>Alnus serrulata</i> <i>Black Alder. Smooth Alder.</i>	541	Alabama	Stockton	C. Mohr	Wet			
	611	Georgia	Altamaha river	A. H. Curtis	Damp	0.106	28	
305. <i>Alnus incana</i> <i>Speckled Alder. Hoary Alder. Black Alder.</i>	374	Vermont	Hinesburgh	C. G. Pringle	Wet loam			
	862	Massachusetts	Danvers	J. Robinson	do			
SALICACEÆ								
306. <i>Salix nigra</i> <i>Black Willow.</i>	232	Vermont	Shelburne	C. G. Pringle	Wet, sandy			
	855	Massachusetts	Topshfield	J. Robinson	Alluvial	0.202	12	50
307. <i>Salix amygdaloides</i> <i>Willow.</i>	884	Utah	Salt Lake City	M. E. Jones	Clay			
	908	Colorado	Canon City	E. Weston				
	911	do	do					
308. <i>Salix laevigata</i> <i>Willow.</i>	690	California	Santa Cruz	G. Engelmann and C. S. Sargent.	Moist, sandy			
309. <i>Salix lasiandra</i> <i>Willow.</i>	1219	do	do	C. L. Anderson				
309. <i>Salix lasiandra</i> , var. <i>lanceifolia</i>	640	do	Strawberry valley ..	G. Engelmann and C. S. Sargent.	Moist, rich			
	981	Oregon	Portland	F. Skinner	Alluvial			
309. <i>Salix lasiandra</i> , var. <i>Fendleriana</i>	889	Utah	City Creek cañon ..	M. E. Jones	Gravelly			
310. <i>Salix longifolia</i> <i>Sand-bar Willow.</i>	1174	Illinois	Rockford	M. S. Bebb		0.120	4	11
310. <i>Salix longifolia</i> , var. <i>exigua</i>	955	Texas	Matagorda bay	C. Mohr	Moist, saline			

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.4988	0.5186		0.5087	0.34	0.35	0.35			809
0.4928	0.4889		0.4908	0.37	0.47	0.42			810
			0.4998			0.39	31.13		
0.4922	0.4851		0.4887	0.39	0.40	0.39			987
0.4855	0.4847		0.4856	0.50	0.49	0.54			991
0.4829	0.4865		0.4897	0.36	0.28	0.32			1026
			0.4813			0.42	29.99		
{ 0.4558 }	0.3722	0.3904	0.4077	{ 0.39 }	0.45 }	0.39		First sp. gr. determination made on sap-wood; second sp. gr. determination made on 0.5 sap-wood.	717
{ 0.4427 }				{ 0.43 }	0.30 }				
0.4172	0.4188		0.4178	0.19	0.25	0.22			979
			0.4127			0.31	25.72		
0.4138	0.3823		0.3981	0.42	0.43	0.42	24.81		694
0.4714	0.4685		0.4675	0.32	0.34	0.33			541
0.4573	0.4738		0.4656	0.47	0.39	0.43			611
			0.4666			0.38	29.08		
0.4029	0.4842		0.4436	0.47	0.40	0.44		All sap-wood	874
0.4980	0.4642	0.4732	0.4778	0.38	0.40	0.39	28.71	0.75 sap-wood	903
			0.4607			0.42			
0.4337	0.4102		0.4215	0.62	0.79	0.71			232
0.4790	0.4621	0.4676	0.4696	0.65	0.71	0.68		First and second sp. gr. determinations made on 0.66 sap-wood; third sp. gr. determination made on 0.5 sap-wood.	855
			0.4456			0.70	27.77		
0.4530			0.4530	0.55	0.89	0.72			884
0.4689	0.4448		0.4488	{ 1.06 }	0.91 }	1.11		One tree	908
0.4612	0.4301			{ 1.38 }	1.08 }				911
			0.4509			0.92	28.10		
0.4916	0.4828		0.4872	0.56	0.60	0.58	30.36		690
0.4810	0.4702		0.4756	0.60	0.59	0.60	29.64		1219
0.450*	0.4596		0.4519	0.61	0.73	0.67			640
0.5063	0.4573	0.4089	0.4575	1.09	0.72	0.91			981
			0.4547			0.79	28.34		
0.4576	0.4619		0.4598	0.53	0.58	0.56	28.65		889
0.4901	0.4959		0.4930	0.46	0.49	0.48	30.72		1174
0.5221	0.5329	0.5482	0.5342	1.02	1.09	1.06	33.29		955

FOREST TREES OF NORTH AMERICA.

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
311. <i>Salix sessilifolia</i>	1143	Oregon	Mouth of Willamette river.	Jos. Howells	Alluvial			
312. <i>Salix discolor</i>	850	Massachusetts	Topshfield	J. Robinson	Loam	0.091	7	6
<i>Glaucous Willow.</i>	1225	New York	Ellenburg	J. H. Sears	do			
313. <i>Salix flavescens</i>	641	New Mexico	Pinos Altos mountains	E. L. Greene				
	721	Montana	Pattee's cañon, Missoula	S. Watson	Rich, moist			
	888	Utah	City Creek cañon ..	M. E. Jones	Moist, gravelly ..	0.060	13	4
313. <i>Salix flavescens</i> , var. <i>Scouleriana</i>	972	Washington territory.	Seattle	G. Engelmann and C. S. Sargent.	Moist, sandy	0.100	22	10
<i>Black Willow.</i>								
314. <i>Salix Hookeriana</i>	966	Oregon	Winchester bay ..	do	Sandy saline			
315. <i>Salix cordata</i> , var. <i>vestita</i>	1175	Nebraska	Brownsville	R. W. Furnas	Alluvial	0.148	5	11
<i>Diamond Willow.</i>	1180	Iowa	Sioux City	W. G. Wright	do	0.117	8	14
316. <i>Salix lasioplepis</i>	609	California	Santa Cruz	G. Engelmann and C. S. Sargent.	Sandy loam	0.128	16	
<i>Willow.</i>								
317. <i>Salix Sitchensis</i>	1161	Oregon	Sauvie's Island ..	Jos. Howells	Alluvial			
<i>Silky Willow.</i>								
318. <i>Populus tremuloides</i>	272 ¹	Colorado	Alpine	T. S. Brandegee	Damp			
<i>Aspen. Quaking Asp.</i>	272 ²	do	do	do	do			
	272 ³	do	do	do	do			
	411	Vermont	Charlotte	C. G. Pringle	Moist			
	1035	Massachusetts	Danvers	J. Robinson	Gravelly	0.187	17	7
319. <i>Populus grandidentata</i>	157	Vermont	Charlotte	C. G. Pringle	Sandy loam			
<i>Poplar.</i>	847	Massachusetts	Danvers	J. Robinson	Gravelly	0.220	41	
320. <i>Populus heterophylla</i>	522	Tennessee	Nashville	A. Gattinger	Alluvial			
<i>River Cottonwood. Swamp Cottonwood.</i>	554	Alabama	Stockton	C. Mohr	do			
321. <i>Populus balsamifera</i>	159	Vermont	Shelburne	C. G. Pringle	Sandy loam			
<i>Balsam. Tacamahac. Balm of Gilead.</i>	961	Alaska	Chilcoot inlet	Paul Schultze	Alluvial			

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.4326	0.4545	0.4319	0.4397	0.49	0.50	0.50	27.40		1148
0.4930 0.4434	0.4184 0.4643		0.4548	0.48	0.49	0.49			859
			0.3974	0.36	0.37	0.37			1225
0.3974			0.4261			0.43	26.55		
0.4912	0.5081		0.4997	0.79	0.68	0.74			641
0.5224	0.4953		0.5094	0.36	0.42	0.39			721
0.4923	0.4707		0.4815	0.67	0.72	0.70			888
			0.4969			0.61	30.97		
0.5820	0.5504		0.5412	0.43	0.34	0.39	33.73		972
0.5471	0.5229		0.5350	0.33	0.32	0.32	33.34		966
0.5750	0.5683		0.5716	0.49	0.44	0.47			1175
0.6998	0.6450		0.6423	0.72	0.71	0.72			1180
			0.6069			0.59	37.83		
0.5704	0.5823	0.5734	0.5587	1.01	0.95	0.98	34.82		669
0.5011	0.5133		0.5072	0.67	0.52	0.59	31.61		1161
0.3785			0.3785	0.76		0.76		All sap-wood	272 ¹
0.3579			0.3579	0.72		0.72		All sap-wood	272 ²
0.3569			0.3569					All sap-wood	272 ³
0.4890	0.5021		0.4951	0.31	0.31	0.31		All sap-wood	411
0.4184 0.4227	0.4255 0.4446		0.4278	0.43	0.39	0.41		All sap-wood	1035
			0.4032			0.55	25.18		
0.5119	0.4872		0.4996		0.43	0.43		All sap-wood	157
0.4205	0.4416	0.4183	0.4268	0.45	0.47	0.46		All sap-wood	847
			0.4632			0.45	28.87		
0.4023	0.4113		0.4068	0.74	0.70	0.72		All sap-wood	522
0.4115	0.4105		0.4110	0.90	0.90	0.90			554
			0.4089			0.81	25.48		
0.3524	0.3464		0.3464	1.12	0.72	0.92		All sap-wood	159
0.3843	0.3768		0.3806	0.39	0.40	0.40			961
			0.3635			0.66	22.65		

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
321. <i>Populus balsamifera</i> , var. <i>candicans</i>	1054	Massachusetts....	Topefield	J. Robinson	Gravelly	0.208	11	19
322. <i>Populus angustifolia</i>	552	Colorado	Manitou Springs....	Robert Douglas....	Sandy loam	0.098	10	11
323. <i>Populus trichocarpa</i>	1012	Oregon	Saint John's Barrel	F. Skinner				
Black Cottonwood. Balsam Cottonwood.	1028	do	Factory, Portland. Portland Furniture Company.	G. Engelmann and C. S. Sargent.				
324. <i>Populus monilifera</i>	199	Ohio	Barney and Smith	E. E. Barney				
Cottonwood. Necklase Poplar. Caro-	234	Vermont.....	Manufacturing Co. Charlotte	C. G. Pringle	Alluvial			
kina Poplar. Big Cottonwood.	255	Missouri.....		G. W. Letterman	do			
.....	304	do	do	do	do			
.....	309	Texas	Dallas	J. Reverchon	do	0.806	5	59
.....	754	Florida	Chattahoochee	A. H. Curtiss	do	0.250	20	
325. <i>Populus Fremontii</i>	659	California.....	Sacramento valley ..	G. R. Vasey	do	0.444	16	19
Cottonwood.	646	do	San Bernardino	W. G. Wright	do			
325. <i>Populus Fremontii</i> , var. <i>Wialisensis</i>								
Cottonwood. White Cottonwood.								
CONIFERÆ.								
326. <i>Libocedrus decurrens</i>	579	do	Saw-mill, Straw-	G. Engelmann and				
White Cedar. Bastard Cedar. Post	634	do	berry valley.	C. S. Sargent.				
Cedar. Incense Cedar.	662	do	Saw-mill, San Ber-	W. G. Wright				
.....			nardino mountains.					
327. <i>Thuja occidentalis</i>	104	Vermont.....	Monkton	C. G. Pringle	Cold, peaty			
White Cedar. Arbor-vita.	379	do	do	do	do			
.....	782	New Brunswick ..		Intercolonial rail-				
.....	783	do	Bridgeton	way. Ed. Sinclair				
.....	790	Province of Quebec	Amqui	A. Grant				
.....	792	do		Grand Trunk rail-				
.....	796	do		way. do				
.....	874	Maine	Mattawamkeag	J. Robinson		0.172	19	71
.....	1099	Wisconsin	Eau Claire	H. C. Putnam	Drift			
328. <i>Thuja gigantea</i>	1017	Oregon	Weidler's saw-mill,	G. Engelmann and				
Red Cedar. Ocmos Cedar.	1021	do	Portland. Portland Furniture Company.	C. S. Sargent. do				
329. <i>Chamaecyparis sphaeroides</i>	350	Alabama	Cottage Hill	C. Mohr	Sandy, wet	0.298	7	20
White Cedar.	350	Massachusetts....	Beverly	J. Robinson	Swampy			
.....	351	do	do	do	do			
.....	352	do	do	do	do			

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.3959	0.4222	0.4301	0.4161	0.48	0.43	0.46	25.08	Cultivated. First and second sp. gr. determinations made on sap-wood.	1054
0.3943	0.3881		0.3912	0.82	0.75	0.79	24.38		552
0.3511	0.3656	0.3567	0.3578	1.47	0.87	1.17		Third sp. gr. determination made on sap-wood.	1012
0.4163	0.3937		0.4050	1.56	1.18	1.37			1028
			0.3814			1.27	23.77		
0.3817	0.3946		0.3882	1.34	1.43	1.39			199
0.3939	0.3302		0.3621	0.68	0.94	0.81			234
0.3315	0.3444		0.3380	1.18	0.92	1.05			255
0.3301	0.3281		0.3216	0.83	0.79	0.81			304
0.4706	0.4778		0.4742	1.09	0.96	1.03			309
0.4355	0.4632		0.4494	0.69	0.60	0.65			754
			0.3889			0.96	24.24		
0.5460	0.4655	0.4597	0.4914	0.77		0.77	30.62	All sap-wood	659
0.4827	0.4176	0.4860	0.4621	1.17	1.09	1.13	23.80		646
0.3901	0.3830		0.3861	0.03	0.04	0.04			579
0.3423	0.3769		0.3599	0.13	0.17	0.15			634
0.4506	0.4506		0.4501	0.08	0.05	0.06			662
			0.4017			0.08	25.08		
0.3043	0.3021		0.3035	0.29	0.25	0.27			104
0.2847	0.2870		0.2859	0.29	0.29	0.29			379
0.3234	0.3275		0.3230	0.49	0.51	0.50			782
0.3163	0.3275	0.3272	0.3237	0.39	0.44	0.42			783
0.3016	0.3007		0.3012	0.29	0.31	0.30			790
0.3452	0.2880		0.3166	0.36	0.44	0.40			792
0.3108	0.3104		0.3106	0.43	0.37	0.40			796
0.3584	0.3529	0.3697	0.3603	0.39	0.34	0.37			874
0.3121	0.3232		0.3177	0.37	0.38	0.38			1099
			0.3164			0.37	19.72		
0.3996	0.4173		0.4087	0.12	0.14	0.13			1017
0.3460	0.3551		0.3506	0.24	0.19	0.22			1021
			0.3796			0.17	23.66		
0.3337	0.3447		0.3392	0.94	0.90	0.92			350
0.3105	0.3644	0.3061	0.3277	0.17	0.15	0.16			350
0.3038	0.3036	0.3182	0.3085	0.13	0.11	0.12			351
0.2990	0.3091	0.4527	0.3536	0.11	0.13	0.12			352
			0.3322			0.33	20.70		

FOREST TREES OF NORTH AMERICA.

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
330. <i>Chamaecyparis Nutkaensis</i> <i>Yellow Cypress. Sitka Cypress.</i>	969	Alaska	Sitka	Paul Schultze				
	983	British Columbia	Saw-mill, Victoria ..	G. Engelmann and C. S. Sargent ..				
	994	Alaska	Peril strait	Paul Schultze				
	1000	do	Weidler's saw-mill, Portland, Oregon.	G. Engelmann and C. S. Sargent ..				
331. <i>Chamaecyparis Lawsoniana</i> <i>Port Orford Cedar. Oregon Cedar. White Cedar. Lawson's Cypress. Ginger Pine.</i>	701	Oregon	Dean & Co.'s saw-mill, Marshfield.	do				
	707	do	do	do				
332. <i>Cupressus macrocarpa</i> <i>Monterey Cypress.</i>	675	California	Monterey	do	Gravelly loam ..			
333. <i>Cupressus Goveniana</i>	691	do	Marin county	G. R. Vasey	Dry ridges	0.233	95	
	1100	do	Callistoga	W. F. Fisher	do	0.234	36	
335. <i>Cupressus Guadalupeensis</i>	798	Eastern Arizona ..	San Francisco mount-ains.	E. L. Greene	Rocky			
	1149	Arizona	Santa Catalina mountains.	C. G. Pringle	do			
336. <i>Juniperus Californica</i> <i>Juniper.</i>	1229	California	San Bernardino county.	do	Gravelly			
336. <i>Juniperus Californica</i> , var. <i>Utahensis</i> .. <i>Juniper.</i>	837	Utah	Lewiston	M. E. Jones	do	0.072	55	12
337. <i>Juniperus pachyphloea</i> <i>Juniper.</i>	555	New Mexico	Silver City	E. L. Greene	Stony			
	592	Arizona	Santa Rita mount-ains.	G. Engelmann and C. S. Sargent ..	Gravelly			
	692	do	do	do	do			
338. <i>Juniperus occidentalis</i> <i>Juniper.</i>	624	California	Yreka plains	do	do			
338. <i>Juniperus occidentalis</i> , var. <i>monosperma</i> .. <i>Juniper.</i>	420	New Mexico	Silver City	E. L. Greene	Stony	0.175	22	64
	527	Colorado	Manitou Springs ..	Robert Douglas ..	Gravelly			
338. <i>Juniperus occidentalis</i> , var. <i>conjugens</i> .. <i>Juniper.</i>	969	Texas	Austin	C. Mohr	Limestone	0.134	28	28
	1102	do	do	S. B. Buckley	do			
339. <i>Juniperus Virginiana</i> <i>Red Cedar. Savin.</i>	14	Massachusetts	Arnold Arboretum ..	C. S. Sargent	Drift	0.230	19	49
	327	Texas	Dallas	J. Reverchon	Calcareous	0.484	16	66
	784	Florida	Chattahoochee	A. H. Curtiss	do			
	800	do	Saint John's river ..	do	Sandy loam			
	837	Massachusetts	Danvers	J. Robinson	Drift			
	924	Florida	Chattahoochee	C. Mohr	Alluvial			
	1055	Massachusetts	Topsfield	J. Robinson	Drift	0.208	15	37

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.4011	0.4484		0.4248	0.30	0.37	0.34			969
0.4987	0.5010		0.4999	0.29	0.34	0.32			983
0.5356	0.5166		0.5261	0.20	0.34	0.32			994
0.4555	0.4686		0.4621	0.39	0.36	0.38			1000
			0.4782			0.34	29.80		
0.4856	0.4487		0.4423	0.07	0.12	0.10			701
0.4390	0.5250		0.4819	0.08	0.09	0.09			707
			0.4621			0.10	28.80		
0.6200	0.6268		0.6261	0.55	0.59	0.57	39.02		975
0.4492	0.4733		0.4613	0.41	0.35	0.38			691
0.4620	0.4910		0.4765	0.54	0.47	0.51			1100
			0.4689			0.45	29.22		
0.4648	0.4643	0.4617	0.4636	0.43	0.38	0.41		All sap-wood.....	798
0.4826	0.5272		0.5049	0.41	0.51	0.46			1149
			0.4843			0.44	30.18		
0.6309	0.6255		0.6282	0.77	0.73	0.75	39.15		1229
0.5504	0.5539		0.5522	0.47	0.51	0.49	34.41	0.8 sap-wood.....	887
0.5744	0.6965		0.6355	0.11	0.16	0.14			555
0.5085	0.5619	}	0.5302	{	0.07	0.07	0.08	One tree.....	{
0.5185	0.5321								
			0.5629			0.11	36.32		692
0.5724	0.5807		0.5765	0.12	0.12	0.12	35.93		624
0.7329	0.7440		0.7385	0.68	0.68	0.68			420
0.6768	0.6935		0.6852	0.89	0.86	0.88			527
			0.7119			0.78	44.36		
	0.6967		0.6967	0.45	0.47	0.46			939
0.6750	0.6840	0.6943	0.6846	0.39	0.53	0.46			1102
			0.6907			0.46	43.04		
0.5112	0.5045		0.5079	0.15	0.14	0.14		0.1 sap-wood.....	14
0.4585	0.4498		0.4539	0.12	0.14	0.13			327
0.5455	0.5310	0.5533	0.5433	0.16	0.17	0.16			734
0.4901	0.4727		0.4814	0.13	0.08	0.11			800
0.4804	0.4693	0.4733	0.4743	0.16	0.14	0.15			837
0.4838	0.4705	0.4740	0.4761	0.16	0.12	0.14			924
0.5290	0.4937	0.5107	0.5111	0.09	0.09	0.09			1055
			0.4926			0.13	30.70		

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
340. <i>Taxodium distichum</i> Bald Cypress. Black Cypress. Red Cypress. White Cypress. Deciduous Cypress.	585	Alabama	Stockton	C. Mohr	Alluvial			
	542	do	Otis & Co., saw-mill.	do	do			
	543	do	do	do	do			
	581	Arkansas	Little Rock	G. W. Letterman	do			
	582	do	do	do	do			
	741	Florida	Chattahoochee	A. H. Curtiss		0.514	55	230
	894 ¹			Department of Agriculture.				
	894 ²			do				
	923	Alabama	Stockton	C. Mohr	Alluvial			
	1212			do				
341. <i>Sequoia gigantea</i> Big Tree.	1213			do				
	1214	Louisiana	Bayou La Battery	do	Alluvial			
342. <i>Sequoia sempervirens</i> Redwood.	657	California	Tulare county	G. Engelmann and C. S. Sargent.	Granite			
	666	do	do	do	do			
343. <i>Taxus brevifolia</i> Yew.	673	do	Russian river	C. S. Sargent				
	710	do	Santa Cruz	Turner, Kennedy & Shaw.				
	711 ¹	do	do	do				
	711 ²	do	do	do				
	712	do	Mendocino county	J. Kentfield & Co				
	713	do	do	do				
	714	do	do	do				
	715	do	do	do				
343. <i>Taxus brevifolia</i> Yew.	978	Oregon	Portland	G. Engelmann and C. S. Sargent.	Moist, rich	0.279	8	46
344. <i>Taxus floridana</i> Yew.	305	Florida	Flat Creek	A. H. Curtiss	Sandy loam			
345. <i>Torreya taxifolia</i> Stinking Cedar. Savin.	62	do	Chattahoochee	C. S. Sargent	Alluvial			
	277	do	do	A. H. Curtiss	Calcareous	0.284	12	78
346. <i>Torreya californica</i> California Nutmeg. Stinking Cedar.	651	California	Marin county	G. R. Vasey	Stony	0.361	20	73
347. <i>Pinus strobus</i> White Pine. Weymouth Pine.	1	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	0.510	19	51
	164	Michigan	Woodsum Machine Company.	E. E. Barney				
	208	do	Barney & Smith	do				
	222	Vermont	Manufacturing Co. Charlotte	C. G. Pringle	Wet, swampy			
	723	Pennsylvania	Williamsport	do				
	777	New Brunswick		Intercolonial railway.				

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.4559	0.4442		0.4501	0.44	0.40	0.42			585
0.5114	0.4896		0.5005	0.35	0.31	0.33		White cypress.....	542
0.5017	0.5224		0.5121	0.66	0.53	0.59		Black cypress.....	543
0.4683	0.4545		0.4614	0.38	0.45	0.42			581
0.4683	0.5031		0.4857	0.34	0.29	0.32			582
0.4159	0.4064	0.3808	0.4020	0.27	0.36	0.32		White cypress sap-wood.....	741
0.2444	0.2013			0.56				Butt of knee.....	804 ¹
0.2362	0.2394			0.55				Top of knee.....	804 ²
0.3442	0.3656							From opposite sides of center of tree next to heart.....	One tree. 928
0.4282	0.3705		0.4072	0.39		0.44		Next outside of preceding.....	
0.4599	0.5022			0.48				do.....	
0.4112	0.4336	0.3492						Outside of tree.....	
0.3860	0.3750		0.3805	0.50	0.50	0.50		Outside wood.....	1212
0.4123	0.4937		0.4540	0.51	0.53	0.52		Heart-wood.....	1213
0.4878	0.4904		0.4891	0.36	0.37	0.37		Black cypress.....	1214
.....			0.4543			0.42	27.65		
0.3234	0.3202		0.3240	0.54	0.50	0.44			607
0.3236	0.3239			0.41	0.33				
0.3524	0.2523		0.2524	0.50	0.64	0.57		All sap-wood.....	606
.....			0.2882			0.50	18.20		
0.3788	0.3965		0.3877	0.12	0.23	0.17			673
0.4242	0.4281		0.4262	0.16	0.19	0.18		Wood from top of tree.....	710
0.3963	0.4444		0.4204	0.11		0.11		Wood from butt of tree.....	711 ¹
0.4912	0.4562		0.4737	0.14		0.14		Wood from butt of tree.....	711 ²
0.3015	0.2982		0.2999	0.06	0.11	0.09			712
0.3610	0.3612		0.3611	0.18	0.15	0.17			713
0.5226	0.4714		0.4970	0.17	0.16	0.17			714
0.4827	0.5181		0.5004	0.10	0.13	0.12		Wood with curled grain.....	715
.....			0.4208			0.14	26.23		
0.6390	0.6553	0.6230	0.6391	0.22	0.22	0.22	39.83		978
0.6340			0.6340	0.20	0.22	0.21	39.51		305
0.6034	0.5969		0.5702	1.08	1.38	1.23			63
0.4728	0.4447		0.4588	0.25	0.21	0.23			277
.....			0.5145			0.73	32.06		
0.4823	0.4696		0.4760	1.27	1.40	1.34	29.66		651
0.3946	0.4106		0.4026	0.11	0.19	0.15			1
0.3481	0.3459		0.3470	0.13	0.17	0.15			164
0.3400	0.3413		0.3409	0.22	0.20	0.21			206
0.4880	0.6461		0.5670	0.23	0.21	0.22			222
0.3690	0.3850		0.3770	0.18	0.12	0.15			723
0.3606	0.3140		0.3373	0.23	0.30	0.26			777

FOREST TREES OF NORTH AMERICA.

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
247. <i>Pinus Strobus</i> —continued	788	New Brunswick ..	Bridgeton	Ed. Sinclair				
	789	Province of Quebec.	Amqui	A. Grant				
	797	do		Grank Trunk railway.				
	1044	Massachusetts ...	Reading	J. Robinson	Drift	0.215	15	16
248. <i>Pinus monticola</i>	975	British Columbia ..	Hasting's saw-mill, Burrard inlet.	G. Engelmann and C. S. Sargent.				
	987	Oregon	Cascade mountains ..	C. S. Sargent	Moist loam			
249. <i>Pinus Lambertiana</i>	688	California	Saw-mill, Strawberry valley.	G. Engelmann and C. S. Sargent.				
	668	do		G. R. Vasey				
	780	do	Laassen's peak	Sierra Lumber Company, San Francisco.				
250. <i>Pinus flexilis</i>	819	Colorado	Forest City	T. S. Brandegee	Gravelly	0.502	38	120
	918	Nevada	Monitor range	A. Triple	do			
251. <i>Pinus albicaulis</i>	992	British Columbia ..	Silver Mountain valley, Fraser river.	G. Engelmann and C. S. Sargent.		0.494	50	180
252. <i>Pinus reflexa</i>	597	Arizona	Santa Rita mountains.	do	Rocky			
	602	New Mexico	Pinos Altos mountains.	E. L. Greene				
	661	Arizona	Santa Rita mountains.	G. Engelmann and C. S. Sargent.				
253. <i>Pinus Parryana</i>	656	California	San Diego county ..	G. R. Vasey				
254. <i>Pinus cembroides</i>	1226	Arizona	Santa Catalina mountains.	C. G. Pringle				
255. <i>Pinus edulis</i>	397	Colorado	Cañon City	E. Weston	Gravelly	0.284	30	79
256. <i>Pinus monophylla</i>	823	Eastern Arizona ..	San Francisco mountains.	E. L. Greene	do			
	882	Utah	Lewiston	M. E. Jones	Rocky	0.164	19	66
	891	Eastern Arizona ..	San Francisco mountains.	E. L. Greene				
	900	California		Department of Agriculture.				
	915	Nevada	Danville	A. Triple	Gravelly			
257. <i>Pinus Balfouriana</i>	577	California	Scott mountains	G. Engelmann and C. S. Sargent.	Rocky	0.368	75	309
	631							
257. <i>Pinus Balfouriana</i> , var. <i>aristata</i>	821	Colorado	Forest City	T. S. Brandegee		0.450	44	136
	914	Nevada	Prospect mountain ..	A. Triple	Rocky			
258. <i>Pinus resinosa</i>	194	Michigan	Barney & Smith Manufacturing Co.	E. E. Barney				

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.3901	0.3970		0.3980	0.20		0.20			788
0.3624	0.3463		0.3543	0.19	0.19	0.19			789
0.3905	0.3725		0.3815	0.21	0.22	0.21			797
0.3503	0.3513	0.3438	0.3485	0.10	0.13	0.12			1044
			0.3854			0.19	24.02		
0.4291	0.4102		0.4197	0.26	0.28	0.27			975
0.3634	0.3554		0.3619	0.15	0.21	0.18		Cut at 3,800 feet elevation.....	987
			0.3908			0.23	24.35		
0.4066	0.4014		0.4040	0.17	0.16	0.17			638
0.3783	0.3889		0.3835	0.29	0.25	0.27			668
0.3153	0.3201		0.3177	0.22	0.24	0.23			730
			0.3684			0.22	22.96		
0.4965	0.4031	0.4526	0.4507	0.20	0.20	0.20		Second sp. gr. determination made on sap-wood. Cut at elevation of 10,000 feet.	819
0.4209			0.4209	0.35		0.35		All sap-wood.....	913
			0.4358			0.28	27.16		
0.4233	0.4097		0.4165	0.30	0.23	0.27	25.96		992
0.4729	0.5253	0.5052	0.4971	0.24	0.24	0.24		First sp. gr. determination made on sap-wood.....	597
0.4986	0.4961	0.4847		0.27	0.20				602
0.4869	0.4696		0.4783	0.30	0.27	0.28			661
			0.4877			0.26	30.39		
0.5628	0.5721		0.5675	0.54		0.54	35.37		656
0.6695	0.6330		0.6512	0.88	0.92	0.90	40.68		1226
0.5813	0.6963		0.6388	0.57	0.68	0.62	39.81	Second sp. gr. specimen contained a knot.....	397
0.6000	0.5568		0.5781	0.78	0.65	0.71			823
0.4499	0.4644		0.4572	0.67	0.77	0.72			883
0.6140	0.7617		0.6878	0.72	0.73	0.72		Second sp. gr. specimen was very resinous.....	891
0.5715	0.5230		0.5473	0.89	0.76	0.83			906
0.5585			0.5585	0.41		0.41			915
			0.5658			0.68	35.26		
0.5434	0.5409	0.5273	0.5434	0.41	0.41	0.40	33.86	One tree.....	577
0.5580	0.5476			0.38	0.42				631
0.4907	0.5319	0.4671	0.4996	0.18	0.18	0.18		Third sp. gr. determination made on sap-wood.....	821
0.6149			0.6149	0.42		0.42		0.75 sap-wood.....	914
			0.5572			0.30	34.72		
0.4872	0.4838		0.4855	0.28	0.18	0.20			194

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
358. <i>Pinus resinosa</i> —continued.....	315	Michigan.....	Hersey.....	W. J. Beal.....				
	413	Vermont.....	Charlotte.....	C. G. Pringle.....	Clay.....			
	779	New Brunswick.....		Intercolonial rail- way.				
	785	do.....	Bridgeton.....	Ed. Sinclair.....				
	1074	Vermont.....	Charlotte.....	C. G. Pringle.....				
	1075	do.....	do.....	do.....				
	1076	do.....	do.....	do.....				
359. <i>Pinus Torreyana</i>	996	California.....	San Diego county.....	G. Engelmann.....	Sandy.....	0.234	82	8
360. <i>Pinus Arizona</i>	1144	Arizona.....	Santa Rita mount- ains.	C. G. Pringle.....	Rocky.....			
<i>Yellow Pine.</i>	1154	do.....	do.....	do.....	do.....			
	1155	do.....	do.....	do.....	do.....			
	1156	do.....	do.....	do.....	do.....			
361. <i>Pinus ponderosa</i>	619	Dakota.....	Deadwood.....	Robert Douglas.....	Gravelly.....			
<i>Yellow Pine. Bull Pine.</i>	626	Oregon.....	Saw-mill, Ashland..	G. Engelmann and C. S. Sargent.				
	630	California.....	Strawberry valley..	do.....	Low, wet, swampy			
	632	do.....	do.....	do.....				
	636	Oregon.....	Saw-mill, Ashland..	do.....				
	689	California.....	Saw-mill, San Ber- nardino.	W. G. Wright.....				
	718	Montana.....	Saw-mill, Missoula..	S. Watson.....				
	731	California.....	Lassen's peak.....	Sierra Lumber Com- pany.				
	907	Colorado.....	Cañon City.....	E. Weston.....				
	910	do.....	do.....	do.....				
	1007	California.....	Saw-mill, San Ber- nardino.	W. G. Wright.....				
362. <i>Pinus Jeffreyi</i>	578	do.....	Scott mountains.....	G. Engelmann and C. S. Sargent.	Dry, gravelly.....			
<i>Bull Pine. Black Pine.</i>	633	do.....	do.....	do.....	do.....			
	667	California.....	Saw-mill, San Ber- nardino.	W. G. Wright.....				
363. <i>Pinus Chihuahuana</i>	593	Arizona.....	Santa Rita mount- ains.	G. Engelmann and C. S. Sargent.	Dry, gravelly.....	0.510	102	53
	664	do.....	do.....	do.....	do.....			
364. <i>Pinus contorta</i>	997	British Columbia.....	Vancouver's island	G. Engelmann and C. S. Sargent.	do.....			
<i>Sorub Pine.</i>								
365. <i>Pinus Murrayana</i>	293	Colorado.....	Forest City.....	T. S. Brandegee.....	Moist, sandy loam.	0.269	43	23
<i>Tamarack. Black Pine. Lodge-pole</i> <i>Pine. Spruce Pine.</i>	563	do.....	do.....	C. S. Sargent.....	do.....			
	625	California.....	Scott mountains.....	G. Engelmann and C. S. Sargent.	do.....			
366. <i>Pinus Sabiniana</i>	571	do.....	Jolon.....	G. B. Vasey.....				
<i>Digger Pine. Bull Pine.</i>	644	do.....	Contra Costa county	do.....	Gravelly.....	0.312	43	

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.5450	0.5416		0.5433	0.20	0.24	0.22			315
0.5121	0.4537		0.4829	0.24	0.24	0.24		All sap-wood	413
0.4555	0.4911		0.4733	0.28	0.29	0.29		All sap-wood	779
0.4250	0.4631		0.4441	0.39	0.34	0.37			785
0.4627	0.4752	0.5227	0.4869	0.28	0.27	0.27		All sap-wood	1074
0.4931	0.4922	0.4945	0.4909	0.26	0.29	0.28		First and third sp. gr. determinations made on 0.5 sap-wood; second sp. gr. determination made on sap-wood.	1075
0.4892	0.4582	0.4858	0.4777	0.24	0.31	0.28		Second sp. gr. determination made on 0.5 sap-wood; third sp. gr. determination made on sap-wood.	1076
			0.4854			0.27	30.25		
0.4659	0.5119	0.4860	0.4879	0.38	0.33	0.35	30.40	First and second sp. gr. determinations made on sap-wood	906
0.5360	0.5079		0.5220	0.14	0.15	0.14			1144
0.5066	0.7844		0.6455	0.21	0.20	0.20		Second sp. gr. specimen very resinous	1154
0.4567	0.4204	0.4611	0.4461	0.19	0.26	0.23			1155
0.4168	0.3964		0.4016	0.25	0.24	0.24			1156
			0.5038			0.20	31.40		
0.4877	0.4813		0.4845	0.27	0.26	0.27			619
0.4552	0.4419		0.4485	0.38	0.41	0.40		All sap-wood	626
0.4571	0.4152		0.4362	0.38	0.36	0.37		All sap-wood	630
0.5264	0.5350		0.5307	0.31	0.30	0.30		All sap-wood	632
0.4184	0.4240		0.4212	0.34	0.43	0.39		All sap-wood	636
0.5144	0.5265		0.5204	0.23	0.23	0.23			689
0.3972	0.4479		0.4225	0.40	0.36	0.38			718
0.4284	0.4437		0.4360	0.28	0.25	0.27		All sap-wood	731
0.4676	0.5305		0.4990	0.58	0.40	0.49		All sap-wood	907
0.4376	0.4459		0.4417	0.45	0.45	0.45		All sap-wood	910
0.5521	0.5412		0.5466	0.30	0.33	0.32		All sap-wood	1007
			0.4715			0.35	29.45		
0.4561	0.4520			0.22	0.20			} One tree. Fourth sp. gr. determination made on 0.5 sap-wood. {	578
0.4985	0.5075		0.4785	0.23	0.36	0.25			633
0.5982	0.5274		0.5628	0.29	0.24	0.27		All sap-wood	867
			0.5206			0.26	32.44		
0.5584	0.5780	}	0.5457	{	0.37 0.41	0.39 0.40	34.01	One tree	593
0.5226	0.5239								664
0.5709	0.5985	0.5750	0.5815	0.21	0.16	0.19	36.24		997
0.3550	0.3551		0.3551	0.36	0.29	0.33			293
0.4291	0.4267		0.4279	0.26	0.25	0.26			563
0.4204	0.4711		0.4457	0.37	0.37	0.37			625
			0.4096			0.32	25.53		
0.4528	0.4531		0.4530	0.43	0.41	0.42		All sap-wood	571
0.4968	0.5312		0.5150	0.40	0.35	0.37		All sap-wood	644
			0.4840			0.40	30.16		

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
367. <i>Pinus Coulteri</i>	1157	California.....	San Bernardino.....	W. G. Wright.....	Dry, gravelly.....			
368. <i>Pinus insignis</i> <i>Monterey Pine.</i>	676	do.....	Monterey.....	G. R. Vasey.....	Gravelly loam.....	0.544	35	
	896	do.....		Department of Ag- riculture.				
369. <i>Pinus tuberculata</i> <i>Knob-cone Pine.</i>	576	do.....	Mount Shasta.....	G. Engelmann and C. S. Sargent.	Gravelly.....	0.418	35	18
370. <i>Pinus Taeda</i> <i>Loblolly Pine. Old-field Pine. Rose- mary Pine.</i>	82	Florida.....	Duval county.....	A. H. Curtiss.....	Moist, sandy.....	0.084	27	6
	855	Alabama.....	Cottage Hill.....	C. Mohr.....	Low, rich.....			
	388	North Carolina.....	Wilmington.....	Edward Kidder.....	Loam.....			
	389	do.....	do.....	do.....	do.....			
371. <i>Pinus rigida</i> <i>Pitch Pine.</i>	13	Massachusetts.....	Arnold Arboretum..	C. S. Sargent.....	Drift.....	0.267	21	15
	1046	do.....	North Reading.....	J. Robinson.....	do.....	0.230	26	8
372. <i>Pinus serotina</i> <i>Pond Pine.</i>	83	Florida.....	Duval county.....	A. H. Curtiss.....	Moist, sandy loam..	0.312	33	19
373. <i>Pinus inops</i> <i>Jersey Pine. Scrub Pine.</i>	621	South Carolina.....	Aiken.....	H. W. Ravenel.....	Dry, sandy.....	0.206	8	42
	622	do.....	do.....	do.....	do.....	0.164	28	34
	1169	Indiana.....	New Albany.....	M. J. Robinson.....				
	1172	New Jersey.....	Mount Holly.....	S. P. Sharples.....				
374. <i>Pinus clausa</i> <i>Sand Pine. Scrub Pine. Spruce Pine.</i>	279	Florida.....	Apalachicola.....	A. H. Curtiss.....	Dry, sandy barrens	0.238	6	36
375. <i>Pinus pungens</i> <i>Table-mountain Pine. Hickory Pine.</i>	321	Virginia.....	Wytheville.....	H. Shriver.....	Clay.....	0.264	20	35
	396	Pennsylvania.....	Colerain Forge.....	J. R. Lowrie.....	Slate.....	0.010	14	
376. <i>Pinus muricata</i> <i>Obispo Pine. Bishop's Pine.</i>	671	California.....	Marin county.....	G. R. Vasey.....	Gravelly.....	0.243	29	
377. <i>Pinus mitis</i> <i>Yellow Pine. Short-leaved Pine. Spruce Pine. Bull Pine.</i>	278	Florida.....	Chattahoochee.....	A. H. Curtiss.....	Clay.....	0.015	33	
	319	Louisiana.....	Amite.....	C. Mohr.....	Sandy loam.....			
	557	Arkansas.....	Texarkana.....	F. L. Harvey.....	do.....			
	558	do.....	do.....	do.....	do.....			
	559	do.....	do.....	do.....	do.....			
378. <i>Pinus glabra</i> <i>Oedar Pine. Spruce Pine. White Pine.</i>	142	South Carolina...	Bonneau's Depot...	H. W. Ravenel.....	Rich upland.....			
	544	Mississippi.....	Gainesville.....	C. Mohr.....	Low, sandy.....			
	764	Florida.....	Chattahoochee.....	A. H. Curtiss.....	Low, wet.....			

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.4272	0.3994		0.4133	0.34	0.40	0.37	25.76	All sap-wood.....	1157
0.4052	0.5244		0.5098	0.25	0.20	0.23			676
0.4082	0.4018		0.4050	0.34	0.37	0.36			896
			0.4574			0.30	28.50		
0.3469	0.3529		0.3499	0.36	0.30	0.33	21.81		576
0.5610	0.5609		0.5609	0.26	0.29	0.27		All sap-wood.....	82
0.5788	0.5466		0.5627	0.25	0.26	0.26		All sap-wood.....	355
0.5458	0.5654		0.5556	0.28	0.26	0.27		Rosemary pine.....	388
0.5045	0.4896		0.4971	0.27	0.23	0.25		All sap-wood.....	389
			0.5441			0.26	33.91		
0.5597	0.5669		0.5633	0.19	0.21	0.20		All sap-wood.....	18
0.4065	0.4154	{	0.4668	{	0.32 0.19	0.26		{ First sp. gr. determination made on sap-wood; second and third sp. gr. determinations made on 0.5 sap-wood. Fourth sp. gr. specimen very resinous.	1046
0.4131	0.6323								
			0.5151			0.23	32.10		
0.7904	0.7931	0.7990	0.7942	0.15	0.18	0.17	49.49		88
0.4906	0.4830		0.4867	0.24	0.27	0.26		All sap-wood.....	621
0.5571	0.5680		0.5626	0.32	0.31	0.32			622
0.5149	0.4603		0.4876	0.32	0.23	0.28		All sap-wood.....	1169
0.5778	0.5841	0.5991	0.5870	0.29	0.34	0.32			1172
			0.5309			0.30	33.09		
0.4696	0.6456		0.5576	0.30	0.31	0.31	34.75		279
0.5235	0.5050		0.5143	0.30	0.30	0.30			321
0.4851	0.4802		0.4727	0.24	0.22	0.23			396
			0.4935			0.27	30.75		
0.5054	0.4831		0.4942	0.26	0.25	0.26	30.80	Second sp. gr. determination made on sap-wood.....	671
0.5739	0.4550		0.5144	0.35	0.35	0.35		All sap-wood.....	278
0.7098	0.7609		0.7333	0.22	0.22	0.22		0.5 sap-wood.....	319
0.6086	0.6129		0.6107	0.23	0.17	0.20			557
0.7262	0.7232		0.7247	0.33	0.31	0.32			558
0.4689	0.4689		0.4689	0.32	0.41	0.37		All sap-wood.....	559
			0.6104			0.29	38.04		
0.4358	0.4235		0.4297	0.40	0.61	0.50		Second sp. gr. determination made on sap-wood.....	142
0.3232	0.3741		0.3487	0.53	0.44	0.49		All sap-wood.....	544
0.4018	0.4002		0.4010	0.35	0.35	0.35		All sap-wood.....	764
			0.3931			0.45	24.50		

FOREST TREES OF NORTH AMERICA.

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
379. <i>Pinus Banksiana</i> <i>Gray Pine. Scrub Pine. Prince's Pine.</i>	394	Michigan	Baldwin	W. J. Beal	Low, sandy			
	780	New Brunswick		Intercolonial rail- way.				
	879	do		do				
380. <i>Pinus palustris</i> <i>Long-leaved Pine. Southern Pine. Georgia Pine. Yellow Pine. Hard Pine.</i>	81	Florida	Duval county	A. H. Curtiss	Sandy loam	0.353	25	119
	85	do	do	do	Moist, sandy	0.355	46	51
	172	Mississippi	Barney & Smith Manufacturing Co	E. E. Barney				
	243	Florida	Saw-mill, Saint John's river.	A. H. Curtiss				
	302	South Carolina	Aiken	H. W. Ravenel	Sandy loam			
	357	Alabama	Cottage Hill	C. Mohr	do			
	358	do	Citronelle	do	do			
	359	do	Chunchula	do	do			
	380	do	do	do	do			
	381	do	do	do	do			
	384	Florida	Saw-mill, Cedar Keys.	A. H. Curtiss				
	385	do	do	do				
	390	North Carolina	Wilmington	E. Kidder				
	562	Texas	Sabine county	G. W. Letterman				
	1096	Alabama	Mobile	C. Mohr				
381. <i>Pinus Cubensis</i> <i>Slash Pine. Swamp Pine. Bastard Pine. Meadows Pine.</i>	84	Florida	Duval county	A. H. Curtiss	Moist, sandy	0.326	58	50
	356	Alabama	Cottage Hill	C. Mohr	do			
	493	Florida	Bay Biscayne	A. H. Curtiss	Coral			
382. <i>Picea nigra</i> <i>Black Spruce.</i>	109	Vermont	Charlotte	C. G. Pringle	Gravelly			
	231	do	do	do	Cold, peaty			
	373	do	Huntingdon	do	Gravelly			
	776	New Brunswick	Bay of Fundy	Intercolonial rail- way.				
	794	Province Quebec	Danville	Grand Trunk rail- way.				
	880	New Brunswick	Bridgeton	Ed. Sinclair				
383. <i>Picea alba</i> <i>White Spruce.</i>	513	New Hampshire	Stratford	C. G. Pringle		0.186	16	22
	620	Dakota	Terry's peak	Robert Douglas	Gravelly			
	773	New Brunswick	Bay of Fundy	Intercolonial rail- way.				
	784	do	Bridgeton	Ed. Sinclair				
	791	Province of Que- bec.	Amqui	A. Grant				
384. <i>Picea Engelmanni</i> <i>White Spruce.</i>	292	Colorado	Forest City	T. S. Brandegee	Damp	0.320	8	76
	575	do	do	C. S. Sargent	Peaty			
	822	do	do	T. S. Brandegee	do	0.318	45	71
	899	do		Department of Agri- culture.				
	905	Utah		do				

THE WOODS OF THE UNITED STATES.

343

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.4622	0.4965		0.4794	0.28	0.33	0.30			394
	0.4588		0.4588	0.20	0.19	0.20		Sap-wood	780
0.4778	0.5065	0.4887	0.4900	0.19	0.18	0.19			879
			0.4761			0.22	29.67		
0.7551	0.7284		0.7418	0.26	0.26	0.26			81
0.7117			0.7117	0.16	0.17	0.17			85
0.6927	0.6974		0.6950	0.26	0.26	0.26			172
0.7569	0.6829		0.7199	0.28	0.21	0.25			243
0.6139	0.5191		0.5665	0.39	0.24	0.31			302
0.6602	0.6616		0.6609	0.27	0.27	0.27			357
0.8652	0.9325		0.8989	0.28	0.20	0.24		Boxed for turpentine, 1852; chipped 10 years; abandoned, 1861.	358
0.8509	0.8450		0.8479	0.15	0.16	0.16		Boxed for turpentine, 1876; chipped 4 years; specimen taken along chip.	359
0.6673	0.7914		0.7294	0.21		0.21		Boxed for turpentine, 1876; chipped 4 years; specimen taken above chip.	360
0.7590	0.7736		0.7663	0.23	0.23	0.23		Boxed for turpentine, 1878; chipped 2 years.	361
0.6163	0.5714		0.5938	0.32	0.32	0.32			364
0.6549	0.5924		0.6236	0.28	0.28	0.28			385
0.4802	0.6186	0.6418	0.5734	0.29	0.34	0.32			390
0.7744	0.6745		0.7245	0.17	0.18	0.18			562
0.6415	0.6490		0.6453	0.19	0.26	0.22		Tree boxed for turpentine 18 or 20 years ago.	1096
			0.6999			0.25	43.62		
0.6533	0.6506		0.6520	0.31	0.31	0.31			84
0.7881	0.7340		0.7611	0.16	0.15	0.16			356
0.8529	0.8889	0.8220	0.8379	0.30	0.31	0.31			498
			0.7504			0.26	46.76		
0.4576	0.4715		0.4646	0.23	0.17	0.20		[P. rubra]	108
0.5289	0.5256		0.5272	0.25	0.27	0.26			231
0.4285	0.4486		0.4386	0.29	0.29	0.29		[P. rubra]	373
0.4730	0.4593		0.4662	0.29	0.27	0.28			776
0.4065	0.4290	0.3907	0.4087	0.29	0.25	0.27			794
0.4830	0.4098	0.4425	0.4451	0.38	0.28	0.33			880
			0.4584			0.27	28.57		
0.4074	0.4194		0.4134	0.26	0.21	0.24			513
0.3848	0.4034		0.3941	0.42	0.38	0.40			620
0.4231	0.4375		0.4303	0.34	0.38	0.36			773
0.3809	0.4188		0.3999	0.32	0.31	0.32			784
0.3737	0.4020		0.3879	0.24	0.29	0.27			791
			0.4051			0.32	25.25		
0.3550	0.3551		0.3551	0.36	0.29	0.33			292
0.3365	0.3217		0.3291	0.35	0.35	0.35			575
0.3597	0.3528		0.3518	0.29	0.24	0.27		Tree with gray bark.	822
0.3717	0.3725		0.3721	0.32	0.33	0.33			899
0.3195	0.3137		0.3166	0.43	0.26	0.34			905
			0.3449			0.32	21.49		

FOREST TREES OF NORTH AMERICA.

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
385. <i>Picea pungens</i> <i>White Spruce. Blue Spruce.</i>	269	Colorado.....	Alpine.....	T. S. Brandegee....	Damp.....			
	270	do.....	do.....	do.....	do.....			
	901	do.....		Department of Agri- culture.				
	906	Utah.....		do.....				
386. <i>Picea Sitchensis</i> <i>Tide-land Spruce.</i>	970	Alaska.....	Sitka.....	Paul Schultze.....				
	977	British Columbia.	Saw-mill, Burrard inlet.	G. Engelmann and C. S. Sargent.				
	1015	Oregon.....	Weidler's mill, Port- land.	do.....				
	1019	do.....	Saw-mill, Astoria...	do.....				
	1026	do.....	Portland Furniture Company.	do.....				
387. <i>Tsuga Canadensis</i> <i>Hemlock.</i>	5	Massachusetts....	Arnold Arboretum..	C. S. Sargent.....	Drift.....	0.382	87	87
	219	Vermont.....	Charlotte.....	C. G. Pringle.....	Gravelly.....			
	726	Pennsylvania....	Williamsport.....	do.....	Damp.....			
	772	New Brunswick..		Intercolonial rail- way.				
	775	do.....	Bay of Fundy.....	do.....				
	778	do.....		do.....				
	787	do.....	Bridgeton.....	Ed. Sinclair.....				
	798	Province Quebec	Danville.....	Grand Trunk rail- way.				
	817	West Virginia...	Grafton.....	C. G. Pringle.....				
	1040	Massachusetts....	Danvers.....	J. Robinson.....	Moist, loam.....	0.280	26	92
	1042	do.....	North Reading.....	do.....		0.260	16	23
388. <i>Tsuga Caroliniana</i> <i>Hemlock.</i>	623	North Carolina...	Hendersonville....	A. H. Curtiss.....	Dry, rocky.....	0.228	51	14
389. <i>Tsuga Mertensiana</i> <i>Hemlock.</i>	971	Washington terri- tory.	Wilkeson.....	G. Engelmann and C. S. Sargent.	Rich loam.....			
	995	Alaska.....	Sitka.....	Paul Schultze.....				
390. <i>Tsuga Pattoniana</i>	980	British Columbia.	Silver peak, near Fraser river.	G. Engelmann and C. S. Sargent.	Gravelly loam...			
391. <i>Pseudotsuga Douglasii</i> <i>Red Fir. Yellow Fir. Oregon Pine. Douglas Fir.</i>	271	Colorado.....	Alpine.....	T. S. Brandegee....	Moist.....	0.276	40	155
	627	California.....	Saw-mill, Strawberry valley.	G. Engelmann and C. S. Sargent.				
	702	Oregon.....	Saw-mill, Marshfield	do.....				
	704	do.....	do.....	do.....				
	705	do.....	E. B. Dean's saw- mill, Marshfield.	do.....				
	706	do.....	do.....	do.....				
	708	do.....	do.....	do.....				
	709	do.....	do.....	do.....				
	720	Montana.....	Saw-mill, Missoula..	S. Watson.....				
	732	California.....	Lassen's peak.....	Sierra Lumber Com- pany.				
	881	Utah.....	Salt Lake.....	M. E. Jones.....	Rocky.....			
	973	British Columbia.	Saw-mill, Burrard inlet.	C. S. Sargent.....				
	974	do.....	do.....	do.....				
	986	do.....	Saw-mill, Victoria..	G. Engelmann and C. S. Sargent.				
	989	Oregon.....	Saw-mill, Portland..	do.....				

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.3480	-----	-----	0.3480	0.22	-----	0.22	-----	-----	269
0.3540	0.3558	-----	0.3549	0.32	0.29	0.31	-----	-----	270
0.4309	0.4274	-----	0.4292	0.49	0.51	0.50	-----	-----	901
0.3764	0.3517	-----	0.3641	0.69	0.28	0.48	-----	-----	906
			0.3740			0.38	23.81	-----	
0.6029	0.6485	-----	0.6257	0.23	0.24	0.24	-----	-----	970
0.3658	0.3974	-----	0.3816	0.16	0.17	0.17	-----	-----	977
0.4280	-----	-----	0.4280	0.16	0.13	0.15	-----	-----	1015
0.3520	0.3423	0.3446	0.3403	0.22	0.13	0.18	-----	-----	1019
0.3661	0.3576	-----	0.3619	0.13	0.12	0.13	-----	-----	1026
			0.4287			0.17	26.72	-----	
0.3880	0.3896	-----	0.3888	0.24	0.25	0.25	-----	-----	5
0.4633	0.4624	-----	0.4629	0.73	0.67	0.70	-----	-----	219
0.3817	0.3823	-----	0.3820	0.43	0.36	0.40	-----	-----	726
0.3963	0.4704	-----	0.4333	0.45	0.46	0.46	-----	Red hemlock	772
0.4839	0.4707	-----	0.4773	0.34	0.42	0.38	-----	White hemlock	775
0.3418	0.3343	-----	0.3381	0.54	0.56	0.55	-----	-----	778
0.4532	0.4538	-----	0.4538	0.44	0.48	0.46	-----	-----	787
0.5033	0.5191	-----	0.5112	0.46	0.46	0.46	-----	-----	798
0.3719	0.3500	-----	0.3610	0.45	0.51	0.48	-----	-----	817
0.4213	0.4554	0.4561	0.4443	0.51	0.60	0.56	-----	-----	1040
0.4112	0.4080	0.4098	0.4097	0.52	0.31	0.42	-----	-----	1042
			0.4239			0.46	26.42	-----	
0.4289	0.4260	-----	0.4275	0.39	0.41	0.40	26.64	-----	628
0.5671	0.4833	-----	0.5252	0.41	0.45	0.43	-----	-----	971
0.4981	0.4975	0.5411	0.5122	0.43	0.39	0.41	-----	-----	995
			0.5182			0.42	32.29	-----	
0.4462	0.4396	0.4503	0.4454	0.34	0.53	0.44	27.76	-----	966
0.4461	0.4674	-----	0.4568	0.16	-----	0.16	-----	-----	271
0.5570	0.5653	-----	0.5612	0.10	0.06	0.08	-----	-----	627
0.4063	0.4364	-----	0.4214	0.08	0.07	0.08	-----	Coarse-grained	702
0.6332	0.6769	-----	0.6551	0.02	0.03	0.03	-----	Grown near the ocean	704
0.5215	0.5256	-----	0.5233	0.03	0.01	0.02	-----	Coarse-grained best quality	705
0.5493	0.5967	-----	0.5715	0.10	0.06	0.08	-----	Fine-grained best quality	706
0.6189	0.5785	-----	0.5987	0.02	0.03	0.03	-----	Coast fir	798
0.4254	0.3859	-----	0.4057	0.04	0.03	0.04	-----	20 miles from coast, clear yellow	709
0.5269	0.5215	-----	0.5242	0.09	0.12	0.11	-----	-----	720
0.4896	0.4941	-----	0.4918	0.12	0.13	0.13	-----	-----	732
0.5251	0.5855	-----	0.5553	0.18	0.18	0.18	-----	-----	881
0.4909	0.4679	-----	0.4744	0.05	0.06	0.06	-----	Red fir	973
0.5028	0.4848	-----	0.4988	0.05	0.04	0.05	-----	Yellow fir	974
0.4511	0.4504	-----	0.4508	0.10	0.08	0.09	-----	-----	986
0.5690	0.5922	-----	0.5806	0.10	0.11	0.11	-----	-----	989

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
391. <i>Pseudotsuga Douglasii</i> —continued	1008	British Columbia	Saw-mill, Burrard inlet.	G. Engelmann and C. S. Sargent.				
	1011	Oregon	Oregon Railway and Navigation Co.	do				
	1016	do	Weidler's saw-mill, Portland.	do				
	1018	do	Saw-mill, Astoria	do				
	1020	do	Portland Furniture Company.	do				
	1022	do	Portland Furniture Company.	do				
391. <i>Pseudotsuga Douglasii</i> , var. <i>macrocarpa</i> . <i>Hemlock.</i>	642	California	Saw-mill, San Bernardino.	W. G. Wright				
392. <i>Abies Fraseri</i> <i>Balsam. She Balsam.</i>	523	North Carolina	Roan mountain	Walcott Gibbs	Peaty loam	0.180	46	
393. <i>Abies balsamea</i> . <i>Balsam Fir. Balm of Gilead Fir.</i>	107	Vermont	Green mountains	C. G. Pringle	Cold, gravelly loam			
	877	do	Monkton	do	Peaty			
394. <i>Abies subalpina</i> <i>Balsam.</i>	449 ¹	Colorado	Forest City	T. S. Brandegee	Moist, sandy loam			
	449 ²	do	do	do	do			
	820	do	do	do	do	0.244	17	155
395. <i>Abies grandis</i> <i>White Fir.</i>	959							
	1009	Oregon	Portland	G. Engelmann and C. S. Sargent.	Rich, alluvial	0.735	56	43
	1010							
396. <i>Abies concolor</i> <i>White Fir. Balsam Fir.</i>	529	Colorado	Engelmann's cañon	Robert Douglas	Rocky	0.196	29	
	580	California	Strawberry valley	G. Engelmann and C. S. Sargent.	Alluvial			
	639	do	do	do	do			
	733	do	Lassen's peak	Sierra Lumber Company.				
397. <i>Abies bracteata</i>	572	do	Santa Lucia mountains	G. R. Vasey				
398. <i>Abies amabilis</i>	1004	British Columbia	Silver peak, near Fraser river.	G. Engelmann and C. S. Sargent.	Rich, sandy loam	0.584	50	120
399. <i>Abies nobilis</i> <i>Red Fir.</i>	965	Oregon	Cascade mountains	do	Rich			
400. <i>Abies magnifica</i> <i>Red Fir.</i>	647	California	Soda Springs	do	Gravelly loam	1.324	71	267
401. <i>Larix Americana</i> . <i>Larch. Black Larch. Tamarack. Hackmatack.</i>	226 ¹	Vermont	Charlotte	C. G. Pringle	Cold, swampy			
	226 ²	do	do	do	do			
	226 ³	do	do	do	do			
	774	New Brunswick	Bay of Fundy	Intercolonial railway.				
	781	do	do	do				
	786	do	Bridgeton	Ed. Sinclair				
	795	do	Danville	Grand Trunk railway.				
	840	Massachusetts	Wenham	J. Robinson	Swampy			

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.5266			0.5266	0.03	0.04	0.04			1008
0.4902	0.4867		0.4885	0.03	0.03	0.03		Railroad tie	1011
1.5831	0.5599		0.5715	0.11	0.07	0.09			1016
0.5660	0.5725		0.5692	0.05	0.04	0.05			1018
0.4331	0.4422		0.4377	0.07	0.11	0.09			1020
0.4362	0.4588		0.4575	0.06	0.07	0.07			1022
			0.5157			0.08	32.14		
0.4064	0.4463		0.4563	0.08	0.08	0.08	28.44		642
0.3699	0.3431		0.3565	0.47	0.60	0.54	22.22		523
0.3386	0.3342		0.3364	0.63	0.49	0.56			107
0.4248	0.4299		0.4273	0.33	0.35	0.34			877
			0.3819			0.45	23.80		
0.3446	0.3515		0.3481	0.66		0.66			449 ¹
0.3442	0.3641		0.3541	0.33		0.33			449 ²
0.3475	0.3397	0.3342	0.3405	0.34	0.38	0.34			820
			0.3476			0.44	21.66		
0.3416	0.3371			0.38	0.32				950
0.3643	0.3842	0.3531	0.3545	0.63	0.57	0.49	21.97	One tree	1009
0.3390	0.3625			0.56	0.47				1010
0.3613	0.3717		0.3665	0.87	0.89	0.88			529
0.3059	0.2910		0.2985	1.12	1.12	1.12			596
0.3286	0.3780		0.3533	0.56	0.48	0.52			639
0.4242	0.4500		0.4371	0.70	1.04	0.87			738
			0.3638			0.85	22.67		
0.4082	0.6483		0.6783	2.09	1.09	2.04	42.27		572
0.4297	0.4159		0.4228	0.25	0.21	0.23	26.35		1004
0.4625	0.4497		0.4561	0.27	0.41	0.34	28.42		965
0.4752	0.4614	0.4738	0.4701	0.30	0.30	0.30	29.30		647
0.6973	0.7075		0.7024	0.26		0.26			226 ¹
0.6945	0.7779		0.7362	0.27		0.27			226 ²
0.6835	0.6069		0.6452	0.32		0.32			226 ³
0.5975	0.5765		0.5870	0.30	0.27	0.28			774
0.5925	0.5740		0.5833	0.47	0.49	0.48			781
0.5389	0.5570		0.5479	0.35	0.30	0.33			786
0.5862	0.5864		0.5873	0.36	0.32	0.34			795
		0.6000	0.6000	0.27	0.48	0.38			840
			0.6236			0.33	33.86		

FOREST TREES OF NORTH AMERICA.

TABLE I.—SPECIFIC GRAVITY, ASH, AND WEIGHT PER CUBIC FOOT

Species.	Office number.	State.	Locality.	Collector.	Soil.	Diameter of tree, in meters.	LAYERS OF GROWTH.	
							Sap-wood.	Heart-wood.
402. <i>Larix occidentalis</i> <i>Tamarack.</i>	719	Montana.....	Missoula.....	S. Watson.....				
	984	Washington ter- ritory.	Fulda.....	William N. Siksdorf.	Moist.....			
	1008	do.....	do.....	do.....				
PALMACEÆ.								
404. <i>Sabal Palmetto</i> <i>Cabbage Tree. Cabbage Palmetto.</i>	242	Florida.....	Sister Island.....	A. H. Curtiss.....	Shell.....			
405. <i>Washingtonia filifera</i> <i>Fan-leaf Palm.</i>	1159	California.....	Agua Caliente.....	W. G. Wright.....	Dry, gravelly.....			
406. <i>Thrinax parviflora</i> <i>Silk-top Palmetto.</i>	504	Florida.....	Long Key.....	A. H. Curtiss.....	Coral.....			
	1107	do.....	Bahia Honda Key.....	do.....	do.....			
407. <i>Thrinax argentea</i> <i>Silver-top Palmetto. Brickley Thatch.</i>	508	do.....	Sugar-loaf Sound.....	do.....	do.....			
	1116	do.....	No-Name Key.....	do.....	do.....			
408. <i>Oreodoxa regia</i> <i>Royal Palm.</i>	505	do.....	Big Cypress.....	do.....	Rich humus.....			
	1119	do.....	Lost Man's river.....	do.....	do.....			
LILIACEÆ.								
409. <i>Yucca canaliculata</i> <i>Spanish Bayonet.</i>	1063	Texas.....	Matagorda.....	C. Mohr.....	Sandy, saline.....			
410. <i>Yucca brevifolia</i> <i>The Joshua. Joshua Tree.</i>	695	California.....	Mohave.....	G. Engelmann and C. S. Sargent.	Dry, gravelly.....			
	696	do.....	do.....	do.....	do.....			
411. <i>Yucca elata</i> <i>Spanish Bayonet.</i>	699	Arizona.....	Tucson.....	do.....	do.....			
412. <i>Yucca baccata</i> <i>Spanish Bayonet. Mexican Banana.</i>	1003	California.....	San Diego.....	G. Engelmann.....	do.....			

OF DRY SPECIMENS OF THE WOODS OF THE UNITED STATES—Continued.

SPECIFIC GRAVITY DETERMINATIONS.				ASH DETERMINATIONS.			Weight, per cubic foot, in pounds (average).	Remarks.	Office number.
First.	Second.	Third.	Average.	First.	Second.	Average.			
0.6663	0.5897		0.6280	0.08	0.09	0.09			719
0.7717	0.7839		0.7778	0.09	0.09	0.09			984
0.8340	0.7984		0.8162	0.07	0.12	0.09			1006
			0.7407			0.09	48.16		
0.5002	0.4020	0.4191	0.4404	7.51	7.80	7.66	27.44		243
0.5597	0.4942	}	0.5173	1.80	1.98	1.89	32.24		1159
0.4700	0.5453								
0.7014	0.7258		0.7136	2.65	2.79	2.72			504
0.4539	0.4987	0.5039	0.4845	5.73	4.81	5.27			1107
			0.5991			3.99	37.34		
0.7857	0.7468		0.7412	1.71	2.19	1.95			596
0.6775	0.6907	0.7118	0.6933	3.90	4.25	4.07			1116
			0.7173			3.01	44.70		
0.2370	0.1887		0.2128	2.96	2.53	2.74		Pth ... } One tree	565
0.8178	0.8807		0.8492	1.31	1.14	1.23		Rind ... }	
0.7451	0.7513		0.7482	2.79	2.53	2.66		Rind ...	1119
			0.6034			2.21	37.60		
0.7162	0.5875	0.6994	0.6677	7.08	5.52	6.27	41.61		1063
0.3473	0.3472		0.3472	5.53	4.75	5.14			695
0.4002	0.4002		0.4002	2.66	3.08	2.87			696
			0.3737			4.00	23.29		
0.2327	0.6511	0.4572	0.4470	8.50	10.06	9.28	27.86		699
0.1990	0.3458		0.2724	8.94		8.94	16.97		1008

FOREST TREES OF NORTH AMERICA.

TABLE II.—ACTUAL FUEL VALUE OF SOME OF THE

Office number.	Catalogue number.	Botanical name.	Common name.	Region.	FUEL VALUE.	
					Per cubic decimeter.	Per kilogram.
165	8	<i>Liriodendron Tulipifera</i>	Tulip Tree. Yellow Poplar. White Wood.....	Atlantic	1425.57	3744.61
908	29	<i>Bursera gummifera</i>	Gum Elemi. Gumbo Limbo. West Indian Birch	Semi-tropical Florida	997.32	2913.58
453	31	<i>Swietenia Mahogoni</i>	Mahogany. Madeira	do	2769.31	3802.95
274	64	<i>Acer saccharinum</i> , var. <i>nigrum</i>	Black Sugar Maple.....	Atlantic	3091.37	4345.48
845	77	<i>Robinia Pseudacacia</i>	Locust. Black Locust. Yellow Locust.....	do	2822.99	3890.02
927	93	<i>Prosopis juliflora</i>	Mesquit. Algaroba. Honey Locust. Honey Pod.	Mexican boundary	3291.21	4352.30
883	115	<i>Cercocarpus ledifolius</i>	Mountain Mahogany.....	Interior Pacific	4234.06	4052.90
1182	139	<i>Liquidambar Styraciflua</i>	Sweet Gum. Star-leaved Gum. Liquidamber. Red Gum. Bilsted.	Atlantic	2255.24	4016.46
128	155	<i>Nyssa uniflora</i>	Large Tupelo. Cotton Gum. Tupelo Gum.....	Southern Atlantic	2332.41	4181.83
1084	184	<i>Diospyros Virginiana</i>	Persimmon.....	Atlantic	2970.45	3781.61
227	193	<i>Fraxinus Americana</i>	White Ash.....	do	2652.84	4217.42
180	207	<i>Catalpa speciosa</i>	Western Catalpa.....	do	1532.42	3036.38
533	224	<i>Ulmus Americana</i>	White Elm. American Elm. Water Elm.....	do	3247.02	4191.87
126	235	<i>Platanus occidentalis</i>	Sycamore. Button Wood. Button-ball Tree. Water Beech.	do	2406.89	4071.83
209	239	<i>Juglans nigra</i>	Black Walnut.....	do	1984.56	3857.26
322	241	<i>Carya oliviformis</i>	Pecan. Illinois Nut	do	2768.72	3954.75
29	242	<i>Carya alba</i>	Shell-bark Hickory. Shag-bark Hickory	do	3851.17	4078.76
539					3319.79	3811.48
72	244	<i>Carya tomentosa</i>	Mocker Nut. Black Hickory. Bull Nut. Big- bud Hickory. White-heart Hickory. King Nut.	do	3380.57	3904.11
1051	245	<i>Carya porcina</i>	Pig Nut. Brown Hickory. Black Hickory. Switch-bud Hickory.	do	3392.12	3922.89
838	246	<i>Carya amara</i>	Bitter Nut. Swamp Hickory	do	2863.42	3903.25
237	247	<i>Carya myristiciformis</i>	Nutmeg Hickory.....	Southern Atlantic	3108.27	3877.58
362	248	<i>Carya aquatica</i>	Water Hickory. Swamp Hickory. Bitter Pecan	do	3140.23	4073.59
1050	251	<i>Quercus alba</i>	White Oak	Atlantic	3197.41	4187.83
988	253	<i>Quercus Garryana</i>	do	Northern Pacific.....	2594.31	3667.39
424	257	<i>Quercus lyrata</i>	Over-cup Oak. Swamp Post Oak. Water White Oak.	Southern Atlantic	3268.92	4105.65
31	260	<i>Quercus Prinus</i>	Chestnut Oak. Rock Chestnut Oak.....	do	2843.69	3997.32
92	272	<i>Quercus rubra</i>	Red Oak. Black Oak	Atlantic	3062.08	4075.16
247	274	<i>Quercus tinctoria</i>	Black Oak. Yellow-bark Oak. Quercitron Oak. Yellow Oak.	do	2595.04	3774.60
339	276	<i>Quercus nigra</i>	Black Jack. Jack Oak	do	2692.51	3713.81
548	277	<i>Quercus falcata</i>	Spanish Oak. Red Oak.....	Southern Atlantic	3193.28	4055.48
511	280	<i>Quercus aquatica</i>	Water Oak. Duck Oak. Possum Oak. Punk Oak	do	2655.82	3718.07
868	290	<i>Castanea vulgaris</i> , var. <i>Americana</i>	Chestnut.....	Atlantic	1868.25	4042.96
55	291	<i>Fagus ferruginea</i>	Beech.....	do	2795.34	3895.04
848	294	<i>Betula alba</i> , var. <i>populifolia</i>	White Birch. Old-field Birch. Gray Birch.....	Northern Atlantic	2509.00	4078.05
225	295	<i>Betula papyrifera</i>	Canoe Birch. White Birch. Paper Birch	do	2582.66	4101.41
272	318	<i>Populus tremuloides</i>	Aspen. Quaking Asp.....	Atlantic and Pacific	1624.64	4292.31
754	324	<i>Populus monilifera</i>	Cottonwood. Necklace Poplar. Carolina Pop- lar. Big Cottonwood.	Atlantic	1906.42	4242.15
874	327	<i>Thuja occidentalis</i>	White Cedar. Arbor-vitæ	Northern Atlantic	1411.57	3917.77
701	331	<i>Chamaecyparis Lawsoniana</i>	Port Orford Cedar. Oregon Cedar. White Ce- dar. Lawson's Cypress. Ginger Pine.	Northern Pacific	2327.52	5263.50
527	338	<i>Juniperus occidentalis</i> , var. <i>monosperma</i>	Juniper	Pacific.....	3143.57	4587.81
923	340	<i>Taxodium distichum</i>	Bald Cypress. Black Cypress. Red Cypress. White Cypress. Deciduous Cypress.	Southern Atlantic	1935.71	4789.73
711	342	<i>Sequoia sempervirens</i>	Redwood	California coast.....	1985.50	4191.47
1044	347	<i>Pinus Strobus</i>	White Pine. Weymouth Pine	Northern Atlantic	1489.03	4272.69
638	349	<i>Pinus Lambertiana</i>	Sugar Pine	Pacific	1785.40	4419.31
900	356	<i>Pinus monophylla</i>	Piton. Nut Pine	Interior Pacific	2248.13	4107.68
194	358	<i>Pinus resinosa</i>	Red Pine. Norway Pine.....	Northern Atlantic	2051.75	4226.05
632	361	<i>Pinus ponderosa</i>	Yellow Pine. Bull Pine	Pacific.....	2441.24	4600.04
625	365	<i>Pinus Murrayana</i>	Tamarack. Black Pine. Lodge-pole Pine. Spruce Pine.	do	1791.32	4019.12
571	366	<i>Pinus Sabiniana</i>	Digger Pine. Bull pine.....	California.....	1804.29	3962.97
389	370	<i>Pinus Taeda</i>	Loblolly Pine. Old-field Pine. Rosemary Pine.	Southern Atlantic	2031.75	4087.20
1046	371	<i>Pinus rigida</i>	Pitch Pine.....	Atlantic coast	3472.26	5491.47
83	372	<i>Pinus serotina</i>	Pond Pine.....	Southern Atlantic	3980.96	5012.54
621	373	<i>Pinus inops</i>	Jersey Pine. Scrub Pine	Atlantic	2008.20	4126.15
321	375	<i>Pinus pungens</i>	Table-mountain Pine. Hickory Pine	Alleghany	2054.78	3995.30
557	377	<i>Pinus mitis</i>	Yellow Pine. Short-leaved Pine. Spruce Pine. Bull Pine.	Atlantic	3091.92	5062.75

MORE IMPORTANT WOODS OF THE UNITED STATES.

RELATIVE FUEL VALUE.		PERCENTAGE IN DRY WOOD.						Specific gravity.	Weight of cubic foot, in pounds.	Office number.
By volume.	By weight.	Ash.	Hydrogen.	Carbon.	Oxygen.	Hydrogen combined with oxygen.	Excess of hydrogen.			
67	66	0.27	6.43	47.29	46.01	5.75	0.68	0.8907	23.72	165
69	70	2.09	6.02	40.80	51.09	6.39	0.36	0.8423	21.33	903
32	62	1.09	6.69	46.76	45.46	5.68	1.01	0.7282	45.88	453
23	19	0.56	6.61	51.55	41.28	5.16	1.45	0.7114	44.32	274
29	58	0.28	6.17	49.19	44.41	5.55	0.62	0.7257	45.22	845
14	18	2.05	6.61	51.08	40.26	5.08	1.58	0.7562	47.12	927
3	42	1.20	5.45	52.14	41.21	5.15	0.30	1.0447	65.10	883
45	46	0.48	5.85	50.99	42.88	5.33	0.52	0.5615	34.99	1182
43	39	0.74	6.97	48.78	43.51	5.44	1.53	0.5645	35.17	128
25	63	0.77	6.44	47.37	45.42	5.67	0.77	0.7855	48.95	1084
36	24	0.30	6.98	49.73	43.04	5.38	1.55	0.6289	39.19	227
65	52	0.47	6.92	47.44	45.17	5.65	1.27	0.4020	25.05	180
16	26	0.74	6.57	50.35	42.34	5.29	1.28	0.7746	48.27	533
42	40	0.57	5.83	51.45	42.15	5.27	0.56	0.5611	36.63	126
58	60	0.56	6.00	49.28	44.16	5.52	0.48	0.5145	32.06	209
33	50	0.95	6.15	49.51	43.39	5.42	0.73	0.7001	43.63	322
7	36	0.73	6.49	49.67	43.12	5.39	1.10	0.9442	58.84	29
13	61	0.83	6.13	48.45	44.59	5.57	0.56	0.8710	54.28	539
11	55	1.04	5.93	49.69	43.84	5.42	0.51	0.8659	53.95	72
10	53	0.74	6.28	48.98	44.00	5.50	0.78	0.8647	53.88	1051
27	56	1.03	5.91	49.71	43.29	5.41	0.50	0.7336	45.71	833
21	59	1.06	6.37	48.26	44.31	5.54	0.83	0.8016	49.95	237
20	38	1.19	6.60	49.16	43.05	5.38	1.22	0.7709	43.04	362
17	28	0.24	6.59	50.44	42.73	5.34	1.25	0.7635	46.58	1060
38	69	0.33	5.73	48.56	45.38	5.67	0.06	0.7074	44.08	968
15	33	0.58	6.75	49.22	43.45	5.43	1.32	0.7963	49.61	424
23	47	0.34	6.33	49.59	43.74	5.47	0.86	0.7114	44.32	31
24	37	0.15	6.62	49.49	43.74	5.47	1.15	0.7514	46.72	92
37	64	0.15	6.09	48.78	44.98	5.62	0.37	0.6875	43.84	247
34	68	1.37	5.73	48.58	44.32	5.54	0.19	0.7250	45.18	339
18	41	0.29	6.14	50.58	42.99	5.37	0.77	0.7874	49.07	548
35	67	0.88	5.75	48.73	44.69	5.58	0.17	0.7143	44.51	511
56	43	0.13	5.70	51.74	42.43	5.30	0.40	0.4621	23.30	868
31	57	0.54	6.11	49.27	44.08	5.51	0.60	0.7175	44.71	55
40	39	0.29	6.49	49.77	43.45	5.43	1.06	0.6160	33.05	848
39	34	0.23	7.12	48.28	44.37	5.54	1.53	0.6297	39.24	225
63	20	0.74	6.58	51.13	41.55	5.19	1.39	0.3785	23.59	272
55	22	0.65	6.26	51.64	41.45	5.18	1.08	0.4494	23.00	754
68	54	0.37	6.37	48.80	44.46	5.56	0.81	0.3603	22.45	874
44	3	0.10	6.28	50.07	33.65	4.21	2.07	0.4422	27.56	701
19	11	0.88	6.03	54.97	38.12	4.76	1.27	0.6852	42.70	527
54	8	0.40	6.54	54.98	38.08	4.76	1.78	0.4084	24.45	923
52	27	0.13	6.01	52.10	41.70	5.22	0.79	0.4737	29.52	711
66	21	0.12	6.08	52.55	41.25	5.15	0.93	0.3485	21.72	1044
59	13	0.19	6.40	52.85	40.56	5.07	1.33	0.4040	25.18	638
46	32	0.83	6.39	50.48	43.30	5.41	0.98	0.5473	34.11	900
49	23	0.20	6.07	52.18	41.55	5.19	0.88	0.4855	30.26	194
41	9	0.31	7.02	52.60	40.07	5.01	2.01	0.5307	33.07	632
58	45	0.37	6.22	50.05	43.86	5.42	0.80	0.4457	27.78	625
57	49	0.42	6.04	50.22	43.32	5.41	0.63	0.4530	28.28	571
50	35	0.25	6.23	50.60	42.92	5.36	0.87	0.4971	30.98	389
9	1	1.12	7.19	59.00	32.68	4.08	3.11	0.6323	39.40	1046
6	6	0.17	6.30	53.55	36.48	4.56	2.24	0.7942	49.49	83
51	31	0.26	6.30	50.74	42.70	5.34	0.96	0.4867	30.33	32
48	48	0.30	5.78	51.07	42.85	5.35	0.43	0.5143	32.05	331
22	5	0.20	6.91	56.64	36.25	4.53	2.38	0.6107	38.06	557

FOREST TREES OF NORTH AMERICA.

TABLE II.—ACTUAL FUEL VALUE OF SOME OF THE

Office number.	Catalogue number.	Botanical name.	Common name.	Region.	FUEL VALUE.	
					Per cubic decimeter.	Per kilogram.
879	879	<i>Pinus Banksiana</i>	Gray Pine. Scrub Pine. Prince's Pine.....	Northern Atlantic.....	2152.66	4393.18
a 359	380	<i>Pinus palustris</i>	{ Long-leaved Pine. Southern Pine. Georgia } Pine. Yellow Pine. Hard Pine.	South Atlantic coast.....	4344.88	5123.64
81					4064.77	5480.35
385					2810.20	4506.42
b 358					4129.95	4594.97
358	381	<i>Pinus Cubensis</i>	Slash Pine. Swamp Pine. Bastard Pine. Meadow Pine.	do.....	4319.67	4806.05
356					3363.40	4418.55
794	382	<i>Picea nigra</i>	Black Spruce.....	Northern Atlantic.....	1614.11	3949.37
1042	387	<i>Tsuga Canadensis</i>	Hemlock.....	do.....	1724.25	4298.58
709	391	<i>Pseudotsuga Douglasii</i>	Red Fir. Yellow Fir. Oregon Pine. Douglas Fir.	Pacific.....	1766.32	4354.84
226	401	<i>Larix Americana</i>	Larch. Black Larch. Tamarack. Hackmatack.	Northern Atlantic.....	2937.46	4182.04
242	404	<i>Sabal Palmetto</i>	Cabbage Tree. Cabbage Palmetto.....	South Atlantic coast.....	1653.35	3754.21
c 565	408	<i>Oreodoxa regia</i>	Royal Palm.....	Semi-tropical Florida.....	859.07	4037.01
d 565					3708.48	4367.03

a Boxed for turpentine 1876; chipped 4 years; specimen taken along chip.
b Boxed for turpentine 1882; chipped 10 years; abandoned 1861.

c Pith.
d Rind.

MORE IMPORTANT WOODS OF THE UNITED STATES—Continued.

RELATIVE FUEL VALUE.		PERCENTAGE IN DRY WOOD.						Specific gravity.	Weight of cubic foot, in pounds.	Office number.
By volume.	By weight.	Ash.	Hydrogen.	Carbon.	Oxygen.	Hydrogen combined with oxygen.	Excess of hydrogen.			
47	15	0.19	6.29	52.93	40.59	5.07	1.23	0.4960	30.54	879
1	4	0.15	7.26	56.19	36.30	4.54	2.72	0.8479	52.84	359
5	2	0.26	7.41	58.61	33.72	4.21	3.20	0.7417	46.22	81
20	12	0.28	6.70	52.70	40.32	5.04	1.66	0.6286	38.86	385
4	10	0.28	6.85	52.99	39.88	4.99	1.86	0.8968	55.96	358
2	7	0.24	6.83	54.78	38.15	4.77	2.06	0.8968	55.96	358
12	14	0.16	6.22	53.33	40.29	5.08	1.19	0.7612	47.44	356
64	51	0.30	6.58	48.45	44.67	5.58	1.00	0.4067	25.47	794
61	25	0.48	5.91	52.88	41.23	5.15	0.76	0.4067	25.58	1042
60	17	0.03	6.42	52.32	41.23	5.15	1.27	0.4066	25.28	709
26	29	0.27	6.03	51.91	41.79	5.22	0.81	0.7024	43.77	226
62	65	7.06	7.06	43.35	41.98	5.24	1.82	0.4404	27.45	242
70	44	2.74	6.82	47.73	42.71	5.94	1.48	0.2128	12.26	565
8	16	1.24	6.98	50.46	41.32	5.16	1.82	0.2492	15.92	565

23 FOR

TABLE III.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Specific gravity of the air-dried specimen.	Direction of grain.	COEFFICIENT OF ELASTICITY.		Modulus of rupture.
								First deflection.	Second deflection.	
MAGNOLIACEÆ.										
1. <i>Magnolia grandiflora</i> <i>Big Laurel. Bull Bay.</i>	346	Alabama	Cottage Hill	C. Mohr	Rich loam	0.7051		1061	957	830
	346	do	do	do	do	0.7006		887	849	761
2. <i>Magnolia glauca</i> <i>Sweet Bay. White Bay. Beaver Tree. White Laurel. Swamp Laurel.</i>	354	do	do	do	Swampy	0.5223		976	980	.33
	354	do	do	do	do	0.5028		976	886	717
3. <i>Magnolia acuminata</i> <i>Ocumber Tree. Mountain Magnolia.</i>	746	Virginia	Wytheville	H. Shriver	Clay limestone	0.5000		904	1061	708
	246	do	do	do	do	0.4755		939	921	665
	261 ¹	do	Fancy Gap	do	Rich, light	0.4862		788	784	574
	261 ¹	do	do	do	do	0.4564		729	723	525
	534	Mississippi	Selvers' mill	C. Mehr	do	0.5682		1061	1050	717
	534	do	do	do	do	0.5852		1109	1085	890
4. <i>Magnolia cordata</i> <i>Ocumber Tree.</i>	1178	Alabama	Winston county	do	do	0.4318		888	864	564
	1178	do	do	do	do	0.4625		976	1017	637
5. <i>Magnolia macrophylla</i> <i>Large-leaved Ocumber Tree.</i>	532	Mississippi	Quitman	do	Rich, low	0.5807		1191	1085	516
	532	do	do	do	do	0.5580		1252	1252	876
6. <i>Magnolia Umbrella</i> <i>Umbrella Tree. Elk Wood.</i>	266 ¹	Virginia	Wytheville	H. Shriver	do	0.4170		800	814	553
	266 ²	do	do	do	do	0.5051		669	673	612
7. <i>Magnolia Fraseri</i> <i>Long-leaved Ocumber Tree.</i>	260	do	Fancy Gap	do	Damp	0.5688		939	976	811
	260 ¹	do	do	do	do	0.5103		887	912	602
8. <i>Liriodendron Tulipifera</i> <i>Tulip Tree. Yellow Poplar. White Wood.</i>	395	Michigan	Lansing	W. J. Beal	do	0.4174		610	610	473
	818	West Virginia	Grafton	C. G. Pringle	do	0.4930		1085	1176	756
	818	do	do	do	do	0.4809		1085	1050	675
	1231	Pennsylvania	Chester county	P. P. Sharples	do	0.4704		976	976	743
	1231	do	do	do	do	0.4911		921	1007	820
	1231	do	do	do	do	0.4619		976	957	755
	1232	do	do	do	do	0.4381		976	848	457
	1232	do	do	do	do	0.4511		976	1028	723
	1232	do	do	do	do	0.4550		904	921	661
	1236	Tennessee	Saw-mill at Nashville.	A. E. Baird	do	0.4614		888	834	604
	1236	do	do	do	do	0.4697		872	888	645
	1236	do	do	do	do	0.4591		888	849	584
	1237	do	do	do	do	0.4386		857	849	584
	1238	do	do	do	do	0.4465		976	976	635
ANONACEÆ.										
9. <i>Asimina triloba</i> <i>Papaw. Oustard Apple.</i>	211	Missouri	Meramec river, Jefferson county.	G. W. Letterman	Alluvial	0.3634		444	467	313
	211	do	do	do	do	0.3575		362	315	238
	332	Tennessee	Cumberland river	A. Gittinger	do	0.4323		696	723	574
10. <i>Anona laurifolia</i> <i>Pond Apple.</i>	479	Florida	Bay Biscayne	A. H. Curtiss	Swampy	0.5705		542	514	638
	479	do	do	do	do	0.5794		530	488	588
CANELLACEÆ.										
12. <i>Canella alba</i> <i>White Wood. Cinnamon Bark. Wild Cinnamon.</i>	1131	do	Elliott's Key	do	Coral	1.1286		1085	1085	905
	1131	do	do	do	do	1.0857		1085	1148	1148
TERNSTROMIACEÆ.										
14. <i>Gordonia Lasianthus</i> <i>Loblolly Bay. Ten Bay.</i>	236	South Carolina	Bonneau's Depot	H. W. Ravenel	Wet pine-barren	0.4844		718	723	499
	236	do	do	do	do	0.4729		787	763	671
	414	do	Aiken	do	Swampy	0.5470		888	921	773
	414	do	do	do	do	0.5005		751	769	729

UNITED STATES UNDER TRANSVERSE STRAIN.

DEFLECTION, IN MILLIMETERS, UNDER A PRESSURE, IN KILOGRAMS, OF—													Ultimate strength: transverse pressure.	Remarks.	Office number.
50	100	150	200	0 (set.)	200	250	300	350	400	450	500	550			
4.6	10.2	16.0	22.8	1.6	22.5	30.5	39.5	54.6					350	Broke suddenly with long splinters	346
5.5	11.5	16.7	23.4	2.0	23.6	31.2	40.5						325	do	346
5.0	10.5	16.0	21.7	1.0	22.0	30.0	44.5						321	Crushed at center bearing; broke suddenly and squarely, splitting to the end.	354
5.0	10.9	16.7	24.0	2.3	24.0	31.5	50.8						306	do	354
5.4	9.2	14.5	20.0	1.0	20.0	28.8	46.0						300	Crushed at center bearing; broke with fine splinters	346
5.2	10.6	17.0	24.0	1.9	25.0	33.7							284	do	346
6.2	13.3	21.0	34.3	4.8	35.0								245	do	361 ¹
6.7	13.5	21.0	31.8	3.4	33.0								224	do	361 ¹
4.6	9.3	14.5	20.0	1.0	20.0	27.0							306	do	384
4.4	9.0	13.7	19.2	0.7	19.2	26.0	35.0	54.0					358	do	384
5.5	11.3	17.3	26.1	2.0	27.0								240	Crushed at center bearing; square break	1178
5.0	9.6	15.2	22.0	1.4	23.0	32.4							272	Crushed at center bearing; splintered break	1178
4.1	9.0	13.8	19.2	1.0	19.5								220	Specimen cross-grained; started at knot	583
3.9	7.8	11.6	15.8	0.3	15.8	20.8	28.5	40.0					374	Broke with fine splinters	583
6.1	12.0	19.4	30.8	4.5	32.4								236	Splinters started at small knots	200 ²
7.3	14.5	23.0	34.8	4.0	36.2	66.0							261	do	200 ²
5.2	10.0	14.4	19.6	1.0	20.0	26.3	37.2						346	Crushed at center bearing; broke with fine splinters	200
5.5	10.7	16.4	24.7	2.0	25.5	36.8							257	Specimen cross-grained; broke with long splinters	200 ¹
8.0	16.0	26.0											202	Square break	306
4.5	8.3	12.0	16.8	0.4	16.9	24.5	34.0						322	Broke with long splinters	818
4.5	9.3	14.3	20.0	0.7	20.2	28.8							288	Crushed at center bearing; broke with long splinters	818
5.0	10.0	15.0	20.6	0.5	20.7	29.0	37.7						317	Crushed at center bearing; broke with large splinters	1261
5.3	9.7	14.4	20.0	0.5	20.5	25.8	34.2	50.0					350	do	1261
5.0	10.2	15.2	20.6	0.5	20.6	27.0	37.7						322	Crushed at center bearing; square break	1261
5.0	11.5	18.0											195	Crushed at center bearing; square break at large knot	1262
5.0	9.5	14.0	19.2	0.5	19.4	25.4	36.0						306	Crushed at center bearing; broke with fine splinters	1262
5.4	10.6	15.7	21.5	0.6	22.0	30.0							283	Crushed at center bearing; broke with long splinters at knot	1332
6.0	11.7	16.0	22.0	1.0	22.4	31.5							296	Square break on tension side, splitting in axis	1236
5.6	11.0	16.0	23.0	1.4	23.5	33.0							275	Specimen cross-grained; broke with the grain	1236
5.5	11.5	17.5	26.6	2.0	27.0								249	Specimen cross-grained; started at the angle of two faces	1236
5.7	11.5	17.2	27.0	2.5	28.0								249	Square break	1237
5.0	10.0	15.2	22.2	1.5	22.4	31.6							271	do	1238
11.0	24.0												133	Broke with large splinters	211
13.5	31.0												123	do	211
7.0	13.5	21.6	31.0	2.9	33.3								245	Broke with splinters at knot	332
9.0	19.0	31.0	50.5	10.0	55.4	94.0							268	Crushed at center bearing; broke with splinters	479
9.2	20.0	32.5	51.5	10.0	54.3	83.0							250	Square break	479
4.5	9.0	13.7	18.5	1.0	19.5	25.0	31.0	38.0					386	Specimen defective, 0.75 sap-wood; square break	1181
4.5	8.5	12.5	17.5	0.8	17.8	22.0	28.0	34.0	42.0	50.0			490	0.75 sap-wood; broke with long, coarse splinters	1181
6.8	13.5	21.5	32.0	3.0	32.5								213	Sap-wood; broke at knot with large splinters	236
6.2	12.8	20.0	29.2	2.5	30.0	40.0							290	Crushed at center bearing; broke with long, fine splinters	236
5.5	10.6	16.3	23.5	1.2	23.7	31.0	44.0						330	do	414
6.5	12.7	19.5	27.0	2.0	28.2	37.0	58.0						311	Broke with long, coarse splinters	414

TABLE III.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Specific gravity of the air-dried specimen.	Direction of grain.	COEFFICIENT OF ELASTICITY.		Modulus of rupture.
								First deflection.	Second deflection.	
TILIACEÆ.										
17. <i>Tilia Americana</i> <i>Lime Tree. Bass Wood. American Linden. Lin. Bee Tree.</i>	2	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	0.4001		814	781	569
	252	Missouri	Allenton	G. W. Letterman	Alluvial	0.4068		976	976	616
	252	do	do	do	do	0.5496		1038	1027	804
	316	Michigan	Hersey	W. J. Beal	Rich loam	0.4517		888	888	525
	1089	Massachusetts	Danvers	J. Robinson	Moist gravel	0.4492		698	729	553
	1089	do	do	do	do	0.4287		668	638	469
17. <i>Tilia Americana</i> , var. <i>pubescens</i>	745	Georgia	Bainbridge	A. H. Curtiss	Low	0.4702		842	834	569
	745	do	do	do	do	0.4555		814	788	548
18. <i>Tilia heterophylla</i> <i>White Bass Wood. Wahoo.</i>	285 ¹	Kentucky	Cliffs Kentucky river	W. M. Linney	Limestone	0.4017		697	634	403
	285 ²	do	Mercer county	do	do	0.3944		751	723	525
	285 ³	do	do	do	do	0.3924		827	814	541
	320	Tennessee	Cumberland river	A. Gittinger	Alluvial	0.4750		1085	1085	703
	320	do	do	do	do	0.4924		976	976	712
MALPIGHIACEÆ.										
19. <i>Byrsonima lucida</i> <i>Tallowberry. Glanberry.</i>	1113	Florida	No-Name Key	A. H. Curtiss	Coral	0.6516		542	525	424
ZYGOPHYLLACEÆ.										
20. <i>Guaicum sanctum</i> <i>Logwood.</i>	476	do	Upper Metacombe Key	do	do	1.1430		800	814	628
	1123	do	Elkott's Key	do	do	1.1582		856	912	947
RUTACEÆ.										
23. <i>Xanthoxylum Clava-Herculis</i> <i>Toothache Tree. Prickly Ash. Sea Ash. Pepper Wood. Wild Orange.</i>	735	do	Chattahoochee	do	Dry, sandy	0.5240		651	698	628
	735	do	do	do	do	0.5353		669	688	708
	1066	Texas	Palestine	C. Mohr	Damp, sandy	0.5698		857	888	757
	1066	do	do	do	do	0.5545		659	630	469
24. <i>Xanthoxylum Caribæum</i> <i>Satin Wood.</i>	1140	Florida	Bahia Honda Key	A. H. Curtiss	Coral	0.9833		814	814	820
	1140	do	do	do	do	0.9545		814	921	687
SIMARUBEÆ.										
25. <i>Simaruba glauca</i> <i>Paradise Tree.</i>	487	do	Bay Biscayne	do	do	0.4580		888	888	586
	487	do	do	do	do	0.4104		1061	976	544
BURSERACEÆ.										
29. <i>Bursera gummiifera</i> <i>Gum Elemi. Gumbo Limbo. West Indian Birch.</i>	462	do	Upper Metacombe Key	do	do	0.3060		417		183
	462	do	do	do	do	0.3086				112
30. <i>Amyris sylvatica</i> <i>Torch Wood.</i>	475	do	do	do	do	1.1140		1038	1085	1244
	475	do	do	do	do	1.1273		976	1085	1366
MELIACEÆ.										
31. <i>Swietenia Mahagoni</i> <i>Mahogany. Madeira.</i>	452	do	do	do	do	0.8364		976	976	895
	452	do	do	do	do	0.8221		1109	1148	1111
ILICINEÆ.										
32. <i>Ilex opaca</i> <i>American Holly.</i>	280	South Carolina	Waverly Mills	W. St. J. Mazyck	Sandy loam	0.6990		651	622	703
	280	do	do	do	do	0.7018		707	664	670
34. <i>Ilex Dahoon</i> <i>Dahoon. Dahoon Holly.</i>	484	Florida	Bay Biscayne	A. H. Curtiss	Low, damp	0.5307		610	610	586
	484	do	do	do	do	0.5333		697	673	558
CYRILLACEÆ.										
37. <i>Cyrilla racemiflora</i> <i>Iron Wood.</i>	341	Alabama	Churchula	C. Mohr	Damp, sandy	0.6916		465	468	314
38. <i>Cliftonia ligustrina</i> <i>Tit. Iron Wood. Buckwheat Tree.</i>	338	do	Cottage Hill	do	Wet	0.6062		698	751	574
	338	do	do	do	do	0.6762		814	814	478

UNITED STATES UNDER TRANSVERSE STRAIN—Continued.

DEFLECTION, IN MILLIMETERS, UNDER A PRESSURE, IN KILOGRAMS, OF—														Ultimate strength: transverse pressure.	Remarks.	Office number.
50	100	150	200	0 (set.)	200	250	300	350	400	450	500	550				
6.0	12.5	20.3	32.0	4.0	34.0									243	Crushed at center bearing; broke with fine splinters.....	2
5.0	10.0	16.5	24.2	2.2	25.5	37.0								263	do.....	252
4.7	9.5	14.0	19.4	1.0	19.8	25.9	33.5							343	do.....	252
5.5	11.0	17.3	27.6	3.5	29.5									224	Crushed at center bearing; broke suddenly.....	316
7.0	13.4	22.3	34.3	4.0	36.0									236	Broke suddenly with large splinters.....	1099
7.3	15.3	26.4	50.5	18.0										200	Broke, on reloading, with large splinters.....	1099
5.8	11.7	18.5	28.6	3.5	30.6									243	Crushed at center bearing; square break.....	745
6.0	12.4	18.8	28.5	3.0	30.1									234	do.....	745
7.0	15.4	24.6												172	do.....	235 ¹
6.5	13.5	20.3	31.0	3.5	33.6									224	do.....	235 ²
5.9	12.0	18.4	29.6	3.5	31.3									231	do.....	235 ³
4.5	9.0	13.7	19.0	0.7	19.5	26.0	39.5							300	do.....	330
5.0	10.0	15.3	21.2	1.5	22.0	29.0	41.0							304	do.....	330
9.0	18.6	31.0												181	Specimen cross-grained; split with the grain.....	1113
6.1	12.0	18.0	25.0	1.2	25.7	33.0								268	do.....	476
5.7	10.7	15.9	21.2	1.0	22.0	27.6	34.0	41.2	49.0					404	do.....	1123
7.5	14.0	21.6	31.0	3.0	32.3	43.0								268	Long fracture with one large splinter.....	735
7.3	14.2	22.5	31.5	2.5	33.0	43.5	70.5							302	Long fracture with two large splinters.....	735
5.7	11.0	17.5	25.5	2.3	27.0	36.0	53.0							323	do.....	1093
7.4	15.5	23.8	35.5											200	Specimen cross-grained; square break.....	1093
6.0	12.0	18.0	24.0	1.3	24.5	31.0	39.0							350	Broke in three pieces.....	1140
6.0	10.6	15.5	21.2	1.0	21.7	28.0								293	Square break on tension side with large flake on compression side; broke at knot.....	1140
5.5	11.0	18.0	26.5	4.0	28.0									250	Square break with large splinters.....	437
4.6	10.0	15.5	21.7	2.0	23.5									232	Square break, the ends splitting.....	437
11.7														78	Square break.....	463
.....														48	do.....	463
4.7	9.0	13.3	18.0	0.3	18.3	22.7	23.0	33.2	38.3	46.3	53.0			531	Broke with long splinters.....	475
5.0	9.0	13.6	18.0	0.3	18.3	23.0	27.7	32.0	33.0	44.0	51.0	61.0		533	Broke with long splinters on compression side.....	475
5.0	10.0	14.6	20.4	1.0	22.0	27.0	32.0	39.0						332	Broke suddenly; stick shattered.....	452
4.4	8.5	13.2	18.4	0.9	18.6	24.1	30.0	35.5	41.4	50.5				474	do.....	452
7.5	15.7	25.0	40.0	5.0	42.0	53.0	64.5							300	Specimen cross-grained; broke with long split.....	230
6.9	14.7	24.0	37.7	5.3	39.0	57.1								236	Square break.....	230
8.0	16.0	26.0	39.2	5.5	41.0									250	Specimen cross-grained; broke with long split.....	434
7.0	14.5	23.5	34.3	5.0	36.3									238	Square break on tension side with large flake on compression side.....	434
10.5	20.0													134	Specimen cross-grained; broke through small knots.....	341
7.0	13.0	20.0	23.8	2.3	30.0									245	Broke at knot.....	333
6.0	12.0	19.8	27.9	2.5	28.8									204	Specimen cross-grained; oblique fracture.....	333

TABLE III.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Specific gravity of the air-dried specimen.	Direction of grain.	COEFFICIENT OF ELASTICITY.		Modulus of rupture.
								First deflection.	Second deflection.	
RHAMNACEÆ.										
42. <i>Reynoldsia latifolia</i> . <i>Red Iron Wood. Darling Plum.</i>	454	Florida	Upper Metacombe Key.	A. H. Curtiss	Coral	1.2012		976	1050	820
43. <i>Condalia ferrea</i> . <i>Black Iron Wood.</i>	460	do	do	do	do	1.2546		1191	1109	1127
	460	do	do	do	do	1.2430		1191	1176	680
45. <i>Rhamnus Caroliniana</i> . <i>Indian Cherry.</i>	803	do	Saint John's river	do	Rich hummock	0.5309		814	794	518
	1094	Arkansas	Jonesboro'	T. B. Kitchens	do	0.5066		718	697	616
47. <i>Rhamnus Purshiana</i> . <i>Bearberry. Bear Wood. Shittim Wood.</i>	993	Oregon	Portland	G. Engelmann and C. S. Sargent.	Rich, alluvial	0.5943		775	912	750
49. <i>Colubrina reclinata</i> . <i>Naked Wood.</i>	1139	Florida	Umbrella Key	A. H. Curtiss	Coral	0.8721		921	976	1216
SAPINDACEÆ.										
50. <i>Æsculus glabra</i> . <i>Ohio Buckeye. Fetid Buckeye.</i>	297	Missouri	Allenton	G. W. Letterman	Rich, moist	0.4653		761	707	496
	297	do	do	do	do	0.4602		687	651	468
	427	Tennessee	Nashville	A. Gattinger	Rich, moist	0.4970		610	574	549
52. <i>Æsculus Californica</i> . <i>California Buckeye.</i>	684	California	Marin county	G. R. Vasey	Rich upland	0.5034		697	669	548
	684	do	do	do	do	0.5228		678	696	649
54. <i>Sapindus marginatus</i> . <i>Wild China. Soapberry.</i>	307	Texas	Dallas	J. Reverchon	Rich, damp	0.7681		888	888	663
	307	do	do	do	do	0.7784		857	948	949
	928	do	Austin	C. Mohr	Limestone	0.5243		842	814	820
	928	do	do	do	do	0.5310		688	697	703
56. <i>Hypelate paniculata</i> . <i>Ink Wood. Iron Wood.</i>	463	Florida	Upper Metacombe Key.	A. H. Curtiss	Coral	1.0405		1135	1206	1261
	463	do	do	do	do	1.0123		939	1017	1118
60. <i>Acer macrophyllum</i> . <i>Broad-leaved Maple.</i>	982	Oregon	Portland	G. Engelmann and C. S. Sargent.	Rich, alluvial	0.5445		697	734	696
	982	do	do	do	do	0.5341		751	751	696
	1023	do	Portland Furniture Company.	do	do	0.4907		697	842	691
	1023	do	do	do	do	0.5053		729	794	656
61. <i>Acer circinatum</i> . <i>Vine Maple.</i>	1013	do	Portland	do	Moist, alluvial	0.6928		626	713	712
	1014	Washington territory.	Wilkeson	do	do	0.7001		634	723	818
64. <i>Acer saccharinum</i> . <i>Sugar Maple. Sugar Tree. Hard Maple. Rock Maple.</i>	299	Missouri	Allenton	G. W. Letterman	Rich upland	0.6381		1526	1457	1219
	376	Vermont	Charlotte	C. G. Pringle	Gravelly	0.6852		888	976	816
	409	New England	Charlestown Navy-yard.	S. H. Pook	do	0.6775		763	930	675
	1233	Vermont	Charlotte	F. H. Horsford	do	0.7447		1683	1953	1404
	1233	do	do	do	do	0.6986		1878	1808	1289
	1234	do	do	do	do	0.7063		1220	1395	1235
	1234	do	do	do	do	0.7108		1395	1480	1284
	1235	do	do	do	do	0.7137		1628	1575	1242
	1235	do	do	do	do	0.7189		1436	1575	1104
64. <i>Acer saccharinum, var. nigrum</i> . <i>Black Sugar Maple.</i>	213	do	do	C. G. Pringle	Clay	0.7241		1039	1149	1055
	2741	Missouri	Allenton	G. W. Letterman	Low, alluvial	0.7355		1085	1221	1055
	440	Tennessee	Nashville	A. Gattinger	Rich	0.6973		888	1149	1024
	757	Florida	Chattahoochee	A. H. Curtiss	Clay	0.6979		257	930	933
	757	do	do	do	do	0.6824		610	688	743
65. <i>Acer dasycarpum</i> . <i>Soft Maple. White Maple. Silver Maple.</i>	1052	Massachusetts	Topsfield	J. Robinson	Low meadow	0.6641		976	1109	1019
65. <i>Acer rubrum</i> . <i>Red Maple. Swamp Maple. Soft Maple. Water Maple.</i>	20	do	Arnold Arboretum	C. S. Sargent.	Drift	0.7148		827	864	738
	530	Mississippi	Kemper's mill	C. Mohr	Rich, swampy	0.6136		871	1039	820

UNITED STATES UNDER TRANSVERSE STRAIN—Continued.

DEFLECTION, IN MILLIMETERS, UNDER A PRESSURE, IN KILOGRAMS, OF—														Ultimate strength: transverse pressure.	Remarks.	Office number.
50	100	150	200	0 (set.)	200	250	300	350	400	450	500	550				
5.0	9.3	13.5	18.0	0.5	19.0	23.0	30.0							350	Specimen cross-grained.....	454
4.1	8.8	11.5	15.5	0.5	16.0	20.0	24.5	30.0	33.7	40.0				481	Long, shattered break.....	460
4.1	8.3	12.1	16.0	0.5	17.0	21.2								290	Specimen cross-grained.....	460
6.0	12.3	18.0	26.5	2.4	27.5									221	Long break, starting at small knot.....	808
6.8	14.2	22.1	33.0	2.5	34.0									263	Shattered.....	1094
6.3	10.7	16.5	22.0	1.0	23.0	28.0	36.6							320	Long, shattered break with large splinters.....	968
5.3	16.0	15.0	19.6	0.5	19.5	24.5	29.0	34.5	48.0	57.0				519	Specimen cross-grained; shattered.....	1139
6.5	13.8	22.4												199	Crushed at center bearing.....	297
7.1	15.0	24.0	41.5											200	do.....	297
8.0	17.0	27.6	44.0	7.5	48.6									234	Crushed at center bearing; broke at knot on tension side.....	427
7.0	14.6	23.0	31.5	1.8	32.0	44.0								265	Long fracture; large splinters.....	684
7.2	14.0	21.7	30.3	2.3	31.8	44.2								277	Long fracture.....	684
5.5	11.0	16.6	23.5	1.5	24.5	32.0	42.0	50.0						383	Long, splintered fracture.....	307
5.7	10.3	15.5	22.4	1.5	23.4	30.7	42.0	58.0	90.0					405	Splintered fracture.....	307
5.8	12.0	19.0	29.0	3.0	29.7	37.0	53.0							350	do.....	928
7.1	14.0	22.0	32.0	4.2	34.2	45.0								300	Long, splintered fracture.....	928
4.3	8.1	12.0	16.5	0.7	17.0	21.5	27.0	33.2	39.5	48.5	62.0			538	Shattered.....	463
5.2	9.6	14.5	19.0	1.0	19.0	24.0	30.5	36.5	44.5	56.0				477	do.....	463
7.0	13.3	21.0	30.5	3.4	32.0	43.0								297	Short break, splitting in axis of stick.....	982
6.5	12.0	19.8	29.5	3.0	31.0	42.0								297	Slightly crushed at center bearing.....	982
7.0	11.6	17.8	25.0	2.0	28.0	35.0								295	Slightly crushed at center bearing; splintered.....	1023
6.7	12.3	19.4	29.0	2.5	30.0	41.0								280	Short break with long, large splinter.....	1023
7.8	13.7	21.5	30.5	2.0	31.4	40.3								304	Long, shattered break.....	1013
7.7	13.5	21.0	29.5	2.0	30.0	39.0	50.5							349	do.....	1014
3.2	6.7	10.2	13.5	0.3	13.8	17.0	21.5	25.8	32.0	39.0	50.5			551	Long splinter on angle of two faces.....	299
5.5	10.9	15.0	20.0	0.7	20.3	26.0	35.0							348	Splintered.....	376
6.4	10.5	15.5	20.5	0.8	21.3	27.0								288	Specimen cross-grained; split with grain.....	409
2.9	5.0	7.7	10.2	0.0	10.2	12.6	15.2	18.2	21.7	25.4	31.4	38.3		599	Broke with fine splinters on back.....	1233
2.6	5.4	8.4	11.2	0.0	11.2	14.0	17.0	20.2	25.3	30.0	39.0	50.0		550	do.....	1233
4.0	7.0	10.2	13.6	0.2	13.6	16.6	20.2	24.6	30.0	36.0	47.0			527	Square break on tension side, splitting in axis.....	1234
3.5	6.6	10.0	13.0	0.2	13.0	16.0	19.5	23.4	27.7	33.0	42.0			548	do.....	1234
3.0	6.2	9.6	13.0	0.0	13.0	15.8	19.7	24.7	30.0	36.8	50.0			530	do.....	1235
3.4	6.2	9.1	12.3	0.0	12.5	15.4	19.0	22.6	27.6	34.5				471	Specimen cross-grained; shattered on angle of two faces.....	1235
4.7	8.5	13.5	18.2	0.5	18.0	23.5	30.6	38.0	49.5	65.0				450	Broke with long splinters.....	213
4.5	8.0	12.0	15.5	0.5	16.3	20.0	26.0	32.0	43.5					450	Crushed slightly at center bearing; broke with fine splinters.....	274
5.5	8.5	13.0	17.6	0.6	18.0	22.5	29.5	37.0	52.0					437	Broke with fine splinters.....	440
5.7	10.5	16.5	23.0	1.5	24.0	30.7	41.5	53.0						398	Splintered on angle, starting at small knot.....	757
8.0	14.2	21.0	30.0	2.0	30.7	40.4	54.0							317	Broke with long, fine splinters.....	757
5.0	8.8	14.0	19.0	1.3	19.2	25.0	32.0	41.0	55.0					435	Broke with long, fine splinters.....	1052
5.9	11.3	17.2	24.5	1.6	26.0	35.0	46.5							315	Specimen sap-wood, cross-grained; broke at knot.....	20
5.6	9.4	14.2	19.5	1.2	20.0	28.0	39.5	58.0						350	Crushed at center bearing; broke with fine splinters.....	530

TABLE III.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Specific gravity of the air-dried specimen.	Direction of grain.	COEFFICIENT OF ELASTICITY.		Modulus of rupture.
								First deflection.	Second deflection.	
66. <i>Acer rubrum</i> —continued.	530	Mississippi	Kemper's mill	C. Mohr	Rich, swampy	0.5999		976	1028	800
	743	Georgia	Bainbridge	A. H. Curtiss	Low	0.5524		888	930	600
	743	do	do	do	do	0.5509		976	1085	797
	878	Massachusetts	Danvers	J. Robinson	do	0.6618		888	888	937
	878	do	do	do	do	0.7102		775	835	830
	1048	do	North Reading	do	do	0.6699		888	904	820
	1048	do	do	do	do	0.6710		842	913	937
67. <i>Negundo aceroides</i> <i>Box Elder. Ash-leaved Maple.</i>	290	Missouri	Allenton	G. W. Letterman	Rich bottom	0.4750		610	618	443
	290	do	do	do	do	0.4585		626	697	572
	311	Texas	Dallas	J. Reverchon	do	0.4773		452	478	500
	311	do	do	do	do	0.4614		514	536	591
68. <i>Negundo Californicum</i> <i>Box Elder.</i>	645	California	Contra Costa county	G. R. Vasey	Rich, moist	0.5227		939	986	785
	645	do	do	do	do	0.5227		921	904	800
ANACARDIACEÆ.										
71. <i>Rhus copallina</i> <i>Dwarf Sumach.</i>	738	Florida	Chattahoochee	A. H. Curtiss	Dry clay	0.4888		634	638	604
	738	do	do	do	do	0.5054		763	835	633
71. <i>Rhus Metopium</i> <i>Poison Wood. Coral Sumach. Mountain Manchineel. Bum Wood. Hog Plum. Doctor Gum.</i>	467	Florida	Upper Metacombe Key.	A. H. Curtiss	Coral	0.7967		921	1050	616
	467	do	do	do	do	0.8105		921	1050	694
LEGUMINOSÆ.										
77. <i>Robinia Pseudacacia</i> <i>Locust. Black Locust. Yellow Locust.</i>	405		Charlestown Navy-yard.	S. H. Pook				1396	1302	1481
	405		do	do		0.8205		857	1302	1862
	405		do	do		0.8148		1221	1268	1514
	815	West Virginia	Grafton	C. G. Pringle		0.6433		1039	1050	1041
	815	do	do	do		0.6433		814	976	851
	1247	New York	Long Island	M. C. Beedle		0.7956		1526	1627	1387
	1247	do	do	do		0.7769		1627	1627	1355
79. <i>Robinia Neo-Mexicana</i> <i>Locust.</i>	1248		do	do		0.8069		976	1123	1069
	1248		do	do		0.8456		1320	1436	1397
	1081	Colorado	Trinidad	W. B. Strong	Low, moist	0.8019		1061	1149	900
80. <i>Olneya Tesota</i> <i>Iron Wood. Arbol de Hierro.</i>	650	California	Lower Colorado valley.	G. Engelmann and C. S. Sargent.	Dry, gravelly	0.9841		976	1039	942
	650	do	do	do	do	1.1966		650	697	558
81. <i>Piscidia Erythrina</i> <i>Jamaica Dogwood.</i>	564	Florida	Upper Metacombe Key.	A. H. Curtiss	Coral	1.0398		740	814	820
	564	do	do	do	do	0.9406		812	888	675
82. <i>Cladrastis tinctoria</i> <i>Yellow Wood. Yellow Ash. Gopher Wood.</i>	38	Kentucky	Mercer county	W. M. Linney	Limestone	0.7852		888	976	969
	38	do	do	do	do	0.6444		937	1028	813
84. <i>Sophora affinis</i>	329	Texas	Dallas	J. Reverchon	Dry, calcareous	0.8697		842	857	820
	982	do	Austin	C. Mohr	do	0.9594		1150	1097	790
85. <i>Gymnocladus Canadensis</i> <i>Kentucky Coffee Tree. Coffee Nut.</i>	519	Tennessee	Nashville	A. Gatteringer	Limestone	0.7143		872	849	703
	1241	Missouri	Allenton	G. W. Letterman	Alluvial	0.6966		1061	1110	813
	1242	do	do	do	do	0.6875		1221	1149	769
	1243	do	do	do	do	0.6670		976	1085	802
86. <i>Gleditsia triacanthos</i> <i>Honey Locust. Black Locust. Three-thorned Acacia. Sweet Locust. Honey Shucks.</i>	531	do	do	do	Low, rich	0.6250		976	1149	968
	532	do	do	do	do	0.6381		1164	1221	907
	444	Tennessee	Nashville	A. Gatteringer	Dry, sandy barrens	0.6969		888	888	893

UNITED STATES UNDER TRANSVERSE STRAIN—Continued.

DEFLECTION, IN MILLIMETERS, UNDER A PRESSURE, IN KILOGRAMS, OF—													Ultimate strength: transverse pressure.	Remarks.	Office number.
50	100	150	200	0 (set.)	200	250	300	350	400	450	500	550			
5.0	9.5	14.5	19.0	0.5	19.2	25.0	35.5						345	Broke with long splinters	530
5.5	10.5	16.5	23.5	2.0	24.5	32.0							256	Specimen 0.75 sap-wood; cross-grained	743
5.0	9.0	14.5	20.0	0.7	20.5	27.0	39.0						340	Sap-wood; crushed at center bearing; broke with fine splinters	743
5.5	11.0	17.0	24.0	1.2	25.0	33.0	43.0	61.0					400	do	878
6.3	11.7	18.3	28.0	3.0	28.4	37.5	52.5	90.0					358	Sap-wood; broke with long, large splinters	878
5.5	10.8	16.0	22.0	1.1	23.0	31.5	42.0						350	do	1048
5.8	10.7	15.5	21.5	1.0	21.8	28.0	37.0	50.0					400	do	1048
8.0	15.8	26.3											189	Broke at knot near center bearing	290
7.8	14.0	22.0	33.5	3.8	35.0								244	Crushed at center bearing; square break	290
10.8	20.6	31.0	44.5	5.0	47.0								217	Square break at knot	311
9.5	18.2	30.0	42.0	3.5	44.0	71.0							232	Flaked on tension side; broke at small knot	311
5.2	9.9	14.2	21.2	1.0	21.2	28.0	37.5						335	Crushed at center bearing; broke into long splinters	645
5.3	10.8	15.8	21.0	0.5	22.0	28.0	36.5						345	Crushed at center bearing; broke into long, fine splinters	645
7.7	15.3	23.2	32.3	2.6	33.5	43.5							296	Shattered, splitting to the end	736
6.4	11.7	18.0	27.0	3.1	28.0	37.3							270	do	736
5.3	9.3	14.5	21.3	1.6	23.0	30.5							263	Square break	467
5.3	9.3	14.5	21.3	1.5	23.0	30.5							296	do	467
3.5	8.0	11.0	14.5	0.8	14.7	18.0	22.0	25.8	30.2	35.6	41.5	47.8	632	60.2 millimeters deflection with 600 kilograms; specimen cross-grained	405
5.7	7.5	12.0	15.5	0.3	15.3	19.0	23.5	28.7	35.0	40.0	47.0	55.0	581	Broke with fine splinters	405
4.0	7.7	11.5	15.6	0.2	15.3	19.0	23.9	28.0	32.0	37.0	44.5	51.8	646	64.3 millimeters deflection with 600 kilograms; broke with fine splinters	405
4.7	9.3	13.8	19.1	0.3	19.0	23.7	30.0	33.0	47.7				444	Broke with coarse splinters	315
6.0	10.0	14.5	19.0	0.5	19.5	24.0	30.0	42.0					363	do	315
3.2	6.0	9.0	12.0	0.0	12.0	15.0	18.0	21.0	24.9	29.3	34.7	42.5	592	Splintered on corners	1247
3.0	6.0	9.0	11.6	0.2	11.8	14.5	17.8	21.2	25.4	31.2	36.0	46.5	578	Broke with fine splinters	1247
5.0	8.7	12.5	16.6	0.4	16.8	20.8	25.5	31.0	36.5	45.6			456	Broke with large splinters on corners	1248
3.7	6.8	10.0	13.2	0.5	13.5	16.4	20.1	24.3	28.0	33.5	39.0	46.4	596	do	1248
4.6	8.5	12.4	16.2	0.4	16.7	20.6	25.0	31.0					338	Broke with long splinters	1031
5.0	9.4	14.2	19.5	0.5	20.0	26.5	33.7	42.0	54.3				402	Cross-grained; split with grain	650
7.4	14.0	21.0	29.9	1.4	31.0								238	Square break	650
6.6	12.0	19.5	26.0	1.8	27.1	33.5	43.0	51.0					354	Broke with long, large splinters	564
5.8	11.0	17.3	24.0	1.4	24.5	30.5							288	do	564
5.5	10.0	15.8	21.6	1.1	22.5	29.0	37.0	47.0	68.0				422	Long, large splinters on angle of two faces	33
5.1	9.5	14.5	20.2	1.0	20.5	28.0							347	Started at small knot; long, oblique fracture	33
5.8	11.4	17.0	23.0	1.5	25.0	33.0	42.0						350	Broke at large knot with long splinters	320
4.4	8.9	13.0	18.2	1.2	18.7	24.0	31.0						341	Long, oblique break	932
5.6	11.5	18.5	23.9	3.0	29.0	37.3	52.0						390	Square break on tension side, splitting in axis of the stick	519
4.6	8.8	13.0	18.4	0.9	18.8	26.0	37.5						347	Splintered on angle of two faces	1241
4.0	8.5	14.0	19.2	1.5	20.0	28.7	46.5						328	Broke with fine splinter	1242
5.0	9.0	13.7	20.0	1.4	20.5	28.3	44.0						342	do	1243
5.0	8.5	12.6	16.8	0.6	18.0	24.0	31.0	41.0	57.0				413	Crushed at center bearing; broke with fine splinter	531
4.2	8.0	12.9	17.6	1.0	18.7	25.0	33.0	47.0					387	Broke with fine splinter	532
5.5	11.0	16.4	22.5	1.0	23.5	31.7	39.5	55.5					381	Broke with long splinter	444

TABLE III.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species	Office number.	State.	Locality.	Collector.	Soil.	Specific gravity of the air-dried specimen.	Direction of grain.	COEFFICIENT OF ELASTICITY.		Modulus of rupture.
								First deflection.	Second deflection.	
87. <i>Gleditsia monosperma</i> <i>Water Locust.</i>	760	Florida	Chattahoochee.	A. H. Curtis	Alluvial	0.7628		1061	1149	1073
	760	do	do	do	do	0.7748		1221	1191	982
88. <i>Parkinsonia Torreyana</i> <i>Green-bark Acacia. Palo Verde.</i>	678	Arizona	Lower Colorado river.	G. Engelmann and C. S. Sargent.	Sandy	0.6705		514	514	511
	678	do	do	do	do	0.6727		568	602	579
91. <i>Cercis Canadensis</i> <i>Redbud. Judas Tree.</i>	436	Tennessee	Nashville	A. Gittinger	Limestone	0.7323		814	904	1008
	1089	Missouri	Allenton	G. W. Letterman	Rich	0.6341		588	610	734
	1090	do	do	do	do	0.6534		452	514	649
	1091	do	do	do	do	0.6250		651	723	516
93. <i>Prosopis juliflora</i> <i>Mesquit. Algaroba. Honey Locust. Honey Pod.</i>	680	Arizona	Tucson	C. S. Sargent		0.7818		542	618	469
	680	do	do	do		0.7614		506	630	574
	927	Texas	Austin	C. Mohr	Rich, calcareous	0.7750		509	501	412
94. <i>Prosopis pubescens</i> <i>Screw Bean. Screw-pod Mesquit. Tornilla.</i>	658	California	Fort Yuma	G. Engelmann and C. S. Sargent.	Sandy	0.8068		814	835	902
	658	do	do	do		0.8568		787	814	886
95. <i>Acacia Greggii</i> <i>Out's Claw.</i>	697	Arizona	Santa Rita mountain.	do	Dry, gravelly	0.8714		1639	1085	792
100. <i>Lysiloma latialiqua</i> <i>Wild Tamarind.</i>	509	Florida	Boca Chica Key	A. H. Curtis	Coral	0.5870		488	461	553
ROSACEÆ.										
102. <i>Chrysobalanus Icaco</i> <i>Cocoa Plum.</i>	480	do	Bay Biscayne	do	Swampy	0.7753		957	1110	961
103. <i>Prunus Americana</i> <i>Wild Plum. Canada Plum. Horse Plum.</i>	68	Missouri	Allenton	G. W. Letterman	Rich upland	0.6003		814	769	640
	68	do	do	do	do	0.5916		651	734	703
	234	Texas	Dallas	J. Reverchon	Rich	0.8045		921	976	1240
104. <i>Prunus angustifolia</i> <i>Chickasaw Plum. Hog Plum.</i>	435	Tennessee	Nashville	A. Gittinger	River bluff	0.6538		634	603	469
107. <i>Prunus emarginata</i> , var. <i>mollis</i>	968	Washington territory.	Wilkeson	G. Engelmann and C. S. Sargent.	Low, rich	0.4699		751	849	687
	968	do	do	do	do	0.4750		751	872	670
108. <i>Prunus serotina</i> <i>Wild Black Cherry. Rum Cherry.</i>	15	Massachusetts	Roxbury	C. S. Sargent	Gravelly	0.7438		775	849	1171
	15	do	do	do	do	0.7386		1017	888	1084
	115 ¹	Michigan	Lansing	W. J. Beal	do	0.5315		651	673	579
	115 ²	do	Dansville	do	do	0.5048		814	849	724
	127	Missouri	Allenton	G. W. Letterman	Rich loam	0.6790		976	976	905
	127	do	do	do	do	0.6670		1221	1028	996
	148	Illinois	Waukegan	It. Douglas	Gravelly	0.6471		976	930	937
	317	Michigan	Hersey	W. J. Beal	Rich	0.5315		775	800	689
	317	do	do	do	do	0.5493		842	828	691
	368	Vermont	Charlotte	C. G. Pringle	Gravelly	0.5648		764	857	769
	406	Virginia or Middle States.	Charlestown Navy-yard.	S. H. Pook		0.5028		651	718	642
	768	Florida	Chattahoochee	A. H. Curtis	Clay	0.6105		976	958	797
	768	do	do	do	do	0.6244		888	976	792
	1053	Massachusetts	Topsfield	J. Robinson	Gravelly	0.6751		740	769	820
	1053	do	do	do	do	0.6716		698	775	820
110. <i>Prunus demissa</i> <i>Wild Cherry.</i>	637	California	Strawberry valley	G. Engelmann and C. S. Sargent.	Low, rich	0.7636		814	769	691
111. <i>Prunus Caroliniana</i> <i>Wild Orange. Mock Orange. Wild Peach.</i>	1082	Florida	Jacksonville	A. H. Curtis	Sandy	0.8785		697	718	586
	1062	Texas	Victoria	C. Mohr	Rich, moist	0.8698		1191	1097	1266
	1062	do	do	do	do	0.8481		976	996	980

UNITED STATES UNDER TRANSVERSE STRAIN—Continued.

DEFLECTION, IN MILLIMETERS, UNDER A PRESSURE, IN KILOGRAMS, OF—													Ultimate strength: transverse pressure.	Remarks.	Office number.
50	100	150	200	0 (set.)	200	250	300	350	400	450	500	550			
4.6	8.5	13.3	18.2	0.6	18.0	23.5	29.0		42.0				458	0.66 sap-wood; broke with long, fine splinters	769
4.0	8.2	12.8	18.5	1.0	19.5	25.5	33.0	40.5	52.5				419	Specimen cross-grained; split with grain	760
9.5	19.0	28.4	43.2	6.4	46.0								218	Specimen cross-grained; broke at knot	672
8.6	16.2	25.4	35.2	3.0	37.7								247	Long, oblique fracture	673
6.0	10.8	15.8	22.9	1.6	22.7	29.7	38.0	50.0	68.0				428	Broke with long, large splinters	436
8.3	16.0	24.5	35.0	2.5	37.0	48.0	74.5						313	 do	1089
10.8	19.0	29.7	43.0	6.0	45.5	61.0							277	 do	1090
7.5	13.5	20.0	28.4	1.5	30.0								220	Broke at knot near support	1091
9.0	15.8	23.0											200	Specimen cross-grained; short, oblique fracture	690
8.2	15.5	22.5	30.7	1.6	31.7								245	 do	690
9.6	19.5	31.5											176	Specimen cross-grained; shattered	927
6.0	11.7	17.7	24.3	1.3	25.0	32.0	41.0						385	Specimen cross-grained; short, oblique fracture	658
6.2	12.0	17.3	24.5	2.2	25.5	33.5	42.0	53.0					378	 do	658
4.7	9.0	14.0	18.1	0.6	18.6	24.0	29.0						338	Broke at knot near the end	697
10.0	21.2	32.7	51.3	8.0	55.3								236	Specimen cross-grained	509
5.1	8.8	13.0	17.8	0.9	18.0	23.0	31.5	40.0	66.0				410	Broke with long splinters, starting at knot	489
6.0	12.7	19.3	27.4	1.7	28.0	37.0							277	Broke with long splinters	68
7.5	13.3	21.0	28.9	1.6	29.0	36.5							300	Broke with long, large splinters	68
5.3	10.0	14.9	20.7	0.5	20.6	27.0	34.0	41.0	50.0	61.5	84.0		529	Broke with fine splinters	334
7.7	16.2	26.0	38.5	4.5									200	Specimen cross-grained, defective; square break on tension side ..	435
6.5	11.5	17.7	25.2	1.5	26.0	36.0							293	Broke with long, coarse splinters	968
6.5	11.2	17.0	24.7	1.7	26.0	36.5							296	Shattered	968
6.3	11.5	17.3	23.6	1.0	24.2	30.0	38.5	46.5	57.5	73.0			500	 do	15
4.8	11.0	15.9	22.0	0.6	22.0	28.0	35.7	44.0	55.0				467	Broke with fine splinters	16
7.5	14.5	21.7	32.0	1.6	32.5								247	Long, oblique fracture	115
6.0	11.5	17.6	24.6	1.0	25.0	32.5	45.0						300	 do	115
5.0	10.0	15.2	21.5	1.3	22.0	28.5	36.5	47.0					386	Broke with long splinters	127
4.0	9.5	15.0	21.0	1.2	22.0	28.0	36.0	45.0	57.5				425	Shattered; long splinters	127
5.0	10.5	16.0	21.5	1.0	22.5	28.0	36.5	45.5					400	Broke with fine splinters	148
6.3	12.2	19.0	26.8	1.1	27.5	36.0							294	Shattered; long splinters on corner	317
5.8	11.8	16.7	23.0	0.6	24.0	30.0							295	Broke with long splinters on corner	317
6.4	11.4	16.7	22.6	1.0	23.0	30.0	40.5						328	Broke with coarse splinters	368
7.5	13.6	21.5	30.7	1.7	31.6	43.5							274	Specimen cross-grained; single fracture	406
5.0	10.2	15.2	21.0	0.6	21.8	26.5	34.0						340	Broke with fine splinters	763
5.5	10.0	15.5	20.8	1.0	21.5	27.0	35.0						338	Broke with long, large splinters	768
6.6	12.7	19.5	27.0	1.5	28.0	36.0	46.0	62.5					350	Specimen cross-grained; long fracture	1053
7.0	12.6	19.5	26.5	1.5	27.0	34.0	46.0	59.5					350	 do	1053
6.0	12.7	19.5	28.4	1.6	28.3	36.5							295	Specimen cross-grained; split with the grain	637
7.0	13.6	22.0	32.3	4.0	34.0	73.0							250	Specimen cross-grained	1032
4.1	8.9	13.5	18.4	0.2	19.0	23.5	29.2	35.7	41.7	53.0	74.0		540	 do	1062
5.0	9.8	14.0	18.5	0.6	19.0	24.5	32.0	37.5					397	Specimen cross-grained; shattered	1062

TABLE III.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Specific gravity of the air-dried specimen.	Direction of grain.	COEFFICIENT OF ELASTICITY.		Modulus of rupture.
								First deflection.	Second deflection.	
113. <i>Prunus ilicifolia</i> <i>Islay.</i>	1158	California	Santa Cruz	C. L. Anderson		0.9734		751	751	820
	1158	do	do	do		0.9678		697	712	745
117. <i>Pyrus coronaria</i> <i>American Crab. Sweet-scented Crab.</i>	1087	Pennsylvania	Nazareth	J. Henry	Moist	0.7857		751	751	441
	1087	do	do	do	do	0.7228		729	723	423
	1088	do	do	do	do	0.7340		508	465	452
	1088	do	do	do	do	0.7382		651	630	623
121. <i>Pyrus sambucifolia</i> <i>Mountain Ash.</i>	410	Vermont	Mount Mansfield	C. G. Pringle	Gravelly	0.5727		575	626	445
125. <i>Crataegus arborescens</i>	607	Georgia	Ogeechee	A. H. Curtiss	Low	0.6818		814	814	799
	607	do	do	do	do	0.7126		788	763	443
126. <i>Crataegus Crus-galli</i> <i>Cockspur Thorn. Newcastle Thorn.</i>	328	Massachusetts	Brookline	J. Robinson	Loam	0.6946		543	575	619
	328	do	do	do	do	0.7166		508	514	566
	1093	Missouri	Allenton	G. W. Letterman	Low, wet	0.7767		751	775	763
	1093	do	do	do	do	0.7946		751	794	766
128. <i>Crataegus subvillosa</i> <i>Scarlet Haw.</i>	949	Texas	Victoria	C. Mohr	Alluvial	0.7565		921	913	860
	1081	Missouri	Saint Louis	H. Eggert		0.8670		842	888	616
129. <i>Crataegus tomentosa</i> <i>Black Thorn. Pear Haw.</i>	426	Tennessee	Nashville	A. Gittinger	Limestone	0.7166		740	740	696
	426	do	do	do	do	0.7527		651	723	720
132. <i>Crataegus spathulata</i> <i>Small-fruited Haw.</i>	926	Louisiana	Webster parish	C. Mohr	Clay	0.7102		718	674	506
134. <i>Crataegus astivalis</i> <i>May Haw. Apple Haw.</i>	239	South Carolina	Bonneau's Depot	H. W. Ravenel	Damp, rich	0.7239		595	592	712
135. <i>Crataegus flava</i> , var. <i>pubescens</i> <i>Summer Haw. Red Haw.</i>	767	Florida	Aspalaga	A. H. Curtiss	Dry clay	0.7973		740	708	724
137. <i>Amelanchier Canadensis</i> <i>Juneberry. Shad Bush. Service Tree. May Cherry.</i>	241	Kentucky	Brumfield Station	W. M. Linney	Waverly shale	0.8312		1191	1221	1256
	849	Massachusetts	Danvers	J. Robinson	Loam	0.8472		1085	1149	1065
	849	do	do	do	do	0.8410		1163	1221	1054
HAMAMELACEÆ.										
139. <i>Liquidambar styraciflua</i> <i>Sweet Gum. Star-leaved Gum. Liquidambar. Red Gum. Bilsted.</i>	546	Alabama	Kemper's mill	C. Mohr	Rich, alluvial	0.5448		603	610	520
	546	do	do	do	do	0.5796		688	674	663
	1095	Arkansas	Little Rock	G. W. Letterman		0.6012		888	930	776
	1095	do	do	do	do	0.5765		976	996	703
	1173	New Jersey	Mount Holly	S. P. Sharples	Clay	0.6080		751	769	750
	1173	do	do	do	do	0.6477		697	496	788
	1181	Mississippi	Yazoo River bottom	R. Abbey	Alluvial	0.5864		751	781	553
	1181	do	do	do	do	0.6001		787	849	544
	1182	do	do	do	do	0.6250		976	1017	670
	1182	do	do	do	do	0.6375		1163	1061	614
	1183	do	do	do	do	0.5409		814	888	661
	1183	do	do	do	do	0.6159		1017	976	616
RHIZOPHORACEÆ.										
140. <i>Rhizophora Mangle</i> <i>Mangrove.</i>	485	Florida	Bay Biscayne	A. H. Curtiss	Salt-marsh	1.1480		1627	1627	1308
	485	do	do	do	do	1.1835		1627	1684	1106
COMBRETACEÆ.										
141. <i>Conocarpus erecta</i> <i>Button Wood.</i>	489	do	do	do	do	1.0240		814	913	830
	489	do	do	do	do	1.0292		1062	1136	1055
142. <i>Laguncularia racemosa</i> <i>White Button Wood. White Mangrove.</i>	507	do	Sugar-Loaf Sound	do	do	0.7384		698	634	272
	507	do	do	do	do	0.7230		775	814	764

UNITED STATES UNDER TRANSVERSE STRAIN—Continued.

DEFLECTION, IN MILLIMETERS, UNDER A PRESSURE, IN KILOGRAMS, OF—													Ultimate strength: transverse pressure.	Remarks.	Office number.
50	100	150	200	0 (set.)	200	250	300	350	400	450	500	550			
6.5	12.0	20.0	29.0	2.0	30.0	38.0	56.0	90.0					350	Broke with large splinters.....	1158
7.0	13.7	22.3	32.5	4.5	35.0	48.0	67.0						318	Broke with long splinters.....	1158
6.5	13.0	21.5											183	Broke at knot.....	1087
6.7	13.5	21.3											180	Specimen cross-grained.....	1087
9.6	21.0	33.5											193	Specimen cross-grained; defective.....	1088
7.5	15.5	24.6	36.5	4.8	37.0	51.5							266	Specimen cross-grained.....	1088
8.5	15.6	25.5											190	Broke at knots.....	410
6.0	12.0	18.3	26.5	1.5	37.1	35.0	47.0						341	Specimen cross-grained; broke with one long splinter.....	607
6.2	12.8	20.5											189	Broke at knot.....	607
9.0	17.0	28.0	39.0	4.9	41.0	58.0							264	Broke with long splinters.....	323
9.6	19.0	28.5	42.3	5.0	44.0								250	Specimen cross-grained; split with grain.....	323
6.5	12.6	20.0	29.0	2.1	29.0	38.0							300	Square break.....	1093
6.5	12.3	18.1	26.0	2.0	26.5	34.2	46.5						302	Broke at knots.....	1093
5.3	10.7	16.0	22.0	1.0	23.0	29.0	37.0	49.5					367	Specimen cross-grained; split with grain.....	949
5.8	11.0	16.6	25.5	2.1	26.0	34.0							263	Broke at knot.....	1061
6.6	13.2	20.2	29.0	2.2	30.5	39.2							298	Broke with a long splinter, starting at knot.....	426
7.5	13.5	21.0	30.3	3.0	32.0	41.0	59.0						307	Broke at knot.....	426
6.8	14.5	22.5	33.0	2.5	34.0								216	do.....	926
8.2	16.5	28.0	39.0	5.0	41.7	57.0	83.0						304	do.....	239
6.6	13.8	23.5	34.0	4.3	35.0	44.5	70.0						309	Broke at knot with a large splinter.....	767
4.1	8.0	11.5	14.6	0.0	14.0	18.5	22.3	28.0	32.5	40.5	51.5		536	Broke with fine splinters.....	241
4.5	8.5	12.6	17.0	0.8	17.4	22.0	29.0	36.0	47.5				463	do.....	849
4.2	8.0	12.4	17.0	0.7	17.7	23.0	30.0	37.0	57.0				450	do.....	849
8.1	16.0	24.5	36.2	3.4	37.9								223	Sap-wood; split lengthwise without breaking.....	546
7.1	14.5	23.5	34.5	4.5	36.2	52.5							283	Sap-wood; crushed and splintered.....	546
5.5	10.5	16.0	22.5	1.0	23.2	30.2	45.0						331	Long, shattered break.....	1095
5.0	9.8	14.5	20.6	0.9	21.0	28.5							300	do.....	1095
6.5	12.7	20.5	31.0	3.5	32.0	45.0	79.0						320	Sap-wood; broke with fine splinters.....	1173
7.0	12.0	19.7	29.0	2.7	30.4	42.0	67.0						315	do.....	1173
6.5	12.5	19.0	32.2	3.6	35.0								236	Broke with long splinters.....	1181
6.2	11.5	17.6	28.7	4.0	31.0								232	do.....	1181
5.0	9.6	14.6	20.4	1.0	21.2	29.2							286	Crushed at center bearing; broke with one long splinter.....	1182
4.2	9.2	14.7	22.0	1.2	23.0	36.0							262	Broke with long, coarse splinters.....	1182
6.0	11.0	17.6	25.0	1.4	25.6	35.0							282	Crushed at center bearing; shattered.....	1183
4.8	10.0	15.5	23.7	2.4	26.0	40.0							263	do.....	1183
3.0	6.0	8.4	11.0	0.2	11.0	13.5	17.0	20.0	24.0	28.0	32.2	38.3	558	Broke with coarse splinters.....	485
3.0	5.8	8.5	11.5	0.2	11.5	14.5	18.0	21.6	25.7	29.6			472	do.....	485
6.0	10.7	16.0	21.6	0.9	22.5	28.3	35.5	47.5					354	Specimen cross-grained; split with grain.....	489
4.6	8.6	12.7	17.5	0.6	17.5	22.0	28.5	35.0	45.0				450	Splintered.....	489
7.0	15.4												116	Broke at knot.....	507
6.3	12.0	19.5	25.7	1.2	26.5	36.0	51.0						326	Shattered.....	507

TABLE III.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Specific gravity of the air-dried specimen.	Direction of grain.	COEFFICIENT OF ELASTICITY.		Modulus of rupture.
								First deflection.	Second deflection.	
MYRTACEÆ.										
144. <i>Eugenia buxifolia</i> <i>Gurgoon Stopper. Spanish Stopper.</i>	1118	Florida	Lost Man's river.	A. H. Curtiss	Humus and coral	1.0635		1637	1575	1055
146. <i>Eugenia monticola</i> <i>Stopper. White Stopper.</i>	1135	do	Umbrella Key	do	Coral	0.9405		1017	1085	1172
148. <i>Eugenia procera</i> <i>Red Stopper.</i>	1127	do	Miami	do	do	0.9066		1191	1206	1179
	1127	do	do	do	do	1.0023		1163	1177	1172
CORNACEÆ.										
151. <i>Cornus florida</i> <i>Flowering Dogwood. Box Wood.</i>	67	Missouri	Allenton.	G. W. Letterman	Upland	0.8987		787	814	872
	67	do	do	do	do	0.8904		787	849	1012
	761	Florida	Chattahoochee.	A. H. Curtiss	Calcareous	0.7795		787	794	830
	812	West Virginia	Grafton.	C. G. Pringle	Dry	0.7980		787	849	951
	812	do	do	do	do	0.7947		763	814	886
	1077	Missouri	Allenton.	G. W. Letterman	Gravelly	0.8647		827	849	1015
	1077	do	do	do	do	0.8490		888	957	966
	1092	do	do	do	Flinty	0.8387		610	647	710
152. <i>Cornus Nuttallii</i> <i>Flowering Dogwood.</i>	960	Oregon	Portland.	G. Engelmann and C. S. Sargent.		0.7763		814	976	990
	960	do	do	do		0.7807		976	1085	1052
153. <i>Nyssa capitata</i> <i>Ogeechee Lims. Sour Tupelo. Gopher Plum.</i>	605	Georgia	Ogeechee river.	A. H. Curtiss	Swampy	0.5739		610	638	694
	605	do	do	do	do	0.6170		697	723	698
154. <i>Nyssa sylvatica</i> <i>Tupelo. Sour Gum. Pepperidge. Black Gum.</i>	235	South Carolina	Bonneau's Depot	H. W. Ravenel	Muck	0.5966		687	781	783
	235	do	do	do	do	0.5735		888	849	745
	517	Tennessee	Cumberland river.	A. Gattinger		0.5979		814	814	689
	759	Florida	Chattahoochee.	A. H. Curtiss	Clay	0.7885		800	849	906
	759	do	do	do	do	0.7986		763	781	952
	813	West Virginia	Grafton.	C. G. Pringle		0.6222		939	888	788
	813	do	do	do		0.6447		888	1039	890
	833	Massachusetts	West Newbury	J. Robinson	Rich	0.7364		740	769	912
	833	do	do	do	do	0.7534		651	814	796
	864	do	do	do	do	0.7233		660	713	827
	894	do	do	do	do	0.6800		814	872	924
	835	do	Chebacco pond.	do	do	0.7914		642	651	717
155. <i>Nyssa uniflora</i> <i>Large Tupelo. Cotton Gum. Tupelo Gum.</i>	128	South Carolina	Bonneau's Depot	H. W. Ravenel	Swampy	0.6648		542	592	701
	128	do	do	do	do	0.6135		561	564	724
	550	Alabama	Stockton.	C. Mohr	Alluvial	0.5455		444	471	628
	550	do	do	do	do	0.5228		595	585	687
	604	Georgia	Ogeechee river.	A. H. Curtiss	Swampy	0.5739		456	444	625
	604	do	do	do	do	0.5841		488	444	553
CAPRIFOLIACEÆ.										
156. <i>Sambucus glauca</i> <i>Elder.</i>	681	California	Contra Costa county.	G. R. Vasey	Gravelly	0.5216		348	305	370
159. <i>Viburnum prunifolium</i> <i>Black Haw. Stag Bush.</i>	1103	Kentucky	Mercer county.	W. M. Linney	Hudson River shale.	0.8352		976	957	729
	1104	do	do	do	Trenton limestone	0.9034		904	1028	1222
	739	Georgia	Bainbridge	A. H. Curtiss	Clay	0.8270		708	734	896
RUBIACEÆ.										
160. <i>Excoecuma Caribæum</i>	466	Florida	Upper Metacombe Key.	A. H. Curtiss	Coral	0.9554		1136	1085	956
	466	do	do	do	do	0.9524		1221	1302	1055
161. <i>Pinckneya pubens</i> <i>Georgia Bark.</i>	381	South Carolina	Bluffton	J. H. Mellichamp	Sandy swamp	0.5425		660	683	496

UNITED STATES UNDER TRANSVERSE STRAIN—Continued.

DEFLECTION, IN MILLIMETERS, UNDER A PRESSURE, IN KILOGRAMS, OF—													Ultimate strength: transverse pressure.	Remarks.	Office number.
50	100	150	200	0 (set.)	200	250	300	350	400	450	500	550			
3.0	6.2	9.5	12.5	0.3	13.0	16.5	21.0	26.0	32.0	43.5			450	Shattered.....	1118
4.8	9.0	13.5	18.5	0.7	18.5	23.5	30.3	37.0	46.5	61.5			500	Splintered.....	1126
4.1	8.1	12.5	17.0	0.9	17.9	23.0	28.5	36.0	46.5	62.0			503	Specimen cross-grained; splintered.....	1127
4.2	8.3	12.5	16.8	0.6	17.5	22.4	29.0	35.0	46.0	61.5			500	Broke with coarse splinters.....	1127
6.2	12.0	18.2	26.9	2.2	27.6	36.0	50.0	70.0					372	Broke at knot.....	67
6.2	11.5	17.0	24.5	2.0	25.5	32.7	43.1	58.2	84.5				432	Maximum deflection, 120 millimeters; broke with large splinters...	67
6.2	12.3	20.0	29.0	3.5	31.2	40.5	56.0						350	Broke with large splinters.....	761
6.2	11.5	18.2	26.7	2.2	28.5	36.5	52.0	68.0	110.0				406	Square break on tension side with large flakes.....	812
6.4	12.0	18.7	27.0	2.5	28.6	39.0	55.0	87.0					378	Square break on tension side with long splinters.....	812
5.9	11.5	16.7	23.6	1.5	24.5	32.0	43.0	60.5	81.5				433	Broke with long, coarse splinters.....	1077
5.5	10.2	16.3	24.0	1.6	24.6	32.0	42.5	57.0					412	Shattered one end.....	1077
8.0	15.2	24.5	35.3	3.6	37.0	48.5	76.0						303	Specimen cross-grained; broke with large splinters.....	1092
6.0	10.0	14.5	20.3	0.9	20.3	25.5	32.0	42.5					397	Broke with long splinters.....	969
5.0	9.0	13.2	17.5	0.4	18.0	23.0	29.0	34.0	43.0				449	do.....	969
8.0	15.3	25.0	37.3	3.3	38.0	52.0							296	Broke short and split in axis.....	605
7.0	13.5	20.3	29.5	2.2	30.5	41.0							285	Shattered.....	605
7.1	12.5	19.6	27.5	1.6	28.2	37.5	51.0						334	do.....	236
5.5	11.5	18.0	25.5	1.5	26.3	34.7	54.5						318	do.....	236
6.0	12.0	19.5	28.7	2.4	29.5	38.0							294	Long split at one end.....	514
6.1	11.5	17.7	25.7	2.0	26.0	34.0	46.0	63.0	93.0				425	Broke into fine splinters.....	750
6.4	12.5	19.0	26.5	2.0	27.5	37.0	48.5	74.0	120.0				406	Deflected 150 millimeters and slipped from the bearing.....	750
5.2	11.0	16.2	22.5	0.4	23.0	29.0	38.0						334	Shattered.....	812
5.5	9.4	13.5	18.0	0.4	18.2	22.0	27.5	35.0					388	do.....	812
6.6	12.7	19.7	28.7	2.3	29.7	38.0	53.0	84.0					389	Sap-wood; broke with fine splinters.....	838
7.5	12.0	19.0	29.0	3.0	30.0	42.0	57.5	95.0					377	do.....	838
7.4	13.7	21.6	32.0	3.5	33.5	45.0	66.0	116.0					353	do.....	894
6.0	11.2	17.0	25.0	2.2	25.2	32.5	44.0	64.0					394	Failed from small splinter on corner.....	894
7.6	15.0	26.0	38.5	5.5	40.6	58.0	100.0						306	Crumpled on compression side at knot; square break with fine splinters.....	895
9.0	16.5	29.0	44.5	7.2	46.0	60.0							299	Broke with long splinters.....	128
8.7	17.3	28.7	43.0	7.5	45.5	66.2	108.0						309	Shattered at the end.....	128
11.0	20.7	33.0	48.5	8.5	53.0	82.0							268	Crushed at center bearing; broke with long splinters.....	550
8.2	16.7	26.7	39.0	5.0	40.8	59.5							293	do.....	550
10.7	22.0	36.5	57.5	11.5	62.5	108.0							271	Failed from large splinter on corner.....	604
10.0	22.0	36.5	58.0	12.2	60.5								236	do.....	604
14.0	32.0	55.0											158	Broke at knot.....	681
5.0	10.2	14.7	20.6	3.5	21.0	27.0	34.0						311	do.....	110 ²
5.4	9.5	14.6	19.7	0.9	21.0	26.0	33.5	41.0	51.0	68.5	92.0		524	Broke at knot with one large splinter.....	110 ⁴
6.9	13.3	20.7	29.0	3.0	31.0	40.0	53.0	71.0					383	Shattered.....	730
4.3	9.0	11.5	15.5	0.4	15.7	20.0	23.7	28.0	33.5				408	Specimen cross-grained; split with grain.....	466
4.0	7.5	11.4	15.2	0.5	15.4	19.0	23.0	28.5	33.0	38.0			450	Broke with fine splinters.....	466
7.4	14.3	22.2											173	Broke at knot with large splinters.....	381

TABLE III.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Specific gravity of the air-dried specimen.	Direction of grain.	COEFFICIENT OF ELASTICITY.		Modulus of rupture.
								First deflection.	Second deflection.	
ERICACEÆ.										
165. <i>Andromeda ferruginea</i>	1083	Florida	Jacksonville	A. H. Curtiss	Hummock	0.7273		814	814	680
166. <i>Arbutus Menziesii</i>	643	California	Contra Costa county	G. R. Vasey	Gravelly	0.7200		740	794	898
	643	do	do	do	do	0.7000		610	626	827
	679	do	Marin county	do	do	0.7454		976	1062	1015
	679	do	do	do	do	0.7546		842	872	898
167. <i>Arbutus Xalapensis</i>	683	Arizona	Santa Rita mount- ains	G. Engelmann and C. S. Sargent		0.7670		651	683	689
	683	do	do	do	do	0.7501		581	549	546
169. <i>Oxydendrum arboreum</i>	353	Alabama	Cottage Hill	C. Mohr	Light, rich	0.7386		787	800	469
	353	do	do	do	do	0.7501		939	976	979
170. <i>Kalmia latifolia</i>	262 ²	Virginia	Fancy Gap	H. Shriver	Moist	0.7642		651	670	696
	262 ²	do	do	do	do	0.6901		465	500	564
171. <i>Rhododendron maximum</i>	263 ¹	do	do	do	do	0.6216		651	673	691
	263 ²	do	do	do	do	0.6408		610	618	683
SAPOTACEÆ.										
175. <i>Chrysophyllum oliviforme</i>	492	Florida	Bay Biscayne	A. H. Curtiss	Coral	0.9663		1136	1252	947
	492	do	do	do	do	0.9583		921	996	1227
176. <i>Sideroxylon Mastichodendron</i>	461	do	Upper Metacombe Key	do	do	0.9872		976	1149	919
	461	do	do	do	do	0.9827		976	1050	1022
177. <i>Diphollis salicifolia</i>	488	do	Bay Biscayne	do	do	0.9802		1356	1436	1263
	488	do	do	do	do	0.9173		1395	1395	1171
	500	do	Umbrella Key	do	do	0.8823		1191	1177	1008
178. <i>Bumelia tenax</i>	746	Georgia	Bainbridge	do	Low	0.7914		751	751	673
179. <i>Bumelia lanuginosa</i>	930	Texas	Austin	C. Mohr	Limestone	0.5847		488	496	567
	930	do	do	do	do	0.5903		478	432	265
	1083	Missouri	Allenton	G. W. Letterman	do	0.7148		488	522	330
181. <i>Bumelia lycioides</i>	333	Tennessee	Nashville	A. Gattlinger	Alluvial	0.8061		697	781	563
182. <i>Bumelia cuneata</i>	1124	Florida	Boca Chica Key	A. H. Curtiss	Coral	0.8630		660	608	516
183. <i>Mimusops Sieberi</i>	458	do	Upper Metacombe Key	do	do	1.0786		996	1028	953
	458	do	do	do	do	1.0191		939	976	874
EBENACEÆ.										
184. <i>Diospyros Virginiana</i>	61	Missouri	Allenton	G. W. Letterman	Rich upland	0.7633		814	794	483
	61	do	do	do	do	0.7862		751	751	818
	425	Tennessee	Nashville	A. Gattlinger	Rich loam	0.8716		1395	1337	1289
	811	West Virginia	Grafton	C. G. Fringle		0.8176		610	622	895
	811	do	do	do		0.8119		610	603	883
	1084	Missouri	Allenton	G. W. Letterman	Rich upland	0.8125		751	814	933
	1084	do	do	do	do	0.8410		574	603	846
187. <i>Balesia diptera</i>	738	Georgia	Bainbridge	A. H. Curtiss	Low	0.6704		642	673	881
	738	do	do	do	do	0.6577		660	697	882
STYRACACEÆ.										
186. <i>Symplocos tinctoria</i>	347	Alabama	Cottage Hill	C. Mohr	Sandy	0.5580		610	622	619
187. <i>Balesia diptera</i>	738	Georgia	Bainbridge	A. H. Curtiss	Low	0.6704		642	673	881
.....	738	do	do	do	do	0.6577		660	697	882

UNITED STATES UNDER TRANSVERSE STRAIN—Continued.

DEFLECTION, IN MILLIMETERS, UNDER A PRESSURE, IN KILOGRAMS OF—													Ultimate strength: transverse pressure.	Remarks.	Office number.
50	100	150	200	0 (set.)	200	250	300	350	400	450	500	550			
6.0	12.0	18.5	26.0	2.0	27.5	35.0							290	Broke at knot	1033
6.6	12.3	19.5	27.0	1.8	28.0	35.0	47.5	62.5					381	Specimen cross-grained	643
8.0	15.6	24.0	33.2	2.2	35.0	44.5	59.0						353	do	643
5.0	9.2	14.0	19.0	0.6	20.0	25.5	32.5	41.5	53.0				433	Square break	679
5.8	11.2	16.3	22.5	1.0	23.5	30.0	38.2	51.0					383	Threw off large flakes on tension side	679
7.5	14.3	22.0	31.5	2.5	32.8	43.0							294	Large flake on tension side	683
9.1	17.8	29.0	43.5	5.5	46.0								233	Broke at knot; flake on compression side	683
6.2	12.2	19.0	31.5										200	Specimen cross-grained; splinter started at large knot	353
5.2	10.0	15.0	21.0	1.2	21.5	28.0	36.0	47.0	65.0				422	Crushed at center bearing; broke with fine splinters	353
7.5	14.6	22.4	31.2	2.6	33.6	44.5							297	Broke at knot	262*
10.5	19.5	31.0	44.0	4.8	47.0								240	Specimen cross-grained; shattered; long split with grain	262*
7.5	14.5	22.8	32.5	2.3	33.5	44.0							205	Threw off flake from tension side	263 ¹
8.0	15.8	25.0	34.2	2.8	35.5	46.0							270	Broke at knot	263*
4.3	7.8	12.0	16.4	0.9	17.4	21.8	28.2	35.0	46.0				404	Specimen cross-grained	492
5.3	9.8	15.5	21.0	1.0	22.0	33.5	37.2						328	do	492
5.0	8.5	12.8	17.5	0.6	18.0	24.0	30.0	41.0					392	Broke with large splinters	461
5.0	9.3	13.6	18.3	0.9	19.0	23.5	30.0	38.0	49.0				436	Broke with coarse splinters	461
3.6	6.8	10.0	13.0	0.8	13.7	17.0	20.0	25.0	30.0	37.0	45.0		539	Broke with long, coarse splinters	483
3.5	7.0	10.6	14.0	0.8	14.5	18.0	22.0	27.0	31.5	37.5	49.0		500	do	483
4.1	8.3	12.3	16.6	0.4	17.0	21.5	27.0	33.5	40.0				430	Broke at knot with long, coarse splinters	500
6.5	13.0	20.0	29.5	2.8	31.5	41.6							287	Specimen cross-grained; broke at knot	746
10.0	19.7	31.5	46.0	7.0	49.2								242	Broke with a large splinter	930
10.2	22.6												113	Broke with large splinter at knot	930
10.0	18.7												141	Defective specimen; broke at knot	1033
7.0	12.5	19.5	28.5	2.4	29.7								240	Broke at knot with one large splinter	333
7.4	16.2	26.7	40.6	7.5	43.5								220	Specimen cross-grained; large flake on compression side	1124
4.9	9.5	14.7	19.7	1.0	20.0	25.2	32.0	39.7	48.7				406	Specimen cross-grained; broke at knot	458
5.2	10.0	15.0	20.2	0.8	21.0	26.5	33.2	42.5					373	Specimen cross-grained; large flake from compression side	458
6.0	12.3	18.0	26.2	2.0	28.0	36.0	51.0						206	Specimen cross-grained; short break on tension side, flake from compression side	61
6.5	13.0	19.4	29.0	3.2	29.7	39.0	55.0						349	Short break on tension side; flake from compression side	61
3.5	7.3	11.0	15.0	0.2	15.0	19.0	24.2	30.0	35.2	43.0	56.0	75.0	550	Broke with fine splinters	425
8.0	15.7	24.3	35.3	4.0	37.9	50.0	70.6	111.5					382	do	811
8.0	16.2	25.7	38.0	4.5	39.0	52.5	79.0	115.5					377	Deflected 170 millimeters before breaking; broke with fine splinters	811
6.5	12.0	19.3	28.5	2.5	30.0	38.0	52.0	73.0					396	Square break on tension side, large flake on compression side	1084
8.5	16.2	26.0	38.0	5.2	40.2	55.0	78.0	135.0					361	do	1084
7.0	13.3	21.0	30.4	3.0	31.8	40.5	58.0	78.0					377	do	1162
8.0	15.7	25.6	40.0	5.5	42.0	63.0							264	Specimen cross-grained	347
7.6	14.5	23.0	32.0	2.5	33.0	44.0	59.0	82.0					376	Shattered at one end	738
7.4	14.0	22.5	32.0	2.6	33.0	43.0	60.0	96.5					355	Broke with large splinters	738

TABLE III.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Specific gravity of the air-dried specimen.	Direction of grain.	COEFFICIENT OF ELASTICITY.		Modulus of rupture.
								First deflection.	Second deflection.	
OLEACEÆ.										
191. <i>Fraxinus pistaciifolia</i> Ash.	660	Arizona	Santa Rita mountains.	G. Engelmann and C. S. Sargent.	Low.....	0.6633		514	542	666
	660	do	do	do	do	0.6995		698	660	578
192. <i>Fraxinus Americana</i> White Ash.	392	Missouri	Allenton	G. W. Letterman	do	0.6592		1110	1220	928
	114 ¹	Michigan	Dansville	W. J. Beal	Clay	0.4983		542	531	586
	114 ¹	do	do	do	do	0.5409		608	679	619
	114 ³	do	Hudson	do	do	0.7535		1061	1163	1041
	114 ⁴	do	Lansing	do	do	0.6622		1191	1284	927
	114 ⁵	do	Dansville	do	Clay	0.5920		814	872	743
	130	South Carolina	Bonneau's Depot	H. W. Ravenel	Wet	0.6914		1191	1284	973
	130	do	do	do	do	0.6864		1136	1221	951
	212	Virginia	Wytheville	H. Shriver	do	0.6569		976	1085	855
	212	do	do	do	do	0.6524		939	1007	907
	227	Vermont	Charlotte	C. G. Pringle	Gravelly	0.6347		814	930	813
	227 ¹	do	do	do	do	0.6613		888	957	860
	227 ²	do	do	do	do	0.7243		1221	1302	1059
	227 ³	do	do	do	do	0.7151		1479	1525	1289
	267 ²	Virginia	Wytheville	H. Shriver	do	0.6250		904	976	820
	431	Tennessee	Nashville	A. Gattinger	Limestone	0.6792		1191	1221	968
	551	Alabama	Kemper's mill	C. Mohr	Alluvial	0.6417		976	1085	803
	551	do	do	do	do	0.6038		1085	976	764
	747	Georgia	Bainbridge	A. H. Curtiss	River-bottom	0.6512		660	642	415
	747	do	do	do	do	0.6250		549	564	693
	937	Texas	Austin	C. Mohr	Rich, calcareous	0.7293		697	708	611
	1045	Massachusetts	Reading	J. Robinson	do	0.7954		814	957	1017
1045	do	do	do	do	0.8205		1221	1191	1101	
192. <i>Fraxinus Americana</i> , var. <i>Texensis</i> .	364	Texas	Dallas	J. Reverchon	Dry, calcareous	0.8194		1085	1028	1115
	364	do	do	do	do	0.7762		1163	1135	1134
193. <i>Fraxinus pubescens</i> Red Ash.	139	Michigan	Lansing	W. J. Beal	Poor	0.6416		976	1122	1073
	229	Vermont	Charlotte	C. G. Pringle	Clay	0.7065		729	752	780
	229	do	do	do	do	0.6812		976	976	1040
	1059	Massachusetts	Topsfield	J. Robinson	River-bottom	0.6080		407	398	581
194. <i>Fraxinus viridis</i> Green Ash.	57	Missouri	Allenton	G. W. Letterman	Rich, wet	0.7273		610	592	816
	308	Texas	Dallas	J. Reverchon	do	0.5977		610	610	544
	308	do	do	do	do	0.6522		888	930	820
	438	Tennessee	Nashville	A. Gattinger	Rich upland	0.7683		976	1085	961
	948	Texas	Victoria	C. Mohr	Rich, wet	0.7817		888	849	944
	948	do	do	do	do	0.8125		904	976	986
	957	do	Matagorda bay	do	do	0.7022		775	781	937
	957	do	do	do	do	0.7684		1628	1575	1179
	957	do	do	do	do	0.6893		697	729	876
195. <i>Fraxinus platycarpa</i> Water Ash.	536	Alabama	Stockton	do	Rich, alluvial	0.4759		443	476	537
196. <i>Fraxinus quadrangulata</i> Blue Ash.	66	Missouri	Allenton	G. W. Letterman	Dry upland	0.8424		814	787	951
	66	do	do	do	do	0.8620		842	888	949
	125	Michigan	Lansing	W. J. Beal	Rich loam	0.5189		740	751	677
	125	do	do	do	do	0.5206		634	687	633

UNITED STATES UNDER TRANSVERSE STRAIN—Continued.

DEFLECTION, IN MILLIMETERS, UNDER A PRESSURE, IN KILOGRAMS, OF—													Ultimate strength: transverse pressure.	Remarks.	Office number.
50	100	150	200	0 (set.)	200	250	300	350	400	450	500	550			
9.5	18.0	29.5	42.8	5.2	44.5	64.0							284	Square break on tension side with large flake on compression side.	660
7.0	14.8	24.2	36.5	4.4	38.0								247	do	660
4.4	8.0	11.5	15.9	0.4	16.5	21.0	28.0	37.0					396	Broke with coarse splinters	393
9.0	18.4	29.5	47.4	6.2	50.6								250	Long, shattered break	1141
7.0	14.4	23.2	35.5	4.2	39.0	54.0							264	Short break on tension side	1141
4.6	8.4	12.3	17.0	1.0	17.5	22.5	30.7	39.0	56.5				444	Sap-wood; broke with fine splinters	1143
4.1	7.6	11.5	15.0	0.4	15.2	19.0	24.0	30.5	41.0				415	Broke with coarse splinters	1144
6.0	11.2	17.8	26.3	2.0	27.5	37.2	58.0						317	Broke with fine splinters	1145
4.1	7.6	11.6	16.0	0.3	16.0	20.7	26.2	33.2	45.5				415	Broke with large splinters on tension side	130
4.3	8.0	11.4	15.8	0.3	16.0	20.4	26.4	35.0	47.0				406	Broke with fine splinters on tension side	130
5.0	9.0	14.0	19.0	0.5	19.5	26.0	34.5	50.0					365	Broke with large splinters on tension side	212
5.2	9.7	14.6	20.0	1.0	20.3	26.6	35.0	50.0					387	do	212
6.0	10.5	15.5	22.0	1.0	22.0	29.3	39.0						347	do	227
5.5	10.2	16.0	22.8	1.1	23.5	31.0	41.5	60.0					367	do	2271
4.0	7.5	11.3	15.2	0.3	15.3	19.3	24.8	30.2	38.2	51.2			452	Specimen cross-grained; broke with long splinters	2272
3.3	6.4	9.8	13.0	0.2	13.0	16.0	19.7	23.7	29.0	35.5	45.0	59.0	550	Failed from splinter on corner	2273
5.4	10.0	14.2	19.5	0.6	20.7	27.0	36.0						350	Square break on tension side with flake on compression side	2673
4.1	8.0	12.2	16.5	0.5	17.0	21.2	27.5	35.0	48.5				413	Square break with large splinters on tension side	431
5.0	9.0	13.0	19.0	1.0	19.5	26.0	35.0						343	Broke with large splinters on tension side	551
4.5	10.0	14.5	21.5	1.3	22.5	30.0	42.0						326	do	551
7.4	15.2	25.3											177	Specimen cross-grained; broke at knot	747
8.9	17.3	27.6	39.2	4.2	41.0	56.2							296	Broke with large splinters on tension side	747
7.0	13.8	22.0	32.3	3.6	33.2	44.5							261	Specimen cross-grained; broke at knot	937
6.0	10.2	15.2	21.5	1.5	22.0	28.2	36.5	47.0	65.0				434	Broke with large splinters on tension side	1045
4.0	8.2	12.8	17.0	0.9	17.5	22.5	29.0	37.0	49.5	72.0			470	Broke with fine splinters on tension side	1045
4.5	9.5	13.0	17.6	0.7	18.3	23.0	30.7	39.5	53.6	79.5			476	Broke with fine splinters	364
4.2	8.6	13.0	18.0	0.5	18.4	23.5	30.5	38.0	46.0	63.5			484	Broke with fine splinters; buckled on compression side	364
5.0	8.7	12.8	17.6	1.0	17.7	22.5	28.5	35.4	45.7	63.2			458	Crushed at center bearing; broke with fine splinters	139
6.7	13.0	21.5	32.2	5.0	33.2	45.5	64.5						333	Broke with large flake on compression side	229
5.0	10.0	15.0	21.0	1.7	21.6	27.3	35.4	47.0	65.7				444	Failed from two large splinters on angles of tension side	229
12.0	24.5	44.2	76.0	18.5	81.0								248	Square break with large flake on compression side	1059
8.0	16.5	27.2	39.5	5.8	41.0	56.0	94.0						348	Broke with small splinters	57
8.0	16.0	24.5	36.0	3.5	37.6								232	Sap-wood; square break, splitting in axis	308
5.5	10.5	15.5	21.0	0.8	21.3	27.5	38.0	51.0					350	Specimen cross-grained; split in axis	308
5.0	9.0	13.5	19.0	1.0	19.5	24.6	32.4	42.0	61.0				410	Broke with fine splinters	438
5.5	11.5	18.0	26.2	2.4	26.5	35.0	47.0	64.5	110.0				403	do	948
5.4	10.0	15.0	21.0	1.1	21.0	28.5	38.0	50.5	90.0				421	do	948
6.3	12.5	19.0	26.1	1.8	26.4	34.0	45.5	60.0					400	do	957
3.0	6.2	9.4	12.7	0.5	12.9	16.0	21.0	26.5	32.5	41.0	58.0		503	do	957
7.0	13.4	20.6	30.0	2.5	30.2	41.6	57.0	84.0					374	Square break on tension side; split in axis	957
11.0	20.5	33.5	52.2	7.0	56.0								229	Short break; shattered	536
6.0	12.4	19.5	27.2	2.0	28.0	38.0	52.0	72.5					406	Broke with fine splinters at small knot on compression side	66
5.8	11.0	17.0	23.2	1.4	24.0	31.2	41.2	58.5					405	do	66
6.6	13.0	20.2	29.4	2.1	30.5	43.0							289	Split and shattered	125
7.7	14.2	22.0	32.0	2.0	33.0	44.0							270	Square break on tension side with large flake on compression side	125

TABLE III.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Specific gravity of the air-dried specimen.	Direction of grain.	COEFFICIENT OF ELASTICITY.		Modulus of rupture.
								First deflection.	Second deflection.	
196. <i>Fraxinus quadrangulata</i> —cont'd	2867	Kentucky	Mercer county	W. M. Linney	Limestone	0.7989		976	1028	1104
	2867	do	do	do	do	0.7914		697	697	715
	291	Missouri	Allenton	G. W. Letterman	Sandy loam	0.6080		651	697	734
	518	Tennessee	Nashville	A. Gattinger	Rich limestone	0.7923		610	651	731
197. <i>Fraxinus Oregona</i> <i>Oregon Ash.</i>	964	Oregon	Portland	G. Engelmann and C. S. Sargent	Low, wet	0.6182		939	948	820
	964	do	do	do	do	0.6057		842	939	757
	1001	do	Woldier's saw-mill	do	do	0.6285		888	913	803
	1001	do	do	do	do	0.6001		740	781	729
	1024	do	Portland Furniture Company	do	do	0.4894		478	474	349
	1024	do	do	do	do	0.4832		425	425	351
	1030	do	do	do	do	0.6430		1062	1085	738
	1030	do	do	do	do	0.6419		1220	1220	766
198. <i>Fraxinus sambucifolia</i> <i>Black Ash. Hoop Ash. Ground Ash.</i>	122	Michigan	Dansville	W. J. Beal	Wet, peaty	0.5215		814	814	613
	147	Illinois	Waukegan	Robert Douglas	Low, wet	0.7371		976	1062	977
	839	Massachusetts	Danvers	J. Robinson	Rich, loamy	0.7365		888	849	820
	839	do	do	do	do	0.7575		814	763	811
199. <i>Forestiera acuminata</i> <i>Privet.</i>	737	Georgia	Bainbridge	A. H. Curtiss	do	0.5875		610	592	698
	737	do	do	do	do	0.5814		872	814	734
201. <i>Osmanthus Americanus</i> <i>Devil Wood.</i>	283	Louisiana	Amite	C. Mohr	Rich, alluvial	0.8762		1221	1328	1202
	283	do	do	do	do	0.8455		1062	1221	1015
	584	Florida	Saint John's river	A. H. Curtiss	Sandy loam	0.7330		1085	1136	937
BORRAGINACEÆ.										
204. <i>Borreria Havanaensis</i> <i>Strong Bark.</i>	1137	do	Key Largo	do	Coral	0.7984		888	996	945
205. <i>Ehretia elliptica</i> <i>Knackaway. Anaqua.</i>	942	Texas	New Braunfels	C. Mohr	Rich, alluvial	0.6649		425	397	722
BIGNONIACEÆ.										
206. <i>Catalpa bignonioides</i> <i>Catalpa. Catawba. Bean Tree. Cigar Tree. Indian Bean.</i>	540	Alabama	Stockton	do	Low, wet	0.4816		669	697	682
	744	Georgia	Bainbridge	A. H. Curtiss	Clay	0.4413		574	610	523
	744	do	do	do	do	0.4352		567	531	469
	744	do	do	do	do	0.4585		814	888	691
207. <i>Catalpa speciosa</i> <i>Western Catalpa.</i>	38	Missouri	Charleston	C. S. Sargent	Wet clay	0.4915		787	849	673
	38	do	do	do	do	0.4915		763	794	585
208. <i>Chilopsis saligna</i> <i>Desert Willow.</i>	682	Arizona	Tucson	G. Engelmann and C. S. Sargent	Moist, gravelly	0.6193		514	514	460
	682	do	do	do	do	0.5716		555	574	680
VERBENACEÆ.										
210. <i>Citharexylum villosum</i> <i>Fiddle Wood.</i>	490	Florida	Bay Biscayne	A. H. Curtiss	Coral	1.0323		1163	1177	708
NYCTAGINACEÆ.										
212. <i>Pisonia obtusata</i> <i>Pigeon Wood. Beef Wood. Cork Wood. Pork Wood.</i>	474	do	Upper Metacombe Key	do	do	0.6884		465	465	296
POLYGONACEÆ.										
213. <i>Coccoloba floridana</i> <i>Pigeon Plum.</i>	473	do	do	do	do	0.9718		1163	1221	937
	473	do	do	do	do	0.9732		1017	1050	896
LAURACEÆ.										
215. <i>Persea carolinensis</i> <i>Red Bay.</i>	585	do	Saint John's river	do	Sandy loam	0.6427		814	864	921
	585	do	do	do	do	0.6489		814	814	888
215. <i>Persea carolinensis</i> , var. <i>palustris</i> .	840	Alabama	Mobile county	C. Mohr	Damp, sandy	0.6216		814	839	820

UNITED STATES UNDER TRANSVERSE STRAIN—Continued.

DEFLECTION, IN MILLIMETERS, UNDER A PRESSURE, IN KILOGRAMS, OF—													Ultimate strength: transverse pressure	Remarks.	Office number.
50	100	150	200	0 (set.)	200	250	300	350	400	450	500	550			
5.0	9.5	14.8	19.3	0.6	20.0	25.0	31.2	38.0	47.0	61.0			471	Specimen cross-grained; split	286 ¹
7.0	14.0	21.0	29.6	2.7	32.0	44.2	66.0						305	Failed at knot	286 ²
7.5	14.0	21.4	30.6	2.3	32.0	41.0	59.0						313	Broke with coarse splinters; shattered	291
8.0	15.0	23.4	33.0	3.5	35.0	46.5	66.5						312	Shattered	518
5.2	10.3	15.0	20.8	0.7	21.5	28.0	39.0						350	Square break on tension side with split in axis	964
5.8	10.4	16.0	22.0	0.9	22.5	30.0	41.0						323	Shattered	964
5.5	10.7	16.2	21.6	0.5	22.5	28.0	36.0						343	Specimen cross-grained; square break on tension side, flake on compression side.	1001
6.6	12.5	18.7	25.7	1.0	26.0	34.0	45.5						311	Specimen cross-grained	1001
10.2	20.6												149	do	1024
11.5	23.0												150	Specimen cross-grained; flake on tension side	1024
4.6	9.0	13.0	18.0	0.6	18.5	23.5	30.2						315	Specimen cross-grained; break started at knot	1030
4.0	8.0	12.5	17.0	0.4	17.3	21.8							327	do	1030
6.0	12.0	19.0	26.5	1.6	28.7	39.0							261	Square break on tension side with coarse splinters	122
5.0	9.2	14.0	20.6	1.7	21.0	28.0	36.0	49.0	74.0				417	Square break with two large splinters	147
5.5	11.5	18.0	26.0	2.5	27.5	35.7	53.0	81.5					350	Square break on tension side with coarse splinters	839
6.0	12.8	22.0	31.2	3.9	34.0	43.0	71.0						346	do	839
8.0	16.5	26.5	38.7	4.5	42.0	61.0							298	Square break on tension side; shattered	737
5.6	12.0	18.0	27.0	2.0	28.2	38.0	64.5						313	do	737
4.0	7.3	11.2	15.3	0.4	17.7	20.0	24.5	31.0	37.0	46.0	60.5		513	Square break on tension side, splitting in the axis with fine splinters.	283
4.6	8.0	12.7	17.8	1.0	18.0	23.5	30.0	37.0	48.0				433	Sap-wood; specimen cross-grained; broke with long splinters	283
4.5	8.6	13.0	18.0	1.0	18.2	23.5	30.0	39.5	55.0				400	do	584
5.5	9.8	14.6	19.8	0.5	20.0	25.0	32.0	40.0	50.0				403	Specimen cross-grained; splintered	1137
11.5	24.6	36.0	53.0	6.9	56.0	77.5	131.0						308	Specimen cross-grained; square break on tension side; split in axis.	948
7.3	14.0	21.5	31.0	2.4	32.5	44.0							291	Broke with coarse splinters	540
8.5	16.0	26.0	37.6	4.5	41.0								223	Crushed at center bearing; broke with coarse splinters	744
8.6	18.4	31.0	63.0	17.5									200	do	744
6.0	11.0	18.0	25.2	1.4	26.0	35.0							295	do	744
6.2	11.5	17.0	23.5	1.2	24.3	33.2							287	Crushed at center bearing; splintered	38
6.4	12.3	20.4	30.0	2.6	30.7	53.0							254	do	38
9.5	19.0	31.6	50.5	6.0									200	Splintered on corner	682
8.8	17.0	26.0	37.0	3.0	38.0	54.0							294	Specimen cross-grained	683
4.2	8.3	12.5	17.5	0.7	18.0	23.0							300	Specimen cross-grained; broke at knots	490
3.7	7.3	11.4	15.0	0.4	15.7	20.0	25.0	30.0	37.0	46.0			499	Broke with fine splinters	490
10.5	21.0												127	Specimen cross-grained; failed from large splinters on corner	474
4.2	8.0	12.4	16.8	0.4	17.5	21.5	26.5	32.0					400	Shattered; large flakes on tension side	473
4.8	9.3	13.5	19.0	0.8	18.7	24.0	31.0	39.0					383	Broke with coarse splinters	473
6.0	11.3	18.0	24.3	1.0	25.0	31.9	40.0	50.0					393	Shattered	585
6.0	12.0	17.6	24.3	1.2	24.2	30.2	40.0	51.0					377	do	585
6.0	11.5	16.6	23.6	1.3	24.0	32.0	42.0	57.5					350	Broke with coarse splinters	240

TABLE III.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Specific gravity of the air-dried specimen.	Direction of grain.	COEFFICIENT OF ELASTICITY.		Modulus of rupture.
								First deflection.	Second deflection.	
217. <i>Sassafras officinale</i> <i>Sassafras.</i>	71	Missouri	Allenton	G. W. Letterman	Low, rich	0.5286		567	561	886
	71	do	do	do	do	0.4569		574	626	708
	387	do	do	do	Alluvial	0.4546		443	465	361
	387	do	do	do	do	0.4818		610	651	703
	440	Tennessee	Nashville	A. Gattinger	Rich	0.4824		651	673	658
	814	West Virginia	Grafton	C. G. Pringle	do	0.5849		567	542	600
	814	do	do	do	do	0.5828		488	501	673
	854	Massachusetts	Danvers	J. Robinson	Rich loam	0.4829		348	337	368
	854	do	do	do	do	0.5034		325	315	464
218. <i>Umbellularia Californica</i> <i>Mountain Laurel. California Laurel. Spice Tree. Cagiput. California Olive. California Bay Tree.</i>	703	Oregon	Coos bay	G. Engelmann and C. S. Sargent	do	0.6813		1085	1149	937
	703	do	do	do	do	0.6755		888	986	675
EUPHORBIACEÆ.										
219. <i>Drypetes crocea</i> <i>Guiana Plum. White Wood.</i>	408	Florida	Upper Metacombe Key.	A. H. Curtiss	Coral	0.9304		976	1039	797
219. <i>Drypetes crocea</i> , var. <i>latifolia</i>	459	do	do	do	do	0.9391		842	872	591
	459	do	do	do	do	0.9218		751	800	825
URTICACEÆ.										
222. <i>Ulmus crassifolia</i> <i>Cedar Elm.</i>	324	Texas	Dallas	J. Reverchon	Rich loam	0.6706		751	781	809
	324	do	do	do	do	0.7348		814	888	733
	929	do	Austin	C. Mohr	do	0.8375		626	610	787
	929	do	do	do	do	0.8682		519	531	762
223. <i>Ulmus fulva</i> <i>Red Elm. Slippery Elm. Moose Elm.</i>	301	Kentucky	Mercer county	W. M. Linney	Limestone	0.5875		976	976	783
	134	Missouri	Allenton	G. W. Letterman	Rich, alluvial	0.7200		888	930	937
	134	do	do	do	do	0.6933		1017	1062	937
	369	Vermont	Charlotte	C. G. Pringle	Gravelly	0.6767		814	939	973
	420	Tennessee	Nashville	A. Gattinger	Clay	0.6821		814	857	712
224. <i>Ulmus Americana</i> <i>White Elm. American Elm. Water Elm.</i>	19	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	0.6920		976	976	989
	19	do	do	do	do	0.7080		921	939	1050
	281	Missouri	Allenton	G. W. Letterman	Alluvial	0.6477		718	718	794
	958	Texas	Colorado river	C. Mohr	Alluvial	0.6596		751	800	813
	958	do	do	do	do	0.6108		729	835	695
	1036	Massachusetts	Danvers	J. Robinson	Gravelly	0.7370		400	391	574
	1036	do	do	do	do	0.7359		542	514	649
	1049	do	North Reading	do	do	0.7660		828	849	1010
225. <i>Ulmus racemosa</i> <i>Rock Elm. Cork Elm. Hickory Elm. White Elm. Olig Elm.</i>	1161	Michigan	Danville	W. J. Beal	do	0.8562		1136	1190	1361
	1161	do	Big Rapids	do	do	0.6245		814	921	886
	1161	do	do	do	Low, gravelly	0.6839		1136	1221	1055
	1161	do	Hudson	do	Alluvial	0.7461		996	1073	1094
	814	do	Hersey	do	Rich loam	0.7677		1220	1302	1165
	314	do	do	do	do	0.7466		1163	1268	1036
	428	Tennessee	Nashville	A. Gattinger	do	0.7442		651	697	869
226. <i>Ulmus alata</i> <i>Wahoo. Winged Elm.</i>	133	South Carolina	Bonneau's Depot	H. W. Ravenel	do	0.8768		452	479	745
	380	Tennessee	Davidson county	A. Gattinger	Loam	0.6615		574	568	703
	533	Mississippi	Kemper's Mill	C. Mohr	Alluvial	0.8243		610	651	820
	533	do	do	do	do	0.7883		697	687	900

UNITED STATES UNDER TRANSVERSE STRAIN—Continued.

DEFLECTION, IN MILLIMETERS, UNDER A PRESSURE, IN KILOGRAMS, OF—													Ultimate strength: transverse pressure.	Remarks.	Office number.
50	100	150	200	0 (set.)	200	250	300	350	400	450	500	550			
8.6	17.4	28.0	40.7	4.5	44.0	68.0							378	Broke short and split.....	71
8.5	15.6	23.4	33.8	2.7	35.5	47.5	80.0						302	Broke with fine splinters.....	71
11.0	21.0	35.5											154	Shattered.....	387
8.0	15.0	23.2	34.0	3.1	36.0								300	do.....	387
7.5	14.5	22.0	32.5	2.0	34.0	47.0							281	Coarse splinters.....	446
8.6	18.0	29.6	45.0	6.2	46.0	74.0							256	Shattered.....	814
10.0	19.5	30.0	42.0	4.9	45.7	63.0							287	do.....	814
14.0	29.0	47.0											157	Square break on tension side with large flake on compression side..	854
15.0	31.0	52.0											198	do.....	854
4.5	8.5	12.5	17.0	0.6	18.0	22.5	29.5	37.0					400	Specimen cross-grained; split.....	703
5.5	9.9	15.7	21.0	1.1	21.5	27.5							288	do.....	703
5.0	9.4	14.6	20.0	1.5	20.6	26.0	33.6						340	Shattered.....	468
5.8	11.2	17.3	24.5	1.6	25.0	32.5							252	Sap-wood; shattered.....	459
6.5	12.2	19.6	27.2	2.0	27.5	36.5	46.5	61.0					352	Specimen cross-grained; sap-wood; shattered.....	459
6.5	12.5	20.0	29.0	2.2	30.0	42.0	56.0						345	Broke with coarse splinters.....	324
6.0	11.0	16.8	23.5	1.3	24.5	31.0	39.2						313	Short break on tension side.....	324
7.8	16.0	27.5	40.5	6.0	42.0	57.0	84.0						336	Sap-wood; shattered.....	929
9.4	18.2	31.0	44.0	6.5	47.0	65.0	97.0						325	do.....	929
5.0	10.0	15.2	23.3	1.7	23.6	30.5	47.0						334	Crushed at center bearing; bent and splintered without breaking..	30 ¹
5.5	10.5	16.0	22.8	1.6	23.5	31.5	43.0	59.2	112.0				400	do.....	134
4.8	9.2	14.0	20.0	1.0	20.8	27.0	36.0	51.2	86.0				400	do.....	134
6.0	10.4	15.5	20.3	1.0	20.3	26.9	35.0	47.5	60.0				415	Thin scale on tension side.....	369
6.0	11.4	17.0	24.2	1.2	25.2	32.6	50.0						304	Broke with coarse splinters; started at knot.....	429
5.0	10.0	15.0	20.8	1.0	21.3	26.5	35.0	45.8	64.0				422	0.5 sap-wood; splintered on corners.....	19
5.3	10.4	14.7	20.4	0.5	21.0	27.0	34.7	45.7	60.0				448	do.....	19
6.8	13.6	21.2	31.0	3.1	31.0	41.0	60.0						339	Broke with coarse splinters.....	281
6.8	13.0	21.0	30.0	3.0	31.0	41.8	61.5						344	do.....	281
6.5	12.2	19.5	28.0	3.0	29.0	40.0	56.6						347	Broke with fine splinters.....	958
6.7	11.7	17.0	25.0	2.2	25.6	34.5	47.7	69.5					382	Square break on tension side, splitting in axis.....	958
12.2	25.0	41.5	69.2	14.4	73.0								245	Specimen cross-grained.....	1036
9.0	19.0	31.0	46.2	6.2	49.2	71.3							277	Broke with fine splinters.....	1036
5.9	11.5	17.5	25.5	2.0	26.2	33.2	45.6	63.0	98.2				431	do.....	1049
6.0	11.4	17.5	24.0	1.9	24.0	31.5	44.0	60.0					394	Buckled on compression side; fine splinters.....	1049
4.3	8.2	12.0	16.7	0.5	17.0	22.0	28.5	34.0	40.8	49.0	64.0	85.0	581	Broke with fine splinters.....	116 ¹
6.0	10.6	16.1	22.0	0.6	23.0	29.0	38.5	53.5					378	Crushed at center bearing; broke with fine splinters.....	116 ²
4.3	8.0	12.2	16.5	0.4	16.7	21.0	26.5	34.0	46.0	65.0			450	Broke with fine splinters.....	116 ³
4.9	9.1	13.7	18.2	0.7	18.4	23.0	30.0	38.0	48.5	67.0			467	do.....	116 ⁴
4.0	7.5	11.0	14.9	0.5	15.0	19.0	24.0	29.4	38.7	52.0			497	do.....	314
4.2	7.7	11.7	15.8	0.5	16.0	20.3	26.0	32.0	44.0				442	do.....	314
7.5	14.0	22.4	33.2	3.5	35.0	45.0	64.0	91.5					371	do.....	428
10.8	20.4	32.0	45.6	5.5	47.0	64.5	94.0						318	Sap-wood; failed at knot.....	133
8.5	17.2	28.8	43.5	7.2	45.5	59.3	85.5						300	0.75 heart wood; splintered at corners.....	380
8.0	15.6	23.7	34.0	3.3	35.3	48.0	64.0	96.0					350	Broke with coarse splinters.....	533
7.0	14.2	23.0	32.3	2.4	33.5	44.5	64.0	97.0					384	Slipped from bearings; buckled large splinter on corner.....	533

TABLE III.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Specific gravity of the air-dried specimen.	Direction of grain.	COEFFICIENT OF ELASTICITY.		Modulus of rupture.
								First deflection.	Second deflection.	
227. <i>Planera aquatica</i>	758	Florida	Chattahoochee.....	A. H. Curtiss	Rich, alluvial	0.5761		509	488	574
	758	do	do	do	do	0.5926		697	751	820
	918	do	do	C. Mohr	do	0.5113		407	415	469
228. <i>Celtis occidentalis</i>	75	Missouri	Allenton	G. W. Letterman	Low, rich	0.5887		610	665	712
<i>Sugarberry. Hackberry.</i>	75	do	do	do	Alluvial	0.6023		787	849	808
	806	Texas	Dallas	J. Reverchon	do	0.7239		555	552	738
	806	do	do	do	do	0.7558		478	528	755
	806	do	do	do	do	0.7706		751	697	846
	873	Massachusetts	Salem	J. Robinson	Loam	0.7727		542	564	771
	873	do	do	do	do	0.7920		626	610	762
	1111	Missouri	Saint Louis	Henry Eggert	Moist loam	0.7613		787	783	891
	1111	do	do	do	do	0.7154		976	939	820
228. <i>Celtis occidentalis</i> , var. <i>reticulata</i> ..	652	Arizona	Santa Rita mount-	G. Engelmann and	Dry	0.7920		669	651	586
<i>Hackberry. Palo Blanco.</i>	652	do	ains.	C. S. Sargent.	do	0.7882		976	1085	1024
229. <i>Ficus aurea</i>	486	Florida	Bay Biscayne	A. H. Curtiss	Coral	0.3215		317	257	278
	486	do	do	do	do	0.3061		222		199
231. <i>Ficus pedunculata</i>	508	do	Boca Chica Key	do	do	0.5085		407		230
<i>Wild Fig. India-rubber Tree.</i>										
232. <i>Morus rubra</i>	132	Missouri	Allenton	G. W. Letterman	Rich loam	0.6876		1039	1062	937
<i>Red Mulberry.</i>	132	do	do	do	do	0.6784		814	904	848
	1244	do	do	do	Upland	0.6516		939	888	766
	1245	do	do	do	do	0.6506		718	814	738
	1246	do	do	do	do	0.6312		697	723	696
	1255	do	do	do	Rich	0.6875		751	697	745
	1255	do	do	do	do	0.6646		634	678	696
234. <i>Maclura aurantiaca</i>	253	Texas	Dallas	J. Reverchon	Bottom	0.8011		857	930	1111
<i>Osage Orange. Bois d'Arc.</i>	253	do	do	do	do	0.7927		939	957	1150
PLATANACEÆ.										
235. <i>Platanus occidentalis</i>	21	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	0.5724		407	454	468
<i>Sycamore. Button Wood. But-</i>	126	Missouri	Allenton	G. W. Letterman	Rich, alluvial	0.6125		888	976	792
<i>ton-ball Tree. Water Beech.</i>	126	do	do	do	do	0.6295		1136	1190	642
236. <i>Platanus racemosa</i>	686	California	Carmel river	G. R. Vasey	Clay	0.5170		626	622	586
<i>Sycamore. Button Wood.</i>	686	do	do	do	do	0.4812		595	626	537
237. <i>Platanus Wrightii</i>	648	Arizona	Santa Rita mount-	G. Engelmann and	Rich, gravelly	0.5170		407	425	468
<i>Sycamore.</i>	648	do	ains.	C. S. Sargent.	do	0.5369		531	488	387
JUGLANDACEÆ.										
238. <i>Juglans cinerea</i>	16	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	0.4829		634	651	368
<i>Butternut. White Walnut.</i>	16	do	do	do	do	0.4579		783	787	668
	76	Missouri	Allenton	G. W. Letterman	Moist, alluvial	0.4318		814	888	687
	76	do	do	do	do	0.4375		976	1017	649
	76	do	do	do	Rich, moist up-	0.4943		1136	1221	696
	123	Michigan	Dansville	W. J. Beal	land. Gravelly clay	0.3864		697	697	560
	393	Michigan	Lansing	do	Gravelly loam	0.3205		488	424	328
	1057	Massachusetts	Topsfield	J. Robinson	Drift	0.5284		814	814	818
239. <i>Juglans nigra</i>	112	Missouri	Allenton	G. W. Letterman	Alluvial	0.5852		904	1039	1029
<i>Black Walnut.</i>	117	Michigan	Dansville	W. J. Beal	Gravelly	0.5652		904	976	766
	149	Illinois	Waukegan	Robert Douglas	Loam	0.6031		872	976	726

UNITED STATES UNDER TRANSVERSE STRAIN—Continued.

DEFLECTION, IN MILLIMETERS, UNDER A PRESSURE, IN KILOGRAMS, OF—													Ultimate strength: transverse pressure.	Remarks.	Office number.
50	100	150	200	0 (set.)	200	250	300	350	400	450	500	550			
9.6	20.0	33.5	48.5	6.0	49.6								245	Sap-wood; broke through knot.....	758
7.0	13.0	19.9	27.2	1.7	29.5	37.2	51.0	80.0					350	Specimen cross-grained.....	758
12.0	23.5	39.0											200	Broke with large splinters; shattered.....	918
8.0	14.7	23.0	33.0	3.0	34.4	45.5	78.0						304	Broke with coarse splinters.....	75
6.2	11.5	17.5	25.3	1.9	25.0	33.5	48.0						345	Failed by bending; fine splinters.....	75
8.8	17.7	27.5	40.5	4.5	41.5	56.0	82.0						315	Specimen cross-grained; splinters on corners at knots.....	306
10.2	18.5	29.5	42.7	5.5	44.5	60.0	87.0						322	Broke with large splinters.....	306
6.5	14.0	21.5	30.0	2.5	32.0	42.0	57.5	87.0					361	Broke with large scale.....	306
9.0	17.3	29.0	42.0	5.7	44.0	58.8	91.6						329	Square break on tension side with coarse splinters.....	873
7.8	16.0	25.2	37.6	4.8	39.0	53.0	82.5						325	Broke with coarse splinters.....	873
6.2	12.8	20.0	28.0	2.3	29.0	37.3	51.0	74.0					380	Shattered; large, coarse splinters.....	1111
5.0	10.4	16.0	21.7	1.1	22.3	30.0	38.2						350	Broke at knot; coarse splinters.....	1111
7.3	15.0	25.0	37.0	5.3	38.0								250	Specimen cross-grained; split with grain.....	652
5.0	9.0	14.8	20.0	1.5	20.2	26.5	35.4	50.0	70.5				437	Specimen cross-grained; broke with large splinters.....	652
15.4	38.0												119	Square break.....	486
22.0													85	do.....	486
12.0													98	Specimen cross-grained; split with grain.....	508
4.7	9.2	14.0	20.5	1.3	20.7	28.0	40.0	55.0					400	Broke with coarse splinters.....	132
6.0	10.8	16.4	23.0	1.0	23.5	31.0	43.0	61.0					362	do.....	132
5.2	11.0	15.8	22.6	2.0	23.5	32.3	46.0						327	Broke with thin flakes.....	1244
6.8	12.0	18.5	26.6	2.3	27.0	38.2	57.0						315	Failed from splinters on the corner.....	1245
7.0	13.5	20.0	29.0	2.6	30.3	42.0							297	Failed from thin scales on tension side.....	1246
6.5	14.0	21.4	32.4	2.6	32.2	43.2	70.0						318	Broke with large splinters.....	1255
7.7	14.4	22.2	32.5	3.0	33.5	44.3							298	do.....	1255
5.7	10.5	15.2	20.2	0.7	20.6	27.0	31.5	38.5	47.0	56.4			474	Broke with fine splinters.....	253
5.2	10.2	15.2	20.4	0.4	21.0	26.4	32.0	38.5	47.8	58.0			491	do.....	253
12.0	23.0	38.0	55.7	9.2									200	Specimen cross-grained; split with grain.....	21
5.5	10.0	14.5	19.6	1.0	20.8	25.0	33.0						338	Square break with coarse splinters.....	126
4.3	8.2	12.3	17.0	0.6	17.5	22.6							274	do.....	126
7.8	15.7	24.4	36.8	3.5	38.3	63.0							250	Broke with fine splinters.....	686
8.2	15.6	24.6	39.0	4.9	41.0								229	Broke with thin flakes from tension side.....	686
12.0	23.0	38.7											200	Specimen cross-grained; split with grain.....	648
9.1	20.0	34.2											165	Specimen cross-grained.....	648
7.7	15.0	23.0											157	Specimen cross-grained; broke at small knots.....	16
6.4	12.4	19.0	27.5	2.0	28.0	38.2							285	Square break on tension side; split in axis.....	16
6.0	11.0	16.8	22.8	0.9	24.0	31.7							293	do.....	76
5.0	9.6	14.5	20.3	0.6	21.0	28.4							277	Crushed at center bearing; broke with long splinters.....	76
4.3	8.0	12.4	17.0	0.9	18.0	24.3							297	Crushed at center bearing; broke with fine splinters.....	76
7.0	14.0	18.5	27.0	2.0	29.0								239	Square break.....	123
10.0	23.0												140	do.....	393
6.0	12.0	17.7	24.0	1.0	24.0	31.0	40.0						349	Square break on tension side, splitting in axis; shattered.....	1057
5.4	9.4	14.0	19.0	0.3	19.2	24.0	29.5	35.6	44.0				439	Specimen cross-grained; shattered.....	112
5.4	10.0	14.8	20.4	0.6	20.5	25.5	32.3						327	do.....	117
5.6	10.0	15.4	21.6	1.3	21.8	29.7	42.0						310	Specimen cross-grained; long break started at small knot.....	149

FOREST TREES OF NORTH AMERICA.

TABLE III.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Specific gravity of the air-dried specimen.	Direction of grain.	COEFFICIENT OF ELASTICITY.		Modulus of rupture.
								First deflection.	Second deflection.	
239. <i>Juglans nigra</i> —continued.	318	Michigan	Lansing	W. J. Beal	Loam	0.6619		976	976	586
	325	Texas	Dallas	J. Reverchon	Alluvial	0.6455		976	930	668
	407		Charlestown Navy-yard.	S. H. Pook		0.5981		976	1136	923
	430	Tennessee	Nashville	A. Gattinger	Limestone	0.5724		857	921	896
	766	Florida	Aspalaga	A. H. Curtiss	Clay	0.6933		1221	1221	820
	766	do	do	do	do	0.6949		1627	1627	1137
	951	Texas	New Braunfels	C. Mohr	Moist, calcareous	0.7483		1221	1191	923
	951	do	do	do	do	0.7386		939	1017	937
240. <i>Juglans rupestris</i> . <i>Walnut.</i>	415	New Mexico	Pinos Altos mountain.	E. L. Greene	Alluvial	0.6125		697	688	462
	415	do	do	do	do	0.7207		660	679	949
	672	California	Contra Costa county	G. R. Vasey	do	0.5977		775	842	468
	672	do	do	do	do	0.6063		626	697	520
241. <i>Carya olivaeformis</i> . <i>Pecan. Illinois Nut.</i>	322	Mississippi	Greenville	C. Mohr	do	0.7444		642	651	457
	322	do	do	do	do	0.7489		904	976	675
	326	Texas	Dallas	J. Reverchon	do	0.7181		470	496	581
	326	do	do	do	do	0.6250		542	542	600
242. <i>Carya alba</i> . <i>Shell-bark Hickory. Shag-bark Hickory.</i>	3	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	0.8295		1395	1356	1245
	3	do	do	do	do	0.7955		1320	1356	1249
	29*	Kentucky	Danville	W. M. Linney	Shale	0.9716		1163	1220	1266
	118*	Michigan	Hudson	W. J. Beal	Clay	0.8085		1356	1395	1057
	118*	do	Lansing	do	do	0.8035		1575	1550	1036
	152	Missouri	Allenton	G. W. Letterman	Rich upland	0.8217		1627	1627	1132
	249	Virginia	Wytheville	H. Shriver	Clay	0.8108		1221	1191	1144
	249	do	do	do	do	0.8041		1085	1163	1160
	249	do	do	do	do	0.8163		1221	1320	1242
	531	Mississippi	Kemper's mill	C. Mohr	Alluvial	0.8835		1320	1479	1289
	531	do	do	do	do	0.8829		1356	1395	1289
	531	do	do	do	do	0.8746		1526	1601	1444
	539	do	do	do	do	0.9148		1285	1526	1242
	539	do	do	do	do	0.7698		1479	1575	1237
	816	West Virginia	Grafton	C. G. Pringle	do	0.7569		1526	1627	1041
	816	do	do	do	do	0.7947		1628	1628	1216
	1056	Massachusetts	Topsfield	J. Robinson	Rich loam	0.8614		800	976	1137
	1056	do	do	do	do	0.8439		888	921	1073
	1097	Missouri	Allenton	G. W. Letterman	Alluvial	0.8261		1395	1502	1331
	1097	do	do	do	do	0.7522		1221	1356	1172
243. <i>Carya sulcata</i> . <i>Big Shell-bark. Bottom Shell-bark.</i>	91	Kentucky	Mercer county	W. M. Linney	do	0.7130		976	1062	1022
	91*	do	do	do	do	0.6603		729	781	755
	383	Missouri	Allenton	G. W. Letterman	do	0.9376		751	842	1223
	383	do	do	do	do	0.9333		660	729	1055
	383	do	do	do	do	0.9421		939	1085	1160
	391	do	do	do	do	0.8886		872	957	1172
	391	do	do	do	do	0.8648		939	1007	1116
	1082	do	do	do	do	0.9000		740	849	1087
	1082	do	do	do	do	0.8834		660	835	1083
	1164	do	do	do	do	0.7927		1221	1191	1144
	1164	do	do	do	do	0.8461		1356	1526	1352
	1164	do	do	do	do	0.7762		1221	1395	1287

UNITED STATES UNDER TRANSVERSE STRAIN—Continued.

DEFLECTION, IN MILLIMETERS, UNDER A PRESSURE, IN KILOGRAMS, OF—													Ultimate strength: transverse pressure.	Remarks.	Office number.
50	100	150	200	0 (sect.)	200	250	300	350	400	450	500	550			
5.0	10.0	15.0	21.6	0.6	21.7								250	Specimen cross-grained; shattered	318
5.0	10.5	16.0	21.9	1.0	21.7	28.0							285	Specimen cross-grained; split at corner	325
5.0	8.6	13.0	17.5	0.2	17.7	22.7	28.5	38.0					394	Crushed at center bearing; broke in long splinters	407
5.7	10.6	16.0	22.0	0.7	13.0	28.2	37.0	49.5					383	Square break; split from end to end	430
4.0	8.0	12.6	18.0	1.2	18.7	24.0	32.0	41.6					350	Square break on tension side; split in axis	766
3.0	6.0	9.8	13.2	0.2	13.5	17.0	21.0	26.0	32.0	40.0			485	 do	766
4.0	8.2	12.7	17.5	1.0	18.0	23.0	29.5	39.2					394	Specimen cross-grained	951
5.2	9.6	15.0	21.0	1.5	21.2	28.0	36.0	49.5					400	Shattered	951
7.0	14.2	23.0											197	Specimen cross-grained; broke at knot	415
7.4	14.4	21.5	29.0	2.0	29.4	38.5	50.0	68.5	112.0				405	Failed from splinters on corners	415
6.3	11.6	17.2	24.0	1.0									200	Specimen cross-grained; broke at knot	672
7.8	14.0	21.5	30.0	1.6	31.2								222	Cross-grained; broke with large splinters	672
7.6	15.0	23.0											195	Defective specimen; square break on tension side	322
5.4	10.0	15.7	21.4	0.9	21.6	28.0							288	Split; did not break	322
10.4	19.7	32.0	51.5	8.8	54.4								248	Specimen cross-grained; square break on tension side; split in axis	326
9.0	18.0	29.3	43.0	5.3	44.0	61.0							256	Square break on tension side; long splinters	326
3.5	7.2	11.4	16.0	0.4	16.2	20.8	27.3	34.5	43.0	58.0	67.0		531	Sap-wood; broke with fine splinters	3
3.7	7.2	10.8	14.4	0.4	14.5	18.2	23.5	30.0	37.0	50.7	75.0		533	 do	3
4.2	8.0	12.2	17.0	0.9	17.6	23.0	29.5	36.0	44.0	60.4	82.0		540	Broke with fine splinters	299
3.6	7.0	10.5	14.0	0.6	14.2	17.6	21.7	26.5	32.0	46.0			451	Broke with long flakes from tension side	1189
3.1	6.3	9.0	12.1	0.2	12.4	15.5	19.0	23.0	29.5				442	Broke with long, fine splinters	1189
3.0	6.0	9.2	12.5	0.2	12.8	16.0	19.5	24.0	29.0	37.0			483	Broke with large, coarse splinters	152
4.0	8.2	13.0	18.2	1.0	19.0	25.0	33.0	43.0	56.0	76.0			428	Sap-wood; broke with fine splinters	249
4.5	8.4	12.7	16.8	0.6	17.3	21.5	28.5	36.6	46.0	64.0			495	 do	249
4.0	7.4	11.2	15.3	0.8	15.4	19.2	25.7	33.5	45.0	57.0	84.0		530	 do	249
3.7	6.6	10.0	13.0	0.2	13.2	16.6	21.0	25.5	33.0	41.0	55.0	100.0	550	Crushed at center bearing; fine splinters	531
3.6	7.0	10.2	13.2	0.3	13.5	17.0	21.0	26.0	33.0	40.6	55.0		550	 do	531
3.2	6.1	9.0	12.0	0.3	12.2	15.3	19.0	23.5	28.4	35.0	44.5	50.0	616	Deflection of 88 millimeters under pressure of 600 kilograms; some crushing at center bearing	531
3.8	6.4	9.6	12.5	0.3	13.0	15.8	20.2	24.0	29.5	39.0	54.0		530	Crushed at center bearing; fine splinters	539
3.3	6.2	9.2	12.6	0.2	13.0	16.2	20.0	25.0	31.5	40.0	59.5		528	0.5 sap-wood; crushed at center bearing; fine splinters	539
3.2	6.0	9.0	12.0	0.2	12.5	16.0	19.7	24.5	30.5				444	Broke with long splinters	816
3.0	6.0	8.6	11.5	0.3	11.5	14.2	18.2	22.5	28.5	36.0	49.5		519	 do	816
6.1	10.0	14.6	20.0	0.9	20.4	26.2	33.0	45.0	58.0	81.0			465	Broke with fine splinters	1056
5.5	10.6	16.0	22.6	1.9	23.0	30.3	40.2	54.0	74.5	105.0			458	Sap-wood; broke with fine splinters	1056
3.5	6.5	9.6	12.5	0.3	13.0	16.0	20.2	25.0	30.7	38.0	49.5	71.2	568	 do	1097
4.0	7.2	10.5	14.2	0.5	14.4	18.0	22.5	28.9	37.0	48.0	70.0		500	0.5 sap-wood; broke with fine splinters	1097
5.0	9.2	14.0	18.5	0.4	19.0	24.2	30.8	41.0	55.6				436	Broke with two large splinters on tension side	91
6.7	12.5	18.5	26.2	1.4	27.0	36.0	50.0						322	Failed from thin flakes on back	919
6.5	11.0	17.6	25.0	2.0	26.1	33.6	45.2	60.0	85.0	132.0			479	Slipped from bearings; splintered	383
7.4	13.4	21.5	30.5	3.2	31.0	40.0	53.0	78.5	102.5	194.0			450	Sap-wood; slipped from bearings; did not break	383
5.2	9.0	13.2	18.3	1.0	19.1	25.0	31.4	41.4	54.5	75.0			495	Failed from thin flakes on back	383
5.6	10.2	15.0	20.5	1.1	21.0	28.0	36.0	46.0	59.0	91.0	185.0		500	Broke with fine splinters	391
5.2	9.7	14.7	20.0	1.0	20.6	26.5	34.0	43.0	61.5	84.0			476	0.5 sap-wood; broke with fine splinters	391
6.6	11.5	16.1	22.2	1.0	22.6	30.0	40.0	54.0	77.0	130.0			464	Did not break; pushed through the bearings	1082
7.4	11.7	16.0	21.3	1.1	22.0	27.5	36.0	47.0	63.0	95.0			462	Broke with fine splinters	1082
4.0	8.2	12.0	16.5	0.6	16.6	21.0	27.5	35.0	45.0	66.5			488	 do	1164
3.6	6.4	9.5	12.8	0.4	12.7	16.0	20.0	24.5	30.3	37.0	46.0	65.0	577	 do	1164
4.0	7.0	10.2	13.6	0.4	14.0	17.4	22.0	27.4	33.2	42.6	56.0		549	 do	1164

TABLE III.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Specific gravity of the air-dried specimen.	Direction of grain.	COEFFICIENT OF ELASTICITY.		Modulus of rupture.
								First deflection.	Second deflection.	
243. <i>Carya sulcata</i> —continued	1165	Massachusetts	North Reading	J. Robinson	Drift	0.7181		1221	1221	780
	1166	do	do	do	do	0.8881		939	1007	1085
	1170	do	do	do	do	0.7636		1221	1395	1020
244. <i>Carya tomentosa</i>	72	Kentucky	Perryville	W. M. Linney	Utica shale	0.7567		904	948	919
Mocker Nut. Black Hickory.	72	do	do	do	do	0.8058		976	1085	1041
Bull Nut. Big-bud Hickory.	254	Missouri	Allenton	G. W. Letterman	Rich upland	0.8693		1479	1628	1411
White-heart Hickory. King Nut.	348	Alabama	Citronelle	C. Mohr	Sandy	0.8659		814	939	1146
245. <i>Carya porcina</i>	6	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	0.8161		888	976	1054
Pig Nut. Brown Hickory. Black Hickory. Switch-bud Hickory.	6	do	do	do	do	0.7818		976	1062	1054
	88	Missouri	Allenton	G. W. Letterman	Rich loam	0.9658		1252	1395	1354
	88	do	do	do	do	0.9369		1039	1149	1171
	121 ¹	Michigan	Dansville	W. J. Beal	Gravelly clay	0.6659		1085	1221	876
	288	Missouri	Allenton	G. W. Letterman	Rich loam	0.8473		976	1085	1146
	442	Tennessee	Nashville	A. Gattinger	Upland	0.8452		751	814	806
	538	Alabama	Kemper's mill	C. Mohr	Alluvial	0.8183		1062	1221	1092
	538	do	do	do	do	0.6932		1062	1097	928
	538	do	do	do	do	0.7461		1085	1221	1110
	1051	Massachusetts	North Reading	J. Robinson	Drift	0.8636		525	734	1164
	1051	do	do	do	do	0.8750		996	1097	1108
	1098	Missouri	Allenton	G. W. Letterman	Flinty	0.8614		888	814	963
	1098	do	do	do	do	0.8113		718	751	935
	1168	do	do	do	do	0.8580		660	713	911
	1168	do	do	do	do	0.8500		800	880	1054
246. <i>Carya amara</i>	153	do	do	do	Rich upland	0.8153		1039	1085	1132
Bitter Nut. Swamp Hickory.	838	Massachusetts	Danvers	J. Robinson	Rich loam	0.7864		763	814	935
	838	do	do	do	do	0.8040		957	1085	1122
	838	do	do	do	do	0.8080		1085	1136	1214
247. <i>Carya myristiciformis</i>	237	South Carolina	Bonneau's Depot	H. W. Ravenel	Rich, swampy	0.8636		1221	1302	1296
Nutmeg Hickory.	237	do	do	do	do	0.8330		1479	1628	1492
248. <i>Carya aquatica</i>	129	do	do	do	Swampy	0.7386		1062	1163	1064
Water Hickory. Swamp Hickory. Bitter Pecan.	362	Mississippi	Vicksburg	C. Mohr	do	0.7546		651	697	921
	362	do	do	do	do	0.7727		976	948	937
	362	do	do	do	do	0.7813		976	976	1099
	740	Georgia	Bainbridge	A. H. Curtiss	Alluvial	0.8198		976	1085	787
	740	do	do	do	do	0.8470		1221	1375	1235
	740	do	do	do	do	0.8199		1085	1028	468
	917	Florida	Chattahoochee	C. Mohr	do	0.6415		729	848	562
MYRICACEÆ.										
249. <i>Myrica cerifera</i>	586	do	Saint John's river	A. H. Curtiss	Sandy loam	0.6227		814	888	955
Bayberry. Wax Myrtle.										
250. <i>Myrica Californica</i>	665	California	Santa Cruz	G. Engelmann and C. S. Sargent.	Rich loam	0.6511		904	976	1055
	665	do	do	do	do	0.6025		976	1007	1017
CUPULIFERÆ.										
251. <i>Quercus alba</i>	8	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	0.7432		679	720	782
White Oak.	8	do	do	do	do	0.8358		763	888	918
	82	Kentucky	Mercer county	W. M. Linney	Limestone	0.8006		1136	1136	1028
	82 ²	do	Boyle county	do	Shale	0.7068		814	930	911
	82 ³	do	do	do	Slate	0.6534		904	948	818

UNITED STATES UNDER TRANSVERSE STRAIN—Continued.

DEFLECTION, IN MILLIMETERS, UNDER A PRESSURE, IN KILOGRAMS, OF—													Ultimate strength: transverse pressure.	Remarks.	Office number.
50	100	150	200	0 (set.)	200	250	300	350	400	450	500	550			
4.0	8.0	11.5	15.6	0.5	16.0	20.0	25.0						333	Broke with coarse splinters.....	1165
5.2	9.7	15.0	20.8	1.3	21.0	28.0	37.2	50.0	69.5	137.0			463	Fine splinters; not broken.....	1166
4.0	7.0	10.5	13.5	0.5	14.0	18.0	24.0	34.0	50.0				435	Broke with fine splinters.....	1170
5.4	10.3	15.3	22.0	1.4	23.0	29.6	41.5	58.0					392	Sap-wood; failed with two thin flakes on back.....	72
5.0	9.0	13.8	19.6	1.1	20.0	26.0	33.0	44.0	62.0				444	Sap-wood; broke with fine splinters.....	72
3.3	6.0	9.0	12.2	0.3	12.2	15.0	19.1	22.7	27.4	33.0	40.0	50.0	602	Broke with coarse splinters.....	254
6.0	10.4	15.6	21.4	1.1	22.0	28.0	37.1	47.0	61.0	89.0			489	Sap-wood; broke with fine splinters.....	848
5.5	10.0	14.5	20.4	1.0	20.5	27.0	34.0	44.0	57.5	79.7			450	0.5 sap-wood; broke with fine splinters.....	6
5.0	9.2	14.2	20.0	1.1	20.7	26.2	35.0	44.2	60.0				450	do.....	6
3.9	7.0	10.9	14.2	0.5	15.0	18.5	23.6	31.0	36.0	46.0	60.2	76.0	578	Broke with fine splinters.....	88
4.7	8.5	12.5	17.5	1.1	18.0	22.5	30.3	37.5	49.0	66.0			500	do.....	88
4.5	8.0	12.0	16.2	0.6	16.6	21.3	28.0	38.2					374	do.....	121 ¹
5.0	9.0	13.7	18.0	0.9	19.0	23.6	30.6	40.0	52.0	65.0			489	Broke with fine splinters.....	288
6.5	12.0	19.0	26.7	2.0	28.0	36.0	47.0						344	Broke at knot with coarse splinters.....	442
4.6	8.0	11.9	16.0	0.6	16.0	21.0	27.0	35.0	44.0	60.0			466	Broke with fine splinters.....	588
4.6	8.9	12.7	17.1	0.6	17.0	23.0	30.0	39.5					396	do.....	538
4.5	8.0	11.6	15.7	0.9	16.4	21.2	28.6	36.4	49.0	68.0			474	Square break on tension side; split in axis.....	588
9.3	12.3	18.0	22.6	6.0	23.3	28.4	36.0	46.5	61.0	84.0			497	Broke with fine splinters.....	1051
4.9	8.9	13.4	18.5	1.3	19.0	24.5	31.0	42.5	56.0	75.5			473	do.....	1051
5.5	12.0	18.8	27.5	3.0	28.5	37.6	52.5	70.0	110.0				411	Broke with coarse splinters; flaked on tension side.....	1098
6.8	13.0	19.2	28.0	2.6	28.8	37.5	51.6	73.0					399	do.....	1098
7.4	13.7	21.0	31.5	5.0	32.5	41.4	57.0	85.0					389	Broke with coarse splinters.....	1168
6.1	11.1	16.1	23.0	2.2	24.2	31.6	41.0	55.0	73.0	105.0			450	do.....	1168
4.7	9.0	13.3	19.0	1.0	18.4	24.0	21.3	41.5	56.4	84.0			483	Broke with long splinters.....	153
6.4	12.0	18.5	26.3	2.2	27.5	37.0	50.5	70.5					399	Broke with fine splinters.....	838
5.1	9.0	14.0	19.0	1.1	19.0	24.7	32.5	42.5	56.7	74.5			479	do.....	838
4.5	8.6	13.7	18.4	0.6	18.7	24.0	30.6	39.0	51.0	68.5	103.0		518	Broke with long splinters.....	838
4.0	7.5	11.2	14.6	0.3	15.0	19.0	23.7	30.0	36.0	45.0	56.5	77.0	553	Broke with fine splinters.....	237
3.3	6.0	9.0	11.6	0.1	12.0	14.7	18.1	22.0	26.0	31.2	38.5	48.0	637	Deflection 63 millimeters with a pressure of 600 kilograms; broke with fine splinters.....	237
4.6	8.4	12.0	16.5	1.0	16.5	22.0	29.0	37.5	52.5	72.0			454	Shattered.....	129
7.5	14.0	21.0	30.5	3.0	31.2	41.2	57.2	86.0					393	0.5 sap-wood; failed from flakes on back.....	362
5.0	10.3	15.2	21.5	1.2	22.0	30.0	42.0	56.5	82.0				400	Broke with long splinters.....	362
5.0	10.0	14.7	20.2	1.1	21.0	27.5	36.5	49.0	68.0	109.0			469	Broke with coarse splinters.....	362
5.0	9.0	13.5	18.0	0.7	18.2	23.5	30.2						336	Failed from long splinter on corner.....	740
4.0	7.1	10.8	14.4	0.4	15.0	18.0	22.2	27.0	33.0	40.0	47.2		527	Broke with coarse splinters.....	740
4.5	9.5	16.5	26.5	4.0									200	Specimen cross-grained.....	740
6.7	11.8	17.0	23.0	0.9	23.6								240	Square break on tension side, splitting in axis.....	917
6.0	11.0	17.4	24.0	1.3	24.0	31.5	42.0						348	Shattered.....	586
5.4	10.0	14.4	19.9	1.0	20.0	25.2	32.0	40.0	50.0				450	Square break on tension side; split in axis; shattered.....	665
5.0	9.7	14.7	20.0	1.0	20.4	26.0	33.0	42.4	60.0				434	do.....	665
7.2	13.5	21.2	30.7	3.1	32.0	42.0	59.0						334	Square break on tension side; split in axis.....	8
6.4	11.0	16.7	23.5	1.5	24.0	31.0	40.0	52.0	70.0				392	Broke with large splinters.....	8
4.3	8.6	13.0	17.6	0.4	18.0	23.0	30.5	38.6	48.5				439	do.....	32
6.0	10.5	15.0	20.7	1.0	21.5	28.2	37.0	47.5					389	Broke with small splinters.....	329
5.4	10.3	15.5	21.5	0.7	21.6	28.6	37.0						349	Square break on tension side, splitting in axis.....	329

TABLE III.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Specific gravity of the air-dried specimen.	Direction of grain.	COEFFICIENT OF ELASTICITY.		Modulus of rupture.
								First deflection.	Second deflection.	
251. <i>Quercus alba</i> —continued	49	Missouri	Allenton	G. W. Letterman	Rich upland	0.7927		976	1028	1024
	49	do	do	do	do	0.7060		740	751	817
	49 ¹	do	do	do	do	0.7784		872	930	937
	49 ²	do	do	do	do	0.7501		814	814	813
	49 ³	do	do	do	do	0.7614		872	913	921
	113	Michigan	Big Rapids	W. J. Beal	Gravelly	0.6001		679	697	684
	113 ²	do	Dansville	do	Sandy	0.7149		814	896	696
	113 ³	do	do	do	do	0.7060		857	775	585
	113 ⁴	do	Hudson	do	do	0.7460		976	976	977
	238	South Carolina	Bonneau's Depot	H. W. Ravenel	Rich, damp loam	0.8085		1356	1356	1055
	238	do	do	do	do	0.7797		1221	1221	1055
	250	Virginia	Wytheville	H. Shriver	Clay	0.7876		1163	1221	1047
	250	do	do	do	do	0.7796		1061	1191	928
	251	do	do	do	do	0.7102		814	842	764
	251	do	do	do	do	0.7170		1163	1221	1059
	259 ¹	do	do	do	Gravelly	0.8091		976	976	1043
	259 ²	do	do	do	do	0.7631		872	921	844
	408	Maryland	Charlestown Navy-yard.	S. H. Pook	do	0.7364		976	976	722
	403	do	do	do	do	0.7143		842	849	653
	443	Tennessee	Nashville	A. Gattinger	Rich bottom	0.6630		567	592	487
	547	Alabama	Kemper's mill	C. Mohr	Alluvial	0.8475		996	1017	1031
	547	do	do	do	do	0.8579		976	1177	1282
	748	Florida	Chattahoochee	A. H. Curtiss	Clay	0.8075		1221	1252	1125
	748	do	do	do	do	0.8099		1320	1302	1153
	749	do	do	do	do	0.8233		751	775	1043
	749	do	do	do	do	0.8655		814	781	750
	895	Massachusetts	do	M. C. Beedle	do	0.7306		740	842	703
	895	do	do	do	do	0.7818		697	723	703
	1050	do	North Reading	J. Robinson	do	0.8001		1110	1191	1158
	1050	do	do	do	do	0.8343		787	751	1040
	1257	do	Charlestown Navy-yard.	S. H. Pook	do	0.8466		1252	1268	1005
	1257	do	do	do	do	0.8773		1061	1085	919
252. <i>Quercus lobata</i> <i>White Oak. Weeping Oak.</i>	670	California	Redding	G. R. Vasey	Gravelly loam	0.7443		679	665	876
	670	do	do	do	do	0.7557		763	769	851
253. <i>Quercus Garryana</i> <i>White Oak.</i>	985	Oregon	Weidler's saw-mill	G. Engelmann and C. S. Sargent	do	0.7864		904	888	1048
	965	do	do	do	do	0.7556		814	842	945
	988	do	Portland	do	Rich loam	0.7142		751	787	919
	988	do	do	do	do	0.7548		814	857	1022
	1027	do	Portland Furniture Company.	do	do	0.7745		697	814	820
	1027	do	do	do	do	0.7535		610	673	703
	1029	do	do	do	do	0.7420		888	930	921
	1029	do	do	do	do	0.6753		697	697	659
254. <i>Quercus obtusiloba</i> <i>Post Oak. Iron Oak.</i>	37 ²	Kentucky	Harrodsburg	W. M. Linney	Shale	0.8795		763	751	1055
	151	South Carolina	Bonneau's Depot	H. W. Ravenel	Rich upland	0.8085		763	849	1005
	151	do	do	do	do	0.8913		976	1017	1048
	256	Missouri	Allenton	G. W. Letterman	Clay	0.8943		763	781	912
	351	Alabama	Citronelle	C. Mohr	do	0.7614		610	630	677
	351	do	do	do	do	0.7688		751	888	893

UNITED STATES UNDER TRANSVERSE STRAIN—Continued.

DEFLECTION, IN MILLIMETERS, UNDER A PRESSURE, IN KILOGRAMS, OF—													Ultimate strength: transverse pressure.	Remarks.	Office number.
50	100	150	200	0 (set.)	200	250	300	350	400	450	500	550			
5.0	9.5	14.4	20.0	1.0	20.0	26.2	34.0	43.5	57.5				437	Broke with fine splinters	49
6.6	13.0	20.4	30.0	8.2	30.5	41.0	57.5						349	Square break on tension side, splitting in axis	49
5.6	10.5	17.0	24.0	2.5	25.4	33.5	45.0	59.5	89.0				400	do	49 ¹
6.0	12.0	19.0	28.0	3.3	28.7	37.0	48.0						347	do	49 ²
5.6	10.7	16.5	23.4	1.8	24.0	31.0	42.5	55.0					393	do	49
7.2	14.0	23.0	33.2	4.0	34.5	49.0							293	do	113
6.0	10.9	16.2	22.7	1.5	23.3	32.5							294	Broke with small splinters; slightly crushed at bearing	113 ²
5.7	12.6	20.0	30.8	3.0	31.5								250	Broke with fine splinters	113 ²
5.0	10.0	16.5	24.2	2.5	25.0	32.0	42.0	58.0	85.0				417	do	113 ²
3.6	7.2	11.0	15.3	0.5	15.4	20.5	26.5	33.0	45.5				450	do	238
4.0	8.0	12.0	16.0	0.5	16.3	21.0	27.5	35.0	46.5	63.0			450	do	238
4.2	8.0	12.0	16.3	0.6	16.6	21.2	28.5	36.0	49.2				447	do	250
4.6	8.2	12.7	17.5	0.7	17.7	22.5	30.0	36.5					396	Broke with large flakes	250
6.0	11.6	18.0	26.0	2.2	26.5	36.3	55.0						326	Broke with fine splinters	251
4.2	8.0	12.0	16.0	0.3	16.1	21.0	27.0	33.5	44.5	61.5			452	Broke with large splinters	251
5.0	10.0	16.0	22.8	1.5	24.0	30.5	39.5	53.5	70.0				445	Broke with fine splinters	259 ¹
5.6	10.6	16.3	23.2	1.5	24.0	32.0	42.2	59.5					360	Broke with large scale	259 ²
5.0	10.0	15.3	21.8	1.5	22.2	29.5	40.8						308	Broke with fine splinters	403
5.8	11.5	17.8	25.3	2.2	27.0	36.5							279	Broke with large scale	403
8.6	16.5	26.0	41.6	5.6	43.5								208	Square break on tension side, splitting in axis	443
4.9	9.6	14.5	20.7	1.3	21.8	28.5	37.5	49.5	69.0				440	Broke with fine splinters	547
5.0	8.3	12.2	16.0	0.6	16.2	20.7	26.0	32.5	40.7	48.7	67.5		547	do	547
4.0	7.8	11.6	16.0	0.5	16.3	20.5	27.0	33.5	43.0	56.0			480	do	748
3.7	7.5	11.0	15.0	0.5	15.5	19.6	25.0	32.0	41.5	51.2			492	do	748
6.5	12.6	18.8	25.5	2.2	27.5	34.5	45.5	65.0	92.0				445	do	749
6.0	12.5	20.0	29.0	3.0	30.0	42.0	73.0						320	Broke at knot	749
6.6	11.8	17.5	24.7	1.0	25.0	33.5							300	Specimen cross-grained	895
7.0	13.5	20.6	29.0	2.0	29.6	39.0							300	Broke with fine splinters	895
4.4	8.2	12.7	17.1	0.7	17.5	22.3	28.5	37.0	47.0	66.0			494	do	1050
6.2	13.0	19.5	28.0	2.4	28.5	36.0	50.0	68.5	100.0				444	do	1050
3.9	7.7	11.7	16.0	0.4	16.5	21.0	26.5	33.5	40.5				420	Broke with large splinters	1257
4.0	9.0	13.3	18.6	0.9	18.7	24.0	32.0	38.0					392	do	1257
7.2	14.7	23.5	33.0	4.0	34.0	45.6	62.0	90.0					374	Specimen cross-grained; splintered on both corners	670
6.4	12.7	18.7	27.3	2.5	28.4	37.0	50.5	75.0					363	Splintered on corner	670
5.4	11.0	15.5	21.5	1.1	21.5	28.5	37.0	47.0	60.5				447	Square break on tension side, splitting in axis	985
6.0	11.6	18.0	25.0	1.7	26.0	33.0	44.0	57.5	75.5				403	Specimen cross-grained	985
6.5	12.4	19.0	27.0	2.0	27.3	34.7	45.6	62.0					392	Square break on tension side, splitting in axis	988
6.0	11.4	16.7	23.0	1.2	23.5	30.0	40.0	52.5	70.0				436	do	988
7.0	12.0	18.0	25.5	1.5	25.6	33.5	44.0						350	do	1027
8.0	14.5	23.0	32.3	2.5	33.5	43.5	59.5						300	do	1027
5.5	10.5	16.0	22.3	1.0	22.3	29.0	36.5	49.0					393	Specimen cross-grained; broke with long splinters	1029
7.0	14.0	22.0	31.0	2.2	31.0	40.7							281	do	1029
6.4	13.0	19.6	27.5	2.0	28.0	36.0	46.7	62.0	76.5	105.0			450	Square break on tension side, splitting in axis	37 ²
6.4	11.5	17.5	24.0	1.1	24.6	32.0	40.0	50.6	67.5				429	do	151
5.0	9.6	14.5	19.5	1.0	19.9	26.0	33.5	43.0	60.0				447	Broke with coarse splinters	151
6.4	12.5	19.6	28.2	3.0	29.2	38.0	50.8	64.0					389	Square break on tension side, splitting in axis	256
8.0	15.5	23.0	32.5	2.7	33.7	43.5							289	do	351
6.5	11.0	17.7	24.7	1.7	25.0	32.0	43.0	56.0					361	do	351

TABLE III.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Specific gravity of the air-dried specimen.	Direction of grain.	COEFFICIENT OF ELASTICITY.		Modulus of rupture.
								First deflection.	Second deflection.	
254. <i>Quercus obtusiloba</i> —continued.	771	Florida	Aspalaga	A. H. Curtiss	Gravelly barrens	0.9172		888	930	809
	771	do	do	do	do	0.8958		814	814	581
255. <i>Quercus undulata</i> , var. <i>Gambellii</i> . <i>Scrub Oak.</i>	417	New Mexico	Pinos Altos mountains.	E. L. Greene		0.8489		610	610	820
	417	do	do	do		0.8092		751	718	998
	525	Colorado	Engelmann's cañon.	Robert Douglas	Rocky	0.7996		448	430	323
	525	do	do	do	do	0.8073		514	528	579
256. <i>Quercus macrocarpa</i> . <i>Burr Oak. Mossy-cup Oak. Over-cup Oak.</i>	79	Kentucky	Mercer county	W. M. Linney	Alluvial	0.7636		1221	1221	1055
	79 ^a	do	do	do	do	0.7358		1085	1163	1092
	137	Missouri	Allenton	G. W. Letterman	Moist upland	0.9418		872	948	1226
	143	Illinois	Waukegan	Robert Douglas	Rich	0.8368		679	679	696
	310	Texas	Dallas	J. Beverchon	Rich, moist	0.8580		1017	1028	968
	310	do	do	do	do	0.8466		939	967	916
	482	Tennessee	Nashville	A. Gattinger	Alluvial	0.7507		976	980	926
	831	Illinois	Winnebago county	M. S. Bebb	Loam	0.7455		1039	1149	1055
	933	Texas	Austin	C. Mohr	Alluvial	0.7864		697	723	914
	933	do	do	do	do	0.8313		729	712	803
	1071	Vermont	Charlotte	C. G. Pringle		0.8523		814	835	1052
	1072	do	do	do		0.8153		904	913	1029
257. <i>Quercus lyrata</i> . <i>Over-cup Oak. Swamp Post Oak. Water White Oak.</i>	545	Mississippi	Kemper's mill	C. Mohr	Alluvial	0.7784		1039	1073	865
	545	do	do	do	do	0.7973		1221	1221	1146
	545	do	do	do	do	0.7898		1285	1302	1041
	762	Florida	Chattahoochee	A. H. Curtiss	do	0.7955		1627	1550	1020
	762	do	do	do	do	0.7955		1744	1526	1055
258. <i>Quercus bicolor</i> . <i>Swamp White Oak.</i>	54	Missouri	Allenton	G. W. Letterman	Alluvial	0.9102		763	835	1106
	54	do	do	do	do	0.7182		976	1085	998
	54 ^a	do	do	do	do	0.7443		976	1085	937
	54 ^a	do	do	do	do	0.7580		1062	1149	1036
	846	Massachusetts	West Newbury	J. Robinson	Low, swampy	0.8752		543	592	581
259. <i>Quercus Michauxii</i> . <i>Basket Oak. Cow Oak.</i>	846	do	Arnold Arboretum	C. S. Sargent	Drift	0.9689		651	693	792
	240	South Carolina	Bonneau's Depot	H. W. Ravenel	Alluvial	0.8432		1085	1085	1064
	240	do	do	do	do	0.8687		957	939	1090
	524	Alabama	Kemper's mill	C. Mohr	do	0.7898		976	930	1116
	524	do	do	do	do	0.7710		1110	1112	1144
	755	Florida	Chattahoochee	A. H. Curtiss	do	0.9199		828	849	1153
260. <i>Quercus prinus</i> . <i>Chestnut Oak. Rock Chestnut Oak.</i>	755	do	do	do	do	0.9289		888	857	1127
	81	Kentucky	Boyle county	W. M. Linney	Shale	0.7313		1221	1252	1219
	81 ^a	do	do	do	do	0.7528		1039	1149	930
	85	do	do	do	Limestone	0.8352		1628	1684	1305
	434	Tennessee	Nashville	A. Gattinger	Rocky upland	0.6552		581	595	546
	925	Alabama	Cullman	C. Mohr	Dry, rocky	0.8207		1628	1628	1130
261. <i>Quercus prinoides</i> . <i>Yellow Oak. Chestnut Oak. Ohin-quapin Oak.</i>	925	do	do	do	do	0.7523		1221	1221	1050
	34	Kentucky	Mercer county	W. M. Linney	Limestone	0.7983		976	976	1146
	34 ^a	do	Boyle county	do	Waverly shale	0.8949		1479	1526	1455
	273	Missouri	Allenton	G. W. Letterman	Limestone	0.9864		976	1050	1291
	287	do	do	do	Flinty	0.9125		1221	1221	1277

UNITED STATES UNDER TRANSVERSE STRAIN—Continued.

DEFLECTION, IN MILLIMETERS, UNDER A PRESSURE, IN KILOGRAMS, OF—													Ultimate strength: transverse pressure.	Remarks.	Office number.
50	100	150	200	0 (sect.)	200	250	300	350	400	450	500	550			
5.5	10.5	15.7	21.0	1.5	21.7	28.0	36.0						345	Splintered on corner	771
6.0	12.0	18.7	26.4	2.5	27.0								248	Square break	771
8.0	16.0	25.3	36.0	4.0	37.0	48.0	66.0						350	Broke with coarse splinters	417
6.5	13.6	21.7	31.0	2.7	31.9	41.5	58.0	84.0	145.0				426	Sap-wood; drew off bearings; fine splinters	417
11.0	22.7												138	Specimen rotten; square break	528
9.5	18.5	30.3	43.5	3.2	45.0								247	Split at one end; square break on tension side	528
4.0	8.0	13.0	17.0	0.3	17.0	22.0	29.0	37.0	47.0	64.0			450	Broke with long splinter on corner	79
4.5	8.4	12.6	17.2	0.5	17.5	22.5	29.7	38.0	48.5	64.0			406	Broke with fine splinters	79 ^a
5.6	10.3	15.4	21.5	1.5	22.0	28.0	36.5	46.0	60.0	79.0	110.0		523	Broke with large splinter on corner	137
7.2	14.4	22.4	32.0	3.4	33.0	44.0							297	Specimen cross-grained; started at knot	143
4.8	9.5	14.0	19.0	1.0	20.0	25.7	33.5	41.5	54.3				413	Specimen cross-grained; split	310
5.2	10.1	15.3	21.6	1.5	21.6	28.2	38.0	49.6					392	Splintered at corners	310
5.0	10.5	15.5	21.4	1.4	21.2	29.0	39.5	54.3					395	Typical	432
4.7	8.5	13.4	17.6	0.6	17.9	23.0	30.5	40.0	54.5	73.0			450	Broke with fine splinters	831
7.0	13.5	20.6	29.0	3.3	29.7	40.2	56.0	73.2					390	do	933
6.7	13.7	21.3	31.4	3.3	33.0	45.0	61.0						343	Square break on tension side	933
6.0	11.7	18.6	25.5	2.5	26.8	34.0	46.0	62.0	79.0				449	Failed from small splinters on corner	1071
5.4	10.7	17.0	23.4	2.0	23.6	30.8	40.5	54.6	73.5				439	Failed from thin flakes on back	1072
6.4	12.0	18.2	26.1	2.5	26.3	34.4	47.0	62.5	88.5				441	Failed from fine splinters on corner	1073
4.7	9.1	13.4	18.0	0.8	19.2	24.0	31.0	41.0					369	Square break; crushed	545
4.0	8.0	12.0	16.0	0.5	16.1	20.4	27.0	33.0	43.0	56.0			489	Broke with fine splinters	545
3.8	7.5	11.7	16.0	0.5	16.0	20.5	26.0	34.0	44.5				444	Failed from long splinters on tension side	545
3.0	6.3	9.5	12.5	0.4	12.9	16.0	21.0	26.2	32.7				435	Broke at knot in center of stick	762
2.8	6.4	9.4	12.5	0.5	12.5	16.0	20.0	25.0	31.4	39.0			450	Square break on tension side	762
6.4	11.7	18.2	25.7	1.4	26.0	33.5	45.0	58.0	78.0	109.5			472	Broke with small splinters; drew off bearing	54
5.0	9.0	14.0	19.0	1.0	19.5	26.0	34.5	45.0	62.0				426	Broke with long splinters	54
5.0	9.0	13.7	19.2	1.0	19.0	25.3	33.0	42.5	57.5				400	Square break on tension side, splitting in axis	54 ^a
4.6	8.5	12.7	17.0	0.6	17.2	22.5	29.5	35.5	47.5				442	Broke with fine splinters	54 ^a
9.0	16.5	24.5	35.0	3.7	36.0								248	Broke at knot near the end	846
7.5	14.1	22.0	31.6	3.8	32.1	41.5	59.0						338	Specimen cross-grained; split with grain	846
4.5	9.0	13.5	18.5	0.9	19.1	24.2	31.6	40.0	52.2	73.0			454	Broke with fine splinters	240
5.1	10.4	15.4	21.0	1.0	21.6	27.6	37.0	49.5	64.5	90.0			469	do	240
5.0	10.5	15.6	21.5	1.1	22.3	28.0	37.0	48.5	65.0	87.0			476	do	524
4.4	8.7	14.0	18.7	1.0	19.0	24.0	31.5	42.5	56.0	75.5			488	do	524
5.9	11.5	17.5	24.5	1.8	25.5	32.3	43.7	54.0	76.0	99.0			492	Failed from large splinter, starting at knot	755
5.5	11.4	17.2	24.0	2.0	24.5	32.0	42.5	58.0	76.0	135.0			481	Did not break; drew from bearing	755
4.0	7.8	11.4	14.7	0.3	15.0	19.0	24.4	30.0	37.5	47.5	61.5		520	Broke with fine splinters	31
4.7	8.5	13.0	17.3	0.5	17.7	21.5	30.0	36.7					397	Square break on tension side, splitting in axis	31 ¹
3.0	5.8	8.5	11.0	0.3	11.6	14.0	17.5	21.5	26.2	31.6	40.0	55.5	557	Crushed at bearing; broke with fine splinters	35
3.4	6.4	9.2	12.5	0.4	12.7	16.0	21.0	27.5	34.5	44.0			233	Square break on tension side, splitting in axis	434
3.0	6.0	9.0	12.6	0.6	12.7	16.0	22.0	27.5	34.5	44.0			482	Square break with large splinters	925
4.0	8.0	12.0	16.0	0.6	16.6	20.7	26.0	33.3	41.0				448	Failed from large splinter on corner	925
5.0	10.0	14.6	20.4	1.4	20.8	26.8	35.0	46.3	58.0	78.5			489	Broke with fine splinters	34
3.3	6.4	10.0	13.0	0.3	13.3	16.5	20.6	25.0	30.0	36.2	45.0	54.0	621	Deflection with a pressure of 600 kilograms, 68 millimeters; broke with fine splinters	34 ^a
4.0	9.3	14.0	19.0	1.0	19.3	25.0	32.5	42.0	51.5	66.5	92.5	135.0	551	Broke with large splinters	273
4.0	8.0	12.0	16.3	0.6	16.4	21.5	27.5	35.0	43.0	55.5	79.0		545	Broke with fine splinters	287

TABLE III.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Specific gravity of the air-dried specimen.	Direction of grain.	COEFFICIENT OF ELASTICITY.		Modulus of rupture.
								First deflection.	Second deflection.	
261. <i>Quercus prinoides</i> —continued.	323	Texas	Dallas	J. Reverchon	Calcareous	0.9205		976	1007	1230
	323	do	do	do	do	0.9966		976	1007	1090
	514	Tennessee	Nashville	A. Gattinger	Alluvial	0.9059		1065	1149	1289
	514	do	do	do	do	0.8755		976	1062	1120
262. <i>Quercus Douglasii</i> . <i>Mountain White Oak. Blue Oak.</i>	688	California	Contra Costa county	G. R. Vasey	Clay	1.0023		888	921	1048
	688	do	do	do	do	1.0284		610	622	937
263. <i>Quercus oblongifolia</i> . <i>White Oak.</i>	655	do	San Diego county	do	Dry, gravelly	1.1408		697	697	968
	655	do	do	do	do	1.0474		1017	1017	469
264. <i>Quercus grisea</i> . <i>White Oak.</i>	698	Arizona	Santa Rita mountains	G. Engelmann and C. S. Sargent.	Dry, rocky	0.9821		718	740	937
266. <i>Quercus Durandii</i>	1103	Texas	Austin	S. B. Buckley	Damp, calcareous	1.0023		888	888	996
	1103	do	do	do	do	1.0420		800	787	987
267. <i>Quercus virens</i> . <i>Live Oak.</i>	404	Florida	Charlestown Navy-yard.	S. H. Pook		1.0469		1436	1502	1055
	799	do	Saint John's river	A. H. Curtiss	Sandy	0.9127		1285	1395	1322
	799	do	do	do	do	0.9307		1221	1252	1212
	919	Alabama	Mobile county	C. Mohr	Rich, sandy	0.9114		1221	1191	1010
	919	do	do	do	do	0.8972		1163	1122	1054
	954	Texas	Matagorda bay	C. Mohr	Sandy loam	1.0114		697	740	816
	954	do	do	do	do	1.0193		751	751	649
268. <i>Quercus chrysolepis</i> . <i>Live Oak. Maui Oak. Valparaiso Oak.</i>	649	California	San Bernardino	W. G. Wright		0.8835		976	1065	1308
	649	do	do	do		0.8989		1039	1149	1268
	653	do	Marin county	G. R. Vasey	Gravelly	0.9386		1221	1221	1284
	653	do	do	do	do	0.9204		1285	1338	1212
269. <i>Quercus Emoryi</i> . <i>Black Oak.</i>	654	Arizona	Santa Rita mountains	G. Engelmann and C. S. Sargent.	Dry, rocky	1.0264		642	638	703
270. <i>Quercus agrifolia</i> . <i>Enceno. Coast Live Oak.</i>	663	California	Marin county	G. R. Vasey	Loam	0.8602		1061	957	937
	663	do	do	do	do	0.8508		976	948	930
271. <i>Quercus Wislizeni</i> . <i>Live Oak.</i>	685	do	Auburn	G. Engelmann		0.8676		904	849	759
	685	do	do	do		0.8653		857	872	877
271. <i>Quercus rubra</i> . <i>Red Oak. Black Oak.</i>	7	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	0.7011		1953	1627	1282
	7	do	do	do	do	0.6872		1285	1356	1171
	7	do	do	do	do	0.6364		1136	1149	1048
	7	do	do	do	do	0.6500		1356	1320	1057
	45 ¹	Kentucky	Mercer county	W. M. Linney	Shale	0.5528		976	996	855
	45 ²	do	do	do	do	0.6159		976	1007	902
	45 ³	do	do	do	do	0.5432		857	1302	787
	92	Kentucky	Mercer county	W. M. Linney	Alluvial	0.6261		814	814	787
	140	Michigan	Dansville	W. J. Beal	Sandy	0.6787		1221	1252	919
	141	do	do	do	do	0.5987		763	769	780
	146	Illinois	Waukegan	Robert Douglas	Gravelly	0.7464		1221	1221	895
	215	Vermont	Charlotte	C. G. Pringle	do	0.7293		1356	1356	1057
	215	do	do	do	do	0.6763		976	1028	877
	217	do	do	do	do	0.7316		888	888	1071
	218	do	do	do	do	0.7254		1221	1302	1242
	218	do	do	do	do	0.6955		1221	1268	1173
	920	Mississippi	do	C. Mohr	Alluvial	0.5841		857	814	680
	920	do	do	do	do	0.5852		976	921	762

UNITED STATES UNDER TRANSVERSE STRAIN—Continued.

DEFLECTION, IN MILLIMETERS, UNDER A PRESSURE, IN KILOGRAMS, OF—													Ultimate strength: transverse pressure.	Remarks.	Office number.
50	100	150	200	0 (set.)	200	250	300	350	400	450	500	550			
5.0	9.7	15.4	20.0	1.1	20.5	26.5	33.5	43.0	54.0	68.0	91.0		525	Broke with long splinters.....	323
5.0	9.7	15.3	21.0	1.5	21.5	28.0	36.0	45.5	59.5	93.0			465	do.....	323
4.5	8.5	12.7	17.3	1.1	17.5	22.3	29.0	36.0	45.0	56.0	71.6	93.0	550	Broke with fine splinters.....	514
5.0	9.2	14.0	19.0	1.0	19.2	25.0	32.0	41.0	51.5	64.0			478	Shattered; cross-grained.....	514
5.5	10.6	16.0	22.0	1.0	22.0	28.0	36.0	46.0	56.0				447	Specimen cross-grained; split with grain.....	688
6.0	15.7	23.2	32.3	2.3	33.0	42.5	55.3	72.0					400	Specimen cross-grained; started at knot.....	688
7.0	14.0	21.0	30.0	2.5	31.0	40.0	52.5	70.0	90.5				413	Specimen cross-grained; split with grain.....	655
4.8	9.6	14.5	20.5	1.2	21.6								200	Sap-wood; broke at knot.....	655
6.8	13.2	20.7	29.0	2.5	29.8	39.0	51.0	65.5					400	0.75 sap-wood; heart defective; typical break.....	608
5.5	11.0	16.5	23.0	1.5	23.4	29.5	38.3	48.5	63.0				426	Specimen cross-grained; split with grain.....	1103
6.1	12.4	19.3	27.0	2.0	27.3	35.0	46.0	61.5	80.0				421	do.....	1103
3.4	6.5	10.0	14.0	0.2	14.0	17.0	21.4	26.5	33.0	43.0			450	Specimen cross-grained; splinter on corner.....	404
3.8	7.0	10.7	14.0	0.4	14.5	18.0	22.7	28.0	34.0	43.5	55.5	73.5	564	Broke with fine splinters.....	799
4.0	7.8	11.5	15.5	0.9	15.7	20.0	25.7	32.0	40.5	50.2	66.7		517	do.....	799
4.0	8.2	12.3	16.3	0.5	16.2	20.6	26.5	33.0	55.0				431	Specimen cross-grained; broke at small knot.....	919
4.2	8.7	12.5	17.1	0.2	17.0	22.0	28.4	35.0	43.0				450	do.....	919
7.0	13.2	20.6	29.5	3.0	30.0	39.0	51.7	69.2					348	do.....	954
6.5	13.0	20.0	29.2	3.1	30.0	40.7							277	Specimen cross-grained; started at knot.....	954
5.0	9.0	13.5	18.2	0.5	18.2	23.0	28.0	35.5	43.0	54.2	73.0	120.0	558	Broke with fine splinters.....	649
4.7	8.5	12.2	16.6	0.5	16.6	20.0	25.5	32.0	38.0	46.5	59.0		541	Failed from large splinter on corner.....	649
4.0	8.0	12.0	16.4	0.6	16.5	21.0	27.0	33.0	40.0	49.0	66.0		548	Specimen cross-grained; large splinter on corner.....	653
3.8	7.3	11.2	14.7	0.6	15.0	19.0	24.5	30.5	36.5	45.0	59.0		517	Broke with coarse splinters.....	653
7.6	15.3	23.7	33.2	3.5	34.7	45.5	62.4						300	Specimen cross-grained; split with grain.....	654
4.6	10.2	15.6	21.8	1.6	22.7	29.6	39.2	52.0	75.0				400	Broke with fine splinters; buckled on compression side.....	663
5.0	10.3	15.0	21.3	1.5	21.9	29.0	38.0	51.0					397	Broke with long scale.....	663
5.4	11.5	17.5	25.4	1.7	25.8	35.0	47.0						324	Broke at knot near the end.....	685
5.7	11.2	16.8	23.3	1.3	24.0	33.0	39.5	51.5					374	Specimen cross-grained; square break on tension side, splitting in axis.....	685
2.5	6.0	9.0	12.0	0.1	12.2	15.3	19.0	23.2	28.2	34.2	45.2		547	Broke with fine splinters.....	7
3.8	7.2	10.7	14.4	0.2	14.7	18.0	22.5	28.0	35.0	45.2	67.0		500	do.....	7
4.3	8.5	12.3	16.6	0.3	16.8	21.5	27.5	35.0	46.0				447	Broke with thin scale and fine splinters.....	7
3.6	7.4	11.3	15.2	0.4	15.3	19.5	25.3	32.0	43.0	59.0			451	do.....	7
5.0	9.8	14.7	20.5	0.7	20.7	27.6	37.6	53.5					365	Broke with fine splinters.....	451
5.0	9.7	15.3	21.7	0.5	20.7	27.0	35.0	46.0					385	Broke with coarse splinters.....	453
5.7	11.5	17.0	23.7	1.2	24.7	32.5	47.5						336	Failed from scale on back.....	453
6.0	12.0	18.0	24.5	1.3	24.6	32.0	42.5						336	0.5 sap-wood; square break on tension side, splitting in axis.....	92
4.0	7.8	11.5	15.4	0.3	15.7	20.0	26.0	33.0					392	Failed from large splinter on each corner.....	140
6.4	12.7	19.2	26.4	1.3	26.7	35.0	47.0						333	Specimen cross-grained.....	141
4.0	8.0	12.5	16.8	0.8	17.0	22.1	28.3	35.5					382	Failed from long splinter on corner.....	146
3.6	7.2	11.0	14.5	0.3	14.6	18.7	23.5	29.0	36.0	48.5			451	do.....	215
5.0	9.5	14.5	20.0	0.6	20.0	26.2	34.5	46.0					374	Specimen cross-grained; broke with large splinters.....	215
5.5	11.0	17.0	24.0	2.0	24.5	31.0	41.0	52.5	68.0	92.0			457	Broke with coarse splinters.....	217
4.0	7.5	11.0	14.6	0.2	14.7	18.5	23.0	28.5	34.3	43.0	54.0		530	Broke with thin scale and fine splinters.....	218
4.0	7.7	11.2	15.4	0.3	15.5	19.7	25.0	30.7	38.0	46.0	60.0		500	Broke with fine splinters.....	218
5.7	12.0	18.0	27.0	2.3	27.7	39.5							290	Failed from long scale on back.....	920
5.0	10.6	16.0	24.0	2.0	24.5	33.0	49.0						325	Broke with long, coarse splinters.....	920

TABLE III.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Specific gravity of the air-dried specimen.	Direction of grain.	COEFFICIENT OF ELASTICITY.		Modulus of rupture.
								First deflection.	Second deflection.	
272. <i>Quercus rubra</i> —continued.....	1043	Massachusetts	North Reading	J. Robinson	Drift	0.7580		1221	1320	1282
	1043	do	do	do	do	0.7426		1221	1302	1172
272. <i>Quercus rubra</i> , var. <i>Texana</i> , <i>Red Oak</i>	981	Texas	Austin	C. Mohr	Calcareous	0.8580		976	1039	1252
	981	do	do	do	do	0.8926		1017	1028	794
273. <i>Quercus coccinea</i> , <i>Scarlet Oak</i>	752	Florida	Aspalaga	A. H. Curtiss	Clay	0.7654		976	1085	1055
274. <i>Quercus tinctoria</i> , <i>Black Oak</i> , <i>Yellow-bark Oak</i> , <i>Quercitron Oak</i> , <i>Yellow Oak</i>	17	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	0.7102		1221	1221	1151
	17	do	do	do	do	0.6847		1085	1149	1054
	36 ¹	Kentucky	Danville Junction	W. M. Linney	Shale	0.7284		976	1028	1043
	36 ²	do	do	do	do	0.7619		1221	1221	1270
	36 ³	do	do	do	Slate	0.7932		607	723	954
	74	Missouri	Allenton	G. W. Letterman	Rich upland	0.8495		921	930	1181
	74	do	do	do	do	0.8408		814	814	930
	86	do	do	do	do	0.7519		888	930	1120
	244	Virginia	Wytheville	H. Shriver	Clay	0.5432		842	820	794
	244	do	do	do	do	0.5026		561	558	551
	247	do	do	do	do	0.7102		1479	1479	1284
	247	do	do	do	do	0.7046		1395	1436	1261
	437	Tennessee	Nashville	A. Gattinger	do	0.6949		1136	1136	937
	628	Oregon	Saw-mill, Ashland	G. Engelmann and C. S. Sargent	do	0.6642		814	849	869
275. <i>Quercus Kelloggii</i> , <i>Black Oak</i>	628	do	do	do	do	0.6841		872	921	844
	963	do	Eugene City	G. H. Collier	do	0.6960		660	651	663
	963	do	do	do	do	0.6875		525	558	696
276. <i>Quercus nigra</i> , <i>Black Jack</i> , <i>Jack Oak</i>	268	Missouri	Allenton	G. W. Letterman	Clay	0.7688		976	976	1043
277. <i>Quercus falcata</i> , <i>Spanish Oak</i> , <i>Red Oak</i>	131	South Carolina	Bonneau's Depot	H. W. Ravenel	Rich loam	0.7521		1395	1302	1172
	181	do	do	do	do	0.7421		1320	1394	1071
	245	Virginia	Wytheville	H. Shriver	Clay	0.6307		1221	1221	1041
	245	do	do	do	do	0.6767		1320	1394	1172
	265 ²	do	Carroll county	do	do	0.7837		1526	1502	1334
	265 ³	do	do	do	do	0.6933		1221	1221	1172
	548	Mississippi	Kemper's mill	C. Mohr	Rich loam	0.7875		1526	1601	1266
	548	do	do	do	do	0.7614		1628	1575	1317
278. <i>Quercus Catesbei</i> , <i>Turkey Oak</i> , <i>Scrub Oak</i> , <i>Forked-leaf Black Jack</i> , <i>Black Jack</i>	342	Alabama	Cottage Hill	do	Barren, sandy	0.6875		888	948	990
	342	do	do	do	do	0.7193		1017	1062	1027
	770	Florida	Aspalaga	A. H. Curtiss	do	0.6395		976	1007	1113
	770	do	do	do	do	0.8442		1163	1122	1053
279. <i>Quercus palustris</i> , <i>Pin Oak</i> , <i>Swamp Spanish Oak</i> , <i>Water Oak</i>	47	Missouri	Allenton	G. W. Letterman	Rich, alluvial	0.7148		1526	1479	1233
	282	do	do	do	do	0.7671		957	976	1055
	282	do	do	do	do	0.7534		904	913	962
280. <i>Quercus aquatica</i> , <i>Water Oak</i> , <i>Duck Oak</i> , <i>Possum Oak</i> , <i>Punk Oak</i>	264 ¹	Virginia	Carroll county	H. Shriver	do	0.6523		1085	1122	969
	264 ²	do	do	do	do	0.6170		1136	1191	1052
	264 ³	do	do	do	do	0.6158		1163	1149	932
	349	Alabama	Cottage Hill	C. Mohr	Sandy loam	0.7142		1163	1285	1155
	349	do	do	do	do	0.7313		1221	1285	1120
	511	Tennessee	Tallahoma	A. Gattinger	do	0.7560		976	1028	919
	511	do	do	do	do	0.7411		1221	1338	1170

UNITED STATES UNDER TRANSVERSE STRAIN—Continued.

DEFLECTION, IN MILLIMETERS, UNDER A PRESSURE, IN KILOGRAMS, OF—													Ultimate strength: transverse pressure.	Remarks.	Office number.
50	100	150	200	0 (set.)	200	250	300	350	400	450	500	550			
4.0	7.4	11.0	14.6	0.3	15.0	19.0	24.0	30.3	37.0	46.0	63.5		547	Broke with fine splinters and scale	1043
4.0	7.5	11.4	15.4	0.3	15.6	20.0	26.0	31.7	40.0	51.5			500	Failed from splinter on each corner	1043
5.0	9.4	14.0	19.0	0.5	19.2	25.0	32.0	40.0	50.7	63.0	86.5		534	Specimen cross-grained; broke with large splinters	931
4.8	9.5	14.7	20.4	1.2	21.0	27.2	38.0						339	Specimen affected with dry rot; broke at knot near the end	931
5.0	9.0	14.0	20.1	1.4	20.5	27.0	36.0	47.0	62.5	90.0			450	Broke with fine splinters	752
4.0	8.0	12.0	16.1	0.3	16.4	21.0	27.0	34.0	42.0	56.5			491	Broke with medium splinters	17
4.5	8.5	12.2	16.5	0.4	16.5	21.1	27.0	34.5	44.0	55.0			450	Broke with scales and fine splinters	17
5.0	9.5	14.5	20.2	1.0	20.4	27.2	35.0	45.5	58.2				445	Broke with coarse splinters on corner	36 ²
4.0	8.0	12.0	16.5	0.6	16.4	20.7	27.0	32.5	41.5	51.7	68.0		542	Broke with fine splinters	36 ²
7.0	13.5	21.4	31.0	3.5	31.5	42.0	58.0	80.0	135.0				407	Broke with split through center parallel to pressure	36 ²
5.3	10.5	15.8	22.6	1.8	23.6	30.6	40.0	51.5	64.0	88.5	140.0		504	Broke with fine splinters	74
6.0	12.0	18.5	25.8	1.8	26.3	34.0	46.2	60.5					397	Specimen cross-grained, knotty	74
5.5	10.5	15.7	21.7	1.5	22.1	28.2	36.7	47.0	57.7	75.0			478	Failed from scale on back and splinter on corner	86
5.8	11.9	18.0	26.0	1.5	26.5	37.0	55.0						339	Square break on tension side, splitting in axis of stick	244
8.7	17.5	28.5	43.4	5.0	44.0								235	do	244
3.3	6.6	10.0	14.0	0.6	14.0	17.7	22.1	27.0	33.0	40.5	50.6		548	Crushed with fine splinters	247
3.5	6.8	10.0	13.4	0.4	13.4	16.7	20.5	25.6	31.7	40.5	51.5		538	do	247
4.3	8.6	12.8	18.0	1.0	18.2	24.5	33.0	44.0	55.0				400	Failed from large scale and coarse splinter	487
6.0	11.5	17.3	23.5	0.9	23.9	31.0	40.3	51.5					371	Specimen cross-grained	628
5.6	10.6	16.0	22.0	0.8	22.1	28.5	36.0	46.7					360	do	628
7.4	15.0	26.0	54.8	10.5	57.0	88.0							283	Failed from coarse splinter on compression side	963
9.3	17.5	27.5	41.5	4.8	43.5	61.5							298	Shaky	963
5.0	10.0	15.0	21.0	1.2	21.4	28.0	36.0	45.0	60.0				445	Broke with fine splinters	266
3.5	7.5	10.5	14.0	0.5	14.0	17.5	22.0	27.5	33.0	40.5	54.0		500	do	131
3.7	7.0	10.2	13.7	0.4	14.0	17.4	21.6	26.5	32.5	41.0	55.0		457	Broke with coarse splinters	131
4.0	8.0	12.0	16.0	0.5	16.0	20.2	25.7	33.0	42.0				444	Square break on tension side, splitting in axis	245
3.7	7.0	10.0	13.7	0.5	13.7	17.0	21.2	26.3	33.0	43.5	60.0		500	Crushed at center bearing; broke with fine splinters	245
3.2	6.5	9.5	12.7	0.4	13.0	16.2	20.7	25.5	30.5	38.5	48.0	65.5	569	do	265 ²
4.0	8.0	12.0	16.0	0.5	16.0	20.0	25.2	31.5	40.0	48.3			500	Broke with long, coarse splinters	265 ²
3.2	6.1	9.0	12.1	0.3	12.4	15.4	19.0	23.0	28.0	34.2	40.0		540	Broke at knot with long flake	548
3.0	6.2	9.3	12.0	0.2	12.0	15.0	18.5	23.0	28.2	35.5	46.0	63.2	562	Crushed slightly at center bearing	548
5.5	10.3	15.4	20.6	0.9	21.1	27.0	34.5	44.0	55.5				423	Broke with large splinters	242
4.8	9.2	14.0	19.0	1.0	19.0	25.0	31.3	39.0	50.2				438	Broke with large splinters; shattered	342
5.0	9.7	14.6	20.2	1.1	20.0	26.0	33.0	41.6	51.5	66.2			475	Specimen cross-grained; split through knot	770
4.2	8.7	13.0	18.5	1.0	18.9	24.2	31.2	39.0	50.0				450	Specimen cross-grained; shattered	770
3.2	6.6	10.2	13.7	0.3	14.2	17.5	22.5	28.0	35.5	44.0	63.0		526	Crushed at center bearing; broke with fine splinters	47
5.1	10.0	15.0	21.0	1.8	22.0	29.0	38.5	49.5	66.0	105.0			450	Square break on tension side, splitting in axis	282
5.4	10.7	17.0	24.0	2.1	24.4	32.5	44.0	60.0	91.0				419	Broke with scale and fine splinters	282
4.5	8.7	13.0	17.7	0.7	18.0	24.0	32.0	44.0	58.5				422	Square break on tension side, splitting in axis	264 ¹
4.3	8.2	11.8	16.0	0.2	16.5	21.0	26.7	34.0	44.6				449	Broke with fine scale	264 ²
4.2	8.5	13.0	18.0	0.9	18.3	24.2	32.5	44.2					398	Square break on tension side, splitting in axis	264 ²
4.2	7.6	11.5	15.5	0.5	15.6	20.0	24.7	31.0	37.0	47.0			493	Broke with fine splinters	349
4.0	7.6	11.5	15.8	0.6	16.0	20.4	26.5	34.0	44.5	58.5			478	do	349
5.0	9.5	14.6	20.4	1.2	20.7	26.5	34.3	44.0					392	Broke with coarse scales	511
4.0	7.3	10.9	14.3	0.5	15.0	19.0	24.0	30.0	38.5	47.0			499	Broke with coarse splinters	511

FOREST TREES OF NORTH AMERICA.

TABLE III.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Specific gravity of the air-dried specimen.	Direction of grain.	COEFFICIENT OF ELASTICITY.		Modulus of rupture.
								First deflection.	Second deflection.	
280. <i>Quercus aquatica</i> —continued	742	Georgia	Bainbridge	A. H. Curtiss	Alluvial	0.7415		1221	1285	806
	742	do	do	do	do	0.8006		1356	1356	1334
281. <i>Quercus laurifolia</i> <i>Laurel Oak.</i>	756	Florida	Saint John's river	do	Sandy loam	0.8034		1221	1177	1055
	756	do	do	do	do	0.7980		1320	1221	1111
	801	do	do	do	do	0.8105		1356	1320	1289
	801	do	do	do	do	0.8204		1221	1320	1266
282. <i>Quercus heterophylla</i> <i>Bartram's Oak.</i>	1171	New Jersey	Mount Holly	S. P. Sharples	Clay	0.7023		1395	1338	1165
	1171	do	do	do	do	0.6818		1110	1085	947
	1171	do	do	do	do	0.6808		1221	1252	1111
283. <i>Quercus cinerea</i> <i>Upland Willow Oak. Blue Jack.</i>	352	Alabama	Citronelle	C. Mohr	Pine-barren	0.7159		697	751	994
284. <i>Quercus hypoleuca</i> <i>Sand Jack.</i>	674	Arizona	Santa Rita mountains	G. Engelmann and C. S. Sargent	Dry, rocky	0.8716		904	967	1120
	674	do	do	do	do	0.8325		930	921	1106
285. <i>Quercus imbricaria</i> <i>Shingle Oak. Laurel Oak.</i>	401	Kentucky	Harrodsburg	W. M. Linney	Utica shale	0.7440		1136	1085	1153
	50	Missouri	Allenton	G. W. Letterman	Rich, moist	0.7477		1252	1302	1284
286. <i>Quercus Phellos</i> <i>Willow Oak. Peach Oak.</i>	512	Tennessee	Tullahoma	A. Gattinger	Moist, siliceous	0.7603		751	781	923
	512	do	do	do	do	0.7557		763	787	1055
287. <i>Quercus densiflora</i> <i>Tanbark Oak. Chestnut Oak. Peach Oak.</i>	687	California	Marin county	G. R. Vasey	Gravelly	0.6910		872	842	846
	687	do	do	do	do	0.7205		970	1085	1048
288. <i>Castanopsis chrysophylla</i> <i>Chinquapin.</i>	729	California	Mendocino county	A. Kellogg		0.5739		996	1007	846
	729	do	do	do		0.5909		1017	1017	635
189. <i>Castanea pumila</i> <i>Chinquapin.</i>	573	Arkansas	Hot Springs	G. W. Letterman	Sandy loam	0.6017		1265	1221	1031
	573	do	do	do	do	0.5960		1062	1062	949
290. <i>Castanea vulgaris</i> , var. <i>Americana</i> . <i>Chestnut.</i>	18	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	0.4123		498	525	415
	18	do	do	do	do	0.5330		888	872	703
	18	do	do	do	do	0.4568		679	651	619
	2582	Virginia	Fancy Gap	H. Shriver	Moist	0.5050		1221	1163	893
	2583	do	do	do	do	0.5053		1085	1039	867
	516	Tennessee	Nashville	A. Gattinger	Sandy	0.4939		872	888	675
291. <i>Fagus ferruginea</i> . <i>Beech.</i>	9	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	0.7333		1221	1302	1312
	9	do	do	do	do	0.7146		1221	1221	1235
	442	Kentucky	Mercer county	W. M. Linney	Hudson River shale	0.6352		1085	1050	1026
	443	do	do	do	do	0.6416		1039	1122	937
	119	Michigan	Dansville	W. J. Beal	Gravelly	0.7571		1221	1302	1284
	119	do	do	do	do	0.6945		1221	1302	1180
	785	Florida	Chattahoochee	A. H. Curtiss	do	0.6892		976	1007	1024
	785	do	do	do	do	0.6770		1163	1136	1048
	853	Massachusetts	Hamilton	J. Robinson	do	0.7000		1221	1221	1106
	853	do	do	do	do	0.7324		1285	1252	1321
	853	do	do	do	do	0.7250		1395	1395	1256
292. <i>Ostrya Virginica</i> <i>Hop Hornbeam. Iron Wood. Lever Wood.</i>	11	do	Arnold Arboretum	C. S. Sargent	Drift	0.7784		1628	1628	1446
	11	do	do	do	do	0.7614		1395	1479	1289
	877	do	Danvers	J. Robinson	Rich loam	0.8400		1395	1395	820
	877	do	do	do	do	0.8591		1320	1395	984
	1047	do	North Reading	do	do	0.8068		1110	1163	1209
	1047	do	do	do	do	0.8182		1186	1177	1057

UNITED STATES UNDER TRANSVERSE STRAIN—Continued.

DEFLECTION, IN MILLIMETERS, UNDER A PRESSURE, IN KILOGRAMS, OF—													Ultimate strength: transverse pressure.	Remarks.	Office number.
50	100	150	200	0 (set.)	200	250	300	350	400	450	500	550			
4.0	7.6	12.0	16.8	1.0	17.0	21.8	29.0						344	Specimen cross-grained	742
3.6	7.2	10.3	13.7	0.4	14.0	17.5	21.3	26.0	31.0	37.0	46.5	58.0	569	Broke with coarse splinters	742
4.0	8.3	12.2	17.0	0.8	17.0	22.5	28.7	36.0	46.3	60.0			450	Broke with large splinters	756
3.7	8.0	11.6	16.0	0.5	16.0	20.6	27.0	33.2	42.5	58.5			474	Broke with large splinters and scale on tension side	756
3.6	7.4	11.0	14.5	0.5	15.0	18.7	23.5	29.5	36.5	46.0	58.5	80.0	550	0.5 sap-wood; broke with fine splinters	801
4.0	7.4	11.0	15.0	0.6	15.5	19.0	24.5	30.5	37.5	46.7	58.5		540	do	801
3.5	7.3	11.0	14.7	0.5	15.0	19.2	23.7	29.5	37.0	48.0			497	Crushed at center bearing; broke with fine splinters	1171
4.4	9.0	12.0	16.5	0.5	17.0	20.9	26.0	33.0	42.0				404	Specimen cross-grained; broke with large splinters	1171
4.0	7.8	11.0	14.7	0.5	15.3	19.0	24.0	30.5	38.7	49.0			474	Broke with fine splinters	1171
7.0	13.0	19.5	26.5	1.5	27.0	35.0	46.0	59.0	76.0				424	Cross-grained; broke with coarse splinters	852
5.4	10.1	15.0	20.8	1.0	21.0	27.0	35.5	44.3	55.7	72.5			478	Broke with coarse splinters	674
5.2	10.6	16.0	22.7	1.2	22.7	30.0	37.0	47.0	58.0	73.0			472	do	674
4.3	9.0	13.5	18.5	0.5	18.6	24.4	31.5	40.0	49.5	60.7			492	0.5 sap-wood; broke with large splinters	401
3.9	7.5	11.2	15.3	0.5	16.0	19.8	25.0	31.5	40.0	50.0	62.0		548	Broke with fine splinters	50
6.5	12.5	19.4	28.0	2.1	29.0	39.0	52.5	69.0					394	Broke with large splinters, starting at knot	512
6.4	12.4	19.2	27.0	1.9	27.5	36.0	47.0	63.0	81.5	110.0			450	Broke with coarse splinters	512
5.6	11.6	17.5	24.5	1.0	24.4	31.0	40.5	55.0					361	Broke with scale; started at knot	687
5.0	9.0	13.6	18.2	0.5	18.4	23.5	29.5	37.0	46.0				447	Broke with coarse splinters	687
4.9	9.7	14.8	20.0	0.4	20.2	25.5	32.0	40.5					361	Specimen cross-grained; shattered	729
4.8	9.6	14.6	20.2	0.5	20.3	26.0							271	Specimen cross-grained; splinter on corner	729
3.8	8.0	12.0	16.2	0.5	16.2	20.5	25.7	32.2	40.7				441	Scale on tension side; broke with coarse splinters	573
4.6	9.2	14.0	19.0	1.2	19.2	24.6	31.5	41.0	62.0				405	Crushed at center bearing; broke with scales on tension side	573
9.8	18.6	30.4											177	Shattered	18
5.5	11.2	17.0	23.6	0.7	24.0	30.7	40.5						300	do	18
7.2	15.0	23.5	33.0	1.7	33.2	45.0							264	Square break; split at end	18
4.0	8.4	13.0	17.5	0.1	17.6	22.0	29.0	38.0					383	Crushed; square break	2589
4.5	9.4	14.4	19.5	0.4	19.6	25.0	33.0	41.0					370	Specimen cross-grained; shattered	2589
5.6	11.0	17.0	23.2	0.6	24.0	31.5							288	Broke with coarse splinters	516
4.0	7.5	11.6	15.7	0.5	15.6	20.0	26.0	30.7	36.5	45.0	55.5		560	do	9
4.0	8.0	12.2	16.2	0.3	16.3	21.0	26.7	32.5	42.0	54.0	75.0		527	Crushed; coarse splinters	9
4.5	9.3	14.0	19.0	0.4	19.0	25.0	31.5	41.0	52.5				438	Square break on tension side, splitting in axis	443
4.7	8.7	13.6	18.0	0.5	18.2	23.7	30.5	38.5	52.0				400	Scale on tension side	443
4.0	7.5	11.0	14.7	0.4	15.0	19.0	24.0	30.5	37.0	47.0	65.5		548	Broke with coarse splinters	119
4.0	7.5	11.2	14.7	0.5	15.0	19.0	24.5	30.0	37.5	46.7	70.0		504	Broke with fine splinters; scale on tension side	119
5.0	9.7	14.2	19.4	0.7	19.5	25.0	32.0	40.0	53.0				437	Broke with coarse splinters	765
4.2	8.6	13.2	18.0	0.6	18.5	23.6	30.0	39.0	51.5				447	Specimen cross-grained; broke at knot	765
4.0	8.0	11.5	16.0	0.5	16.5	19.7	25.0	31.0	40.0	49.5			472	Failed from long splinter on corner	853
3.8	7.8	11.6	15.4	0.5	15.5	20.0	25.5	31.0	40.0	50.0	76.5		521	Crushed with fine splinters	853
3.5	7.0	10.0	13.6	0.4	14.0	17.5	21.7	27.0	32.5	42.0	56.5		536	Crushed with long splinters	853
3.0	6.0	8.6	11.5	0.3	12.0	15.0	18.5	22.0	27.0	33.5	40.0	52.0	617	Sap-wood; broke with fine splinters; deflection with 600 kilograms pressure, 64 millimeters	11
3.5	6.6	10.0	13.5	0.2	13.7	17.0	22.0	28.0	34.0	43.0	60.0	75.0	550	Sap-wood; broke with fine splinters	11
3.5	7.0	10.4	14.0	0.4	14.0	18.0	22.3	28.0					350	Broke at knot	877
3.7	7.0	11.0	14.5	0.5	14.5	18.5	23.4	30.0	37.5				420	Square break on tension side, splitting in axis	877
4.4	8.4	13.0	17.5	1.0	18.0	23.7	30.5	40.0	51.5	67.5	102.0		516	Specimen cross-grained; splintered on corner	1047
4.3	8.3	12.6	17.5	1.0	18.0	23.5	31.0	40.0	52.0	73.5			451	Failed from scale on back	1047

TABLE III.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Specific gravity of the air-dried specimen.	Direction of grain.	COEFFICIENT OF ELASTICITY.		Modulus of rupture.
								First deflection.	Second deflection.	
298. <i>Carpinus Caroliniana</i> <i>Hornbeam. Blue Beech. Water Beech. Iron Wood.</i>	46	Missouri.....	Allenton.....	G. W. Letterman..	Damp, alluvial....	0.7798		921	957	1134
	73	Kentucky.....	Mercer county.....	W. M. Linney.....	Trenton limestone	0.7707		1221	1285	1355
	73 ¹	do.....	do.....	do.....	do.....	0.7852		1395	1395	1172
	73 ²	do.....	do.....	do.....	do.....	0.7641		898	939	1055
	73 ³	do.....	do.....	do.....	do.....	0.7983		1285	1395	1456
BETULACEÆ.	1038	Massachusetts....	Danvers.....	J. Robinson.....	Gravelly.....	0.7307		857	921	772
294. <i>Betula alba</i> , var. <i>populifolia</i> <i>White Birch. Old-field Birch. Gray Birch.</i>	10	do.....	Arnold Arboretum.	C. S. Sargent.....	Drift.....	0.6121		814	814	909
	10	do.....	do.....	do.....	do.....	0.6168		465	503	598
	848	do.....	Danvers.....	J. Robinson.....	Gravelly.....	0.5049		872	872	837
295. <i>Betula papyrifera</i> <i>Canoes Birch. White Birch. Paper Birch.</i>	722	Montana.....	Miasoula.....	Sereno Watson...	Wet.....	0.6244		1110	1149	1083
	722	do.....	do.....	do.....	do.....	0.5048		1221	1149	1008
	836	Massachusetts....	Townsend.....	J. Robinson.....	do.....	0.7084		1221	1320	1181
	990	Alaska.....	Chilcoot inlet.....	Paul Schultze.....	do.....	0.6136		1039	1065	816
	990	do.....	do.....	do.....	do.....	0.6239		1231	1191	994
	1065	Vermont.....	Charlotte.....	C. G. Pringle.....	do.....	0.6021		1526	1502	1165
	1065	do.....	do.....	do.....	do.....	0.6053		1628	1575	1099
	1065	do.....	do.....	do.....	do.....	0.6614		1395	1526	1228
	1066	do.....	do.....	do.....	do.....	0.6136		1085	1191	1090
	1066	do.....	do.....	do.....	do.....	0.6136		1320	1338	1017
	1067	do.....	do.....	do.....	do.....	0.6409		1356	1356	1134
	1067	do.....	do.....	do.....	do.....	0.6028		1285	1285	982
296. <i>Betula occidentalis</i> <i>Black Birch.</i>	528	Colorado.....	Engelmann's cañon.	Robert Douglas...	Wet, sandy.....	0.5996		888	872	738
	629	California.....	Strawberry valley..	G. Engelmann and C. S. Sargent.	Wet, peaty.....	0.5805		976	976	874
297. <i>Betula lutea</i> <i>Yellow Birch. Gray Birch.</i>	843	Massachusetts....	Danvers.....	J. Robinson.....	Gravelly.....	0.7147		1628	1650	1202
	843	do.....	do.....	do.....	do.....	0.7215		1575	1526	1289
	1068	Vermont.....	Charlotte.....	C. G. Pringle.....	do.....	0.6579		1575	1601	1216
	1068	do.....	do.....	do.....	do.....	0.6647		1628	1628	1219
	1069	do.....	do.....	do.....	do.....	0.6578		1479	1628	1320
	1069	do.....	do.....	do.....	do.....	0.6647		1628	1684	1212
	1070	do.....	do.....	do.....	do.....	0.6710		1479	1550	1245
	1070	do.....	do.....	do.....	do.....	0.6056		1953	1775	1282
298. <i>Betula nigra</i> <i>Red Birch. River Birch.</i>	136	Missouri.....	Allenton.....	G. W. Letterman..	Moist loam.....	0.5835		1039	1085	1041
	136	do.....	do.....	do.....	do.....	0.5815		1221	1221	975
	841	Massachusetts....	North Andover.....	J. Robinson.....	Alluvial.....	0.6335		1039	1085	954
	841	do.....	do.....	do.....	do.....	0.6273		1221	1356	1005
	842	do.....	do.....	do.....	do.....	0.6346		921	904	937
	842	do.....	do.....	do.....	do.....	0.6149		1039	1028	926
299. <i>Betula lenta</i> <i>Cherry Birch. Black Birch. Sweet Birch. Mahogany Birch.</i>	4	do.....	Arnold Arboretum.	C. S. Sargent.....	Drift.....	0.7393		1395	1436	1289
	4	do.....	do.....	do.....	do.....	0.7454		1479	1502	1282
	221	Vermont.....	Charlotte.....	C. G. Pringle.....	Gravelly.....	0.7002		1356	1302	1294
	844	Massachusetts....	Danvers.....	J. Robinson.....	do.....	0.8579		1395	1415	1052
	844	do.....	do.....	do.....	do.....	0.8215		1575	1502	1165
301. <i>Alnus rubra</i> <i>Alder.</i>	967	Alaska.....	Sitka.....	Paul Schultze.....	do.....	0.5000		976	1028	701
	967	do.....	do.....	do.....	do.....	0.5017		1017	1017	703
	901	Washington territory.	Puyallup.....	G. Engelmann and C. S. Sargent.	do.....	0.5190		842	888	830
	901	do.....	do.....	do.....	do.....	0.5244		842	807	820

UNITED STATES UNDER TRANSVERSE STRAIN—Continued.

DEFLECTION, IN MILLIMETERS, UNDER A PRESSURE, IN KILOGRAMS, OF—													Ultimate strength: transverse pressure.	Remarks.	Office number.
50	100	150	200	0 (set.)	200	250	300	350	400	450	500	550			
5.3	10.2	15.2	21.0	1.2	21.5	28.0	36.4	48.0	60.5	80.0			484	Square break on tension side, splitting in axis	46
4.0	7.6	11.0	15.0	0.4	15.0	19.0	24.0	29.7	35.5	45.0	56.0	68.0	578	Broke with fine splinters	73
8.5	7.0	10.5	14.0	0.6	14.4	18.6	23.5	29.0	34.8	41.5			500	Specimen cross-grained; failed from large splinter on corner	73 ¹
5.5	10.4	15.7	21.7	1.5	22.4	29.0	37.0	50.5	64.0	93.5			450	Square break on tension side, splitting in axis	73 ²
3.8	7.0	10.0	13.7	0.4	14.0	17.7	21.7	27.2	33.0	38.5	48.0	59.0	600	Deflection with 600 kilograms pressure, 82 millimeters; broke with fine splinters.	73 ³
5.7	10.6	17.0	24.0	2.2	24.8	33.0	45.0						330	Specimen cross-grained; started at knot	1038
6.0	12.0	18.0	25.4	0.4	26.0	34.2	48.5	67.0					388	Crushed at center bearing; square break on tension side	10
10.5	19.4	30.6	46.0	6.5	47.8								250	Broke at knot	10
5.6	11.2	17.6	25.5	2.5	26.8	36.2	50.0	82.0					357	Crushed at center bearing; broke with fine splinters	848
4.4	8.5	13.0	17.2	0.5	17.5	22.6	28.4	35.7	46.0	65.0			462	Square break on tension side, splitting in axis	722
4.0	8.5	13.4	19.0	1.0	19.2	25.0	33.0	44.0	58.5				430	Broke with coarse splinters	722
4.0	7.4	11.0	15.0	0.5	15.0	19.0	24.0	30.0	37.5	49.0	68.7		504	Failed from scale and long splinter on corner	836
4.7	9.0	14.0	19.5	1.0	20.0	26.3	35.0						348	Specimen cross-grained; broke on corner	990
4.0	8.2	12.4	17.0	0.5	17.0	22.0	28.0	36.0	50.0				424	Broke with coarse splinters	990
3.2	6.5	10.0	13.0	0.5	13.4	17.0	21.0	26.5	33.0	43.0			497	Crushed at center bearing; broke with fine splinters	1065
3.0	6.2	9.5	13.3	0.7	13.3	17.0	21.5	26.5	33.0	43.0			469	Crushed at center bearing; broke with large splinter on corner	1065
3.5	6.4	10.0	13.4	0.6	13.7	17.0	22.0	27.0	33.0	39.0	51.5		534	Square break on tension side; broke with splinters	1065
4.5	8.2	12.6	17.0	0.4	17.8	21.7	27.5	35.0	44.0				456	Specimen cross-grained; broke with coarse splinters	1066
3.7	7.3	11.2	15.0	0.4	15.5	19.7	25.0	32.0	41.6				434	Crushed at center bearing; broke with fine scales	1066
3.6	7.2	10.7	14.3	0.3	14.3	18.0	23.0	28.0	35.5	43.0			484	Failed from scales and coarse splinters	1067
3.8	7.6	12.0	16.5	0.5	16.5	21.5	28.0	36.0	52.0				419	Crushed at center bearing; splintered	1067
5.5	11.2	18.0	27.0	2.9	27.4	37.5	54.0						315	Specimen cross-grained; shattered	528
5.0	10.0	15.5	21.4	0.9	21.6	29.2	40.5	60.0					373	Crushed at center bearing; square break on tension side, splitting in axis.	629
3.0	6.3	9.3	12.7	0.5	12.7	16.2	20.0	24.5	30.5	36.5	50.0		513	Sap-wood; specimen cross-grained; shattered	843
3.1	6.4	9.2	12.5	0.4	12.6	15.7	19.5	24.5	30.0	36.0	47.5	60.0	550	Sap-wood; crushed at center bearing; broke with coarse splinter on corner.	843
3.1	6.1	9.4	12.4	0.3	12.2	15.4	19.2	23.2	29.0	37.5	57.0		519	Sap-wood; crushed at center bearing; broke with coarse scales	1068
3.0	6.0	9.0	11.5	0.3	11.5	14.5	18.0	21.7	27.0	34.5	49.5		520	Crushed at center bearing; broke with fine splinters	1068
3.3	6.0	8.7	11.6	0.2	11.5	14.0	17.0	20.7	25.0	31.0	40.0	55.0	563	0.5 sap-wood; crushed at center bearing; broke with fine splinters	1069
3.0	5.8	8.8	11.6	0.3	11.7	14.5	17.7	21.7	26.5	33.0	45.5		517	0.75 sap-wood; crushed at center bearing; broke with fine splinters	1069
3.3	6.3	9.4	12.4	0.3	12.4	15.0	19.5	24.0	30.5	38.0			531	Sap-wood; crushed at center bearing; broke with fine splinters on corner.	1070
2.5	5.5	8.3	11.3	0.1	11.4	14.5	18.0	22.0	27.0	33.7	45.5		547	Heart-wood; crushed at center bearing; broke with fine splinters	1070
4.7	9.0	13.3	18.2	0.5	18.6	23.5	30.0	39.0	64.5				444	Sap-wood; broke with coarse splinters at corners	136
4.0	8.0	12.0	15.9	0.7	16.3	21.0	27.0	35.5	51.0				416	Sap-wood; crushed at center bearing; broke with coarse splinters at corners.	136
4.7	9.0	14.0	19.0	0.9	19.4	25.0	32.5	45.0	66.0				407	Sap-wood; crushed at center bearing; splintered at corners	841
4.0	7.2	11.0	14.7	0.6	14.7	19.0	24.0	31.0	41.0				429	Sap-wood; crushed at center bearing; broke with fine splinters on tension side.	841
5.3	10.8	17.0	23.6	1.5	24.0	32.0	45.0	61.5	100.0				400	Sap-wood; crushed at center bearing; shattered on corner	842
4.7	9.5	14.0	19.2	0.5	19.4	25.0	33.0	45.0					395	Sap-wood; specimen cross-grained; started at knot	843
3.5	6.8	10.0	13.5	0.4	13.5	17.5	21.6	27.0	34.0	40.7	51.5	72.0	550	Sap-wood; crushed at center bearing; square break on tension side, splitting in axis.	4
3.3	6.5	10.0	13.0	0.3	13.5	16.7	20.8	26.0	32.0	40.0	51.0		547	Sap-wood; splintered on corners	4
3.6	7.5	10.4	13.3	0.2	13.6	16.5	19.7	24.0	28.5	34.5	43.7	60.0	552	Square break on tension side, splitting in axis	221
3.5	6.9	10.0	13.6	0.4	13.5	17.2	22.0	27.0	34.0				448	Splintered on corners	844
3.1	6.5	10.0	13.7	0.4	14.0	18.0	22.6	28.0	35.5	45.5			497	Specimen cross-grained	844
5.0	9.5	14.4	19.0	0.5	19.5	25.5							299	Square break on tension side, splitting in axis	967
4.8	9.6	14.6	19.8	0.5	20.0	26.0							300	do	967
5.6	11.0	16.7	22.5	0.7	22.7	30.0	37.5	50.0					354	do	991
5.8	12.1	19.0	27.0	1.9	27.5	37.0	50.0	72.0					350	do	991

TABLE III.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Specific gravity of the air-dried specimen.	Direction of grain.	COEFFICIENT OF ELASTICITY.		Modulus of rupture.
								First deflection.	Second deflection.	
301. <i>Alnus rubra</i> —continued.	1025	Oregon	Portland Furniture Company.	G. Engelmann and C. S. Sargent.		0.4739		1320	1285	900
	1025	do	do	do		0.5084		1221	1338	905
302. <i>Alnus rhombifolia</i> Alder.	635	do	Ashland	do		0.4857		904	948	738
	717	Montana	Missoula	Sereno Watson	Wet	0.4545		679	669	567
	717	do	do	do	do	0.4886		729	751	701
	979	Oregon	Drain	C. S. Sargent.	Moist loam	0.4465		904	948	698
	979	do	do	do	do	0.4375		888	913	703
303. <i>Alnus oblongifolia</i> Alder.	694	California	San Bernardino	W. G. Wright		0.4170		787	751	696
	694	do	do	do		0.4049		787	787	677
305. <i>Alnus incana</i> Speckled Alder. Hoary Alder. Black Alder.	802	Massachusetts	Danvers	J. Robinson	Wet loam	0.5005		1163	1085	820
SALICACEÆ.										
306. <i>Salix nigra</i> Black Willow.	855	do	Topsfield	do	Alluvial	0.5330		407	391	424
307. <i>Salix amygdaloides</i> Willow.	908	Colorado	Cañon City	E. Weston		0.4675		610	514	511
	908	do	do	do		0.4676		488	476	530
	911	do	do	do		0.4670		554	514	609
308. <i>Salix laevigata</i> Willow.	690	California	Santa Cruz	G. Engelmann and C. S. Sargent.	Moist, sandy	0.5341		479	488	734
	690	do	do	do	do	0.5284		509	488	553
309. <i>Salix lasiandra</i> , var. <i>lanceifolia</i>	640	do	Strawberry valley	do	Moist, rich	0.4705		542	525	584
	981	Oregon	Portland	F. Skinner	Alluvial	0.4375		1163	1085	701
	981	do	do	do	do	0.4614		976	1028	743
309. <i>Salix lasiandra</i> , var. <i>Fendleri</i> ana	880	Utah	City Creek cañon	M. E. Jones	Gravelly	0.4406		330	305	469
313. <i>Salix flavescens</i>	721	Montana	Pattée's cañon, Missoula.	Sereno Watson	Rich, moist	0.5057		1221	1085	809
313. <i>Salix flavescens</i> , var. <i>Scouleriana</i> Black Willow.	972	Washington territory.	Seattle	G. Engelmann and C. S. Sargent.	Moist, sandy	0.5719		1285	1375	928
	972	do	do	do	do	0.5278		1221	1149	888
316. <i>Salix lasiolepis</i> Willow.	669	California	Santa Cruz	do	Sandy loam	0.5760		921	888	813
318. <i>Populus tremuloides</i> Aspen. Quaking Asp.	272 ^a	Colorado	Alpine	T. S. Brandegee	Damp	0.3955		740	781	562
	272 ^a	do	do	do	do	0.3885		651	651	541
	1035	Massachusetts	Danvers	J. Robinson	Gravelly	0.4590		976	913	720
	1035	do	do	do	do	0.4828		888	913	881
319. <i>Populus grandidentata</i> Poplar.	847	do	do	do	do	0.4443		976	996	696
	847	do	do	do	do	0.4386		904	930	748
320. <i>Populus heterophylla</i> River Cottonwood. Swamp Cottonwood.	522	Tennessee	Nashville	A. Gittinger	Alluvial	0.4155		740	723	642
321. <i>Populus balsamifera</i> Balsam. Tucamahac. Balm of Gilead.	961	Alaska	Chilcoot inlet	Paul Schultze	do	0.4080		857	842	574
	961	do	do	do	do	0.4034		888	872	527
321. <i>Populus balsamifera</i> , var. <i>candicans</i>	1054	Massachusetts	Topsfield	J. Robinson	Gravelly	0.4085		740	697	626
	1054	do	do	do	do	0.4506		751	763	593
322. <i>Populus angustifolia</i> Black Cottonwood.	552	Colorado	Manitou Springs	Robert Douglas	Sandy loam	0.4488		488	458	401
323. <i>Populus trichocarpa</i> Black Cottonwood. Balsam Cottonwood.	1012	Oregon	Saint John's Barrel Factory, Portland.	F. Skinner		0.3750		1062	1050	682
	1012	do	do	do		0.3898		1085	1062	637
	1028	do	Portland Furniture Company.	G. Engelmann and C. S. Sargent.		0.4108		1221	1221	694
	1028	do	do	do		0.4108		1163	1136	745

UNITED STATES UNDER TRANSVERSE STRAIN—Continued.

DEFLECTION, IN MILLIMETERS, UNDER A PRESSURE, IN KILOGRAMS, OF—													Ultimate strength: transverse pressure.	Remarks.	Office number.
50	100	150	200	0 (set.)	200	250	300	350	400	450	500	550			
8.7	7.6	11.5	15.0	0.1	15.7	20.0	25.2	33.0					384	Crushed at center bearing; broke with fine splinters.	1025
4.0	7.3	11.0	14.7	0.0	14.8	18.5	23.5	30.0					386	Specimen cross-grained; scaled on corner.	1025
5.4	10.3	15.5	20.6	0.5	21.2	27.0	35.7						315	Square break on tension side, splitting in axis.	635
7.2	14.6	23.0	34.0	3.5	35.0								242	Sap-wood; cross-grained.	717
6.7	13.0	20.4	30.0	3.0	31.0	44.0							299	Square break.	717
5.4	10.3	15.7	21.8	0.9	22.0	30.2							298	Square break on tension side, splitting in axis.	979
5.5	10.7	16.5	22.2	1.0	23.0	30.6	50.0						300	Specimen cross-grained; square break on tension side, splitting in axis.	979
6.2	13.0	19.6	28.3	1.3	29.0	39.0							297	Crushed at center bearing; square break on tension side, splitting in axis.	694
6.2	12.4	19.0	27.3	1.5	28.0	38.7							289	Crushed at center bearing; splintered on corner.	694
4.2	9.0	14.0	19.6	0.8	19.6	26.0	35.0						350	Broke with coarse splinters.	862
12.0	25.0	44.0											181	Specimen cross-grained; split with grain.	855
8.0	19.0	36.0	64.5	16.2	73.0								218	Crushed at center bearing; splintered on compression side.	908
10.0	20.5	36.0	58.0	11.0	63.0								226	do.	908
8.8	19.0	32.5	55.0	11.5	57.0	91.0							260	Crushed at center bearing; splintered on tension side.	911
10.2	20.0	31.2	46.0	5.0	47.5	64.0	110.0						313	Square break on tension side, splitting in axis; shattered.	690
9.6	20.0	32.0	46.0	4.8	48.5								236	Specimen cross-grained; flaked on tension side.	690
9.0	18.6	31.0	46.0	5.0	48.0								240	Specimen cross-grained; broke at knot.	640
4.2	9.0	14.2	21.0	1.2	21.5	29.5	45.0						299	Crushed at center bearing; broke with fine splinters.	961
5.0	9.5	14.3	19.5	0.5	20.0	26.0	36.0						317	Crushed at center bearing; scaled on tension side.	961
14.8	32.0	57.0	125.0	32.5									200	Crushed at center bearing; shattered.	889
4.0	9.0	14.0	19.0	1.1	20.0	26.5	36.5						345	Crushed at center bearing; splintered; square break on tension side, splitting in axis.	721
3.8	7.1	10.5	14.2	0.5	14.7	18.3	24.0	31.5					396	Crushed at center bearing; broke with fine splinters.	972
4.0	8.5	12.5	16.8	0.3	17.2	22.0	28.0	35.0					379	Crushed at center bearing; broke with coarse splinters.	972
5.3	11.0	16.6	24.5	1.5	25.0	33.0	46.0						347	Crushed at center bearing; broke with fine splinters.	669
6.6	12.5	19.3	27.7	2.4	28.7								240	Broke with long scale.	272 ^a
7.5	15.0	23.2	33.0	3.8	37.2								231	Specimen cross-grained; broke at knot.	272 ^a
5.0	10.7	17.0	24.0	1.5	24.5	32.0	50.0						309	Crushed at center bearing.	1035
5.5	10.7	17.0	25.0	2.9	26.0	35.0	48.0	71.5					376	Broke with many fine splinters.	1035
5.0	9.8	15.0	21.4	1.5	22.0	30.0							297	Splintered on corner.	847
5.4	10.5	15.5	21.7	1.0	22.0	28.7	41.0						319	Crushed at center bearing; broke with fine splinters.	847
6.6	13.5	21.4	30.7	3.3	32.0	47.0							274	do.	522
5.7	11.6	17.4	25.0	1.3	26.2								245	Specimen affected with dry rot; square break.	961
5.5	11.2	17.6	25.5	1.7	26.2								225	do.	961
6.6	14.0	22.4	34.2	4.1	35.8	52.0							267	Crushed at center bearing; square break.	1054
6.5	12.8	21.0	34.8	5.9	36.0	62.0							253	do.	1054
10.0	21.3	35.2											171	Specimen cross-grained; broke at knot.	558
4.6	9.3	14.0	20.8	1.0	20.8	28.4							291	Square break with small splinters.	1012
4.5	9.2	14.0	19.6	0.9	20.0	27.5							272	do.	1012
4.0	8.0	12.0	18.0	1.1	17.6	24.0							296	do.	1028
4.2	8.6	12.6	17.6	0.5	17.6	25.0							275	do.	1028

TABLE III.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Specific gravity of the air-dried specimen.	Direction of grain.	COEFFICIENT OF ELASTICITY.		Modulus of rupture.
								First deflection.	Second deflection.	
324. <i>Populus monilifera</i> <i>Cottonwood. Necklace Poplar. Carolina Poplar. Big Cottonwood.</i>	255	Missouri	Allenton	G. W. Letterman	Alluvial	0.3552		669	669	548
	304	do	do	do	do	0.3824		888	857	661
	304	do	do	do	do	0.3477		751	723	605
	309	Texas	Dallas	J. Reverchon	do	0.5136		1136	1149	893
	309	do	do	do	do	0.4858		842	842	759
	754	Florida	Chattahoochee	A. H. Curtiss	do	0.5001		1221	1302	902
	754	do	do	do	do	0.4909		1221	1221	937
	754	do	do	do	do	0.4925		1221	1191	851
325. <i>Populus Fremontii</i> <i>Cottonwood.</i>	659	California	Sacramento valley	G. R. Vasey	do	0.4977		1017	1017	584
	659	do	do	do	do	0.5000		1085	1085	813
325. <i>Populus Fremontii</i> , var. <i>Wializeni</i> . <i>Cottonwood. White Cottonwood.</i>	646	do	San Bernardino	W. G. Wright	do	0.5079		651	665	743
	646	do	do	do	do	0.5023		669	679	687
	909	Colorado	Cañon City	E. Weston		0.4867		904	1149	795
	909	do	do	do		0.4451		976	939	703
	909	do	do	do		0.4499		872	835	579
	909	do	do	do		0.5107		763	794	642
CONIFERÆ.										
326. <i>Libocedrus decurrens</i> <i>White Cedar. Bastard Cedar. Post Cedar. Incense Cedar.</i>	634	California	Saw-mill, Strawberry valley.	G. Engelmann and C. S. Sargent.		0.3748		697	751	619
	634	do	do	do		0.3979		814	814	702
	662	do	Saw-mill, San Bernardino mountains.	W. G. Wright		0.4818		872	896	783
	662	do	do	do		0.4989		872	888	654
	662	do	do	do		0.4869		828	888	654
327. <i>Thuja occidentalis</i> <i>White Cedar. Arbor-vita.</i>	879	Vermont	Monkton	C. G. Pringle	Cold, peaty	0.3023		488	514	455
	879	do	do	do	do	0.2902		444	452	441
	782	New Brunswick		Intercolonial railway.		0.3580		519	542	544
	783	do	Bridgeton	Ed. Sinclair		0.3439		651	610	527
	783	do	do	do		0.3413		626	651	448
	790	Province of Quebec	Amqui	A. Grant		0.3216		588	610	584
	790	do	do	do		0.3336		610	581	586
	792	do		Grand Trunk railway.		0.2714		331	337	394
	792	do		do		0.2612		279	294	352
	792	do		do		0.2805		269	269	354
	874	Maine	Mattawamkeag	J. Robinson		0.3835		763	781	586
328. <i>Thuja gigantea</i> <i>Red Cedar. Oance Cedar.</i>	1099	Wisconsin	Eau Claire	H. C. Putnam	Drift	0.3560		542	558	586
	1099	do	do	do	do	0.4121		514	542	654
328. <i>Thuja gigantea</i> <i>Red Cedar. Oance Cedar.</i>	1017	Oregon	Weidler's saw-mill, Portland.	G. Engelmann and C. S. Sargent.		0.4312		1221	1149	818
	1017	do	do	do		0.4256		976	1028	820
	1021	do	Portland Furniture Company.	do		0.3623		872	930	635
	1021	do	do	do		0.3783		1017	1028	722
329. <i>Chamaecyparis spheroides</i> <i>White Cedar.</i>	850	Alabama	Cottage Hill	C. Mohr	Sandy, wet	0.3626		271	257	328
	850	do	do	do	do	0.3153		341	334	280
	850	Massachusetts	Beverly	J. Robinson	Swampy	0.3625		496	488	483
	850	do	do	do	do	0.3602		425	415	551
	851	do	do	do	do	0.3523		444	444	469
	851	do	do	do	do	0.3407		519	514	509
	862	do	do	do	do	0.3324		388	376	469

UNITED STATES UNDER TRANSVERSE STRAIN—Continued.

DEFLECTION, IN MILLIMETERS, UNDER A PRESSURE, IN KILOGRAMS, OF—													Ultimate strength: transverse pressure.	Remarks.	Office number.
50	100	150	200	0 (set.)	200	250	300	350	400	450	500	550			
7.3	14.6	23.0	33.5										234	Crushed at center bearing; square break on tension side, splitting in axis.	255
5.5	11.4	17.2	24.2	1.4	25.3	33.2							282	Crushed at center bearing; square break	304
6.5	13.5	21.0	30.0	2.6	30.7	41.5							258	do	304
4.3	8.5	13.3	18.8	1.1	19.1	26.0	35.2	51.5					381	Crushed at center bearing; broke with fine splinters	309
5.8	11.6	18.0	25.6	3.0	26.5	36.0	48.5						324	Crushed at center bearing; shattered	309
4.0	7.5	12.0	16.0	0.4	16.0	20.7	25.8	33.0					385	Crushed at center bearing; broke with fine splinters	754
4.0	8.0	12.5	16.8	0.3	17.0	21.5	27.5	35.5					400	Shattered	754
4.0	8.2	12.8	17.2	0.6	17.5	22.4	29.0	39.0					363	do	754
4.8	9.6	14.8	21.4	1.7	21.8								249	Started at knot; splintered at corner	659
4.5	9.0	14.0	19.4	0.9	19.4	25.3	33.3						347	Specimen cross-grained; splintered on corner	659
7.5	14.7	22.0	32.4	2.4	33.0	43.0	62.5						317	Specimen cross-grained; split	646
7.3	14.4	23.0	32.6	2.5	34.0	47.0							293	Crushed at center bearing; square break on tension side, splitting in axis.	646
5.4	8.5	13.4	19.0	1.3	19.5	25.5	34.6						339	do	909
5.0	10.4	16.4	23.0	1.5	23.2	31.0	42.5						300	Square break on tension side, splitting in axis	909
5.6	11.7	18.4	26.7	2.1	27.2								247	Shattered	909
6.4	12.3	20.0	31.0	5.5	30.4	43.5							274	Broke with fine splinters	909
7.0	13.0	20.7	28.0	1.2	29.0	41.5							264	Square break	634
6.0	12.0	18.0	25.0	0.6	25.5	32.5	45.0						300	do	634
5.6	10.9	16.5	22.5	0.5	23.0	29.0	35.5						334	Specimen cross-grained; shattered	662
5.6	11.0	16.5	22.0	0.7	22.0	29.0							279	do	662
5.9	11.0	16.6	22.0	0.4	22.5	28.5							279	do	662
10.0	19.0	30.0											194	Square break; splintered	379
11.0	21.6	34.6											188	do	379
9.4	18.0	28.0	38.7	2.0	40.7								232	Square break on tension side, splitting in axis	783
7.5	16.9	24.0	33.5	1.2	34.6								225	do	783
7.8	15.0	23.2											191	Broke with coarse splinters	783
8.3	16.0	24.5	35.5	2.0	37.0								249	Square break; splintered	790
8.0	16.8	26.0	36.2	1.6	37.2								250	Square break; splintered	790
14.7	29.0	50.5											168	Square break	792
17.5	33.2												150	Square break; split to end	792
18.2	37.5	63.0											151	Square break	792
6.4	12.5	19.5	27.5	1.6	28.7								250	do	874
7.3	13.4	20.5	28.5	1.4	29.0	39.0							284	do	874
9.0	17.5	27.0	38.0	2.4	40.5								250	do	1099
9.5	18.0	28.5	39.0	2.1	40.8	55.0							279	Square break with long splinters	1099
4.0	8.5	13.0	17.8	0.4	18.0	23.0	30.0						349	Shattered	1017
5.0	9.5	14.0	19.0	0.5	19.0	24.0	31.0						350	Square break	1017
5.6	10.5	15.5	21.7	0.8	22.2	31.0							271	Shattered	1021
4.8	9.5	14.5	20.0	0.8	20.0	26.7	37.0						308	Square break on tension side, splitting in axis	1021
18.0	38.0												140	do	350
14.3	29.2	50.5											162	Square break	350
9.8	20.0	32.5	47.5	5.2	50.2								206	Square break on tension side, splitting in axis	850
11.5	23.5	36.0	54.0	5.6	55.0								235	Square break	850
11.0	22.0	33.2											200	Square break on tension side, splitting in axis	851
9.4	19.0	29.6	46.0	5.2	49.5								217	Square break	851
12.6	26.0	38.0	63.0	8.2									200	do	853

TABLE III.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Specific gravity of the air-dried specimen.	Direction of grain.	COEFFICIENT OF ELASTICITY.		Modulus of rupture.
								First deflection.	Second deflection.	
330. <i>Chamaecyparis Nutkaensis</i> <i>Yellow Cypress. Sitka Cypress.</i>	969	Alaska	Sitka	Paul Schultze		0.5697		814	814	851
	969	do	do	do		0.4229		888	976	696
	983	British Columbia	Saw-mill, Victoria	G. Engelmann and C. S. Sargent.		0.5267		1221	1221	1020
	983	do	do	do		0.5114		1110	1191	930
	994	Alaska	Peril strait	Paul Schultze		0.5074		1163	1149	872
	994	do	do	do		0.4913		1085	1122	937
	994	do	do	do		0.5267		1330	1221	930
	1000	do	Weidner's saw-mill, Portland.	G. Engelmann and C. S. Sargent.		0.5078		814	842	696
331. <i>Chamaecyparis Lawsoniana</i> <i>Port Orford Cedar. Oregon Cedar. White Cedar. Lawson's Cypress. Ginger Pine.</i>	701	Oregon	Dean & Co.'s saw-mill, Marshfield.	do		0.5239		904	930	830
	707	do	do	do		0.4682		1221	1221	818
	707	do	do	do		0.5335		1628	1502	1029
332. <i>Cupressus macrocarpa</i> <i>Monterey Cypress.</i>	675	California	Monterey	do	Gravelly loam	0.6307		976	1063	1041
	675	do	do	do	do	0.6512		1085	1085	1052
333. <i>Cupressus Goveniana</i> <i>Juniper.</i>	691	do	Marin county	G. R. Vasey	Dry ridges	0.5580		452	456	630
	691	do	do	do	do	0.5563		514	542	650
	1100	do	Calistoga	W. F. Fisher	do	0.4834		610	593	635
	1100	do	do	do	do	0.5078		457	407	342
337. <i>Juniperus pachyphloea</i> <i>Juniper.</i>	692	Arizona	Santa Rita mountains.	G. Engelmann and C. S. Sargent.	Gravelly	0.5630		610	630	783
	692	do	do	do	do	0.5419		595	595	738
338. <i>Juniperus occidentalis</i> , var. <i>conjugens</i> . <i>Juniper.</i>	1102	Texas	Austin	S. B. Buckley	Limestone	0.7347		751	734	460
339. <i>Juniperus Virginiana</i> <i>Red Cedar. Savin.</i>	14	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	0.5316		509	501	701
	14	do	do	do	do	0.5302		514	514	743
	327	Texas	Dallas	J. Reverchon	Calcareous	0.5357		595	610	851
	327	do	do	do	do	0.5239		610	630	562
	734	Florida	Chattahoochee	A. H. Curtiss	do	0.5723		787	787	1052
	734	do	do	do	do	0.5354		814	849	1031
	800	do	Saint John's river	do	Sandy loam	0.5558		787	781	755
	800	do	do	do	do	0.5385		904	888	806
	924	do	Chattahoochee	C. Mohr	Alluvial	0.4997		868	868	888
	1249	Tennessee	Wilson county	A. E. Baird		0.7373		588	688	750
	1250	do	do	do		0.6029		444	474	394
	1251	do	do	do		0.5671		488	542	469
	1252	do	do	do		0.6398		718	697	860
340. <i>Taxodium distichum</i> <i>Bald Cypress. Black Cypress. Red Cypress. White Cypress. Deciduous Cypress.</i>	535	Alabama	Stockton	C. Mohr	Alluvial	0.4907		1163	1122	675
	535	do	do	do	do	0.4923		1110	1163	666
	741	Florida	Chattahoochee	A. H. Curtiss		0.4167		904	904	687
	741	do	do	do		0.4334		939	939	698
341. <i>Sequoia gigantea</i> <i>Big Tree.</i>	657	California	Tulare county	G. Engelmann and C. S. Sargent.	Granite	0.3426		595	595	511
	657	do	do	do	do	0.3506		425	444	469
	666	do	do	do	do	0.2687		359	315	401
342. <i>Sequoia sempervirens</i> <i>Redwood.</i>	673	do	Russian river	C. S. Sargent		0.4103		574	558	637
	673	do	do	do		0.		787	751	708

UNITED STATES UNDER TRANSVERSE STRAIN—Continued.

DEFLECTION, IN MILLIMETERS, UNDER A PRESSURE, IN KILOGRAMS, OF—													Ultimate strength: transverse pressure.	Remarks.	Office number.
50	100	150	200	0 (set.)	200	250	300	350	400	450	500	550			
6.0	12.0	18.0	24.7	1.0	24.5	31.5	41.0	60.7					363	Square break; splintered.....	969
5.5	10.0	15.0	21.0	1.0	21.2	29.5							297	Crushed at center bearing; square break.....	969
4.0	8.0	12.0	16.0	0.4	16.0	20.0	25.5	31.7	42.0				435	Crushed at center bearing; square break, splitting in axis.....	968
4.4	8.2	12.5	17.0	0.5	17.5	21.5	28.0	36.0					397	do.....	963
4.2	8.5	12.5	17.2	0.4	17.0	22.0	28.5	39.5					372	do.....	964
4.5	8.7	13.5	17.6	0.2	18.0	22.5	29.0	37.0	53.0				400	Shattered.....	964
3.7	8.0	11.5	15.4	0.0	15.0	19.0	24.0	31.5					397	do.....	964
6.0	11.6	17.5	24.2	1.0	24.4	31.5							298	do.....	1000
6.4	13.5												120	Specimen cross-grained; started at knot.....	1000
5.4	10.5	15.6	21.6	0.6	22.4	29.0	37.0						350	Square break; shattered.....	701
4.0	8.0	12.0	16.4	0.4	16.5	21.0	28.0						349	Crushed at center bearing; square break on tension side, splitting in axis.....	707
3.0	6.5	9.6	12.7	0.2	13.0	16.0	20.0	25.4	32.0				439	do.....	707
5.0	9.2	13.5	18.0	1.0	18.0	22.5	28.0	33.0	40.0				444	Square break on tension side, splitting in axis; shattered from end to end.....	675
4.5	9.0	13.5	17.6	0.3	18.0	22.5	27.5	34.0	42.0				449	do.....	675
10.3	21.4	35.0	49.0	5.0	49.0								222	Shattered.....	691
9.5	18.0	27.8	40.0	3.4	41.7	53.0							281	do.....	691
8.0	16.5	25.5	35.0	1.5	35.7	46.4							271	Specimen cross-grained; shattered.....	1100
10.7	24.0												146	Specimen cross-grained; splinter on corner.....	1100
8.0	15.5	24.0	32.5	1.0	32.4	42.5	55.0						334	Shattered.....	692
8.2	16.4	25.0	34.0	1.1	34.0	44.0	58.5						315	do.....	692
6.5	13.3	20.5	28.0	1.6									200	Specimen cross-grained; shattered.....	1102
9.6	19.5	29.0	40.6	1.9	40.0	52.0							299	Shattered.....	14
9.5	19.0	29.0	40.0	2.0	40.5	53.0	70.0						317	do.....	14
8.2	16.0	25.0	33.2	1.2	33.5	43.0	56.0	73.5					363	do.....	327
8.0	15.5	24.0	31.0	0.8	32.0								240	do.....	327
6.2	12.4	18.2	24.5	0.7	24.5	31.7	38.5	47.2	59.5				449	0.25 sap-wood; square break on tension side, splitting in axis.....	734
6.0	11.5	17.0	23.0	0.4	23.0	29.0	35.7	45.5	54.5				440	0.5 sap-wood; square break; split end to end.....	734
6.2	12.5	19.2	26.5	1.1	26.7	35.0	43.5						322	Square break at knot; split to end.....	800
5.4	11.0	16.5	21.7	0.3	22.0	28.2	35.0						344	Square break on tension side, splitting in axis.....	800
5.5	11.0	16.1	21.5	0.2	21.8	27.7	34.5	43.5					379	Square break; split parallel to pressure.....	924
8.3	14.2	21.4	28.0	0.8	29.0	34.7	43.5						320	Specimen not seasoned; shattered.....	1249
11.0	20.6	31.0											163	Specimen not seasoned; cross-grained.....	1250
10.0	18.0	26.0	34.0	1.5									200	Specimen not seasoned; cross-grained; split to end.....	1251
6.8	14.0	21.5	28.7	1.3	30.0	36.5	45.2	58.0					381	Specimen not seasoned; shattered.....	1252
8.4	16.4	25.6	37.0	4.1	39.0								222	Specimen not seasoned; square break on tension side, splitting in axis.....	1253
8.7	16.2	26.0	36.0	2.4	37.0	43.0	66.0						300	Specimen not seasoned; shattered.....	1254
4.2	8.7	12.9	17.5	0.5	18.2	24.5							288	Square break.....	535
4.4	8.4	13.0	17.9	0.6	18.2	24.0							284	Crushed at center bearing with flakes on tension side.....	535
5.4	10.8	16.7	23.5	1.0	24.5	33.0							293	Square break on tension side, splitting in axis.....	741
5.2	10.4	16.0	22.8	1.1	23.0	30.8							298	Broke with coarse splinters.....	741
8.2	16.4	25.0	34.0	1.2	35.0								218	Square break on tension side, splitting in axis.....	657
11.5	22.0	34.0	49.0	3.6									206	Square break.....	657
13.6	31.0	54.0											171	Sap-wood; shattered.....	666
8.5	17.5	27.0	38.0	2.0	38.5	52.5							272	Square break; split to end.....	673
6.2	13.0	19.0	26.2	0.5	26.5	34.5							300	do.....	673

TABLE III.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Specific gravity of the air-dried specimen.	Direction of grain.	COEFFICIENT OF ELASTICITY.		Modulus of rupture.
								First deflection.	Second deflection.	
342. <i>Sequoia sempervirens</i> —continued.	710	California	Santa Cruz	Turner, Kennedy & Shaw.		0.4443		697	697	654
	711	do	do	do		0.4525		751	751	581
	711	do	do	do		0.4410		542	558	501
	712	do	Mendocino county	J. Kentfield & Co.		0.3239		542	568	394
	712	do	do	do		0.3239		651	651	586
	713	do	do	do		0.3841		814	849	558
	713	do	do	do		0.3648		751	708	633
	714	do	do	do		0.5289		751	751	734
	714	do	do	do		0.5243		651	666	804
	715	do	do	do		0.5533		651	610	295
343. <i>Taxus brevifolia</i> . <i>Yew.</i>	978	Oregon	Portland	G. Engelmann and C. S. Sargent.	Moist, rich	0.6743		814	849	1148
	978	do	do	do	do	0.7110		697	673	1010
345. <i>Torreya taxifolia</i> . <i>Stinking Cedar. Savin.</i>	62	Florida	Chattahoochee	C. S. Sargent	Alluvial	0.5968		651	651	945
	277	do	do	A. H. Curtiss	Calcareous	0.4693		872	921	844
	277	do	do	do	do	0.4639		939	913	872
346. <i>Torreya Californica</i> . <i>California Nutmeg. Stinking Cedar.</i>	651	California	Marin county	G. R. Vasey	Stony	0.5068		407	450	469
	651	do	do	do	do	0.5135		444	444	698
347. <i>Pinus Strobus</i> . <i>White Pine. Weymouth Pine.</i>	1	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	0.4301		751	849	623
	1	do	do	do	do	0.4090		921	930	682
	222	Vermont	Charlotte	C. G. Pringle	Wet, swampy	0.6204		574	592	804
	777	New Brunswick		Intercolonial railway.		0.3671		651	638	424
	777	do		do		0.3932		1085	1039	635
	788	do	Bridgeton	Ed. Sinclair		0.4022		1085	1062	654
	788	do	do	do		0.4088		1039	1039	687
	789	Province of Quebec	Amqui	A. Grant		0.3698		763	751	170
	789	do	do	do		0.3671		775	751	327
	797	do		Grand Trunk railway.		0.4092		976	976	330
	797	do		do		0.3938		842	849	621
	1044	Massachusetts	Reading	J. Robinson	Drift	0.3710		688	697	390
348. <i>Pinus monticola</i> . <i>White Pine.</i>	975	British Columbia	Hastings' saw-mill, Burrard inlet.	G. Engelmann and C. S. Sargent.		0.4323		1163	1191	684
	987	Oregon	Cascade mountains	C. S. Sargent	Moist loam	0.3682		751	740	627
	987	do	do	do	do	0.3699		957	921	616
349. <i>Pinus Lambertiana</i> . <i>Sugar Pine.</i>	638	California	Saw-mill, Strawberry valley.	G. Engelmann and C. S. Sargent.		0.4301		888	888	687
	668	do		G. R. Vasey		0.3927		740	740	577
	668	do		do		0.3971		814	814	656
	730	do	Laassen's peak	Sierra Lumber Company.		0.3363		763	787	541
	730	do	do	do		0.3307		740	740	530
350. <i>Pinus flexilis</i> . <i>White Pine.</i>	819	Colorado	Forest City	T. S. Brandegee	Gravelly	0.4602		751	787	687
	819	do	do	do	do	0.4642		651	697	635
	819	do	do	do	do	0.4733		787	835	776
	913	Nevada	Danville	A. Triple	do	0.4664		381	381	396
351. <i>Pinus albicaulis</i>	992	British Columbia	Silver Mountain valley, Fraser river.	G. Engelmann and C. S. Sargent.		0.5066		370	348	464
	992	do	do	do		0.4587		679	673	701

UNITED STATES UNDER TRANSVERSE STRAIN—Continued.

DEFLECTION, IN MILLIMETERS, UNDER A PRESSURE, IN KILOGRAMS, OF—													Ultimate strength: transverse pressure.	Remarks.	Office number.
50	100	150	200	0 (set.)	200	250	300	350	400	450	500	550			
7.0	14.0	21.2	29.5	1.0	30.0	39.0							279	Square break	710
6.5	13.0	19.4	26.0	0.5	26.4	33.5							248	Square break; shattered	711
9.0	17.5	26.6	36.3	1.7	37.0								252	Square break	711
9.0	17.2	26.6											168	Shattered with flakes on tension side	712
7.5	15.0	22.4	32.0	1.5	33.5								250	Square break	712
6.0	11.5	17.2	23.5	0.7	24.0								238	Square break on tension side, splitting in axis	712
6.5	13.8	20.0	27.7	1.0	28.4	38.0							270	Square break	712
6.5	13.0	20.0	26.6	0.6	27.0	34.0	50.0						312	Shattered from end to end	714
7.5	14.7	22.0	29.5	0.9	30.0	38.0	48.0						343	Square break	714
7.5	16.0												126	Specimen with curly grain; square break on tension side, splitting in axis.	715
6.0	11.5	17.8	24.0	0.9	24.0	31.5	38.0	47.0	59.3	75.0			490	Square break on tension side, splitting in axis; shattered	978
7.0	14.5	21.5	29.7	1.7	30.0	39.0	49.0	61.6	84.0				431	do	978
7.5	15.0	22.5	31.0	2.4	31.0	41.0	52.0	66.5	92.5				403	Shattered	63
5.6	10.6	16.5	22.8	0.4	22.5	28.7	35.5	47.0					360	Square break on tension side, splitting in axis; shattered	277
5.2	10.7	16.0	22.6	0.5	22.2	28.5	35.5	46.2					372	do	277
12.0	22.7	34.5	50.5										200	Broke at small knot at point of compression	651
11.0	22.0	33.0	46.5	3.1	49.5	68.0							298	Square break on tension side, splitting in axis; also broke at knot near the end.	651
6.5	11.5	17.2	23.8	1.2	24.2	32.0							266	Square break with scale on tension side	1
5.3	10.5	16.4	22.8	0.5	23.2	31.0							291	Square break on tension side, splitting in axis; shattered	1
8.5	16.5	25.0	34.3	2.3	35.0	46.5	62.0						343	Square break; shattered	222
7.5	15.3	22.6											181	Cross-grained	777
4.5	9.4	14.0	20.0	1.0	20.4	29.0							271	Started at knot	777
4.5	9.2	14.2	19.8	0.6	21.0	29.0							279	Crushed at center bearing; square break	788
4.7	9.4	14.8	20.5	0.3	21.0	28.0							293	do	788
6.4	13.0	20.7	30.0	2.0	31.0								243	Square break	789
6.3	13.0	20.0	32.3	3.6	33.5								225	do	789
5.0	10.0	15.4	21.6	1.0	21.8	30.0							397	Square break; split to one end	797
5.8	11.5	17.8	25.2	1.8	25.2	34.0							265	Square break	797
7.1	14.0	22.4											167	Specimen cross-grained; broke at large knot	1044
5.6	11.0	17.0	24.0	1.3	24.6								249	Crushed at center bearing; splinter on corner	1044
4.2	8.2	12.5	17.1	0.7	17.5	24.0							292	Crushed at center bearing; broke with fine splinters	975
6.5	13.2	20.2	32.1	3.8	34.2								225	Square break	987
5.1	10.6	16.2	23.5	1.3	24.0	35.0							263	do	987
5.5	11.0	17.0	23.0	0.6	23.0	30.2							293	Square break on tension side, splitting in axis	638
6.6	13.2	20.3	29.9	1.2	29.0								246	Shattered	668
6.0	12.0	18.2	26.0	1.0	26.0	35.0							280	Square break	668
6.4	12.4	18.7	28.0	2.4	29.5								231	Crushed at center bearing; square break	730
6.6	13.2	20.6	32.0	3.0	32.8								226	Square break	730
6.5	12.4	18.9	26.0	1.0	27.0	35.0							293	Square break; shattered	819
7.5	14.0	21.6	31.0	2.0	31.4	46.2							271	do	819
6.2	11.7	17.5	24.0	0.8	24.2	32.0	43.0						331	Specimen cross-grained; square break on tension side, splitting in axis.	819
12.8	25.6	41.5											169	Square break at knot	913
13.2	28.0	46.0											198	Square break; shattered	993
7.2	14.5	22.5	31.0	1.2	32.0	42.0							299	do	993

TABLE III.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Specific gravity of the air-dried specimen.	Direction of grain.	COEFFICIENT OF ELASTICITY.		Modulus of rupture.
								First deflection.	Second deflection.	
352. <i>Pinus reflexa</i> <i>White Pine.</i>	661	Arizona	Santa Rita mountains.	G. Engelmann and C. S. Sargent.		0.5261		814	849	689
	661	do	do	do		0.5392		839	976	851
353. <i>Pinus Parryana</i> <i>Piñon. Nut Pine.</i>	656	California	San Diego county...	G. R. Vasey.....		0.6862		390	849	347
	656	do	do	do		0.6342		444	407	504
355. <i>Pinus edulis</i> <i>Piñon. Nut Pine.</i>	397	Colorado.....	Cañon City	E. Weston	Gravelly	0.6704		429	421	448
356. <i>Pinus monophylla</i> <i>Piñon. Nut Pine.</i>	882	Utah	Lewiston	M. E. Jones.....	Rocky	0.5894		421	415	281
	915	Nevada.....	Danville.....	A. Triple	Gravelly.....	0.6579		519	454	295
357. <i>Pinus Balfouriana</i>	631	California	Scott mountains ...	G. Engelmann and C. S. Sargent.	Rocky	0.5623		542	528	291
	631	do	do	do	do	0.5714		651	660	558
357. <i>Pinus Balfouriana</i> , var. <i>aristata</i> <i>Foxtail Pine. Hickory Pine.</i>	821	Colorado.....	Forest City	T. S. Brandegee...		0.5113		568	574	469
	821	do	do	do		0.4987		888	888	771
	821	do	do	do		0.4811		842	763	689
	914	Nevada.....	Prospect mountain..	A. Triple	Rocky	0.6240		651	634	694
358. <i>Pinus resinosa</i>	815	Michigan	Hersey	W. J. Beal		0.5482		1356	1395	595
	815	do	do	do		0.5511		1320	1375	735
	785	New Brunswick ..	Bridgeton	Ed. Sinclair		0.4951		1039	1039	806
	785	do	do	do		0.4191		888	849	670
	1074	Vermont.....	Charlotte	C. G. Pringle.....		0.4886		1221	1110	773
	1075	do	do	do		0.4789		976	1067	703
	1076	do	do	do		0.5221		1039	1062	813
	1076	do	do	do		0.5164		1163	1221	799
359. <i>Pinus Torreyana</i>	996	California	San Diego county...	G. Engelmann	Sandy	0.5471		634	630	809
	996	do	do	do	do	0.5800		542	454	703
360. <i>Pinus Arizonae</i>	1154	Arizona	Santa Rita mountains.	C. G. Pringle.....	Rocky	0.6570		768	746	703
	1154	do	do	do	do	0.7913		814	835	553
	1155	do	do	do	do	0.4749		814	857	598
	1155	do	do	do	do	0.4696		872	888	769
	1156	do	do	do	do	0.4250		729	794	647
361. <i>Pinus ponderosa</i>	619	Dakota	Deadwood	Robert Douglas...	Gravelly.....	0.5034		1221	1285	881
	626	Oregon	Saw-mill, Ashland ..	G. Engelmann and C. S. Sargent.		0.4788		1356	1252	930
	630	California	Strawberry valley ..	do	Low, wet, swampy	0.4814		191	190	443
	630	do	do	do	do	0.4748		266	264	382
	632	do	Saw-mill, Strawberry valley.	do		0.5502		1221	1221	905
	636	do	do	do		0.4429		814	814	780
	689	do	Saw-mill, San Bernardino.	W. G. Wright		0.5079		1285	1302	1041
	718	Montana.....	Saw-mill, Missoula ..	S. Watson		0.4502		1085	1110	766
	718	do	do	do		0.4750		1163	1163	818
	731	California.....	Lassen's peak	Sierra Lumber Company.		0.4795		1221	1221	783
	731	do	do	do		0.4803		1163	1163	795
	907	Colorado.....	Cañon City	E. Weston		0.5206		813	305	391
	910	do	do	do		0.5277		271	247	438
362. <i>Pinus Jeffreyi</i>	633	California	Scott mountains ...	G. Engelmann and C. S. Sargent.	Dry, gravelly	0.5678		498	506	585
	633	do	do	do	do	0.5758		417	394	316
	667	do	Saw-mill, San Bernardino.	W. G. Wright		0.5948		1221	1320	902
	667	do	do	do		0.6329		1637	1479	1171

UNITED STATES UNDER TRANSVERSE STRAIN—Continued.

DEFLECTION, IN MILLIMETERS, UNDER A PRESSURE, IN KILOGRAMS, OF—													Ultimate strength; transverse pressure.	Remarks.	Office number.
50	100	150	200	0 (set.)	200	250	300	350	400	450	500	550			
6.0	11.5	17.5	24.2	1.0	25.0	32.0							294	Specimen cross-grained; failed with long split.....	661
5.2	10.0	15.0	20.0	0.8	20.2	25.6	33.0	43.0					363	do	661
12.5	28.0												148	Broke at knot near end	656
11.0	24.0	37.2	55.0	7.0	59.2								215	do	656
11.4	23.2	39.2											191	Broke at knot.....	397
11.0	23.5												120	Specimen cross-grained; broke at knot	882
9.4	21.5												126	do	915
9.0	18.5												124	do	631
7.5	14.8	23.2	32.5	3.2	34.5								238	Square break	631
8.6	17.0	25.7	36.0	2.5									200	do	821
5.5	11.0	17.3	23.5	0.9	23.5	31.0	40.0						329	0.25 sap-wood; specimen cross-grained.....	821
5.8	12.8	19.0	26.3	1.1	27.0	36.0							290	Square break with split at end	821
7.5	15.4	24.0	34.2	3.0	35.7	46.2							296	Shattered	914
3.6	7.0	10.7	14.5	0.3	14.6	18.8	24.0	31.0					382	Crushed at center bearing; square break on tension side, splitting in axis.....	315
3.7	7.1	10.5	14.0	0.4	14.0	17.8	22.5	28.5					399	Crushed at center bearing; square break	315
4.7	9.4	14.0	19.2	0.8	19.5	25.7	35.0						344	Specimen cross-grained; shattered.....	785
5.5	11.5	16.8	23.5	0.8	24.0	32.0							286	Square break	785
4.0	8.8	13.0	17.6	0.3	17.6	23.0	30.0						330	Square break on tension side, splitting in axis.....	1074
5.0	9.7	15.0	20.0	0.6	20.3	26.0	40.2						300	Square break on tension side, splitting in axis; crushed at center bearing.....	1075
4.7	9.2	14.0	19.5	0.7	19.5	25.0	33.0						347	Square break; crushed at center bearing	1076
4.2	8.0	12.3	16.6	0.5	17.0	21.5	28.4						341	Broke with long scale	1076
7.7	15.5	22.4	30.8	1.9	31.5	41.5	55.5						345	Shattered	996
9.0	21.5	34.6	50.2	7.5	51.7	73.0							300	do	996
6.4	13.1	20.0	27.5	1.0	28.2	36.0	48.0						300	Square break on tension side, splitting in axis.....	1154
6.0	11.7	17.9	25.0	1.5	25.3								236	Specimen cross-grained; broke at knot	1154
6.0	11.4	16.6	23.2	1.1	24.0	31.0							253	Square break on tension side, splitting in axis	1155
5.6	11.0	16.6	22.1	0.8	22.1	29.0	39.0						328	Square break on tension side, splitting in axis; shattered	1155
6.7	12.3	19.0	26.5	1.1	27.0	35.0							276	Square break; split to one end.....	1156
4.0	7.6	11.5	15.5	0.3	15.3	19.5	25.0	31.5					376	0.75 sap-wood; crushed at center bearing; square break on tension side, splitting in axis.....	619
3.6	7.8	11.6	15.5	0.2	15.5	20.0	24.2	30.5					397	Square break; shattered.....	626
25.6	51.5	80.0											189	Broken at end; shattered	630
18.4	37.0	57.0											163	Short break; shattered	630
4.0	8.0	12.0	16.0	0.4	16.2	20.0	25.0	31.6					386	Square break on tension side, splitting in axis	632
6.0	12.0	18.0	24.8	1.0	25.3	33.5	44.0						333	Square break	636
3.8	7.5	11.2	14.7	0.2	15.0	18.5	23.0	28.0	35.0				444	Square break on tension side, splitting in axis.....	689
4.5	8.8	13.3	18.0	0.4	18.0	23.0	30.7						327	do	718
4.2	8.4	12.7	17.0	0.5	17.0	22.4	28.0						340	Square break; split to one end.....	718
4.0	8.0	12.5	16.8	0.5	16.7	21.5	28.5						334	Square break on tension side, splitting in axis	731
4.2	8.4	12.4	17.0	0.6	17.5	21.5	27.5						339	Specimen cross-grained	731
15.6	32.0	54.0											167	Sap-wood; specimen cross-grained; broke at knot.....	907
18.0	39.5	71.0											187	Cross-grained	910
9.8	19.3	30.0	40.0	3.8	41.8	55.5							250	Square break on tension side, splitting in axis; shattered	633
11.7	24.8												135	Square break at knot	633
4.0	7.4	11.0	15.0	0.5	15.0	19.0	24.0	32.5					385	Crushed at center bearing; broke with fine splinters	667
3.0	6.6	9.6	13.0	0.4	13.2	16.0	20.0	24.5	30.0				500	Square break on tension side, splitting in axis	667

TABLE III.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Specific gravity of the air-dried specimen.	Direction of grain.	COEFFICIENT OF ELASTICITY.		Modulus of rupture.
								First deflection.	Second deflection.	
363. <i>Pinus Chihuahuana</i>	664	Arizona	Santa Rita mountains.	G. Engelmann and C. S. Sargent.	Dry, gravelly	0.5801		626	651	905
	664	do	do	do	do	0.5883		872	800	750
364. <i>Pinus contorta</i>	997	British Columbia.	Vancouver's island.	do	do	0.6456		1806	1775	1048
	997	do	do	do	do	0.6198		1395	1395	937
365. <i>Pinus Murrayana</i>	298	Colorado	Forest City	T. S. Brandegee	Moist, sandy loam.	0.4551		542	501	286
	298	do	do	do	do	0.4265		525	514	347
	563	do	do	C. S. Sargent	do	0.4546		976	976	743
	625	California	Scott mountains	G. Engelmann and C. S. Sargent.	do	0.4600		857	888	675
366. <i>Pinus Sabiniana</i>	644	do	Contra Costa county	G. R. Vasey	Gravelly	0.5460		514	528	738
	644	do	do	do	do	0.5426		651	642	820
367. <i>Pinus Coulteri</i>	1157	do	San Bernardino	W. G. Wright	Dry, gravelly	0.4443		1221	1221	818
	1157	do	do	do	do	0.4322		1062	1062	703
368. <i>Pinus insignis</i>	676	do	Monterey	G. R. Vasey	Gravelly loam	0.4835		888	872	670
	676	do	do	do	do	0.5095		1136	1085	809
369. <i>Pinus tuberculata</i>	576	do	Mount Shasta	G. Engelmann and C. S. Sargent.	Gravelly	0.4071		854	362	298
	576	do	do	do	do	0.3901		519	496	525
370. <i>Pinus Taeda</i>	82	Florida	Duval county	A. H. Curtiss	Moist, sandy	0.0068		1627	1627	996
	82	do	do	do	do	0.6147		1744	1684	1402
	355	Alabama	Cottage Hill	C. Mohr	Low, rich	0.5914		688	638	790
	355	do	do	do	do	0.5846		838	864	666
	388	North Carolina ..	Wilmington	E. Kidder	Loam	0.5600		1395	1302	1012
	388	do	do	do	do	0.4306		888	888	562
	389	do	do	do	do	0.4675		921	939	773
	389	do	do	do	do	0.4937		1136	1085	795
371. <i>Pinus rigida</i>	13	Massachusetts	Arnold Arboretum ..	C. S. Sargent	Drift	0.5670		697	651	797
	13	do	do	do	do	0.5943		775	769	851
	1046	do	North Reading	J. Robinson	do	0.4371		542	537	647
	1046	do	do	do	do	0.5140		376	368	663
372. <i>Pinus serotina</i>	88	Florida	Duval county	A. H. Curtiss	Moist, sandy loam.	0.7592		1136	1103	1158
	88	do	do	do	do	0.7518		1221	1177	1172
373. <i>Pinus inops</i>	621	South Carolina	Aiken	H. W. Ravenel	Dry, sandy	0.5403		405	465	682
	622	do	do	do	do	0.5537		697	697	562
	1169	Indiana	New Albany	M. J. Robinson	do	0.5610		751	781	855
	1169	do	do	do	do	0.5703		543	528	499
	1172	New Jersey	Mount Holly	S. P. Sharples	Clay	0.5920		452	428	722
	1172	do	do	do	do	0.7088		373	358	637
374. <i>Pinus clausa</i>	279	Florida	Apalachicola	A. H. Curtiss	Dry, sandy barren.	0.5341		542	528	429
	279	do	do	do	do	0.5218		595	558	574
375. <i>Pinus pungens</i>	321	Virginia	Wytheville	H. Shriver	Clay	0.5180		679	630	771
	321	do	do	do	do	0.5300		976	976	680
376. <i>Pinus muricata</i>	671	California	Marin county	G. R. Vasey	Gravelly	0.5873		1039	1065	1005
	671	do	do	do	do	0.5883		1221	1302	1059

UNITED STATES UNDER TRANSVERSE STRAIN—Continued.

DEFLECTION, IN MILLIMETERS, UNDER A PRESSURE, IN KILOGRAMS, OF—													Ultimate strength: transverse pressure.	Remarks.	Office number.
50	100	150	200	0 (set.)	200	250	300	350	400	450	500	550			
7.8	15.0	23.0	31.5	1.6	32.0	42.0	55.0	71.0					388	Square break on tension side, splitting in axis	664
5.0	12.2	18.2	25.7	1.3	25.8	32.4	43.7						324	Failed from large splinter on tension side	664
2.7	5.5	8.2	11.5	0.2	11.6	14.4	18.0	21.6	23.0				447	Crushed at center bearing; failed from tension side	997
3.5	7.0	10.2	14.0	0.3	14.0	18.0	21.3	27.0					400	Failed from large splinter on corner	997
9.0	19.5												122	Broke at knot	293
9.3	19.0												148	Square break at knot	293
5.0	10.0	15.0	20.7	0.5	21.0	27.0	37.5						317	Square break; splintered	563
5.7	11.0	16.8	23.7	1.0	23.5	32.0							288	Square break	625
5.0	10.0	15.0	20.2	0.5	20.6	27.5	35.5						328	do	625
9.5	18.5	29.0	43.0	4.6	44.5	61.0	95.0						315	Square break on tension side, splitting in axis; shattered	644
7.5	15.2	22.5	32.0	2.0	33.0	44.0	62.5						350	do	644
4.0	8.0	12.2	16.2	0.2	16.6	21.3	27.5						349	do	1157
4.6	9.2	14.6	20.0	0.5	20.0	27.0							300	Broke at knot	1157
5.5	11.2	17.0	23.5	1.0	23.7	31.0							286	Specimen cross-grained; shattered	676
4.3	9.0	14.3	19.6	0.4	19.8	26.0	33.4						345	do	676
13.8	27.0												125	Sap-wood; square break	576
9.4	19.7	30.6	45.0	4.8	48.0								224	do	576
3.0	6.0	9.0	12.0	0.4	12.3	15.6	19.0	23.0	29.0				426	Crushed at center bearing; square break on tension side, splitting in axis	82
2.8	5.8	8.7	11.7	0.2	11.8	14.5	17.6	21.3	25.7	30.2	37.2		624	do	82
7.1	15.3	23.3	33.0	2.5	33.0	44.0	57.5						337	Long split at one end	355
5.5	11.3	17.6	24.0	1.5	24.5	32.6							284	Long split	355
3.5	7.5	11.0	15.0	0.3	15.0	18.6	23.5	29.2	38.0				432	0.33 sap-wood; square break	388
5.5	11.0	17.0	24.0	1.1	24.0								240	Square break on tension side, splitting in axis; shattered	388
5.3	10.4	16.0	22.4	1.0	22.5	29.4	38.5						330	do	389
4.3	9.0	13.0	18.0	0.5	18.0	23.0	30.2						339	do	389
7.0	15.0	23.6	33.5	3.0	34.0	45.8	63.5						340	Square break on tension side, splitting in axis	13
6.3	12.7	20.0	28.5	2.4	29.2	37.5	50.0	68.0					363	do	13
9.0	18.2	27.5	41.6	4.0	42.7	60.0							276	Shattered	1046
13.0	26.5	40.0	60.0	8.0	63.0	88.0							283	Square break on tension side, splitting in axis	1046
4.3	8.4	12.3	17.0	0.6	17.0	21.5	27.0	33.0	39.0	48.5			494	Broke with coarse splinters	83
4.0	8.3	12.6	17.0	0.5	17.4	22.2	27.5	34.5	42.5	52.6			500	do	83
10.5	21.0	32.3	49.9	6.0	51.0	71.0							291	Square break on tension side, splitting in axis	621
7.0	14.0	22.0	31.0	0.5	33.0								240	Oblique fracture; started at knot	622
6.5	12.5	19.5	27.0	1.6	27.5	35.8	47.6	63.5					365	Shattered	1169
9.0	18.5	29.5	43.0	5.3	45.0								213	Square break	1169
10.8	22.8	36.0	50.0	6.0	52.5	71.0	114.0						308	Specimen cross-grained; shattered	1172
13.1	27.3	41.0	60.0	6.3	63.5	88.0							272	do	1172
9.0	18.5	32.0											183	Specimen cross-grained; started at knot	279
8.2	17.5	28.0	40.0	3.5	41.2								245	Shattered	279
7.2	15.5	24.0	35.0	3.5	36.5	47.0	64.0						329	do	321
5.0	10.0	16.5	23.2	1.3	23.2	30.8							290	Specimen cross-grained; broke near end	321
4.7	9.0	13.2	18.5	0.3	18.0	23.2	29.0	36.0	47.0				429	Square break	671
4.0	7.5	11.8	15.0	0.0	15.2	19.0	24.0	28.5	35.0	44.0			452	do	671

TABLE III.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Specific gravity of the air-dried specimen.	Direction of grain.	COEFFICIENT OF ELASTICITY.		Modulus of rupture.
								First deflection.	Second deflection.	
377. <i>Pinus mitis</i> <i>Yellow Pine. Short-leaved Pine. Spruce Pine. Full Pine.</i>	278	Florida	Chattahoochee	A. H. Curtiss	Clay	0.6524		1221	1252	874
	278	do	do	do	do	0.6490		1395	1415	1055
	319	Louisiana	Amite	C. Mohr	Sandy loam	0.6516		1479	1458	1188
378. <i>Pinus glabra</i> <i>Cedar Pine. Spruce Pine. White Pine.</i>	142	South Carolina	Bonneau's Depot	H. W. Ravenel	Rich upland	0.4850		634	660	460
	142	do	do	do	do	0.4035		642	660	570
	544	Mississippi	Gainesville	C. Mohr	Low, sandy	0.3535		220	208	262
	544	do	do	do	do	0.3500		244	238	240
	544	do	do	do	do	0.4268		244	233	431
	764	Florida	Chattahoochee	A. H. Curtiss	Low, wet	0.5089		561	542	766
	764	do	do	do	do	0.4713		595	592	683
379. <i>Pinus Banksiana</i> <i>Gray Pine. Scrub Pine. Prince's Pine.</i>	394	Michigan	Baldwin	W. J. Beal	Low, sandy	0.4912		574	574	574
	394	do	do	do	do	0.4917		814	709	649
	780	New Brunswick		Intercolonial railway.		0.5191		1320	1356	947
	780	do		do		0.4297		872	939	570
	879	do		do		0.4839		976	930	448
	879	do		do		0.5129		1039	1085	722
380. <i>Pinus palustris</i> <i>Long-leaved Pine. Southern Pine. Georgia Pine. Yellow Pine. Hard Pine.</i>	81	Florida	Duval county	A. H. Curtiss	Sandy loam	0.9375		1628	1628	1160
	81	do	do	do	do	0.8125		1526	1575	1266
	81	do	do	do	do	0.6950		976	1062	820
	81	do	do	do	do	0.9822		1628	1628	1240
	85	do	do	do	Moist, sandy	0.6164		814	888	820
	85	do	do	do	do	0.6632		1628	1575	1036
	85	do	do	do	do	0.7568		1039	1085	937
	85	do	do	do	do	0.8800		642	630	935
	243	do	Saw-mill, Saint John's river.	do		0.7087		1953	1775	1266
	243	do	do	do		0.7066		2123	1713	1289
	357	Alabama	Cottage Hill	C. Mohr	Sandy loam	0.6643		1628	1550	1172
	357	do	do	do	do	0.8410		1628	1713	1472
	358	do	Citronelle	do	do	0.8906		1436	1395	1289
	358	do	do	do	do	0.8706		1628	1775	1392
	359	do	Chunchula	do	do	0.7986		1628	1684	1233
	359	do	do	do	do	0.8728		1628	1628	1397
	360	do	do	do	do	0.7654		1684	1776	1380
	360	do	do	do	do	0.7242		1221	1302	1170
	361	do	do	do	do	0.7585		2035	1053	1287
	361	do	do	do	do	0.7480		2035	2035	1322
	384	Florida	Saw-mill, Cedar Keys.	A. H. Curtiss		0.6005		957	967	642
	384	do	do	do		0.6438		1628	1655	1064
	385	do	do	do		0.7038		1221	1221	1045
	390	North Carolina	Wilmington	E. Kidder		0.6739		1808	1776	1137
	390	do	do	do		0.6293		1628	1550	1062
	1096	Alabama	Mobile	C. Mohr		0.6375		1085	1163	1165
	1096	do	do	do		0.6852		1479	1459	1095
381. <i>Pinus Cubensis</i> <i>Slash Pine. Swamp Pine. Bastard Pine. Meadow Pine.</i>	84	Florida	Duval county	A. H. Curtiss	Moist, sandy	0.6934		1479	1302	1179
	84	do	do	do	do	0.6646		1163	1191	1048
	84	do	do	do	do	0.8314		1479	1479	1181
	356	Alabama	Cottage Hill	C. Mohr	do	0.7716		1878	1953	1279
	356	do	do	do	do	0.7710		1953	1958	1465

UNITED STATES UNDER TRANSVERSE STRAIN—Continued.

DEFLECTION, IN MILLIMETERS, UNDER A PRESSURE, IN KILOGRAMS, OF—														Ultimate strength: transverse pressure.	Remarks.	Office number.
50	100	150	200	C (set.)	200	250	300	350	400	450	500	550				
4.0	7.8	11.2	15.1	0.2	15.2	19.0	24.0	29.2					373	Specimen cross-grained; started at knot	278	
3.5	6.9	10.8	14.2	0.3	14.2	18.0	22.0	27.0	32.0	45.0			450	Square break on tension side, splitting in axis	278	
3.3	6.7	10.0	13.1	0.0	13.2	16.5	20.2	25.0	29.5	34.0	43.5		507	do	319	
7.7	14.8	23.2	32.0	3.0	33.2								200	Specimen cross-grained; broke with long split with grain	142	
7.6	14.8	22.5	32.0	2.4	32.2								243	Square break on tension side, splitting in axis	142	
22.2	47.0												112	do	544	
20.0	41.0												145	Shattered	544	
20.0	42.0	70.5											184	Square break at knot	544	
8.7	18.0	29.0	42.0	5.2	43.0	60.0	89.0						327	Shattered	764	
8.2	16.5	26.0	37.0	3.8	38.0	51.0							270	do	764	
8.5	17.0	27.0	39.0	5.0	40.2								245	Square break	394	
6.0	12.7	19.6	27.5	1.4	27.6	38.0							277	Long, shattered break	394	
3.7	7.2	10.8	14.3	0.3	14.3	18.2	23.0	28.6	38.0				404	Broke with small splinters	780	
5.6	10.4	16.4	20.5	1.0	21.0								243	Broke at small knots	780	
5.0	10.5	17.0											191	Broke at knot	879	
4.7	9.0	13.7	18.4	0.4	18.4	30.0	37.0						308	Broke at small knots; shattered	879	
3.0	6.0	9.0	12.4	0.2	12.5	15.6	19.0	22.6	27.0	32.5			495	Square break on tension side, splitting in axis	81	
3.2	6.2	9.3	12.4	0.3	12.7	15.5	19.0	23.0	27.2	31.7	37.5		540	Crushed at center bearing; square break	81	
5.0	9.2	14.0	18.2	0.4	18.2	23.0	28.0						350	Square break with large splinters	81	
3.0	6.0	9.0	12.6	0.3	12.3	15.6	19.0	22.0	26.2	31.0	39.0		529	Square break on tension side, splitting in axis with large splinters ..	81	
6.0	11.0	17.0	23.2	1.5	24.0	31.5	41.5	55.7					350	Square break on tension side, splitting in axis	85	
3.0	6.2	9.2	12.3	0.3	12.3	15.5	19.0	23.6	28.2				442	Crushed at center bearing; square break	85	
4.7	9.0	13.7	18.8	1.0	18.8	24.0	31.0	37.2					400	0.5 sap-wood; splintered break	85	
7.6	15.5	23.5	32.5	1.8	33.0	42.3	55.0	70.0					399	Shattered; split to one end	85	
2.5	5.5	8.0	10.7	0.0	10.9	13.9	16.6	19.2	23.0	26.7	32.3		540	Broke with large flakes on back	243	
2.3	5.7	8.2	11.0	0.0	11.0	13.6	16.6	20.0	23.0	27.5	32.5	39.6	550	do	243	
3.0	6.3	9.2	12.3	0.3	12.5	15.5	19.0	23.0	27.0	31.0	37.5		500	Square break with flakes on back	357	
3.0	5.7	8.5	11.3	0.3	11.4	14.0	17.0	20.6	23.5	28.3	33.0	38.5	628	45 millimeters deflection with 600 kilograms; broke with flakes on back.	357	
3.4	7.0	10.0	13.4	0.4	13.7	16.8	20.6	25.5	30.3	36.0	45.0		550	Square break. (a) Boxed 1852; chipped ten years; abandoned 1861 ..	358	
3.0	5.5	8.6	11.4	0.2	11.5	14.0	17.0	23.0	27.0	32.0	38.0	46.8	594	Shattered. (a) Boxed 1852; chipped ten years; abandoned 1861 ..	358	
3.0	5.8	9.0	11.6	0.3	12.0	14.5	17.5	21.0	25.0	29.0	34.0		526	Square break with large flakes on corners. (a) Boxed 1876; chipped four years; specimen taken along chip.	359	
3.0	6.0	9.5	12.6	0.2	12.6	15.6	19.0	23.0	27.0	31.5	37.0	44.0	596	Square break on tension side, splitting in axis. (a) Boxed 1876; chipped four years; specimen taken along chip.	359	
2.9	5.5	8.4	11.4	0.0	11.4	14.0	17.0	20.0	23.0	27.0	31.5	37.0	589	Broke with thin flakes on back. (a) Boxed 1876; chipped four years; specimen taken above chip.	360	
4.0	7.5	11.0	14.6	0.4	14.6	18.4	22.5	28.0	33.0	40.0			499	Square break, somewhat shattered. (a) Boxed 1876; chipped four years; specimen taken above chip.	360	
2.4	5.0	7.2	10.0	0.0	10.0	12.5	15.0	17.8	20.7	24.0	28.0		549	Square break on tension side, splitting in axis. (a) Boxed 1878; chipped two years.	361	
2.4	4.8	7.3	9.8	0.3	10.0	12.0	15.0	17.0	20.2	23.6	28.0	34.0	564	Broke with many splinters. (a) Boxed 1878; chipped two years ..	361	
5.1	10.1	15.2	20.3	0.6	20.5	26.7							274	Specimen cross-grained; split	384	
3.0	5.9	8.5	11.5	0.2	11.6	14.2	17.5	20.8	24.5	29.0	36.5		454	Broke with flakes on back	384	
4.0	8.0	11.4	15.5	0.4	15.4	19.6	24.0	29.2	35.0				446	Broke with thick flakes on back	385	
2.7	5.5	8.5	11.0	0.0	11.2	14.2	17.0	21.0	25.0	30.5			485	do	390	
3.0	6.3	9.2	12.4	0.2	12.8	15.5	19.2	23.7	28.5	38.0			453	do	390	
4.5	8.4	12.6	16.8	0.6	16.8	21.5	26.5	32.5	39.0	48.0			497	Shattered. (a) Tree boxed eighteen or twenty years ago	1096	
3.3	6.7	9.5	13.0	0.3	13.2	16.5	20.5	24.5	29.0	35.4			487	Broke with large flake. (a) Tree boxed eighteen or twenty years ago.	1096	
3.3	7.5	11.2	15.3	0.2	15.5	19.5	24.5	30.0	35.7	44.0			499	Specimen cross-grained; split	84	
4.2	8.2	12.4	16.5	0.4	16.4	20.8	25.3	31.0	36.5				447	Specimen cross-grained; broke with large splinters	84	
3.3	6.6	10.0	13.0	0.2	13.2	16.2	20.0	24.0	28.5	33.0	39.0		504	Square break	84	
2.6	5.0	7.8	10.3	0.2	10.2	13.0	15.5	18.2	21.3	25.0	30.0		542	Slightly splintered	356	
2.5	5.0	7.3	10.0	0.3	10.0	12.0	15.0	17.5	21.0	24.0	28.0	31.5	625	39.2 millimeters deflection with 600 kilograms; broke with large flat splinters on corners.	356	

a In the manufacture of turpentine.

FOREST TREES OF NORTH AMERICA.

TABLE III.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Specific gravity of the air-dried specimen.	Direction of grain.	COEFFICIENT OF ELASTICITY.		Modulus of rupture.
								First deflection.	Second deflection.	
361. <i>Pinus Cubensis</i> —continued.....	493	Florida	Bay Biscayne.....	A. H. Curtiss.....	Coral	0.8406		1479	1479	1064
	493	do	do	do	do	0.8942		1744	1684	996
362. <i>Picea nigra</i> <i>Black Spruce.</i>	231	Vermont	Charlotte	C. G. Pringle	Cold, peaty	0.5604		1526	1479	930
	231	do	do	do	do	0.5429		1221	1221	902
	373	do	Huntington	do	Gravelly	0.4587		904	921	698
	776	New Brunswick	Bay of Fundy	Intercolonial rail- way.	do	0.4698		1062	1062	504
	776	do	do	do	do	0.4919		1221	1221	834
	794	Province of Que- bec.	Danville	Grand Trunk rail- way.	do	0.4296		996	1062	717
	794	do	do	do	do	0.4153		1221	1085	701
	880	New Brunswick	Bridgeton	Ed. Sinclair	do	0.4425		976	888	687
	880	do	do	do	do	0.4785		939	872	741
363. <i>Picea alba</i> <i>White Spruce.</i>	513	New Hampshire	Stratford	C. G. Pringle	Gravelly	0.4455		787	769	736
	513	do	do	do	do	0.4963		679	688	703
	773	New Brunswick	Bay of Fundy	Intercolonial rail- way.	do	0.4579		1285	1252	813
	773	do	do	do	do	0.4596		1163	1136	792
	784	do	Bridgeton	Ed. Sinclair	do	0.4411		939	976	699
	784	do	do	do	do	0.4530		976	976	731
	791	Province of Que- bec.	Amqui	A. Grant	do	0.4319		1221	1236	811
	791	do	do	do	do	0.4163		1136	1149	703
	364. <i>Picea Engelmanni</i> <i>White Spruce.</i>	292	Colorado	Forest City	T. S. Brandegge	Damp	0.4325		976	1028
292		do	do	do	do	0.3642		554	588	420
575		do	do	C. S. Sargent	Peaty	0.3411		751	751	548
823		do	do	T. S. Brandegge	do	0.3805		868	864	577
365. <i>Picea pungens</i> <i>White Spruce. Blue Spruce.</i>	269	do	Alpine	do	Damp	0.3939		542	574	441
	270 ¹	do	do	do	do	0.4036		444	444	387
	270 ²	do	do	do	do	0.3810		610	642	539
366. <i>Picea Sitchensis</i> <i>Tide land Spruce.</i>	970	Alaska	Sitka	Paul Schultze	do	0.4676		921	957	784
	970	do	do	do	do	0.4568		1062	1085	741
	977	British Columbia	Saw-mill, Burrard inlet.	G. Engelmann and C. S. Sargent.	do	0.4038		1085	1149	682
	977	do	do	do	do	0.3882		1039	1062	635
	1015	Oregon	Weidler's saw-mill, Portland.	do	do	0.4398		1221	1302	776
	1019	do	Saw-mill, Astoria	do	do	0.3517		996	976	614
	1019	do	do	do	do	0.3688		1110	1085	673
	1026	do	Portland Furniture Company.	do	do	0.3891		642	651	553
	1026	do	do	do	do	0.3810		610	642	431
367. <i>Tsuga Canadensis</i> <i>Hemlock.</i>	5	Massachusetts.....	Arnold Arboretum..	C. S. Sargent	Drift	0.4210		763	814	574
	5	do	do	do	do	0.3989		787	721	586
	219	Vermont	Charlotte	C. G. Pringle	Gravelly	0.4716		1085	1062	738
	219	do	do	do	do	0.4699		1017	1007	797
	772	New Brunswick	do	Intercolonial rail- way.	do	0.5124		1136	1136	900
	772	do	do	do	do	0.5129		1136	1177	1031
	775	do	Bay of Fundy.....	do	do	0.4922		488	479	677
	775	do	do	do	do	0.4458		432	432	635
	778	do	do	do	do	0.3496		679	688	541
	778	do	do	do	do	0.3675		669	665	530
	787	do	Bridgeton	Ed. Sinclair	do	0.4811		976	976	745
	787	do	do	do	do	0.4905		1085	1085	790

UNITED STATES UNDER TRANSVERSE STRAIN—Continued.

DEFLECTION, IN MILLIMETERS, UNDER A PRESSURE, IN KILOGRAMS, OF—													Ultimate strength, transverse pressure.	Remarks.	Office number.
50	100	150	200	0 (set.)	200	250	300	350	400	450	500	550			
3.3	6.6	10.0	13.6	1.0	14.1	18.2	22.0	26.5	31.5	37.0			454	Square break	493
2.8	5.8	8.9	12.5	0.5	12.6	16.0	20.6	25.0	29.2				426	Square break on tension side, splitting in axis	493
3.2	6.6	10.0	13.5	0.3	13.8	17.0	21.2	28.0					397	Square break with scales on back	231
4.0	8.0	12.5	16.5	0.4	17.0	22.0	28.7	39.5					385	Crushed at center bearing; square break	231
5.4	10.6	16.4	22.7	1.0	23.0	30.0							298	Square break on tension side, splitting in axis	373
4.6	9.2	14.5	21.0	1.2	21.5								215	Square break at large knot	776
4.0	8.0	12.0	16.4	0.5	16.5	22.0	29.0	45.0					356	Square break	776
4.9	9.2	13.8	19.0	0.6	19.1	26.0	41.0						306	Square break on tension side, splitting in axis	794
4.0	8.0	13.5	19.0	0.5	19.5	27.0							299	Square break	794
5.0	10.0	15.0	21.2	0.9	21.5	30.0							293	do	880
5.2	11.2	17.4	23.6	1.0	24.4	31.5	47.0						316	Square break; split to end	880
6.2	12.7	19.2	27.0	1.6	27.4	37.5	55.5						314	Broke with flat scales on back	513
7.2	14.2	22.4	32.7	4.0	33.5	45.7							300	Square break; shattered	513
3.8	7.8	12.0	16.0	0.5	16.2	21.0	29.2						347	Crushed at center bearing; failed from flakes on tension side	773
4.2	8.6	13.0	17.6	0.5	18.0	23.0	31.5						338	Square break	773
5.2	10.0	15.5	21.3	1.0	21.8	29.6							294	do	784
5.0	10.0	15.3	21.3	0.6	21.5	29.5	43.2						312	do	784
4.0	7.9	12.0	17.0	0.7	17.0	22.5	29.5						346	Crushed at center bearing; failed from thin scale on tension side	791
4.3	8.5	13.0	17.5	0.4	18.0	24.0							360	Crushed; square break	791
5.0	9.5	14.0	19.0	0.6	19.0	25.0	36.0						323	Broke with thin scale on back	292
8.8	16.6	26.4											179	Square break at knot near end	292
6.5	13.0	19.0	28.0	2.4	29.6								234	Square break	575
5.5	11.3	17.4	26.0	2.2	27.2								246	Square break with short splinters	822
9.0	17.0	28.0											188	Started at knot	269
11.0	22.0	35.2											165	do	270 ¹
8.0	15.2	24.7	39.5	5.5	41.2								230	Square break	270 ²
5.3	10.2	15.2	22.0	1.6	22.5	31.0	46.0						313	Crushed at center bearing; broke with few thin splinters	979
4.6	9.0	13.4	18.2	0.7	18.7	25.0	34.0						316	do	979
4.5	8.5	13.2	19.0	1.0	19.0	27.0							291	Crushed at center bearing; broke with fine splinters	977
4.7	9.2	14.3	21.0	1.5	21.8	32.5							271	Crushed at center bearing; broke with thin flake	977
4.0	7.5	11.4	15.4	0.4	15.8	21.0	28.5						331	Crushed at center bearing; broke with fine splinters	1015
4.9	10.0	14.8	22.0	1.5	22.6	35.3							262	Crushed at center bearing; broke with thin flake	1019
4.4	9.0	13.6	19.7	1.0	20.0	28.0							287	Crushed at center bearing; broke with fine splinters	1019
7.6	15.0	24.4	36.4	4.3	38.0								236	Shattered	1026
8.0	15.2	24.7											184	Specimen cross-grained; split	1026
6.4	12.0	17.8	25.0	1.0	25.4								245	Shattered	5
6.2	12.5	19.2	27.0	2.0	27.9								250	Square break on tension side, splitting in axis	5
4.5	9.2	14.0	19.0	0.6	19.0	24.8	34.5						315	Broke with large flakes on back	219
4.8	9.7	14.3	19.5	0.7	19.8	25.0	33.0						340	Square break	219
4.3	8.6	12.8	17.0	0.3	17.3	22.0	28.5	36.4					384	Square break with long split in center	772
4.3	8.3	12.5	17.0	0.4	17.2	22.0	27.0	36.0					449	do	772
10.0	20.4	34.0	49.0	7.2	51.0	69.5							289	Square break on tension side, splitting in axis	775
11.3	22.6	35.5	53.2	7.8	54.0	72.0							271	Shattered	775
7.2	14.2	21.8	32.8	4.0	33.0								231	Square break; shattered	778
7.3	14.7	22.7	34.5	3.0	36.0								226	Square break	778
5.8	10.0	15.0	20.5	0.6	21.0	27.5	35.7						318	Square break; somewhat shattered	787
4.5	9.0	13.5	18.5	0.6	16.7	24.0	32.5						337	Square break	787

FOREST TREES OF NORTH AMERICA.

TABLE III.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Specific gravity of the air-dried specimen.	Direction of grain.	COEFFICIENT OF ELASTICITY.		Modulus of rupture.
								First deflection.	Second deflection.	
387. <i>Tsuga Canadensis</i> —continued	793	Province of Quebec	Danville	Grand Trunk railway.		0.5264		1136	1177	839
	793	do	do	do		0.5248		1221	1285	1024
	817	West Virginia	Grafton	C. G. Pringle.		0.4041		957	986	649
	817	do	do	do		0.3748		814	814	558
	1040	Massachusetts	Danvers	J. Robinson	Moist loam	0.5096		763	746	848
	1040	do	do	do	do	0.5114		751	763	858
	1042	do	North Reading	do		0.4248		1062	1028	703
388. <i>Tsuga Caroliniana</i> . <i>Hemlock.</i>	623	North Carolina	Hendersonville	A. H. Curtiss	Dry, rocky	0.5335		697	713	462
	971	Washington territory.	Wilkeson	G. Engelmann and C. S. Sargent.	Rich loam	0.5318		1017	1017	570
	995	Alaska	Sitka	Paul Schultze.		0.5902		1628	1628	1104
389. <i>Tsuga Mertensiana</i> . <i>Hemlock.</i>	995	do	do	do		0.5472		1526	1479	1055
	980	British Columbia	Silver peak, near Fraser river.	G. Engelmann and C. S. Sargent.	Gravelly loam	0.4590		751	763	720
	980	do	do	do	do	0.4715		775	787	720
390. <i>Tsuga Pattoniana</i>	980	British Columbia	Silver peak, near Fraser river.	G. Engelmann and C. S. Sargent.	Gravelly loam	0.4590		751	763	720
	980	do	do	do	do	0.4715		775	787	720
	980	do	do	do	do	0.4715		775	787	720
391. <i>Pseudotsuga Douglasii</i> . <i>Red Fir. Yellow Fir. Pina. Douglas Fir.</i> Oregon	2712	Colorado	Alpine	T. S. Brandegee	Moist	0.4852		814	842	776
	2712	do	do	do	do	0.4786		857	888	666
	2712	do	do	do	do	0.4874		1110	1149	848
	627	California	Saw-mill, Strawberry valley.	G. Engelmann and C. S. Sargent.		0.5735		1744	1628	1050
	627	do	do	do		0.5382		1221	1356	937
	702	Oregon	Saw-mill, Marshfield	do		0.4373		888	930	572
	704	do	do	do		0.6590		1526	1628	1181
	705	do	E. B. Dean's saw-mill, Marshfield.	do		0.5795		1136	1221	933
	706	do	do	do		0.5477		1395	1395	937
	708	do	do	do		0.6687		1575	1628	1249
	708	do	do	do		0.6892		1808	1808	1317
	709	do	do	do		0.4375		1085	1085	766
	709	do	do	do		0.4448		1163	1085	802
	720	Montana	Saw-mill, Missoula	S. Watson		0.5345		1062	1062	703
	720	do	do	do		0.5448		1221	1221	921
	732	California	Lassen's peak	Sierra Lumber Company.		0.5227		1221	1252	820
	732	do	do	do		0.5090		1163	1163	703
	881	Utah	Salt Lake	M. E. Jones	Rocky	0.5601		996	976	933
	881	do	do	do	do	0.5705		1163	1110	975
	973	British Columbia	Saw-mill, Burrard inlet.	C. S. Sargent		0.5005		1221	1221	771
	973	do	do	do		0.4909		1163	1149	820
	974	do	do	do		0.5306		1356	1436	902
	986	do	Saw-mill, Victoria	G. Engelmann and C. S. Sargent.		0.4596		1085	1122	787
	989	Oregon	Saw-mill, Portland	do		0.6016		1221	1191	1008
	1008	British Columbia	Saw-mill, Burrard inlet.	do		0.4879		1252	1238	764
	1008	do	do	do		0.5501		1628	1628	898
	1011	Oregon	Oregon Railway and Navigation Co.	do		0.5386		1221	1356	900
	1011	do	do	do		0.5325		1191	1191	696
	1016	do	Weidler's saw-mill, Portland.	do		0.5960		1628	1713	900
	1016	do	do	do		0.6085		1628	1628	696
	1018	do	Saw-mill, Astoria	do		0.6129		1628	1628	947
	1018	do	do	do		0.6135		1808	1744	998
	1020	do	Portland Furniture Company.	do		0.4632		1110	1085	1048
	1020	do	do	do		0.4485		974	974	1055

UNITED STATES UNDER TRANSVERSE STRAIN—Continued.

DEFLECTION, IN MILLIMETERS, UNDER A PRESSURE, IN KILOGRAMS, OF—													Ultimate strength: transverse pressure.	Remarks.	Office number.
50	100	150	200	0 (set.)	200	250	300	350	400	450	500	550			
4.3	8.3	13.0	17.5	0.5	18.0	23.0	29.5	39.0					358	Crushed at center bearing with thin flake on tension side	798
4.0	7.6	11.5	15.3	0.2	15.6	19.5	24.3	30.0	42.4				437	Square break; long split in center	793
5.1	9.9	15.4	22.0	0.8	22.2	30.5							273	Square break	817
6.0	12.0	18.2	27.0	1.5	28.0								238	do	817
6.4	13.1	20.4	28.0	2.0	28.0	36.0	47.0	64.0					362	Broke with flakes on tension side	1040
6.5	12.8	18.7	25.8	1.3	26.0	33.5	43.0	61.0					366	Square break on tension side, splitting in axis	1040
4.6	9.5	14.4	20.3	0.7	20.5	27.5							300	Crushed at center bearing; failed with flakes on corner	1042
5.3	10.9	16.4	23.2	1.3	23.7	32.0							300	Square break; shattered	1042
7.0	13.7	21.2											197	Specimen cross-grained; started at knot	933
4.8	9.6	14.4	21.0	1.4	21.5								243	Started at knots	971
3.0	6.0	9.3	12.6	0.4	12.9	16.0	19.7	24.5	30.5	39.0			471	Crushed at center bearing; failed from flakes on back	995
3.2	6.6	9.6	13.2	0.2	13.2	16.5	20.5	25.0	34.0				450	Crushed; splintered	995
4.5	12.8	19.5	27.0	1.1	27.9	36.0	55.5						307	Square break	980
4.3	12.4	18.9	26.4	1.4	26.7	35.5	48.2						307	do	980
6.0	11.6	17.0	23.3	0.9	23.7	32.0	44.0						331	Square break and split at end	271 ^a
5.7	11.0	16.5	24.0	1.0	24.0	33.0							284	Square break on tension side, splitting in axis	271 ^a
4.4	8.5	13.0	17.5	0.6	18.0	23.0	29.5	41.2					362	Did not break; split through the center	271 ^a
2.8	6.0	9.2	12.6	0.0	12.6	16.0	19.5	24.0	30.5				448	Square break; slightly splintered	627
4.0	7.2	11.0	14.5	0.2	14.6	18.2	23.0	29.6					400	do	627
5.5	10.5	17.0	22.5	1.5	23.0								244	Flaked on tension side	702
3.2	6.0	9.0	12.1	0.3	12.2	15.0	18.5	23.0	27.0	33.0	38.5		504	Square break; large splinters	704
4.3	8.0	11.6	15.8	0.5	16.0	20.3	25.5	31.5					398	Specimen cross-grained; split with grain	705
3.5	7.0	10.4	13.6	0.3	14.0	17.0	22.0	27.8	39.5				400	Crushed at center bearing; square break	706
3.1	6.0	9.0	12.3	0.2	12.5	16.0	18.6	22.7	27.5	34.0	45.6		533	Square break on tension side, splitting in axis	708
2.7	5.4	8.2	11.0	0.0	11.2	14.0	17.0	20.0	24.5	29.0	34.5	44.5	562	do	708
4.5	9.0	13.6	18.7	0.4	19.0	24.7	32.2						327	Failed from large splinter on corner	709
4.2	9.0	13.3	17.8	0.5	18.0	23.5	30.0						342	Specimen cross-grained; splintered	709
4.6	9.2	14.0	19.6	0.5	20.4	26.5	35.8						300	Specimen cross-grained; shattered	720
4.0	8.0	12.0	16.2	0.4	16.8	21.0	27.5	34.2					393	do	720
4.0	7.8	11.7	16.0	0.4	16.0	20.8	26.5	36.0					350	Failed from large splinters on corners	732
4.2	8.4	13.0	17.8	0.6	18.0	23.4	30.5						300	Failed from large splinters on one corner	732
4.9	10.0	15.6	22.0	1.0	22.0	29.0	36.0	45.7					398	Square break on tension side, splitting in axis	881
4.2	8.8	13.4	18.6	0.5	18.5	24.6	30.6	38.5	51.5				416	Failed from large splinter on corner	881
4.0	8.0	12.2	17.0	0.4	17.5	22.0	29.2						329	Square break	973
4.2	8.5	12.7	17.0	0.3	17.5	22.5	28.7						350	do	973
2.6	6.8	10.1	13.5	0.3	13.6	17.5	22.2	28.0					385	Failed from large splinter on each corner	974
4.5	8.7	13.2	18.0	0.6	18.0	24.2	31.0						336	Square break on tension side, splitting in axis	986
4.0	8.2	12.5	17.0	0.4	17.2	21.5	27.5	35.0	45.0				430	Crushed at center bearing; broke with fine splinters	989
3.9	7.3	11.0	14.7	0.3	15.0	19.0	25.5						326	Crushed at center bearing; broke with flakes on back	1008
3.0	6.0	9.0	12.5	0.0	12.2	15.5	19.5	26.6					383	Crushed at center bearing; square break; splintered	1008
4.0	7.2	11.0	14.5	0.5	14.8	18.7	24.0	31.5					384	Crushed at center bearing; broke with fine splinters	1011
4.1	8.2	12.3	16.8	0.4	16.9	22.2							297	Started at knot; splintered on corner	1011
3.0	5.7	8.8	12.0	0.2	12.3	15.0	19.0	24.0	32.0				384	Crushed at center bearing; broke with fine splinters	1016
3.0	6.0	9.0	11.6	0.2	12.0	14.8	18.4	23.4	32.5				207	Cracked at knot	1016
3.0	6.0	8.8	11.8	0.2	11.8	14.6	18.0	22.0	28.0				404	Crushed at center bearing and scaled on tension side	1018
2.7	5.6	8.6	11.6	0.2	12.0	14.5	18.5	23.0	31.0				426	Splintered	1018
4.4	9.0	13.3	18.0	0.5	18.0	24.0	34.0						447	do	1020
5.0	10.0	15.2	21.2	0.7	21.5	28.5							450	do	1020

TABLE III.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Specific gravity of the air-dried specimen.	Direction of grain.	COEFFICIENT OF ELASTICITY.		Modulus of rupture.
								First deflection.	Second deflection.	
381. <i>Pseudotsuga Douglasii</i> —cont'd....	1022	Oregon	Portland Furniture Company.	G. Engelmann and C. S. Sargent.		0.4899		1221	1221	769
	1022	do	do	do		0.4735		1163	1136	698
391. <i>Pseudotsuga Douglasii</i> , var. <i>macrocarpa</i> . <i>Hemlock</i> .	642	California.....	Saw-mill, San Bernardino.	W. G. Wright.....		0.5357		1062	1050	783
	642	do	do	do		0.5397		1085	1050	909
392. <i>Abies Fraseri</i>	523	North Carolina ...	Roan mountain	Walcott Gibbs....	Peaty loam	0.3602		976	1017	654
	523	do	do	do	do	0.3523		976	976	621
	523	do	do	do	do	0.3636		976	921	642
393. <i>Abies balsamea</i>	377	Vermont	Monkton	C. G. Pringle.....	Peaty	0.4455		740	734	445
	377	do	do	do	do	0.4419		651	638	584
394. <i>Abies subalpina</i>	449 ¹	Colorado.....	Forest City	T. S. Brandegee ...	Moist, sandy loam	0.3941		775	787	548
	449 ¹	do	do	do	do	0.3750		775	723	370
	449 ²	do	do	do	do	0.3662		872	849	586
	449 ²	do	do	do	do	0.3807		679	642	347
	820	do	do	do	do	0.3358		688	679	518
	820	do	do	do	do	0.3350		751	740	462
	820	do	do	do	do	0.3673		787	787	492
	820	do	do	do	do	0.3622		976	888	460
395. <i>Abies grandis</i>	1009	Oregon	Portland	G. Engelmann and C. S. Sargent.	Rich, alluvial	0.4002		763	751	333
	1010	do	do	do	do	0.3597		976	1085	527
	1010	do	do	do	do	0.3641		976	1039	621
396. <i>Abies concolor</i>	529	Colorado.....	Engelmann's cañon	Robert Douglas...	Rocky	0.3602		660	697	555
	529	do	do	do	do	0.3861		775	787	490
	639	California	Strawberry valley	G. Engelmann and C. S. Sargent.	Alluvial	0.4019		688	655	703
	639	do	do	do	do	0.4744		651	673	764
	733	do	Laassen's peak	Sierra Lumber Company.		0.4969		1285	1320	904
	733	do	do	do	do	0.4504		1252	1320	796
398. <i>Abies amabilis</i>	1004	British Columbia.	Silver peak, near Fraser river.	G. Engelmann and C. S. Sargent.	Rich, sandy loam..	0.4754		1221	1252	769
	1004	do	do	do	do	0.5168		1221	1268	816
399. <i>Abies nobilis</i>	965	Oregon	Cascade mountains	do	Rich	0.5074		1221	1285	905
	965	do	do	do	do	0.4932		1221	1268	820
400. <i>Abies magnifica</i>	647	California	Soda Springs	C. Engelmann and C. S. Sargent.	Gravelly loam	0.4608		542	514	580
	647	do	do	do	do	0.5134		976	939	516
	647	do	do	do	do	0.4965		542	534	703
401. <i>Larix Americana</i>	226 ¹	Vermont.....	Charlotte	C. G. Pringle.....	Cold, swampy	0.7381		2325	1953	1169
	226 ¹	do	do	do	do	0.7295		1526	1628	1055
	774	New Brunswick..	Bay of Fundy.....	Intercolonial railway.		0.6147		1221	1356	937
	774	do	do	do	do	0.6336		1395	1479	790
	781	do	do	do	do	0.5973		697	734	703
	781	do	do	do	do	0.5742		688	688	717
	786	do	Bridgeton	Ed. Sinclair		0.5964		1430	1395	923
	786	do	do	do	do	0.6204		1285	1356	1066
	795	do	Danville	Grand Trunk railway.		0.6170		1479	1436	937
	795	do	do	do	do	0.6030		1526	1395	670
	840	Massachusetts....	Wenham	J. Robinson	Swampy	0.5988		842	872	820
	840	do	do	do	do	0.6272		814	843	823

UNITED STATES UNDER TRANSVERSE STRAIN—Continued.

DEFLECTION, IN MILLIMETERS, UNDER A PRESSURE, IN KILOGRAMS, OF—													Ultimate strength: transverse pressure.	Remarks.	Office number.
50	100	150	200	0 (set.)	200	250	300	350	400	450	500	550			
4.0	8.0	12.0	16.4	0.4	16.5	22.0	31.0						328	Square break; split at end	1022
4.2	8.6	12.5	17.0	0.4	17.1	22.0	29.0						298	Square break	1022
4.6	9.3	14.0	19.0	0.4	19.3	25.2	32.0						384	Square break on tension side, splitting in axis; shattered	643
4.5	9.3	14.3	19.4	0.4	19.7	25.7	33.0	42.8					388	do	643
5.0	9.6	15.0	21.0	1.0	21.5	31.0							279	Square break; splintered	528
5.0	10.0	15.0	22.0	1.5	22.2	33.5							265	do	528
5.0	10.6	15.5	22.6	1.1	23.7	34.0							274	do	528
6.6	13.3	21.0											190	Square break	377
5.8	10.8	16.8	22.5	0.5	22.5								249	Specimen cross-grained	377
6.3	12.4	19.5	29.0	2.5	29.3								234	Crushed at center bearing; square break on tension side, splitting in axis.	449 ^a
6.3	13.5	21.5											158	Square break at knot	449 ^a
5.6	11.5	16.7	24.5	1.5	25.6	39.2							250	Crushed at center bearing; flaked on back	449 ^a
7.2	15.2												148	Specimen cross-grained; broke at knot	449 ^a
7.1	14.4	22.0	32.0	2.4	35.5								221	Square break	820
6.5	13.2	21.0											197	do	820
6.2	12.4	20.5	32.5	4.4	35.0								210	Square break and split	820
5.0	11.0	17.0											200	Broke with long, oblique split	820
6.4	13.0												142	Specimen shaky; broke with long split	1009
5.0	9.0	14.0	20.5	1.0	20.7								225	Crushed at center bearing	1010
5.0	9.4	14.5	21.0	1.6	22.0	32.0							265	Square break with long, thin splinter	1010
7.4	14.0	21.5	32.0	2.5	34.5								237	Shattered	529
6.3	12.4	20.0	29.0	2.0	30.0								213	Square break; shattered	529
7.1	14.9	23.0	32.2	1.9	33.4	44.7	64.0						300	Square break with flakes on tension side	639
7.5	14.5	22.4	31.5	2.2	32.5	43.7	63.5						326	Thin flake on back	639
3.8	7.4	11.0	14.9	0.4	15.0	18.5	23.4	29.5					386	Crushed at center bearing with thin flakes	733
3.9	7.4	11.3	15.0	0.2	16.0	21.0	28.0						340	Crushed at center bearing; splintered	733
4.0	7.8	12.0	16.4	0.6	16.8	22.0	32.5						328	Square break	1004
4.0	7.7	11.9	16.2	0.5	16.8	22.0	30.0						348	Crushed at center bearing; square break	1004
4.0	7.6	11.4	15.4	0.4	16.0	21.0	27.0	33.0					386	Crushed at center bearing; flaked	965
4.0	7.7	11.7	16.0	0.4	16.2	21.8	29.0	47.0					350	Crushed at center bearing; splintered	965
9.0	19.0	30.7	46.0	5.5	49.4	94.0							250	Square break; split	647
5.0	10.4	15.7	22.0	1.0	22.5	30.0	41.0						348	Specimen cross-grained	647
9.0	18.3	29.5	40.8	4.0	43.5	62.0							300	Square break on tension side, splitting in axis	647
2.1	5.0	7.4	10.0	0.1	10.2	12.7	16.0	19.0	24.0	29.0			499	Crushed at center bearing; broke with fine splinters	226 ^a
3.2	6.0	9.2	12.0	0.2	12.0	15.0	18.5	23.0	29.0	38.0			450	Crushed at center bearing; broke with thin scales	226 ^a
4.0	7.2	10.8	14.5	0.8	14.6	18.0	23.5	30.0	46.0				400	Crushed at center bearing; broke with fine splinters	774
3.5	6.6	10.0	13.2	0.4	13.5	16.5	23.0						337	Large scale on tension side	774
7.0	13.3	21.0	29.7	2.8	30.5	40.4	55.5						300	Shattered at one end	781
7.1	14.2	22.4	32.0	2.9	32.8	44.0	63.5						306	Specimen cross-grained; square break; split	781
3.4	7.0	11.0	14.5	0.5	14.6	18.2	23.7	30.0					394	Crushed at center bearing; broke with fine splinters	786
3.8	7.2	10.8	14.5	0.4	14.7	18.5	23.4	29.0	36.0	46.0			455	Square break; splintered	786
3.3	6.8	10.0	13.2	0.3	13.5	17.0	21.5	28.5	40.0				400	Crushed at center bearing; scaled on tension side	795
3.2	7.0	10.3	14.0	0.4	14.5	18.8	24.2	32.5					371	Crushed at center bearing; scaled	795
5.8	11.2	18.2	26.3	3.0	27.0	35.0	49.0						350	Failed from large splinter on corner	840
6.0	11.6	17.8	25.7	2.0	26.5	35.5	47.0	75.0					351	Square break on tension side, splitting in axis	840

TABLE III.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Specific gravity of the air-dried specimen.	Direction of grain.	COEFFICIENT OF ELASTICITY.		Modulus of rupture.
								First deflection.	Second deflection.	
402. <i>Larix occidentalis</i> <i>Tamarack.</i>	719	Montana.....	Missoula.....	S. Watson.....		0.6255		1221	1221	914
	719	do.....	do.....	do.....		0.6066		1356	1395	1106
	984	Washington territory.....	Fulda.....	W. Siksdorf.....	Moist.....	0.8136		1628	1713	1289
	984	do.....	do.....	do.....	do.....	0.8364		1628	1713	1481
	1006	do.....	do.....	do.....	do.....	0.8426		1878	1953	1287
	1006	do.....	do.....	do.....	do.....	0.8376		1953	1953	1287
PALMACEÆ.										
405. <i>Washingtonia filifera</i> <i>Fan-leaf Palm.</i>	1159	California.....	Agua Caliente.....	W. G. Wright.....	Dry, gravelly.....	0.6012		460	408	271
	1159	do.....	do.....	do.....	do.....	0.6307		697	723	586

TABLE IV.—BEHAVIOR OF SOME OF THE WOODS OF THE UNITED STATES

Species.	Office number.	State.	Locality.	Collector.	Soil.	Specific gravity.	Direction of pressure.	COEFFICIENT OF ELASTICITY.		Modulus of rupture.
								Second deflection.	Fourth deflection.	
MAGNOLIACEÆ.										
1. <i>Magnolia grandiflora</i> <i>Big Laurel. Bull Bay.</i>	346	Alabama	Cottage Hill	C. Mohr	Rich loam	0.7347		1191	1097	974
2. <i>Magnolia glauca</i> <i>Sweet Bay. White Bay. Beaver Trees. White Laurel. Swamp Laurel.</i>	354	do	do	do	Swampy	0.5262		1185	1028	816
SAPINDACEÆ.										
54. <i>Sapindus marginatus</i> <i>Wild China. Soapberry.</i>	307	Texas	Dallas	J. Reverchon	Rich, damp	0.7838		1109	1062	851
LEGUMINOSÆ.										
77. <i>Robinia Pseudacacia</i> <i>Locust. Black Locust. Yellow Locust.</i>	1248	New York.....	Long Island	M. C. Beedle		0.8476		1221	1221	1160
HAMAMELACEÆ.										
139. <i>Liquidambar styraciflua</i> <i>Sweet Gum. Star-leaved Gum. Liquidamber. Red Gum. Bilsted.</i>	546	Alabama	Kemper's mill	C. Mohr	Rich, alluvial	0.6537		939	921	923
OLEACEÆ.										
192. <i>Fraxinus Americana</i> , var. <i>Texensis</i>	364	Texas	Dallas	J. Reverchon	Dry, calcareous...	0.8198		1221	1191	1172
BIGNONIACEÆ.										
207. <i>Catalpa speciosa</i> <i>Western Catalpa.</i>	38	Missouri.....	Charleston.....	C. S. Sargent.....	Wet clay	0.4783		957	940	692
	38	do	do	do	do	0.4757		939	888	703
URTICACEÆ.										
224. <i>Ulmus Americana</i> <i>White Elm. American Elm. Water Elm.</i>	19	Massachusetts...	Arnold Arboretum	do	Drift	0.7534		1085	1050	1118
228. <i>Celtis occidentalis</i> <i>Sugarberry. Hackberry.</i>	306	Texas	Dallas	J. Reverchon	Alluvial	0.7491		888	800	916
JUGLANDACEÆ.										
239. <i>Juglans nigra</i> <i>Black Walnut.</i>	951	do	New Braunfels	C. Mohr	Moist, calcareous.	0.7108		976	996	936
243. <i>Carya aquatica</i> <i>Water Hickory. Swamp Hickory. Bitter Pecan.</i>	362	Mississippi.....	Vicksburg	do		0.7739		976	957	956
CUPULIFERÆ.										
251. <i>Quercus alba</i> <i>White Oak.</i>	8	Massachusetts....	Arnold Arboretum	C. S. Sargent.....	Drift	0.6958		708	787	892
	8	do	do	do	do	0.6820		763	600	658

UNITED STATES UNDER TRANSVERSE STRAIN—Continued.

DEFLECTION, IN MILLIMETERS, UNDER A PRESSURE, IN KILOGRAMS, OF—													Ultimate strength: transverse pressure.	Remarks.	Office number.
50	100	150	200	0 (set.)	200	250	300	350	400	450	500	550			
4.0	8.0	12.0	16.2	0.5	16.5	21.0	27.5	36.0					390	Crushed at center bearing; broke with fine splinters	719
3.6	7.0	10.5	14.0	0.4	14.0	18.0	22.5	28.0	34.0	45.0			472	do	719
3.0	5.7	8.4	11.0	0.3	11.4	14.0	17.5	20.6	24.7	28.5	35.7	47.6	550	do	964
3.0	5.7	8.2	11.0	0.2	11.0	13.6	16.5	19.5	23.0	26.8	32.0	36.5	632	Deflection with 600 kilograms = 45.5 millimeters; broke with fine splinters.	934
2.6	5.0	7.6	10.4	0.2	10.4	13.0	16.0	19.0	22.7	27.2	34.0		549	Crushed at center bearing; flaked	1006
2.5	5.0	7.5	10.0	0.2	10.0	12.6	15.5	18.5	22.0	26.5			549	Failed from large splinter on corner	1006
10.6	24.2												116	Twisted and split	1159
7.0	13.5	20.0	29.4	1.8	33.5								250	Square break	1159

UNDER TRANSVERSE STRAIN: SPECIMENS EIGHT CENTIMETERS SQUARE.

DEFLECTION, IN MILLIMETERS, UNDER A PRESSURE, IN KILOGRAMS, OF—											Ultimate strength: transverse pressure.	Remarks.	Office number.
400	800	1200	1600	0 (set.)	1600	2000	2400	2800	3200	3600			
2.0	4.1	6.4	8.9	0.2	9.0	11.2	15.1	19.5	26.0		3325	Broke with large splinters on back	346
2.2	4.3	6.9	9.5	0.3	9.5	12.4	16.2				2785	Broke with large splinters	354
2.2	4.4	6.6	9.2	0.4	9.2	12.2	16.0				2903	Broke with large flake on back	307
2.0	4.0	6.1	8.0	0.0	8.1	10.2	12.6	15.2	19.2	25.0	3992	Broke with large splinters on back	1248
2.9	5.2	7.7	10.6	0.7	10.7	14.2	19.3	26.0			3184	Broke with many fine splinters	546
2.2	4.0	6.0	8.2	0.3	8.2	10.2	13.0	16.5	20.6	26.5	4000	Broke with large splinters	364
2.5	5.1	7.8	11.0	0.3	11.0	15.5					2361	Crushed and split to the end	38
2.8	5.2	7.9	11.0	0.4	11.0	15.7	22.6				2400	Broke with fine splinter on one corner	38
2.3	4.5	6.8	9.3	0.2	9.4	12.0	15.2	18.5	25.0	35.0	3815	Broke with fine splinters	19
2.9	5.5	8.6	12.2	0.8	12.3	17.0	23.2	33.0			3125	Broke with large splinter on one corner	306
2.6	5.0	7.5	9.8	0.4	10.0	12.5	15.5	19.5			3193	Broke with large splinters on corners	351
2.5	5.0	7.4	10.2	0.5	10.3	13.6	18.2	24.4	36.0		3270	Broke with large splinters	363
2.5	6.9	9.8	12.4			15.6	19.2	27.3			3043	Broke with large splinters on back	8
3.1	6.4	10.0	14.8	1.2	15.0	21.0					2245	do	8

TABLE IV.—BEHAVIOR OF SOME OF THE WOODS OF THE UNITED STATES UNDER

Species.	Office number.	State.	Locality.	Collector.	Soil.	Specific gravity.	Direction of pressure.	COEFFICIENT OF ELASTICITY.		Modulus of rupture.
								Second deflection.	Fourth deflection.	
259. <i>Quercus Michauxii</i> <i>Basket Oak. Cow Oak.</i>	524	Alabama	Kemper's mill	C. Mohr	Alluvial	0.8107		775	697	884
	524	do	do	do	do	0.8348		800	745	984
260 <i>Quercus Prinus</i> <i>Chestnut Oak. Rock Chestnut Oak.</i>	925	do	Cullman	do	Dry, rocky	0.8158		1085	1028	582
275. <i>Quercus Kelloggii</i> <i>Black Oak.</i>	963	Oregon	Eugene City	G. H. Collier		0.7625		626	564	736
280. <i>Quercus aquatica</i> <i>Water Oak. Duck Oak. Possum Oak. Punk Oak.</i>	349	Alabama	Cottage Hill	C. Mohr	Sandy loam	0.7253		1366	1320	1144
BETULACEÆ.										
301. <i>Alnus rubra</i> <i>Alder.</i>	991	Washington territory.	Puyallup	G. Engelmann and C. S. Sargent.		0.5381		729	610	524
CONIFERÆ.										
329. <i>Chamaecyparis sphaeroides</i> <i>White Cedar.</i>	350	Alabama	Cottage Hill	C. Mohr	Sandy, wet	0.8719		364		376
370. <i>Pinus Taeda</i> <i>Loblolly Pine. Old-field Pine. Rosemary Pine.</i>	82	Florida	Duval county	A. H. Curtiss	Moist, sandy	0.5802		1061	1017	792
	82	do	do	do	do	0.7614		1285	1302	820
372. <i>Pinus serotina</i> <i>Pond Pine.</i>	83	do	do	do	Moist, sandy loam	0.7614		989	921	933
	83	do	do	do	do	0.8271		787	751	904
380. <i>Pinus palustris</i> <i>Long-leaved Pine. Southern Pine. Georgia Pine. Yellow Pine. Hard Pine.</i>	81	do	do	do	Sandy loam	0.8609		1436	1550	1057
	85	do	do	do	Moist, sandy	0.7213		1065	1007	940
	243	do	Saw-mill, Saint John's river.	do		0.6788		1221	1252	820
	243	do	do	do		0.6193		1163	1221	949
381. <i>Pinus Cubensis</i> <i>Slash Pine. Swamp Pine. Bastard Pine. Meadow Pine.</i>	84	do	Duval county	do	Moist, sandy	0.7633		1163	1221	1029

TRANSVERSE STRAIN: SPECIMENS EIGHT CENTIMETERS SQUARE—Continued.

DEFLECTION, IN MILLIMETERS, UNDER A PRESSURE, IN KILOGRAMS, OF—											Ultimate strength: transverse pressure.	Remarks.	Office number.
400	800	1200	1600	0 (set.)	1600	2000	2400	2800	3200	3600			
3.2	6.3	10.0	14.0	1.5	14.5	19.5	26.5	39.0			3016	A large season crack at one corner; splintered on opposite corner.	524
3.2	6.1	9.3	13.1	1.3	13.3	18.0	24.2	33.0	49.0		3393	Splinters on corners.....	524
2.5	4.5	7.0	9.5	0.6	9.5	13.3					1987	Split lengthwise with small splinters on corners.	925
4.0	7.8	12.0	17.3	1.4	17.5	24.0	40.6				2513	Broke with large splinters on corners.....	963
2.0	3.5	5.5	7.4	0.2	7.3	9.3	11.8	14.6	19.0	25.5	3905	Crushed at center bearing; broke with fine splinters...	349
3.2	6.7	10.7	16.0	2.0	16.7						1789	Broke with large splinters on back; shattered.....	991
6.8	13.4	21.0									1284	Short, square break; no splinters.....	350
2.3	4.6	7.1	9.6	0.2	9.7	12.6	17.0				2703	Shattered.....	83
2.0	3.8	5.6	7.5	0.2	7.6	10.0	12.0	21.5			2800	do.....	83
2.7	5.2	8.0	10.6	0.4	11.0	13.6	17.3	22.0			3184	0.75 sap-wood; broke with large splinter on corner.....	83
3.0	6.2	9.5	13.0	0.9	13.4	18.0	23.0	30.0			3085	0.5 sap-wood; broke with large splinter on corner.....	83
1.9	3.4	5.1	6.3	0.0	7.0	8.7	10.5	12.5	14.6	17.5	3600	Square break on tension side, splitting in axis.....	81
2.1	4.5	7.0	9.7	0.3	9.9	12.6	16.4	20.5	27.5		3207		85
2.0	4.0	5.9	7.8	0.0	7.9	10.0	12.2				2789	Broke with large splinter on corner; somewhat shattered.	243
2.2	4.2	6.0	8.0	0.4	8.0	9.8	12.0	16.0	23.5		3239	Square break on tension side, splitting in axis.....	243
2.3	4.2	6.0	8.0	0.4	8.0	10.2	12.6	15.5	19.5		3513	Split between rings at one end.....	84

TABLE V.—BEHAVIOR OF THE PRINCIPAL WOODS

Species	Office number.	State.	Locality.	Collector.	Soil.	Longitudinal compression: Ultimate strength, in kilos.	Remarks.
MAGNOLIACEÆ.							
1. <i>Magnolia grandiflora</i>	346	Alabama	Cottage Hill	C. Mohr.	Rich loam	7853	Crushed fibers at 10 millimeters knot 28 millimeters from end.
Big Laurel. Bull Bay.	346	do	do	do	do	8056	Triple flexure
2. <i>Magnolia glauca</i>	354	do	do	do	Swampy	7212	Fibers crushed at 51 millimeters from end.
Sweet Bay. White Bay. Beaver Tree. White Laurel. Swamp Laurel.	354	do	do	do	do	6369	Fibers crushed at middle
3. <i>Magnolia acuminata</i>	746	Virginia	Wytheville	H. Shriver	Clay limestone	6795	Fibers crushed at 76 millimeters from end.
Cucumber Tree. Mountain Magnolia.	246	do	do	do	do	6582	Triple flexure
	261 ¹	do	Fancy Gap	do	Rich, light	5126	Crushed in vicinity of knots at middle.
	261 ²	do	do	do	do	5570	Crushed fibers at 63 millimeters from end.
	534	Mississippi	Selvers' mill	C. Mohr.	do	7389	Crushed fibers at 76 millimeters from end.
	534	do	do	do	do	8333	Crushed fibers at 102 millimeters from end and on opposite side at end.
4. <i>Magnolia cordata</i>	1178	Alabama	Winston county	do		6577	Crushed fibers at 38 millimeters from middle.
Cucumber Tree.	1178	do	do	do		6527	Crushed fibers in vicinity of knots 63 millimeters from end.
5. <i>Magnolia macrophylla</i>	582	Mississippi	Quitman	do	Rich, low	7357	Opened longitudinal crack between rings.
Large-leaved Cucumber Tree.	582	do	do	do	do	8301	Crushed fibers at 63 millimeters from end.
6. <i>Magnolia Umbrella</i>	266 ¹	Virginia	Wytheville	H. Shriver		5647	Crushed fibers at 128 millimeters from end.
Umbrella Tree. Elk Wood.	266 ²	do	do	do		6073	Triple flexure; developed intersecting "Cooper lines".
7. <i>Magnolia Fraseri</i>	260	do	Fancy Gap	do	Damp	5806	Failed at 6 millimeters knot 51 millimeters from end.
Long-leaved Cucumber Tree.	260	do	do	do	do	7575	Fibers crushed at 51 and at 128 millimeters from end.
8. <i>Liriodendron Tulipifera</i>	395	Michigan	Lansing	W. J. Beal		4663	Failed at knot at middle.
Tulip Tree. Yellow Poplar. White Wood.	818	West Virginia	Grafton	C. G. Pringle		6341	Fibers crushed at 51 millimeters from end; angle of crushing, 55°.
	818	do	do	do		6636	Fibers crushed at 128 millimeters from end; angle of crushing, 65°.
	1231	Pennsylvania	Chester county	P. P. Sharples		6514	Fibers crushed at 76 millimeters from end; angle of crushing, 75°.
	1231	do	do	do		6169	Fibers crushed at middle
	1232	do	do	do		6305	do
	1232	do	do	do		5874	Fibers crushed in vicinity of knot 51 millimeters from end.
	1236	Tennessee	Saw-mill at Nashville.	A. E. Baird		5489	Fibers crushed at 63 millimeters from end.
	1236	do	do	do		5806	do
ANONACEÆ.							
9. <i>Asimina triloba</i>	211	Missouri	Meramec river, Jefferson county.	G. W. Letterman	Alluvial	3402	Fibers crushed at 114 millimeters from end.
Papaw. Oustard Apple.	211	do	do	do	do	3388	Crushed at knot 63 millimeters from end.
10. <i>Anona laurifolia</i>	479	Florida	Bay Escayne	A. H. Curtiss	Swampy	4690	Fibers crushed at 63 millimeters from end; angle of crushing, 65°.
Pond Apple.	479	do	do	do	do	4967	Fibers crushed on one side at middle.
CANELACEÆ.							
12. <i>Canella alba</i>	1131	do	Elliott's Key	do	Coral	12746	Fibers crushed at middle and split along grain.
White Wood. Cinnamon Bark. Wild Cinnamon.	1131	do	do	do	do	12292	Fibers crushed near middle; deflected from heart.
TERNSTROMIACEÆ.							
14. <i>Gordonia Lasianthus</i>	236	South Carolina	Bonneau's Depot	H. W. Ravenel	Wet pine-barren	5842	Fibers crushed at knots near end.
Loblolly Bay. Tan Bay.	236	do	do	do	do	5752	Fibers crushed at 63 and at 127 millimeters from end at knots 10 millimeters in diameter.
	414	do	Alken	do	Swampy	6790	Fibers crushed at 63 millimeters from end.
	414	do	do	do	do	6396	Fibers crushed at 10 millimeters knot near end.
TILIACEÆ.							
17. <i>Tilia Americana</i>	2	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	4287	Fibers crushed at 19 and at 103 millimeters from end.
Lime Tree. Bass Wood. American Linden. Lin. Bee Tree.	2	do	do	do			
	124	Michigan	Big Rapids	W. J. Beal	Gravel	4944	Fibers crushed at 127 millimeters from end.

OF THE UNITED STATES UNDER COMPRESSION.

Direction of grain.	PRESSURE, IN KILOGRAMS, REQUIRED TO PRODUCE AN INDENTATION, IN MILLIMETERS, OF—												Remarks.	Office number.
	0.25	0.51	0.76	1.02	1.27	1.52	1.78	2.03	2.28	2.54	4.81	5.08		
	2041		8221	8447	8497	8670	8847	8978	4078	4209			Began to shear fibers.....	346
	1724	2388	2556	2676	2805	2921	3016	3112	3189	3291			Slight shearing of fibers.....	346
	1152	1834	1433	1547	1651	1783	1805	1846	1900	1978			Sheared fibers.....	354
	1080	1829	1488	1533	1603	1667	1742	1760	1796	1878			Slight shearing of fibers.....	354
	975	1343	1429	1533	1588	1642	1696	1733	1758	1792			do.....	346
	1016	1420	1535	1651	1787	1805	1889	1941	2037	2111			Splintered at pith; slight shearing of fibers.....	246
	1080	1315	1420	1506	1588	1660	1674	1758	1787	1833			Indented without shearing fibers.....	261
	1157	1379	1465	1579	1665	1715	1774	1824	1864	1901			Sheared fibers.....	261
	1016	1442	1583	1660	1751	1860	1910	1982	2068	2114			Slight shearing of fibers.....	584
	1125	1610	1760	1828	1932	2019	2105	2245	2304	2359			do.....	584
	884	1098	1170	1228	1315	1343	1406	1483	1524	1538			do.....	1178
	1111	1323	1402	1470	1538	1606	1660	1719	1740	1819			Fibers sheared.....	1178
	1325	1742	1882	2005	2123	2223	2318	2418	2486	2635			do.....	582
	1384	1647	1765	1887	2025	2078	2191	2263	2336	2404			do.....	582
	616	975	1098	1111	1148	1216	1247	1297	1329	1343			Indented without shearing fibers.....	200
	953	1270	1388	1470	1542	1619	1678	1742	1787	1855			Fibers sheared.....	200
	1080	1524	1665	1810	1901	1996	2082	2178	2245	2318			Indented without shearing fibers.....	200
	1016	1015	1787	1914	2032	2082	2146	2195	2223	2295			Slight shearing; short specimen, 120 millimeters long; split at both ends.	200
	907	1066	1125	1143	1175	1202	1234	1252	1297	1315			Sheared fibers.....	895
	1025	1297	1388	1470	1533	1606	1674	1737	1792	1846			Slight shearing of fibers.....	818
	749	1025	1170	1229	1311	1370	1415	1474	1524	1565			do.....	818
	630	844	925	980	1048	1125	1152	1193	1220	1266			Sheared fibers.....	1231
	1052	1288	1397	1433	1533	1601	1656	1706	1746	1801			do.....	1231
	794	1080	1152	1216	1257	1302	1343	1379	1384	1442			do.....	1232
	885	1012	1084	1120	1186	1234	1270	1335	1352	1388			do.....	1232
	916	1157	1229	1293	1361	1429	1479	1520	1565	1601			do.....	1236
	885	1116	1166	1202	1261	1306	1352	1397	1456	1515			do.....	1236
	607	894	957	1016	1089	1075	1111	1116	1129	1152			do.....	211
	602	943	1052	1111	1157	1198	1229	1261	1288	1297			do.....	211
	943	1429	1610	1805	1941	2068	2159	2254	2304	2336			Slight shearing of fibers.....	479
	1034	1660	1882	2019	2182	2227	2341	2386	2477	2563			do.....	479
	3674	6577	7621	8346	8578	8822	8936	8981	9844	9571			Sheared fibers and opened grain.....	1181
	3311	6668	8523	9458	9758	10206	10614	10886	10990	11227			Sheared fibers and opened grain at middle and at end.	1181
	935	1125	1248	1343	1388	1452	1533	1579	1642	1701			Sheared fibers.....	226
	535	1021	1116	1211	1298	1347	1433	1488	1538	1597			do.....	226
	1089	1519	1642	1746	1867	1932	2005	2068	2146	2218			do.....	414
	1134	1583	1665	1760	1846	1923	1991	2059	2123	2177			do.....	414
	934	1157	1297	1335	1393	1442	1497	1542	1578	1619			do.....	2
	703	880	925	1007	1048	1111	1157	1189	1225				do.....	2
	717	852	894	957	1034	1043	1071	1093	1120	1123			do.....	124

TABLE V.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Longitudinal compression: Ultimate strength, in kilos.	Remarks.
17. <i>Tilia Americana</i> —continued	124	Michigan	Big Rapids	W. J. Beal	Gravel	4944	
	252	Missouri	Allenton	G. W. Letterman	Alluvial	7167	Fibers crushed at 114 millimeters from end.
	252	do	do	do	do	7235	Fibers crushed at 45 millimeters from end.
	316	Michigan	Hersey	W. J. Beal	Rich loam	6314	Fibers crushed at 51 millimeters from end.
	316	do	do	do	do		
	1089	Massachusetts	Danvers	J. Robinson	Moist gravel	5080	Fibers crushed at 76 millimeters from end.
	1089	do	do	do	do	4527	Triple flexure
17. <i>Tilia Americana</i> , var. <i>pubescens</i> .	745	Georgia	Bainbridge	A. H. Curtiss	Low	6305	Fibers crushed at 38 millimeters from end.
	745	do	do	do	do	6668	Fibers crushed at 127 millimeters from end.
18. <i>Tilia heterophylla</i> <i>White Bass Wood. Wahoo.</i>	2851	Kentucky	Cliffs Kentucky river.	W. M. Linney	Limestone	4917	Fibers crushed at 102 millimeters from end.
	2853	do	Mercer county	do	do	5548	Fibers crushed at 51 millimeters from end.
	2853	do	do	do	do	5126	Fibers crushed at 152 millimeters from end.
	320	Tennessee	Cumberland river.	A. Gattinger	Alluvial	7666	Fibers crushed at middle
	320	do	do	do	do	8278	Fibers crushed at 114 millimeters from end.
MALPIGHIACEÆ.							
19. <i>Byrsionia lucida</i> <i>Tallowberry. Glamberry.</i>	1113	Florida	No-Name Key	A. H. Curtiss	Coral	6260	Fibers crushed near middle in vicinity of small knots.
	1113	do	do	do	do		
ZYGOPHYLLACEÆ.							
20. <i>Guaiaecum sanctum</i> <i>Lignum-vita.</i>	476	do	Upper Metacombe Key.	do	do	11930	Longitudinal split; stick warped before test.
	1133	do	Elliott's Key	do	do	11648	Crushed fibers and split along grain near end.
RUTACEÆ.							
23. <i>Xanthoxylum Clava-Herculis</i> <i>Toothache Tree. Prickly Ash. Sea Ash. Pepper Wood. Wild Orange.</i>	735	do	Chattahoochee	do	Dry, sandy	7372	Triple flexure; developed intersecting "Cooper lines".
	735	do	do	do	do	7698	Crushed at middle; deflected from heart.
	1086	Texas	Palestine	C. Mohr	Damp, sandy	7707	Crushed fibers at 127 millimeters from end.
	1086	do	do	do	do	5978	Crushed fibers at knot 6 millimeters in diameter; angle of crushing, 40°.
24. <i>Xanthoxylum Caribæum</i> <i>Satin Wood.</i>	1140	Florida	Bahia Honda Key.	A. H. Curtiss	Coral	11158	Brittle; crushing of fibers; splintered along grain.
	1140	do	do	do	do	10751	Fractured suddenly; shattered along grain.
SIMARUBEÆ.							
25. <i>Simaruba glauca</i> <i>Paradise Tree.</i>	487	do	Bay Biscayne	do	do	7303	Fibers crushed at middle; angle of crushing, 90°.
	487	do	do	do	do	6328	Split end to end along grain
BURSERACEÆ.							
26. <i>Bursera gummifera</i> <i>Gum Elemi. Gumbo Limbo. West Indian Birch.</i>	462	do	Upper Metacombe Key.	do	do	2350	Fibers crushed in vicinity of knots.
	462	do	do	do	do	2595	Fibers crushed at 38 millimeters from end.
20. <i>Amyris sylvatica</i> <i>Torch Wood.</i>	475	do	do	do	do	11975	Split along grain
	475	do	do	do	do		
MELIACEÆ.							
31. <i>Swietenia Mahagoni</i> <i>Mahogany. Madeira.</i>	452	do	do	do	do	11204	Fibers crushed at 89 millimeters from end; threw off splinters.
	452	do	do	do	do	10115	Failed at 13 millimeters from end; knot 76 millimeters from end.
ILICINEÆ.							
32. <i>Ilex opaca</i> <i>American Holly.</i>	280	South Carolina	Waverly Mills	W. St. J. Mazyck	Sandy loam	7339	Triple flexure; knot near end
	280	do	do	do	do	6078	Failed at knot 63 millimeters from end.
34. <i>Ilex Dahoon</i> <i>Dahoon. Dahoon Holly.</i>	484	Florida	Bay Biscayne	A. H. Curtiss	Low, damp	5652	Failed in vicinity of knots 76 millimeters from end.
	484	do	do	do	do	5512	Fibers crushed at 114 millimeters from end.
CYRILLACEÆ.							
35. <i>Chiftonia ligustrina</i> <i>Tit. Iron Wood. Buckwheat Tree.</i>	368	Alabama	Cottage Hill	C. Mohr	Wet	6169	Fibers crushed at end and at 64 millimeters from end.
	338	do	do	do	do	5706	Triple flexure; fibers crushed at 127 millimeters from end; grain wavy.

UNITED STATES UNDER COMPRESSION—Continued.

Direction of grain.	PRESSURE, IN KILOGRAMS, REQUIRED TO PRODUCE AN INDENTATION, IN MILLIMETERS, OF—												Remarks.	Office number.
	0.25	0.51	0.76	1.02	1.27	1.52	1.78	2.03	2.28	2.54	4.51	5.08		
	472	694	753	767	794	817	852	871	918	966			Sheared fibers.....	134
	826	935	916	1008	1139	1184	1247	1302	1338	1365			do.....	232
	544	794	835	862	912	953	980	1025	1061	1095			Slight shearing of fibers.....	252
	612	749	803	857	930	953	998	1039	1080	1111			do.....	316
	662		857	903	948	1002	1034	1066	1107	1148			do.....	316
	612	812	839	894	925	948	989	1021	1052	1071			do.....	1069
	640	821	862	903	944	984	1020	1043	1080	1107			Sheared fibers.....	1089
	612	789	866	899	907	948	980	1016	1089	1075			do.....	745
	635	889	889	934	993	1043	1066	1120	1166	1220			do.....	745
	481	694	767	817	898	889	898	912	966	990			do.....	285
	857	989	1057	1071	1125	1175	1202	1229	1275	1311			do.....	285
	798	857	898	953	989	1012	1052	1080	1139	1170			do.....	285
	866	1093	1161	1216	1261	1320	1374	1420	1470	1504			do.....	320
	749	1039	1111	1161	1207	1247	1297	1343	1406	1438			do.....	320
	1769	2926	3198	3339	3475	3529	3620	3697	3765	3819			do.....	1113
	1610	2404	2858	3153	3257	3393	3515	3652	3710	3756			do.....	1113
	4854	8663	11022	12565	13562	14289	14742	14946	14969	15105	16103	16194	Sheared fibers; split along grain.....	476
	1769	5421	8392	10614	11817	12565	13245	13653	14198	14590	15531	15740	Sheared fibers; split along grain; did not take even bearing.	1133
	1293	1996	2173	2245	2259	2304	2318	2468	2522	2567	2680	3094	Sheared fibers.....	735
	1111	1819	2082	2068	2164	2223	2277	2322	2368	2400	2790	2994	do.....	735
	2041	3016	3157	3307	3429	3543	3647	3742	3856	3919	4626	4786	do.....	1096
	1179	1882	2041	2259	2341	2363	2395	2422	2499	2563	3016	2287	do.....	1096
	1905	4128	5285	5693	5987	6396	6423	6646	6895	6985	7892	7802	Sheared fibers; split along grain.....	1140
	2717	4513	5262	5670	5942	6214	6419	6600	6804	6976	7802	8165	Sheared fibers.....	1140
	861	1134	1220	1311	1315	1356	1442	1456	1488	1501	1746	1928	do.....	467
	866	1184	1352	1388	1452	1520	1569	1579	1615	1647	1951	2078	do.....	467
	563	720	753	803	812	843	862	885	903	984	1111	1157	Sheared fibers; specimen worm-eaten.....	462
	485	640	662	671	685	708	712	721	739	749	852	903	do.....	462
	2767	5851	7417	8097	8698	9117	9426	9707					Split along grain at ends.....	475
	2926	5829	7187	8210	8890	9208	9435	9753	10024	10433	11431		Split at ends; sheared fibers.....	475
	2087	3697	4060	4332	4559	4799	4899	5035	5193				Sheared fibers.....	452
	2531	4309	4831	5035	5343	5670	5851	5974	6110	6985	7349		do.....	452
	1179	1792	2023	2182	2363	2490	2608	2694	2790	2980	3639	3878	Slight shearing of fibers.....	289
	1551	2518	2835	3075	3289	3429	3602	3769	3910	4073	4831	5171	Sheared fibers.....	289
	1243	1805	1905	1973	2032	2123		2232	2313	2395	2744	2948	do.....	484
	885	1343	1452	1533	1583	1687	1774	1805	1846	1878	2200	2449	do.....	484
	1338	1987	2186	2390	2576	2726	2840	2971	3065	3166	3900	4287	do.....	333
	1202	1678	1882	2014	2136	2254	2350	2390	2536	2717	3193	3674	do.....	333

TABLE V.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Longitudinal compression: Ultimate strength, in kilos.	Remarks.
RHAMNACEÆ.							
42. <i>Reynoldsia latifolia</i>	454	Florida	Upper Metacombe Key.	A. H. Curtiss	Coral	13426	Split along grain
<i>Red Iron Wood. Darling Plum.</i>	454	do	do	do	do		
43. <i>Condalia ferrea</i>	460	do	do	do	do	13290	Fibers crushed at knots and split along grain.
<i>Black Iron Wood.</i>	460	do	do	do	do	12406	Fibers crushed at knots 64 millimeters from end.
45. <i>Rhamnus Caroliniana</i>	803	do	Saint John's river.	do	Rich hummock	7112	Fibers crushed at 140 millimeters from end.
<i>Indian Cherry.</i>	803	do	do	do	do		
47. <i>Rhamnus Purshiana</i>	993	Oregon	Portland	G. Engelmann and C. S. Sargent.	Rich, alluvial	9934	Fibers crushed near middle
<i>Bearberry. Bear Wood. Skittim Wood.</i>							
SAPINDACEÆ.							
50. <i>Æsculus glabra</i>	297	Missouri	Allenton	G. W. Letterman	Rich, moist	4531	Fibers crushed at 76 and at 102 millimeters from end; split along grain.
<i>Ohio Buckeye. Fetid Buckeye.</i>	297	do	do	do	do	4985	Fibers crushed near middle; grain wavy.
	386	do	do	do	Alluvial	5344	Fibers crushed at 19 millimeters from end.
	386	do	do	do	do	5207	Fibers crushed at 127 millimeters from end.
52. <i>Æsculus Californica</i>	684	California	Marin county	G. R. Vasey	Rich upland	6328	Fibers crushed at 51 millimeters from end.
<i>California Buckeye.</i>	684	do	do	do	do	5044	Fibers crushed at middle in vicinity of knot.
53. <i>Ungnadia speciosa</i>	944	Texas	New Braunfels	C. Mohr	Limestone	6305	Crushed at knot 51 millimeters from end.
<i>Spanish Buckeye.</i>	944	do	do	do	do		
54. <i>Sapindus marginatus</i>	305	do	Dallas	J. Reverchon	Rich, damp	7802	Fibers crushed near middle
<i>Wild China. Soapberry.</i>	307	do	do	do	do	8641	Triple flexure; developed intersecting "Cooper lines" at middle.
	928	do	Austin	C. Mohr	Limestone	7212	Triple flexure; deflected about 16 millimeters without crushing.
	928	do	do	do	do	6436	Triple flexure; split along grain.
56. <i>Hypelate paniculata</i>	463	Florida	Upper Metacombe Key.	A. H. Curtiss	Coral	10931	do
<i>Ink Wood. Iron Wood.</i>	463	do	do	do	do	10387	Fibers crushed at 102 millimeters from end; angle of crushing, 60°.
57. <i>Hypelate trifoliata</i>	464	do	do	do	do	6872	Deflected and split along grain
<i>White Iron Wood.</i>	464	do	do	do	do	7167	Triple flexure
59. <i>Acer macrophyllum</i>	282	Oregon	Portland	G. Engelmann and C. S. Sargent.	Rich, alluvial	6541	Fibers crushed at knot 76 millimeters from end.
<i>Broad-leaved Maple.</i>	982	do	do	do	do	6418	Triple flexure; knot near end
	1023	do	Portland Furniture Company.	do		5520	Fibers crushed at 25 and at 76 millimeters from end.
	1023	do	do	do		5919	Fibers crushed near middle and near end.
61. <i>Acer circinatum</i>	1013	do	Portland	do	Moist, alluvial		
<i>Vine Maple.</i>	1014	Washington territory.	Wilkeson	do	do	7349	Fibers crushed at knot at middle
64. <i>Acer saccharinum</i>	298	Missouri	Allenton	G. W. Letterman	Rich upland	10931	Fibers crushed near middle and at 25 millimeters from end.
<i>Sugar Maple. Sugar Tree. Hard Maple. Rock Maple.</i>	298	do	do	do	do		
	299	do	do	do	do	11186	Fibers crushed at 127 millimeters from end.
	376	Vermont	Charlotte	C. G. Pringle	Gravelly	8302	Fibers crushed at 76 millimeters from end.
	376	do	do	do	do		
	409	New England	Charlestown Navy-yard.	S. H. Pook		9063	Fibers crushed at 25 millimeters from end.
	1233	Vermont	Charlotte	F. H. Horsford		11000	Fibers crushed at 39 millimeters from end at 6 millimeters knot.
	1233	do	do	do		9970	Fibers crushed at end; cross-grained.
	1234	do	do	do		9957	Fibers crushed at end
	1234	do	do	do		9707	Fibers crushed at 25 millimeters from end.
	1235	do	do	do		8890	Fibers crushed at 70 millimeters from end; specimen split before testing.

UNITED STATES UNDER COMPRESSION—Continued.

Direction of grain.	PRESSURE, IN KILOGRAMS, REQUIRED TO PRODUCE AN INDENTATION, IN MILLIMETERS, OF—												Remarks.	Office number.
	0.25	0.51	0.76	1.02	1.27	1.52	1.78	2.03	2.28	2.54	2.81	3.08		
	3016	6260	7892	8890	9753	10524	11158	11658	12112	12610			Sheared fibers; split at ends	454
	3062	6396	8119	9617	10705	11865	12383	13109	13789	14470	17237		do.....	454
	3447	6033	7689	9004	10115	11113	11612	12474	13245	13903	17509		do.....	460
	2994	6260	8097	9480	10660	11794	12337	13789	14765	15603			do.....	460
	1338	1837	2032	2073	2195	2218	2304	2390	3468	2518	2880		do.....	803
	1247	1860	1982	2073	2153	2214	2277	2309	2350	2386	2858	3039	Sheared fibers.....	803
	1656	2404	2758	2980	3075	3202	3280	3339	3352	3470	3765		Slightly sheared fibers; split at ends	993
	490	796	885	921	971	1021	1071	1116	1157	1179	1429		Sheared fibers slightly	297
	975	1107	1216	1247	1298	1347	1397	1429	1515	1547	1778	1928	Sheared fibers; split at end	297
	680	1179	1452	1579	1701	1733	1801	1846	1928	1987	2331	2658	Sheared fibers.....	684
	1066	1433	1569	1651	1742	1810	1892	1973	2068	2127	2540	2835	do.....	684
	1061	1592	1801	1896	2078	2109	2195	2259	2327	2431	2744	3166	do.....	944
	1225	2182	2495	2576	2699	2785	2867	2939	2976	3075			Sheared fibers; split at ends	944
	1769	3158	3289	3520	3742	3874	4037	4155	4284	4468	5262	5738	Sheared fibers.....	307
	2073	3108	3348	3520	3701	3874	4037	4241	4355	4518	5298	6033	Slight shearing of fibers	307
	2109	3692	4513	4914	5307	5398	5579	5974	6105	6214	7190	7553	do.....	928
	2064	3366	3955	4404	4649	4872	5062	5216	5425	5566	6759	7439	do.....	928
	2767	4944	5715	6078	6419	6759	6931	7158	7376	7557	8210	8346	Sheared fibers; split at sides	464
	1796	3901	4899	5489	5874	6205	6432	6654	6895	7167	8074	8641	do.....	464
	1796	2313	2422	2513	2604	2672	2812	2835	2926	2980	3289	3583	Sheared fibers.....	982
	1950	2617	2785	2903	3012	3116	3216	3248	3339	3493	3792	4105	do.....	982
	1633	2313	2495	2563	2613	2703	2767	2858	2985	3012	2976	3697	do.....	1023
	1338	1765	1906	2073	2159	2282	2359	2381	2440	2504	2880	3129	do.....	1023
	1590	2576	2835	3085	3252	3420	3574	3701	3828		4854	5307	do.....	1013
	1225	2168	2758	3039	3157	3280		3674	3720	3747	4468	5085	Slight shearing of fibers	1014
	2849	4355	4944	5307	5579	5874	6023	6101	6260	6328	7530	8074	Sheared fibers.....	298
	2359	3357	3720	4060	4287	4513	4740	4899	4967				Slight shearing of fibers; split at end	298
	2858		4626	4967	5262	5489	5693	5906	6101	6328	7485	7802	Sheared fibers.....	299
	2064	3157	3398	3606	3751	3892	4060	4160	4228	4364	4899	5239	do.....	376
	2087	2926	3239	3420	3602	3748	3819	3983	4055	4164	4763	5194	do.....	376
	1724	2313	2472	2613	2794	2849	2944	3057	3107	3202	3647	3901	do.....	409
	2250	3302	3606	3801	3983	4169	4287	4418	4581	4708	5443	5987	do.....	1233
	2132	2767	3057	3257	3357	3502	3611	3706	3801	3969	4436	4899	do.....	1233
	2586	3583	3792	4060	4246	4382	4423	4495	4581	4699	5648	6169	do.....	1234
	2540	4128	4653	4745	4940	5149	5330	5434	5530	5625	6260	6736	do.....	1234
	2540	3652	3928	4069	4200	4332	4427	4522	4672	4831	5579	6078	do.....	1236

TABLE V.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Longitudinal compression: Ultimate strength, in kilos.	Remarks.
64. <i>Acer saccharinum</i> —continued..	1235	Vermont.....	Charlotte.....	F. H. Horsford.....		9970	Fibers crushed at end.....
64. <i>Acer saccharinum</i> , var. <i>nigrum</i> . <i>Black Sugar Maple</i> .	213	do	do	C. G. Pringle.....	Clay	8890	Fibers crushed at 10 millimeters knot 51 millimeters from end.
	213	do	do	do	do		
	274 ^a	Missouri.....	Allenton.....	G. W. Letterman..	Low, alluvial	10070	Fibers crushed at 76 millimeters from end.
	274 ^a	do	do	do	do		
	757	Florida	Chattahoochee...	A. H. Curtiss	Clay	8255	Triple flexure; developed intersecting "Cooper lines".
	757	do	do	do	do	6895	Fibers crushed at knot 102 millimeters from end.
	1167	Missouri.....	Allenton.....	G. W. Letterman..	Rich, alluvial	9321	Fibers crushed at middle; developed "Cooper lines".
	1167	do	do	do	do	9390	Triple flexure; "Cooper lines".
65. <i>Acer dasycarpum</i> <i>Soft Maple. White Maple. Silver Maple.</i>	1052	Massachusetts....	Topsfield.....	J. Robinson	Low meadow	7711	Fibers crushed at 51 and at 76 millimeters from end; knots.
66. <i>Acer rubrum</i> <i>Red Maple. Swamp Maple. Soft Maple. Water Maple.</i>	20	do	Arnold Arboretum	C. S. Sargent.....	Drift	7688	Fibers crushed near middle.....
	20	do	do	do	do		
	530	Mississippi	Kemper's mill	C. Mohr.....	Rich, swampy.....	7303	Fibers crushed at 89 millimeters from end.
	530	do	do	do	do	8437	Fibers crushed at 25 millimeters from end.
	743	Georgia	Bainbridge	A. H. Curtiss	Low	7235	Fibers crushed near middle.....
	743	do	do	do	do	6759	Fibers crushed at middle and at end.
	878	Massachusetts....	Danvers	J. Robinson	do	6903	Triple flexure; "Cooper lines".
	878	do	do	do	do	7380	Triple flexure; "Cooper lines"; middle bend 115 millimeters from end.
	1048	do	North Reading	do	do	7485	Fibers crushed at 89 millimeters from end.
	1048	do	do	do	do	7371	Fibers crushed at 25 and at 102 millimeters from end.
67. <i>Negundo aceroides</i> <i>Box Elder. Ash-leaved Maple.</i>	290	Missouri.....	Allenton.....	G. W. Letterman..	Rich bottom	5080	Fibers crushed at 76 millimeters from end.
	290	do	do	do	do	5298	Fibers crushed near middle and split longitudinally.
	311	Texas	Dallas	J. Reverchon	do	5148	Fibers crushed at 25 millimeters from middle.
	311	do	do	do	do	5080	Fibers crushed at middle.....
68. <i>Negundo Californicum</i> <i>Box Elder.</i>	645	California.....	Contra Costa	G. R. Vasey.....	Rich, moist	6613	Fibers crushed at knot near middle.
	645	do	do	do	do	7530	Fibers crushed at 32 millimeters from middle.
ANACARDIACEÆ							
71. <i>Rhus copallina</i> <i>Dwarf Sumach.</i>	736	Florida	Chattahoochee...	A. H. Curtiss	Dry clay		
	736	do	do	do	do	6033	Fibers crushed at knot 76 millimeters from end.
71. <i>Rhus copallina</i> , var. <i>lanceolata</i> ..	830	Texas	Dallas	J. Reverchon	Dry, gravelly	7666	Fibers crushed at 89 millimeters from end at 3 millimeters from knot.
	830	do	do	do	do		
73. <i>Rhus Metopium</i> <i>Poison Wood. Coral Sumach. Mountain Manchineel. Plum Wood. Hog Plum. Doctor Gum.</i>	467	Florida	Upper Metacombe Key.	A. H. Curtiss.....	Coral	7847	Fibers crushed at knot 51 millimeters from middle.
	467	do	do	do	do	9199	Fibers crushed at 127 millimeters from end.
LEGUMINOSÆ							
77. <i>Robinia Pseudacacia</i> <i>Locust. Black Locust. Yellow Locust.</i>	405		Charlestown Navy-yard.	S. H. Pook		13426	Fibers crushed at end and at 76 millimeters from end.
	405		do	do		12996	Fibers crushed near end and split along grain.
	441		do	do		10229	Fibers crushed at 10 millimeters from knot and at 51 millimeters from end.
	441		do	do			
	815	West Virginia....	Grafton.....	C. G. Pringle		12497	Fibers crushed in vicinity of knots
	815	do	do	do		8369	
	1247	New York	Long Island	M. C. Beedle		9934	Fibers crushed in vicinity of small knot at end.
	1247	do	do	do		11294	Fibers crushed in vicinity of small knot 102 millimeters from end.
	1248	do	do	do		10274	Fibers crushed at end of specimen
	1248	do	do	do		11385	do

UNITED STATES UNDER COMPRESSION—Continued.

Direction of grain.	PRESSURE, IN KILOGRAMS, REQUIRED TO PRODUCE AN INDENTATION, IN MILLIMETERS, OF—												Remarks.	Office number.
	0.25	0.51	0.76	1.02	1.27	1.52	1.78	2.03	2.28	2.54	2.81	3.08		
	1941	2703	2971	3175	3379	3584	3656	3788	3933	4037	4672	5358	Sheared fibers	1235
	2313	3357	3747	3924	4110	4332	4478	4617	4781	4890	5715	6283	do.....	213
	1542	2-99	3392	3579	3751	3907	4028	4164	4336	4441	5262	5715	do.....	213
	2812	4219	4559	4844	5080	5262	5401	5543	5670	5784	6396	6849	do.....	274*
	1860	2744	2948	3106	3334	3465	3615	3801	3892	4028	4808	5208	do.....	274*
	2087	3379	3611	3837	4046	4173	4377	4527	4581	4753	5353	5851	do.....	757
	2223	3243	3475	3652	3792	3907	4019	4069	4300	4346	4944	5389	do.....	757
	3026	3978	4377	4599	4744	4962	5080	5239	5339	5452	6328	6782	do.....	1167
	1951	2840	3066	3329	3479	3629	3797	3933	4033	3724	4831	4881	do.....	1167
	1315	2250	2376	2790	2899	3035	3157	3293	3397	3520	4219	4717	do.....	1062
	1951	2518	2703	2840	2985	3085	3252	3381	3538	3615	4264	4536	do.....	20
	2540	3837	4327	4500	4622	4753	4872	4985	5203	5298	6028	6568	do.....	20
	2073	2976	3248	3379	3479	3606	3742	3833	3928	4024	4626	4879	do.....	530
	1533	1896	2041	2123	2254	2427	2449	2472	2567	2685	3175		Sheared fibers; split at end	530
	1179	1633	1715	1842	1887	1946	2028	2087	2141	2177	2581	2722	Sheared fibers	743
	1361	1792	1880	1932	2028	2141	2214	2254	2350	2400	2722	3039	do.....	743
	1706	2649	2803	3016	3121	3257	3384	3479	3665	3760	4332	4808	do.....	878
	1469	2032	2268	2427	2449	2580	2703	2776	2890	2935	3493	3742	Slight shearing of fibers	878
	1805	2259	2386	2508	2613	2758	2794	2939	107	3198	3652	4037	do.....	1048
	1361	2168	2404	2563	2703	2794	2858	2971	3039	3139	3901	4332	do.....	1048
	862	1071	1166	1315	1352	1433	1463	1556	1619	1678	1973	2177	Sheared fibers	290
	984	1447	1610	1624	1710	1774	1855	1860	1905	1951	2168	2308	do.....	290
	993	1352	1479	1529	1574	1660	1683	1733	1774	1819	2064	2223	do.....	311
	1542	2250	2449	2486	2490	2531	2658	2672	2790	2794	3063	3311	do.....	311
	1179	1479	1574	1633	1683	1801	1896	1982	2005	2082	2295		Sheared fibers; split at end	645
	1202	1520	1619	1674	1756	1860	1951	1996	2073	2114	2449	2790	Sheared fibers	645
	1184	1397	1479	1551	1637	1665	1733	1801	1805	1835	2067	2177	Sheared fibers; split at end	736
	1270	1610	1678	1805	1851	1969	1991	2046	2100	2150	2381		do.....	736
	1111	1438	1637	1765	1878	1946	2059	2150	2159	2159	1973	2449	Sheared fibers; split at end	330
	1089	1609	1928	2028	2104	2168	2177	2218	2268	2259			do.....	330
	1384	1928	2395	2749	2830	3193	3293	3434	3525	3570	3969	4346	Sheared fibers; crushed obliquely at each end	467
	1633	2858	3289	3529	3765	3901	4060	4095	4296	4395	4944	5353	Sheared fibers; split at sides	467
	2926	3856	4346	4740	5126	5353	5706	5920	6283	6396	7485	8074	Sheared fibers; split at end	405
	2676	3878	4155	4436	4763	5058	5416	5615	5869	6056			Slight shearing; split at end	405
	2291	3343	3742	4064	4296	4522	4699	4844	4980	5112	5987		Sheared fibers; split at end	441
	2944	4264	4559	4808	4967	5103	5307	5421	5534	5670			do.....	441
	771	2019	2722	3016	3108	3207	3343	3438		3620	4037		do.....	815
	1814	2341	2486	2622	2703	2835	2894	2926	3094	3216	3765	4037	Sheared fibers	815
	2132	3257	3615	3783	3978	4069	4264	4386	4500	4604			Slight shearing; split at end	1247
	1905	3243	3674	3878	4060	4237	4377	4482	4604	4662			Slight shearing; split at ends	1247
	2223	3302	3606	3792	3963	4160	4346	4427	4608	4672	5307		Sheared fibers; split at end	1244
	2495	3529	4128	4219	4332	4513	4626	4831	4967	5048			Sheared fibers; split at ends	1248

TABLE V.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Longitudinal compression: Ultimate strength, in kilos.	Remarks.
79. <i>Robinia Neo-Mexicana</i> . <i>Locust</i> .	1081	Colorado	Trinidad	W. B. Strong	Low, moist	10931	Fibers crushed at 51 millimeters from end.
	1081	do	do	do	do		
80. <i>Olneya Tesota</i> . <i>Iron Wood. Arbol de Hierro</i> .	650	California	Lower Colorado valley.	G. Engelmann and C. S. Sargent.	Dry, gravelly		
	650	do	do	do	do	5851	Split along grain; oblique fracture
81. <i>Piscidia Erythrina</i> . <i>Jamaica Dogwood</i> .	564	Florida	Upper Metacombe Key.	A. H. Curtiss	Coral	9548	Shattered one end; cross-grained.
	564	do	do	do	do		
82. <i>Cladrastis tinctoria</i> . <i>Yellow Wood. Yellow Ash. Gopher Wood</i> .	83 ^a	Kentucky	Mercer county	W. M. Linney	Limestone	9924	Crushed near middle and at end
	33	do	do	do	do		
	489	Tennessee	Nashville	A. Gattinger	Alluvial	7167	Crushed at end and at 102 millimeters from end.
84. <i>Sophora affinis</i>	329	Texas	Dallas	J. Reverchon	Dry, calcareous	7734	Split along grain from end to end.
	982	do	Austin	C. Mohr	do	10524	Triple flexure; took reversed bend near middle.
85. <i>Gymnocladus Canadensis</i> . <i>Kentucky Coffee Tree. Coffee Nut</i> .	296	Missouri	Allenton	G. W. Letterman	Low, rich	9208	Crushed at 13 and at 114 millimeters from end.
	519	Tennessee	Nashville	A. Gattinger	Limestone	6169	Crushed and split along grain in vicinity of knots.
	1241	Missouri	Allenton	G. W. Letterman	Alluvial	5874	
	1242	do	do	do	do	5424	Crushed at end and at 102 millimeters from end.
	1243	do	do	do	do	5343	Triple flexure
86. <i>Gleditsia triacanthos</i> . <i>Honey Locust. Black Locust. Three-thorned Acacia. Sweet Locust. Honey Shucks</i> .	53 ^a	do	do	do	Low, rich	8119	do
	53 ^a	do	do	do	do	7711	Crushed at 76 millimeters from end.
	444	Tennessee	Nashville	A. Gattinger	Dry, sandy barren	8174	Triple flexure
87. <i>Gleditsia monosperma</i> . <i>Water Locust</i> .	760	Florida	Chattahoochee	A. H. Curtiss	Alluvial	9889	Crushed at 89 millimeters from end at 10 millimeters from knot.
	760	do	do	do	do	8799	Crushed at middle at knot 5 millimeters in diameter.
88. <i>Parkinsonia Torreyana</i> . <i>Green-bark Acacia. Palo Verde</i> .	678	Arizona	Lower Colorado river.	G. Engelmann and C. S. Sargent.	Sandy	6078	Crushed at 6 millimeters knot at middle.
	678	do	do	do	do	7280	Crushed at 64 millimeters from end.
91. <i>Cercis Canadensis</i> . <i>Redbud. Judas Tree</i> .	436	Tennessee	Nashville	A. Gattinger	Limestone	8119	Crushed at knot near middle
	1089	Missouri	Allenton	G. W. Letterman	Rich	8369	Crushed near middle
	1090	do	do	do	do	6704	Crushed at knot 102 millimeters from end.
	1091	do	do	do	do	6849	Crushed in vicinity of knots at middle.
93. <i>Prosopis juliflora</i> . <i>Mesquit. Algaroba. Honey Locust. Honey Pod</i> .	680	Arizona	Tucson	C. S. Sargent		9934	Split along grain from end to end; crushed near middle.
	680	do	do	do		10841	Triple flexure
	927	Texas	Austin	C. Mohr	Rich, calcareous	7462	Crushed near middle; opened cracks along grain; split before testing.
94. <i>Prosopis pubescens</i> . <i>Screw Bean. Screw-pod Mesquit. Tornilla</i> .	658	California	Fort Yuma	G. Engelmann and C. S. Sargent.	Sandy	10034	Crushed at 6 millimeters knot at middle.
	658	do	do	do	do	11431	Crushed near end; cross-grained
98. <i>Acacia Greggii</i> . <i>Cat's Claw</i> .	697	Arizona	Santa Rita mountains.	do	Dry, gravelly	11885	Split along grain from end to end.
100. <i>Lysiloma latifolia</i> . <i>Wild Tamarind</i> .	509	Florida	Boca Chica Key	A. H. Curtiss	Coral	7053	Crushed near middle
	1112	do	Key Largo	do	do	8337	Crushed at 10 millimeters knot 102 millimeters from end.
	1112	do	do	do	do		
ROSACEÆ.							
102. <i>Chrysobalanus Icaco</i> . <i>Cocoa Plum</i> .	480	do	Bay Biscayne	do	Swampy		
103. <i>Prunus Americana</i> . <i>Wild Plum. Canada Plum. Horse Plum</i> .	68	Missouri	Allenton	G. W. Letterman	Rich upland	8063	Crushed at 25 millimeters from end in vicinity of small knots.
	68	do	do	do	do	8799	Crushed at 19 and at 89 millimeters from end.
	334	Texas	Dallas	J. Reverchon	Rich	10796	Crushed at 102 millimeters from end and at end.
104. <i>Prunus angustifolia</i> . <i>Chickasaw Plum. Hog Plum</i> .	435	Tennessee	Nashville	A. Gattinger	River bluff	6441	

UNITED STATES UNDER COMPRESSION—Continued.

Direction of grain.	PRESSURE, IN KILOGRAMS, REQUIRED TO PRODUCE AN INDENTATION, IN MILLIMETERS, OF—												Remarks.	Office number.
	0.23	0.51	0.76	1.00	1.27	1.52	1.78	2.03	2.28	2.54	2.81	3.08		
	1905	3498	3878	4155	4427	4614	4799	5013	5126	5298	5924	6192	Sheared fibers	1081
	2041	3557	3910	4037	4241	4400	4559	4786	4808	4831	5602		Sheared fibers; split at end	1031
	1860	3765	5398	6849	7983	9526	10614	11521	12474	13245			Slight shearing; split at end	650
	4309	8256	10897	11885	12973	14062	14787	15467	16057	16510	18008		do	654
	2586	4173	4695	5126	5353	5693	5920	6033	6237	6386			Sheared fibers; opened seasoning crack	564
	2313	3992	4626	5013	5420	5702	6010	6237	6396	6600	7530		Sheared fibers; split at end	564
	1656	2078	2250	2350	2440	2540	2703	2762	2840	2985	3402	3756	Sheared fibers	38
	2518	2840	3216	3348	3434	3543	3624	3720	3847	3946	4527	4926	do	57
	2676	2427	4808	5035	5262	5439	5715	5897	6074	6260	7190	7756	Sheared fibers	859
	2041	3720	4581	5062	5434	5761	5874	6105	6559	6655	7576	7847	Sheared fibers; split at end; compressed area contained 3 millimeters knot	858
	1610	2926	3153	3357	3529	3697	3847	3933	4105	4209	4695	5162	Sheared fibers; split at end	110
	1656	2028		2254	2322	2472	2540	2608	2676	2835	5289	3652	Sheared fibers	1341
	1429	1928	2114	2282	2369	2427	2522	2622	2672	2758			Slight shearing; split at ends	1242
	1352	1787	1928	1991	2082	2168	2245	2318	2390	2454	2880		Sheared fibers; split at ends	1243
	1796	2245	2313	2404	2495	2654	2812	2939	3075	3180	3720		Slight shearing; split at end	53 ^t
	1170	1905	2168	2268	2404	2495	2608	2694	2776	2862	3357		Sheared fibers; split at end	53 ^b
	2041	2903	2903	3071	3193	3379	3561	3847	3946	4014		5512	Sheared fibers	444
	2182	3674	4219	4332	4577	4763	5035	5162	5252	5389	5934	6350	do	760
	2019	3447	3856	4073	4264	4468	4658	4799	5026	5203	6078	6350	do	760
	1452	2223	2699	2948	3620	3298	3484	3652	3801	3896	4527	5080	Sheared fibers and splintered at 6 millimeters knot	678
	1833	2939	3266	3479	3020	3701	3788	3901	4024	4082	4717	5035	Sheared fibers	678
	1116	1769	2132	2395	2663	2844	3044	3207	3357	3538	4173	4527	Slight shearing, caused by uneven loading; seasoning crack	466
	1760	2291	2522	2703	2880	3026	3184	3298	3388	3520	4201	4699	Sheared fibers	1089
	1638	2527	2794	3057	3252	3352	3538	3583	3810	3882	4536		Sheared fibers; split at end	1090
	1452	2263	2676	2767	2875	3071	3153	3293	3343	3425	3765	4105	Slight shearing	1091
	2713	4219	4518	4790	4931	5216	5513	5648	5802	6028	6849	7394	Sheared fibers	680
	2132	3007	4468	4699	4990	5103	5330	5425	5521	5702	6486		Sheared fibers; split at end	680
	3611	5874	6214	6332	6532	6804	6967	7067	7117	7244	8210	8483	Sheared fibers	827
	1769	4105	4604	4967	5239	5421	5738	5860	6169	6214	6940	7508	do	658
	2381	3924	4536	5080	5294	5625	5851	5965	6114	6314	7849	7963	do	658
	1452	1860	1928	2037	2159	2241	2331	2368	2445	2496	2880	3130	Sheared fibers	509
	1497	2341	2685	2821	2967	3066	3171	3248	3339	3438	3937	4241	do	1112
	1724	2654	2858	2930	3107	3252	3357	3456	3606	3652	4196	4491	do	1112
	1724		3153	3484	3538	3810	3892	4150	4241	4355	5058		Sheared fibers; split at end	489
	1860	2522	2880	3153	3162	3257	3343	3484	3579	3697	4241	4473	Sheared fibers	68
	1796	2449	2703	2890	3039	3221	3356	3443	3620	3683	4219	4672	do	68
	2132	3198	3620	3628	4014	4150	4332	4527	4662	4795	5670		Sheared fibers; split at end	334
	1124	1565	1801	1982	2182	2291	2359	2413	2540	2608	3130	3536	Sheared at corner; 6 millimeters knot	426

TABLE V.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Longitudinal compression: Ultimate strength, in kilos.	Remarks.
105. <i>Prunus Pennsylvanica</i> . <i>Wild Red Cherry. Pin Cherry. Pigeon Cherry.</i>	233	Vermont	Charlotte	C. G. Pringle	Cold, gravelly	6532	Crushed at 3 millimeters knot 102 millimeters from end.
	233	do	do	do	do	6486	Crushed at 102 millimeters from end.
106. <i>Prunus umbellata</i> . <i>Sloe. Black Sloe.</i>	606	Georgia	Altamaha river	A. H. Curtiss	Clay	7960	Cross-grained; split at end
	606	do	do	do	do		
107. <i>Prunusemarginata</i> , var. <i>mollis</i> .	968	Washington territory.	Wilkeson	G. Engelmann and C. S. Sargent.	Low, rich	8051	Crushed fibers near middle
	968	do	do	do	do	6603	Crushed at 6 millimeters knot 25 millimeters from end.
108. <i>Prunus serotina</i> . <i>Wild Black Cherry. Bum Cherry.</i>	15	Massachusetts	Roxbury	C. S. Sargent	Gravelly	10138	Failed at knot at end and split along grain.
	15	do	do	do	do	10256	Triple flexure.
	115	Michigan	Lansing	W. J. Beal	do	7235	Crushed at middle at 3 millimeters knot.
	115	do	Dansville	do	do	8732	Crushed at 32 millimeters from end.
	127	Missouri	Allenton	G. W. Letterman	Rich loam	9979	Crushed at 25 millimeters from middle.
	127	do	do	do	do	8890	Triple flexure
	148	Illinois	Waukegan	R. Douglas	Gravelly	9199	Crushed at middle and at end
	317	Michigan	Hersey	W. J. Beal	Rich	7802	Crushed at end
	317	do	do	do	do	8324	Crushed at 127 millimeters from end.
	368	Vermont	Charlotte	C. G. Pringle	Gravelly	9085	Crushed at middle
	406	Virginia or Middle States.	Charlestown Navy-yard.	S. H. Pook		6504	Crushed at end and split obliquely along grain.
	763	Florida	Chattahoochee	A. H. Curtiss	Clay	8773	Crushed at 51 millimeters from end and split obliquely along grain.
	763	do	do	do	do	9571	Crushed fibers at 38 millimeters from end.
	1053	Massachusetts	Topsfield	J. Robinson	Gravelly	8029	Crushed fibers near middle of one side of specimen.
	1053	do	do	do	do	8009	Triple flexure
109. <i>Prunus Capuli</i> . <i>Wild Cherry.</i>	418	New Mexico	Pinos Altos mountains.	E. L. Greene	Alluvial	8809	Split along grain from end to end; knotty.
	418	do	do	do	do		
110. <i>Prunus demissa</i> . <i>Wild Cherry.</i>	637	California	Strawberry valley	G. Engelmann and C. S. Sargent.	Low, rich	8165	Crushed at middle; 3 millimeters knot.
111. <i>Prunus Caroliniana</i> . <i>Wild Orange. Mock Orange. Wild Peach.</i>	1032	Florida	Jacksonville	A. H. Curtiss	Sandy	7847	Crushed near middle at small knots.
	1062	Texas	Victoria	C. Mehr	Rich, moist	9503	Crushed at 16 millimeters knot 102 millimeters from end.
	1062	do	do	do	do	9617	Shattered at end; cross-grained
113. <i>Prunus ilicifolia</i> . <i>Islay.</i>	1158	California	Santa Cruz	C. L. Anderson		8709	Crushed at ends; knot 4 millimeters in diameter.
	1158	do	do	do	do		
115. <i>Cercocarpus ledifolia</i> . <i>Mountain Mahogany.</i>	883	Utah	City Creek cañon	M. E. Jones	Rocky	10478	Cross-grained; shattered
	883	do	do	do	do		
117. <i>Pyrus coronaria</i> . <i>American Crab. Sweet-scented Crab.</i>	808	Delaware	Kiamensi	W. M. Canby	Clay	7485	Triple flexure.
	808	do	do	do	do	7756	Crushed at knot 6 millimeters in diameter 127 millimeters from end.
	1087	Pennsylvania	Nazareth	J. Henry	Moist	5851	Crushed at knot 10 millimeters in diameter 25 millimeters from end.
	1087	do	do	do	do	6940	Crushed 76 millimeters from end in vicinity of 4 millimeters knot.
	1088	do	do	do	do	5874	Crushed at 102 millimeters from end; cross-grained.
	1088	do	do	do	do	6328	Crushed at two 6 millimeters knots 76 millimeters from end.
120. <i>Pyrus Americana</i> . <i>Mountain Ash.</i>	214	Vermont	Charlotte	C. G. Pringle	Gravelly	6305	Crushed at middle; knots 3 millimeters in diameter.
	365	do	Huntington	do	do	5851	Crushed at 3 millimeters knot 102 millimeters from end.
121. <i>Pyrus sambucifolia</i> . <i>Mountain Ash.</i>	410	do	Mount Mansfield	do	do	6123	Triple flexure
125. <i>Crataegus arborescens</i> .	663	South Carolina	Aiken	H. W. Ravenel	Rich	6672	Crushed at 16 millimeters knots 38 and 76 millimeters from end.
	607	Georgia	Ogeechee	A. H. Curtiss	Low	3346	Crushed at middle; knot 3 millimeters in diameter.
	607	do	do	do	do	8890	Crushed at 102 millimeters from end.

UNITED STATES UNDER COMPRESSION—Continued.

Direction of grain.	PRESSURE, IN KILOGRAMS, REQUIRED TO PRODUCE AN INDENTATION, IN MILLIMETERS, OF—												Remarks.	Office number.
	0.25	0.51	0.76	1.02	1.27	1.52	1.78	2.03	2.28	2.54	2.81	3.08		
	980	1257	1347	1442	1501	1538	1628	1680	1710	1805	2078		Sheared fibers; split at end	232
	885	1483	1507	1724	1796	1932	1982	2073	2114	2164			Sheared fibers	233
	2099	4649	5280	5851	6214	6555	6759	6872	7145	5276	8029	8483	do	606
	2313	3583	4073	4527	4740	5126	5285	5416	5535				Slight shearing; split at ends	606
	975	1021	1084	1161	1216	1306	1347	1393	1470	1483	1715	1951	Sheared fibers; split at end	908
	612	1111	1216	1252	1343	1452	1483	1542	1619	1687	2019	2078	do	908
	3112	5398	6237	6646	6885	7063	7221	7326	7372	7876	8165	8618	Sheared fibers and splintered	15
	3107	5307	5489	5920	6237	6468	6646	6795	6972	7112	7530	8210	Sheared fibers	15
	1973	2241	2468	2540	2699	2749	2849	2999	3048	3094	3620	3878	do	115 ²
	2019	2649	2880	3075	3202	3334	3393	3493	3674	3788	4491	4944	Sheared fibers	127
	1588	2177	2341	2536	2667	2771	2844	3026	3071	3166	3538		Sheared fibers; split at end	127
	1837	2381	2753		2880	2880	2948	3075	3198	3202	3674	3901	Sheared fibers	317
	1384	1588	1656	1765	1860	1941	2019	2087	2173	2214			Sheared fibers; split at end	317
	1225	1660	1837	1969	2078	2164	2259	2313	2427	2481	2939	3153	do	368
	1125	2749	1483	1569	1615	1660	1710	1700	1805	1851	2159	2268	Sheared fibers	406
	1588	2078	2259	2336	2390	2513	2576	2631	2722	2753	3107	3334	do	763
	1542	2123	2350	2495	2581	2713	2853	2890	2939	3048			Sheared fibers; split at end	763
	1724	2195	2713	2812	2971	3093	3184	3302	3420	3479	4173	4581	do	1053
	2313	3538	3007	4173	4432	4717	4799	4931	5062	5208	5978	6341	Sheared fibers	1053
	2041	2708	4196	4445	4241	4999	5126	5376					Split at ends	418
	2041	3311	3738	4200	4445	4699	4940	5207	5466	5661	6740	7802	Sheared fibers; split at end	418
	2177	3221	3561	3792	3937	4087	4219	4346	4477	4531	5443		do	637
	2313	4060	4699	5216	5534	5793	6141	6386	6568	7230			Slight shearing; split at end; short specimen, 12 centimeters long.	1032
	2223	4105	4786	5307	5557	5869	6105	6292	6522	6613	7892		Sheared fibers	1063
	2132	3606	3674	3969	4191	4518	4740	4890	4990	5153	6033		Sheared fibers; split at end	1062
	3221	4513	5126	5625	5874	6205	6482	6759	7021	7235	8392	9934	Slight shearing; 4 millimeters knot	1156
	1452	2676	3198	3611	3901	4128	4445	4740	4899	5071	6305		Split at end	1156
	3153	5171	6314	7122	7372	7892	8460	8850	9199	9531	12247	13598	Sheared fibers	883
	2948	5534	6894	7462	7983	8523							Split at end	883
	1678	2404	2622	2858	3030	3198	3289	3357	3561	3674	4241	4753	Sheared fibers	808
	2395	3357				4191			4536	4649	5434	5987	do	808
	1792	2835	3171	3370	3583	3683	3856	4028	4173	4237	4990		Slight shearing; split at end	1087
	1905	3788	4513	4831	5017	5207	5334	5507	5661	5719	6396	7076	Sheared fibers; 13 millimeters knot at corner of compression area.	1087
	2449	3632	4128	4400	4572	4786	4890	5080	5298	5325	6169	6532	Slight shearing	1088
	1588	2699	3166	3470	3792	4055	4173	4318	4481	4563	5080	5693	Sheared fibers	1088
	1043	1533	1656	1751	1869	1887	2019	2096	2223	2268	2676	2812	Sheared fibers; split at ends	214
	680	1406	1678	1701	1715	1805	1941	2019	2032	2214	2608	2939	Sheared fibers	410
	957	1746	2254	2522	2703	2840	3016	3116	3248	3438	4150	4763	do	363 ²
	1905	2699	3030	3207	3438	3574	3701	3901	4024	3729	4980	5398	do	607
	1343	1082	2359	2567	2713	2830	2985	3193	3339	3407	4128		Sheared fibers; split at end	607

TABLE V.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Longitudinal compression: Ultimate strength, in kilos.	Remarks.
114. <i>Crataegus Crus-galli</i> <i>Cockspur Thorn. Newcastle Thorn.</i>	328	Massachusetts....	Brookline.....	J. Robinson.....	Loam.....	6033	Crushed at 64 millimeters from end.
	328	do.....	do.....	do.....	do.....	5806	Shattered at end.....
	1093	Missouri.....	Allenton.....	G. W. Letterman.....	Low, wet.....	7349	Triple flexure; small knots.....
	1093	do.....	do.....	do.....	do.....	8346	Crushed at 127 millimeters from end and split along grain.
123. <i>Crataegus subvillosa</i> <i>Scarlet Haw.</i>	949	Texas.....	Victoria.....	C. Mohr.....	Alluvial.....	8605	Crushed at 19 millimeters from end and along one face.
	1081	Missouri.....	Saint Louis.....	H. Eggert.....		8618	Crushed at middle at knot 3 millimeters in diameter.
129. <i>Crataegus tomentosa</i> <i>Black Thorn. Pear Haw.</i>	426	Tennessee.....	Nashville.....	A. Gattinger.....	Limestone.....	7434	Crushed at each end and at middle.
	426	do.....	do.....	do.....	do.....	6795	Crushed at middle at knot 3 millimeters in diameter.
132. <i>Crataegus spathulata</i> <i>Small-fruited Haw.</i>	926	Louisiana.....	Webster parish.....	C. Mohr.....	Clay.....	7280	Crushed at 16 millimeters knot 102 millimeters from end.
133. <i>Crataegus aestivalis</i> <i>May Haw. Apple Haw.</i>	239	South Carolina.....	Bonneau's Depot.....	H. W. Ravenel.....	Damp, rich.....	7122	Crushed at middle on one corner.....
134. <i>Crataegus flava</i> , var. <i>pubescens</i> <i>Summer Haw. Red Haw.</i>	767	Florida.....	Aspalaga.....	A. H. Curtiss.....	Dry clay.....	8437	Crushed along one corner; deflected from crushed side.
135. <i>Amelanchier Canadensis</i> <i>Juneberry. Shad Bush. Service Tree. May Cherry.</i>	241	Kentucky.....	Brumfield Station.....	W. M. Linney.....	Waverly shale.....	11294	Crushed at 115 millimeters from end.
	849	Massachusetts.....	Danvers.....	J. Robinson.....	Loam.....	10433	Sap-wood; triple flexure.....
	849	do.....	do.....	do.....	do.....	10410	Crushed at 25 millimeters from middle.
HAMAMELACEÆ.							
138. <i>Liquidambar styraciflua</i> <i>Sweet Gum. Star-leaved Gum. Liquidamber. Red Gum. Bilsted.</i>	540	Alabama.....	Kemper's mill.....	C. Mohr.....	Rich, alluvial.....	6795	Crushed near middle; angle of crushing, 50°.
	546	do.....	do.....	do.....	do.....	6486	Triple flexure.....
	1095	Arkansas.....	Little Rock.....	G. W. Letterman.....		7847	Crushed at 25 and at 102 millimeters from end.
	1095	do.....	do.....	do.....		7892	Crushed at 51 millimeters from end.
	1173	New Jersey.....	Mount Holly.....	S. P. Sharples.....	Clay.....	6337	Crushed at 64 millimeters from end and at middle on opposite face.
	1173	do.....	do.....	do.....	do.....	6306	Crushed and split at end; cross-grained.
	1181	Mississippi.....	Yazoo River bottom.....	R. Abbey.....	Alluvial.....	6940	Crushed at 64 millimeters from end; 0.75 sap-wood.
	1181	do.....	do.....	do.....	do.....	7870	Crushed at 64 millimeters from end.
	1182	do.....	do.....	do.....	do.....	8573	Crushed at middle.....
	1182	do.....	do.....	do.....	do.....	8210	do.....
	1183	do.....	do.....	do.....	do.....	7938	Crushed at 25 millimeters from middle.
	1183	do.....	do.....	do.....	do.....	8256	Crushed at middle.....
RHIZOPHORACEÆ.							
140. <i>Rhizophora Mangle</i> <i>Mangrove.</i>	485	Florida.....	Bay Biscayne.....	A. H. Curtiss.....	Salt-marsh.....	13517	Crushed at end in vicinity of knot.
	485	do.....	do.....	do.....	do.....	14016	Crushed at middle; cross-grained.
COMBRETACEÆ.							
141. <i>Conocarpus erecta</i> <i>Button Wood.</i>	489	do.....	do.....	do.....	do.....	9571	Crushed at 38 millimeters from end; worm-eaten.
	489	do.....	do.....	do.....	do.....	9614	Triple flexure.....
142. <i>Laguncularia racemosa</i> <i>White Button Wood. White Mangrove.</i>	507	do.....	Sugar-Loaf Sound.....	do.....	do.....	7212	Crushed near middle; cross-grained and worm-eaten.
	507	do.....	do.....	do.....	do.....	7167	Crushed at 76 millimeters from end; small knots.
MYRTACEÆ.							
144. <i>Eugenia buxifolia</i> <i>Gurgeon Stopper. Spanish Stopper.</i>	1118	do.....	Lost Man's river.....	do.....	Humus and coral.....	14198	Crushed at 127 millimeters from end.
	1118	do.....	do.....	do.....	do.....		
146. <i>Eugenia monticola</i> <i>Stopper. White Stopper.</i>	1135	do.....	Umbrella Key.....	do.....	Coral.....	8754	Split along grain, opening season cracks; split in seasoning cracks.
	1135	do.....	do.....	do.....	do.....	8636	Split and crushed near end.....
148. <i>Eugenia procera</i> <i>Red Stopper.</i>	1127	do.....	Miami.....	do.....	do.....	10931	Triple flexure; developed intersecting "Cooper lines".
	1127	do.....	do.....	do.....	do.....	10567	do.....
CORNACEÆ.							
451. <i>Cornus florida</i> <i>Flowering Dogwood. Box Wood.</i>	67	Missouri.....	Allenton.....	G. W. Letterman.....	Upland.....	9276	Triple flexure; opened cracks at end.
	67	do.....	do.....	do.....	do.....	9571	do.....
	761	Florida.....	Chattahoochee.....	A. H. Curtiss.....	Calcareous.....	7031	Crushed at 102 millimeters from end; cross-grained.

UNITED STATES UNDER COMPRESSION—Continued.

Direction of grain.	PRESSURE, IN KILOGRAMS, REQUIRED TO PRODUCE AN INDENTATION, IN MILLIMETERS, OF—												Remarks.	Office number.
	0.25	0.51	0.76	1.02	1.27	1.53	1.78	2.03	2.28	2.54	4.81	5.08		
	1315	2313	2495	2676	2803	2980	3085	3307	3307	3307	3988	4296	Sheared fibers	528
	1314	2155	2490	2713	2908	2976	3071	3171	3198	3484	4004	4763	do	528
	2041	3153	3656	3937	4255	4482	4605	4850	5085	5171			Slight shearing; split at end	1008
	1701	2676	3085	3379	3611	3828	3902	4119	4264	4301	5353	6033	Sheared fibers	1008
	1315	2495	3130	3484	3788	3983	4219	4391	4581	4753	4806	6350	do	949
	2449	3674	4119	4445	4626	4922	5013	5162	5439	5534	6759	7212	Sheared fibers; split at end	1061
	1787	3257	3765	4055	4309	4482	4658	4781	4944	5035	5851	6486	Sheared fibers	426
	1293	2359	2948	3130	3379	3638	3742	3892	4119	4287	4035	5625	do	426
	1315	2617	3066	3329	3484	3674	3910	4055	4164	4287	5262	5761	do	926
	1433	2699	3130	3402	3583	3742	3856	4023	4101	4264	4090	5579	Sheared fibers; split at end	239
	1905	3720	4445	4831	5108	5398	5625	5874	6056	6200	7394	7983	Sheared fibers; indented on 6 millimeters knot	767
	2041	3221	3674	3992	4423	4436	4877	4990	5194	5334	6350		Sheared fibers; split at end	241
	1860	3221	3788	4110	4491	4708	4922	5112	5316	5489	6646	7349	Sheared fibers	849
	2449	3515	3946	4291	4536	4863	5098	5303	5484	5684	6782		Sheared fibers; split at end	849
	1424	1914	2019	2100	2254	2381	2477	2536	2645	2717	3166	3329	Sheared fibers	546
	1588	1-83	2073	2223	2313	2395	2518	2604	2703	2799	3289	3620	do	546
	1334	1996	2191	2250	2277	2377	2413	2486	2527	2540	2926	3121	do	1096
	771	1315	1474	1542	1574	1674	1719	1769	1860	1932			Sheared fibers; split at end	1096
	1198	1814	1996	2096	2186	2350	2472	2531	2676	2708	3289		do	1173
	3078	2944	3026	3162	3302	3438	3561	3661	3801	3942	4445	4881	Sheared fibers	1173
	1089	1506	1588	1633	1715	1760	1928	1941	1973	2014	2313	2440	do	1181
	952	1452	1574	1669	1760	1851	1887	1941	2028	2050	2404	2531	do	1182
	1384	1792	2000		2227	2300	2350	2449	2586	2649	3075	3302	do	1182
	1043	1520	1650	1814	1914	2019	2091	2141	2291	2345	2708	2899	Sheared fibers; split at end	1182
	907	1452	1633	1923	2041	2091	2168	2273	2381	2445	2722	2926	Sheared fibers	1182
	1157	1488	1678	1837	1896	1941	2073	2155	2223	2273			Sheared fibers; split at end	1182
	1311	4173	5806	6940	7576	8006	8464	8822	9140	9453			Slight shearing; split at end	481
	2227	4763	5987	6713	7212	7689	8052	8387	8663	8913	10637		Sheared fibers; split at end	481
	2404	4173	4854	5398	5751	6078	6396	6600	6895	7085			do	481
	1903	4491	5216	5761	6101	6432	6759	7099	7226	7530	8799		do	481
	817	1515	1796	2050	2132	2359	2436	2531	2626	2790			Sheared fibers; split at end	501
	1270	1973	2250	2472	2640	2708	2844	2948	3035	3116	3674	3901	Sheared fibers	501
	2336	3765	4763	5398	5851	6169	6613	6963	7294	7576			Opened seasoning cracks	1116
	4691	5761	5942	6373	6849	7076	7303	7553	7802	8029	9526	9753	Sheared fibers; split at end	1116
	2586	4527	5648	6033	6532	6795	7258	7475	7779	7983	9026	9753	Sheared fibers	1116
	3538	5090	5761	6432	6963	7439	7792	8119	8506	8786	10614		Sheared fibers; split at end	1127
	2313	4672	5942	6668	7235	7689	8119	8419	8714	8990	10841	11930	do	1127
	2948	4581	5715	6146	6386	6754	7076	7326	7485	7729	8890	9798	Sheared fibers; 7 millimeters knot in compression surface.	67
	1860	3720	4513	4953	5180	5464	5761	5965	6192	6396	7802	8702	Sheared fibers; split at end	67
	2833	3162	3493	3765	4037	4241	4418	4563	4667	4844	5625	6169	do	761

FOREST TREES OF NORTH AMERICA.

TABLE V.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Longitudinal compression: Ultimate strength, in kilos.	Remarks.
151. <i>Cornus florida</i> —continued	812	West Virginia	Grafton	C. G. Pringle	Dry	8732	Sap-wood; triple flexure; intersecting "Cooper lines"; split at end.
	812	do	do	do	do	7621	do
	1077	Missouri	Allenton	G. W. Letterman	Gravelly	9004	Triple flexure.
	1077	do	do	do	do	8981	Triple flexure; split at ends; intersecting "Cooper lines".
	1092	do	do	do	Flinty	8210	Triple flexure; split at ends.
152. <i>Cornus Nuttallii</i> <i>Flowering Dogwood.</i>	960	Oregon	Portland	G. Engelmann and C. S. Sargent.		10387	Crushed at middle.
	960	do	do	do		10819	Crushed at 102 millimeters from end.
153. <i>Nyssa capitata</i> <i>Ogeechee Lime. Sour Tupelo. Gopher Plum.</i>	605	Georgia	Ogeechee river	A. H. Curtiss	Swampy	6895	Crushed at 114 millimeters from end; split along grain.
	605	do	do	do	do		
154. <i>Nyssa sylvatica</i> <i>Tupelo. Sour Gum. Pepperidge. Black Gum.</i>	517	Tennessee	Cumberland river	A. Gattinger		7349	Crushed on one face at 25 millimeters from middle.
	750	Florida	Chattahoochee	A. H. Curtiss	Clay	7892	Triple flexure; split at ends.
	750	do	do	do	do	8119	Crushed at 3 millimeters knots at middle.
	813	West Virginia	Grafton	C. G. Pringle		8414	Crushed near middle; split along one corner.
	813	do	do	do		8210	Crushed at 102 millimeters from end.
	823	Massachusetts	West Newbury	J. Robinson	Rich	7689	Triple flexure; split at ends.
	823	do	do	do	do	6623	Crushed at knots 64 millimeters from end.
	824	do	do	do	do	6577	Crushed at 6 millimeters knot 25 millimeters from middle.
	824	do	do	do	do	7394	Triple flexure; split at ends.
	835	do	Chebacco pond	do		7022	Triple flexure, deflected diagonally; split at ends.
	835	do	do	do		7176	Triple flexure; split at ends; intersecting "Cooper lines".
155. <i>Nyssa uniflora</i> <i>Large Tupelo. Cotton Gum. Tupelo Gum.</i>	128	South Carolina	Bonneau's Depot	H. W. Ravenel	Swampy	6896	Crushed at 12 and at 127 millimeters from end on opposite sides.
	128	do	do	do	do	6328	Triple flexure; split at ends.
	550	Alabama	Stockton	C. Mohr	Alluvial	5035	Crushed near middle.
	550	do	do	do	do	5715	Triple flexure.
	604	Georgia	Ogeechee river	A. H. Curtiss	Swampy	6123	Triple flexure 102 millimeters from end.
	604	do	do	do	do	5489	Triple flexure; split at end.
CAPRIFOLIACEÆ.							
156. <i>Sambucus glauca</i> <i>Elder.</i>	681	California	Contra Costa county.	G. R. Vasey	Gravelly	4400	Shattered at end; 10 millimeters knot.
158. <i>Viburnum Lentago</i> <i>Sheepberry. Nannyberry.</i>	370	Vermont	Hinesburg	C. G. Pringle	Swampy	8890	Crushed near middle; grain wavy.
159. <i>Viburnum prunifolium</i> <i>Black Hazz. Stag Bush.</i>	110 ⁹	Kentucky	Mercer county	W. M. Linney	Hudson River shale.	10160	Crushed at ends in vicinity of knots; split along grain.
	110 ⁴	do	do	do	Trenton limestone.	10329	Crushed 38 millimeters from middle.
	729	Georgia	Bainbridge	A. H. Curtiss	Clay	7938	Crushed 26 millimeters from middle; cross-grained.
RUBIACEÆ.							
160. <i>Exostemma Caribæum</i>	466	Florida	Upper Metacombe Key.	do	Coral	13331	Shattered from end to end.
	466	do	do	do	do	10660	Crushed at knot at middle; split along grain.
161. <i>Pinckneya pubens</i> <i>Georgia Bark.</i>	381	South Carolina	Bluffton	J. H. Mellichamp	Sandy swamp	4355	Crushed at knot 25 millimeters from middle; opened between rings.
ERIOACEÆ.							
164. <i>Vaccinium arboreum</i> <i>Farkleberry.</i>	343	Alabama	Citronelle	C. Mohr	Sandy	5874	Split and crushed at end; cross-grained.
	343	do	do	do	do	6895	Triple flexure, deflected diagonally.
165. <i>Andromeda ferruginea</i>	1083	Florida	Jacksonville	A. H. Curtiss	Hummock	7802	Crushed at 76 millimeters from end; cross-grained.
166. <i>Arbutus Menziesii</i> <i>Madroña.</i>	643	California	Contra Costa county.	G. R. Vasey	Gravelly	7303	Triple flexure; split at ends.
	643	do	do	do	do	7756	do
	679	do	Marin county	do	do	7530	Shattered and crushed at ends.
	679	do	do	do	do	9548	Crushed fibers at middle.

UNITED STATES UNDER COMPRESSION—Continued.

Direction of grain.	PRESSURE, IN KILOGRAMS, REQUIRED TO PRODUCE AN INDENTATION, IN MILLIMETERS, OF—												Remarks.	Office number.
	0.25	0.51	0.76	1.02	1.27	1.52	1.78	2.03	2.28	2.54	4.81	5.08		
	2041	3130	3493	3761	3983	4146	4355	4527	4677	4831	5761	6328	Sheared fibers.....	812
	1951	3130	3606	3892	4069	4250	4400	4536	4717	4899	5987	6396	do.....	812
	862	2291	2606	4418	4944	5325	5715	5878	6146	6419	7599	8663	do.....	1077
	1951	3856	4491	4944	5243	5552	5711	5883	6114	6373	7394		Sheared fibers; split at end.....	1077
	2223	3633	4509	4944	5158	5398	5606	5779	5929	6123	7212	7938	Sheared fibers.....	1092
	2068	2994	3843	3615	3819	4014	4132	4323	4436	4604	5489	5829	do.....	960
	1814	3198	3556	3742	3946	4164	4287	4491	4572	4708	5579	6169	do.....	960
	1406	2177	2386	2522	2713	2858	2971	3085	3143	3289	3878		do.....	605
	1452	1814		2150	2254		2449	2608	2622	3039	3153		do.....	605
	1338	1796	1941	2064	2150	2245	2331	2400	2499	2622	3016		Sheared fibers; split at ends.....	517
	1792	2767	3130		3538	3756	3892	4073	4219	4400	5307	5851	Sheared fibers.....	750
	2291	3130	3329	3606	3792	3992	4241	4423	4536	4726	5851	6350	do.....	750
	1951	3749	2921	3075	3243	3375	3438	3497	3588	3670	4196	4391	do.....	813
	1179	1352	2019	2259	2336	2404	2495	2531	2567	2617	3039	2894	do.....	813
	1633	2676	2948	3062	3243	3583	3697	3919	4055	4137	5126	5887	Sheared fibers on one edge.....	833
	1724	2495	2744	2971	3148	3307	3447	3556	3683	3810	4581	5171	do.....	833
	1257	2531	2713	2899	3030	3171	3311	3420	3502	3615	4418	5013	Slight shearing.....	834
	1384	2245	2513	2849	2921	3107	3193	3311	3484	3579	4196	4581	Slight shearing.....	834
	1901	2699	3130	3357	3515	3697	3856	3983	4105	4241	5035	5851	do.....	835
	1837	2740	3130	3311	3529	3656	3942	3946	4064	4264	5216	5615	do.....	835
	2132	2622	2875	3003	3148	3329	3475	3574	3715	3828	4401	4944	Sheared fibers.....	128
	2318	2948	3130	3207	3348	3438	3593	3742	3842	3946	4545	4944	do.....	128
	1043	1529	1588	1678	1765	1869	1960	2014	2059	2132	2436	2667	do.....	550
	1343	1905	2041	2168	2254	2381	2472	2586	2664	2708	3085	3334	do.....	550
	1497	2313	2654	2803	3012	3139	3216	3348	3425	3493	4219	4491	Sheared fibers; split at end.....	604
	1202	1610	1783	1805	1923	1996	2118	2195	2232	2308	2699	3039	Sheared fibers.....	604
	1452	1928	2068	2150	2218	2291	2379	2436	2486	2536			Sheared fibers; split at end.....	681
														370
	1996	2019	4649	4976	5252	5498	5715	5874	6033	6283			Sheared fibers; split at ends.....	110 ^a
	1882	3992	4854	5207	5513	5724	5965	6232	6441	6595			Slight shearing; split at ends.....	110 ^a
	2449	3438	3806	4033	4264	4463	4690	4840	4981	5162	6305		Sheared fibers on one edge; split at ends.....	739
	2948	5806	6940	7439	7938	8142	8460	8936	8981	9026	9889	10705	Sheared fibers; radial split.....	466
	3447	5761	6668	7067	7475	7756	7915	8074	8192	8301			do.....	466
	590	1021	1406	1574	1678	1792	1851	1896	1941	1982	2214	2422	Sheared fibers; 4 millimeters knot on indented area.....	381
	1941	3402	3833	4164	4432	4899	5171	5398	5512	6305	6918		Sheared fibers.....	343
	1905	3243	3742	4241	4522	4749	4990	5162	5302	5489	6577	7076	Slight shearing; 10 millimeters knot at edge.....	343
	1792	2699	3193	3402	3611	3638	4024	4173	4300	4491	5216	5715	Sheared fibers.....	1033
	1815	2404	2694	2967	3048	3266	3393	3574	3729	3846	4468	5035	Sheared fibers.....	643
	2087	2685	2980	3243	3443	3620	3828	3992	4119				do.....	643
	1941	2858	3243	3515	3720	3851	3996	4173	4309	4418	5126	5579	do.....	679
	1361	2313	2640	2908	3075	3243	3393	3561	3692	3810	4491		Sheared fibers; split at ends.....	679

FOREST TREES OF NORTH AMERICA.

TABLE V.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Longitudinal compression: Ultimate strength, in kilob.	Remarks.
167. <i>Arbutus Xalapensis</i>	683	Arizona	Santa Rita mount- ains.	G. Engelmann and C. S. Sargent.		7123	Crushed near middle at knots 3 millimeters in diameter.
.....	683	do	do	do		5715	Split obliquely from middle to end, crushing fibers at middle.
169. <i>Oxydendrum arboreum</i>	853	Alabama	Cottage Hill	C. Mohr	Light, rich	7430	Crushed at 10 millimeters knot at middle.
<i>Borrel Tree. Sour Wood.</i>	853	do	do	do	do	7847	Crushed at 10 millimeters knot 25 millimeters from middle.
.....	515	Tennessee	Nashville	A. Gattinger	Sandy rock	8799	Triple flexure
.....	515	do	do	do	do		
170. <i>Kalmia latifolia</i>	262 ^a	Virginia	Fancy Gap	H. Shriver	Molst	6931	Crushed at two 6 millimeters knots at end.
<i>Laurel. Calico Bush. Spoon Wood. Ivy.</i>	262 ^a	do	do	do	do	6849	Crushed at knot 51 millimeters from end.
171. <i>Rhododendron maximum</i>	263	do	do	do	do	7462	Crushed at 25 millimeters from middle.
<i>Great Laurel. Rose Bay.</i>	263	do	do	do	do	6577	Crushed at 3 millimeters knot at middle.
SAPOTACEÆ.							
175. <i>Chrysophyllum oliviforme</i>	492	Florida	Bay Biscayne	A. H. Curtiss	Coral	10433	Crushed at 25 millimeters from middle, deflecting diagonally.
.....	492	do	do	do	do	8709	Crushed at knot at middle; split obliquely.
176. <i>Sideroxylon Mastichodendron</i>	461	do	Upper Metacombe Key.	do	do	10932	Split along grain at end; slight crushing.
<i>Mastic.</i>	461	do	do	do	do	9889	Crushed at end; oblique split
177. <i>Dipholis salicifolia</i>	488	do	Bay Biscayne	do	do	11952	Triple flexure
<i>Bustia. Cassada.</i>	488	do	do	do	do	12565	Crushed near middle
.....	500	do	Umbrella Key	do	do	11272	Crushed at 64 millimeters from end and split along grain.
.....	500	do	do	do	do	10931	Crushed at 102 millimeters from end in vicinity of knots.
178. <i>Bumelia tenax</i>	746	Georgia	Bainbridge	do	Low	7235	Split obliquely from middle to end.
179. <i>Bumelia lanuginosa</i>	930	Texas	Austin	C. Mohr	Limestone	5489	Crushed at end
<i>Gum Elastic. Shittim Wood.</i>	930	do	do	do	do	5012	Crushed at 102 millimeters from end at 5 millimeters knot.
.....	1083	Missouri	Allenton	G. W. Letterman	do	6895	Deflected at middle; split at ends.
181. <i>Bumelia lycioides</i>	333	Tennessee	Nashville	A. Gattinger	Alluvial	7825	Split at end; cross-grained; season- ing crack at middle.
<i>Iron Wood. Southern Buck- thorn.</i>	1124	Florida	Boca Chica Key	A. H. Curtiss	Coral	7643	Triple flexure, deflecting from knots.
182. <i>Bumelia cuneata</i>	458	do	Upper Metacombe Key.	do	do	8913	Cross-grained; split obliquely from end to end.
<i>Ants' Wood. Downward Plum. Saffron Plum.</i>	458	do	do	do	do	5806	Cross-grained; split along season- ing cracks.
183. <i>Mimusops Sieberi</i>		do	do	do	do		
<i>Wild Dilly.</i>		do	do	do	do		
EBENACEÆ.							
184. <i>Diospyros Virginiana</i>	61	Missouri	Allenton	G. W. Letterman	Rich upland	7892	Triple flexure, deflected diago- nally.
<i>Persimmon.</i>	61	do	do	do	do	7485	do
.....	425	Tennessee	Nashville	A. Gattinger	Rich loam	9005	Crushed at end
.....	811	West Virginia	Grafton	C. G. Pringle		7394	Triple flexure
.....	811	do	do	do		7892	Triple flexure, deflected diago- nally.
.....	1084	Missouri	Allenton	G. W. Letterman	Rich upland	8301	Deflected at 23 millimeters from middle.
.....	1084	do	do	do	do	8029	do
.....	1162	do	do	do	Rich	8415	Triple flexure, deflected diago- nally; intersecting "Cooper lines".
.....	1162	do	do	do	do	7901	
STYKACEÆ.							
186. <i>Symplocos tinctoria</i>	347	Alabama	Cottage Hill	C. Mohr	Sandy	6146	Crushed at 6 millimeters knot 76 millimeters from end and at 3 millimeters knot at end.
<i>Horse Sugar. Sweet Leaf.</i>	347	do	do	do	do		
187. <i>Halosia diptera</i>	738	Georgia	Bainbridge	A. H. Curtiss	Low	6486	Failed at 6 millimeters knot 127 millimeters from end and split along grain.
<i>Snow-drop Tree. Silver-bell Tree.</i>	738	do	do	do	do	7394	Triple flexure
OLEACEÆ.							
191. <i>Fraxinus pistaciifolia</i>	660	Arizona	Santa Rita mount- ains.	G. Engelmann and C. S. Sargent.	do	6441	Crushed at middle on one face
<i>Ash.</i>	660	do	do	do	do	5874	Crushed at middle in vicinity of 3 millimeters knot.

UNITED STATES UNDER COMPRESSION—Continued.

Direction of grain.	PRESSURE, IN KILOGRAMS, REQUIRED TO PRODUCE AN INDENTATION, IN MILLIMETERS, OF—												Remarks.	Office number.
	0.25	0.51	0.76	1.02	1.27	1.52	1.78	2.03	2.28	2.54	2.81	3.08		
	1520	2586	2976	3221	3411	3629	3801	3946	4105	4219	4922	5421	Sheared fibers	683
	2359	3588	3983	4250	4482	4662	4854	4990	5112	5207	6123	6623	do.	683
	1951	3062	3479	3751	3946	4237	4414	4554	4695	4831	5670	6350	do.	358
	1211	2223	2676	2971	3153	3284	3479	3633	3760	3856	4491	5058	do.	358
	1407	2214	2527	2604	2849	3021	3193	3325	3493	3611	4287	4626	Sheared fibers; split at end	515
	1560	2254	2586	2767	2921	3003	3157	3207	3393	3438	3946		do.	675
	2268	3289	3674	3909	4105	4237	4350	4491	4581	4690	5534		do.	262*
	2223	3447	3892	4173	4287	4432	4581	4690	4808	4890	5625	6078	Sheared fibers	263*
	2041	2767	2980	3107	3252	3352	3434	3502	3597	3674	4264	4491	do.	263
	1951	2468	2708	2830	2880	2994	3085	3166	3252	3302	3810	4150	Sheared fibers; split at end	263
	4219	5579	6033	6464	6791	7145	7430	7693	7974	8219	9753		do.	492
	1800	3765	4626	5126	5425	5742	6014	6250	6532	6736	8119	9026	do.	492
	2313	4536	5171	5586	5874	6201	6382	6677	6940	7117			Slight shearing; split at end	461
	2767	4332	4890	5307	5489	5761	5987	6123	6296	6464	7248	7847	Sheared fibers; split at end; short specimen, 121 millimeters long.	461
	2087	3420	4082	4436	4717	5035	5434	5670					Split at end	468
	2177	3493	3983	4327	4608	4844	4649	5396	5479	5648	6441		Sheared fibers on one edge; split at end	468
	975	2486	3425	3810	4114	4364	4604	4808		5103			Sheared fibers; split at end	500
	907	2269	3239	3833	4073	4400	4527	4844	5013	5207	6123		do.	500
	1474	2155	2431	2712	2894	3116	3298	3497	3710	3882	4654	5002	Sheared fibers	746
	1384	1923	2055	2168	2259	2381	2472	2572	2635	2703	3121	3443	do.	930
	975	1497	1606	1692	1801	1896	1978	2073	2177	2268	2703	3085	do.	930
	1769	2835	3148	3425	3633	3942	4024	4191	4350	4518	5396	5851	Sheared fibers; split at corner	1063
	880	2522	2527	3289	3529	3751	3964	4178	4364	4536			Split at ends	333
	658	2313	3720	4281	4581	4799	4944	5207	5479	5666	6577	7462	Sheared fibers	1124
	2676	4355	5307	6023	6532	6895	7235	7521	7811	8029	8936	9163	do.	458
	2359	3697	4491	5058	5470	5851	6155	6396	6713	6940	8279	9026	do.	458
	1905	3130	3674	4001	4214	4332	4527	4667	4786	4890	5625	6146	do.	61
	2041	3402	4060	4436	4695	4971	5067	5239	5362	5534	6486	7076	do.	61
	1960	2602	4450	4908	5289	5561	5869	6083	6310	6477	7666	8392	do.	425
	2767	4536	5353	5738	6005	6260	6486	6849	7049	7235	8256	8799	do.	811
	1905	3266	3901	4341	4604	4854	5080	5285	5434	5615	6668	7530	do.	811
	1746	3402	4264	4699	5035	5307	5421	5661	5811	5965	7031	7711	do.	1084
	2676	4092	4766	5194	5484	5715	5853	6042	6187	6319	7258	7756	do.	1084
	2540	4493	5557	6005	6323	6568	6736	6881	7049	7221	8170	8663	do.	1162
	1293	3039	4000	4695	5080	5421	5606	5851	6069	6246	7417	8301	Slight shearing of fibers	1163
	1837	2608	2744	2871	2967	3003	3039	3057	3065	3162	3720	3856	Sheared fibers	347
	1170	1796	1960	2064	2127	2214	2295	2381	2445	2564	2903	3239	do.	347
	1397	2336	2753	2998	3153	3366	3497	3683	3810	3910	4491	5171	do.	738
	1802	2227	2522	2767	2948	3080	3212	3343	3501	3674	4355	4763	Sheared fibers	666
	2132	3039	3425	3529	3787	3901	4150	4382	4509	4604	5443	6078	Sheared fibers along one edge	666

TABLE V.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Longitudinal compression: Ultimate strength, in kilos.	Remarks.
192. <i>Fraxinus Americana</i> White Ash.	39 ²	Missouri	Allenton	G. W. Letterman	Low	7530	Triple flexure
	114 ¹	Michigan	Dansville	W. J. Beal	Clay	5851	do
	114 ¹	do	do	do	do	5579	Triple flexure; split at end
	114 ³	do	Hudson	do	do	9326	Deflected diagonally; crushed at 25 millimeters from middle.
	114 ⁴	do	Lansing	do	do	9934	Crushed at 25 millimeters from end.
	114 ⁵	do	Dansville	do	Clay	5761	Crushed at 38 millimeters from middle; "Cooper lines."
	130	South Carolina	Bonneau's Depot	H. W. Ravenel	Wet	9208	Crushed at 64 millimeters from end.
	130	do	do	do	do	8709	Crushed at 114 millimeters from end.
	212	Virginia	Wytheville	H. Shriver	do	7983	Crushed at middle
	212	do	do	do	do	7756	Crushed at 32 millimeters from middle.
	227 ¹	Vermont	Charlotte	C. G. Pringle	Gravelly	7281	Crushed at 51 millimeters from middle; deflected diagonally.
	227 ³	do	do	do	do	7485	Crushed at 3 millimeters knot 12 millimeters from middle.
	227 ³	do	do	do	do	7349	Triple flexure
	207 ¹	Virginia	Wytheville	H. Shriver	do	6441	Crushed at 38 millimeters from middle.
	207 ³	do	do	do	do	6509	Deflected diagonally; maximum bend 38 millimeters from middle.
	431	Tennessee	Nashville	A. Gattinger	Limestone	8346	Triple flexure
	551	Alabama	Kemper's mill	C. Mohr	Alluvial	7439	do
	551	do	do	do	do	6736	Deflected and fibers crushed at 76 millimeters from end.
	747	Georgia	Bainbridge	A. H. Curtiss	River-bottom	5057	Crushed at 3 millimeters knot at middle.
	747	do	do	do	do	5080	Triple flexure; split at ends
	937	Texas	Austin	C. Mohr	Rich, calcareous	7167	Crushed at end and at 76 millimeters from end in vicinity of knots.
	1045	Massachusetts	Reading	J. Robinson	do	8890	Triple flexure; middle bend 19 millimeters from center.
	1045	do	do	do	do	8686	do
193. <i>Fraxinus pubescens</i> Red Ash.	229 ²	Vermont	Charlotte	C. G. Pringle	Clay	8301	Failed at knot 64 millimeters from end; cross-grained.
	229 ³	do	do	do	do	8063	Triple flexure
194. <i>Fraxinus viridis</i> Green Ash.	1050	Massachusetts	Topsfield	J. Robinson	River-bottom	4513	Triple flexure, deflected diagonally.
	57	Missouri	Allenton	G. W. Letterman	Rich, wet	6486	Deflected at middle
195. <i>Fraxinus platycarpa</i> Water Ash.	308	Texas	Dallas	J. Reverchon	do	7349	Crushed at middle on one face
	308	do	do	do	do	5829	Triple flexure
	438	Tennessee	Nashville	A. Gattinger	Rich upland	8618	Crushed at 89 millimeters from end, splitting between rings.
	948	Texas	Victoria	C. Mohr	Rich, wet	8097	Deflected diagonally
	948	do	do	do	do	9753	Triple flexure; middle bend 38 millimeters eccentric.
	957	do	Matagorda bay	do	do	7892	Crushed at 6 millimeters knot 51 millimeters from end.
	957	do	do	do	do	7666	Triple flexure
196. <i>Fraxinus quadrangulata</i> Blue Ash.	536	Alabama	Stockton	do	Rich, alluvial	4014	Crushed at 76 millimeters from end.
	66	Missouri	Allenton	G. W. Letterman	Dry upland	8754	Triple flexure
	66	do	do	do	do	9299	Triple flexure, deflected diagonally.
	125	Michigan	Lansing	W. J. Beal	Rich loam	6228	do
	125	do	do	do	do	6327	Triple flexure; developed intersecting "Cooper lines".
	286 ¹	Kentucky	Mercer county	W. M. Linney	Limestone	9344	Deflected at middle and at 3 millimeters knot 76 millimeters from end.
	286 ²	do	do	do	do	8428	Crushed at 3 millimeters knot 51 millimeters from end and near middle.
	291	Missouri	Allenton	G. W. Letterman	Sandy loam	6849	Crushed on one face 19 millimeters from middle.
	518	Tennessee	Nashville	A. Gattinger	Rich limestone	8609	Crushed at knots 102 millimeters from end.

UNITED STATES UNDER COMPRESSION—Continued.

Direction of grain.	PRESSURE, IN KILOGRAMS, REQUIRED TO PRODUCE AN INDENTATION, IN MILLIMETERS, OF—												Remarks.	Office number.
	0.25	0.51	0.76	1.02	1.27	1.52	1.78	2.03	2.28	2.54	2.81	3.08		
	1334	2078	2400	2486	2527	2744	2930	2980	3130	3207			Split at ends; short specimen, 120 millimeters long..	90 ^a
	1134	1332	1488	1597	1610	1696	1719	1783	1810	1883	2155	2413	Sheared fibers.....	114 ¹
	749	1302	1334	1542	1619	1619	1724	1787	1833	1842	2205	2472	do.....	114 ¹
	1996	2948	3121	3298	3488	3633	3833	3964	4033	4128	4944	5579	Split at corner of specimen; fibers not sheared.....	114 ²
	1542	1882	1982	2068	2205	2304	3368	2468	2518	2576	3039	3130	do.....	114 ²
	1315	1842	2041	2164	2279	2427	2499	2586	2603	2690	3130	3357	Split at corner; slight shearing.....	114 ²
		2109	2395	2622	2717	2939	3075	3162	3334	3488	4219		Slight shearing; split at end.....	130
	1529	2654	2976	3148	3284	3434	3529	3665	3800	3937	4401	4990	Sheared fibers.....	130
	1588	1769	1987	2273	2400	2486	2590	2762	2848	2985	3742	4491	Split at end; sheared fibers.....	212
	1588	1741	2023	2186	2336	2504	2667	2830	2894	3057	3810	4296	Sheared fibers.....	212
	1588	1973	2218	2277	2440	2536	2631	2776	2899	3089	3882	4382	do.....	227 ¹
	1882	2985	3388	3611	3742	3904	4042	4137	4250	4350	5171	5285	do.....	227 ²
	1560	1619	2708	2903	2989	3162	3307	3425	3543	3946			Sheared fibers; split at end.....	227 ²
	1860	2481	2744	2890	3103	3248	3388	3525	3583	3706	3856		Split at end.....	267 ¹
	2449	2948	3221	3379	3574		3842	3946	4092	4132	4899	5443	Slight shearing of fibers.....	267 ²
	1551	2159	2250	2463	2579	2708	2867	2978	3134	3221	3924	4300	do.....	431
	1179	1642	1801	1932	2028	2132	2195	2241	2400	2341	2926	3130	Slight shearing of fibers; split at end.....	551
	1352	1579	1746	1887	1991	2123	2218	2336	2438	2522	3062	3429	Slight shearing of fibers.....	551
	1538	2376	2976	3157	3234	3334	3425		3529	3579	3901	3901	Sheared fibers.....	747
	1433	1801	1982	2014	2214	2313	2331	2436	2461	2531	2976	3089	Slight shearing of fibers.....	747
	1905	2468	2694	2948	3152	3334	3620	3788	3901	3987	4900	5489	Slight shearing.....	937
	2023	2622	2885	3198	3470	3751		4196	4350	4604	5489	6010	Slight shearing of fibers.....	1045
	2041	3357	3878	4082	4160	4305	4445	4522	4196	4300	5434	5851	Indented without shearing.....	1045
	1325	1996	2291	2608	2685	2866	3116	3207	3420	3529	4559	5353	do.....	364
	1837	2944	3384	3615	3669	4033	4287	4418	4473	4491	4763		Slight shearing; split stick.....	364
	1533	2518	2799	3021	3162	3343	3484	3597	3679	3751	4377	4699	Sheared fibers.....	229 ^a
	1384	2427	2753	2948	3221	3311	3425	3484	3529	3674	4196	4423	do.....	229 ^a
	1660	2867	3148	3334	3434	3620	3774	3837	3983	4073	4626	5035	Indented without shearing.....	1059
	1579	2767	3243	3357	3593	3710	3878	3933	4037	3810	4717	5035	do.....	57
	1633	1800	1951	2109	2223	2155	2064	2563	2713	2580	3583	3992	Slight shearing of fibers.....	308
	1588	2245	2409	2486	2567	2676	2749	2830	2890	2935	3348	3463	Sheared fibers.....	308
	2041	2708	2894	3175	3379	3647	3828	3910	4055	4205	5035	5625	Slight shearing of fibers.....	438
	2041	3198	3638	3969	4200	4482	4672	4844	5121	5330	6237	6940	do.....	948
	1860	3402	3992	4281	4454	4717	4808	5035	5207	5289	6078	6595	Fibers did not shear.....	948
	2109	3026	3293	3470	3720	3856	3987		4196	4332	4935	5262	Sheared fibers.....	957
	2177	3202	3611	3842	4028	4164	4309	4454	4572	4672	5421	5987	do.....	957
	1179	1760	1932	2096	2209	2318	2377	2459	2522	2549	2912	3157	do.....	536
	2680	3901	4264	4418	4590	4722	4849	5022	5158	5316	6373	7031	Slight shearing of fibers.....	66
	2359	3620	4191	4513	4886	5180	5489	5706	5965	6128	7485	8029	do.....	66
	1452	1633	1656	1705	1724	1842	1896	1973	2029	2105	2427	2586	Sheared fibers.....	125
	1179	1506	1574	1619	1674	1778	1873	1982	1982	2037	2381	2567	do.....	125
	1542		3701	3974	4150	4332				4967	5942	6509	Slight shearing of fibers.....	286 ¹
	1225	2718	3334	3720	3882	4082	4173	4296	4427	4545	5149		Sheared fibers; split at end.....	286 ²
	1973	2740	2894	2989	3071	3202	3357	3420	3479	3615	4128	4445	Indented section covers unsound knot 10 millimeters in diameter.	291
	1656	3171	3819	4164	4377	4527	4744	4835	5108	5216	6123	6849	Slight shearing of fibers.....	518

TABLE V.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector	Soil.	Longitudinal compression: Ultimate strength, in kilos.	Remarks.
197. <i>Fraxinus Oregona</i> <i>Oregon Ash.</i>	964	Oregon	Portland	G. Engelmann and C. S. Sargent.	Low, wet	9753	Crushed at middle
	964	do	do	do	do	7938	Crushed 25 millimeters from middle on one face.
	1001	do	Weldler's saw-mill	do		8799	Triple flexure, deflected diagonally.
	1001	do	do	do		9063	Crushed at end
	1024	do	Portland Furniture Company.	do		6441	Crushed at middle; side of specimen split off.
	1024	do	do	do		5557	Crushed at middle; angle of crushing, 70°.
	1030	do	do	do		9268	Crushed at middle; angle of crushing, 53°.
	1030	do	do	do		9798	Crushed at 51 and at 114 millimeters from end.
198. <i>Fraxinus sambucifolia</i> <i>Black Ash. Hoop Ash. Ground Ash.</i>	122	Michigan	Dansville	W. J. Beal	Wet, peaty	6237	Crushed on one face 25 millimeters from middle.
	839	Massachusetts	Danvers	J. Robinson	Rich, loamy	7021	Triple flexure; middle bend 25 millimeters eccentric.
	839	do	do	do	do	6441	do
199. <i>Forestiera acuminata</i> <i>Privet.</i>	737	Georgia	Bainbridge	A. H. Curtiss		6118	Split from end to end
	737	do	do	do			
201. <i>Osmanthus Americanus</i> <i>Devil Wood.</i>	283	Louisiana	Amite	C. Mohr	Rich, alluvial	9058	Crushed along one face
	283	do	do	do	do	9063	Crushed at knot 64 millimeters from end.
	584	Florida	Saint John's river	A. H. Curtiss	Sandy loam	8777	Crushed at 76 millimeters from end.
	584	do	do	do	do	8119	Crushed at 102 millimeters from end.
BORRAGINACEÆ.							
204. <i>Borreria Havanaensis</i> <i>Strong Bark.</i>	1137	do	Key Largo	do	Coral	10614	Crushed fibers at 38 millimeters from end.
	1137	do	do	do	do	7779	Cross-grained; oblique fracture along grain.
205. <i>Ehretia elliptica</i> <i>Knackaway. Anagua.</i>	942	Texas	New Braunfels	C. Mohr	Rich, alluvial	6078	Crushed at one corner near middle.
	942	do	do	do	do	6305	Triple flexure, deflected diagonally.
BIGNONIACEÆ.							
206. <i>Catalpa bignonioides</i> <i>Catalpa. Catawba. Bean Tree. Cigar Tree. Indian Bean.</i>	540	Alabama	Stockton	do	Low, wet	6827	Crushed at end
	744	Georgia	Bainbridge	A. H. Curtiss	Clay	5625	Crushed at 38 millimeters from middle on one face.
	744	do	do	do	do	5012	Crushed at 38 millimeters from end.
207. <i>Catalpa speciosa</i> <i>Western Catalpa.</i>	38	Missouri	Charleston	C. S. Sargent	Wet clay	6668	Crushed at 19 and at 102 millimeters from end.
	38	do	do	do	do	6373	Triple flexure
208. <i>Chilopsis saligna</i> <i>Desert Willow.</i>	682	Arizona	Tucson	G. Engelmann and C. S. Sargent.	Moist, gravelly	4755	Failed in vicinity of 10 millimeters knot.
VERBENACEÆ.							
210. <i>Citharexylum villosum</i> <i>Fiddle Wood.</i>	490	Florida	Bay Biscayne	A. H. Curtiss	Coral	11067	Crushed near middle; angle of crushing, 55°.
	490	do	do	do	do	11090	Crushed near middle
NYCTAGINACEÆ.							
212. <i>Pisonia obtusata</i> <i>Pigeon Wood. Beef Wood. Cork Wood. Perk Wood.</i>	474	do	Upper Metacombe Key.	do	do	5398	Split along grain from middle to end.
	474	do	do	do	do	4527	Crushed at knots 102 millimeters from end.
POLYGONACEÆ.							
213. <i>Coccoloba floridana</i> <i>Pigeon Plum.</i>	473	do	do	do	do	12292	Crushed at 38 millimeters from middle and at end.
	473	do	do	do	do	12382	Crushed at middle
214. <i>Coccoloba uvifera</i> <i>Sea Grape.</i>	453	do	do	do	do	4355	Split from end to end; badly cracked in seasoning.
	453	do	do	do	do	3900	do
LAURACEÆ.							
215. <i>Persea Carolinensis</i> <i>Red Bay.</i>	585	do	Saint John's river	do	Sandy loam	9548	Crushed at 5 millimeters knot near middle.
	585	do	do	do	do	8799	Crushed at end
215. <i>Persea Carolinensis</i> , var. <i>palustris</i> .	340	Alabama	Mobile county	C. Mohr	Damp, sandy	5216	Crushed at 64 millimeters from end.
	340	do	do	do	do	6532	do
217. <i>Sassafras officinale</i> <i>Sassafras.</i>	71	Missouri	Allenton	G. W. Letterman	Low, rich	6069	Crushed at 51 millimeters from middle and at ends.
	71	do	do	do	do	6496	Crushed at 3 millimeters knot 38 millimeters from middle.

UNITED STATES UNDER COMPRESSION—Continued.

Direction of grain.	PRESSURE, IN KILOGRAMS, REQUIRED TO PRODUCE AN INDENTATION, IN MILLIMETERS, OF—												Remarks.	Specimen number.
	0.25	0.51	0.76	1.02	1.27	1.52	1.78	2.03	2.28	2.54	2.81	3.08		
	1452	2200	2504	2685	2849	2971	3075	3175	3248	3379	3478		Slight shearing of fibers; split at end.....	904
	1669	2359	2576	2703	2835	2935	3089	3180	3266	3357	3478		do.....	904
	1633	2436	2595	2626	2803	2980	3021	3121	3221	3298	3429		do.....	1001
	1814	2223	2313	2440	2632	2749	2894	3026	3130	3243	3356	4128	do.....	1001
	1457	1842	1960	2118	2214	2313	2431	2531	2622	2654	3148	3470	Sheared fibers.....	1024
	1315		2064	2096	2141	2245	2345	2381	2486	2536	2903	3107	do.....	1024
	1579	2223	2472	2685	2880	3094	3289	3637	3468	3525	4128	4355	Sheared fibers; split at end.....	1030
	1370	2277	2286	2654	2880	3016	3212	3302	3438	3561	4150	4708	Sheared fibers.....	1030
	1189	1674	1746	1819	1842	1887	1937	1982	2003	2082	2295	2427	Sheared fibers; split at end.....	123
	2273	3348	3601	3983	4119	4264	4336	4527	4613	4713	5376	5489	Slight shearing of fibers.....	830
	2132	2699	2819	3084	3337	3583	3819	3992	4153	4318	5285	5625	do.....	830
	871	1837	2205	2331	2368	2481	2576	2635	2717	2753	2912	3583	Sheared fibers.....	737
	1397	2381	2749	2930	3066	3184	3243	3375	3434	3484	4078	4287	do.....	737
	2744	4355	4990	5398	5648	5920	6123	6373	6518	6659	7756	8523	do.....	298
	2449	2640	2903	3212	3465	3742	3924	4150	4287	4432	5489		Sheared fibers; split at end; short specimen, 103 millimeters long.	298
	2223	2867	3021	3334	3506	3656	3788	4014	4110	4246	5058	5625	Sheared fibers.....	584
	2041	2486	2703	3062	3212	3425	3570	3710	3842	3969	4763	5262	do.....	584
	2087	3756	4482	4877	4967	5216	5371	5489	5620	5702	6577	7076	do.....	1187
	2132	3334	3837	4146	4436	4681	4872	4990	5171	5343	6237	6718	do.....	1187
	1610	2790	3339	3611	3878	4037	4155	4336	4477	4572	5239	5851	do.....	942
	1542	2608	3066	3293	3447	3651	3774	3901	4037	4150	4926	5443	do.....	942
	930	1216	1334	1429	1488	1524	1579	1606	1642	1728	2028		Sheared fibers; split at end; specimen 120 millimeters long.	540
	735	848	871	925	944	980	1025	1066	1080	1089	1302	1433	Sheared fibers.....	744
	717	1061	1152	1207	1257	1302	1347	1393	1402	1438	1701	1878	do.....	744
	998	1234	1352	1429	1488	1551	1619	1660	1674	1719	1960		Sheared fibers; split at end.....	38
	703	1098	1170	1229	1266	1320	1356	1402	1442	1488	1724	1941	Sheared fibers.....	38
	1270	1851	2105	2205	2304	2468	2558	2658	2731	2803	3375	3593	do.....	693
	2676	3652	3991	4264	4491	4704	4895	5080	5162	5216			Slight shearing of fibers; split at end.....	490
	2048	4241	4808	5153	5362	5561	5632	5942					do.....	490
	749	1302	1515	1600	1746	1778	1787	1801	1796	1805	1978	2064	Sheared fibers; split at end.....	474
	880	1325	1533	1609	1728	1765	1851	1878	1928	2005	2177		do.....	474
	4128	5398	6260	7031	7303	7485	7621	8074	8392	8663	10115	10796	Sheared fibers.....	473
	2404	3810	4409	4980	5316	5625	5860	6123	6532	6759			Slight shearing; split at end; short specimen, 120 millimeters long.	473
	2019	3039	3230	3339	3429	3465	3534	3624	3724	3788	4309	4522	Sheared fibers.....	586
	1792	2608	2763	2803	2926	2994	3075	3125	3212	3253	3856	4196	do.....	586
	1951	2880	3157	3339	3515			3878	3937	3987	4491	4854	do.....	340
	1551	2205	2481	2558	2631	2717	2794	2858	2980	3066	3538	3878	do.....	340
	1810	1805	2019	2037	2127	2155	2168	2218	2295	2318	2694	2835	do.....	71
	1111	1792	2014	2068	2118	2164	2214	2322	2381	2406	2767	2808	do.....	71

TABLE V.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Longitudinal compression: Ultimate strength, in kilos.	Remarks.
217. <i>Sassafras officinale</i> —continued.	387	Missouri	Allenton	G. W. Letterman	Alluvial	5751	Crushed at 51 millimeters from end.
	387	do	do	do	do	6328	Crushed at middle and at 64 millimeters from end.
	446	Tennessee	Nashville	A. Gattinger	Rich	6713	Crushed at 32 and at 89 millimeters from end.
	814	West Virginia	Grafton	C. G. Pringle		6418	Triple flexure, deflected diagonally.
	814	do	do	do		6659	do
	854	Massachusetts	Danvers	J. Robinson	Rich loam	4559	Triple flexure
	854	do	do	do	do	5851	do
	1163	Missouri	Allenton	G. W. Letterman	Low, alluvial	6341	Triple flexure; developed intersecting "Cooper lines".
	1163	do	do	do	do	6038	Triple flexure
218. <i>Umbellularia Californica</i> . <i>Mountain Laurel. California Laurel. Spice Tree. Oagiput. California Olive. California Bay Tree.</i>	708	Oregon	Coos bay	G. Engelmann and C. S. Sargent.		9485	Crushed at 25 millimeters from middle.
	708	do	do	do		8754	Crushed at 38 millimeters from end.
EUPHORBIACEÆ.							
219. <i>Drypetes crocea</i> . <i>Guiana Plum. White Wood.</i>	468	Florida	Upper Metacombe Key.	A. H. Curtiss	Coral	10410	Crushed in vicinity of small knots
219. <i>Drypetes crocea</i> , var. <i>latifolia</i> .	459	do	do	do	do	8256	Split suddenly from end to end
	459	do	do	do	do	8392	Split suddenly; oblique fracture.
URTICACEÆ.							
222. <i>Ulmus crassifolia</i> . <i>Cedar Elm.</i>	324	Texas	Dallas	J. Reverchon	Rich loam	7847	Crushed at 51 millimeters from end.
	324	do	do	do	do	8414	Crushed in vicinity of small knots 103 millimeters from end.
	929	do	Austin	C. Mohr	do	5951	Crushed at end; cross-grained.
	929	do	do	do	do	6781	Triple flexure
223. <i>Ulmus fulva</i> . <i>Red Elm. Slippery Elm. Moose Elm.</i>	303	Kentucky	Mercer county	W. M. Linney	Limestone	7847	Deflected and crushed at end and at 102 millimeters from end.
	304	do	do	do	do	8573	Crushed at middle
	120	Michigan	Dansville	W. J. Beal	Gravelly	9889	Crushed at 102 millimeters from end.
	134	Missouri	Allenton	G. W. Letterman	Rich, alluvial	8437	Triple flexure
	134	do	do	do	do	8392	Crushed near middle in vicinity of knot.
224. <i>Ulmus Americana</i> . <i>White Elm. American Elm. Water Elm.</i>	19	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	8641	Deflected; crushed at middle and end.
	19	do	do	do	do	8573	Triple flexure
	281	Missouri	Allenton	G. W. Letterman	Alluvial	7698	Triple flexure; middle bend eccentric.
	281	do	do	do	do	6895	Crushed at middle; deflected diagonally.
	958	Texas	Colorado river	C. Mohr	do	5851	Crushed at knot 102 millimeters from end.
	958	do	do	do	do	4990	Crushed at 10 millimeters knot near middle.
	1036	Massachusetts	Danvers	J. Robinson	Gravelly	7022	Triple flexure, deflected diagonally.
	1036	do	do	do	do	5579	do
	1049	do	North Reading	do	do	9049	Triple flexure; developed intersecting "Cooper lines".
225. <i>Ulmus racemosa</i> . <i>Rock Elm. Cork Elm. Hickory Elm. White Elm. Old Elm.</i>	116	Michigan	Dansville	W. J. Beal	do	11385	Triple flexure; middle deflection 25 millimeters eccentric.
	116	do	Big Rapids	do	do	9571	do
	116	do	do	do	Low, gravelly	7847	Crushed fibers at 82 millimeters from end.
	116	do	Hudson	do	Alluvial	9571	Triple flexure
	814	do	Hersey	do	Rich loam	10387	Crushed at 25 and at 114 millimeters from end.
	814	do	do	do	do	10206	Triple flexure, deflected diagonally.
	428	Tennessee	Nashville	A. Gattinger	do	7349	Crushed at small knot at middle.
226. <i>Ulmus alata</i> . <i>Wahoo. Winged Elm.</i>	138	South Carolina	Bonneau's Depot	H. W. Ravenel	do	6895	Crushed at 13 millimeters knot at middle.
	138	do	do	do	do	7847	Triple flexure, deflected diagonally.
	880	Tennessee	Davidson county	A. Gattinger	Loam	6280	Crushed at 10 millimeters knot 64 millimeters from end.
	533	Mississippi	Kemper's mill	C. Mohr	Alluvial	7008	Triple flexure
	533	do	do	do	do	7892	Triple flexure, deflected diagonally.

UNITED STATES UNDER COMPRESSION—Continued.

Direction of grain.	PRESSURE, IN KILOGRAMS, REQUIRED TO PRODUCE AN INDENTATION, IN MILLIMETERS, OF—												Remarks.	Office number.
	0.25	0.51	0.76	1.02	1.27	1.52	1.78	2.03	2.28	2.54	4.81	5.08		
	1111	1488	1615	1665	1710	1760	1846	1873	1937	1982	2250	2440	Sheared fibers	287
	1071	1442	1588	1642	1689	1710	1765	1801	1801	1887	2155		Sheared fibers; split at end	287
	1452	1892	1914	1978	2055	2123	2155	2209	2259	2304	2608	2758	Sheared fibers	446
	1111	1882	2109	2195	2254	2350	2413	2472	2490	2513	2884	3121	do	814
	1542	1778	1982	2164	2250	2377	2531	2672	2776	2862	3561	3946	do	814
	1270	2068	2381	2468	2522	2558	2621	2658	2699	2744	3139	3334	do	854
	1778	1960	2014	2078	2132	2177	2250	2336	2427	2473	2858	3062	do	854
	1275	1833	2068	2164	2259	2413	2468	2531	2595	2635	3198	3538	Slight shearing of fibers	1163
	1665	2250	2368	2440	2490	2531	2576	2622	2667	2690	2908	3026	Sheared fibers	1168
	2019	2699	2971	3066	3193	3329		3561	3674	3756	4300	4672	do	703
	1978	2744	2926	3057	3198	3334	3357	3561	3628	3674	4037	4445	do	703
	2177	3856	4854	5466	5797	6092	6373	6577	6804	6949	8340	8845	do	468
	3652	5670	6328	6768	7145	7439	7756	7983	8155	8237	9480	10160	do	459
	3130	4509	5108	5615	5874	6110	6396	6599	6749	6918	8256	8799	Sheared fibers; split at end	459
	1315	1878	2259	2477	2640	2803	3030	3166	3334	3484	4300	4808	Sheared fibers	824
	2676	3393	3620	3637	4046	4237	4359	4531	4690	4899	5851	6192	do	824
	2676	3856	4296	4635	4890	4999	5353	5516	5797	5965	7212	7993	Slight shearing of fibers	929
	2109	3765	4196	4482	4744	4953	5163	5380	5611	5797	7031	8029	Sheared fibers	929
	1678	1814	1910	2032	2223	2322	2427	2522	2667	2744	2948		Slight shearing of fibers; split at end	30 ^a
	1093	1479	1619	1765	1855	1989	2064	2164	2277	2359	2858	3153	do	30 ^a
	1565	1932	2223	2440	2576	2667	2758	2866	2935	2994			do	120
	1379	2277	2386	2527	2685	2799	2976	3112	3207	3266	3937		do	134
	1551	2073	2300	2481	2658	2958		3069	3207	3334	4014		do	134
	1592	2350	2699	2875	2971	3121	3212	3348	3438	3529	4250	4513	Sheared fibers	19
	1338	1905	2168	2377	2504	2645	2753	2844	2971	3057	3674	3992	do	19
	1610	1960	2041	2186	2336	2372	2504	2567	2663	2744			Split at ends; fibers not sheared; specimen 120 millimeters long.	281
	1293	1769	1914	2064	2196	2313	2431	2549	2698	2726	3221		do	281
	1610	1669	1715	1787	1882	1982	2118	2214	2331	2354	2971	3357	Sheared fibers	958
	1324	1669	1896	2014	2109	2214	2304	2409	2459	2522	2971	3266	do	958
	1656	2449	2744	2958	3139	3261	3452	3556	3697	3774	4740	5171	Slight shearing of fibers	1036
	1796	2835	3266	3515	3697	3810	3964	4046	4199	4281	4907	5353	do	1036
	2096	2890	3198	3434	3697	3828	4028	4182	4281	4468	5489	5874	do	1049
	2291	3289	3674	4037	4241	4527	4695	4896	5035	5198			Split at end; fibers did not shear	116
	1338	1887	2041	2164	2268	2381	2477	2576	2658	2708	3266		Split at end; slight shearing of fibers	116 ^a
	1179	1987	2254	2468	2613	2794	2898	3085	3125	3243			Slight shearing of fibers; split at end	116 ^a
	1111	2295	2740	2971	3184	3402	3583	3738	3882	4019	4808		do	116 ^a
	2291	2853	3075	3206	3452	3638	3765	3919	4033	4196	4808		do	314
	1906	2568	3016	3198	3366	3556	3665	3747	3946	4119	4854		do	314
	1678	2875	3343	3620	3842	4073	4200	4400	4581	4672	5625	6123	Slight shearing of fibers	428
	2586	3901	4377	4877	5013	5243	5557	5697	5942	6069	7393	7689	do	183
	1474	2912	3606	4092	4404	4644	4881	5116	5389	5579	6895	7621	do	183
	1678	2495	2617	2708	2869	2971	3030	3202	3293	3348	3860	4196	Sheared fibers	380
	2381	3030	3257	3465	3620	3810	3992	4150	4287	4400	5353	5806	Slight shearing of fibers	532
	2336	3498	3965	4264	4522	4758	4937	5194	5358	5587	6441	7285	do	532

TABLE V.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Longitudinal compression: Ultimate strength, in kilos.	Remarks.
227. <i>Planera aquatica</i>	758	Florida	Chattahoochee...	A. H. Curtiss	Rich, alluvial	6373	Triple flexure.....
	758	do	do	do	do	6528	Triple flexure, deflected diagonally.
	918	do	do	C. Mohr	do	6214	Crushed at 38 and at 102 millimeters from end.
228. <i>Celtis occidentalis</i>	75	Missouri	Allenton	G. W. Letterman	Low, rich	5679	Crushed on one face at 25 millimeters from middle.
<i>Sugarberry. Hackberry.</i>	75	do	do	do	Alluvial	6895	Crushed at knot 51 millimeters from end.
	306	Texas	Dallas	J. Reverchon	do	5579	Crushed at middle, deflected diagonally.
	306	do	do	do	do	6169	Triple flexure, deflected diagonally.
	375	Tennessee	Davidson county	A. Gattinger	Loam	6895	Triple flexure
	375	do	do	do	do	7031	Triple flexure, deflected diagonally.
	873	Massachusetts	Salem	J. Robinson	do	5489	Triple flexure
	873	do	do	do	do	7303	Triple flexure, deflected diagonally.
	1111	Missouri	Saint Louis	Henry Eggert	Moist loam	8278	Deflected; crushed at middle
	1111	do	do	do	do	8074	Crushed at 6 millimeters knot 102 millimeters from end.
228. <i>Celtis occidentalis</i> , var. <i>reticulata</i> .	652	Arizona	Santa Rita mountains.	G. Engelmann and C. S. Sargent.	Dry	6985	Crushed at 12 millimeters knots at end and at 102 millimeters from end.
<i>Hackberry. Palo Blanco.</i>							
229. <i>Ficus aurea</i>	486	Florida	Bay Biscayne	A. H. Curtiss	Coral	3198	Triple flexure
	486	do	do	do	do	1996	Crushed at middle and at 38 millimeters from end.
231. <i>Ficus pedunculata</i>	508	do	Boca Chica Key	do	do	4491	Split obliquely.....
<i>Wild Fig. India-rubber Tree.</i>							
232. <i>Morus rubra</i>	132	Missouri	Allenton	G. W. Letterman	Rich loam	8754	Crushed at middle and at 25 millimeters from end.
<i>Red Mulberry.</i>	132	do	do	do	do	8483	Crushed at middle and at end; deflected diagonally.
	433	Tennessee	Nashville	A. Gattinger	do	6827	Failed at 12 millimeters knot 102 millimeters from end.
	1244	Missouri	Allenton	G. W. Letterman	Upland	6056	Crushed at 25 millimeters from middle at 3 millimeters knot.
	1245	do	do	do	do	6169	Crushed at 19 millimeters from end in vicinity of small knots.
	1246	do	do	do	do	5987	Triple flexure.....
	1255	do	do	do	Rich	5829	Triple flexure, deflected diagonally.
	1255	do	do	do	do	5661	Triple flexure
234. <i>Maclura aurantiaca</i>	253	Texas	Dallas	J. Reverchon	Bottom	14107	Crushed at middle; angle of crushing, 50°.
<i>Osage Orange. Bois d'Arc.</i>	253	do	do	do	do	11771	Triple flexure.....
PLATANACEÆ.							
235. <i>Platanus occidentalis</i>	21	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	6759	Crushed at end and at 102 millimeters from end.
<i>Sycamore. Button Wood.</i>	21	do	do	do	do	5579	Crushed at end and at 127 millimeters from end.
<i>Button-ball Tree. Water Beech.</i>	126	Missouri	Allenton	G. W. Letterman	Rich, alluvial	8233	Crushed at 102 millimeters from end.
	126	do	do	do	do	8256	Crushed at 38 and at 127 millimeters from end.
236. <i>Platanus racemosa</i>	686	California	Carmel river	G. R. Vasey	Clay	4491	Crushed at 13 and at 76 millimeters from end and split along grain.
<i>Sycamore. Button Wood.</i>	686	do	do	do	do	5888	Crushed at middle of one face.
237. <i>Platanus Wrightii</i>	648	Arizona	Santa Rita mountains.	G. Engelmann and C. S. Sargent.	Rich, gravelly	5398	Triple flexure, deflected diagonally.
<i>Sycamore.</i>	648	do	do	do	do	5058	Triple flexure
JUGLANDACEÆ.							
238. <i>Juglans cinerea</i>	16	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	6169	Crushed at 51 and at 127 millimeters from end.
<i>Butternut. White Walnut.</i>	16	do	do	do	do	6396	Crushed at 102 millimeters from end.
	76	Missouri	Allenton	G. W. Letterman	Moist, alluvial	7621	Crushed at 64 millimeters from end.
	76	do	do	do	do	7066	Crushed at 51 millimeters from end.
	76	do	do	do	Rich, moist upland	6849	Crushed at 25 millimeters from end.
	123	Michigan	Dansville	W. J. Beal	Gravelly clay	6123	Crushed at 19 millimeters from end.
	145	Illinois	Waukegan	Robert Douglas	Alluvial	5080	Crushed at 13 millimeters from middle and at 38 millimeters from end.
	393	Michigan	Lansing	W. J. Beal	Gravelly loam	4150	Crushed at 114 millimeters from end and at end.
	1057	Massachusetts	Topsfield	J. Robinson	Drift	6373	Crushed at middle

UNITED STATES UNDER COMPRESSION—Continued.

Direction of grain.	PRESSURE, IN KILOGRAMS, REQUIRED TO PRODUCE AN INDENTATION, IN MILLIMETERS, OF—												Remarks.	Office number.
	0.25	0.51	0.76	1.02	1.27	1.52	1.78	2.03	2.28	2.54	2.81	3.08		
	1397	1996	2254	2386	2549	2749	2858	2994	3107	3212	3324	4355	Sheared fibers.....	758
	1406	1941	2141	2350	2481	2622	2744	2830	2971	3094	3810	4273	Slight shearing of fibers.....	758
	1134	1633	1769	1796	1973		2109	2182	2245	2313	2699	2903	Sheared fibers.....	918
	1415	1860	1987	2123	2232	2341	2459	2563	2640	2722			Slight shearing; split at end.....	75
	1466	2041	2245	2427	2576	2672	2799	2894	3016	4080	3652		Sheared fibers.....	75
	2608	3493	3819	4037	4327	4531	4735	4899	4980	5180	6337	6940	Slight shearing of fibers.....	306
	1111	2404	3003	3216	3488	3683	3869	4053	4209	4341	5421	6033	do.....	306
	1724	2699	3107	3402	3583	3837	3951	4105	4204	4350	5216	5829	Fibers did not shear.....	375
	1746	2639	3583	3865	4073	4332	4450	4617	4817	4944	5851	6396	do.....	375
	1610	2703	3016	3175	3393	3562	3715	3924	4064	4110	4877	5307	do.....	373
	1751	2749	3026	3320	3505	3742	3974	4119	4314	4445	5126	5625	Slight shearing of fibers.....	373
	1606	2812	3243	2976	3574	3629	3742	3837	3987	3940	4649		Slight shearing; split at end.....	1111
	1579	2903	3339	3620	3910	4009	4309	4527	4626	4758	5670	6192	Sheared fibers.....	1111
	2223	3493	3946	4200	4373	4536	4699	4994	5071	5262	6078	6350	do.....	651
	522	817	862	880	885	885	894	894	898	903	903	1039	Sheared fibers; specimen 120 millimeters long.....	486
	640	907	980	1030	1075	1116	1161	1175	1211	1229	1429	1479	Sheared fibers.....	486
	975	1080	1243	1329	1905	2019	2064	2123	2191	2313	2694	2840	do.....	508
	1751	2440	2609	2840	2926	3062	3162	3266	3320	3393	3833	4105	do.....	132
	1769	2376	2753	2880	2969	3116	3266	3334	3479	3543	4087	4518	do.....	132
	1524	1973	2078	2166	2218	2286	2409	2481	2608	2681	3157	3620	Sheared fibers; split at end.....	1244
	1960	2640	2790	2926	3016	3184	3248	3379	3534	3674	4309	4786	Sheared fibers.....	1245
	1905	2313	2531	2676	2875	3066	3252	3515	3652	3715	4445	4944	do.....	1246
	1960	2640	2790	2926	3016	3184	3248	3379	3534	3674	4309	4786	do.....	1255
	1905	2313	2531	2676	2875	3066	3252	3515	3652	3715	4445	4944	do.....	1256
	2903	5080	5670	5720	5761	6010	6019	6192	6328	6423	7167	7608	do.....	253
	2586	4740	5421	5606	5851	6069	6214	6396	6432	6577	6985	6895	Sheared fibers; split along grain from end to end.....	253
	1678	2105	2331	2486	2622	2703	2880	2985	3094	3166	3107	4423	Sheared fibers.....	21
	1492	1982	2177	2341	2440	2581	2527	2685	2985	3066	3583	4082	Slight shearing of fibers.....	21
	1202	2404	2858	3039	3085	3134	3289	3348	3429	3488	3788	3946	Sheared fibers.....	126
	1505	2136	2277	2377	2440	2549	2635	2708	2762	2826	3259		Sheared fibers; split at end.....	126
	1039	1365	1501	1579	1665	1740	1814	1846	1901	1978	2313	2486	Sheared fibers.....	686
	753	1071	1157	1257	1306	1393	1479	1533	1578	1628	2019	2200	do.....	686
	1134	1676	1855	1987	2114	2191	2263	2359	2404	2495	2894	3044	do.....	648
	934	1338	1428	1520	1619	1683	1765	1842	1887	1932	2313	2540	do.....	648
	998	1542	1588	1669	1742	1801	1833	1905	1987	2037	2381	2590	do.....	16
	1089	1452	1506	1542	1610	1678	1719	1769	1810	1860	2127	2291	do.....	16
	793	1098	1134	1202	1257	1306	1343	1356	1393	1461	1609	1792	do.....	76
	1025	1257	1361	1474	1615	1656	1710	1751	1783	1846	2087		Sheared fibers; split at end.....	76
	839	1293	1361	1433	1479	1592	1637	1710	1769	1833	2168	2381	Sheared fibers.....	76 ^a
	703	998	1102	1134	1170	1211	1243	1266	1297	1302	1479	1533	do.....	123
	749	1179	1315	1406	1460	1492	1588	1628	1669	1715	1996		Sheared fibers; split at end.....	145
	408	508	599	617	635	667	690	708	717	739	866	934	Sheared fibers.....	398
	1021	1701	1860	1946	2023	2078	2127	2173	2223	2268	2481	2626	do.....	1067

FOREST TREES OF NORTH AMERICA.

TABLE V.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Longitudinal compression: Ultimate strength, in kiloe.	Remarks.
239. <i>Juglans nigra</i> <i>Black Walnut.</i>	112	Missouri	Allenton	G. W. Letterman	Alluvial	9026	Crushed at middle
	117	Michigan	Danaville	W. J. Beal	Gravelly	9957	Crushed at 102 millimeters from end.
	318	do	Lansing	do	Loam	9095	Split obliquely from end to end
	325	Texas	Dallas	J. Reverchon	Alluvial	8346	Crushed at 19 millimeters from middle.
	407		Charlestown Navy-yard.	S. H. Pook		9934	Crushed at 25 millimeters from middle.
	766	Florida	Aspalaga	A. H. Curtiss	Clay	9979	Crushed at small knot 32 millimeters from middle.
	766	do	do	do	do	9684	Crushed at middle
	951	Texas	New Braunfels	C. Mohr	Moist, calcareous	8301	Crushed at 51 millimeters from end.
	951	do	do	do	do	9662	Crushed at 32 millimeters from end.
240. <i>Juglans rupestris</i> <i>Walnut.</i>	415	New Mexico	Pinos Altos mountains.	E. L. Greene	Alluvial	5942	Shattered stick at end
	415	do	do	do	do	6532	Split along grain in oblique direction.
	672	California	Contra Costa county.	G. R. Vasey	do	8256	Crushed at 51 millimeters from end.
	672	do	do	do	do	7258	Crushed at 6 millimeters knot at middle.
241. <i>Carya olivaeformis</i> <i>Pecan. Illinois Nut.</i>	322	Mississippi	Greenville	C. Mohr	do	6940	Split at ends
	322	do	do	do	do	8256	Triple flexure; developed intersecting "Cooper lines".
	326	Texas	Dallas	J. Reverchon	do	6033	Triple flexure
	326	do	do	do	do	6577	Triple flexure, deflected diagonally.
242. <i>Carya alba</i> <i>Shell-bark Hickory. Shag-bark Hickory.</i>	8	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	9095	Triple flexure; 60 per cent. sap-wood; deflected from sap side.
	8	do	do	do	do	8301	Split at end
	29 ^a	Kentucky	Danville	W. M. Linney	Shale	8808	Failed at 10 millimeters knot near middle; sap-wood.
	118 ^a	Michigan	Hudson	W. J. Beal	Clay	11431	Crushed at end
	118 ^a	do	Lansing	do	do	11567	Crushed at 114 millimeters from end.
	152	Missouri	Allenton	G. W. Letterman	Rich upland	10160	Crushed at 38 millimeters from middle.
	249	Virginia	Wytheville	H. Shriver	Clay	10478	Triple flexure
	249	do	do	do	do	9934	do
	531	Mississippi	Kemper's mill	C. Mohr	Alluvial	10342	Crushed at 38 millimeters from middle.
	531	do	do	do	do	8890	Crushed at knots; cross-grained
	589	do	do	do	do	10660	Crushed at 38 and at 70 millimeters from end.
	589	do	do	do	do	10514	Crushed near middle
	816	West Virginia	Grafton	C. G. Pringle	do	11204	Crushed at 51 millimeters from end.
	816	do	do	do	do	11022	Crushed at 25 millimeters from middle.
	1056	Massachusetts	Topshfield	J. Robinson	Rich loam	9026	Triple flexure, deflected diagonally toward heart; 80 per cent. sap-wood.
	1056	do	do	do	do	8600	Triple flexure, deflected diagonally; 90 per cent. sap-wood
	1097	Missouri	Allenton	G. W. Letterman	Alluvial	10015	Triple flexure; split from end to end; 45 per cent. sap-wood.
243. <i>Carya sulcata</i> <i>Big Shell-bark. Bottom Shell-bark.</i>	91 ^a	Kentucky	Mercer county	W. M. Linney	do		
	383	Missouri	Allenton	G. W. Letterman	do	8301	Triple diagonal flexure; 60 per cent. sap-wood on concave side.
	383	do	do	do	do	7802	Triple flexure; 80 per cent. sap-wood on convex side.
	391	do	do	do	do	9707	Triple flexure; 80 per cent. sap-wood on concave side.
	391	do	do	do	do	9571	Triple diagonal flexure
	1082	do	do	do	do	7983	Triple flexure; 50 per cent. sap-wood.
	1082	do	do	do	do	8369	Triple flexure; 5 per cent. sap-wood.
	1164	do	do	do	do	10592	Crushed at 25 millimeters from end.
	1165	do	do	do	do	8845	Crushed at 25 millimeters from end and split along grain.
	1166	do	do	do	do	8961	Triple flexure; 95 per cent. sap-wood.
	1170	do	do	do	do	9281	Crushed at 6 millimeters from end; sap-wood.
244. <i>Carya tomentosa</i> <i>Mocker Nut. Black Hickory. Bull Nut. Big-bud Hickory. White-heart Hickory. King Nut.</i>	72	Kentucky	Perryville	W. M. Linney	Utica shale	8608	Triple flexure; sap-wood
	72	do	do	do	do	8981	do

UNITED STATES UNDER COMPRESSION—Continued.

Direction of grain.	PRESSURE, IN KILOGRAMS, REQUIRED TO PRODUCE AN INDENTATION, IN MILLIMETERS, OF—												Remarks.	Office number.
	0.35	0.51	0.76	1.02	1.27	1.52	1.78	2.03	2.28	2.54	2.81	3.08		
	1452	2268	2518	2672	2767	2890	2985	3089	3184	3266	3361	3424	Sheared fibers.....	112
	1560	2223	2895	2481	2576	2740	2776	2849	2912	2985	3447	3615	do.....	117
	1746	2676	2894	3071	3212	3302	3497	3579	3710	3869	4400	4740	do.....	318
	1769	2740	2921	3221	3388	3597	3783	3827	3955	4042	4423		Sheared fibers; split at end.....	325
	1838	1851	1909	2073	2185	2250	2350	2427	2504	2545	2980	3180	Sheared fibers.....	407
	1479	1951	2449	2713	2890	3008	3121	3253	3393	3456	4128	4536	do.....	706
	2812	3701	3969	4082	4255	4377	4568	4649	4749	4799	5489	5738	Sheared fibers; indented section includes two 2 millimeters knots.	706
	1905	2903	3212	3438	3638	3797	3978	4118	4264	4432	5216	5897	Sheared fibers.....	951
	1769	2667	2903	3175	3338	3502	3697	3783	3969	4128	4899	5534	Slight shearing of fibers.....	951
	2019	3470	3924	4196	4336	4500	4604	4740	4840	4931	5625	6078	Sheared fibers.....	415
	953	1928	2245	2486	2200	2830	2953	3039	3153	3289	3882		Sheared fibers; split at end.....	415
	1043	1941	2214	2336	2440	2545	2622	2713	2762	2894	3289	3538	do.....	672
	1170	2155	2468	2576	2658	2749	2808	2912	2976	3035	3438	3742	Sheared fibers.....	672
	1951	2967	3289	3456	3652	3765	3905	4028	4119	4228	4990		Slight shearing of fibers; split at end.....	322
	1928	3221	3878	4173	4423	4753	4944	5080	5398	5557	6192		Slight shearing of fibers; split at end; shaky stick.....	322
	794	1882	2404	2744	2971	3103	3202	3366	3447	3538	4105	4400	Indented without shearing fibers.....	326
	1792	3039	3447	3674	3810	4037	4196	4354	4436	4672	5285	5942	Slight shearing of fibers.....	326
	2200	3857	3847	4060	4363	4559	4717	4877	5035	5180	6078	6668	Slight shearing of fibers; 60 per cent. sap-wood.....	3
	2048	4105	4536	5035	5239	5534	5851	6078	6250	6419	7303	7756	do.....	3
	2245	4042	4581	5080	5363	5652	5851	5978	6260	6373	7439	7938	Indented without shearing fibers; sap-wood.....	29*
	2245	2713	3293	3620	3819	3983	4150	4281	4427	4491	5239	5706	Sheared fibers; split at end.....	118
	2205	3243	3615	3837	4014	4150	4346	4482	4554	4695	5058	5376	Sheared fibers.....	118
	1406	2699	3207	3529	3797	3955	4146	4355	4463	4608	5396	6033	Slight shearing of fibers.....	249
	2812	4060	4300	4559	4717	4886	5017	5189	5294	5452	6486	6804	do.....	249
	2019	3130	3783	4173	4495	4786	4944	5167	5394	5534			Fibers did not shear; split along grain.....	531
	1429	2005	3171	3620	3933	4164	4364	4608	4744	4844	5715		Fibers did not shear; split at end.....	531
	2404	3302	3652	3851	3978	4287	4495	4695	4808	4940	5579		Slight shearing of fibers; split at end.....	539
	1656	2576	2926	3103	3293	3502	3652	3756	3978	4105	4854	5396	Slight shearing of fibers.....	539
	1565	2613	2939	3103	3243	3447	3529	3652	3765	3882	4536	4944	Sheared fibers.....	816
	1657	2590	3016	3198	3366	3511	3661	3710	3797	3887	4527	4877	do.....	816
	2703	4014	4513	4917	5149	5421	5575	5761	5987	6114	7417	8142	Indented without shearing fibers; sap-wood.....	1056
	2994	4264	4649	4985	5353	5670	5920	6214	6477	6704	7825	8754	do.....	1056
	2631	3992	4626	5071	5389	5715	5851	6033	6123	6396	7439	7825	Slight shearing of fibers; 40 per cent. sap-wood.....	1087
	1837	3311	3765	3924	4060	4205	4341	4427	4617	4726	5489	5693	Sheared fibers; 90 per cent. sap-wood.....	91*
	3062	4626	5171	5579	5965	6250	6555	6849	7212	7439	8437	9163	Sheared fibers; 85 per cent. sap-wood.....	383
	2132	3837	4346	4563	4967	5243	5516	5670	5756	5987	7031	7576	Slight shearing of fibers.....	391
	3334	4355	4854	5239	5751	5797	6159	6341	6532	6704	7825	8618	do.....	391
	2245	3697	4241	4559	4817	5013	5163	5398	5525	5756	6895	7530	Fibers did not shear.....	1082
	1497	3003	3470	3774	3847	4209	4495	4640	4863	5058	6101	6691	Fibers sheared; 80 per cent. sap-wood.....	1082
	2200	2790	4055	4382	4581	4872	5058	5203	5298	5280	6441	6713	Slight shearing of fibers.....	1164
	1769	2495	2744	2840	3016	3085	3289	3425	3479	3529	4037	4491	Sheared fibers.....	1165
	2381	3610	4332	4626	4831	5060	5271	5421	5579	5806	6940	7576	Slight shearing of fibers; 80 per cent. sap-wood.....	1166
	2381	3493	3819	4073	4255	4509	4690	4877	4985	5116	6033	6782	Slight shearing of fibers; sap-wood.....	1170
	3107	4037	4336	4527	4708	4976	5163	5294	5475	5662	6668	7145	Sheared fibers; sap-wood.....	72
	2118	3221	3701	3910	4128	4291	4482	4626	4790	4908	5761	6509	do.....	72

TABLE V.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Longitudinal compression: Ultimate strength in kilos.	Remarks.
244. <i>Carya tomentosa</i> —continued.	254	Missouri	Allenton	G. W. Letterman	Rich upland	11249	Triple flexure; middle deflection 25 millimeters eccentric; 0.5 sap-wood.
	289	do	do	do	do	9163	Crushed at 13 millimeters knot 102 millimeters from end.
	248	Alabama	Citronelle	C. Mohr	Sandy	9367	Triple diagonal flexure; sap-wood.
245. <i>Carya porcina</i> Fig Nut. Brown Hickory. Black Hickory. Switch-bud Hickory.	6	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	8165	Failed at knot at middle; 80 per cent. sap-wood.
	6	do	do	do	do	9095	Crushed at 32 millimeters from end.
	88	Missouri	Allenton	G. W. Letterman	Rich loam	10934	Triple flexure; sap-wood
	88	do	do	do	do	9880	Triple flexure, diagonal direction; sap-wood.
	121	Michigan	Dansville	W. J. Beal	Gravelly clay	9117	Triple flexure; middle deflection 25 millimeters from center.
	288	Missouri	Allenton	G. W. Letterman	Rich loam	8754	
	442	Tennessee	Nashville	A. Gattinger	Upland	8845	Triple flexure
	538	Alabama	Kemper's mill	C. Mohr	Alluvial	10424	do
	1051	Massachusetts	North Reading	J. Robinson	Drift	9526	Triple flexure; middle deflection 25 millimeters from center; 0.2 sap-wood.
	1168	Missouri	Allenton	G. W. Letterman	Flinty	8392	Triple flexure; sap-wood
	1168	do	do	do	do	8392	do
246. <i>Carya amara</i> Bitter Nut. Swamp Hickory.	153	do	do	do	Rich upland	8210	Triple flexure, deflected diagonally.
	153	do	do	do	do	9117	Crushed at 3 millimeters knot 25 millimeters from middle.
	838	Massachusetts	Danvers	J. Robinson	Rich loam	7983	Triple flexure
	838	do	do	do	do	8119	do
247. <i>Carya myristiceiformis</i> Nutmeg Hickory.	237	South Carolina	Bonneau's Depot	H. W. Ravenel	Rich, swampy	9253	Crushed at knots near middle
	237	do	do	do	do	11158	Crushed at 51 millimeters from middle; cross-grained.
248. <i>Carya aquatica</i> Water Hickory. Swamp Hickory. Bitter Pecan.	129	do	do	do	Swampy	8437	Triple flexure, diagonal deflection: 80 per cent. sap-wood on convex side.
	862	Mississippi	Vicksburg	C. Mohr		7280	Triple flexure, deflected diagonally; middle bend 25 millimeters from center; sap-wood.
	862	do	do	do		7439	Triple flexure; sap-wood
	740	Georgia	Bainbridge	A. H. Curtiss	Alluvial	9208	Fibers crushed at 102 millimeters from end.
	740	do	do	do	do	7892	Fibers crushed at 6 millimeters knots 102 millimeters from end.
	917	Florida	Chattahoochee	C. Mohr	do	6509	Triple flexure, diagonal deflection.
	917	do	do	do	do	7666	Fibers crushed at 64 millimeters from end.
MYRICACEÆ.							
249. <i>Myrica cerifera</i> Bayberry. Wax Myrtle.	586	do	Saint John's river	A. H. Curtiss	Sandy loam	7122	Triple flexure
250. <i>Myrica California</i>	665	California	Santa Cruz	G. Engelmann and C. S. Sargent.	Rich loam	7983	Crushed fibers at 51 millimeters from middle.
	665	do	do	do	do	9049	Crushed fibers at middle.
CUPULIFERÆ.							
251. <i>Quercus alba</i> White Oak.	8	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	5579	Deflected and split along grain
	8	do	do	do	do	5353	Triple flexure; split along grain.
	32	Kentucky	Mercer county	W. M. Linney	Limestone	9163	Triple flexure
	32 ^a	do	Boyle county	do	Shale	8074	do
	32 ^b	do	do	do	Slate	7779	Crushed fibers at end.
	49	Missouri	Allenton	G. W. Letterman	Rich upland	9163	Triple flexure 19 millimeters from end.
	49	do	do	do	do	7702	Triple flexure
	49 ¹	do	do	do	do	8437	Triple flexure, deflected diagonally.
	49 ²	do	do	do	do	7825	Triple flexure
	49 ³	do	do	do	do	8301	Crushed fibers at end.
	118	Michigan	Big Rapids	W. J. Beal	Gravelly	6781	Triple flexure
	112 ^a	do	Dansville	do	Sandy	8437	do
	112 ^b	do	do	do	do	7392	Triple flexure, deflected diagonally.

UNITED STATES UNDER COMPRESSION—Continued.

Direction of grain.	PRESSURE, IN KILOGRAMS, REQUIRED TO PRODUCE AN INDENTATION, IN MILLIMETRES, OF—												Remarks.	Office number.
	0.25	0.51	0.76	1.02	1.27	1.52	1.78	2.03	2.28	2.54	4.81	5.08		
	2404	3788	4246	4401	4662	4831	5013	5149	5280	5412	6305	6623	Sheared fibers; sap-wood.....	254
	2268	3243	3547	3756	3983	4200	4332	4486	4604	4708			Sheared fibers; split at end; specimen 120 millimeters long.	289
	2350	3742	4150	4445	4662	4831	5013	5216	5421	5588	6509	7167	Sheared fibers; sap-wood.....	348
	2540	4119	4559	4854	5171	5307	5512	5670	5851	5933	6782	7167	Slight shearing of fibers; 40 per cent. sap-wood.....	6
	1903	3198	3611	3810	4073	4300	4527	4672	4854	4980		6159	do.....	6
	2948	4400	4990	5276	5661	5924	6105	6386	6577	6759	7802	8523	Fibers did not shear; sap-wood.....	88
	2971	4423	4990	5307	5602	5806	6010	6169	6341	6508	7756	8483	do.....	88
	1170	1760	1928	2182	2259	2413	2531	2613	2703	2849	3348	3583	Slight shearing of fibers; split at end.....	121
														288
	2041	3583	4191	4527	4831	5058	5285	5516	5702	5853	6849	7303	Slight shearing of fibers.....	442
	3130	4473	5017	5294	5525	5738	5969	6149	6314	6382	7521	8047	do.....	538
	2586	4173	4717	5035	5307	5557	5779	5933	6159	6373	7485	8097	do.....	1051
	2041	3742	4281	4527	4649	4899	5067	5248	5512	5606	6509	7054	do.....	1168
	2449	4173	4695	4926	5149	5262	5579	5770	5878	6069	7031	7648	do.....	1168
	1315	2744	3379	3697	3828	4032	4146	4314	4445	4545	5376	5851	do.....	153
	2032	3130	3484	3751	4073	4359	4427	4672	4804	4805	5738	6292	do.....	153
	2132	2994	3311	3475	3620	3751	3910	4037	4150	4219	4980	5489	do.....	839
	1111	2789	3397	3797	3992	4296	4527	4681	4904	5035			Slight shearing of fibers; split at end.....	838
	1702	3674	4486	4899	5104	5421	5634	5738	5996	6164	7303	8006	Slight shearing of fibers.....	237
	2812	4082	4409	4667	4890	5198	5389	5525	5715	5851	6895	7394	Sheared fibers.....	237
	1474	2667	3089	3420	3538	3720	3915	3974	4223	4296	4831	5479	do.....	129
	2890	4450	4990	5285	5470	5851	6023	6101	6310	6423	7190	7756	Sheared fibers; sap-wood.....	303
	2109	3366	3633	3910	4057	4196	4386	4522	4672	4877	5693	6169	do.....	362
	2449	4028	4740	5225	5479	5696	5946	6214		6509	7553	7892	Sheared fibers.....	740
	2590	4028	4572	5013	5316	5625	5887	6023	6292	6419	7553	8210	do.....	740
	1678	2903	3221	3465	3583	3738	3819	3910	3983	4119	4672	5035	do.....	917
	1406	2608	3026	3216	3339	3475	3593	3620	3697	3738	4219	4513	do.....	917
	1293	1973	2122	2223	2204	2486	2545	2649	2713	2758	3311	3706	do.....	586
	1325	2449	2767	2976	3198	3348	3479	3615	3710	3797	4400	4740	Sheared fibers; 3 millimeters knot in indented section.	605
	1474	2232	2518	2713	2835	2926	3057	3130	3257	3379	3847	4173	Sheared fibers.....	605
	1343	1946	2059	2218	2313	2354	2499	2581	2672	2722	3166	3529	do.....	8
	2041	3239	3388	3468	3576	3633	3742	3842	3951	4046	4445	4672	do.....	8
	2404	3588	4042	4445	4608	5080	5330	5539	5711	5906	6713	7212	do.....	32
	1384	2041	2291	2440	2563	2699	2867	2980	3075	3139			Slight shearing of fibers; split at end.....	32 ^a
	1951	2563	3035	3206	3379	3447	3583	3692	3729	3819	4219	4653	Sheared fibers.....	32 ^a
	1542	2767	3130	3329	3552	3774	3837	4119	4268	4409	5080		Slight shearing of fibers; split at end.....	49
	1951	3357	3464	3751	3946	4114	4332	4495	4672	4872	5851	5987	Sheared fibers.....	49
	1442	2722	3027	3311	3506	3701	3937	4105	4309	4482	5262	5579	Fibers did not shear.....	49 ^a
	2313	3221	3476	3718	3938	3865	3978	4110	4200	4296	4922	5353	Sheared fibers.....	49 ^a
	2313	3298	3556	3801	3969	4173	4309	4495	4686	4826	5670	6056	do.....	49 ^a
	997	1774	1932	2050	2109	2168	2263	2400	2368	2427	2740	2953	do.....	113
	1678	2486	2694	2850	3012	3207	3298	3397	3484	3588	4219	4513	do.....	113 ^a
	1515	2441	2513	2581	2654	2713	2776	2890	3030	3075	3434	3810	do.....	113 ^a

FOREST TREES OF NORTH AMERICA.

TABLE V.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Longitudinal compression. Ultimate strength, in kilos.	Remarks.
251. <i>Quercus alba</i> —continued.....	238	South Carolina	Bonneau's Depot	H. W. Ravenel	Rich, damp loam	10024	Triple flexure.....
	238	do	do	do	do	9617	Crushed at 89 millimeters from end.
	250	Virginia	Wytheville	H. Shriver	Clay	10026	Crushed at 13 millimeters from middle.
	250	do	do	do	do	9889	Crushed at middle; split perpendicular to rings.
	251	do	do	do	do	7576	Crushed on one face at 25 millimeters from middle.
	251	do	do	do	do	6940	Crushed at 16 millimeters knot 61 millimeters from end.
	259 ¹	do	do	do	Gravelly	9730	Triple diagonal flexure.....
	259 ²	do	do	do	do	8437	do
	403	Maryland	Charlestown Navy-yard.	S. H. Pook	do	8799	Fibers crushed in oblique lines at middle and at ends.
	403	do	do	do	do	9049	Fibers crushed at 32 millimeters from end.
	443	Tennessee	Nashville	A. Gattinger	Rich bottom	6532	Deflected at 102 millimeters from end; split at end.
	547	Alabama	Kemper's mill	C. Mohr	Alluvial	8690	Triple flexure; middle bend 25 millimeters from center.
	547	do	do	do	do	9528	do
	748	Florida	Chattahoochee	A. H. Curtiss	Clay	8663	Crushed fibers at 32 millimeters from end.
	749	do	do	do	do	8119	Triple flexure, deflected diagonally.
	749	do	do	do	do	7847	Triple flexure, deflected diagonally; knot at middle.
	895	Massachusetts		M. C. Beedle		7621	Crushed at 102 millimeters from end; split from end to end perpendicular to rings.
	895	do		do		6396	Split from end to end, opening season cracks.
	1050	do	North Reading	J. Robinson		7425	Crushed fibers at 32 millimeters from end.
	1050	do	do	do		7303	Triple flexure.....
	1257		Charlestown Navy-yard.	S. H. Pook		9480	do
	1257		do	do		8392	Crushed fibers at 32 millimeters from end.
252. <i>Quercus lobata</i> White Oak. Weeping Oak.	670	California	Redding	G. R. Vasey	Gravelly loam	7053	Crushed at 6 millimeters knots 31 millimeters from end.
	670	do	do	do	do	6532	Crushed at end in vicinity of knots.
253. <i>Quercus Garryana</i> White Oak.	985	Oregon	Weidler's saw-mill	G. Engelmann and C. S. Sargent.		7892	Triple flexure; bearing defective.
	985	do	do	do		8256	do
	988	do	Portland	do	Rich loam	7847	Triple flexure, deflected toward heart.
	988	do	do	do	do	7430	Triple flexure, deflected from heart.
	1027	do	Portland Furniture Company.	do		8483	Crushed at 127 millimeters from end and split along grain.
	1027	do	do	do		8483	Triple flexure.....
	1029	do	do	do		8754	do
	1029	do	do	do		7530	do
254. <i>Quercus obtusiloba</i> Post Oak. Iron Oak.	37 ¹	Kentucky	Harrodsburg	W. M. Linney	Shale	7462	do
	37 ²	do	do	do	do	6541	Crushed at 6 millimeters knot at middle.
	151	South Carolina	Bonneau's Depot	H. W. Ravenel	Rich upland	10002	Triple flexure.....
	151	do	do	do	do	8799	do
	256	Missouri	Allenton	G. W. Letterman	Clay	8437	do
	351	Alabama	Citronelle	C. Mohr	do	6078	Triple flexure; split along grain.
	351	do	do	do	do	6305	Crushed at 102 millimeters from end and split along grain.
	771	Florida	Aspalaga	A. H. Curtiss	Gravelly barrens	8618	Crushed at 6 millimeters knot 102 millimeters from end.
	771	do	do	do	do	7870	Triple flexure.....
255. <i>Quercus undulata</i> , var. <i>Gambellii</i> . Scrub Oak.	417	New Mexico	Pinos Altos mountains.	E. L. Greene		7439	do
	417	do	do	do		7961	Crushed at 25 millimeters from middle at 3 millimeters knot.
	525	Colorado	Engelmann's cañon	Robert Douglas	Rocky	6691	Triple flexure; split along grain.
	525	do	do	do	do	4581	Crushed at knots 32 millimeters from middle (dry rot).
256. <i>Quercus macrocarpa</i> Burr Oak. Mossy-oak. Overcup Oak.	79	Kentucky	Mercer county	W. M. Linney	Alluvial	8392	Triple flexure.....
	79 ²	do	do	do	do	8119	do

UNITED STATES UNDER COMPRESSION—Continued.

Direction of grain.	PRESSURE, IN KILOGRAMS, REQUIRED TO PRODUCE AN INDENTATION, IN MILLIMETERS, OF—												Remarks.	Office number.
	0.25	0.51	0.76	1.02	1.27	1.52	1.78	2.03	2.28	2.54	2.81	3.08		
	2744	3574	4014	4250	4545	4735	4976	5153	5325	5489	5613	7031	Slight shearing of fibers	238
	1633	2341	2617	2820	3094	3248	3434	3629	3765	3891	4500		Slight shearing of fibers; split at end	238
	2008	3447	3760	4019	4200	4468	4658	4799	5017	5194	6214	6759	do	250
	1633	2200	2504	2703	2894	3153	3329	3465					Fibers did not shear; split at end	250
	2313	3085	3311	3447	3488	3615	3738	3833	3887	3964	4355	4899	Sheared fibers	251
	1438	3815	3865	2567	2713	2858	3048	3166	3248	3357	3901	4400	do	251
	1860	2767	3016	3357	3574	3751	3933	4078	4223	4296	5103	5625	Slight shearing of fibers; split at end	250
	1882	2495	2622	2849	3026	3162	3334	3465	3583	3692			do	250
	2132	2449	2753	2948	3021	3166	3293	3411	3488	3583	4250		Sheared fibers; split at end	403
	1483	1987	2205	2023	2150	2303	2367	3048	3157	3252			Split at end	403
	1996	2080	3071	3162	3266	3343	3429	3479	3570	3633	4037		Sheared fibers	448
	2427	3130	3402	3633	3978	4223	4445	4790	4990	4740	6260	7045	Slight shearing of fibers	547
	1951	2812	3134	3429	3683	3901	4119	4273	4450	4595	5625	6033	Slight shearing of fibers; split at end	547
	989	1969	2631	2976	3239	3497	3692	3856					Split at end	748
	2499	3579	3974	4341	4672	4881	5076	5334	5525	5738	6759	7391	Indented without shearing fibers	749
	1542	2004	2367	3248	3502	3697	3842	3983	4092	4240	4944	5380	do	749
	2109	3311	3311	3389	3561	3656	3751	3833	3887	3960	4545	4854	Sheared fibers; split at side of stick	895
	1179	1951	2186	2336	2440	2527	2645	2731	2817	2903	3434	3765	Sheared fibers	895
	2313	3639	4001	4300	4536	4763	5035	5225	5271	5470	6704	7235	Slight shearing of fibers	1050
	1941	2699	3039	3284	3538	3720	3905	4105	4255	4404	5398	5942	Fibers did not shear	1050
	1438	2985	5212	3447	3574	3701	3842	4010	4114	4246	4890	5376	Sheared fibers	1257
	2223	2858	3289	3470	3611	3751	3892	3983	4042	4128	4513	4890	do	1257
	1384	2233	2581	2799	3016	3230	3388	3574	3724	3842	4617		Did not shear fibers; split at end	670
	1860	2450	2685	2862	3012	3180	3307	3438	3588	3710	4332	4854	Slight shearing of fibers; split at end	670
	2313	3352	3615	3837	4028	4182	4346	4405	4626	4758	5670	6214	Sheared fibers	985
	2041	2998	3289	3484	3742	3905	4155	4300	4427	4527	5579	5878	do	985
	4482	3493	3742	3882	4024	4246	4305	4391	4527	4653	5330	5897	do	988
	1896	2676	2794	2980	3157	3257	3425	3497	3674	3792	4377	4795	do	988
	2214	3797	4119	4291	4491	4695	4831	4990	5126	5252	6123	6600	Sheared fibers; specimen split into two pieces	1027
	1588	2767	3302	3567	3788	3946	4110	4204	4427	4495	5398		Sheared fibers; split at end	1027
	1896	3833	4110	4827	4454	4613	4753	4922	5062	5194	6069	6386	Sheared fibers	1029
	885	2295	2812	2912	3080	3125	3261	3402	3525	3647	3856		Sheared fibers; split at end	1029
	2245	3284	3515	3742	3992	4264	4436	4581	4749	4890	5761	6214	Slight shearing of fibers	37
	1928	3375	3963	4309	4604	4886	5062	5398	5570	5747	6804	7621	do	37
	3357	4563	5017	5234	5579	5823	6110	6314	6482	6664	7756	8119	do	151
	1619	2676	3316	3765	4095	4454	4763	5035	5339	5579			Split at end; fibers did not shear	151
	2123	3379	3842	4228	4477	4740	4944	5144	5298	5466	6396	6872	Slight shearing of fibers	256
	2586	3810	4237	4482	4681	4872	5062	5271	5398	5489	6396		Fibers did not shear; split at end	351
	1547	2449	2767	2921	3107	3316	3425	3538	3710	3819	4473	4990	Slight shearing of fibers	351
	2250	3479	3887	4110	4305	4527	4658	4744	4853	5002	5606	5860	Sheared fibers	771
	2495	3928	4400	4704	4899	5089	5307	5466	4643	5711	6359	6800	do	771
	1778	2971	3257	3529	3792	4037	4219	4423	4581	4704	5398	5992	Slight shearing of fibers	417
	1860	2785	3112	3438	3724	3933	4250	4432	4599	4708	5584	6010	do	417
	2032	3447	3833	4042	4246	4436	4559	4736	4863	5013	5942	6305	Sheared fibers	525
	2291	3747	4146	4377	4527	4644	4813	4953	5067	5216	6105	6532	do	525
	2141	2803	3021	3275	3484	3647	3801	3910	4033	4101	4763	5171	do	79
	2214	3057	3334	3470	3633	3747	3937	4105	4273	4350	5216	5606	Slight shearing of fibers	79

TABLE V.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Longitudinal compression: Ultimate strength, in kilos.	Remarks.
256. <i>Quercus macrocarpa</i> —cont'd.	137	Missouri	Allenton	G. W. Letterman	Moist upland	8754	Triple flexure
	143	Illinois	Waukegan	Robert Douglas	Rich	7076	do
	310	Texas	Dallas	J. Reverchon	Rich, moist	8609	Triple flexure; split at end
	310	do	do	do	do	8709	Crushed at middle of one face
	432	Tennessee	Nashville	A. Gattinger	Alluvial	7255	Triple flexure, deflected diagonally.
	831	Illinois	Winnebago county	M. S. Bebb	Loam	8256	Crushed at 19 millimeters from end.
	933	Texas	Austin	C. Mohr	Alluvial	7053	Crushed in vicinity of 3 millimeters knot.
	933	do	do	do	do	6613	Triple flexure, deflected diagonally.
	1071	Vermont	Charlotte	C. G. Pringle		7983	do
	1072	do	do	do		7326	Triple flexure
	1073	do	do	do		7938	Triple flexure, deflected diagonally.
257. <i>Quercus lyrata</i> . Over-cup Oak. Swamp Post Oak. Water White Oak.	424	Tennessee	Nashville	A. Gattinger	Low	5511	Split obliquely across the grain; cross-grained.
	424	do	do	do	do	7303	Crushed and split at ends.
	545	Mississippi	Kemper's mill	C. Mohr	Alluvial	8523	Crushed at middle of one face
	545	do	do	do	do	8754	Triple flexure, deflected diagonally.
	762	Florida	Chattahoochee	A. H. Curtiss	do	7756	Crushed near middle; stick worm-eaten.
	762	do	do	do	do	9344	Crushed near middle.
258. <i>Quercus bicolor</i> . Swamp White Oak.	12	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	7530	Triple flexure; split along grain.
	12	do	do	do	do	7212	Crushed fibers at end
	54	Missouri	Allenton	G. W. Letterman	Alluvial	8596	Triple flexure; split along grain.
	54	do	do	do	do	8596	Triple flexure.
	54 ^a	do	do	do	do	7983	Triple flexure, deflected diagonally.
	54 ^a	do	do	do	do	8437	Fibers crushed at 64 millimeters from end.
	846	Massachusetts	West Newbury	J. Robinson	Low, swampy	7022	Triple flexure, deflected diagonally.
	846	do	Arnold Arboretum	C. S. Sargent	Drift	7421	Triple flexure.
259. <i>Quercus Michauxii</i> . Basket Oak. Cow Oak.	240	South Carolina	Bonneau's Depot	H. W. Ravenel	Alluvial	6418	Triple flexure; split along grain.
	240	do	do	do	do	7756	do
	524	Alabama	Kemper's mill	C. Mohr	do	7847	Triple flexure.
	524	do	do	do	do	7983	Triple flexure, deflected diagonally.
	755	Florida	Chattahoochee	A. H. Curtiss	do	7938	Triple flexure at 8 millimeters knot 102 millimeters from end of concave side; split at end.
	755	do	do	do	do	8346	Triple flexure, deflected diagonally.
260. <i>Quercus prinus</i> . Hickory Oak. Rock Chestnut Oak.	31	Kentucky	Boyle county	W. M. Linney	Shale	8777	Crushed fibers at middle.
	31 ^a	do	do	do	do	9208	Crushed fibers at end
	35	do	do	do	Limestone	10560	Crushed fibers at 32 and at 127 millimeters from end.
	434	Tennessee	Nashville	A. Gattinger	Rocky upland	5942	Crushed and split at end.
	925	Alabama	Cullman	C. Mohr	Dry, rocky	9299	Crushed fibers at 76 millimeters from end.
	925	do	do	do	do	7892	Split at end; cross-grained
261. <i>Quercus prinoides</i> . Yellow Oak. Chestnut Oak. Chinquapin Oak.	34	Kentucky	Mercer county	W. M. Linney	Limestone	7938	Triple flexure; middle bend 25 millimeters from center; deflected from heart.
	34 ^a	do	Boyle county	do	Waverly shale	11022	Crushed at 5 millimeters knot 89 millimeters from end.
	273	Missouri	Allenton	G. W. Letterman	Limestone	8663	Crushed and split at end.
	287	do	do	do	Flinty	9276	Triple flexure, deflected from heart.
	323	Texas	Dallas	J. Reverchon	Calcareous	9163	Deflected 76 millimeters from end and split along grain.
	514	Tennessee	Nashville	A. Gattinger	Alluvial	9117	Triple flexure, deflected toward heart.
	514	do	do	do	do	9253	do
262. <i>Quercus Douglasii</i> . Mountain White Oak. Blue Oak.	688	California	Contra Costa county	G. R. Vasey	Clay	8709	Crushed fibers at 51 millimeters from middle
	698	do	do	do	do	9117	Crushed fibers near middle

UNITED STATES UNDER COMPRESSION—Continued.

Direction of grain.	PRESSURE, IN KILOGRAMS, REQUIRED TO PRODUCE AN INDENTATION, IN MILLIMETERS, OF—												Remarks.	Office number.
	0.25	0.51	0.76	1.02	1.27	1.52	1.78	2.03	2.28	2.54	2.81	3.08		
	2676	3887	4341	4817	5167	5403	5622	5915	6033	6260	7530	7938	Slight shearing of fibers	187
	749	2676	3184	3420	3588	3792	3933	4042	4178	4355			Short specimen, 120 millimeters long; split at ends ..	148
	1179	2254	2654	2812	3080	3230	3434	3602	3801	3910	4717	5126	Slight shearing; split at ends	310
	1678	2790	2980	3134	3356	3535	3692	3792	3937	4028	4653	5153	Slight shearing	310
	1960	2835	2935	3166	3393	3525	3638	3792	3919	3983	4881	5353	Sheared fibers	432
	1542	2549	2858	2967	3134	3248	3456	3624	3756	3901	4690		Slight shearing of fibers; split at end	331
	1769	2849	3148	3361	3484	3665	3779	2882	3964	4064	4763	5058	Slight shearing of fibers	933
	1678	2926	3366	3574	3715	3850	4028	4155	4237	4314	5062	5307	 do	933
	2078	3556	3996	4336	4604	4944	5035	5252	5430	5579	6577	7145	 do	1071
	2744	3456	3652	3937	4150	4341	4518	4613	4799	4969	6078	6668	 do	1072
	1996	2885	3280	3470	3701	3882	4114	4291	4391	4513	5307	5806	 do	1073
	2858	4500	5189	5606	5847	6196	6378	6500	6600	6722	7847	8119	 do	424
	2481	4105	4749	5162	6010	5869	6112	6864	6559	6695	7666	8074	Sheared fibers	424
	1452	1796	2068	2254	2422	2617	2790	2935	3062	3157	3720		Slight shearing of fibers; split at end	545
	2078	2971	3320	3574	3833	3996	4114	4296	4436	4590	5443	5806	Sheared fibers	545
	2041	2631	2926	3180	3366	3529	3661	3751	3882	3933	4559	4990	 do	762
	1461	2050	2313	2563	2722	2985	3157	3329	3438	3593	4336		Sheared fibers; split at end	762
	1905	2640	2740	2908	3094	3230	3370	3534	3615	3729	4346		Slight shearing of fibers; split at end	12
	1343	2073	2322	2495	2676	2844	2976	3125	3261	3348			do	12
	1687	2776	3071	3311	3579	3742	4014	4178	4350	4554	5479		Fibers did not shear; split at end	54
		2921	3329	3683	3955	4166	4500	4717	4944	5149	6169	6459	Slight shearing of fibers	54
	2313		3529	3683	3792	3942	3983	4237	4332	4441	5035	5625	Sheared fibers	54 ²
	1628	2368	2586	2744	2935	3094	3248	3388	3529	3661	4355		Slight shearing of fibers; split at end	54 ²
	2091	3239	3652	3919	4196	4445	4604	4735	4944	5098	6078	6459	Fibers did not shear	846
	1951	3198	3538	3856	4046	4287	4626	4786	4971	5158	6123	6759	 do	846
	1940	2767	3044	3339	3547	3674	3878	4028	4146	4309	5080		Slight shearing of fibers; split at end	240
	2359	3030	3402	3720	3978	4150	4486	4658	4849	5008	6033	6577	Fibers did not shear	240
	1896	2540	2867	2948	3130	3280	3447	3606	3720	3842	4382		Fibers did not shear; split at end	524
	1202	2050	2341	2572	2776	2930	3116	3261	3375	3461	4028	4382	Fibers did not shear	524
	3057	3656	4092	4482	4744	4990	5171	5416	5615	5793	6849	7394	Slight shearing of fibers	755
	1860	2994	3434	3847	4173	4409	4672	4908	5126	5294	6260	6927	 do	755
	2155	3266	3583	3674	3901	3992	4119	4191	4300	4366	4795		Slight shearing of fibers; split at end	31
	1542	2177	2504	2713	2921	3075	3239	3379	3479	3638			Split at ends	31 ¹
	1860	3452	3670	4042	4300	4531	4626	4886	4967	5262	5652		Sheared fibers	35
	2245	3329	3397	3674	3842	3983	4037	4146	4191	4246	4990	5285	 do	434
	1452	2676	3039	3270	3434	3574	3720	3856	3978	4073	4626		Sheared fibers; split at end	925
	1656	2980	3325	3520	3720	3878	4033	4160	4287	4436	5035		do	925
	1588	2989	3311	3402	3561	3701	3842	3928	4024	4150	4831	5298	Slight shearing of fibers	34
	1624	3166	3598	3783	3992	4196	4382	4559	4708	4872	5706		Slight shearing of fibers; split at end	34 ²
	2613	3847	4219	4495	4854	5103	5262	5498	5670	5838	7031	7756	Sheared fibers	273
	2223	3348	3697	3951	4164	4396	4563	5753	4904	5107	5851	6600	Sheared fibers; indented section covers 3 millimeters knot.	287
														323
	1170	2767	3425	3765	3969	4164	4436	4527	4717	4877	5829	6169	Slight shearing of fibers	514
	2404	3892	4261	4559	4804	5044	5216	5398	5579	5747	6985	7439	 do	514
	2767	4808	5642	6373	6713	7008	7248	7512	7756	7928	9026	9634	 do	688
	1769	3720	4491	4990	5262	5496	5797	5942	6314	6396	7666	8392	Fibers did not shear	688

TABLE V.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Longitudinal compression: Ultimate strength, in kilos.	Remarks.
263. <i>Quercus oblongifolia</i> <i>White Oak.</i>	655	California.....	San Diego county.	G. R. Vasey.....	Dry, gravelly.....	6260	Stick shattered at seasoning cracks.
	655	do	do	do	do	7021	Stick shattered at knots and seasoning cracks.
264. <i>Quercus grisea</i> <i>White Oak.</i>	698	Arizona	Santa Rita mountains.	G. Engelmann and C. S. Sargent.	Dry, rocky	7666	Crushed at 3 millimeters knot 25 millimeters from end.
265. <i>Quercus Durandii</i>	1103	Texas	Austin.....	S. B. Buckley.....	Damp, calcareous.	9140	Triple flexure; split along grain..
	1103	do	do	do	do	7892	Crushed at knots 51 millimeters from end.
	1103	do	do	do	do	8618	Failed at knots 51 and 70 millimeters from end.
267. <i>Quercus virens</i> <i>Live Oak.</i>	404	Florida	Charlestown Navy-yard.	S. H. Pook.....		10478	Crushed at middle and split; cross-grained.
	799	do	Saint John's river.	A. H. Curtiss.....	Sandy	9707	Crushed fibers at 25 millimeters from middle and at 25 millimeters from end.
	799	do	do	do	do	9934	Triple flexure; developed intersecting "Cooper lines."
	919	Alabama	Mobile county ..	C. Mohr	Rich, sandy.....	8165	Crushed fibers at end; split along side.
	919	do	do	do	do	8936	Crushed fibers at middle, 6 millimeters from knot.
	954	Texas	Matagorda bay ..	do	Sandy loam.....	6577	Crushed in vicinity of knots 102 millimeters from end.
	954	do	do	do	do	7439	Triple flexure.....
268. <i>Quercus chrysolepis</i> <i>Live Oak. Maul Oak. Val-paraiso Oak.</i>	649	California.....	San Bernardino ..	W. G. Wright.....		8845	Crushed fibers at 32 millimeters from middle.
	649	do	do	do		9480	Crushed fibers at end.....
	653	do	Marin county.....	G. R. Vasey.....	Gravelly.....	9072	Crushed at knot 51 millimeters from middle.
	653	do	do	do	do	7485	Crushed at knot 64 millimeters from end.
269. <i>Quercus Emoryi</i> <i>Black Oak.</i>	654	Arizona.....	Santa Rita mountains.	G. Engelmann and C. S. Sargent.	Dry, rocky.....	6759	Crushed at end and split.....
270. <i>Quercus agrifolia</i> <i>Encino. Coast Live Oak.</i>	663	California.....	Marin county.....	G. R. Vasey.....	Loam	6985	Crushed at two 6 millimeters knots 19 and 57 millimeters from end.
	663	do	do	do	do	7847	Crushed fibers at 57 millimeters from middle.
271. <i>Quercus Wislizeni</i> <i>Live Oak.</i>	685	do	Auburn.....	G. Engelmann		8052	Crushed and splintered at end ..
	685	do	do	do		9004	Crushed at end
272. <i>Quercus rubra</i> <i>Red Oak. Black Oak.</i>	7	Massachusetts....	Arnold Arboretum	C. S. Sargent.....	Drift.....	9617	Crushed at 25 millimeters from end; opened grain.
	7	do	do	do	do	10093	Crushed at 32 millimeters from end.
	45	Kentucky	Mercer county.....	W. M. Linney.....	Shale	6895	Triple flexure, deflected parallel to rings.
	45	do	do	do	do	7621	Triple flexure, deflected parallel to rings.
	45	do	do	do	do	6623	Crushed at 34 millimeters from end.
	89	Missouri.....	Allenton.....	G. W. Letterman..	Rich loam.....	10705	Crushed at 102 millimeters from end.
	89	do	do	do	do	10524	Crushed at 19 and at 89 millimeters from end.
	92	Kentucky	Mercer county.....	W. M. Linney.....	Alluvial	6917	Triple flexure, deflected perpendicular to rings.
	140	Michigan.....	Dansville	W. J. Beal.....	Sandy	7122	Crushed at end; splitting of wedge-shaped piece; cross-grained.
	141	do	do	do	do	7031	Crushed at end and at 25 millimeters from middle.
	146	Illinois.....	Waukegan.....	Robert Douglas ..	Gravelly.....	5625	Crushed at 6 millimeters knot 51 millimeters from end.
	215	Vermont.....	Charlotte	C. G. Pringle.....	do	8663	Triple flexure, deflected toward heart.
	215	do	do	do	do	9208	Crushed fibers at end
	217	do	do	do	do	9026	Triple flexure.....
	217	do	do	do	do	7802	Triple flexure, deflected parallel to rings.
	218	do	do	do	do	8981	Crushed fibers at end
	920	Mississippi.....	Enterprise.....	C. Mohr.....	Alluvial	6396	Crushed fibers at 25 millimeters from middle.
	920	do	do	do	do	5579	Crushed fibers at 51 millimeters from end.
	1043	Massachusetts....	North Reading ..	J. Robinson.....	Drift.....	9957	Triple flexure; middle bend 25 millimeters from center.
	1043	do	do	do	do	9063	Crushed at end and at 102 millimeters from end.
272. <i>Quercus rubra</i> , var. <i>Texana</i> ... <i>Red Oak.</i>	931	Texas.....	Austin.....	C. Mohr.....	Calcareous.....	9321	Crushed at knots 114 millimeters from end.
	931	do	do	do	do	9299	Triple flexure; middle deflection 25 millimeters from center.
273. <i>Quercus coccinea</i> <i>Scarlet Oak.</i>	752	Florida	Aspalaga	A. H. Curtiss.....	Clay	8074	Triple flexure, diagonal deflection.

UNITED STATES UNDER COMPRESSION—Continued.

Direction of grain.	PRESSURE, IN KILOGRAMS, REQUIRED TO PRODUCE AN INDENTATION, IN MILLIMETERS, OF—												Remarks.	Office number.
	0.25	0.51	0.76	1.02	1.27	1.52	1.78	2.03	2.28	2.54	2.81	3.08		
	1724	4717	6305	7349	7983	8437	8981	9390	9707	9979	1157	1293	Sheared fibers.....	655
	2359	4309	5171	5761	6078	6477	6795	7081	7372	7566			Split at ends; sap-wood.....	655
	2472	4219	4990	5398	5829	5987	6214	6419	6623	6777	7870		Split at end.....	698
	2200	3828	4354	4626	4940	5071	5353	5489	5657	5806	6377	7031	Slight shearing of fibers.....	1103
	2631	3856	4332	4667	4904	5216	5466	5643	5806	5987	7122	7606	do.....	1103
														1103
	3130	5534	6441	6782	7122	7358	7576	7802	7970	8265	9117	10047	Sheared fibers.....	404
	1724	3742	4445	4922	5176	5470	5706	5920	6105	6382	7606	8415	Slight shearing of fibers.....	799
	1701	2971	3579	3892	4259	4491	4753	4999	5285	5543	6340	7066	do.....	799
	2449	4468	5060	5398	5657	5851	6060	6223	6332	6513	7462	7802	Slight shearing of fibers; split at end.....	919
	1769	3311	3856	4146	4436	4563	4922	5053	5280	5362			Split at end.....	919
	1879	3629	4400	4908	5252	5602	5874	6128	6314	6577	8097	8845	Slight shearing of fibers.....	954
	1610	2994	3789	4150	4391	4744	5053	5339	5584	5856	7439	8188	do.....	954
	2563	4001	4445	4677	4971	5289	5557	5720	5878	6078	6985	7621	do.....	649
	1633	4128	4922	5421	5625	5965	6141	6332	6609	6745	7892	8483	do.....	649
	2087	3002	4187	4495	4844	5107	5280	5512	5733	5929	7212	7924	do.....	653
	1910	3456	3978	4364	4877	4971	5248	5570	5702	5929	7054	8006	do.....	653
	2758	4672	5693	6283	6646		7303	7621	7870	8192	9694		Slight shearing of fibers; split at end.....	654
	1424	2676	3393		4060	4377	4626	4881	5116	5312	6577	7258	Slight shearing of fibers.....	663
	1520	2604	3030	3212	3470	3665	3797	3960	4092	4223	5013	5670	do.....	663
	2813	3856	4454	4854	5058	5421	5643	5851	6005	6250	7457	7910	do.....	685
	1415	2685	3116	3402	3665	3896	4082	4264	4482	4695	5398	6305	do.....	685
	1637	2295	2495	2694	2899	3157	3266	3479	3692	3783	4445		Slight shearing of fibers; split at end.....	7
	1043	2046	2395	2622	2744	2939	3075	3289	3434	3488	4033		do.....	7
	1905	2168	2286	2400	2440	2586	2690	2785	2817	2875	3366	3674	Sheared fibers.....	45
	871	1651	1955	2028	2136	2286	2313	2468	2536	2599	3021		Sheared fibers; split at end.....	45
	1084	1851	1955	1987	2123	2232	2295	2350	2481	2536	2958	3357	Sheared fibers.....	45
	1796	2976	3393	3683	3892	4033	4359	4500	4633	4854	5715		Slight shearing of fibers; split at end.....	89
	2109	2880	3153	3357	3543	3783	3983	4209	4355	4491	5398		do.....	89
	2132	2540	2862	2971	3171	3370	3543	3656	3851	3951	4559	4990	Sheared fibers.....	92
	1869	2014	2164	2322	2422	2626	2803	2890	3003	3075	3774		Slight shearing of fibers; split at end.....	140
	1678	2254	2372	2518	2681	2785	2944	3021	3130	3175	3783	3901	Sheared fibers.....	141
	1778	2576	2722	2858	3026	3171	3293	3420	3525	3611	4309		Split at end.....	146
	1951	2404	2558	2835	3003	3588	3397	3568	3756	3937	5058	5851	Sheared fibers.....	215
	1111	2245	2531	2767	2899	3075	3193	3316	3393	3461	4164		Sheared fibers; split at end.....	215
	1837	2522	2699	2753	2944	3075	3166	3243	3388	3570	4264		Split at end.....	217
	1769	2454	2672	2808	2969	3121	3212	3343	3452	3570	4264		do.....	217
	1905	2631	3012	3166	3329	3543	3665	3701	3797	3946			Split at end; short specimen, 120 millimeters long ..	218
	1560	1910	1951	2028	2141	2209	2263	2318	2354	2440	2894	3166	Sheared fibers.....	920
	1084	1678	1787	1833	1869	1978	2041	2105	2164	2209	2622	2869	do.....	920
	1565	2118	2418	2549	2767	2908	2985	3134	3225	3302			Split at end.....	1043
	1724	2409	2703	3334	3484	3175	3307	3438	3570	3629	4445	4854	Split at end; fibers did not shear.....	1043
	2495	3310	4332	4772	4971	5252	5407	5489	5711	5856	6568	7349	Sheared fibers; split at end.....	931
	2109	3438	3882	4146	4359	4563	4844	5062	5252	5380	6608	7439	Slight shearing of fibers.....	931
	1120	2527	2899	3071	3234	3438	3674	4291	4687	4160	5112	5715	Fibers did not shear.....	753

FOREST TREES OF NORTH AMERICA.

TABLE V.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Longitudinal compression: Ultimate strength, in kilos.	Remarks.
274. <i>Quercus tinctoria</i> Black Oak. Yellow-bark Oak. Quercitron Oak. Yellow Oak.	17	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	8890	Crushed at end, also at 102 millimeters from end.
	17	do	do	do	do	9081	Crushed fibers near middle
	36	Kentucky	Danville Junction	W. M. Linney	Shale	7530	Crushed 51 millimeters from end at 3 millimeters knot.
	36 ²	do	do	do	do	9617	Crushed fibers on one face at 13 millimeters from middle.
	36 ³	do	do	do	Slate	7326	Deflected and split along grain from end to middle.
	74	Missouri	Allenton	G. W. Letterman	Rich upland	8890	Triple flexure, deflected parallel to rings.
	74	do	do	do	do		
	86	do	do	do	do	8256	Triple flexure, deflected from heart.
	244	Virginia	Wytheville	H. Shriver	Clay	5534	Triple flexure, deflected parallel to rings.
	244	do	do	do	do	5103	Crushed and split at end; brittle.
	247	do	do	do	do	8663	Crushed at 13 and at 76 millimeters from end.
	247	do	do	do	do	9562	Crushed at end.
275. <i>Quercus Kelloggii</i> Black Oak.	437	Tennessee	Nashville	A. Gattinger	do	7689	Crushed at 19 millimeters from end.
	628	Oregon	Saw-mill, Ashland	G. Engelmann and C. S. Sargent.		8233	Crushed at 51 millimeters from end.
	628	do	do	do		8301	Triple flexure.
	963	do	Eugene City	G. H. Collier		6396	Crushed and splintered at end
276. <i>Quercus nigra</i> Black Jack. Jack Oak.	963	do	do	do		5806	Splintered at end
	268	Missouri	Allenton	G. W. Letterman	Clay	7766	Triple flexure.
277. <i>Quercus falcata</i> Spanish Oak. Red Oak.	339	Alabama	Citronelle	C. Mohr	Sandy	8142	Crushed in vicinity of small knots.
	131	South Carolina	Bonneau's Depot	H. W. Ravenel	Rich loam	9208	Crushed at 64 millimeters from end.
278. <i>Quercus Catesbeii</i> Turkey Oak. Scrub Oak. Forked-leaf Black Jack. Black Jack.	131	do	do	do		9163	Crushed at 25 and at 102 millimeters from end.
	245	Virginia	Wytheville	H. Shriver	Clay	9730	Crushed at 19 millimeters from end.
	245	do	do	do	do	10093	Crushed at 25 millimeters from end.
	265 ²	do	Carroll county	do	do	9698	Crushed on one face at 25 millimeters from middle and at end.
	265 ³	do	do	do	do	9081	Crushed at 102 millimeters from end.
	548	Mississippi	Kemper's mill	C. Mohr	Rich loam	10006	Crushed at end and at 114 millimeters from end.
	548	do	do	do	do	9276	Crushed at 89 millimeters from end.
	342	Alabama	Cottage Hill	do	Barren, sandy	6809	Triple flexure, deflected diagonally.
	342	do	do	do	do	7734	Triple flexure; split along grain.
	770	Florida	Aspalaga	A. H. Curtiss	do	7212	Triple flexure; knot at middle.
279. <i>Quercus palustris</i> Pin Oak. Swamp Spanish Oak. Water Oak.	770	do	do	do		7507	Crushed at end and at 3 millimeters knot 51 millimeters from end.
	47	Missouri	Allenton	G. W. Letterman	Rich, alluvial	8437	Crushed at end
	47	do	do	do	do	8156	do
	282	do	do	do	do	6895	Triple flexure
280. <i>Quercus aquatica</i> Water Oak. Duck Oak. Possum Oak. Punk Oak.	282	do	do	do	do	7961	do
	264 ¹	Virginia	Carroll county	H. Shriver		7974	do
	264 ²	do	do	do		7802	Crushed at 25 millimeters from middle on one face.
	264 ³	do	do	do		7530	do
	349	Alabama	Cottage Hill	C. Mohr	Sandy loam	8709	Triple flexure
	349	do	do	do	do	8799	Crushed at 64 millimeters from end.
	511	Tennessee	Tullahoma	A. Gattinger	do	7167	Crushed at knots at middle and at 51 millimeters from end.
	511	do	do	do	do	7371	Crushed at 6 millimeters knot 76 millimeters from end.
	742	Georgia	Bainbridge	A. H. Curtiss	Alluvial	8256	Crushed fibers at middle
	742	do	do	do	do	8506	Crushed at 89 millimeters from end in vicinity of small knots.
281. <i>Quercus laurifolia</i> Laurel Oak.	756	Florida	Saint John's river	do	Sandy loam	8506	Triple flexure
	756	do	do	do	do	7884	Crushed in vicinity of 3 millimeters knot 102 millimeters from end.

UNITED STATES UNDER COMPRESSION—Continued.

Direction of grain.	PRESSURE, IN KILOGRAMS, REQUIRED TO PRODUCE AN INDENTATION, IN MILLIMETERS, OF—												Remarks.	Office number.
	0.25	0.51	0.76	1.02	1.27	1.52	1.78	2.03	2.28	2.54	4.81	5.08		
	2032	2313	2495	2812	2880	3048	3075	3221	3361	3470	4173		Slight shearing of fibers; split at end	17
	1542	2449	2681	2849	3012	3134	3348	3434	3529	3656	4346		do	17
	1637	2522	4150	3012	3162	3379	3525	3593	3747	3856	4491		do	36
	2313	3166	3357	3543	3674	3742	3983	4037	4110	4187	4980		Split at end; fibers did not shear	36 ²
	2518	3407	3665	3842	4064	4123	4287	4400	4626	4896		6532	Fibers indented without shearing	36 ²
	1769	2903	3529	3819	4028	4300	4518	4649	4935	5103	6283		Split at end; fibers did not shear	74
	1565	3266	3992	4436	4717	5013	5398	5670	5915	6146	7212	7965	Indented without shearing fibers	74
	1928	2858	3130	3230	3383	3588	3810	3892	4064	4196	4944		Split at end	86
	1343	2291	2322	2386	2395	2454	2518	2563	2649	2685	3085	3248	Sheared fibers; split at end	244
	835	1569	1633	1674	1751	1842	1878	1928	1951	2078	2404	2654	Sheared fibers	244
	1315	2313	2576	2790	2976	3153	3252	3368	3574	3665			Split at ends	247
	2019	2422	2667	2867	3035	3212	3338	3443	3566	3647	4377		Slight shearing of fibers; split at end	247
	1547	2277	2481	2726	2880	2130	3193	3407	3547	3683		4899	do	437
	1769	2672	2790	2803	2917	3003	3026	3026	3139	3225	3710	4037	Sheared fibers	628
	1642	2495	2681	2713	2776	2890	2994	3075	3193	3221	3856		Slight shearing of fibers; split at end	628
	980	2232	2840	2967	3066	3184	3352	3452					Shaky stick; split at ends	963
	1229	1880	2141	2313	2372	2513	2586	2617	2694	2758			do	963
	2948	4024	4277	4495	4672	4854	5026	5262	5407	5579	6486	7021	Slight shearing of fibers	268
	2767	4110	4010	4246	4491	4699	4881	5067	5262	5407	6392		Sheared fibers; split at end	339
	2336	3438	3701	4042	4264	4572	4922	5044	5216	5407	6214		Slight shearing of fibers; split at end	131
	975	2250	2799	3053	3284	3493	3665	3828	3933	4110	4877		do	131
	1987	2395	2572	2722	2803	2958	3098	3230	3302	3425			Split at end	245
	1406	1814	1887	2087	2245	2440	2622	2767	2980	3153	3993		Slight shearing of fibers; split at end	246
	1728	2563	2793	2885	3021	3130	3298	3402	3497	3574			Split at end; fibers not sheared	265 ²
	1384	2400	2617	2744	2835	2994	3125	3207	3330	3438	4128		Slight shearing of fibers; split at end	265 ²
	2436	2953	3293	3488	3638	3828	3942	4082	4178	4327			do	548
	1751	2867	6248	3536	3697	3983	4110	4264	4386	4522	5216		do	548
	2068	2753	2890	3066	3302	3429	3615	3751	3882	3983	4786	5262	Sheared fibers	342
	1792	2477	2703	2853	3026	3180	3302	3447	3579	3688	4468		Sheared fibers; split at end	342
	1597	2776	3221	3561	3756	4060	3815	4463	4717	4831	5851		Slight shearing of fibers; split at end	770
	2118	3411	3905	4259	4500	4817	5071	5303	5067	5706	6918	7666	Sheared fibers	770
	1338	2268	2640	2821	2935	3130	3202	3316	3470	3538	4287		Slight shearing of fibers; split at end	47
														47
	1456	2068	2295	2486	2703	2880	3039	3202	3352	3393	4150		Slight shearing of fibers; split at end	282
	2359	2939	3075	3343	3484	3715	3847	4055	4160	4327			Fibers did not shear; split at end	282
	1624	2109	2241	2359	2486	2527	2676	2753	2840	2980	3511		Slight shearing of fibers; split at end	264 ¹
	1406	1842	1932	2087	2155	2214	2341	2400	2454	2495	2948		do	264 ²
	1678	1955	2005	2096	2173	2295	2413	2481	2595	2676	3121		Sheared fibers; split at end	264 ²
	1769	2676	2790	3139	3311	3484	3674	3901	4060	4173			Fibers did not shear; split at end	349
	1506	2245	2490	2744	2866	3062	3221	3388	3538	3652	4377		do	349
	1542	2685	3130	3475	3692	3937	4128	4191	4445	4626	5534		Indented without shearing fibers	511
	1293	2313	2749	3012	3248	3393	3606	3783	3910	4037	5080		Slight shearing of fibers; split at end	511
	2858	3511	3851	4196	4495	4781	5013	5239	5416	5615	6849	7349	Sheared fibers	742
	1542	2926	3547	3429	4092	4814	4500	4690	4790	4949	5761		Slight shearing of fibers; split at end	742
	2858	3765	4219	4462	4735	5035	5262	5461	5661	5847	6736	7521	Sheared fibers	756
	1610	3039	3429	3720	3942	4164	4346	4500	4192	4364			Split at end; fibers did not shear	756

TABLE V.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Longitudinal compression: Ultimate strength, in kilos.	Remarks
281. <i>Quercus laurifolia</i> —continued.	801	Florida	Saint John's river.	A. H. Curtiss		9480	Crushed at 25 millimeters from middle.
	801	do	do	do		7734	Crushed at 89 millimeters from end and at end.
282. <i>Quercus heterophylla</i> — <i>Bartram's Oak.</i>	1171	New Jersey	Mount Holly	S. P. Sharples	Clay	5171	Crushed at 25 millimeters knot 51 millimeters from end.
	1171	do	do	do	do	8029	Crushed at 102 millimeters from end.
283. <i>Quercus cinerea</i> — <i>Upland Willow Oak. Blue Jack. Sand Jack.</i>	352	Alabama	Citronelle	C. Mohr	Pine-barren	7167	Split at end and splintered at 102 millimeters from end; brittle.
284. <i>Quercus hypoleuca</i>	674	Arizona	Santa Rita mountain.	G. Engelmann and C. S. Sargent.	Dry, rocky	7167	Triple flexure, deflected diagonally; split at end.
	674	do	do	do	do	2222	Cross-grained; oblique split 152 millimeters long.
285. <i>Quercus imbricaria</i> — <i>Shingle Oak. Laurel Oak.</i>	402	Kentucky	Harrodsburg	W. M. Linney	Utica shale	8029	Crushed at end
	402	do	do	do	do	9026	Triple flexure, deflected parallel to rings.
	50	Missouri	Allenton	G. W. Letterman	Rich, moist	8845	Triple flexure
	135	do	do	do	Rich loam	9458	Crushed at 76 millimeters from end.
286. <i>Quercus Phellos</i> — <i>Willow Oak. Peach Oak.</i>	512	Tennessee	Tullahoma	A. Gattinger	Moist, siliceous	5987	Crushed at 19 millimeters knot at end.
	512	do	do	do	do	6486	Triple flexure
287. <i>Quercus densiflora</i> — <i>Tanbark Oak. Chestnut Oak. Peach Oak.</i>	687	California	Marin county	G. R. Vasey	Gravelly	6464	Crushed at 38 millimeters from middle at 5 millimeters knot.
	687	do	do	do	do	8754	Triple flexure; middle bend 25 millimeters from middle.
288. <i>Castanopsis chrysophylla</i> — <i>Chinquapin.</i>	729	do	Mendocino county	A. Kellogg		5651	Crushed at end at 3 millimeters knot.
	729	do	do	do		8256	Crushed at end
289. <i>Castanea pumila</i> — <i>Chinquapin.</i>	573	Arkansas	Hot Springs	G. W. Letterman	Sandy loam	8156	Crushed at 51 millimeters from end.
	573	do	do	do	do	7689	Crushed at 5 millimeters knot 51 millimeters from middle.
290. <i>Castanea vulgaris</i> , var. <i>Americana</i> . <i>Chestnut.</i>	18	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	4137	Triple flexure, deflected parallel to rings.
	18	do	do	do	do	5298	Crushed 25 millimeters from middle at 3 millimeters knot.
	2581	Virginia	Fancy Gap	H. Shriver	Moist	7235	Crushed at 25 millimeters from end.
	2581	do	do	do	do	7485	Crushed at 102 millimeters from end.
	516	Tennessee	Nashville	A. Gattinger	Sandy	6373	Crushed at 44 millimeters from end.
291. <i>Fagus ferruginea</i> — <i>Beech.</i>	9	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	7847	Crushed at 32 millimeters from middle and split along grain.
	9	do	do	do	do	7076	Crushed at 64 millimeters from end.
	449	Kentucky	Mercer county	W. M. Linney	Hudson River shale	7566	Crushed at 76 millimeters from end.
	449	do	do	do	do	8006	Crushed at 25 and at 127 millimeters from end.
	119	Michigan	Danversville	W. J. Beal	Gravelly	8822	Crushed at end
	119	do	do	do	do	8346	Crushed at middle and at end.
	765	Florida	Chattahoochee	A. H. Curtiss	do	6496	Crushed at middle in vicinity of 13 millimeters knot.
	765	do	do	do	do	6827	Crushed at end
	853	Massachusetts	Hamilton	J. Robinson	do	8278	Crushed at 25 millimeters from end.
	853	do	do	do	do	7235	Crushed at 19 millimeters from end.
292. <i>Ostrya Virginica</i> — <i>Hop Hornbeam. Iron Wood. Lever Wood.</i>	11	do	Arnold Arboretum	C. S. Sargent	Drift	9390	Crushed at 51 millimeters from middle.
	11	do	do	do	do	9934	Triple flexure; middle bend 25 millimeters eccentric.
	877	do	Danvers	J. Robinson	Rich loam	9707	Triple flexure
	877	do	do	do	do	6359	Crushed at 89 millimeters from end.
	1047	do	North Reading	do		7983	Triple flexure, deflected diagonally perpendicular to rings.
	1047	do	do	do		8641	Crushed at 51 millimeters from end; opened grain.
293. <i>Carpinus Caroliniana</i> — <i>Hornbeam. Blue Beech. Water Beech. Iron Wood.</i>	46	Missouri	Allenton	G. W. Letterman	Damp, alluvial	6963	Deflected at middle and split at ends.
	73	Kentucky	Mercer county	W. M. Linney	Trenton limestone	9390	Triple flexure
	73	do	do	do	do	8573	do
	1038	Massachusetts	Danvers	J. Robinson	Gravelly	6949	Cross-grained; split at knots.

UNITED STATES UNDER COMPRESSION—Continued.

Direction of grain.	PRESSURE, IN KILOGRAMS, REQUIRED TO PRODUCE AN INDENTATION, IN MILLIMETERS, OF—												Remarks.	Office number.
	0.25	0.51	0.76	1.02	1.27	1.52	1.78	2.03	2.28	2.54	2.81	3.08		
	1805	2980	3348	3674	3901	4000	4291	4463	4527	4631	5652	6056	Slight shearing of fibers	801
	1438	2685	3157	3438	3647	3847	3978	4164	4309	4436	5262		Slight shearing of fibers; split at end.....	801
	1973	2713	2921	3157	3339	3493	3652	3828	4001	4137	5013		do.....	1171
	998	1878	2155	2291	2477	2617	2718	2844	3007	3116	3788		do.....	1171
		2132	2622	2958	3221	3375	3593	3738	3878	4019			Split at end	852
	1610	2948	3525	3769	3978	4191	4332	4463	4559	4672	5512		Slight shearing of fibers; split at end	674
	1842	3402	4128	4486	4717	4944	5153	5343	5489	5643	6623	7167	Sheared fibers.....	674
	1851	3153	3402	3611	3788	3960	4164	4341	4513	4699	5579	6419	Indented without shearing fibers	40 ^a
	1315	2586	3112	3434	3629	3878	4037	4196	4400	4572	5625	6214	do.....	40 ^a
	1270	2245	2744	2921	3139	3343	3538						Split at ends	50
	2358	3216	3402	3665	3937	4205	4414	4626	4844	5035	5874		do.....	185
	1610	2835	3248	3429	3611	3801	3951	4078	4205	4300	5171	5679	Indented without shearing fibers	512
	1148	2540	2890	3125	3293	3438	3602	3756	3878	3887	4631	5398	Slight shearing of fibers	512
	1792	2939	3153	3384	3583	3701	3810	3960	4082	4164	4899	5443	Sheared fibers.....	687
	1860	3012	3366	3525	3602	3806	4001	4191	4318	4436	5262	5738	do.....	687
	1089	1452	1610	1733	1837	1892	2005	2073	2105	2173	2586		Sheared fibers; split at end.....	729
	1325	1606	1778	1892	1987	2136	2227	2286	2354	2450	2994		do.....	729
	1225	1452	1746	1774	1960	2028	2182	2223	2295	2395			Slight shearing of fibers; split at end	573
	575	1338	1524	1665	1814	1914	1978	2041	2122	2214	2608		do.....	573
	1338	1474	1642	1746	1914	2019	2118	2204	2227	2250	2903	3130	Sheared fibers.....	18
	925	1506	1706	1787	1846	1910	1978	2028	2073	2123	2413	2554	do.....	18
	889	1284	1424	1524	1610	1687	1756	1810	1887	1932	2227		Sheared fibers; split at ends	258 ^a
	871	1461	1637	1715	1774	1860	1901	1982	2037	2091			Sheared fibers; split at end.....	258 ^b
		1080	1229	1297	1347	1452	1529	1597	1647	1706	2041		do.....	516
	2223	2808	3057	3212	3407	3652	3774	3905	4037	4169	4854	5625	Sheared fibers.....	9
	1452	2223	2536	2713	2858	3121	3288	3420	3565	3665	4491		Slight shearing of fibers; split at end.....	9
	1111	2744	2971	3162	3284	3470	3583	3701	3797	3874	4391	4877	Sheared fibers.....	44 ^a
	2631	3193	3329	3525	3652	3774	3874	3951	3987	4132	4740	5149	do.....	44 ^a
	1973	2740	3134	3248	3411	3552	3611	3697	3915	4009	4581	4990	do.....	119
	1860	2767	3016	3293	3493	3593	3710	3856	3928	4033			Sheared fibers; split at end	119
	1542	2291	2513	2703	2866	2994	3121	3280	3348	3456	4150	4581	Sheared fibers.....	765
	1379	2123	2359	2490	2658	2753	2875	3048	3089	3193	3720		Slight shearing of fibers; split at end.....	765
	1083	2427	2640	2855	2985	3121	3207	3338	3443	3579	4173		do.....	853
	1452	2241	2486	2685	2840	2994	3071	3202	3438	3532	4264		do.....	853
	2223	3352	3692	3964	4146	4327	4527	4658	4854	5080	6078		do.....	11
	1678	2391	2713	2894	3130	3202	3438	3615	3774	3924	4854		do.....	11
	2132	3470	3874	4155	4386	4622	4849	5080	5252	5479	6713		do.....	877
	1973	3039	3402	3720	3983	4219	4423	4604	4803	4908	5942		do.....	877
	1547	2586	2885	3103	3348	3536	3729	3833	3987	4119	4899	5579	Slight shearing of fibers	1047
	1315	2381	2740	3017	3184	3398	3588	3742	3937	4046	4990	5579	do.....	1047
	1542	2072	2935	3139	3329	3493	3606	3665	3792	3910			Split at ends	46
	1792	2767	3012	3206	3515	3706	3955	4060	4273	4355	5489	6010	Slight shearing of fibers	73
	1501	2454	2785	3066	3438	3593	3801	3978	4119	4246	5307		Sheared fibers; split at ends	73
	1247	2576	2935	3193	3339	3598	3774	3892	4087	4146	4944	5470	Sheared fibers.....	1038

TABLE V.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Longitudinal compression: Ultimate strength, in kilos.	Remarks.
BETULACEÆ.							
294. <i>Betula alba</i> , var. <i>populifolia</i> <i>White Birch. Old-field Birch. Gray Birch.</i>	10	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	6123	Triple flexure parallel to rings....
	10	do	do	do	do	5307	Crushed at 6 millimeters knot 102 millimeters from end.
	848	do	Danvers	J. Robinson	Gravelly	5262	Crushed and split at middle
295. <i>Betula papyrifera</i> <i>Canoe Birch. White Birch. Paper Birch.</i>	223	Vermont	Charlotte	C. G. Pringle			
	223	do	do	do			
	722	Montana	Missoula	Sereno Watson	Wet	8346	Crushed at 102 millimeters from end.
	722	do	do	do	do	6713	Crushed at 25 millimeters from middle; deflected diagonally.
	836	Massachusetts	Townsend	J. Robinson		7983	Crushed at 32 millimeters from middle on one face.
	836	do	do	do		6890	Crushed at 25 millimeters from end.
	990	Alaska	Chilcoot inlet	Paul Schultze		6486	Crushed at middle at 3 millimeters knot.
	990	do	do	do		6577	Crushed 76 millimeters from end at 3 millimeters knot.
	1065	Vermont	Charlotte	C. G. Pringle		8346	Crushed at 8 millimeters knot at middle.
	1065	do	do	do		7485	Crushed at 6 millimeters knot at middle.
	1066	do	do	do		7485	Crushed at 38 millimeters from end.
	1066	do	do	do		7552	Crushed at 6 millimeters knot 38 millimeters from end.
	1067	do	do	do		7847	Crushed at 76 millimeters from end and at end.
	1067	do	do	do		9662	Crushed at 38 millimeters from middle.
296. <i>Betula occidentalis</i> <i>Black Birch.</i>	528	Colorado	Engelmann's cañon	Robert Douglas	Wet, sandy	5670	Crushed at middle; opened grain three-fourths the length of stick.
	629	California	Strawberry valley	G. Engelmann and C. S. Sargent	Wet, peaty	6849	Crushed at middle and split along grain.
297. <i>Betula lutea</i> <i>Yellow Birch. Gray Birch.</i>	843	Massachusetts	Danvers	J. Robinson	Gravelly	8279	Crushed at 51 millimeters from end.
	843	do	do	do	do	10070	do
	1068	Vermont	Charlotte	C. G. Pringle	do	9934	do
	1068	do	do	do	do	10093	Triple diagonal flexure perpendicular to rings.
	1069	do	do	do	do	10623	Crushed at 61 millimeters from end.
	1069	do	do	do	do	10841	do
	1070	do	do	do	do	9208	Crushed at 3 millimeters knot 25 millimeters from middle.
	1070	do	do	do	do	10206	Crushed at middle
298. <i>Betula nigra</i> <i>Red Birch. River Birch.</i>	136	Missouri	Allenton	G. W. Letterman	Moist loam	7339	Crushed at 5 millimeters knot at middle.
	136	do	do	do	do	7122	Crushed at 10 millimeters knot 32 millimeters from end.
	841	Massachusetts	North Andover	J. Robinson	Alluvial	7249	Triple flexure perpendicular to rings.
	841	do	do	do	do	7666	Crushed at 44 millimeters from middle.
	842	do	do	do	do	6600	Triple flexure toward heart.
	842	do	do	do	do	6069	Crushed at 6 millimeters knot 38 millimeters from end.
299. <i>Betula lenta</i> <i>Cherry Birch. Black Birch. Sweet Birch. Mahogany Birch.</i>	4	do	Arnold Arboretum	C. S. Sargent	Drift	9072	Crushed at 31 millimeters from middle; deflected diagonally.
	4	do	do	do	do	8823	Triple flexure; middle bend 32 millimeters eccentric.
	221	Vermont	Charlotte	C. G. Pringle	Gravelly	11022	Crushed at 89 millimeters from end.
	221	do	do	do	do	10931	Crushed at middle
	844	Massachusetts	Danvers	J. Robinson	do	9480	Crushed at 102 millimeters from end.
	844	do	do	do	do	10115	Crushed at 19 millimeters from middle.
300. <i>Alnus maritima</i> <i>Seaside Alder.</i>	810	Delaware	Pepper's mills	W. M. Canby	Wet, sandy		
301. <i>Alnus rubra</i> <i>Alder.</i>	967	Alaska	Sitka	Paul Schultze		6976	Crushed at 32 millimeters from middle.
	967	do	do	do		6033	Crushed at middle
	991	Washington territory.	Puyallup	G. Engelmann and C. S. Sargent		6010	Crushed at 76 millimeters from end.
	991	do	do	do		6829	Triple flexure; middle bend 61 millimeters eccentric.
	1025	Oregon	Portland Furniture Company.	do		6750	Crushed at 25 millimeters from middle.

UNITED STATES UNDER COMPRESSION—Continued.

Direction of grain.	PRESSURE, IN KILOGRAMS, REQUIRED TO PRODUCE AN INDENTATION, IN MILLIMETERS, OF—												Remarks.	Office number.
	0.25	0.51	0.76	1.02	1.27	1.52	1.78	2.03	2.28	2.54	2.81	3.08		
1179	1619	1778	1846	1955	2055	2168	2223	2322	2390	2971	3198		Sheared fibers.....	10
1048	1547	1597	1701	1769	1877	1937	1991	2078	2173				Slight shearing of fibers; split at end.....	10
1406	2223	2332	2409	2495	2549	2604	2654	2703	2748	3311			do.....	848
1179	1660	1787	1868	1932	2014	2105	2150	2209	2254	2676	2903		Sheared fibers; indented section covers 3 millimeters knot.	223
830	1093	1170	1325	1356	1460	1565	1615	1698	1783	2087			Slight shearing of fibers; split at end.....	223
1656	1973	2114	2236	2295	2391	2531	2649	2731	2799	3334	3705		Sheared fibers.....	722
1189	1978	2223	2350	2495	2649	2740	2844	2948	3016	3529	3946		Slight shearing of fibers; indented section covers 3 millimeters knot.	722
1134	1701	1878	2028	2118	2250	2345	2431	2545	2635	3198	3629		Slight shearing of fibers; split at end.....	836
1343	1941	2064	2214	2295	2440	2549	2676	2771	2867	3470	3901		do.....	836
1384	2028	2168	2304	2391	2531	2676	2785	2862	2935	3543	3847		Sheared fibers.....	990
930	1470	1583	1687	1796	1864	1941	2023	2091	2177	2676	2994		do.....	990
1270		2064	2313	2427	2490	2554	2676	2748	2799	3288			Sheared fibers; split at end.....	1065
1261	1746	1883	2005	2077	2182	2295	2391	2513	2595	3207			do.....	1065
975	1338	1529	1592	1710	1796	1883	1932	2028	2114	2563	2858		do.....	1066
1021	1488	1633	1715	1819	1892	1951	2046	2073	2200	2667			Slight shearing of fibers; split at end.....	1066
1179	1533	1656	1769	1892	1978	2150	2209	2304	2391	2939	3357		Slight shearing of fibers.....	1067
794	1325	1447	1579	1683	1765	1846	1941	2023	2082	2567			Slight shearing of fibers; split at end.....	1067
930	1837	2168	2340	2459	2599	2685	2744	2862	2930	3348	3652		Sheared fibers; split at end.....	528
749	1320	1424	1533	1597	1669	1751	1805	1892	1937	2395	2699		Slight shearing of fibers.....	629
1746	2359	2586	2722	2867	3030	3121	3243	3357	3561	4332	4536		do.....	843
1742	2622	2858	3062	3130	3311	3470	3574	3661	3774	4423	4590		do.....	843
1520	1987	2214	2404	2477	2576	2731	2794	2880	2998	3652			Sheared fibers; split at end.....	1068
1315	1769	1987	2168	2282	2345	2504	2572	2669	2776				Slight shearing of fibers; split at end.....	1068
1216	2028	2254	2449	2567	2708	2758	2894	2998	3071	3606			Sheared fibers; split at end.....	1069
1343	1887	2118	2254	2368	2481	2590	2654	2749	2799				Split at ends; fibers did not shear.....	1069
1642	2495	2699	2858	2948	3107	3221	3343	3456	3574	4241	4672		Sheared fibers.....	1070
1021	1542	1733	1851	2000	2132	2241	2345	2463	2567	3130			Split at end; fibers did not shear.....	1070
1225	2108	2472	2685	2790	2889	3016	3085	3162	3216	3742	4014		Slight shearing of fibers.....	136
1216	1497	1637	1760	1851	1982	2064	2205	2308	2345		2880		Sheared fibers; split at end.....	136
889	1343	1642	1801	1923	1982	2046	2159	2182	2232	2581	2912		Sheared fibers.....	841
894	1379	1542	1669	1765	1855	1910	1987	2037	2082	2563	2799		Slight shearing of fibers.....	841
1547	2123	2254	2377	2522	2626	2717	2803	2894	2989	3674	4105		do.....	842
1120	1515	1624	1737	1851	1896	2000	2087	2141	2205	2209	2948		do.....	842
2136	3561	4114	4386	4572	4753	4990	5112	5257	5443	6396	6922		do.....	4
1497	2440	2713	2894	3130	3230	3397	3574	3656	3815	4626			do.....	4
1665	2685	3089	3334	3497	3720	3856	4073	4205	4305	5216	5761		Slight shearing of fibers.....	844
1179	2336	2781	3030	3261	3411	3583	3801	3955	4101	5022	5761		do.....	844
1270	1810	1932	2023	2068	2164	2218	2259	2341	2400	2812			Sheared fibers; split at end.....	810
1089	1270	1347	1397	1474	1492	1547	1619	1628	1665	1933	2041		Sheared fibers.....	967
1002	1724	1923	1996	2032	2127	2177	2254	2308	2354	2803	3039		do.....	967
1774	2064	2223	2377	2481	2572	2636	2753	2808	2930	3574	3910		do.....	991
975	1257	1351	1410	1482	1568	1632	1736	1780	1827	2198	2427		do.....	991
767	1361	1515	1615	1699	1756	1896	1946	1987	2078	2449	2723		do.....	1025

TABLE V.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Longitudinal compression: Ultimate strength, in kilos.	Remarks.
301. <i>Alnus rubra</i> —continued	1025	Oregon	Portland Furniture Company.	G. Engelmann and C. S. Sargent.		7258	Crushed at 102 millimeters from end.
302. <i>Alnus rhombifolia</i>	635	do	Ashland	do		7022	Crushed at 25 millimeters from middle.
<i>Alder.</i>	717	Montana	Missoula	Sereno Watson ..	Wet	4990	Crushed at 114 millimeters from end at 3 millimeters knot.
	717	do	do	do	do	4899	Crushed at 102 millimeters from end on one face.
	979	Oregon	Drain	C. S. Sargent ..	Moist loam	5171	Crushed at 25 millimeters from middle.
	979	do	do	do	do	6396	Triple flexure
303. <i>Alnus oblongifolia</i>	694	California	San Bernardino ..	W. G. Wright ..		4445	Crushed at 89 millimeters from end at 3 millimeters knot.
<i>Alder.</i>	694	do	do	do		4459	Crushed at 102 millimeters from end at 3 millimeters knot.
305. <i>Alnus incana</i>	374	Vermont	Hinesburg	C. G. Pringle ..	Wet loam	4617	Crushed at 25 millimeters from middle.
<i>Speckled Alder. Hoary Alder. Black Alder.</i>							
SALICACEÆ.							
306. <i>Salix nigra</i>	232	do	Shelburne	do	Wet, sandy	4545	Triple flexure perpendicular to rings.
<i>Black Willow.</i>	855	Massachusetts ..	Topsfield	J. Robinson	Wet loam	2277	Crushed at 5 millimeters knot 51 millimeters from end; cross-grained.
307. <i>Salix amygdaloides</i>	908	Colorado	Cañon City	E. Weston		3498	Crushed at 6 millimeters knot at middle.
<i>Willow.</i>	908	do	do	do		4355	Triple flexure; split along grain between rings.
	911	do	do	do		5171	Crushed at 51 millimeters from end; split along grain between rings.
	911	do	do	do		3878	Crushed at 5 millimeters knot 51 millimeters from middle.
308. <i>Salix laevigata</i>	690	California	Santa Cruz	G. Engelmann and C. S. Sargent.	Moist, sandy	5012	Triple flexure
<i>Willow.</i>	690	do	do	do	do	5216	do
309. <i>Salix lasiandra</i> , var. <i>lancifolia</i> .	640	do	Strawberry valley	do	Moist, rich	4378	Crushed at 51 millimeters from end in vicinity of knots.
	981	Oregon	Portland	F. Skinner	Alluvial	5942	Crushed 13 millimeters from middle at 2 millimeters knot.
	981	do	do	do	do	6056	Crushed at 89 millimeters from end and at end.
309. <i>Salix lasiandra</i> , var. <i>Fendleri</i> .	889	Utah	City Creek cañon	M. E. Jones	Gravelly	4581	Crushed at 19 millimeters from middle at 3 millimeters knot.
313. <i>Salix flavescens</i>	721	Montana	Pattée's cañon, Missoula.	Sereno Watson ..	Rich, moist	6532	Crushed near middle on one face.
	721	do	do	do	do	6532	Crushed at 3 millimeters knot 76 millimeters from end.
313. <i>Salix flavescens</i> , var. <i>Scouleriana</i> .	972	Washington territory.	Seattle	G. Engelmann and C. S. Sargent.	Moist, sandy	8074	Crushed at middle
<i>Black Willow.</i>	972	do	do	do	do	6895	Crushed at 5 millimeters knot near middle.
314. <i>Salix Hookeriana</i>	966	Oregon	Winchester bay ..	do	Sandy saline	7031	Triple flexure, deflected diagonally.
	966	do	do	do	do	6623	Triple flexure
316. <i>Salix lasiolepis</i>	669	California	Santa Cruz	G. Engelmann and S. C. Sargent.	Sandy loam	6169	Crushed near middle on one face.
<i>Willow.</i>							
318. <i>Populus tremuloides</i>	272	Colorado	Alpine	T. S. Brandegee ..	Damp	4219	Crushed at 51 and at 114 millimeters from end; split along grain.
<i>Ash. Quaking Asp.</i>	272	do	do	do	do	4717	Triple flexure parallel to rings.
	1035	Massachusetts ..	Danvers	J. Robinson	Gravelly	6260	Crushed at 38 millimeters from middle on one face.
	1035	do	do	do	do	5942	do
319. <i>Populus grandidentata</i>	847	do	do	do	do	5625	Crushed at 89 millimeters from end.
<i>Poplar.</i>	847	do	do	do	do	5829	Triple flexure; middle bend 38 millimeters eccentric.
320. <i>Populus heterophylla</i>	522	Tennessee	Nashville	A. Gattinger	Alluvial	4527	Crushed at 16 millimeters knot 102 millimeters from end.
<i>River Cottonwood. Swamp Cottonwood.</i>							
321. <i>Populus balsamifera</i>	961	Alaska	Chilcoot inlet ..	Paul Schultze ..	do	4672	Crushed at 6 millimeters knot 25 millimeters from end.
<i>Balsam. Tacamahac. Balm of Gilead.</i>	961	do	do	do	do	5379	Triple flexure; middle bend 32 millimeters eccentric.
321. <i>Populus balsamifera</i> , var. <i>canadica</i> .	1054	Massachusetts ..	Topsfield	J. Robinson	Gravelly	4527	Crushed at 76 millimeters from end.
	1054	do	do	do	do	4309	Triple flexure perpendicular to rings.

UNITED STATES UNDER COMPRESSION—Continued.

Direction of grain.	PRESSURE, IN KILOGRAMS, REQUIRED TO PRODUCE AN INDENTATION, IN MILLIMETERS, OF—											Remarks.	Office number.
	0.25	0.51	0.76	1.02	1.27	1.52	1.78	2.03	2.28	2.54	4.81	5.08	
	953	1243	1352	1461	1529	1628	1715	1709	1805	1910	2254	Sheared fibers; split at end	1025
	916	1529	1665	1769	1851	1937	1996	2068	2132	2200	2486	Sheared fibers	635
	631	953	1043	1071	1080	1120	1152	1179	1234	1279	1442	do	717
	499	857	903	948	998	1043	1084	1129	1166	1220	1497	do	717
	500	944	1052	1084	1125	1193	1270	1315	1347	1388	1669	do	979
	667	1030	1111	1175	1229	1288	1347	1397	1415	1483	1678	do	979
	930	1089	1120	1202	1257	1302	1361	1402	1443	1489	1765	do	694
	658	939	998	1080	1120	1207	1225	1266	1311	1343	1624	do	694
	907	1293	1397	1569	1628	1719	1824	1896	1941	2032	2395	Slight shearing of fibers; split at end; specimen 120 millimeters long.	232
	644	993	1143	1297	1361	1442	1529	1574	1642	1733	2155	Slight shearing of fibers	855
	953	1084	1202	1252	1347	1393	1447	1524	1574	1619	2019	do	908
	1043	1220	1361	1438	1479	1547	1642	1674	1733	1814	2168	do	908
	499	907	1016	1075	1106	1234	1306	1347	1393	1452	1774	do	911
	953	1016	1080	1094	1184	1216	1257	1306	1352	1624	1833	do	911
	885	1215	1424	1488	1569	1651	1719	1814	1923	1951	2305	Sheared fibers	690
	1202	1851	1987	2123	2218	2286	2350	2482	2504	2590	3029	do	690
	1134	1497	1560	1619	1674	1760	1801	1846	1892	1928	2214	do	640
	1030	1207	1270	1388	1447	1547	1569	1624	1674	1715	2032	do	981
	635	880	943	1016	1080	1120	1170	1220	1266	1302	1570	do	981
	993	1107	1170	1266	1311	1384	1438	1488	1533	1579	1905	Slight shearing of fibers	889
	671	1229	1379	1479	1565	1615	1724	1787	1833	1865	2132	Slight shearing of fibers; split at end	721
	771	1252	1397	1565	1597	1674	1729	1801	1855	1928	2313	Sheared fibers	721
	862	1021	1384	1615	1678	1719	1810	1960	2019	2078	2481	Slight shearing of fibers; split at end	972
	1021	1687	1982	2168	2359	2486	2567	2685	2731	2858	3311	Sheared fibers; split at end; indented section covers 6 millimeters knot.	972
	1066	1529	1633	1706	1769	1851	1937	1966	2069	2118	2541	Slight shearing of fibers; split at end	966
	1406	1851	2064	2168	2241	2291	2391	2454	2518	2595	2971	Sheared fibers	669
	948	1134	1202	1257	1279	1311	1352	1388	1393	1397	1610	do	272*
	658	735	739	758	771	807	839	848	862	898	1025	do	272*
	821	1397	1565	1637	1710	1756	1824	1905	1962	1991	2268	do	1035
	862	1267	1261	1315	1365	1411	1470	1524	1569	1578	1860	do	1035
	640	885	934	980	1030	1075	1120	1161	1170	1211	1474	do	847
	658	844	880	903	957	998	1034	1080	1129	1170	1402	do	847
	885	1084	1184	1288	1384	1438	1511	1547	1633	1669	1960	Sheared fibers; split at end; specimen 120 millimeters long.	523
	817	1089	1125	1179	1238	1288	1348	1384	1393	1415	1651	Sheared fibers	961
	885	1030	1075	1125	1166	1166	1179	1220	1261	1306	1488	do	961
	889	943	1052	1116	1157	1198	1261	1302	1320	1361	1656	Slight shearing of fibers	1054
	522	730	807	852	903	957	1002	1043	1084	1125	1315	do	1054

TABLE V.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Longitudinal compression: Ultimate strength, in kilos.	Remarks.
322. <i>Populus angustifolia</i> <i>Black Cottonwood.</i>	552	Colorado.....	Manitou Springs..	Robert Douglas..	Sandy loam.....	4332	Failed at 6 millimeters knot 25 millimeters from middle.
323. <i>Populus trichocarpa</i> <i>Black Cottonwood. Balsam Cottonwood.</i>	1012	Oregon.....	Saint John's Barrel Factory, Portland.	F. Skinner.....		5851	Crushed at 38 millimeters from middle.
	1012	do.....	do.....	do.....		6214	Crushed at 51 millimeters from end.
	1028	do.....	Portland Furniture Company.	G. Engelmann and C. S. Sargent.		6192	Crushed at 38 millimeters from middle.
	1028	do.....	do.....	do.....		6713	Crushed at 25 millimeters from middle.
324. <i>Populus monilifera</i> <i>Cottonwood. Necklace Poplar. Carolina Poplar. Big Cottonwood.</i>	255	Missouri.....	Allenton.....	G. W. Letterman..	Alluvial.....	4763	Triple flexure.....
	304	do.....	do.....	do.....	do.....	4264	Crushed at middle and opened along grain.
	304	do.....	do.....	do.....	do.....	4541	Crushed at 102 millimeters from end.
	309	Texas.....	Dallas.....	J. Reverchon.....	do.....	6260	Crushed at 38 millimeters from end.
	309	do.....	do.....	do.....	do.....	6214	Crushed at middle.....
	754	Florida.....	Chattahoochee....	A. H. Curtis.....	do.....	6192	Crushed at 6 millimeters knot 38 millimeters from middle.
	754	do.....	do.....	do.....	do.....	7326	Crushed at 25 millimeters from middle.
325. <i>Populus Fremontii</i> <i>Cottonwood.</i>	659	California.....	Sacramento valley	G. R. Vasey.....	do.....	5967	Crushed at 38 millimeters from middle at 3 millimeters knot.
	659	do.....	do.....	do.....	do.....	6123	Crushed at 38 millimeters from end.
325. <i>Populus Fremontii</i> , var. <i>Wislizeni</i> . <i>Cottonwood. White Cottonwood.</i>	646	do.....	San Bernardino...	W. G. Wright.....	do.....	5625	Crushed at 25 millimeters from middle.
	646	do.....	do.....	do.....	do.....	5216	Triple flexure; middle bend 19 millimeters eccentric.
	909	Colorado.....	Cañon City.....	do.....		6759	Crushed at 89 millimeters from end.
	909	do.....	do.....	do.....		6713	Crushed at 5 millimeters knot 64 millimeters from end.
	909	do.....	do.....	do.....		5466	Crushed at 25 millimeters from end in vicinity of 5 millimeters knot.
	909	do.....	do.....	do.....		5919	Triple flexure.....
CONIFERÆ.							
326. <i>Libocedrus decurrens</i> <i>White Cedar. Bastard Cedar. Post Cedar. Incense Cedar.</i>	634	California.....	Saw-mill, Strawberry valley.	G. Engelmann and C. S. Sargent.		5618	Crushed at 25 millimeters from middle.
	634	do.....	do.....	do.....		6115	Crushed at 102 millimeters from end; split whole length of specimen.
	662	do.....	Saw-mill, San Bernardino mountains.	W. G. Wright.....		8754	Crushed at middle.....
	662	do.....	do.....	do.....		5299	Crushed at end; opened along grain.
327. <i>Thuja occidentalis</i> <i>White Cedar. Arbor-vita.</i>	379	Vermont.....	Monkton.....	C. G. Pringle.....	Cold, peaty.....	4626	Crushed at 25 millimeters from end.
	379	do.....	do.....	do.....	do.....	4545	Crushed at 13 and at 57 millimeters from end.
	782	New Brunswick..		Intercolonial railway.		5534	Crushed at 25 and at 102 millimeters from end.
	782	do.....	do.....	do.....		5035	Triple flexure.....
	783	do.....	Bridgeton.....	Ed. Sinclair.....		5579	Crushed at 13 and at 102 millimeters from end.
	783	do.....	do.....	do.....		5398	Triple flexure; middle bend 25 millimeters eccentric.
	790	Province of Quebec	Amqui.....	A. Grant.....		4355	Crushed at 89 millimeters from end and at end.
	790	do.....	do.....	do.....		4785	Triple flexure; small knots at middle.
	792	do.....		Grand Trunk railway.		2994	Crushed at 51 millimeters from end.
	792	do.....	do.....	do.....		3221	Crushed at 25 millimeters from end.
	796	do.....	do.....	do.....		4545	Crushed at 32 millimeters from end.
	796	do.....	do.....	do.....		5103	Triple diagonal flexure.....
	874	Maine.....	Mattawamkeag...	J. Robinson.....		5965	Crushed at 89 millimeters from end.
	874	do.....	do.....	do.....		6532	Triple flexure.....
328. <i>Thuja gigantea</i> <i>Red Cedar. Canoe Cedar.</i>	1099	Wisconsin.....	Eau Claire.....	H. C. Putnam.....	Drift.....	4423	Split at knot at middle; sides swelled.
	1099	do.....	do.....	do.....	do.....	5806	Crushed at 51 millimeters from end; split along grain.
	1017	Oregon.....	Weidler's saw-mill, Portland.	G. Engelmann and C. S. Sargent.		6895	Crushed at worm-hole 51 millimeters from end; split along grain.
	1017	do.....	do.....	do.....		8301	Crushed at 32 millimeters from middle; sudden fracture.
	1021	do.....	Portland Furniture Company.	do.....		6396	Crushed at 83 millimeters from end.
	1021	do.....	do.....	do.....			

UNITED STATES UNDER COMPRESSION—Continued.

Direction of grain.	PRESSURE, IN KILOGRAMS, REQUIRED TO PRODUCE AN INDENTATION, IN MILLIMETERS, OF—												Remarks.	Office number.
	0.25	0.51	0.76	1.02	1.27	1.52	1.78	2.03	2.28	2.54	2.81	3.08		
	454	944	1111	1166	1225	1306	1343	1356	1397	1442	1669	1905	Slight shearing of fibers	552
	749	907	1002	1043	1080	1120	1161	1184	1225	1261	1474		Sheared fibers	1012
	504	712	793	817	889	934	948	993	1030	1039			Slight shearing of fibers; split at end	1012
	930	998	1093	1166	1216	1279	1338	1365	1429	1474			do	1028
	449	712	789	852	889	939	998	1043	1084	1125	1384		do	1028
		595	653	680	726	766	807	848	852	875	1030	1179	Sheared fibers	255
	666	898	939	930	998	1043	1080	1120	1166	1198	1315	1452	do	304
	408	608	685	726	780	830	875	903	948	984	1179		Slight shearing of fibers; split at end	304
	1588	2254	2395	2436	2490	2567	2622	2703	2753	2807	3302		do	309
	522	1043	1157	1229	1315	1356	1447	1488	1538	1597	2032		do	309
	862	1320	1488	1542	1628	1687	1801	1851	1896	1946	2223		Sheared fibers; split at end	754
	839	1143	1220	1306	1352	1397	1442	1488	1529	1574	1837			754
	934	1216	1311	1352	1438	1515	1569	1619	1665	1728	2087		Sheared fibers; split at end	650
	703	998	1179	1252	1325	1388	1442	1556	1588	1647	2023		do	650
	2064	2971	3143	3216	3307	3375	3420	3479	3529	3574	4037		do	646
	653	1125	1220	1338	1397	1470	1574	1660	1724	1805	2177	2495	Sheared fibers	646
	1120	1311	1411	1438	1533	1637	1719	1756	1801	1869	2177		Slight shearing of fibers; split at end	909
	658	925	1061	1107	1189	1239	1279	1352	1397	1442	1701	1833	Sheared fibers; split at end	909
	635	835	880	930	975	1016	1071	1120	1166	1211	1433	1610	do	912
	817	1052	1116	1179	1243	1288	1338	1397	1447	1488	1760		do	912
	1071	1179	1216	1270	1315	1365	1402	1438	1442	1465	1628	1769	Sheared fibers	634
	1007	1202	1284	1447	1470	1492	1538	1574	1583	1597	1787	1896	do	634
	956	1334	1393	1493	1533	1578	1619	1669	1710	1751	1941	2046	Sheared fibers; split along indented face	662
	1071	1805	1828	1864	1928	2000	2087	2182	2250	2400	2622		Sheared fibers; split at end	662
	522	703	744	785	812	835	875	894	903	984	1034	1134	Sheared fibers	379
	508	703	717	721	730	739	753	762	798	807	934	993	do	379
	821	1012	1098	1116	1161	1175	1202	1225	1261	1270	1420	1542	do	782
	490	857	896	934	939	948	966	979	989	1007	1116	1229	Sheared fibers; split along indented face	782
	621	880	948	1007	1080	1034	1066	1075	1089	1112	1243	1288	Sheared fibers	783
	494	821	975	1034	1071	1080	1102	1116	1120	1134	1216	1248	Sheared fibers; indented face covers 3 millimeters knot	788
	631	989	1016	1030	1057	1075	1107	1116	1129	1161	1311	1488	Sheared fibers	700
	680	839	934	957	989	1016	1030	1043	1075	1098	1215	1325	do	790
	549	721	767	789	812	852	875	884	894	912	1043	1075	do	792
	404	599	626	635	649	662	676	680	694	708	794	839	do	792
	671	866	907	962	1002	1043	1075	1116	1157	1170	1397		do	706
	703	871	896	939	964	1025	1039	1057	1075	1102	1261	1325	do	796
	594	789	807	866	893	907	930	939	957	980			Sheared fibers; split at end	874
	717	953	993	1025	1039	1048	1057	1075	1089	1116	1216	1293	Sheared fibers	874
	572	944	1034	1075	1084	1125	1161	1193	1220	1270	1497	1610	do	1009
	658	889	953	1016	1057	1107	1152	1207	1252	1266	1447	1542	do	1009
	712	803	875	916	964	1034	1071	1093	1116	1161			Slight shearing of fibers; split at end	1017
	1080	1089	1117	1315	1415	1501	1497	1778	1406	1333	1746	1769	Sheared fibers	1017
	567	685	753	803	844	857	896	912	939	966	1116		Sheared fibers; split at end	1021
	980	1148	1107	1143	1211	1266	1288	1334	1370	1384	1579	1728	Sheared fibers	1021

FOREST TREES OF NORTH AMERICA.

TABLE V.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Longitudinal compressive strength, in kilos.	Remarks.
329. <i>Chamaecyparis sphaeroides</i> <i>White Cedar.</i>	850	Alabama	Cottage Hill	C. Mohr	Sandy, wet.	4400	Crushed at 25 millimeters from end.
	850	do	do	do	do	4105	do
	850	Massachusetts	Beverly	J. Robinson	Swampy	4060	Crushed at 76 millimeters from end at 3 millimeters knot.
	850	do	do	do	do	3756	Crushed at middle in vicinity of 3 millimeters knots.
	851	do	do	do	do	4014	Crushed at end; opened grain
	851	do	do	do	do	4581	Crushed at 3 millimeters knot 51 millimeters from end.
	852	do	do	do	do	4173	Crushed at 32 millimeters from end; opened between rings.
	852	do	do	do	do	4105	Crushed at 25 and at 127 millimeters from end.
330. <i>Chamaecyparis Nutkaensis</i> <i>Yellow Cypress. Sitka Cypress.</i>	909	Alaska	Sitka	Paul Schultze		5897	Triple diagonal flexure parallel to rings.
	909	do	do	do		7031	Crushed at 38 millimeters from end.
	983	British Columbia	Saw-mill, Victoria.	G. Engelmann and C. S. Sargent.		8210	Crushed at 64 millimeters from end.
	983	do	do	do		7779	Triple flexure; middle bend 38 millimeters eccentric.
	904	Alaska	Peril strait	Paul Schultze		7711	Crushed at 25 millimeters from end.
	994	do	do	do		7439	Crushed at 13 millimeters knot 38 millimeters from middle.
	1000	do	Weidler's saw-mill, Portland.	G. Engelmann and C. S. Sargent.		7217	Crushed at 31 millimeters from end.
	1000	do	do	do		6967	Crushed at 25 and at 102 millimeters from end.
331. <i>Chamaecyparis Lawsoniana</i> <i>Port Orford Cedar. Oregon Cedar. White Cedar. Lawson's Cypress. Ginger Pine.</i>	701	Oregon	Dean & Co.'s saw-mill, Marshfield.	do		7235	Crushed at 25 and at 51 millimeters from end; split along grain.
	707	do	do	do		7462	Crushed at end and at 102 millimeters from end.
	707	do	do	do		7666	Crushed at 51 millimeters from end.
332. <i>Cupressus macrocarpa</i> <i>Monterey Cypress.</i>	675	California	Monterey	do	Gravelly loam		
	675	do	do	do	do		
333. <i>Cupressus Goveniana</i>	691	do	Marin county	G. R. Vasey	Dry ridges	5693	Triple flexure
	691	do	do	do	do	7349	do
	1100	do	Calistoga	W. F. Fisher	do	8253	Crushed at middle; end shattered; split along grain.
	1100	do	do	do	do	4672	Split along grain; shattered
338. <i>Juniperus occidentalis</i> <i>Juniper.</i>	624	do	Yreka plains	G. Engelmann and C. S. Sargent.			
	624	do	do	do			
338. <i>Juniperus occidentalis, var. conjugens.</i> <i>Juniper.</i>	939	Texas	Austin	C. Mohr	Limestone	9049	Shattered stick at 127 millimeters from end and at end.
	939	do	do	do	do		
	1102	do	do	S. B. Buckley	do	9662	Triple diagonal flexure perpendicular to rings.
	1102	do	do	do	do	6804	Split obliquely; cross-grained and knots.
339. <i>Juniperus Virginiana</i> <i>Red Cedar. Savin.</i>	827	do	Dallas	J. Reverchon	Calcareous	7172	Crushed at 102 millimeters from end and split along grain.
	827	do	do	do	do	7439	Triple flexure; middle bend 32 millimeters eccentric.
	734	Florida	Chattahoochee	A. H. Curtiss	do	7915	Crushed at 25 and at 127 millimeters from end.
	734	do	do	do	do	7249	Crushed at 3 millimeters knot 25 millimeters from end.
	800	do	Saint John's river	do	Sandy loam	7031	Crushed at 10 millimeters knot 13 millimeters from end.
	800	do	do	do	do	5715	Opened grain at 6 millimeters knot near middle.
	837	Massachusetts	Danvers	J. Robinson	Drift	5126	Triple diagonal flexure parallel to rings.
	837	do	do	do	do	4944	do
	1055	do	Topshfield	do	Gravelly	5851	Crushed in vicinity of 13 millimeters knot at middle.
	1055	do	do	do	do	7076	Crushed at middle; deflected
	1249	Tennessee	Wilson county	A. E. Baird		8256	Crushed at knot at middle.
	1250	do	do	do		5879	Crushed at knot near end.
	1253	do	do	do		6577	Triple diagonal flexure perpendicular to rings; opened grain.

UNITED STATES UNDER COMPRESSION—Continued.

Direction of grain.	PRESSURE, IN KILOGRAMS, REQUIRED TO PRODUCE AN INDENTATION, IN MILLIMETERS, OF—												Remarks.	Office number.
	0.25	0.51	0.76	1.02	1.27	1.52	1.78	2.03	2.28	2.54	2.81	3.08		
	340	581	726	762	803	812	848	857	875	889	1007	1125	Sheared fibers	850
	640	984	1034	1075	1116	1166	1211	1234	1270	1270	1452	1574	do	850
	680	880	939	962	993	1030	1061	1084	1111	1120	1306	1452	do	850
	626	953	1139	1184	1243	1279	1293	1302	1315	1329	1429	1429	do	850
	403	735	862	871	880	894	903	921	944	962	1034	1089	do	851
	522	608	803	821	835	839	844	857	880	889	998	1080	do	851
	644	1025	1193	1315	1365	1415	1474	1506	1565	1610	1805	1973	Slight shearing of fibers	852
	544	975	1125	1311	1356	1411	1483	1551	1579	1642	1928	2132	do	852
	549	885	971	1012	1025	1057	1093	1120	1129	1157	1306	1406	Sheared fibers	900
	522	7.6	903	1048	1080	1120	1166	1175	1216	1234	1406	1520	Slight shearing of fibers	900
	721	1170	1302	1393	1488	1547	1588	1642	1724	1760	2028		Sheared fibers; split at end	903
	930	1592	1787	1842	1860	1919	1960	2009	2032	2087	2313		Slight shearing of fibers; split at end	903
	749	1247	1320	1393	1442	1492	1547	1592	1647	1687	1996		do	994
	1125	1442	1515	1597	1683	1833	1774	1819	1860	1896	2214		Slight shearing of fibers; split at end; 3 millimeters knot.	994
	1320	1837	1982	2073	2114	2164	2214	2259	2400	2331	2586	2790	Sheared fibers	1000
	1542	2032	2118	2205	2254	2400	2354	2404	2481	2513	2635	2994	do	1000
	866	1429	1520	1529	1579	1579	1651	1687	1728	1746	1928	2155	do	701
	690	953	1080	1120	1170	1211	1229	1275	1306	1329			Split along grain	707
	640	989	1116	1161	1202	1207	1266	1311	1325	1361			do	707
	1656	3311		3742	3860	3951	4051	4164	4187	4214	4717	4967	Sheared fibers	675
	1709	3266	3511	3620	3724	3797	3860	3901	4014	4078			Sheared fibers; split at ends	675
	1406	2540	3016	3311	3425	3515	3647	3729	3783	3842	4241	4527	Slight shearing of fibers; indented section covers two 3 millimeters knots.	691
	1805	2413	2558	2572	2619	2672	2703	2739	2803	2821			Slight shearing of fibers; split at corner	691
	594	1574	2118	2404	2563	2626	2681	2731	2799	2817	3166	3348	Sheared fibers	1100
	1728	2427	2558	2622	2771	2858	2944	3012	3094	3134	3606	3901	do	1100
	1348	2431	2925	3134	3202	3307	3393	3434	3484	3515	3882		Sheared fibers; split at ends; worm-eaten	624
	1610	2459	2622	2681	2758	2826	2899	2953	3016	3094	3574	4001	do	624
	1778	3593	4291	4653	4817	4944	5103	5158	5221	5294	5987	6305	Sheared fibers	939
	1474	3134	4046	4359	4513	4649	4854	4953	5080	5068			Sheared fibers; split at end	939
	1746	3402	4083	4264	4414	4626	4726	4795	4899	4962	5570		Sheared fibers; split at ends	1102
														1102
	1087	2014	2668	2186	2250	2304	2340	2395	2449	2504	2994	3202	Sheared fibers	827
	1207	1932	2132	2182	2214	2236	2250	2304	2341	2350	2586		Sheared fibers; split at end	827
	1587	2381	2699	2844	2890	2935	3030	3066	3094	3139	3438		do	734
														
	1384	2091	2291	2341	2363	2454	2486	2549	2590	2608	2989	3221	Sheared fibers	800
	1043	1700	1909	2028	2062	2118	2159	2173	2214	2236	2527	2672	do	800
	839	1438	1583	1705	1733	1765	1778	1796	1810	1824	1951	2074	do	837
	817	1279	1424	1511	1533	1574	1579	1628	1651	1665	1878	1996	do	837
	1001	1923	2245	2.50	2518	2540	2572	2617	2649	2681	3030		Sheared fibers; split at end	1056
														
	1588	2790	3066	3188	3334	3443	3565	3588	3674	3720	4264		Slight shearing of fibers; split at end	1248
	1202	2404	2694	2840	2965	3039	3153	3202	3293	3348	3901	4196	Sheared fibers	1250

FOREST TREES OF NORTH AMERICA.

TABLE V.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Longitudinal compression; Ultimate strength, in kilos.	Remarks.
340. <i>Taxodium distichum</i> . <i>Bald Cypress. Black Cypress. Red Cypress. White Cypress. Deciduous Cypress.</i>	535	Alabama	Stockton	C. Mohr	Alluvial	8029	Crushed at 64 millimeters from end.
	535	do	do	do	do	7031	Crushed at 51 millimeters from end.
	542	do	Mobile	do		6759	Crushed at 38 millimeters from middle.
	542	do	do	do		6759	Crushed at 25 millimeters from end.
	741	Florida	Chattahoochee	A. H. Curtiss		6328	do
	741	do	do	do		5697	Crushed at end.
341. <i>Sequoia gigantea</i> . <i>Big Tree.</i>	657	California	Tulare county	G. Engelmann and C. S. Sargent	Granite	6341	Crushed at middle
	657	do	do	do	do	6078	do
342. <i>Sequoia sempervirens</i> . <i>Redwood.</i>	673	do	Russian river	C. S. Sargent		6917	Crushed at 25 millimeters from middle.
	673	do	do	do		7394	Crushed at middle.
	710	do	Santa Cruz	Turner, Kennedy & Shaw		7122	Crushed at 76 millimeters from end.
	711	do	do	do		6523	Crushed at middle; shattered
	711	do	do	do		7979	Crushed at 64 millimeters from end; threw off two splinters.
	712	do	Mendocino county	J. Kentfield & Co.		5262	Crushed at 51 millimeters from end.
	712	do	do	do		5307	Crushed at 38 millimeters from middle.
	713	do	do	do		7122	do
	713	do	do	do		5942	Crushed at 25 millimeters from middle in vicinity of 2 millimeters knots.
	714	do	do	do		7349	Crushed at middle and at 38 millimeters from end; opened grain.
	715	do	do	do		6305	Crushed at 102 millimeters from end; split from end to end; grain curly.
343. <i>Taxus brevifolia</i> . <i>Yew.</i>	978	Oregon	Portland	G. Engelmann and C. S. Sargent	Moist, rich	6668	Failed at 19 millimeters knot at middle.
	978	do	do	do	do	8799	Crushed near middle at 3 millimeters knot.
345. <i>Torreya taxifolia</i> . <i>Stinking Cedar. Savin.</i>	62	Florida	Chattahoochee	C. S. Sargent	Alluvial	7258	Crushed at 6 millimeters knot 51 millimeters from middle; opened grain.
	62	do	do	do	do	7349	Crushed at 19 millimeters from end.
	277	do	do	A. H. Curtiss	Calcareous	7008	Triple diagonal flexure perpendicular to rings.
	277	do	do	do	do	7838	Triple diagonal flexure
346. <i>Torreya Californica</i> . <i>California Nutmeg. Stinking Cedar.</i>	651	California	Marin county	G. R. Vasey	Stony	5579	Crushed at 64 millimeters from middle at 5 millimeters knot.
	651	do	do	do	do	5670	Triple flexure
347. <i>Pinus Strobus</i> . <i>White Pine. Weymouth Pine.</i>	1	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	5398	Crushed at 51 millimeters from middle.
	1	do	do	do	do	5239	Crushed at 64 millimeters from end.
	222	Vermont	Charlotte	C. G. Pringle	Wet, swampy	6214	Crushed at 76 millimeters from end.
	777	New Brunswick		Intercolonial railway.		4427	Crushed at 10 millimeters knot 38 millimeters from middle.
	777	do		do		4210	Crushed at 10 millimeters knot 51 millimeters from end.
	788	do	Bridgeton	Ed. Sinclair		6169	Crushed at 51 end at 114 millimeters from end.
	788	do	do	do		6305	Crushed at 102 millimeters from end.
	789	Province of Quebec	Amqui	A. Grant		5806	Crushed at 13 millimeters from end.
	789	do	do	do		4095	Crushed at end
	797	do		Grand Trunk railway.		5842	Crushed at 64 millimeters from middle.
	797	do		do		5025	Crushed at end and at 32 millimeters from middle.
	1044	Massachusetts	Reading	J. Robinson	Drift	5534	Crushed at 45 millimeters from middle.
	1044	do	do	do	do	4967	Crushed at 83 millimeters from end.
348. <i>Pinus monticola</i> . <i>White Pine.</i>	975	British Columbia	Hastings' saw-mill, Burrard inlet.	G. Engelmann and C. S. Sargent		6123	Crushed at 76 millimeters from end.
	987	Oregon	Cascade mountains.	C. S. Sargent	Moist loam	4981	Triple flexure parallel to rings
	987	do	do	do	do	4944	Crushed at 13 millimeters from end.
349. <i>Pinus Lambertiana</i> . <i>Sugar Pine.</i>	638	California	Saw-mill, Strawberry valley.	G. Engelmann and C. S. Sargent		6441	Crushed at three places near ends.

UNITED STATES UNDER COMPRESSION—Continued.

Direction of grain.	PRESSURE, IN KILOGRAMS, REQUIRED TO PRODUCE AN INDENTATION, IN MILLIMETERS, OF—												Remarks.	Office number.
	0.25	0.51	0.76	1.02	1.27	1.52	1.78	2.03	2.28	2.54	4.81	5.08		
	494	785	886	916	939	1002	1066	1093	1139	1175	1334		Sheared fibers; split at end	536
	449	767	839	889	921	944	1007	1034	1052	1080	1247	1338	Slight shearing of fibers	536
	885	1424	1597	1678	1742	1787	1824	1864	1905	1941	2205		Slight shearing of fibers; split at end	542
	889	1220	1306	1352	1415	1483	1529	1551	1588	1619	1682	2087	Sheared fibers; split at corner	542
	463	1045	1325	1452	1483	1529	1569	1610	1619	1642	1796	1932	Sheared fibers	741
	880	1120	1202	1257	1320	1356	1415	1447	1483	1529	1792	1987	do.....	741
	839	953	980	1030	1084	1098	1134	1148	1161	1184	1320	1460	do.....	687
	862	963	971	1052	1098	1166	1225	1288	1329	1442	1524	1701	do.....	687
	563	852	921	953	984	1002	1025	1080	1034	1061	1179		Sheared fibers; split at end	719
	780	1012	1066	1075	1080	1084	1107	1157	1198	1221	1370	1452	Sheared fibers; split along grain	711
	626	762	860	839	948	998	1089	1035	1039	1048			Slight shearing of fibers; split along grain	711
	617	780	821	866	898	934	944	962	984	1007	1116		Sheared fibers; split along grain	713
	703	1048	1061	998	963	975	962	1084	1075	1098	1216	1306	Sheared fibers	713
	857	1306	1111	1116	1189	1325	1393	1442	1470	1501	1693	1792	do.....	713
	749	839	871	921	957	1002	1043	1080	1098	1130	1296		Sheared fibers; split at end	713
	1774	2440	2731	2853	2926	3003							Split at ends	715
	2223	3052	4037	4150	4296	4360	4436	4513	4581	4631	5103	5489	Sheared fibers	978
	2359	3665	3878	4024	4150	4255	4386	4473	4613	4726	5421	5851	do.....	978
	1656	2331	2613	2825	2971	3121	3266	3357	3488				Split at ends	62
	1021	2341	2840	3035	3125	3270	3365	3456	3515	3661	4287	4537	Sheared fibers; split at ends	62
	1257	1923	2118	2209	2254	2313	2345	2363	2386	2440	2713		Sheared fibers; split at end	277
	1043	1529	1633	1683	1742	1796	1851	1896	1941	1964	2245	2413	Sheared fibers	277
	1021	1452	1628	1710	1805	1842	1896	1941	2046	2087	2481	2676	Sheared fibers; split at end	651
	1071	1293	1941	2050	2118	2205	2250	2259	2304	2341	2596	2767	Sheared fibers	651
	953	1135	1211	1261	1297	1347	1370	1398	1429	1447	1588	1778	do.....	1
	640	1111	1243	1393	1447	1488	1511	1524	1538	1551	1724	1769	Sheared fibers; split at end	1
	098	1542	1883	1950	2087	2227	2304	2427	2454	2563	2971	3130	Slight shearing of fibers	222
	521	885	1048	1075	1111	1116	1120	1134	1152	1161	1270	1329	Sheared fibers	777
			953	962	980	1012	1021	1052	1068	1111	1270	1384	do.....	777
	671	962	1043	1080	1125	1166	1207	1220	1247	1297	1520	1660	do.....	798
	789	930	1030	1075	1120	1161	1207	1247	1270	1293	1424		Sheared fibers; split at end	798
	409	708	762	803	844	852	866	889	896	921	1043	1157	Sheared fibers	799
	526	780	885	912	944	957	980	993	1025	1039	1175	1270	do.....	799
	930	1075	1152	1207	1261	1320	1361	1411	1488	1520	1792	2023	do.....	797
	531	862	944	971	984	1025	1084	1096	1116	1148	1293	1397	do.....	797
	866	1216	1252	1302	1329	1347	1370	1384	1397	1420	1529	1601	do.....	1044
	500	903	962	980	1002	1048	1071	1075	1098	1116	1243	1361	do.....	1044
	630	953	1052	1093	1139	1189	1234	1275	1311	1347	1547		Sheared fibers; split at end	975
	712	871	962	1025	1075	1120	1161	1196	1184	1220	1424	1565	Sheared fibers	987
	602	857	925	975	998	1034	1075	1096	1120	1143	1293	1384	do.....	987
	535	1034	1216	1275	1311	1338	1356	1398	1424	1456	1610	1724	do.....	688

FOREST TREES OF NORTH AMERICA.

TABLE V.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Longitudinal compression: Ultimate strength, in kilos.	Remarks.
349. <i>Pinus Lambertiana</i> —cont'd	668	California		G. R. Vasey		5080	Crushed at 38 millimeters from end.
	780	do	Lassen's peak	Sierra Lumber Company.		4626	Triple flexure
350. <i>Pinus flexilis</i> . <i>White Pine.</i>	819	Colorado	Forest City	T. S. Brandegee	Gravelly	6123	Crushed at 25 millimeters from end.
	819	do	do	do	do	6123	Crushed at 76 millimeters from end.
	918	Nevada	Danville	A. Triple	do	4527	Crushed at 10 millimeters knot 76 millimeters from end.
351. <i>Pinus albicaulis</i>	992	British Columbia	Silver Mountain valley, Fraser river.	G. Engelmann and C. S. Sargent.		4740	Triple flexure
	992	do	do	do		5851	do
352. <i>Pinus reflexa</i> . <i>White Pine.</i>	661	Arizona	Santa Rita mountain.	G. Engelmann and C. S. Sargent.		8029	Crushed at 38 and at 89 millimeters from end; opened grain.
	661	do	do	do		7621	Crushed at 32 millimeters from end; split obliquely along grain.
353. <i>Pinus Parryana</i> . <i>Piñon. Nut Pine.</i>	656	California	San Diego county	G. R. Vasey		5262	Crushed at 64 millimeters from end at 16 millimeters knot.
	656	do	do	do		5579	Triple flexure.
355. <i>Pinus edulis</i> . <i>Piñon. Nut Pine.</i>	397	Colorado	Cañon City	E. Weston	Gravelly	5579	Crushed at end at 5 millimeters knot.
356. <i>Pinus monophylla</i> . <i>Piñon. Nut Pine.</i>	882	Utah	Lewiston	M. E. Jones	Rocky	4037	Split obliquely along grain.
	915	Nevada	Danville	A. Triple	Gravelly	4740	Failed at 13 millimeters knots at middle.
357. <i>Pinus Balfouriana</i>	681	California	Scott mountains	G. Engelmann and C. S. Sargent.	Rocky	4763	Failed at 19 millimeters knot 76 millimeters from end.
	681	do	do	do	do	6033	Crushed at 3 millimeters knots at middle.
357. <i>Pinus Balfouriana</i> , var. <i>aristata</i> . <i>Foothill Pine. Hickory Pine.</i>	821	Colorado	Forest City	T. S. Brandegee		5489	Crushed at 32 millimeters from end; cross-grained.
	821	do	do	do		5126	Crushed at end
	914	Nevada	Prospect mountain	A. Triple	Rocky	5012	Crushed at middle and at 3 millimeters knot 25 millimeters from middle.
358. <i>Pinus resinosa</i> . <i>Red Pine. Norway Pine.</i>	815	Michigan	Hersey	W. J. Beal		7756	Crushed at 51 millimeters from end.
	815	do	do	do		8301	Crushed at 64 millimeters from end.
	785	New Brunswick	Bridgeton	Ed. Sinclair		7167	Crushed at 10 millimeters knot 25 millimeters from end.
	785	do	do	do		8165	Crushed at end
	1074	Vermont	Charlotte	C. G. Pringle		6023	Crushed at 25 and at 114 millimeters from end.
	1075	do	do	do		5942	Crushed at 25 and at 76 millimeters from end.
	1076	do	do	do		7349	Crushed at 51 millimeters from end.
359. <i>Pinus Torreyana</i>	990	California	San Diego county	G. Engelmann	Sandy	4400	Crushed at 25 and at 102 millimeters from end.
	990	do	do	do	do	4676	Triple flexure; middle bend 51 millimeters eccentric.
	990	do	do	do	do		
360. <i>Pinus Arizona</i> . <i>Yellow Pine.</i>	1154	Arizona	Santa Rita mountain.	C. G. Pringle	Rocky	7485	Crushed at 25 millimeters from end.
	1154	do	do	do	do	7349	Crushed at end
	1155	do	do	do	do	5330	Shattered at end
	1155	do	do	do	do	6250	Crushed at 51 millimeters from end.
	1156	do	do	do	do	5060	Crushed at 25 millimeters knot at end.
	1156	do	do	do	do	4044	do
361. <i>Pinus ponderosa</i> . <i>Yellow Pine. Bull Pine.</i>	619	Dakota	Deadwood	Robert Douglas	Gravelly	7015	Crushed at 51 millimeters from end.
	626	Oregon	Saw-mill, Ashland	G. Engelmann and C. S. Sargent.		7530	Crushed at 76 millimeters from end.
	630	California	Strawberry valley	do	Low, wet, swampy	3629	Triple flexure perpendicular to rings; knots.
	630	do	do	do	do	3765	do
	632	do	Saw-mill, Strawberry valley.	do		8256	Triple flexure
	636	do	do	do		5942	Crushed at 51 millimeters from end.
	689	do	Saw-mill, San Bernardino.	W. G. Wright		7756	do

UNITED STATES UNDER COMPRESSION—Continued.

Direction of grain.	PRESSURE, IN KILOGRAMS, REQUIRED TO PRODUCE AN INDENTATION, IN MILLIMETERS, OF—												Remarks.	Offer number.
	0.25	0.51	0.76	1.02	1.27	1.52	1.78	2.03	2.28	2.54	2.81	3.08		
	1116	1311	1338	1406	1438	1483	1497	1524	1547	1579	1637	1682	Sheared fibers.....	688
	658	871	944	982	984	1021	1061	1084	1120	1139	1261	1234	do.....	796
	676	1302	1525	1619	1637	1696	1728	1760	1774	1796	1973	2032	do.....	819
	662	1270	1406	1483	1529	1574	1615	1642	1687	1724	1941	2078	do.....	819
	1098	1669	1824	1955	2014	2119	2173	2227	2400	2341	2607	2858	Sheared fibers; opened grain at end and along one face.	913
	866	1325	1406	1474	1501	1542	1583	1624	1665	1687	1982	2096	Sheared fibers.....	992
	1066	1642	1796	1892	1982	2023	2073	2114	2159	2209	2495	2735	do.....	992
	1261	1710	1905	1973	2032	2073	2123	2159					Split at end; indented section covers 9 millimeters knot.	661
	1134	1809	1982	2005	2053	2118	2182	2209	2223	2273	2376	2758	Sheared fibers.....	661
	2023	2676	3157	3243	3411	3520	3611	3679	3738	3833	4400	4854	Slight shearing of fibers.....	686
	1497	2386	2608	2735	2840	2926	2996	3085	3216	3266	3356	4264	Sheared fibers.....	656
	1905	2790	3030	3216	3388	3493	3674	3856	3992	4146	5013		Sheared fibers; split at end.....	897
	1792	2123	2345	2481	2540	2654	2717	2753	2821	2994			do.....	892
	958	2169	2505	2744	2885	3021	3125	3212	3311	3367	3892	3720	Sheared fibers.....	915
	871	2014	2459	2658	2713	2753	2880	2921	2953	2985	3339	3538	do.....	631
	1211	1811	1860	1923	1987	2023	2050	2091	2136	2168	2032	2676	do.....	631
	1442	2136	2286	2391	2477	2576	2645	2703	2771	2799	3311	3515	do.....	821
	1116	1506	1500	1619	1674	1705	1751	1801	1833	1842	2168	2381	do.....	831
	1397	2000	2105	2209	2268	2359	2431	2527	2622	2681	2948	3470	do.....	914
	671	1034	1157	1220	1288	1338	1379	1415	1442	1470	1678		Slight shearing of fibers; split at end.....	315
	862	1170	1275	1325	1365	1424	1483	1547	1574	1619	1860		do.....	315
	744	1116	1243	1433	1479	1497	1551	1584	1624	1665	1901	2023	Sheared fibers.....	785
	898	1406	1583	1647	1706	1724	1765	1787	1801	1837	2032	2168	do.....	785
	594	930	1025	1080	1134	1166	1207	1234	1270	1311	1497		Sheared fibers; split at end.....	1074
	703	989	1034	1111	1157	1202	1238	1266	1298	1329	1520	1633	Sheared fibers.....	1075
	953	1243	1379	1429	1461	1520	1569	1601	1633	1669	1923	2087	do.....	1076
	862	1025	1111	1170	1234	1288	1343	1370	1415	1452	1706		Sheared fibers; split at end.....	1076
	1270	2000	2177	2259	2327	2386	2431	2449	2499	2554	2948	3166	Sheared fibers.....	996
	1338	2486	2165	2223	2291	2350	2400	2477	2572	2595	3016	3261	do.....	996
	1452	2132	2291	2359	2440	2486	2527	2495	2518	2536			Sheared fibers; split at end.....	996
	1021	1987	1710	1796	1864	1946	1991	2041	2118	2141	1973		do.....	1154
	1034	1669	1837	1892	1941	1991	2028	2073	2100	2576	2835	2926	Sheared fibers.....	1154
	1089	1365	1429	1488	1524	1500	1615	1660	1692	1719	1990	2205	do.....	1155
	1016	1574	1687	1724	1774	1805	1833	1882	1896	1905			Sheared fibers; split at end.....	1155
	980	1397	1524	1579	1597	1601	1610	1637	1651	1656	1778	1892	Sheared fibers.....	1156
	907	1361	1388	1406	1429	1456	1497	1524	1538	1565	1805	1919	do.....	1156
	680	971	1093	1161	1207	1257	1302	1338	1361	1402	1610		Sheared fibers; split at end.....	619
	1157	1724	1851	1879	1905	1978	2046	2082	2118	2155	2336	2540	Sheared fibers.....	626
	1452	2406	2468	2518	2608	2667	2749	3803	2853	2903	3266	3438	do.....	630
	698	1098	1765	1833	1851	1888	1932	1946	1969	1982	2214	2381	do.....	630
	1134	1343	1456	1533	1601	1669	1733	1787	1851	1901	2177		Slight shearing of fibers; split at end.....	632
	1121	1179	1189	1275	1325	1343	1365	1411	1438	1529	1700	1796	Sheared fibers.....	636
	998	1429	1542	1624	1687	1700	1796	1846	1890	1932	2200		do.....	636

TABLE V.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Longitudinal compression: Ultimate strength, in kilos.	Remarks.
361. <i>Pinus ponderosa</i> —continued.	718	Montana	Saw-mill, Missoula	S. Watson		6463	Crushed at end
	718	do	do	do		6477	Crushed at 51 millimeters from end; 13 millimeters knot at end.
	731	California	Lassen's peak	Sierra Lumber Company.		7349	Crushed at middle
	907	Colorado	Cañon City	E. Weston		3892	Triple diagonal flexure; knots near middle.
	910	do	do	do		4173	Triple diagonal flexure; 6 millimeters knot at middle bend.
362. <i>Pinus jeffreyi</i> <i>Bull Pine. Black Pine.</i>	633	California	Scott mountains	G. Engelmann and C. S. Sargent.	Dry, gravelly	5443	Crushed at middle at 6 millimeters knot.
	633	do	do	do	do	6577	Triple diagonal flexure parallel to rings; middle bend 6 millimeters eccentric.
	667	do	Saw-mill, San Bernardino.	W. G. Wright		6750	Oblique split 178 millimeters long, separating stick.
	667	do	do	do		7938	Crushed at 51 millimeters from end.
363. <i>Pinus Chihuahuana</i>	664	Arizona	Santa Rita mountains.	G. Engelmann and C. S. Sargent.	Dry, gravelly	5035	Triple diagonal flexure
	664	do	do	do	do	5761	Shattered end; crushed at 102 millimeters from end.
364. <i>Pinus contorta</i> <i>Scrub Pine.</i>	997	British Columbia	Vancouver's island	do	do	7802	Crushed at 102 millimeters from end.
	997	do	do	do	do	9934	Crushed at 51 millimeters from end.
365. <i>Pinus Murrayana</i> <i>Tamarack. Black Pine. Lodge-pole Pine. Spruce Pine.</i>	293	Colorado	Forest City	T. S. Brandegee	Moist, sandy loam	4626	Crushed at 38 millimeters from end.
	293	do	do	do	do	4037	Crushed at 10 millimeters knot 19 millimeters from end.
	563	do	do	C. S. Sargent.	do	6214	Crushed at 89 millimeters from end.
	625	California	Scott mountains	G. Engelmann and C. S. Sargent.	do	5648	Crushed at 10 millimeters knot 102 millimeters from end.
	625	do	do	do	do	6115	Crushed at 5 millimeters knot at middle.
366. <i>Pinus Sabiniana</i> <i>Digger Pine. Bull Pine.</i>	644	do	Contra Costa county.	G. R. Vasey	Gravelly	5262	Triple flexure; middle bend 25 millimeters eccentric.
	644	do	do	do	do	5511	Failed at 13 millimeters knot at middle.
367. <i>Pinus Coulteri</i>	1157	do	San Bernardino	W. G. Wright	Dry, gravelly	4808	Crushed at 16 millimeters knot 38 millimeters from middle.
	1157	do	do	do	do	6940	Crushed at 76 millimeters from end.
368. <i>Pinus insignis</i> <i>Monterey Pine.</i>	676	do	Monterey	G. R. Vasey	Gravelly loam	6827	Crushed at 38 millimeters from end at 5 millimeters knot.
	676	do	do	do	do	6532	Crushed at 51 to 71 millimeters from end.
369. <i>Pinus tuberculata</i> <i>Knob-cone Pine.</i>	576	do	Mount Shasta	G. Engelmann and C. S. Sargent.	Gravelly	4672	Crushed at 102 millimeters from end.
	576	do	do	do	do	3742	Crushed at knots at middle and near end.
370. <i>Pinus Troda</i> <i>Loblolly Pine. Old-field Pine. Rosemary Pine.</i>	82	Florida	Duval county	A. H. Curtiss	Moist, sandy	9154	Crushed at 76 millimeters from end.
	82	do	do	do	do	9185	Crushed at 51 millimeters from end.
	355	Alabama	Cottage Hill	C. Mohr	Low, rich	4445	Triple flexure
	355	do	do	do	do	4876	do
	388	North Carolina	Wilmington	E. Kidder	Loam	7194	Crushed at 76 millimeters from end.
	388	do	do	do	do	8437	Crushed at 38 millimeters from end in vicinity of knots.
371. <i>Pinus rigida</i> <i>Pitch Pine.</i>	389	do	do	do	do	5398	Crushed at middle in vicinity of 5 millimeters knot.
	389	do	do	do	do	5783	Crushed at middle in vicinity of knot.
	13	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	6123	Crushed at 102 millimeters from end.
	13	do	do	do	do	7076	Triple flexure parallel to rings
	1046	do	North Reading	J. Robinson	do	4808	Crushed at 64 millimeters from end.
	1046	do	do	do	do	4740	Triple flexure
372. <i>Pinus serotina</i> <i>Pond Pine.</i>	83	Florida	Duval county	A. H. Curtiss	Moist, sandy loam	7038	Triple flexure perpendicular to rings.
	83	do	do	do	do	8219	Triple flexure parallel to rings
373. <i>Pinus inops</i> <i>Jersey Pine. Scrub Pine.</i>	621	South Carolina	Aiken	H. W. Ravenel	Dry, sandy	5080	Triple flexure, contained 7 knots 3 to 6 millimeters in diameter.
	622	do	do	do	do	6577	Crushed in vicinity of 3 millimeters knot 51 millimeters from end.
	1169	Indiana	New Albany	M. J. Robinson		5829	Crushed at end at 13 millimeters knot.
	1169	do	do	do		5489	Crushed at 13 millimeters knot 76 millimeters from end.

UNITED STATES UNDER COMPRESSION—Continued.

Direction of grain.	PRESSURE, IN KILOGRAMS, REQUIRED TO PRODUCE AN INDENTATION, IN MILLIMETERS, OF—												Remarks.	Office number.
	0.35	0.51	0.76	1.02	1.27	1.52	1.78	2.03	2.28	2.54	2.81	3.08		
	1021	1166	1211	1261	1297	1311	1347	1361	1397	1438	1678	1878	Sheared fibers.....	718
	857	1225	1302	1325	1365	1402	1452	1479	1542	1579	1769	1928	do.....	718
	1202	1769	1905	1982	2023	2068	2114	2155	2186	2218	2540	2685	do.....	731
	885	1538	1769	1819	1860	1928	2000	2096	2123	2182	2495	2694	do.....	907
	685	1492	1728	1833	1901	1987	2028	2091	2141	2182	2136	2849	do.....	910
	1261	1879	2023	2173	2259	2309	2345	2368	2418	2431	2099		Sheared fibers; split at end.....	638
	1579	1769	1882	1946	1969	2014	2064	2082	2118	2150			do.....	638
	862	1315	1379	1479	1547	1606	1651	1687	1751	1773	2064	2223	do.....	667
	771	1293	1483	1569	1624	1665	1715	1755	1796	1842	2087	2277	Sheared fibers.....	667
	249	953	1769	1937	2000	2105	2168	2223	2286	2354	2812		Sheared fibers; split at ends.....	664
	1021	2177	2676	2853	2939	3013	3071	3116	3175	3293	3765		Slight shearing of fibers; split at ends.....	604
	1565	1787	1905	2000	2078	2164	2250	2309	2372	2440			Slight shearing of fibers; split at end.....	907
	1202	1996	2341	2586	2685	2830	2894	2994	3075	3166			Sheared fibers; split at end.....	907
	885	1071	1202	1229	1252	1270	1279	1288	1293	1311	1497	1699	Sheared fibers.....	298
	667	1061	1143	1170	1198	1220	1243	1252	1275	1288	1497	1579	do.....	298
	1270	1374	1501	1588	1665	1715	1774	1846	1910	1964	2223	2404	do.....	563
	898	1266	1438	1492	1529	1565	1579	1601	1647	1665	1814		Sheared fibers; split at end.....	625
	867	1052	1152	1193	1252	1302	1329	1374	1406	1433	1610	1787	Sheared fibers.....	625
	1202	1778	1910	1982	2041	2087	2168	2206	2259	2313	2667	2849	do.....	644
	1479	2014	2214	2332	2363	2413	2499	2545	2599	2654	3107		Slight shearing of fibers; split at end.....	644
	1689	1225	1352	1397	1470	1533	1574	1615	1651	1687	2023	2214	Sheared fibers.....	1157
	753	1247	1374	1424	1479	1547	1574	1619	1669	1715	1973	2186	do.....	1157
	817	1496	1569	1624	1665	1701	1715	1737	1746	1769	1969	2068	do.....	676
	953	1456	1619	1674	1710	1746	1778	1819	1883	1869	2118		Sheared fibers; split at end.....	676
	599	1093	1306	1379	1424	1456	1489	1524	1588	1569	1769	1905	Sheared fibers.....	576
	767	1166	1261	1297	1320	1347	1379	1397	1429	1470		1746	do.....	576
		1179	1347	1393	1456	1520	1574	1637	1656	1699			Split at ends; fibers did not shear.....	82
	1207	1982	2218	2373	2496	2518	2563	2585	2672	2713	2994		Sheared fibers; split at ends.....	82
	1157	1447	1529	1592	1666	1706	1787	1796	1819	1855			Split at end; fibers did not shear.....	355
	1243	1764	1919	2050	2118	2223	2268	2322	2368	2418	2756	2953	Indented without shearing of fibers.....	355
	880	1093	1161	1216	1275	1347	1442	1497	1547	1597	1905		Sheared fibers; split at ends.....	368
	862	1120	1198	1275	1325	1406	1456	1547	1597	1637	1996		Split at end; fibers did not shear.....	368
	1247	1483	1497	1560	1683	1674	1701	1733	1774	1796	2019	2182	Sheared fibers.....	389
	925	1520	1678	1801	1855	1901	1928	1973	2005	2037	2223	2336	do.....	389
	1828	2223	2295	2341	2449	2531	2595	2658	2740	2844	3289	3526	do.....	13
	1542	2518	2844	3089	3207	3302	3357	3488	3579	3692	4219	4491	do.....	13
	939	1270	1343	1374	1393	1402	1433	1452	1492	1520	1678	1796	do.....	1046
	703	1179	1270	1424	1442	1483	1501	1529	1531	1565	1787	1883	do.....	1046
	2654	3479	3751	3901	4019	4150	4250	4359	4409	4522	5216	5851	do.....	83
	2658	4255	4854	5289	5461	5679	5729	5825	5834	5852	5761		Sheared fibers; split along grain.....	83
	1520	2132	2318	2395	2477	2558	2608	2658	2703	2749	3030	3202	Sheared fibers.....	621
	1043	1678	2023	2127	2186	2241	2277	2309	2341	2368	2613		Sheared fibers; split at end.....	622
	1211	1796	2032	2096	2164	2195	2236	2250	2288	2286	2572	2635	Sheared fibers.....	1169
	1116	2073	2350	2431	2504	2527	2599	2649	2690	2731			Slight shearing of fibers; split at end.....	1169

TABLE V.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Longitudinal compression: Ultimate strength, in kilob.	Remarks.
873. <i>Pinus inops</i> —continued	1172	New Jersey	Mount Holly	S. P. Sharples	Clay	6033	Crushed at middle on one face
	1172	do	do	do	do	5579	Triple diagonal flexure
874. <i>Pinus clausa</i>	279	Florida	Apalachicola	A. H. Curtiss	Dry, sandy barrens	5715	Triple flexure parallel to rings; knots at middle.
	279	do	do	do	do	6341	Crushed at knots at middle
875. <i>Pinus pungens</i>	321	Virginia	Wytheville	H. Shriver	Clay	5987	Triple flexure perpendicular to rings.
	321	do	do	do	do	5353	Crushed at 89 millimeters from end; split at end.
876. <i>Pinus muricata</i>	671	California	Marin county	G. R. Vasey	Gravelly	8663	Crushed at 114 millimeters from end at 3 millimeters knot.
	671	do	do	do	do	7621	Crushed at 102 millimeters from end.
877. <i>Pinus mitis</i>	278	Florida	Chattahoochee	A. H. Curtiss	Clay	4763	do
	278	do	do	do	do	7963	Crushed at small 2 millimeters knots 25 millimeters from middle.
	319	Louisiana	Amite	C. Mohr	Sandy loam	10138	Crushed at 6 millimeters knot 25 millimeters from middle.
878. <i>Pinus glabra</i>	142	South Carolina	Bonneau's Depot	H. W. Ravenel	Rich upland	5466	Triple flexure
	142	do	do	do	do	5398	Crushed at 51 millimeters from end.
	544	Mississippi	Gainesville	C. Mohr	Low, sandy	3130	Crushed at 76 millimeters from end at 16 millimeters knot.
	544	do	do	do	do	2903	Triple flexure
	764	Florida	Chattahoochee	A. H. Curtiss	Low, wet	5353	Triple flexure parallel to rings.
	764	do	do	do	do	5375	Crushed 32 millimeters from middle in vicinity of knot.
879. <i>Pinus Banksiana</i>	394	Michigan	Baldwin	W. J. Beal	Low, sandy	5104	Crushed at end at 13 millimeters knot.
	394	do	do	do	do	4944	Triple flexure; opened grain
	780	New Brunswick	Intercolonial railway.	do	do	5080	Crushed at 6 millimeters knot 76 millimeters from end.
	780	do	do	do	do	7756	Crushed and opened between rings at end.
	879	do	do	do	do	8188	Crushed at 89 millimeters from end.
	879	do	do	do	do	6813	Crushed at 3 millimeters knot at middle.
880. <i>Pinus palustris</i>	81	Florida	Duval county	A. H. Curtiss	Sandy loam	8392	Split obliquely; stick broke in two pieces.
	81	do	do	do	do	10977	Crushed at 13 to 76 millimeters from end.
	85	do	do	do	Moist, sandy	8618	Triple flexure parallel to rings
	85	do	do	do	do	9321	Crushed at 13 millimeters from end.
	243	do	Saw-mill, Saint John's river.	do	do	11204	Crushed at 19 millimeters from end.
	243	do	do	do	do	10160	Crushed at middle
	302	South Carolina	Aiken	H. W. Ravenel	Dry, sandy	7847	Crushed near middle
	302	do	do	do	do	8165	Crushed at 51 millimeters from end.
	357	Alabama	Cottage Hill	C. Mohr	Sandy loam	10478	Crushed at 38 millimeters from end.
	357	do	do	do	do	10047	Crushed at 38 millimeters from middle.
	358	do	Citronelle	do	do	10569	Crushed at end
	358	do	do	do	do	11839	Triple flexure
	359	do	Chunchula	do	do	11930	Crushed and splintered at end
	359	do	do	do	do	10931	Crushed at middle
	360	do	do	do	do	11068	Crushed at 32 millimeters from end.
	360	do	do	do	do	10931	Crushed at 64 millimeters from middle.
	361	do	do	do	do	10660	Crushed at 38 millimeters from end.
	361	do	do	do	do	10931	Crushed at 76 millimeters from end.
	384	Florida	Saw-mill, Cedar Keys.	A. H. Curtiss	do	9775	Crushed at 76 millimeters from end and shattered.
	384	do	do	do	do	9034	Crushed at 51 millimeters from end.
	385	do	do	do	do	8663	Crushed at 38 millimeters from end.
	385	do	do	do	do	9017	Crushed at 89 millimeters from end.

UNITED STATES UNDER COMPRESSION—Continued.

Direction of grain.	PRESSURE, IN KILOGRAMS, REQUIRED TO PRODUCE AN INDENTATION, IN MILLIMETERS, OF—												Remarks.	Office number.
	0.25	0.51	0.76	1.02	1.27	1.52	1.78	2.03	2.28	2.54	2.81	3.07		
	1247	2404	2812	3075	3248	3343	3411	3456					Split at end	1173
	1884	2062	2254	2304	2400	2440	2513	2572	2604	2640	2694	3198	Sheared fibers	1173
	1243	1941	2100	2218	2400	2377	2436	2477	2536	2576	2680		Sheared fibers; split at ends	279
	1134	1429	1656	1715	1801	1814	1846	1860	1869	1923	2132	2359	Sheared fibers	279
	1252	1687	1883	1910	1951	2082	2118	2191	2250	2331	2767		Sheared fibers; split along grain	321
	953	1452	1615	1705	1733	1756	1787	1801	1824	1910	2291	2495	Sheared fibers	321
	1429	1901	2050	2136	2232	2296	2345	2386	2431	2490	2767		Sheared fibers; split at end	671
	862	1325	1515	1601	1669	1756	1787	1842	1901	1946	2304		do	671
	1080	1429	1551	1624	1706	1756	1810	1837	1951	2009	2341		Slight shearing of fibers; split at end	278
	1398	2177	2481	2613	2676	2762	2781	2790	2785	2790	3139		Sheared fibers	278
	1157	1538	1624	1724	1810	1864	1928	1987	2037	2096			Slight shearing of fibers; split at end	319
	1043	1760	1941	2028	2096	2154	2209	2250	2295	2345	2699	2917	Sheared fibers	142
	1030	1470	1506	1565	1606	1619	1647	1665	1687	1701	1800	1951	do	142
	962	1198	1311	1335	1415	1442	1479	1506	1533	1565	1746	1879	do	544
	690	1148	1284	1338	1393	1429	1438	1470	1483	1515	1633	1851	do	544
	1043	1515	1592	1669	1728	1760	1810	1851	1901	1941	2186	2336	do	764
	1370	1674	1787	1864	1928	1982	2005	2014	2068	2096	2404	2785	do	764
	707	1247	1393	1470	1529	1569	1615	1660	1683	1715	1996	2141	do	394
	544	1452	1633	1778	1851	1978	2019	2064	2091	2146	2454	2653	do	394
	1134	1442	1501	1568	1637	1710	1778	1814	1882	1923	2336	2585	Sheared fibers; 3 millimeters knot in indented section	780
	807	1315	1520	1628	1687	1719	1760	1810	1842	1855	2032		Sheared fibers; split at end	780
	857	1306	1406	1415	1424	1438	1524	1597	1628	1669	1769	1951	Sheared fibers	879
	1071	1293	1361	1443	1529	1574	1642	1687	1765	1792	2064		Sheared fibers; split at end	879
	1760	2268	2499	2622	2681	2799	2885	2948	3021	3062	3221	3866	Sheared fibers	81
	1388	1860	1996	2087	2168	2259	2313	2386	2440	2486			Slight shearing of fibers; split at end	81
	1442	1624	1728	1778	1873	1941	1991	2050					Split at end; fibers did not shear	85
	996	1615	1778	1860	1919	1987	2046	2150	2236	2286	2699		Sheared fibers; split at end	85
	1588	1960	2064	2109	2191	2273	2322	2363	2409	2440			Slight shearing of fibers; split at end	243
	1951	2803	3012	2994	2812	2880	3075	3193	3257	3302	3946		Sheared fibers; split at end	243
	1479	1882	1882	1906	2023	2259	2522	2622	2703	2903			Sheared fibers; opened seasoning cracks	302
	1125	1452	1515	1637	1710	1792	1864	2431	2486	2581	2971		Slight shearing of fibers; split at end	302
	1134	1442	1569	1746	1796	1892	1973	2041	2114	2168			do	357
	1760	2395	2558	2622	2635	2649	2676	2903	3025	3134			Sheared fibers; split at end	357
	1406	1683	1833	1941	2087	2127	2182	2254	2295	2350			Split at end; fibers did not shear	358
	1860	2767	2971	3039	3139	3289	3388	3484	3529	3633	4150		Slight shearing of fibers; split at end	358
	1951	2296	2495	2640	2808	2890	3071	3198	3270				Split at end; fibers did not shear	359
	2223	3202	3447	3484	3502	3543	3683	3756	3842	3896			Split at end; slight shearing of fibers	359
	1531	1837	2037	2214	2422	2540	2676	2758	2885	2998			do	360
	2041	2921	2971	2812	3153	2817	2904	3193	3316	3452			Sheared fibers; split at end	360
	1883	3048	3438	3661	3751	3887	3878	3933	4014	4101	4808	5216	Sheared fibers	361
	1610	1900	1996	2136	2270	2341	2422	2486	2581	2654	3030		Slight shearing of fibers; split at end	361
	1406	1542	1706	1778	1914	2000	2082	2132	2177	2263	2740		do	384
	1452	2540	2885	3026	3116	3106	3216	3298	3357	3411			Sheared fibers; split at end	384
	1270	1905	2078	2218	2295	2359	2436	2522	2572	2635	3107		Slight shearing of fibers; split at ends	385
	1311	1810	2023	2155	2259	2300	2359	2440	2495	2595	2813		do	385

TABLE V.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Longitudinal compression: Ultimate strength, in kilos.	Remarks.
380. <i>Pinus palustris</i> —continued...	390	North Carolina	Wilmington	E. Kidder		10206	Crushed at middle and at 25 millimeters from middle.
	390	do	do	do		9957	Crushed at 51 millimeters from end.
381. <i>Pinus Cubensis</i> <i>Slash Pine. Swamp Pine.</i> <i>Bastard Pine. Meadow Pine.</i>	84	Florida	Duval county	A. H. Curtiss	Moist, sandy	9063	Crushed at 51 millimeters from end and at end.
	84	do	do	do	do	8637	Triple flexure.
	356	Alabama	Cottage Hill	C. Mohr	do	12792	Crushed at 32 and at 102 millimeters from end.
	356	do	do	do	do	13585	Crushed at 76 millimeters from end.
	493	Florida	Bay Biscayne	A. H. Curtiss	Coral	8550	Split obliquely along grain; seasoning cracks.
	493	do	do	do	do	11136	Crushed at 25 millimeters from middle.
382. <i>Picea nigra</i> <i>Black Spruce.</i>	231	Vermont	Charlotte	C. G. Pringle	Cold, peacy	8210	Crushed at 3 millimeters knot 76 millimeters from end.
	231	do	do	do	do	6668	Crushed at 3 millimeters knots at middle and at end.
	373	do	Huntington	do	Gravelly	6242	Crushed at 83 millimeters from end.
	776	New Brunswick	Bay of Fundy	Intercolonial railway.		6305	Crushed at 6 millimeters knot 51 millimeters from middle.
	776	do	do	do		6305	Crushed at middle at 6 millimeters knot.
	794	Province of Quebec	Danville	Grand Trunk railway.		5851	Triple flexure; middle bend 25 millimeters eccentric.
	794	do	do	do		6373	Crushed at 25 millimeters from end.
	880	New Brunswick	Bridgeton	Ed. Sinclair		6464	Crushed at 13 millimeters from middle.
	880	do	do	do		6260	Crushed at middle on one face
383. <i>Picea alba</i> <i>White Spruce.</i>	513	New Hampshire	Stratford	C. G. Pringle	Gravelly	4672	Crushed at 5 millimeters knot 51 millimeters from end.
	513	do	do	do	do	5117	Crushed at 13 millimeters knot at middle.
	620	Dakota	Terry's peak	R. Douglas	do	5330	Crushed at 51 millimeters from middle.
	620	do	do	do	do	5443	Crushed at 3 millimeters knot 51 from end.
	773	New Brunswick	Bay of Fundy	Intercolonial railway.		5715	Crushed at 16 millimeters knot 25 millimeters from middle.
	773	do	do	do		4518	Crushed at 16 millimeters knot 102 millimeters from end.
	784	do	Bridgeton	Ed. Sinclair		5987	Crushed at 45 millimeters from middle.
	784	do	do	do		5978	Crushed at 51 millimeters from end.
	791	Province of Quebec	Amqui	A. Gault		5670	Crushed at 76 millimeters from end.
384. <i>Picea Engelmanni</i> <i>White Spruce.</i>	292	Colorado	Forest City	T. S. Brandegee	Damp	4037	Triple flexure parallel to rings.
	292	do	do	do	do	3883	Crushed at 10 millimeters knot 89 millimeters from end.
	575	do	do	C. S. Sargent	Peaty	5262	Triple flexure.
	822	do	do	T. S. Brandegee	do	3901	Crushed at 102 millimeters from end; cross-grained
385. <i>Picea pungens</i> <i>White Spruce. Blue Spruce.</i>	269	do	Alpine	do	Damp	4046	Crushed at 102 millimeters from end at 10 millimeters knot.
	270	do	do	do	do	3792	Crushed at 16 millimeters knot 76 millimeters from end.
	270	do	do	do	do	4545	Crushed at 38 millimeters from end and at middle at 3 millimeters knots.
386. <i>Picea Sitchensis</i> <i>Tide-land Spruce.</i>	970	Alaska	Sitka	Paul Schultze		6568	Triple flexure parallel to rings.
	970	do	do	do		6146	Triple flexure perpendicular to rings.
	977	British Columbia	Saw-mill, Burrard inlet.	G. Engelmann and C. S. Sargent.		5715	Crushed at 64 millimeters from end.
	977	do	do	do		5579	Crushed at 76 millimeters from end.
	1015	Oregon	Weidler's saw-mill, Portland	do		6486	Crushed at middle; angle of crushing, 85°.
	1019	do	Saw-mill, Astoria	do		5216	Crushed at 76 millimeters from end; angle of crushing, 85°.
	1019	do	do	do		5625	Crushed at 76 millimeters from end.
	1026	do	Portland Furniture Company.	do		5017	Crushed at 13 and at 102 millimeters from end.
387. <i>Tsuga Canadensis</i> <i>Hemlock.</i>	1026	do	do	do		4527	Triple flexure.
	5	Massachusetts	Arnold Arboretum	C. S. Sargent	Drift	5987	Triple flexure parallel to rings.
	5	do	do	do	do	5706	Crushed at 51 millimeters from end.
	219	Vermont	Charlotte	C. G. Pringle	Gravelly	7702	Triple flexure.

UNITED STATES UNDER COMPRESSION—Continued.

Direction of grain.	PRESSURE, IN KILOGRAMS, REQUIRED TO PRODUCE AN INDENTATION, IN MILLIMETERS, OF—												Remarks.	Office number.
	0.25	0.51	0.76	1.02	1.27	1.52	1.78	2.03	2.28	2.54	2.81	3.08		
	1315	1474	1637	1814	1887	1973	2155	2223	2277	2350			Slight shearing of fibers; split at ends	390
	1542	1941	1987	2141	2313	2431	2563	2654	2785	2944			Sheared fibers; split at ends	390
	1111	1406	1515	1560	1592	1687	1710	1737	1796	1824			Slight shearing of fibers; split at ends	84
	1293	1769	1846	1892	1932	1982	2078	2159	2277	2331	2785		do	84
	1229	1452	1538	1706	1796	1951	2041	2123	2236	2304			do	356
	2313	3393	3783	3983	4078	4237	4264	4309	4418	4527			Sheared fibers; split at ends	356
	2880	3892	4400	4786	5080	5252	5579	5761	5851	6214			Sheared fibers; split at end; indented section covers 10 millimeters knot.	493
	1637	2722	3130	3298	3438	3456	3710	3801	3687	3901	4146	4246	Sheared fibers.	493
	826	1261	1406	1442	1429	1520	1569	1606	1628	1660	1851		Sheared fibers; split at ends	231
	925	1225	1488	1542	1558	1637	1678	1719	1778	1810			Slight shearing of fibers; split at ends	231
	590	807	944	989	1066	1120	1166	1202	1234	1257	1452	1619	do	373
	644	1089	1225	1261	1279	1302	1325	1347	1370	1397	1560	1724	Sheared fibers	776
	749	1089	1238	1302	1352	1379	1402	1438	1479	1511	1742		Sheared fibers; split at end	776
	725	962	1030	1071	1102	1125	1170	1211	1225	1247	1429	1506	Sheared fibers	704
	671	1007	1061	1071	1098	1129	1161	1202	1220	1243	1388	1506	do	794
	635	962	1039	1080	1116	1143	1166	1202	1225	1252	1370	1488	do	880
	635	989	1034	1093	1125	1161	1193	1207	1238	1275	1488	1579	do	880
	776	1061	1152	1179	1202	1225	1261	1293	1315	1334	1588		do	513
	735	1120	1257	1325	1388	1420	1438	1456	1506	1542	1678	1824	Slight shearing of fibers	513
	726	1139	1311	1397	1447	1488	1524	1542	1565	1579	1724		Sheared fibers; 3 millimeters knot covered by indented section.	620
	676	1084	1229	1343	1398	1438	1498	1533	1574	1687	1987	3132	Sheared fibers	620
	568	798	880	907	957	1012	1043	1080	1116	1152	1325		Slight shearing of fibers; split at end	773
	631	939	1061	1098	1125	1143	1175	1220	1266	1302	1438	1569	Sheared fibers	773
	662	817	894	989	1007	1030	1080	1096	1111	1124	1379		Sheared fibers; split at end	784
	644	916	962	966	968	1039	1061	1071	1098	1125	1261	1379	Sheared fibers	784
	817	966	1021	1071	1125	1170	1220	1270	1320	1347	1533		Sheared fibers; split at end	791
	640	984	1043	1071	1139	1184	1229	1261	1302	1325	1492	1637	Sheared fibers	791
	567	1080	1170	1284	1275	1325	1361	1393	1415	1447	1579	1733	do	293
	953	1379	1515	1597	1665	1710	1751	1765	1778	1814			Sheared fibers; 10 millimeters knot covered by indented section.	293
	572	835	844	857	862	866	896	944	948	953	1043	1139	Sheared fibers	575
	590	903	984	1025	1066	1098	1120	1152	1175	1207	1320		Sheared fibers; split at end	822
	907	1279	1402	1461	1524	1565	1619	1656	1699	1683	1832	2009	Sheared fibers	269
	640	912	948	962	984	1016	1084	1052	1066	1098	1275	1370	do	270 ¹
	1016	1148	1234	1257	1298	1334	1379	1397	1443	1470	1637	1778	do	270 ²
	889	1184	1302	1415	1479	1529	1606	1660	1723	1760	2087		Sheared fibers; split at end	970
	717	1325	1574	1628	1706	1765	1829	1892	1969	2068	2395	2631	Sheared fibers	970
	626	749	807	871	944	966	993	1068	1161	1252			Slight shearing of fibers; split at end	977
	758	1148	1257	1320	1347	1388	1406	1424	1433	1456	1715		Sheared fibers	977
	458	626	690	749	789	830	866	903	939	966			Split at end; fibers did not shear	1015
	721	794	848	898	957	1021	1048	1120	1139	1161	1433		Slight shearing of fibers; split at end	1019
	726	907	925	934	971	1016	1048	1080	1116	1161	1352		Sheared fibers; split at end	1019
	866	1043	1120	1170	1198	1220	1279	1325	1356	1415	1633		Sheared fibers	1026
	631	907	980	1025	1052	1071	1111	1125	1152	1166	1302	1397	do	1026
	871	1007	1084	1198	1275	1297	1306	1352					Slight shearing of fibers; split at end	5
	771	817	848	894	921	1002	1107	1170	1216	1257	1542	1732	Sheared fibers	5
	568	771	866	963	993	1016	1052	1129	1120	1161	1311		Slight shearing of fibers; split at end	219

TABLE V.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Longitudinal compression: Ultimate strength, in kilos.	Remarks.
287. <i>Tsuga Canadensis</i> —continued.	219	Vermont	Charlotte	C. G. Pringle	Gravelly	7892	Crushed at 51 millimeters from middle.
	772	New Brunswick		Intercolonial railway.		4649	Crushed at middle at 3 millimeters knot.
	772	do		do		5307	Triple flexure perpendicular to rings.
	775	do	Bay of Fundy	do		5489	Crushed at 45 millimeters from middle.
	775	do	do	do		5738	Triple flexure; middle bend 51 millimeters eccentric.
	778	do		do		5035	Crushed and shattered at end in vicinity of knots.
	778	do		do		5058	Crushed at middle at 10 millimeters knot.
	787	do	Bridgeton	Ed. Sinclair		5851	Crushed at 76 millimeters from end; opened grain from end to end.
	787	do	do	do		6301	Triple flexure perpendicular to rings.
	798	Province of Quebec	Danville	Grand Trunk railway.		7403	Crushed at 6 millimeters knot 64 millimeters from end.
	798	do	do	do		8346	Crushed at 13 to 25 millimeters from end.
	817	West Virginia	Grafton	C. G. Pringle		6321	Crushed at 19 millimeters from middle.
	817	do	do	do		6101	Crushed at 32 millimeters from middle.
	1040	Massachusetts	Danvers	J. Robinson	Moist loam	6827	Crushed at 76 millimeters from end.
	1040	do	do	do	do	5516	Triple flexure
	1042	do	North Reading	do		6486	Crushed at 76 millimeters from end.
	1042	do	do	do		5126	Crushed at 51 millimeters from middle.
388. <i>Tsuga Caroliniana</i> Hemlock.	623	North Carolina	Hendersonville	A. H. Curtis	Dry, rocky	6450	Crushed at 5 millimeters knot near middle.
389. <i>Tsuga Mertensiana</i> Hemlock.	971	Washington territory.	Wilkeson	G. Engelmann and C. S. Sargent.	Rich loam	9185	Crushed at 76 millimeters from end.
	995	Alaska	Sitka	Paul Schultze		7021	Crushed at 6 millimeters knot 45 millimeters from end.
	995	do	do	do		9435	Crushed at 51 millimeters from middle.
390. <i>Tsuga Pattoniana</i> .	980	British Columbia	Silver peak, near Fraser river.	G. Engelmann and C. S. Sargent.	Gravelly loam	5806	Triple flexure; middle bend 36 millimeters eccentric at 3 millimeters knot.
	980	do	do	do	do	6341	Crushed at 3 millimeters knot at middle.
391. <i>Pseudotsuga Douglasii</i> Red Fir. Yellow Fir. Oregon Pine. Douglas Fir.	2713	Colorado	Alpine	T. S. Brandegee	Moist	6613	Triple flexure
	627	California	Saw-mill, Strawberry valley.	G. Engelmann and C. S. Sargent.		9117	Crushed at 51 millimeters from middle.
	627	do	do	do		9390	Crushed at end and at 102 millimeters from end.
	702	Oregon	Saw-mill, Marshfield.	do		5489	Triple flexure perpendicular to rings.
	704	do	do	do		9026	Crushed at 51 millimeters from end.
	705	do	E. B. Dean's saw-mill, Marshfield.	do		8709	Triple flexure
	706	do	do	do		20550	Crushed at 89 millimeters from end.
	708	do	do	do		9979	Crushed at 45 millimeters from end.
	708	do	do	do		11022	Crushed at 25 millimeters from end at 3 millimeters knot.
	709	do	do	do		7303	Crushed at 76 millimeters from end.
	709	do	do	do		7212	Crushed at 76 millimeters from middle.
	720	Montana	Saw-mill, Missoula	S. Watson		7249	Crushed at 51 millimeters from end; shattered end.
	720	do	do	do		8936	Triple flexure; middle bend 38 millimeters eccentric.
	732	California	Lassen's peak	Sierra Lumber Company.		8029	Crushed at 19 millimeters from end.
	732	do	do	do		7802	Crushed at end
	881	Utah	Salt Lake	M. E. Jones	Rocky	6895	Crushed at 3 millimeters knots 38 millimeters from end.
	881	do	do	do	do	6305	Crushed at 18 millimeters knot at middle.
	973	British Columbia	Saw-mill, Burrard inlet.	C. S. Sargent		7235	Crushed at middle
	973	do	do	do		7779	Crushed at 38 millimeters from end.
	974	do	do	do		8600	Crushed at 51 millimeters from end.
	974	do	do	do		9026	do
	986	do	Saw-mill, Victoria.	G. Engelmann and C. S. Sargent.		7578	Triple flexure
	989	Oregon	Saw-mill, Portland	do		7689	Crushed at group of knots 25 millimeters from middle.
	1008	British Columbia	Saw-mill, Burrard inlet.	do		7756	Crushed and shattered at end.

UNITED STATES UNDER COMPRESSION—Continued.

Direction of grain.	PRESSURE, IN KILOGRAMS, REQUIRED TO PRODUCE AN INDENTATION, IN MILLIMETERS, OF—												Remarks.	Office number.
	0.25	0.51	0.76	1.02	1.27	1.52	1.78	2.03	2.28	2.54	4.81	5.08		
	667	852		894	975	1002	1066	1116	1152	1179	1220	1429	Slight shearing of fibers; split at end.....	210
	885	998	1120	1320	1397	1415	1438	1400	1511	1628	1760		Sheared fibers; split at end.....	772
	767	1106	1257	1315	1433	1492	1551	1637	1710	1778	2078		Sheared fibers; split from end to end.....	772
	1159	1343	1474	1538	1628	1724	1778	1851	1910	1969	2404	2722	Slight shearing of fibers; split at end.....	775
	993	1198	1288	1352	1442	1547	1615	1687	1737	2096	2313		do.....	775
	730	944	1021	1075	1107	1157	1170	1211	1234	1252	1361	1497	Sheared fibers.....	778
	794	1811	1415	1411	1415	1447	1479	1538	1628	1674	1837	1996	do.....	778
	925	1474	1461	1461	1465	1583	1637	1705	1751	1760	1996	2141	do.....	787
	499	857	964	998	1048	1089	1143	1170					Split at end.....	787
	1338	1833	2041	2218	2241	2245	2245	2250	2254	2241	2282		Slight shearing of fibers; split at end.....	798
	1111	1452	1551	1660	1733	1792	1833	1828					do.....	798
	544	706	780	803	844	857	871	907	921	939	1034		Sheared fibers; split from end to end.....	817
	522	763	785	826	852	871	898	948	969	1002	1120	1184	Sheared fibers.....	817
	572	1034	1129	1193	1247	1288	1325	1365	1402	1456	1710		Slight shearing of fibers; split at end.....	1040
	1338	1600	1837	1937	2068	2141	2259	2309	2354	2381	2676		Sheared fibers; split at ends.....	1040
	476	867	1093	1225	1306	1352	1397	1415	1415	1420	1520	1687	Sheared fibers.....	1042
	658	771	817	871	894	939	989	1021	1034	1071			Slight shearing of fibers; split at ends.....	1042
	1184	1189	1842	1928	1996	2032	2078	2091	2123	2141	2336	2405	Slight shearing of fibers; split at end.....	622
	563	848	971	1048	1098	1139	1170	1220	1261	1302	1529		Slight shearing of fibers; split at ends.....	971
	912	1179	1315	1397	1461	1533	1597	1656	1733	1760	2223		do.....	965
	1406	2032	2132	2228	2313	2395	2495	2558	2640	2681	3053		Sheared fibers; split at ends.....	965
	862	1315	1438	1501	1551	1592	1647	1687	1737	1799	2019	2227	Sheared fibers.....	980
	1089	1488	1533	1665	1778	1828	1910	1960	1996	2068	2381	2595	do.....	980
	1406	1951	2118	2277	2345	2436	2467	2531	2576	2599	2899		Sheared fibers; split at end.....	271*
	866	1170	1279	1311	1325	1361	1406	1447	1479	1501	1715		Slight shearing of fibers; split at end.....	627
	739	1069	1106	1221	1284	1320	1356	1406	1434	1461	1685		do.....	627
	1306	2141	2440	2576	2687	2758	2799	2899	2930				Slight shearing of fibers; split at end; 10 millime- knot on indented section.	702
	1338	2096	2209	2322	2391	2472	2245	2590	2645	2731			Slight shearing of fibers; split at end.....	704
	1161	1583	1656	1710	1742	1778	1833	1896	1933	1987	2223	2359	Sheared fibers; split at end.....	705
	866	1120	1189	1257	1293	1352	1374	1411	1465	1501	1742		do.....	706
	1769	2391	2522	2678	2817	2971	3094	3257	3411	3557	4105		do.....	708
	1134	2078	2308	2409	2499	2595	2672	2740	2821	2917			do.....	708
	1060	1207	1247	1297	1352	1370	1384	1406	1483	1511	1724	1833	Sheared fibers.....	709
	1229	1179	1089	1293	1429	1529	1642	1706	1765	1824	1951	2023	do.....	709
	1035	1198	1238	1306	1370	1411	1447	1497	1536	1574	1814	2005	do.....	720
	971	1438	1513	1506	1515	1547	1628	1674	1719	1774			do.....	720
	862	1107	1220	1296	1320	1361	1433	1483	1506	1556			Slight shearing of fibers; split at end.....	732
	1533	2200	2295	2313	2245	2109	2132	2246	2350	2436			Sheared fibers; split at end.....	732
	1833	1941	2482	2545	2595	2635	2713	2758	2821	2912	3296	3561	do.....	881
	1315	1992	2118	2254	2300	2345	2409						Slight shearing of fibers; split at end.....	881
	862	957	993	916	1012	1016	1084	1161	1116	1143	1460	1579	Sheared fibers.....	973
	1098	1814	1982	1941	1800	1778	1569	1565	1637	1733	2495	2722	do.....	973
	895	1216	1225	1202	1210	1338	1411	1401	1524	1374	1824		Sheared fibers; split at end.....	974
	767	1089	1152	1202	1252	1302	1343	1365	1402	1433	1588	1687	do.....	974
	780	862	953	1002	1048	1080	1120	1139	1170	1193	1311		do.....	986
	983	1320	1470	1547	1588	1647	1674	1728	1765	1801	2130		do.....	989
	839	1060	1297	1302	1347	1397	1452	1488	1533	1599			Fibers did not shear.....	1608

TABLE V.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Longitudinal compression: Ultimate strength, in kilos.	Remarks.
391. <i>Pseudotsuga Douglasii</i> —cont'd	1008	British Columbia	Saw-mill, Burrard inlet.	G. Engelmann and C. S. Sargent.		8981	Crushed at 38 millimeters from middle.
	1011	Oregon	Oregon Railway and Navigation Company.	do		8799	Crushed at 3 millimeters knot at middle.
	1011	do	do	do		6804	Crushed at 16 millimeters knot 76 millimeters from end.
	1016	do	Weidler's saw-mill, Portland.	do		8981	Crushed at 80 millimeters from end.
	1016	do	do	do		9798	Crushed at 5 millimeters from middle.
	1018	do	Saw-mill, Astoria	do		10002	Crushed at 51 millimeters from end.
	1018	do	do	do		9662	Crushed at 38 millimeters from end.
	1020	do	Portland Furniture Company.	do		7892	do
	1020	do	do	do		7666	Triple flexure
	1022	do	do	do		8346	Crushed at 51 millimeters from end.
	1022	do	do	do		8392	Crushed at 51 millimeters from middle.
391. <i>Pseudotsuga Douglasii</i> , var. <i>macrocarpa</i> . <i>Hemlock</i> .	642	California	Saw-mill, San Bernardino.	W. G. Wright		6872	Split obliquely; pieces flew from machine.
	642	do	do	do		7938	Crushed at 103 millimeters from end; opened grain.
392. <i>Abies Fraseri</i> <i>Balsam. She Balsam.</i>	523	North Carolina	Roan mountain	Walcott Gibbs	Peaty loam	5674	Crushed at 6 millimeters knot 31 millimeters from end.
	523	do	do	do	do	5239	Crushed at 10 millimeters knot 25 millimeters from middle.
393. <i>Abies balsamea</i> <i>Balsam Fir. Balm of Gilead Fir.</i>	377	Vermont	Monkton	C. G. Pringle	Peaty	5851	Crushed at 25 millimeters from middle.
394. <i>Abies subalpina</i> <i>Balsam.</i>	449	Colorado	Forest City	T. S. Brandegee	Moist, sandy loam	4854	Crushed at middle at 25 millimeters knot.
	449 ¹	do	do	do	do	4400	Crushed at 6 millimeters knot 38 millimeters from middle; split along grain.
	449 ¹	do	do	do	do	3856	Crushed at 10 millimeters knot at middle; split along grain.
	449 ²	do	do	do	do	5661	Crushed at 38 millimeters from end.
	820	do	do	do	do	5398	Crushed at middle
	820	do	do	do	do	4808	Crushed and shattered at end.
395. <i>Abies grandis</i> <i>White Fir.</i>	1010	Oregon	Portland	G. Engelmann and C. S. Sargent.	Rich, alluvial	6033	Crushed at 25 millimeters from middle.
	1010	do	do	do	do	6477	Crushed at 19 millimeters from middle.
396. <i>Abies concolor</i> <i>White Fir. Balsam Fir.</i>	529	Colorado	Engelmann's cañon	Robert Douglas	Rocky	4309	Crushed 51 millimeters from end (at end bearing transverse test); split along rings.
	529	do	do	do	do	4967	Split at 10 millimeters knots 25 millimeters from middle.
	639	California	Strawberry valley	G. Engelmann and C. S. Sargent.	Alluvial	6350	Triple flexure.
	639	do	do	do	do	5670	Triple flexure perpendicular to rings; opened grain.
	733	do	Lassen's peak	Sierra Lumber Company.		8618	Crushed at middle
	733	do	do	do		7507	Crushed at 38 millimeters from end.
398. <i>Abies amabilis</i>	1004	British Columbia	Silver peak, near Fraser river.	G. Engelmann and C. S. Sargent.	Rich, sandy loam	7838	Crushed at 25 millimeters from middle.
	1004	do	do	do	do	7122	Crushed at middle
399. <i>Abies nobilis</i> <i>Red Fir.</i>	965	Oregon	Cascade mountains.	do	Rich	7235	Crushed at 51 millimeters from end.
	965	do	do	do	do	7276	Crushed at 25 millimeters from middle.
400. <i>Abies magnifica</i> <i>Red Fir.</i>	647	California	Soda Springs	G. Engelmann and C. S. Sargent.	Gravelly loam	7258	Crushed at 51 millimeters from end.
	647	do	do	do	do	6668	Triple flexure; middle bend 25 millimeters eccentric.
401. <i>Larix Americana</i> <i>Larch. Black Larch. Tamarack Hackmatack.</i>	226 ¹	Vermont	Charlotte	C. G. Pringle	Cold, swampy	12021	Crushed at 25 millimeters from middle.
	226 ¹	do	do	do	do	10387	Triple flexure; middle bend 25 millimeters eccentric.
	226 ²	do	do	do	do	6849	Crushed at 6 millimeters knot at middle; split from end to end.
	226 ²	do	do	do	do	8165	Crushed at middle; split
	774	New Brunswick	Bay of Fundy	Intercolonial railway.		7802	Crushed at 13 millimeters knot 51 millimeters from end; opened between rings.
	774	do	do	do		9095	Crushed at 64 millimeters from middle at 25 millimeters knot.

UNITED STATES UNDER COMPRESSION—Continued.

Direction of grain.	PRESSURE, IN KILOGRAMS, REQUIRED TO PRODUCE AN INDENTATION, IN MILLIMETERS OF—												Remarks.	Office number.
	0.25	0.51	0.76	1.02	1.27	1.52	1.78	2.03	2.28	2.54	2.81	3.08		
	1315	2087	2254	2241	1906	1978	1980	2073	2227	1987	2876	2903	Sheared fibers.....	1008
	771	1306	1433	1479	1520	1556	1588	1006	1633	1080	1452	1565	do.....	1011
	925	1166	1216	1275	1334	1388	1429	1447	1488	1524	1724		Sheared fibers; split at end.....	1011
	895	1107	1157	1202	1257	1302	1352	1393	1429	1438	1642		Fibers did not shear; split at end.....	1016
	817	998	1048	1075	1125	1184	1234	1284	1343	1388			Slight shearing of fibers; split at end.....	1016
	1048	1211	1297	1343	1393	1456	1537	1615	1660	1737	2078		do.....	1018
	889	1129	1189	1229	1284	1329	1384	1447	1483	1506			do.....	1018
	730	780	807	839	944	1002	1039	1071	1102	1184	1361		do.....	1020
	671	862	953	930	1002	1111	1157	1198	1211	1252	1429	1624	Sheared fibers.....	1020
	817	848	885	907	962	993	1080	1080	1107	1143	1306		Slight shearing of fibers; split at end.....	1022
	1179	1452	1497	1406	1293	1315	1470	1651	1751	1805	1987	2313	Sheared fibers.....	1022
	1080	1551	1633	1610	1615	1692	1778	1846	1910	1960	2177	2409	do.....	642
	1187	1520	1583	1624	1669	1728	1746	1760	1824	1842	2062	2186	do.....	642
	540	830	964	1061	1093	1125	1157	1170	1193	1202	1379	1442	do.....	523
	499	835	939	975	1002	1039	1080	1116	1129	1166	1306	1379	do.....	523
	644	866	993	1107	1202	1257	1306	1365	1397	1447			Slight shearing of fibers; split at end; 4 millimeters knot in indented section.....	877
	780	880	993	1048	1093	1125	1184	1211	1238	1266	1442	1588	Sheared fibers.....	449
	726	1016	1080	1111	1139	1175	1211	1247	1261	1284	1474	1597	do.....	449 ¹
	794	925	980	944	980	1007	1030	1061	1116	1116	1270	1374	do.....	449 ¹
	594	821	894	939	975	1012	1034	1052	1080	1102	1270		Sheared fibers; split at end.....	449 ²
	590	817	866	889	894								Split at end.....	820
	563	826	957	989	1007	1030	1039	1052	1089	1057	1170	1225	Sheared fibers; split at ends.....	820
	631	817	839	839	857	912	962	1012	1066	1068	1225	1379	do.....	1010
	499	617	669	712	762	803	817	844	866	885			Slight shearing of fibers; split at end.....	1010
	769	1002	1093	1125	1143	1166	1170	1179	1193	1202	1302		Sheared fibers; split at end.....	529
	522	776	894	971	1062	1025	1052	1064	1098	1116	1225		do.....	529
	749	1134	1189	1198	1216	1247	1279	1320	1338	1361	1610	1769	Sheared fibers.....	639
	852	1116	1207	1270	1320	1347	1402	1438	1465	1511	1746	1905	do.....	639
	1134	1243	1338	1438	1533	1592	1692	1769	1824	1869			Fibers did not shear; split at end.....	733
	1134	1179	1229	1270	1275	1325	1470	1533	1592	1624	1800		Sheared fibers; split at end.....	733
	504	844	880	939	998	1080	1116	1157	1216	1257	1429		do.....	1004
	680	862	930	1002	1061	1098	1129	1166	1193	1234	1452		do.....	1004
	1356	1565	1701	1842	1941	2105	2218	2263	2345	2413	2858	3221	Slight shearing of fibers; split at end.....	965
	1184	1610	1696	1778	1802	1969	2050	2123	2173	2214	2586		Sheared fibers; split at end.....	965
	844	1016	1093	1148	1175	1207	1252	1275	1311	1325	1474	1651	Sheared fibers.....	647
	1021	1583	1760	1860	1914	1982	2059	2073	2118	2141	2404	2586	do.....	647
	1270	1709	2087	2241	2391	2472	2549	2581	2640	2717	2948	3130	Slight shearing of fibers; split at ends.....	226 ¹
	1225	1533	1628	1778	1869	1946	1987	2050	2114	2164	2449		Slight shearing of fibers; split at end; specimen 120 millimeters long.....	226 ²
	1039	1533	1728	1860	1982	2073	2150	2263	2381				Fibers did not shear; split at end; specimen 120 millimeters long.....	226 ³
	1724	2358	2699	2875	2939	3026	3039	3502	3506	3612			Slight shearing of fibers; shattered stick; specimen 120 millimeters long.....	226 ⁴
	852	1143	1247	1348	1388	1429	1442	1470	1497	1538	1746		Slight shearing of fibers; split at end.....	774
	794	1184	1279	1352	1447	1470	1474	1483	1506	1538	1769		do.....	774

FOREST TREES OF NORTH AMERICA.

TABLE V.—BEHAVIOR OF THE PRINCIPAL WOODS OF THE

Species.	Office number.	State.	Locality.	Collector.	Soil.	Longitudinal compression: Ultimate strength, in kilos.	Remarks.
401. <i>Larix Americana</i> —continued.	781	New Brunswick	Bay of Fundy	Intercolonial railway.		8147	Triple flexure
	781	do	do	do		8142	Crushed at 25 millimeters from middle.
	786	do	Bridgeton	Ed. Sinclair		9390	do
	786	do	do	do		7666	Crushed at 51 millimeters from middle.
	795	do	Danville	Grand Trunk railway.		9480	Crushed at 64 millimeters from middle.
	795	do	do	do		8437	Crushed at middle
	840	Massachusetts	Wenham	J. Robinson	Swampy	6967	Triple flexure parallel to rings; intersecting "Cooper lines".
	840	do	do	do	do	7530	Crushed at 102 millimeters from end on one face.
402. <i>Larix occidentalis</i> . <i>Tamarack.</i>	719	Montana	Missoula	S. Watson		9367	Triple flexure
	719	do	do	do		10047	Crushed at 64 millimeters from end.
	984	Washington territory.	Fulda	W. Siksdorf	Moist	11648	Crushed at 51 millimeters from end at 3 millimeters knot.
	984	do	do	do	do	12633	Crushed at 3 millimeters knot 19 millimeters from end.
	1006	do	do	do	do	11785	Crushed at 25 millimeters from middle.
	1006	do	do	do	do	10660	Crushed at 96 millimeters from end; opened grain.
PALMACEÆ.							
405. <i>Washingtonia filifera</i> . <i>Fan-leaf Palm.</i>	1150	California	Agua Caliente	W. G. Wright	Dry, gravelly	1461	Split obliquely; fracture 228 millimeters long.
	1150	do	do	do	do	5806	Split obliquely; fracture 305 millimeters long.

UNITED STATES UNDER COMPRESSION—Continued.

Direction of grain.	PRESSURE, IN KILOGRAMS, REQUIRED TO PRODUCE AN INDENTATION, IN MILLIMETERS, OF—												Remarks.	Office number.
	0.25	0.51	0.76	1.02	1.27	1.52	1.78	2.03	2.28	2.54	4.81	5.08		
	993	1125	1220	1311	1388	1461	1538	1601	1674	1774	2313	2586	Slight shearing of fibers.....	781
	1270	1765	1905	2041	2159	2254	2345	2422	2504	2572	3198		Slight shearing of fibers; split at end	781
	862	934	1025	1075	1125	1169	1229	1261	1302	1325			do.....	788
	703	1202	1270	1325	1347	1433	1533	1624	1692	1778	2087	2313	do.....	788
	730	1111	1189	1288	1329	1388	1429	1443	1461	1501			do.....	788
	1002	1315	1429	1506	1551	1656	1769	1860	1905	1941			do.....	795
	1315	1783	1842	1987	2082	2168	2227	2313	2422	2472	3085		Sheared fibers; split at end; indented section covers 6 millimeters knot.	840
	1179	1769	1892	1946	2028	2168	2259	2390	2490	2586	3030		Slight shearing of fibers; split at end	840
	953	1374	1774	1880	1928	2000	2019	2050	2118	2186	2541	2713	Sheared fibers.....	719
	1211	1452	1579	1696	1796	1890	1946	1987	2082	2118	2449		Slight shearing of fibers; split at end	719
	1315	1451	1547	1619	1710	1792	1910	1987	2046	2123	2563		do.....	984
	1633	2396	2350	2345	2400	2545	2681	2808	2048	3094			do.....	984
	1656	2681	3134	3348	3375	3470	3652	3756	3901	4051			do.....	1006
	1406	1669	1905	2046	2159	2300	2391	2500	2549	2672			Fibers did not shear.....	1006
	318	508	576	617	658	690	708	735	758	776	945	1048	Sheared fibers.....	1159
	812	1207	1325	1397	1443	1497	1551	1610	1628	1656	1982	1996	Slight shearing of fibers.....	1159

31 FOR

PART III.

THE FORESTS OF THE UNITED STATES
IN THEIR
ECONOMIC ASPECTS.

THE FORESTS OF THE UNITED STATES IN THEIR ECONOMIC ASPECTS.

GENERAL REMARKS.

The maps of relative average forest density joined to this report are intended to illustrate the present productive capacity of the forest covering of the country (map No. 16, portfolio). They are based, except in the case of the extreme western states and territories, upon the returns of enumerators. In states originally wooded all land not accounted for in the returns as cleared or treeless, or otherwise known to be destitute of tree covering, is treated as forest. The county is taken as the unit, and is seldom divided, unless varied topography or different natural features in different parts makes further subdivision desirable. In the western states and territories, where topography determines forest distribution, county lines are disregarded, and the estimates are based upon special reports of census experts, or upon the published reports of the various government surveys, maps, etc. The condition and productive capacity of the forest covering have been carefully investigated at many points in each county or unit region, and the area covered with forest, obtained in the manner described above, is multiplied by the average stand of timber or other useful wood. The results thus obtained are necessarily greatly generalized to conform to the scale of the maps used.

The following statement represents the value of the forest crop of the United States for the census year, so far as it has been possible to obtain it:

Saw logs	\$139,836,869	Charcoal used as fuel—		Wood used in the manufacture of—	
Wood used for domestic purposes as fuel (estimated)	306,950,040	In manufacture of iron	\$4,726,114	Handles	\$897,170
Wood used by railroads as fuel	5,126,714	In manufacture of precious metals ..	29,806	Wheel stock	1,360,892
Wood used by steamboats as fuel	1,812,083	In the twenty largest cities	521,316	Wood pulp	1,974,074
Wood used as fuel—		Naval stores	5,000,000	Baskets	814,125
In the manufacture of brick and tile ..	3,978,331	Southern moss	500,000	Excelsior	150,800
In the manufacture of wool	425,239	Railroad ties (29,554,694)	9,806,247	Oars	81,000
In the manufacture of salt	121,681	Fence posts (for fencing railroads) ..	180,000	Shoe pegs	72,000
In the production of precious metals ..	2,874,593	Uncultivated vegetable substances used		Hand-made shingles	47,952
In other mining operations	673,692	in the manufacture of medicines	587,000		
		Uncultivated nuts	78,540	Total	490,073,094
		Hoop-poles	1,947,816		

These returns are incomplete and often unsatisfactory. Many important items are omitted entirely. It was found impossible to obtain statistics of the amount and value of the wood (posts, split rails, etc.) used in fencing, with the exception of posts used by railroads. The amount of material thus consumed annually must be very large, probably exceeding \$100,000,000 in value. No returns of the amount and value of the bark of different trees used in tanning leather have been received, and there are no statistics of the amount and value of the unsawed timber produced—spars, piles, telegraph and other poles, hewed timber, hard wood exported in the log, ships' knees, etc.—that is, all timber not manufactured in saw-mills into lumber. The value of the timber of this sort cut in the United States every year must be very large. The returns include the railway ties laid down by completed roads, and do not embrace those used in the construction of some 10,000 miles of new road built during the census year. It was found impossible to obtain even an estimate of the amount and value of the cooperage stock produced outside of regular saw-mills, and the returns of hand-made shingles only include those made from cypress at a few points in the south Atlantic region. Maple sugar to the amount of 36,576,061 pounds and 1,796,048 gallons of molasses were produced in the forests of the United States during the year 1879. No statistics of the value of these products have, however, been received. Statistics of the value of material consumed in the manufacture of excelsior, wood pulp, wheel stock, handles, shoe pegs, baskets, oars, and hoop-poles are incomplete, and do not fully represent the value of the wood used. The statistics of the value of wild nuts and wild vegetable substances collected are very incomplete, and it has been found impossible to separate the value of the imported from that of the native wood used in the manufacture of veneers, an industry consuming a large amount of high-priced hard wood. Could complete returns of the forest crop of the census year have been obtained it is not improbable that it would be found to exceed \$700,000,000 in value.

THE LUMBER INDUSTRY.

The following table represents the volume, by states and territories, of the lumber industry of the United States for the census year, as derived from the returns of the enumerators on the schedule of manufactures, and from the reports of special agents for manufactures in cities having at the time of the Ninth Census 8,000 or more inhabitants. No distinction between the different kinds of wood sawed was attempted in the enumeration:

THE FORESTS OF THE UNITED STATES.

STATISTICS OF THE LUMBERING INDUSTRY OF THE

	States and Territories.	Number of establishments.	Capital.	AVERAGE NUMBER OF HANDS EMPLOYED.				LABOR.	MATERIALS.	
				Maximum at any one time in the year.	Males above 16 years.	Females above 15 years.	Children and youth.	Wages paid during the year.	Value of logs.	Value of mill supplies.
			Dollars.					Dollars.	Dollars.	Dollars.
	The United States	25,708	181,186,122		141,564	425	5,967	31,845,974	139,836,869	6,314,516
1	Alabama	354	1,545,655	2,798	1,611		86	424,156	1,517,986	90,649
2	Alaska									
3	Arizona	13	102,450	126	77		2	33,375	126,486	5,300
4	Arkansas	310	1,067,840	2,985	1,690		54	237,394	1,009,954	60,441
5	California	251	6,454,718	4,945	3,423		11	1,095,736	2,055,635	188,868
6	Colorado	96	481,200	1,605	870		7	112,931	654,500	45,794
7	Connecticut	300	657,300	1,262	699		8	178,336	609,024	32,545
8	Dakota	39	113,750	451	286		4	54,974	269,235	12,640
9	Delaware	86	259,250	646	378		13	40,694	229,763	13,612
10	District of Columbia	1	25,000	35	25			6,000	32,000	2,060
11	Florida	135	2,219,550	3,240	1,945		85	562,249	1,763,617	103,596
12	Georgia	655	3,101,452	4,971	3,298		94	554,085	3,049,435	147,720
13	Idaho	48	192,460	311	169		4	33,367	213,661	16,875
14	Illinois	640	3,295,433	5,799	3,652		199	787,867	2,959,537	185,368
15	Indiana	2,022	7,048,088	16,252	9,926		413	1,571,740	9,290,428	336,669
16	Indian territory									
17	Iowa	328	4,946,390	4,155	2,526		463	825,244	4,023,661	118,224
18	Kansas	146	262,975	831	507		9	66,757	421,738	25,711
19	Kentucky	670	2,290,558	5,140	2,506	1	94	671,939	2,238,888	171,855
20	Louisiana	175	903,950	1,514	943		33	200,063	1,106,280	80,779
21	Maine	848	6,339,396	9,836	6,480		188	1,161,142	4,754,613	197,344
22	Maryland	369	1,237,694	1,769	1,216		23	223,786	1,041,836	64,959
23	Massachusetts	606	2,480,340	3,130	1,940		30	431,612	1,827,497	76,608
24	Michigan	1,649	39,260,422	30,866	22,732	143	1,300	6,967,905	30,819,003	1,432,369
25	Minnesota	234	6,771,145	3,772	2,732	22	100	924,473	4,408,468	120,587
26	Mississippi	295	922,595	2,113	1,123		47	197,867	1,190,902	23,214
27	Missouri	881	2,867,970	6,678	3,408		95	669,644	3,113,049	102,243
28	Montana	36	208,200	374	143			47,945	257,320	20,778
29	Nebraska	38	93,375	295	136		4	29,213	153,823	11,055
30	Nevada	9	132,000	66	35			9,892	151,790	11,020
31	New Hampshire	680	3,745,790	4,765	3,056		48	548,556	2,159,461	113,530
32	New Jersey	284	1,657,395	1,066	760		8	179,693	942,752	47,227
33	New Mexico	26	74,675	282	172			24,240	100,145	16,910
34	New York	2,822	13,230,934	17,509	11,056		389	2,162,972	8,628,874	490,389
35	North Carolina	776	1,743,217	5,334	2,938		91	447,431	1,490,616	86,523
36	Ohio	2,352	7,944,412	15,277	8,769		548	1,708,300	8,603,127	292,979
37	Oregon	228	1,577,875	1,185	566		13	242,154	1,294,703	36,639
38	Pennsylvania	2,827	21,418,588	21,160	14,443	8	463	2,918,459	13,378,599	576,841
39	Rhode Island	49	144,250	260	139		13	33,143	116,085	4,803
40	South Carolina	420	1,056,265	2,338	1,431		37	221,963	1,170,088	67,273
41	Tennessee	755	2,004,503	5,587	3,577		141	549,222	2,006,124	136,761
42	Texas	324	1,660,952	4,579	3,136	1	49	732,914	1,909,794	186,961
43	Utah	107	272,750	845	375		10	65,175	216,619	21,655
44	Vermont	688	3,274,250	4,501	2,411		100	426,953	1,939,775	82,093
45	Virginia	907	2,122,925	5,812	3,922		89	540,231	1,884,288	119,489
46	Washington	37	2,456,450	891	495		4	200,539	1,174,005	14,070
47	West Virginia	472	1,068,920	3,765	2,057		126	459,945	1,307,843	67,529
48	Wisconsin	704	19,824,050	14,079	7,748	250	467	2,257,218	12,219,097	262,376
49	Wyoming	7	26,700	68	38			6,380	24,725	2,625

THE FORESTS OF THE UNITED STATES.

487

UNITED STATES FOR THE YEAR ENDING MAY 31, 1880.

PRODUCTS.								
Lumber (board measure).	Laths.	Shingles.	Staves.	Sets of headings.	Spool and bobbin stock (board measure).	Value of all other products.	Total value of all products.	Rank according to value of products.
<i>Feet.</i> 18,091,356,000	<i>Number.</i> 1,761,788,000	<i>Number.</i> 5,555,046,000	<i>Number.</i> 1,248,228,000	<i>Number.</i> 146,528,000	<i>Feet.</i> 84,076,000	<i>Dollars.</i> 2,682,668	<i>Dollars.</i> 233,268,729	
251,851,000	14,147,000	5,427,000	2,857,000	437,000			2,649,684	23 1
10,715,000	150,000	1,760,000	300,000			1,010	215,918	44 3
172,503,000	6,527,000	61,758,000	1,640,000	350,000			1,793,848	29 4
304,795,000	2,420,000	188,718,000	2,063,000	1,203,000		3,000	4,428,950	13 5
63,792,000	4,925,000	27,214,000				700	1,051,295	34 6
64,427,000	1,719,000	7,192,000	270,000	12,000	33,000	12,980	1,076,455	33 7
29,286,000	564,000	4,823,000				500	435,702	37 8
31,572,000	817,000	506,000	4,510,000	550,000			411,060	38 9
4,000,000	1,000,000					2,000	50,000	46 10
a 247,627,000	20,101,000	3,061,000	791,000	110,000		13,999	3,080,291	21 11
451,798,000	17,438,000	25,832,000	2,014,000	984,000	4,000	3,975	4,875,810	12 12
18,204,000	750,000	4,235,000				60,200	349,635	40 13
b 334,244,000	b 25,977,000	b 15,306,000	24,448,000	1,385,000	30,000	16,807	5,063,037	11 14
915,943,000	28,081,000	26,634,000	283,071,000	26,889,000	1,957,000	145,750	14,260,830	5 15
c 412,578,000	c 79,924,000	c 128,100,000	5,335,000	650,000		88,343	6,185,628	9 16
45,281,000	25,000	835,000			150,000	3,400	682,697	35 17
305,684,000	26,856,000	25,253,000	23,148,000	8,174,000	883,000	75,655	4,064,361	14 18
133,472,000	7,745,000	30,195,000	220,000	33,000		15,470	1,764,640	30 19
506,656,000	184,820,000	426,580,000	62,376,000	3,312,000	13,426,000	182,618	7,933,868	7 20
123,336,000	7,955,000	4,429,000	16,227,000			149,894	1,813,332	28 21
d 205,244,000	d 16,947,000	d 19,667,000	21,062,000	1,860,000	572,000	44,895	3,120,184	20 22
4,172,572,000	461,805,000	2,584,717,000	199,821,000	21,897,000	6,038,000	531,406	53,449,928	1 23
563,974,000	88,088,000	194,566,000	7,825,000	547,000		21,100	7,806,038	8 24
168,747,000	7,908,000	5,355,000	60,000			5,849	1,920,335	27 25
e 399,744,000	e 20,839,000	e 8,832,000	21,426,000	3,363,000		7,097	5,265,617	10 26
21,420,000	2,620,000	9,627,000				1,900	527,695	36 27
18,585,000						1,100	265,062	41 28
21,545,000		485,000					243,200	42 29
292,267,000	49,454,000	67,086,000	31,354,000	3,491,000	3,072,000	58,612	3,842,012	15 30
109,679,000	8,948,000	10,717,000	40,000	155,000	883,000	40,885	1,627,640	32 31
11,195,000	107,000	722,000	20,000				178,930	45 32
1,184,220,000	79,399,000	305,711,000	62,654,000	22,196,000	1,003,000	285,263	14,856,910	4 33
241,822,000	13,340,000	8,707,000	45,000	571,000	1,258,000	7,195	2,672,796	32 34
910,832,000	50,625,000	24,876,000	214,245,000	25,779,000	25,000	196,788	13,864,460	6 35
177,171,000	18,245,000	5,040,000				10,500	2,030,463	26 36
1,733,844,000	183,740,000	288,561,000	80,062,000	10,401,000	326,000	398,044	22,457,359	2 37
8,469,000	10,000	1,986,000	365,000		3,700,000	174	240,579	43 38
185,772,000	23,133,000	10,036,000	385,000	93,000		41,700	2,061,507	25 39
302,673,000	21,275,000	14,205,000	4,342,000	570,000	6,000	72,998	3,744,905	16 40
328,968,000	14,131,000	112,523,000		140,000		10,850	3,673,449	17 41
25,709,000	1,363,000	9,293,000				1,765	375,164	39 42
322,942,000	19,745,000	55,711,000	13,219,000	1,572,000	415,000	2,575	3,258,816	19 43
315,939,000	14,402,000	8,223,000	14,333,000	929,000	800,000	30,355	3,434,168	18 44
160,176,000	6,550,000	3,610,000	23,666,000				1,734,742	31 45
180,112,000	12,071,000	3,695,000	41,992,000	1,952,000		40,195	2,431,857	24 46
1,542,021,000	215,132,000	862,922,000	82,545,000	7,498,000		152,171	17,952,347	3 47
2,960,000	300,000	805,000					40,990	47 48

a Including 77,500,000 feet manufactured from logs cut in Alabama.

b Including 73,700,000 feet lumber, 13,041,000 laths, and 11,226,000 shingles, manufactured from logs cut in Wisconsin.

c Including 334,199,000 feet lumber, 78,728,000 laths, and 127,591,000 shingles, manufactured from logs cut in Wisconsin.

d Including 20,000,000 feet lumber, 11,982,000 laths, and 800,000 shingles, manufactured from logs cut in New Hampshire and Vermont.


e Including 27,000,000 feet lumber, 12,400,000 laths, and 5,300,000 shingles, manufactured from logs cut in Wisconsin.


THE FORESTS OF THE UNITED STATES.

In the following table the average importance of the saw-mills located in the different states and territories is shown:

AVERAGE SIZE AND PRODUCT OF SAW-MILLS IN EACH STATE AND TERRITORY OF THE UNITED STATES.

States and Territories.	Total number of establishments.	Capital.	NUMBER OF HANDS EMPLOYED.		MATERIALS AND LABOR.			PRODUCTS.							
			Maximum at any one time in the year.	Average.	Value of logs.	Value of mill supplies.	Wages paid during the year.	Lumber (board measure).	Laths.	Shingles.	Staves.	Sets of head-ings.	Spool and bobbin stock.	Value of other products.	Total value of all products.
		Dolls.			Dolls.	Dolls.	Dolls.	Feet.	No.	No.	No.	No.	Feet.	Dolls.	Dolls.
The United States..	25,708	7,048	8.8	5.8	5,436	246	1,235	703,000	68,000	216,000	48,000	5,700	1,800	104	9,078
Alabama.....	354	4,366	7.9	4.6	4,288	256	1,198	712,000	40,000	15,000	6,000	1,000			7,485
Alaska.....															
Arizona.....	13	7,880	9.7	6.0	9,729	408	2,567	824,000	11,000	136,000	23,000			77	16,609
Arkansas.....	819	3,847	9.0	5.5	3,166	190	744	541,000	20,000	194,000	5,000	1,000			5,623
California.....	251	25,716	19.7	13.7	8,190	744	4,365	1,214,000	10,000	553,000	8,000	5,000		12	17,645
Colorado.....	96	5,013	16.6	9.0	6,808	476	1,176	664,000	51,000	283,000				7	10,951
Connecticut.....	300	2,119	4.0	2.0	2,030	108	594	215,000	6,000	24,000	1,000			43	359
Dakota.....	39	2,917	11.6	7.5	6,903	324	1,410	751,000	15,000	124,000				13	11,174
Delaware.....	86	3,015	7.5	4.5	2,672	158	473	367,000	4,000	6,000	52,000	6,000			4,780
District of Columbia.....	1	25,000	35.0	25.0	32,000	2,000	6,000	4,000,000	1,000,000					2,000	50,000
Florida.....	135	16,441	24.0	15.0	13,064	768	4,165	1,834,000	150,000	23,000	6,000	1,000		104	22,668
Georgia.....	655	4,735	7.5	5.0	4,655	225	845	690,000	27,000	39,000	8,000	1,000		6	7,443
Idaho.....	48	4,009	6.5	3.6	4,452	852	700	380,000	16,000	88,000				1,234	9,246
Illinois.....	640	5,149	9.0	6.0	4,624	290	1,230	522,000	40,000	24,000	38,000	2,000		26	7,911
Indiana.....	2,022	3,485	8.0	5.0	4,600	166	777	453,000	14,000	13,000	140,000	18,000		72	7,052
Indian territory.....															
Iowa.....	328	15,080	12.6	9.0	13,267	360	2,513	1,258,000	244,000	890,000	16,000	2,000		117	19,000
Kansas.....	146	1,891	5.7	3.5	2,888	176	457	810,000	40,000	6,000				23	4,700
Kentucky.....	670	3,418	7.7	3.8	3,341	256	1,003	456,000	40,000	87,000	34,000	12,000		113	6,066
Louisiana.....	175	5,165	8.6	5.6	6,321	462	1,143	762,000	44,000	170,000	1,000			88	10,083
Maine.....	848	7,475	11.6	7.8	5,607	232	1,369	668,000	218,000	503,000	73,000	4,000		215	9,356
Maryland.....	369	3,354	4.8	3.0	2,623	176	606	334,000	21,000	12,000	44,000			406	4,914
Massachusetts.....	606	4,093	5.0	3.0	3,015	126	712	838,000	28,000	82,000	35,000	3,000		73	5,149
Michigan.....	1,649	23,808	18.7	14.7	18,700	868	4,225	2,530,000	280,000	1,568,000	121,000	13,000		322	31,867
Minnesota.....	234	28,936	16.0	12.0	18,839	515	3,950	2,410,000	376,000	881,000	33,000	2,000		90	31,478
Mississippi.....	295	3,127	7.0	4.0	4,037	95	671	572,000	27,000	18,000				18	6,509
Missouri.....	861	3,255	7.6	4.0	3,534	116	760	453,000	23,000	10,600	24,000	4,000		8	6,000
Montana.....	36	5,788	11.0	4.0	7,148	577	1,332	595,000	73,000	267,000				53	14,658
Nebraska.....	38	2,457	8.0	3.7	4,048	290	771	357,000						29	6,975
Nevada.....	9	14,666	7.0	4.0	16,865	1,224	1,099	2,394,000		54,000					27,022
New Hampshire.....	680	5,508	7.0	4.5	3,175	167	806	429,000	73,000	98,800	46,000	5,000		86	5,650
New Jersey.....	284	5,836	8.7	2.7	3,319	166	633	386,000	31,000	38,000				142	5,731
New Mexico.....	26	2,870	10.8	6.7	3,851	650	932	430,000	4,000	27,000					6,690
New York.....	2,622	4,688	6.0	4.0	3,057	173	766	419,000	28,000	108,000	22,000	8,000		101	5,087
North Carolina.....	776	2,246	7.0	4.0	1,921	111	576	311,000	17,000	11,000				9	3,445
Ohio.....	2,352	3,378	6.5	4.0	3,658	124	726	387,000	21,000	10,000	91,000	11,000		83	5,896
Oregon.....	228	6,920	5.0	2.5	5,678	160	1,062	777,000	80,000	2,000				46	8,905
Pennsylvania.....	2,827	7,576	7.5	5.0	4,732	204	1,032	613,000	65,000	102,000	28,000	3,000		139	7,944
Rhode Island.....	49	2,944	5.0	3.0	2,369	98	676	172,000		40,000	7,000			8	4,000
South Carolina.....	420	2,515	5.5	3.5	2,785	160	528	442,000	55,000	24,000				99	4,837
Tennessee.....	755	2,655	7.0	5.0	2,657	181	727	400,000	28,000	18,000	5,000			96	4,960
Texas.....	324	5,126	14.0	9.8	5,894	577	2,262	1,015,000	43,000	347,000				82	11,338
Utah.....	107	2,549	7.9	3.5	2,024	202	609	240,000	14,000	87,000				16	3,506
Vermont.....	688	4,759	6.5	3.6	2,819	119	620	469,000	28,000	80,000	19,000	2,000		3	4,736
Virginia.....	907	2,340	6.0	4.0	2,055	131	593	348,000	16,000	9,000	15,000	1,000		33	3,786
Washington.....	37	68,390	24.0	13.5	31,730	380	5,420	4,329,000	177,000	97,000	639,000				40,885
West Virginia.....	472	3,535	8.0	4.0	2,770	143	974	381,000	25,000	8,000	89,000	4,000		85	5,562
Wisconsin.....	704	28,159	20.0	12.0	17,366	358	3,206	2,190,000	805,000	1,226,000	117,000	10,000		216	25,600
Wyoming.....	7	3,814	9.7	5.5	3,632	375	911	423,000	43,000	123,000					5,855





Michigan is the greatest lumber-producing state in the Union. The value of its lumber product, with that of Wisconsin and Minnesota, exceeds one-third of the total value of all the lumber manufactured in the United States. This enormous development of the lumber business in the lake region is due to the excellence of its forests, the natural advantages of the country for manufacturing lumber, and the easy communication between these forests and the treeless agricultural region west of the Mississippi river.

The extinction of the forests of the lake region may be expected to seriously affect the growth of population in the central portion of the continent. The country between the Mississippi river and the Rocky mountains, now largely supplied with lumber from Michigan, Wisconsin, and Minnesota, must for building material soon depend upon the more remote pine forests of the Gulf region or those of the Pacific coast. A great development in the now comparatively unimportant lumber-manufacturing interests in these regions may therefore be expected. New centers of distribution must soon supplant Chicago as a lumber market, and new transportation routes take the place of those built to move the pine grown upon the shores of the great lakes. It is not probable, however, that any one point will ever attain the importance now possessed by Chicago as a center for lumber distribution. With the growth of the railroad system and the absence of good water communication from the great forests remaining in the country toward the center of the continent, lumber will be more generally shipped direct by rail from the mills to the consumer than in the past. In this way the pine of Mississippi, Louisiana, and Arkansas will reach Kansas, Nebraska, and the whole country now tributary to Chicago. Western Texas and northern Mexico will be supplied by rail with the pine of eastern Texas, and the prairies of Minnesota and Dakota must draw their lumber by rail, not as at present from the pine forests covering the shores of lake Superior, but from the fir and redwood forests of the Pacific coast.

FUEL.

The following table represents the consumption of forest products as fuel during the census year. The estimates of the amount and value of the wood used for domestic fuel are based upon answers to letters of inquiry addressed to persons living in every town in the United States. The average amount and value of the wood used by a family of five persons, taken as a unit, is multiplied by the number of families in each state using wood for fuel, and the result thus obtained is taken as the total state consumption:

WOOD USED AS FUEL FOR VARIOUS PURPOSES.

Use.	Cords.	Value.	Use.	Cords.	Value.
For domestic purposes as below	140,537,439	\$306,950,040	In the manufacture of brick and tile.....	1,157,522	\$3,078,331
By railroads.....	1,971,813	5,126,714	In the manufacture of salt.....	540,448	121,681
By steamboats.....	787,862	1,812,083	In the manufacture of wool.....	158,208	425,239
In mining and amalgamating the precious metals.....	358,074	2,874,568	Total.....	145,778,137	321,962,373
In other mining operations.....	266,771	673,692			

ESTIMATED CONSUMPTION OF WOOD FOR DOMESTIC PURPOSES.

Number of persons using wood for domestic fuel, 32,375,074.

States and Territories.	Cords.	Value.	States and Territories.	Cords.	Value.	States and Territories.	Cords.	Value.
Alabama.....	6,076,754	\$8,727,377	Kansas.....	2,095,439	\$7,328,723	North Carolina.....	7,434,690	\$9,019,569
Alaska.....			Kentucky.....	7,994,813	13,313,220	Ohio.....	8,191,543	16,492,574
Arizona.....	170,017	734,572	Louisiana.....	1,944,858	4,607,415	Oregon.....	482,254	1,254,511
Arkansas.....	3,922,400	5,065,821	Maine (a).....	1,215,881	4,078,137	Pennsylvania.....	7,361,992	15,067,651
California.....	1,748,062	7,693,731	Maryland.....	1,152,919	3,170,941	Rhode Island.....	154,953	706,011
Colorado.....	426,719	1,638,783	Massachusetts (a).....	890,041	4,613,263	South Carolina.....	3,670,959	11,505,997
Connecticut.....	525,639	2,371,532	Michigan.....	7,838,904	13,197,240	Tennessee.....	8,084,611	10,674,722
Dakota.....	422,948	3,028,300	Minnesota.....	1,669,568	5,873,421	Texas.....	4,883,852	10,177,311
Delaware.....	177,308	751,311	Mississippi.....	5,090,758	7,145,116	Utah.....	171,923	418,289
District of Columbia.....	26,902	80,706	Missouri.....	4,016,373	8,633,405	Vermont.....	782,338	2,509,189
Florida.....	609,046	1,230,412	Montana.....	119,947	460,638	Virginia.....	5,416,112	10,404,134
Georgia.....	5,910,045	8,279,245	Nebraska.....	908,188	3,859,843	Washington.....	184,226	499,904
Idaho.....	99,910	382,686	Nevada.....	155,276	972,712	West Virginia.....	2,241,069	3,374,791
Illinois.....	5,200,104	14,136,662	New Hampshire.....	567,719	1,964,660	Wisconsin.....	7,206,126	11,863,739
Indiana.....	7,059,874	13,334,729	New Jersey.....	642,598	2,787,216	Wyoming.....	40,213	224,848
Indian territory.....			New Mexico.....	169,946	1,063,360	Total.....	140,537,439	\$306,950,040
Iowa.....	4,090,649	14,611,280	New York.....	11,290,975	37,599,364			

(a) Including a small amount imported from Canada.

CONSUMPTION OF CHARCOAL.

Domestic and manufacturing purposes.	Bushels.	Value.
In the twenty largest cities.....	4,319,194	\$521,316
In the manufacture of iron.....	69,502,091	4,726,114
In the production of precious metals.....	97,687	29,306
Total.....	74,008,972	5,276,736

The forests of the United States, in spite of the great and increasing drains made upon them, are capable of yielding annually for many years longer a larger amount of material than has yet been drawn from them, even with our present reckless methods of forest management. The great pine forest of the north has already, it is true, suffered fatal inroads. The pine which once covered New England and New York has already disappeared. Pennsylvania is nearly stripped of her pine, which once appeared inexhaustible. The great northwestern pineries are not yet exhausted, and with newly-introduced methods, by which logs once supposed inaccessible are now profitably brought to the mills, they may be expected to increase the volume of their annual product for a few years longer in response to the growing demands of the great agricultural population fast covering the treeless midcontinental plateau. The area of pine forest, however, remaining in the great pine-producing states of Michigan, Wisconsin, and Minnesota is dangerously small in proportion to the country's consumption of white pine lumber, and the entire exhaustion of these forests in a comparatively short time is certain. The wide areas now covered in New England by a vigorous second growth of white pine, although insignificant in extent and productiveness in comparison with the forests it replaces, must not be overlooked in considering the pine supply of the country. These new forests, yielding already between two and three hundred million feet of lumber annually, are capable of great future development.

The pine belt of the south Atlantic region still contains immense quantities of timber unequalled for all purposes of construction, although unsuited to take the place of the white pine of the north. The southern pine forests, although stripped from the banks of streams flowing into the Atlantic, are practically untouched in the Gulf states, especially in those bordering the Mississippi river. These forests contain sufficient material to long supply all possible demands which can be made upon them.


The hard-wood forests of the Mississippi basin are still, in certain regions at least, important, although the best walnut, ash, cherry, and yellow poplar have been largely culled. Two great bodies of hard-wood timber, however, remain, upon which comparatively slight inroads have yet been made. The most important of these forests covers the region occupied by the southern Alleghany Mountain system, embracing southwestern Virginia, West Virginia, western North and South Carolina, and eastern Kentucky and Tennessee. Here oak unequalled in quality abounds. Walnut is still not rare, although not found in any very large continuous bodies, and cherry, yellow poplar, and other woods of commercial importance are common. The second great body of hard wood, largely oak, is found west of the Mississippi river, extending from central Missouri to western Louisiana. The forests of Michigan, especially those of the northern peninsula, still abound in considerable bodies of hard wood, principally maple. Throughout the remainder of the Atlantic region the hard-wood forests, although often covering considerable areas, have everywhere lost their best timber, and are either entirely insufficient to supply the local demand of the present population, or must soon become so.

In the Pacific region the great forests of fir which extend along the coast region of Washington territory and Oregon are still practically intact. Fire and the ax have scarcely made a perceptible impression upon this magnificent accumulation of timber. Great forests of pine still cover the California sierras through nearly their entire extent; the redwood forest of the coast, however, once, all things considered, the most important and valuable body of timber in the United States, has already suffered seriously, and many of its best and most accessible trees have been removed. This forest still contains a large amount of timber, although its extent and productive capacity has been generally exaggerated. The demand for redwood, the only real substitute for white pine produced in the forests of the United States, is rapidly increasing, and even at the present rate of consumption the commercial importance of this forest must soon disappear.

The pine forests which cover the western slopes of the northern Rocky mountains and those occupying the high plateau and inaccessible mountain ranges of central Arizona and southwestern New Mexico have not yet suffered serious damage at the hands of man. The remaining forests of the Pacific region, of little beyond local importance, are fast disappearing. The area of these interior forests is diminished every year by fire and by the demands of a careless and indifferent population; and their complete extermination is probably inevitable.


The forest wealth of the country is still undoubtedly enormous. Great as it is, however, it is not inexhaustible, and the forests of the United States, in spite of their extent, variety, and richness, in spite of the fact that the climatic conditions of a large portion of the country are peculiarly favorable to the development of forest growth, cannot always continue productive if the simplest laws of nature governing their growth are totally disregarded.

The judicious cutting of a forest in a climate like that of the Atlantic or Pacific Coast regions entails no serious or permanent loss. A crop ready for the harvest is gathered for the benefit of the community; trees which have reached their prime are cut instead of being allowed to perish naturally, and others take their place. The permanence of the forest in regions better suited for the growth of trees than for general agriculture may thus be insured. Two causes, however, are constantly at work destroying the permanence of the forests of the country and threatening their total extermination as sources of national prosperity—fire and browsing animals inflict greater permanent injury upon the forests of the country than the ax, recklessly and wastefully as it is generally used against them.



LEGEND

- I. Less than 0.1 per Cent
- II. 0.1 to 1.0
- III. 1.0 to 10
- IV. More than 10



MAP OF THE
UNITED STATES

SHOWING THE PROPORTION OF WOODLAND WITHIN THE SETTLED AREA,
BURNED OVER DURING THE CENSUS YEAR
COMPILED UNDER THE DIRECTION OF
C. S. SARGENT, SPECIAL AGENT

0 100 200 300 400 500 Miles

FOREST FIRES.

The extent of the loss which the country sustains every year from injury to woodlands by fire is enormous. An attempt was made to obtain, by means of circulars of inquiry addressed to enumerators of the census and other persons living in every town of the United States, some estimate of the actual destruction of forest material in this way. More than 30,000 of these circulars were sent out. The information obtained, often vague and unsatisfactory, after a most critical examination, in which all doubtful or contradictory returns were entirely thrown out, is presented in the following table and accompanying map. It must be borne in mind that estimates based upon information obtained in this manner are liable to very considerable error, and due allowance must therefore be made for inaccurate or incomplete returns. Many towns, and even counties, in which forest fires are known to have occurred during the year 1880, made no returns whatever, and the returns of other counties were excluded. It is therefore fair, perhaps, to assume that the following table, inaccurate and unsatisfactory as it no doubt is in many respects, at least does not exaggerate the annual loss inflicted upon the country by forest fires:

TABLE OF FOREST FIRES OCCURRING DURING THE CENSUS YEAR.

States and Territories.	Areas burned, in acres.	Value of property destroyed.	CAUSES OF FIRE.															
			Improving pas- ture.	Clearing land.	Locomotives.	Hunters.	Camp fires.	Smokers.	Malice.	Prairie fires.	Coal pits.	Lightning.	Indians.	Prospectors.	Travelers.	Spontaneous combustion.	Wood cutters.	Carelessness.
The United States	10, 274, 089	\$25, 462, 250	197	1, 152	508	628	72	85	262	12	9	82	56	10	2	2	3	3
Alabama	569, 160	121, 225	34	16	4	20	3											
Alaska																		
Arizona	10, 240	56, 000					3		2				2					
Arkansas	858, 115	259, 470		27		20						1						
California	356, 815	440, 750		9		23	23		5				4					
Colorado	113, 820	935, 500				7	10		1				5	2	2			
Connecticut																		
Dakota			4		2	2			1									
Delaware	3, 305	15, 675		6	6				2									
District of Columbia																		
Florida	105, 320	69, 900	11	2		2			3									
Georgia	705, 351	167, 620	21	15	2	16												
Idaho	21, 000	202, 000				3						2	10	6				
Illinois	48, 091	45, 775		20		27	12		3									
Indiana	90, 427	130, 335		52	20	23			4									
Indian territory	1, 000																	
Iowa	11, 017	45, 470		26	5	8				7								
Kansas	7, 080	14, 700			1		3		1	5								
Kentucky	556, 647	237, 635		51	12	33			10									
Louisiana	64, 410	6, 800	2	2		2												
Maine	35, 230	123, 315		39	14	20			3									
Maryland	41, 076	37, 425		31	16	14			5									
Massachusetts	13, 899	102, 262		40	52	37		19	8		3							
Michigan	238, 271	985, 985		161	43	59		3					1					
Minnesota	250, 805	1, 395, 110		40	13	14			9				8					
Mississippi	222, 800	78, 505	12	8	1	17			1									
Missouri	783, 646	294, 865	27	14	16	29		1	10									
Montana	88, 020	1, 128, 000			1	1		1					1	1				
Nebraska																		
Nevada	8, 710	19, 000				3							3					
New Hampshire	5, 954	63, 610		7	12	6		1	1									
New Jersey	71, 074	252, 240		7	28	6			7		6							
New Mexico	64, 034	142, 075		37	1	2			2				3			2		
New York	149, 491	1, 210, 785		37	43	22												
North Carolina	546, 102	357, 980		115	11	34	10	4	25			22						
Ohio	74, 114	797, 170		94	27	57		3	11									
Oregon	132, 320	593, 850		7		12			4				4					
Pennsylvania	685, 738	3, 043, 723		129	133	17			102									
Rhode Island																		
South Carolina	431, 730	291, 225	22	17	1	25						2						

TABLE OF FOREST FIRES OCCURRING DURING THE CENSUS YEAR—Continued.

States and Territories.	Areas burned, in acres.	Value of property destroyed.	CAUSES OF FIRE.															
			Improving pasturage.	Clearing land.	Locomotives.	Hunters.	Camp fires.	Smokers.	Malice.	Prairie fires.	Coal pits.	Lightning.	Indians.	Prospectors.	Travelers.	Spontaneous combustion.	Wood cutters.	Carelessness.
Tennessee	985,430	\$5,254,960	19	19	6	14		1	14									
Texas	599,359	273,990	19	3		7		2	16			4						
Utah	42,865	1,042,800					3						4				3	3
Vermont	3,941	48,466		10	5	2			1									
Virginia	272,319	326,944		26	13	12												
Washington	37,910	713,200		5		3			2			1	8					
West Virginia	476,775	155,280	6	22	7	13			6									
Wisconsin	406,298	725,610	20	58	13	15			3									
Wyoming	83,780	3,255,000				1							3	1				

The largest number of these fires of any one class was traced to farmers clearing land and allowing their brush fires to escape into the forest. The carelessness of hunters in leaving fires to burn in abandoned camps, next to farmers, was the cause of the greatest injury. The railroads were responsible, too, for serious damage to the forest from fires set by sparks from locomotives, while the intentional burning of herbage in the forest to improve pasturage often caused serious destruction of timber.

Only the value of the material actually destroyed by fire is included in these estimates. The loss of timber by fire, great as it is, is insignificant in comparison with the damage inflicted upon the soil itself, or with the influence of fire upon subsequent forest growth. If a forest is destroyed by fire all trees, old and young, giants ready for the ax, and germinating seedlings—the embryo forests of succeeding centuries—are swept away. Undergrowth essential to protect the early growth of trees, the roots of perennial herbage, and the seeds of all plants are consumed. The fertility, or rather the ability of the burned soil to produce again spontaneously a similar crop of trees to the one destroyed, is lost, and the subsequent recovering of burned land with the species of the original forest is only accomplished, if accomplished at all, through the restoration of fertility following the slow growth and decay of many generations of less valuable plants. A northern pine and spruce forest when destroyed by fire is succeeded by a growth of brambles, in time replaced by dwarf birch, poplar, and bird cherries, of no economic value; scrub oaks and various hard woods follow these, and pine rarely reappears except upon land long mellowed in the various operations of agriculture.

In the south Atlantic region a gradual change in the composition of the pine forests is steadily going on under the influence of fire. Less valuable species now occupy the ground once covered with forests of the long-leaved pine, through which annual fires have been allowed to run to improve the scanty pasturage they afford. Stockmen have been benefited at the expense of the permanency of the forest. Fire, too, changes the composition of the broad-leaved forests of the Atlantic region, although its influence is here less marked than upon forests of conifers, which, unlike deciduous trees, rarely grow from stump shoots, and must depend entirely upon the germination of seeds for their reproduction. Still, in regions continually burned over during a long period of time and then covered again with forests, as is the case in some portions of Kentucky and Tennessee, valuable species, like the white oak and the yellow poplar, are rare or entirely wanting in the new forest growth.

The forests of the north Pacific coast offer an exception to the law, otherwise general, for this continent at least, that a change of forest crop follows a forest fire. The fir forests of western Washington territory and Oregon when destroyed by fire are quickly replaced by a vigorous growth of the same species, and the fires which have consumed great bodies of the California redwood have not prevented the reproduction of this species by seeds and shoots. In the interior Pacific region forests destroyed by fire either do not reproduce themselves, or when, under exceptionally favorable climatic conditions, a growth of trees recovers the burned surface, poplars and scrub pines replace the more valuable species of the original forest.

The damage inflicted upon the permanency of the forests of the country by browsing animals is only surpassed by the injury which they receive from fire.

The custom of turning domestic animals into the forest to pick up a scanty and precarious living, common in all parts of the country, is universal in the southern and central portions of the Atlantic region and in California. Sheep, cattle, and horses devour immense quantities of seedling trees, the future forests of the country. They bark the trunks and destroy the vigor and often the life of larger trees. Hogs root up young pines and other plants to feed upon their succulent roots, and devour the edible fruit of many trees. In this way not only is the permanence of the forest endangered, but in the case of deciduous forests their composition is often seriously affected. Species with thin-shelled edible seeds, pines, white oaks, chestnuts, and beeches, are unable to hold their own against species with bitter or unpalatable fruit, on account of the excessive destruction of their seeds by hogs and other animals.

In the central portions of the Atlantic region the general replacement of the sweet-fruited valuable white oaks in the young forest growth by the less valuable bitter-fruited black oaks is noticeable, and seriously endangers the future value of the forests of this whole region. The damage inflicted upon the California mountain forests by sheep is immense; they threaten the complete extermination of these noble forests, and with them the entire agricultural resources of the state.

The pasturage of the forest is not only enormously expensive in the destruction of young plants and seeds, but this habit induces the burning over every year of great tracts of woodland, which would otherwise be permitted to grow up naturally, in order to hasten the early growth of spring herbage. Such fires, especially in the open pine forests of the south, do not necessarily consume the old trees. All undergrowth and seedlings are swept away, however, and not infrequently fires thus started destroy valuable bodies of timber. This is especially true, also, in the coniferous forests of the Pacific region.

The railroads of the country, using in the construction and maintenance of their permanent ways vast quantities of timber, inflict far greater injury upon the forests than is represented by the consumption of material. Railway ties, except in California, are almost invariably cut from vigorous young trees from 10 to 12 inches in diameter; that is, from trees which twenty or thirty years ago escaped destruction by fire or browsing animals, and which, if allowed to grow, would at the end of fifty or one hundred years longer afford immense quantities of valuable timber. The railroads of the United States, old and new, consume every year not far from 60,000,000 ties; the quantity of lumber in 60,000,000 ties is comparatively not very great, and would hardly be missed from our forests; but the destruction of 30,000,000 vigorous, healthy young trees, supposing that an average of two ties is cut from each tree, is a serious drain upon the forest wealth of the country and should cause grave apprehensions for the future, especially in view of the fact that in every part of the country there are now growing fewer seedling trees of species valuable for railway ties than when the trees now cut for this purpose first started.

The condition of the forests of Maine is interesting. They show that forest preservation is perfectly practicable, in the Atlantic region at least, when the importance of the forest to the community is paramount. The prosperity of this state, born of the broad forests of pine and spruce which once covered it almost uninterruptedly, was threatened by the prospective exhaustion of these forests, in danger of extermination by fire and the ill-regulated operations of the lumbermen. The very existence of the state depended upon the maintenance of the forest. The great forests of pine could not be restored, but the preservation of the few remnants of these forests was not impossible. Fires do not consume forests upon which a whole community is dependent for support, and methods for securing the continuance of such forests are soon found and readily put into execution. The forests of Maine, once considered practically exhausted, still yield largely and continuously, and the public sentiment which has made possible their protection is the one hopeful symptom in the whole country that a change of feeling in regard to forest property is gradually taking place. The experience of Maine shows that where climatic conditions are favorable to forest growth the remnants of the original forest can be preserved and new forests created as soon as the entire community finds forest preservation really essential to its material prosperity.

The production of lumber is not, however, the only function of forests; and the future extent and condition of those of the United States cannot, in every case, be safely regulated by the general law which governs the volume of other crops by the demand for them. Forests perform other and more important duties in protecting the surface of the ground and in regulating and maintaining the flow of rivers. In mountainous regions they are essential to prevent destructive torrents, and mountains cannot be stripped of their forest covering without entailing serious dangers upon the whole community. Such mountain forests exist in the United States. In northern Vermont and New Hampshire they guard the upper waters of the Connecticut and the Merrimac; in New York they insure the constant flow of the Hudson. Such forests still cover the upper slopes of the Alleghany mountains and diminish the danger of destructive floods in the valleys of the Susquehanna and the Ohio. Forests still cover the upper water-sheds of the Missouri and the Columbia, the Platte and the Rio Grande, and preserve the California valleys from burial under the *débris* of the sierras. The great mountain forests of the country still exist, often almost in their original condition. Their inaccessibility has preserved them; it cannot preserve them, however, much longer. Inroads have already been made into these forests; the ax, fire, and the destructive agency of browsing animals are now everywhere invading them. Their destruction does not mean a loss of material alone, which sooner or later can be replaced from other parts of the country; it means the ruin of great rivers for navigation and irrigation, the destruction of cities located along their banks, and the spoliation of broad areas of the richest agricultural land. These mountain forests once destroyed can only be renewed slowly and at enormous cost, and the dangers, actual and prospective, which threaten them now offer the only real cause for general alarm to be found in the present condition of the forests of the United States. Other forests may be swept away and the country will experience nothing more serious than a loss of material, which can be produced again if the price of lumber warrants the cultivation of trees as a commercial enterprise; but if the forests which control the flow of the great rivers of the country perish, the whole community will suffer widespread calamity which no precautions taken after the mischief has been done can avert or future expenditure prevent.

NORTH ATLANTIC DIVISION.

MAINE.

The forests of the Northern Pine Belt once extended over the state of Maine. Pine and spruce, with which were mingled maple, birch, and other deciduous trees, covered the entire state, with the exception of the immediate coast region between the Kennebec and the Penobscot rivers, a region of hard-wood forest; hemlock was common.


The original pine and spruce forests of the state have been practically destroyed. Pine has been cut in every township, and the largest spruce everywhere culled, except from the inaccessible region about the headwaters of the Allagash river. Scattered bodies of the original pine, often of considerable extent and generally connected with farms, exist in the southern, and especially in the southeastern, counties, and fine hemlock of large size is still an important element of the forest in the central and southern portions of the region west of the Penobscot river. Birch, maple, and oak, too heavy for transport by raft, are still common, except in the neighborhood of manufacturing centers and the lines of railroad. Hard-wood timber is particularly fine and abundant through the central portion of the state; farther north the forest is more generally composed of coniferous trees.


The lumber business of southern and central Maine attained its greatest importance as early as 1850. In that year spruce was for the first time driven down the Kennebec with pine, and the proportion of spruce to pine has since steadily increased, until, in the season of 1879-'80, only 20 per cent. of the lumber cut on that river was pine. The lowest point of productive capacity of the forests of Maine has probably been passed. The reckless disregard of forest property which characterized the early lumbering operations of the state has been replaced by sensible methods for preserving and perpetuating the forest. This change in public sentiment in regard to the forests has followed naturally the exhaustion of the forest wealth of the state. As this disappeared the importance of preserving some part, at least, of the tree covering, the source of the state's greatest prosperity, forced itself upon public attention; for unless the forests could be perpetuated, the state must lose forever all commercial and industrial importance. It has followed that the forests of Maine, as compared with those in other parts of the country, are now managed sensibly and economically. They are protected from fire principally through the force of public sentiment, and only trees above a certain size are allowed to be cut by loggers buying stumpage from the owners of land. In the southern counties the young pine now springing up freely on abandoned farming lands is carefully protected, and large areas are planted with pine in regions where the natural growth has not covered the soil. The coniferous forests, under the present management, may be cut over once in every fifteen or twenty years, producing at each cutting a crop of logs equivalent to 1,000 feet of lumber to the acre, of which from 5 to 7 per cent. is pine, the rest spruce.

Forest fires, which formerly inflicted every year serious damage upon the forests of the state, are now of comparatively rare occurrence. During the census year only 35,230 acres of woodland were reported destroyed by fire, with an estimated loss of \$123,315. These fires were set by farmers in clearing land, by careless hunters, and by sparks from locomotives.

The following estimates of the amount of pine and spruce standing in the state May 31, 1880, were prepared by Mr. Cyrus A. Packard, of Augusta, land agent of the state. They were made up from the results of actual surveys, and have been reviewed by a large number of experts most familiar with the condition of the forests in different parts of the state:

Basin of—	Pine (<i>Pinus Strobus</i>).	Spruce (<i>Picea nigra</i>).
	<i>Feet, board measure.</i>	<i>Feet, board measure.</i>
Saint John river and tributaries.....	75,000,000	1,400,000,000
Penobscot river and tributaries.....	100,000,000	1,600,000,000
Kennebec river and tributaries.....	50,000,000	1,000,000,000
Androscoggin river and tributaries.....	50,000,000	500,000,000
Saint Croix, Machias, Narragausus, and Union rivers and other small streams ..	200,000,000	500,000,000
Total	475,000,000	5,000,000,000
Cut for the census year ending May 31, 1880	188,825,000	301,020,000





DENSITY OF FORESTS

COMPILED UNDER THE DIRECTION OF
C. S. SARGENT, SPECIAL AGENT
1883.

Scale
0 1 2 3 4 5 6 7 8 9 10 MILES

Quantities of logs cut in Aroostook county are driven down the Saint John river and manufactured in New Brunswick. During the season of 1879-'80 there were handled in this way 70,000,000 feet of spruce, 4,500,000 feet of pine, 2,800,000 feet of cedar, 1,500,000 feet of squared pine timber, 1,000,000 feet of squared birch timber, 110,000 feet of squared larch timber. Of this 70 per cent. of the spruce and 80 per cent. of the pine were returned to the United States manufactured into lumber, and the whole of the cedar in the form of shingles.

Important industries dependent for material upon a supply of hard wood have long flourished in the state. Large quantities of cooperage stock, woodenware, handles, spools, bobbins, etc., are manufactured, and more recently the production of wood pulp and excelsior, principally from poplar and other soft woods, has assumed important proportions. Manufacturers from nearly every part of the state report a deterioration and scarcity of the best timber, especially oak, which is now largely imported from Canada or replaced by southern hard pine. Birch, however, is still abundant, and is largely exported in the form of spool and bobbin stock. The manufacture of potash, once an important industry of the state, has been abandoned as unprofitable. Several establishments engaged in the manufacture of tanning extracts from hemlock bark are located in the state, and the numerous tanneries upon the Penobscot river consume large quantities of the same material. The demand for hemlock lumber is now good, and the logs, after being stripped of their bark, are manufactured into lumber and not allowed, as in other parts of the country, to rot upon the ground. A recently-established industry is the manufacture of kegs, barrels, and woodenware from pulp made from chips, brush, and other waste material of the forest. Partial estimates of the hoop-pole industry give a product of 5,449,200, valued at \$75,612. During the year 1879 153,334 pounds of maple sugar were produced in the state.

ANDROSCOGGIN COUNTY.—One-half of this county is reported covered with woods, largely second growth; it contains, however, considerable bodies of fine first-growth white pine. Manufacturers of cooperage stock report oak exhausted, other hard woods scarce and of inferior quality, and that no second-growth timber is of sufficient size for use. A large amount of excelsior is manufactured, principally from poplar.

AROOSTOOK COUNTY.—Nine-tenths of this county is reported covered with forests, the clearings being confined to the neighborhoods of the rare settlements along the river bottoms. Logs cut in this county are largely rafted down the Saint John river, and little lumber in proportion to the cut is manufactured within its limits. The production of cooperage stock and other articles requiring hard wood in their manufacture is rapidly increasing, and with abundant material such industries seem destined to great development.

CUMBERLAND COUNTY.—One-half of this county is reported covered with woods, principally of second growth. Manufacturers of cooperage stock report a general deterioration and scarcity of material, especially hard woods, now nearly exhausted. Spruce and poplar in large quantities are manufactured into wood pulp.

FRANKLIN COUNTY.—Three-fourths of this county is reported covered with woods, principally confined to the northern portion. Staves, hoop-poles, handles, and excelsior are manufactured in large quantities.

HANCOCK COUNTY.—Seven-eighths of this county is reported covered with woods, largely composed, toward the coast, of second-growth white pine. The northern portions contain fine bodies of large hemlock. Manufacturers of cooperage stock report deterioration of material; ash especially has become scarce.

KENNEBEC COUNTY.—Four-tenths of this county is reported covered with woods, largely second growth. Merchantable spruce and pine have been everywhere removed. Considerable areas are again covered with pine, and the wooded area is increasing. Next to Penobscot this is the most important lumber manufacturing county in the state. Numerous mills located on the Kennebec river saw logs driven from its upper waters and from beyond the limits of the county. Large quantities of poplar and spruce are consumed annually in the manufacture of wood pulp, excelsior, handles, etc. The supply of hard wood is small and of inferior quality. The poplar now used is nearly all second growth.

KNOX COUNTY.—One-half of this county is reported covered with woods, generally of second growth. Heavy timber, however, still exists in the towns of Washington, Appleton, and Union. White pine is scarce, and great deterioration in timber of all kinds is reported. Scarcity in the near future is apprehended by manufacturers. A large amount of cord-wood is consumed annually in burning lime.

LINCOLN COUNTY.—About one-half of this county is reported covered with woods, nearly all second growth.

OXFORD COUNTY.—From one-half to two-thirds of this county is reported covered with woods. The northern portion still contains large areas of original forest, although pine and spruce have been culled everywhere. In the southern part of the county there are considerable bodies of second-growth white pine, and the wooded area is increasing. Cooperage stock, handles, and wood pulp are largely manufactured. Manufacturers report that timber of all kinds has deteriorated in quality and become scarce, with the exception of oak, which is still abundant and of good quality.

PENOBSCOT COUNTY.—Nine-tenths of this county is reported covered with woods. The merchantable pine and spruce have been removed from the southern portion and everywhere culled. In the northern townships hemlock is still abundant and of fine quality. Penobscot is the great lumber manufacturing county of the state, Bangor, once the principal market in the United States for pine lumber, being still the most important saw-mill center. Spruce and not pine, however, except in insignificant quantities, is now manufactured upon the Penobscot. Manufacturers using hard woods report an abundant supply of excellent material.

PISCATAQUIS COUNTY.—From eight- to nine-tenths of this county is reported covered with forests, the southern portion only being cleared of the original tree growth.

SAGADAHOC COUNTY.—One-half of this county is reported covered with woods, principally second growth. Considerable second-growth white pine is now growing up upon abandoned farm lands, and the wooded area of the county is increasing. Manufacturers report all timber of sufficient size for use scarce and of inferior quality, and apprehend early exhaustion of hard woods suitable for mechanical purposes.

SOMERSET COUNTY.—Five-sixths of this county is reported covered with woods, the southern portion only being cleared of its forests of spruce and pine. Excelsior, handles, woodenware, etc., are largely manufactured. Hard-wood timber of all sorts is abundant and of excellent quality, with the exception of black ash, now scarce and in great demand.

WALDO COUNTY.—From one-quarter to one-half of this county is reported covered with woods, generally of second growth. The wooded area is now gradually increasing by the growth of white pine on abandoned farming lands. Manufacturers report a scarcity and deterioration of timber of all kinds of sufficient size for use.

WASHINGTON COUNTY.—From eight- to nine-tenths of this county is reported covered with woods. In the southern portion considerable areas contain scattered bodies of large pine, and through the center of the county are large tracts of first-growth hemlock forests. No future scarcity of lumber is apprehended.

YORK COUNTY.—From one-third to one-half of this county is reported covered with woods; it contains large quantities of scattered pine. Second-growth pine is spreading on abandoned agricultural land, and the forest area is increasing. Wood pulp, cooperage stock, and handles are largely manufactured. Timber of all sorts is reported as depreciating in both quality and quantity. No immediate scarcity, however, is apprehended.

NEW HAMPSHIRE.

The forests of New Hampshire were originally composed of a belt of spruce, mixed with maple, birch, and other hard-wood trees, occupying all the northern part of the state and extending southward through the central portion; the southeastern part of the state and the region bordering the Connecticut river were covered with forests of white pine, through which considerable bodies of hard wood were scattered. The original white-pine forests of New Hampshire are practically exhausted, although in the northern counties of the state there still remain a few scattered bodies remote from streams and of small size; once of great extent and importance, these forests have disappeared before the ax of the settler and lumberman, or have been wasted by forest fires. Large areas, however, once covered with forests of pine, have grown up again, especially in the southern part of the state, with this tree. No estimate of the amount of this second-growth pine standing in the state has been possible; it furnished during the census year a cut of 99,400,000 feet of lumber, board measure. The remaining forests of the state, considered as a source of lumber supply, are composed of spruce, more or less mixed with hard woods, of which the sugar maple and the birch are the most valuable. In the northern part of the state large areas of the original spruce forest remain, although these bodies of timber are now only found at a considerable distance from streams.


Fires, which at different times have destroyed vast areas of forest, especially in the northern part of the state, are now less frequent and destructive. During the year 1880 but 5,954 acres were reported stripped of their tree covering by fires. Of such fires twelve were set by sparks from locomotives, seven by the escape into the forest of fires originally set in clearing land for agricultural purposes, six by sportsmen, one through malice, and one by the careless use of tobacco.

The basis of the following estimate of the amount of merchantable black spruce (*Picea nigra*) lumber standing May 31, 1880, in Carroll, Coos, and Grafton counties, where alone the spruce forests of the state are now of commercial importance, was furnished by Mr. G. T. Crawford, of Boston, and verified by the testimony of other experts:


BLACK SPRUCE (*Picea nigra*).

Counties.	Feet, board measure.
Carroll.....	60,000,000
Coos.....	1,000,000,000
Grafton.....	450,000,000
Total.....	1,510,000,000
Cut for the census year ending May 31, 1880 (including 26,000,000 feet sawed on the Connecticut river, in Massachusetts).	153,175,000

It is roughly estimated that the spruce forests of the state contain over 33,750,000 cords of hard wood and 165,000,000 feet of hemlock.







Julius E. Smith & Co. 1881

Scale

25 20 15 10 5 0 5 10 15 20 25 MILES

Partial returns of the hoop-pole industry give a production during the census year of 4,225,000, valued at \$29,280. New Hampshire is fourth among the states in the importance of its maple-sugar product. During the year 1879 it produced 2,731,945 pounds.

BELKNAP COUNTY.—From one-third to three-eighths of this county is reported covered with woods.

CARROLL COUNTY.—Five-eighths of this county is reported covered with woods. In the northern portion there are still large areas covered with an original growth of spruce. Large quantities of charcoal are manufactured in this county, and the usual method of lumbering adopted here and very generally in northern New Hampshire is first to cut the spruce large enough for saw-logs, taking all trees 6 inches in diameter 25 feet from the ground, and then cut for charcoal all the remaining growth, hard wood and soft, even the young spruce. As the land cleared is of little value for agricultural purposes, it is allowed to grow up again with wood. Deciduous trees come up at first, and these are sometimes, but not always, followed by spruce. It is necessary to exercise great care in order to prevent the newly-cleared tracts from suffering from fire, as the material for charcoal, cut into cord-wood, is often left on the ground until the second season. Mr. C. G. Pringle, who studied the forests of this region, furnishes the following notes upon the forests of Carroll county:

"The forests on the mountain sides between Crawford's and Bartlett are composed principally of the yellow and paper birch, the sugar maple, the red maple, poplars, the black spruce, and the balsam fir. About Bartlett scattering specimens of white pine make their appearance. In the more level part of North Conway the red and the pitch pine and the hemlock become common, while on the more sterile, sandy plains farther down the Saco these pines with the white birch constitute the principal arborescent growth.

"The tract known as Hart's location, lying partly in the White Mountain notch, includes 10,000 acres, 2,000 of which bear 15,000 feet per acre of spruce and hemlock—rather more of hemlock than of spruce; 10,000 acres in this tract will cut 25 cords of hard wood per acre. The town of Bartlett, partly cleared, still has 40,000 acres of woodland, which will yield an average of 5,000 feet per acre of spruce and hemlock and 15 cords of hard wood. Sargent's grant covers mount Crawford, Stair mountain, and a part of mount Washington. On this tract are 15,000 acres of timber-land, carrying 20,000 feet per acre, chiefly spruce. The Thompson and Meserve purchase comprises portions of mounts Washington, Jefferson, and Madison, and covers 12,000 acres. Two thousand acres of this will yield 30,000 feet of spruce and hemlock per acre in nearly equal proportions. The remaining 10,000 acres will cut 25 cords of hard wood per acre. The Bean purchase lies north of the town of Jackson, and covers 40,000 acres. It is occupied by a dense forest, amounting to 20,000 feet of spruce and hemlock and 20 cords of hard wood per acre. Originally there was considerable pine on the streams and sides of the mountains in this vicinity, particularly on mount Kearsarge, but now there is little left. Twelve and twenty-five years ago much of the town of Bartlett was burned over, and a different growth has come up—white birch, poplar, bird cherry, etc."

A large amount of cooperage stock, excelsior, and an average of 1,000 cords of shoe pegs (from birch and maple) are annually made in this county. Considerable damage to oak and poplar caused by the ravages of the army-worm [?] are reported. The natural increase of timber is said, however, nearly to equal the present consumption by local industries, and scarcity is not apprehended.

CHESHIRE COUNTY.—About one-half of this county is reported covered with woods.

COOS COUNTY.—Nine-tenths of this county is reported covered with forests. The following is extracted from Mr. Pringle's notes upon the forests of this county:

"Everything east of the Connecticut lakes and about the upper portions of Indian and Perry streams is original forest. Such also is the condition of the Gilmanton, Atkinson, and Dartmouth College grants and the towns of Dixville, Odell, and Kilkenny. All the eastern portions of Clarksville, Stewartstown, Colebrook, Columbia, and Stratford are forest, and nearly all of Wentworth's location, Millsfield, Errol, Dummer, Cambridge, and Success. In these forests the spruce will cut 5,000 feet and the hard wood about 50 cords per acre. There is considerable hemlock, but even less pine than in Essex county, Vermont. Not much of the region has been burned over, and spruce comes into the soil again but slowly after clearings and fires.

"In the township of Kilkenny, in the mountains east of Lancaster, there are 16,000 acres of forest still untouched, though a branch railroad from Lancaster into this forest has been surveyed, and may be constructed in a few years, for the purpose of bringing the lumber down to the mills at Lancaster. Lowe and Burbank's grant is a wilderness, three-fourths well timbered and the remainder a mountain ridge of nearly bare rock. Bean's purchase is nearly inaccessible and but little lumbered. Stark, on the upper Ammonoosuc, is badly cut over, only about one-quarter remaining in virgin forest. About one-half of Berlin is uncut; also the northern half of Randolph, the south half of Gorham, and the south quarter of Shelburne. Considerable land in Success was burned over some years ago, as well as some in Stark and in the eastern part of Berlin, but fires have not lately been very destructive in the New Hampshire forests."

A large amount of cooperage stock, handles, wood pulp, shoe pegs, etc., is manufactured in this county. Abundant material, with the exception of ash, is reported.

GRAFTON COUNTY.—One-half of this county is reported covered with woods, mostly confined to the northern and central portions. Shoe pegs, cooperage stock, wood pulp, and excelsior are largely manufactured. The amount of material is considered abundant for the present consumption.

HILLSBOROUGH COUNTY.—One-half of this county is reported covered with woods, mostly second growth. A large amount of cooperage and wheel stock is manufactured. No deterioration in the quality of material is reported, although at the present rate of consumption it must soon become exhausted.

MERRIMACK COUNTY.—One-half of this county is reported covered with woods. Cooperage stock, handles, and excelsior are largely manufactured. A slight deterioration in the quality of material is reported.

ROCKINGHAM COUNTY.—From one-quarter to five-eighths of this county is reported covered with woods, mostly second growth.

STRAFFORD COUNTY.—Four-tenths of this county is reported covered with woods, mostly second growth. Hoop-poles, cooperage stock, etc., are largely manufactured. Wood of all sorts is reported scarce and rapidly increasing in value.

VERMONT.

The forests of Vermont, as compared with those of New Hampshire and Maine, are varied in composition. About the shores of lake Champlain several western trees first appear, and throughout the state the forest is more generally composed of deciduous than coniferous species. Forests of spruce, however, spread over the high ridges of the Green mountains, their foot-hills being covered with hard-wood trees and little pine or hemlock occurring in the valleys. A forest of white pine once stretched along the banks of the Connecticut, and great bodies of this tree occurred in the northwestern part of the state, adjacent to lake Champlain. The original white-pine forests of the state are now practically exhausted. They are represented by a small amount of second-growth pine only, which furnished during the census year a cut of 6,505,000 feet of lumber, board measure.

The forests of Vermont now suffer comparatively little from fire, although at different periods during the last fifty years very serious fires have laid waste great areas of forest in the Green Mountain region. During the year 1880 3,941 acres of woodland were reported destroyed by fire, with an estimated loss of \$48,466. Of such fires ten escaped from farms into the forest, five were set by locomotives, two were traced to the carelessness of hunters, and one to malice.

Large amounts of cooperage stock, woodenware, furniture, paper-pulp, excelsior, veneers, etc., are manufactured throughout the state. Material for these industries is fast disappearing, and a great deterioration in quality, especially of oak, ash, and chestnut, is reported by manufacturers.

Vermont surpasses all other states in the manufacture of maple sugar. During the year 1879 11,261,077 pounds were produced in the state.

The following estimate of the spruce standing in the state May 31, 1880, has been prepared from Mr. Pringle's report, and is based upon the statements of numerous timber-land owners and experts in different parts of the state:

BLACK SPRUCE (*Picea nigra*).

Regions.	Feet, board measure.
Green Mountain range.....	380,000,000
Valley of the Connecticut river.....	375,000,000
Total	755,000,000
Cut for the census year ending May 31, 1880 (excluding 16,191,000 feet imported from Canada).	199,086,000

Partial returns of the hoop-pole industry give a production during the census year of only 43,900, valued at \$470.

ADDISON COUNTY.—About one-third of this county is reported covered with woods. Spruce and ash are scarce and rapidly disappearing. Oak of sufficient size for the manufacture of cooperage stock is exhausted.

BENNINGTON COUNTY.—Two-thirds of this county is reported covered with woods. Manufacturers of woodenware and cooperage stock consider the prospects for future local supply favorable.

CALEDONIA COUNTY.—From one-third to three-eighths of this county is reported covered with woods, mostly confined to the northern and western portions.

CHITTENDEN COUNTY.—About one-fifth of this county is reported as woodland. The following extracts are made from Mr. Pringle's note upon the forests of Vermont:

"Except on the summits of a few of the higher peaks of the Green mountains, where black spruce and balsam fir grow to the exclusion of other trees, the arboreal growth is composed of a large number of species. In the valleys and on the foot-hills, and even on the slopes of the higher mountains in their lower portions, hemlocks mingle with spruce, beech, maple, and birch (yellow birch chiefly, for there is little white birch seen in northern Vermont); basswood, butternut, the ashes, red oaks, etc., are confined to the lower elevations and are less abundant than the trees first mentioned. Between the isolated patches of spruce and fir about the summits of the mountains and the region where hemlock is found, rock maple, yellow birch, and black spruce are the predominating species.

"To estimate the area of valuable original forest still standing in the Green mountains is not an easy task. The belt extends from the Canada line to Massachusetts, and even into that state. The outlines of this belt are made very irregular by the cleared and settled valleys which run up among the mountains, and by reason of forest clearings, so that its width is constantly varying as we proceed from one end to the other.

"The woodlands of the plateau, some 10 miles broad and elevated from 200 to 300 feet above lake Champlain, lying between the foot-hills of the Green mountains and the lower plain beside the lake, occupy, for the most part, rocky hills, and are composed principally of sugar maple, beech, basswood, white ash, black birch, and red oak. Certain limestone hills offer a favorable situation for the butternut, the ironwood, the slippery elm, and the bitter hickory. The swamps and other lowlands yield the red maple, the black ash, the white elm, and the black willow. The latter, especially along streams, is associated with alders and the sheepberry. The colder, sphagnous swamps are covered with a growth, more or less dense, of yellow cedar, black spruce, balsam, and larch; sometimes in the higher portions the white pine mingles with these, scattered or in groves. When grown in such soil this wood is liable to be extremely hard and brittle. The poplars occupy hillsides and ridges where the soil is a light, cold, sandy loam; with them the bird cherry is perpetually associated. The black cherry is scattered in a diversity of soils. White oak and hickory attain their best development on clayey soil or glades of slight elevation; on the red sand-rock hills they are smaller. Certain slopes of cold clay are still here heavily wooded with hemlock, while warm clay lands are the favored site of the burr oak. In the vicinity of the lake and its tributaries low, wet shores are scattered over with the swamp white oak and the burr oak. The chestnut oak is common on the thin, poor soil of the red sand-rock hills, ranging through the valley from the lake as far back in some places as the foot-hills of the Green mountains. The red pine appears on the sandy shores of lake Champlain, and extends far up the Winooski river. The moister and more fertile portions of the sandy plain are still occupied to some extent by white pine, the poorer portions by pitch pine. The white birch occurs on cold, wet, sandy soil near the lake; and in the mountains the black spruce becomes the most common tree; with it in stronger soil are associated the yellow birch and the sugar maple.

"*Burlington.*—This place is believed to rank as third, or next to Albany, among the lumber markets of the United States. More lumber may enter some ports, as Oswego and Tonawanda, for transshipment, but all lumber brought to this market is stored and sold here. The kind is chiefly white pine brought up the lake from Canada, a little of it being cut in Michigan (perhaps one-tenth); all the rest is of Canadian growth. A few of the lumber companies here own lands of limited extent among the Green mountains, from which they obtain spruce for clapboards, etc. The general direction which the lumber sent from here takes is to the older portions of New England, Massachusetts, Rhode Island, and Connecticut, considerable pine being sent even to Maine, which once supplied to commerce so much of this material. Much lumber is dressed here and sent to Boston for shipment to foreign countries. The business still enjoys the highest prosperity, and during the census year, under the stimulus of general commercial prosperity, it was especially active. As yet no lack in the supply is felt, the loggers only having to go farther back in the Canadian forests than formerly to obtain timber enough to meet the demand. The proportion of lumber worked up here is small, there being merely a few factories producing doors, sash, blinds, packing boxes, etc."

ESSEX COUNTY.—Five-sixths of this county is reported covered with forest. The following is extracted from Mr. Pringle's report:

"Four-fifths of that part of the county of Essex lying north of Guildhall and Victory is still in virgin forest, which will yield 5,000 feet of spruce per acre. The towns of Lewis and Averill are entirely unlogged, and so is Avery's Gore. Colton is mostly covered with forest, and so is Ferdinand. Timber-lands compose about two-thirds of Granby and East Haven, and cover the back parts of the river towns and those crossed by the Grand Trunk railroad. South of Guildhall and Victory the towns of Concord and Lunenburg are mostly cleared and settled. The proportion of hemlock in these forests is not large; there is considerable yellow cedar and a large amount of maple, birch, and beech—probably 50 cords per acre. There is but little pine in all this region, principally confined to the township of Lewis; elsewhere only occasional pine trees occur."

FRANKLIN COUNTY.—From one-fourth to three-tenths of this county is reported covered with forest, mostly confined to the hills in the northeastern and northern portions. In the village of Montgomery a large establishment for the manufacture of butter tubs is located, and at East Richford birch is largely manufactured into turned ware.

GRAND ISLE COUNTY.—About a quarter of this county is reported covered with woods.

LAMOILLE COUNTY.—About one-third to one-half of this county is reported covered with woods, very generally distributed over its entire surface.

ORANGE COUNTY.—One-quarter of this county is reported covered with forest.

ORLEANS COUNTY.—One-half of this county is reported covered with woods. The following is extracted from Mr. Pringle's notes:

"At Newport, situated at the southern extremity of lake Memphremagog, are several mills for cutting veneering from birch. The product of these mills is closely packed in boxes, so that it cannot warp, and sent to the manufactories near the large cities, to be used for chair bottoms and other purposes. Southward from Newport, in the valleys of the Barton and Black rivers, which flow northward into lake Memphremagog, and of the Passumpsic river, which runs southward and joins the Connecticut, are almost continuous swamps of yellow

cedar, black spruce, and larch, from which the cedar timber is now being largely drawn to be sawed into shingles. At Barton the hard woods are largely cut into material for furniture, which is shipped toward the sea-board before being put together.

"The valley of the Clyde river from Newport to Island Pond is cleared for the most part and improved for farms. The usual species of the northern forest occupy the summits of the low hills on either side of the valley. Eastward from Island Pond, down the Neipegan river to the Connecticut by the line of the Grand Trunk railroad, we pass through the wild region from which the lumbermen have only taken some of the spruce and pine. Here, beginning 2 or 3 miles back from the railroad, or in some places much nearer to it, a virgin and unbroken forest stretches over the slopes and summits of the hills for many miles to the northward and southward; black spruce, yellow birch, sugar maple, and beech are its chief component species. In a few places, where the soil is sandy, white pine occurs in straggling groves or isolated specimens, and the swamps, as well as those of all of northern Vermont, are occupied by the black spruce, yellow cedar, and by a few scattering pines. The pine being the kind of lumber first secured, is seldom found now in these Vermont swamps. The cedars are now cut and manufactured into shingles, fence posts, railway ties, etc., for which purposes the lasting quality of the wood makes it eminently suited. There is little hemlock in northeastern Vermont, and it is believed to indicate poor soil wherever it occurs. The soil of this entire region presents a marked contrast to that of northern New York, being fertile and in other respects well adapted to agriculture. On this account land once lumbered over is generally occupied by the farmer and not allowed to come up again to forest, except in the more hilly portions."

Staves, tubs, pails, buckets, and hoops are largely manufactured from spruce, cedar, and ash. The quality of the material used is said to have deteriorated, and manufacturers report that at the present rate of consumption it will soon be consumed.

RUTLAND COUNTY.—Four-tenths of this county is reported covered with woods, principally in the eastern portion. Elm, formerly largely used in manufacture of tubs, etc., is reported exhausted, and basswood has become scarce.

WASHINGTON COUNTY.—One-third of this county is reported covered with woods, principally situated in belts along its eastern and western borders. The following is extracted from Mr. Pringle's report:

"Reaching Montpelier from the west we have left behind the Green Mountain gneiss and entered a granitic formation. Here is an extensive burned region; the fire, in consuming the forest and vegetable mold upon the surface of the land, has exposed granite boulders thickly imbedded in the soil. To replace the forest growth thus removed there is only an occasional little spruce or balsam to be found among the thickets of bird cherry. The hilltop and hillside forests east of Montpelier show hemlocks everywhere mingled with sugar maples, yellow birches, and spruce; farther east the spruce and birch predominate. Approaching the Connecticut river, hemlocks and maples again appear and second-growth white pine and paper birches take the place of the other species."

WINDHAM COUNTY.—Three-eighths of this county is reported covered with woods, mostly confined to ridges of the Green mountains. Ash and white pine are reported very scarce.

WINDSOR COUNTY.—From one-fourth to one-third of this county is reported covered with woods, quite generally distributed over the hills. Tubbs, barrels, kegs, and buckets of white and red oak, white pine, spruce, and ash are manufactured. Oak is reported by manufacturers to be already practically exhausted, spruce to be fast disappearing, and ash very scarce and in danger of speedy extermination.

MASSACHUSETTS, RHODE ISLAND, AND CONNECTICUT.

The original forest which once covered these states has disappeared and been replaced by a second, and sometimes by a third and fourth growth of the trees of the Northern Pine Belt. The area covered by tree growth in these states is slowly increasing, although, with the exception of the young forests of white pine, the productive capacity of their woodlands is, in view of the heavy demands continually made upon them, especially by the railroads, rapidly diminishing. Abandoned farming land, if protected from fire and browsing animals, is now very generally, except in the immediate vicinity of the coast, soon covered with a vigorous growth of white pine. The fact is important, for this new growth of pine promises to give in the future more than local importance to the forests of this region.

These states sustain a considerable annual loss from forest fires. In Massachusetts during the year 1880 13,899 acres of woodland were reported destroyed by fire, with a loss of \$102,262. Of these fires fifty-two were set by locomotives, forty by fires started on farms and escaping to the forest, thirty-seven by hunters, nineteen by the careless use of tobacco, eight through malice, and three by carelessness in the manufacture of charcoal. No returns in regard to forest fires in Rhode Island and Connecticut have been received, but it is believed that in proportion to their forest area such fires are not less destructive in these states than in Massachusetts. Numerous important industries using hard wood have been driven from these states or forced to obtain their material from beyond their limits. On the other hand, industries like the manufacture of certain sorts of woodenware, using second-growth pine, are rapidly increasing in volume. The principal forests now found in these states are situated in Berkshire, Hampden, and Worcester counties, Massachusetts.

BERKSHIRE COUNTY, MASSACHUSETTS.—From one-third to one-half of this county is reported covered with woods, largely second growth. The high ridges of the hills are still covered with forests of black spruce, their slopes and intervening valleys with hard woods or hemlock, now often replaced by a growth of young white pine. Cooperage stock, baskets, and wood pulp are largely manufactured. Spruce is reported to have deteriorated in quality; manufacturers consider the supply of material, however, abundant for all present local demands.

FRANKLIN COUNTY, MASSACHUSETTS.—One-half of this county is reported covered with woods, largely second-growth white pine.

WORCESTER COUNTY, MASSACHUSETTS.—One-half of this county is reported covered with woods, largely second-growth white pine. Winchendon, the most important point in the United States for the manufacture of woodenware, small cooperage, etc., is supplied with material from the young pine forests of this and the neighboring counties. Timber is reported to have deteriorated. The supply of pine is not equal to the demand, and is rapidly increasing in value.

In Barnstable county, Massachusetts, numerous experiments in forest planting have been made. In South Orleans and neighboring towns fully 10,000 acres of sandy, barren soil have been successfully and profitably planted with pitch pine. Similar plantations have been made upon the island of Nantucket; and many large groves of white pine planted many years ago in Bristol and Plymouth counties demonstrate the entire practicability of forest culture in this whole region.

The only important lumber manufacturing establishments found in these states are situated upon the Connecticut river, in Massachusetts and Connecticut. They are entirely supplied with material from the forests of northern New Hampshire and Vermont. Partial returns of the hoop-pole industry give a production during the census year in Massachusetts of 11,507,600, valued at \$95,009; in Connecticut, of 191,000, valued at \$9,660.

NEW YORK.

That portion of the state north of the forty-third degree of latitude, including within its limits the elevated Adirondack region, was once covered with a dense forest of maple, birch, basswood, and other northern deciduous trees, through which were scattered spruce and pine. The low hills bordering the Hudson and extending along the southern boundary of the state west of that river were covered with the coniferous species of the Northern Pine Belt. Over the remainder of the state the broad-leaved forests of the Mississippi basin spread almost uninterruptedly, except where an occasional sandy plain or high elevation favored the growth of pines. The original forest still covers large areas in the northern counties, and protects the hills through which the Delaware river forces its way in crossing the southern part of the state. With these exceptions, however, the forests of New York are now almost exclusively of second growth.

The forests of the state, especially in the north, have at different times suffered great damage from fire. During the census year 149,491 acres of woodland were reported destroyed by fire, with a loss of \$1,210,785. Of these fires thirty-seven were set by farmers clearing land for agricultural purposes and allowing them to escape to the forest, forty-three were set by locomotives, and twenty-two by the carelessness of sportsmen.

With the exception of the spruce of the Adirondack region, the forests of the state are no longer important as a source of general lumber supply; and many industries depending upon hard woods have in late years decreased in importance, owing to the want of sufficient material, or have been forced to obtain their supply of timber from the west. White oak, largely consumed by the railroads, has become scarce, and has advanced at least 50 per cent. in value during the last twelve years. Elm, ash, hickory, and other woods are reported scarce in all parts of the state. Partial returns of the hoop-pole industry give a production during the census year of 10,948,258, valued at \$155,764.

New York is only surpassed by Vermont in the amount of maple sugar produced by its forests. During the year 1879 10,693,619 pounds were manufactured in the state.

The following extracts are taken from Mr. Pringle's report upon the forests of northern New York:

"One who enters northeastern New York at Port Kent, and takes stage by way of Keeseville to the Saranac lakes, finds himself, as long as his route runs up the Au Sable river, which is as far as the Au Sable forks, passing through a region which gives evidence of having been formerly covered with pine. The white, the red, and the pitch pine are all represented here. The pitch pine is confined chiefly to the sterile sandy plains between the Au Sable and the Saranac rivers. The red pine mingles with this species, and grows on the rocky hills of the region and on the river cliffs, while the abundance of white pine in nearly all situations must have made this quarter of the state, like the region of Vermont lying opposite, a valuable pinery in former times. But fifty or seventy-five years have passed since the pine of the Champlain valley was harvested and shipped to England by way of the Saint Lawrence.

"In the valleys of the Au Sable and the Saranac rivers white pines spring up numerously whenever permitted to do so, and I am told that farmers, realizing that much of their soil is not suitable for profitable agriculture, are seriously considering whether it be not to their highest advantage to surrender much of their land to timber growing, and encourage the growth of the more valuable species, such as white pine, white oak, etc. Of non-coniferous trees

the white, red, and black oaks are conspicuous among the pines, and in the colder and wetter sands the white birch is common. But through all this region the trees are all of second growth, and lumber for building purposes is largely imported.

"The forest on the upper waters of the Au Sable and of the divide between this river and the Saranac is principally devoted to supplying fuel to numerous iron furnaces. The best butt logs only of spruce are sorted out and sent to the saw-mills as the forests are mowed down; the hemlock bark is removed for the tanneries, but everything else, young pine, spruce, and poplar, fall clean with maple and birch. Here and there, even far up on the hillsides, are seen the charcoal kilns, and around and about them, quite to the crest of the foot-hills of the Adirondacks, the woods are cut down in great swaths to feed them. Lands once cut over are left to grow up to timber again, though fires originating in the dead brushwood and consuming the sun-dried vegetable mold on the surface of the soil generally interfere with any new growth of trees.

"Little Tupper lake is situated in the heart of the Adirondack wilderness, and is surrounded by some of the most valuable timberlands to be found in all this region. The woods about the lake have never heard the lumberman's ax. The stream which connects it with Tupper lake, by way of Round pond, is not adapted to driving, and before lumber could be brought down it would be necessary to clear out the stream by blasting away much rock and building a dam with flood-gates at the foot of Round pond. The shores of this beautiful lake present a marked contrast to those of any I have as yet visited. On other shores and river banks I had seen scattering pines, but on all the points and bluffs of this lake throughout its entire circuit, and even following the ravines far back in the hills, are great groves and belts of white pine with straight and clean shafts towering high above all other trees, unless is excepted the red pine, of which a few specimens are mingled with them on the gravelly banks of the lake, vying with the white pines in height and beauty of trunk. At certain places on the shores of this lake, and particularly along the sluggish streams connecting it with Round pond below, are considerable swamps occupied chiefly by larch. It is pleasing to observe and to learn from guides that this lake region of the Adirondack woods has suffered but little from forest fires. It is only limited areas here and there on the shores of the lakes and ponds or along the rivers that have been devastated by fires originally started in hunters' camps. Seldom do these fires spread far back from the water, a fact which is to be attributed, it is believed, to the wet and mossy condition of these woods; yet, when they have been lumbered, as is the case lower down the Racket river, and a considerable proportion of the trees have been removed so as to expose the brushwood, etc., to the drying influences of the sun, much the usual liability to fire exists here.

"It is safe to assume that 2,500 square miles fairly represent the area of the virgin forests of the Adirondack wilderness. This area will average 3,000 feet of spruce (board measure) per acre, or about five billion feet in the aggregate. The amount of hemlock, variously estimated from 300 to 10,000 feet per acre, will cut at least 2,000 feet per acre, or 3,000,000,000 feet in the aggregate, or its equivalent; when the bark alone is considered, 3,000,000 cords of bark. The pine hardly, if at all, exceeds 200 feet per acre, or 320,000,000 feet in all. The hard wood growing over this entire region will fairly average 40 cords per acre, or 64,000,000 cords.

"Glens Falls is the great sawing center for the lumber cut upon the upper Hudson. This business here has passed the point of maximum prosperity and begun to decline; not that there was any necessity for a diminution of the yearly crop of logs from this field, if the forest could be protected from devastating fires. The lumberman leaves standing, as far as possible, the spruce trees too small for the ax, and these, the overshadowing growth being removed, grow with increased vigor, so that good crops of timber could be harvested from the soil every thirty or forty years, were it not that over at least one-half of the area lumbered fire follows the ax, burning deep into the woody soil and inducing an entire change of tree covering. Poplars, birches, and bird cherries, if anything, succeed the spruces and firs. From this cause alone the lumbering industry of the region must dwindle. A large area utterly unadapted to agriculture is being made desolate and nearly valueless, and its streams, the feeders of the water privileges and canals below, become every year more and more slender and fitful. These fires are largely set by reckless sportsmen and hunters, with whom this region peculiarly abounds in summer. They are careless in their smoking; they neglect to watch and properly extinguish the fires lighted for camp and cooking purposes, and sometimes they even delight to set fire to the dry brushwood of lumbered land in lawless sport. Again, to some extent, a class of petty pioneers follow the lumberman, obtaining for a trifling sum a title to a little land, or, squatting without rights, set fire to the dry brushwood left by the lumberers, and allow the fire to spread at will, devastating thousands of dollars' worth of property for the mere convenience of saving themselves the trouble of burning boundary strips around their fields, which might not cost them labor to the amount of \$10. The laws of New York in respect to the setting of forest fires are totally inadequate to protect the forests. The opinion prevails in the forest region of northern New York that a growth of trees removed is followed by a similar growth, the result of young seedling trees left in the soil, except in the case of pine. 'Pine once cleared off is never renewed,' was the invariable remark. This of course presumes that fire is kept out of the clearing, for after a fire has consumed the brushwood and much of the 'duff' or vegetable mold, and with this all the young seedling trees, and even the seeds of trees that may be in the soil, an entirely different growth from the hemlock and spruce springs up. Raspberry bushes are the first to appear, the seeds of which are dropped by birds flying over the clearing. Bird cherries generally appear among the first trees, the seeds being dropped everywhere in a new country by birds;

poplars and small willows also appear early in a burned district, their downy seeds being widely distributed by the wind. It is only through the agency of the wind that the seeds of birches and conifers can be disseminated, and spruces and hemlocks must needs appear, if they return at all, as tardy stragglers.

"Not many miles above Glens Falls the Hudson flows out from among the lowest outposts of the Adirondacks and winds through a plain which reaches from near Troy to the vicinity of the southern ends of lakes George and Champlain. The soil of this plain is sand deposited by the waters of former periods. The hills which bound this plain on the northwest are piles of sand, gravel, and boulders, evidently the moraines of a glacier which once flowed through the course of the Hudson. All this region, from Troy to Luzerne, among the foot-hills of the Adirondacks, must formerly have been covered with pine; among the hills and near the streams white pine, and in the more sterile central portions of the plain, red and pitch pine. To-day there exists of these species scarcely more than a scanty and scattered second growth.

"Thirty or forty years ago it was thought that all the accessible spruce in the valley of the upper Hudson had been harvested, but there is to-day nearly as much sawed at Glens Falls as there was at that time. At that time nearly all the timber standing near this river and its larger tributaries had been cut. Such as stood 5 or 10 miles back from these streams and all that was growing in the valleys of the smaller streams, or higher up the mountain slopes, would not pay the cost of hauling to the larger streams; but it is this timber which now furnishes the present supply. Logs are now driven out of streams which were then thought incapable of being driven. By damming streams so small that they may almost dry up in midsummer, throwing the logs into their courses during the winter, either above or below the dams, and in spring-time, when the dams are pouring with the floods resulting from the melting of deep mountain snows, tipping the planks of the dams and letting loose the torrents, the logs from remote places are got out to the large rivers where they can be driven. All the rivers of this region, however, are steep and rocky. The logs come down with their ends badly battered, and often with gravel and fragments of rock driven into the ends in a manner to injure the saws. They must, therefore, be 'butted' before being sawed; that is, a thin section is cut from each end, and on this account the logs are cut in the woods 4 inches or, for the worst streams, 6 or more inches longer than the standard length. The standard length for all logs brought down the Hudson is 13 feet. The character of these streams is such that long logs, for spars or other purposes, cannot safely be driven through them. Such sticks are certain to get fastened among rocks and cause bad jams. As already stated, the lumber business upon the upper Hudson is well advanced in its decline, and a score of years hence it must become insignificant under the practices now pursued, and the future of this valley gives little promise of prosperity; the soil is inferior in quality and not adapted to agriculture, while the timber, once the chief source of its prosperity, is nearly exhausted.

"As a lumber market Albany ranks second in the United States, or next to Chicago. White pine is the variety of lumber most largely handled here, and two-thirds of it comes from Michigan by way of the Erie canal, the remaining one-third coming from Canada through lake Champlain, the white pine contributed by New York being an inappreciable quantity. Most of the lumber firms here are merely commission dealers, although in two large mills considerable lumber is dressed before being shipped. The region supplied by this market includes the banks of the Hudson, New York city, New Jersey, and the shores of Long Island sound. A little reaches Philadelphia, and much is shipped to foreign ports from the city of New York. A great deal of the lumber handled by Albany dealers, however, does not go to Albany at all, but, sold by runners, is sent direct by railroad from the Michigan mills to points south of New York. The lumber trade here is still in full prosperity.

"Leaving the beautiful Mohawk valley at Rome, the traveler by the Rome and Watertown railroad soon notes a less improved region, and one, indeed, less capable of improvement. For a long time the road stretches over a sandy plain; in the higher portions of this plain, not far from Rome, the red and pitch pines are seen, and in the wetter places hemlocks and black spruces appear, with white birch, black ash, etc. On the higher, undulating lands, 20 or 30 miles north of Rome, white pine and hemlock seem once to have been the most abundant species of the forest; they now exist only in broken and scattered ranks, although numerous stumps give evidence of a former heavy growth of these two species. Northward from Albion the country gradually rises, hard wood becoming more and more common until on the limestone banks of the Black river at Watertown the patches of woodland are mainly composed of birch and maple. Yet the soil continues sandy, and at a little distance from the river is favorable to the growth of pine, and I can readily believe that all this sandy tract east of lake Ontario was originally covered with a heavy growth, principally of pine and hemlock. The pine was long since harvested, and now the mills and tanneries are consuming the hemlock. On each of the small streams that flow into lake Ontario are established saw-mills which cut quantities of hemlock yearly. Little, however, is sawed at Watertown, although a limited amount of logs is driven down to Dexter at the mouth of the Black river, and there sawed; yet once the neighborhood of Watertown and Dexter was a great center for the production of pine lumber. This region (chiefly its swamps) still yields a little black spruce. The lumber sawed along the Rome and Watertown railroad at Williamstown, Richmond, etc., is mostly sent southward to Syracuse and other places to meet the demand there for coarse lumber. The lumber yards at Watertown are mostly filled with Canadian pine.

"Carthage, in Jefferson county, was once an important lumber center. The 'Long falls' of the Black river furnished unlimited water-power. Immense quantities of pine and hemlock lined the banks of the river and covered

the plains of the vicinity; northward lay a heavy pinery. Canal-boats laden with lumber were towed through the river to Lyon's falls and thence by canal to Utica. Now the pine is nearly all gone from this region, the saw-mills are rotting down and only a little hemlock is sawed here.

"That portion of the state which lies along the Saint Lawrence river as far east as the vicinity of Malone, and extending some 25 miles back from the river, seldom exceeds 250 feet above the sea-level and is, for the most part, clayey loam, flat and well adapted to agriculture. This tract is now pretty well settled. Proceeding to the southeastward and rising to an altitude of 250 feet a wide region of sandy soil is entered, cold, damp, and unfit for agricultural purposes. This is the region of forest lying northwestward of the mountains in the southern portions of Saint Lawrence and Franklin counties, and has not yet been badly encroached upon by the ax and fire. The destruction of this forest would be a public calamity, so useless is the soil for any other purpose than the production of timber, and so harmful to the settled country below would be the consequences resulting from clearing it. This forest is, no doubt, capable of yielding, perpetually, an annual crop double that now drawn from it. This estimate, of course, is based upon the supposition that fires are prevented. But this side of the forest is less invaded by fires than the valley of the Hudson river, and fires do not burn so deeply into the soil nor consume so much of the vegetable matter; they are, consequently, less fatal to the continuance of timber growth.

"At Canton, in Saint Lawrence county, and in its vicinity as far down as Buck's bridge, below Morley, is sawed all the lumber cut on the Grass river. From this point the lumber is shipped principally to Massachusetts and Connecticut by rail, both via Rome and via Plattsburgh and Rouse's Point.

"Colonel Colton, of Norwood upon the Racket river, explained to me at length the methods employed by him in the lumber business, and, as nearly the same methods are pursued throughout this region, I give his account. Several weeks of the summer he devotes to exploring the lands of his company, to decide from what tract the stock of logs for the following year shall be drawn. In the settlements near the margin of the forest are men whose business it is to cut and haul onto the ice of the river during winter the timber desired by the lumber companies. Contracts are made with these men to harvest the timber above a certain diameter on certain specified tracts belonging to the company. The contractors go to their respective fields of labor as soon as the snow is of sufficient depth, taking into the woods a force of men, horses, and supplies, and building camps in the vicinity of their work. When a full stock of logs is placed on the river, and the spring floods break up the ice and set the logs going, other contracts are made with the same or other men to drive the logs into the booms of the different mills at a stipulated price per log. If, as is usually the case, logs of several different companies are on the same river, all are driven down in common, and the drive is called a 'union drive'. Arrived at the uppermost boom—formed by chaining together logs floating on the surface of the water and held in place by occasional piers, strong but rude structures of logs filled in with rocks, located above the first sawing station—the logs belonging to these mills are sorted out and turned into the different booms, while those belonging below are sent on their way down the channel. Once within the boom of the mills to which they belong, they are again assorted; the pine, hemlock, and the spruce are separated, and the different grades are floated into separate booms or pockets which lead down to the different mills or saws which are to cut up each separate class. At the mills inclined planes lead down to the water from each gang of saws, up which, chains being attached to the logs, they are drawn by the machinery into the mill. After sawing, the sorting of the lumber into different grades is completed with care. The boards are run through planing-mills which smooth both sides, then through other machines which tongue and groove their edges, and finally fine saws neatly trim their ends. This dressing of the lumber at the mills makes a saving in freight when it is shipped, besides greatly facilitating sales. Colonel Colton invited me to accompany him 20 or 30 miles up the river to see the 'drive' which was just coming out of the woods. The highway by which we drove led near the river, and we could see the logs everywhere coming down, advancing endwise with the current. In many places of still water the entire breadth of the river for some distance was closely covered with them. These were not so small as those usually seen in the Maine rivers, but were from full-grown trees of the original forest—spruce from 1 foot to 2 feet in diameter. With the spruce logs were a few hemlocks, usually of larger size; a few pine logs, sometimes 2 or 3 feet in diameter, floated with the others. As the water was lowering, stranded logs were seen everywhere along the shore. They covered gravel banks and bars in the middle of the river, and were piled in disorder on the rocks of the rapids, or, pushing over the waterfalls, stood on end in the midst of the white, pouring torrent.

"A few miles above Potsdam we entered upon a sandy soil; the farms appeared less productive and the farm-buildings and fences gave evidence of less thrift. As we advanced toward Colton, a region near the borders of the forest some twenty years settled, less and less prosperity among the settlers was manifest. The tilled fields appeared incapable of yielding even passably good crops; some of them could do no more than give a small crop of rye once in three years. The grass lands were red with sorrel, which comes up everywhere over this region as soon as the forest is cleared and the ground burned over. The sandy soil is cold and sour, in some places so light as to be blown about by the wind. Above South Colton we drove over sandy plains utterly incapable of sustaining the meager population, which ekes out a wretched existence by means of fishing and lumbering. My companion affirmed that settlements had been pushed farther into the forest than they can be maintained, and that they must in most places be abandoned and the land given up to forest again. All along our way the woodlands were

straggling and sadly ravaged by the ax, fire, and wind. The spruce and pine had been culled out and most of the hemlock had been cut down and barked. Half-burned stumps and logs and gaunt and blackened trunks still standing disfigured the landscape on every side.

"The species of trees observed embraced all those common in northern woodlands. In one locality black cherry was remarkably abundant. Formerly the saw-mills of Colton cut pine, as there was a larger proportion of this lumber upon the Racket than is usually found in northern New York; now they do little business in any lumber.

"As we passed up along the river I saw small squads of 'drivers' stationed in a few places where the character of the river was such that it was liable to become obstructed with logs. By assisting the logs to pass such places great jams are prevented. The main body of the men, however, worked at the rear of the drive, scrambling over the disordered piles of logs which accumulate upon the shore or lodge against the rocks in the midst of the current. With their cant-hooks the men pry and roll the logs into the current, springing about on the pile as the logs roll from under their feet. Not unfrequently logs are left by the subsiding waters among the rocks at some distance from the main channel of the river. Files of men on each side then seize them with their cant-hooks and, splashing through the shallow water, bring them by main force into the channel. Sometimes logs become fastened among the rocks where the current is so swift that they cannot be reached by a boat or in any other way. Then hooks attached to ropes are thrown out from the shore; the logs are grappled and thus hauled off into the current. The drivers work Sundays and week days, fair weather or foul; their occupation is full of peril, and men are lost every year. Such are usually, as a driver assured me, 'men who do not know where it is safe to go.' But sometimes the most careful men become mixed with the rolling logs or seized by the current of the waterfalls and are swept away.

"Franklin county contains 995,279 acres, and 347,500 acres are still believed to be timbered. The timbered portion lies in the south end of the county, and because it is not watered through much of its area by streams of sufficient size for driving out the logs, much of the timber is inaccessible, or rather, the prices of lumber do not yet warrant hauling the logs long distances. The country across the line of the Ogdensburg and Lake Champlain railroad appears exhausted of its spruce and hemlock. Some tracts of hard wood are still standing, but the poplars, whose young growth often conceals the stumps and prostrate trunks of dead hemlocks, really seem in many places the most common species. But little timber land remains in Clinton county and, until the present season, lumbering on the Saranac had been for several years nearly suspended. This year, however, a company was cutting a few million feet of lumber drawn from the woods of Essex and Franklin counties. The lumber of the eastern side of the Adirondack wilderness mostly comes out by the way of the Saranac and the Hudson rivers. The mountain sides about lake George are being denuded of their spruce, which is sawed in the vicinity of Ticonderoga, and here, as elsewhere, fires follow the ax in their usual fashion."

The forests of the Adirondack region have suffered severe loss at different times, particularly in 1878, by the sudden death of great blocks of black spruce. Mr. Pringle carefully studied the extent of this destruction and the causes which produced it. In regard to these, great diversity of opinion exists among woodsmen and others familiar with the Adirondack forests. It has been generally supposed that the trees were killed by an unusually severe summer drought, or by the attacks of a boring insect working under the bark; but the testimony gathered by Mr. Pringle points to other causes of destruction. The spruce occupies dry mountain slopes and ridges and deep wet swamps never greatly affected by drought. It is noticed that as many trees have died in the swamps as upon the dry slopes. It is evidently not drought, then, which has caused them to perish. The opinion, too, is firmly held by the most intelligent observers that insects do not attack the trees until they are dead or nearly dead, and are never found in vigorous living specimens.

The black spruce is not a long-lived tree, and this dying out may indicate that the old trees of this forest, probably all of nearly the same age, had so nearly reached the limits of their natural existence as to be unable to withstand some unusual or severe climatic state, such as a period of intense winter cold or late spring frost. The following extracts from Mr. Pringle's report will indicate the opinions of those best able perhaps to form an opinion upon this subject:

"Mr. Mark Moody, residing at the foot of Tupper lake, a hunter and woodsman who has passed his life in the forest, testifies as follows: 'The spruce died fearfully in his vicinity about two years ago; he tried to learn the cause. Sixteen years ago the spruce had died out much in the same way as it has been doing lately. It is the older trees which die. They seem to die by crops, successively. Under the large trees were always springing up small trees to take the places of those that perish. There seems to be a narrower limit to the life of the spruce than to that of any other species. Other trees do not die in the same manner, by crops. The spruce does not seem to enjoy the same green old age, long drawn out, as other trees do, but when it has reached its full growth seems to relinquish its vitality without any apparent or sufficient cause, and before giving evidence of decay or any diminution of vigor.'

"Mr. Wardner, of Bloomingdale, Essex county, an old hunter, woodsman, and guide, testified as follows: 'The spruce timber on this side of the forest has failed clear through to its northern borders, in the same manner and during the same seasons as in other portions of the region.' Mr. Wardner first noticed the leaves falling and covering the ground in 1878; the destruction was continued through 1879, but during the past season he had met

with very few trees that were dying. Spruce timber had perished in this manner before, and he pointed out a broad valley in which most of the trees were dead and falling when he came into this region, twenty-five years before. He had carefully endeavored to ascertain the cause; was positive that insects either under the bark or upon the leaves had nothing to do with the death of the spruce trees, and he is sure that it is not due to drought, as he has seen the greatest destruction on the northern slopes. No active destructive agent being apparent, he inclines to the opinion that the spruce trees die because they have reached the limit of their life, and that it is some peculiarity of the winter rather than the summer that turns the scale against them; for this reason they perish in quantities, sometimes in sections. He has counted the rings of many trees, and considers 100 to 150 years the average lifetime of the spruce."

Whatever has caused the destruction of these forests, the damage thus occasioned, both in the loss of valuable timber and in the increased danger of forest fires from the presence of such a body of dead wood is enormous. It is believed by Mr. Pringle that from one-third to one-half of the fully-grown spruce timber left in the Adirondack region is dead.

NEW JERSEY.

The original forests of New Jersey have disappeared, except from some of the highest and most inaccessible ridges situated in the northwestern part of the state, and these, with the increased demands of the railroads for ties and other material, are now fast losing their forest covering. The forests of New Jersey are insufficient to supply the wants of the population of the state, and nearly all the lumber it consumes is brought from beyond its limits. The forests of pitch pine, which once covered large areas in the southern counties, have now generally been replaced by a stunted growth of oaks and other broad-leaved trees.

The forests of New Jersey, especially those on the dry sandy soil of the southern part of the state, have long suffered from destructive fires. During the census year 71,074 acres of forest were reported destroyed by fire, causing a loss of \$252,240. Of these fires twenty-eight were set by locomotives, seven through malice, seven by fires set on farms escaping to the forest, and six each by the carelessness of hunters and charcoal-burners.

The manufacture of cooperage stock and other industries using hard woods have been largely abandoned, owing to the decrease of the local supply of timber.

PENNSYLVANIA.

Pennsylvania once possessed vast forests of white pine and hemlock stretching over both flanks of the Alleghany mountains and extending from the northern boundaries of the state to its southern limits. East and west of the Alleghany region the whole country was covered with a heavy growth of broad-leaved trees mixed with hemlocks and occasional groves of pines. Merchantable pine has now almost disappeared from the state, and the forests of hard wood have been either replaced by a second growth or have been so generally culled of their best trees that comparatively little valuable hard-wood timber now remains. Large and valuable growths of hemlock, however, are still standing in northwestern Pennsylvania. From all parts of the state manufacturers using hard wood report great deterioration and scarcity of material, and Pennsylvania, which during the census year was only surpassed by Michigan in the value of its forest crop, must soon lose, with its rapidly disappearing forests, its position as one of the great lumber-producing states.

The following estimates of merchantable pine and hemlock standing in Pennsylvania May 31, 1880, have been prepared by Mr. H. C. Putnam. They are based upon the reports of a large number of timber-land owners and experts familiar with the forests of the state:


WHITE PINE (*Pinus Strobus*).


Regions.	Feet, board measure.
Alleghany river and tributaries	500,000,000
West Branch of the Susquehanna river and tributaries	1,300,000,000
Total	1,800,000,000
Estimated amount cut for the census year ending May 31, 1880	880,000,000

HEMLOCK (*Tsuga Canadensis*).

Estimated amount of hemlock standing May 31, 1880	4,500,000,000
Estimated amount cut for the census year, exclusive of trees cut for their bark alone.	300,000,000

Of lumber of all kinds 1,848,304,000 feet, including 288,561,000 shingles and 183,740,000 laths, were manufactured in the state during the census year; the nature of the returns, however, prevents anything beyond an estimate, based upon extended correspondence, of the amount of pine and hemlock sawed.





Numerous bodies of pine too small to be indicated on the map, of no great commercial importance and not included in these estimates, still remain scattered over the region originally occupied by pine forest.

The forests of Pennsylvania, especially through the mountain regions, have long suffered from destructive fires. During the census year 685,738 acres of forest were reported destroyed by fire, with a loss of \$3,043,723. Of these fires a large proportion were traced to locomotives and the escape of fires from farms to the forest.

The forests of Pennsylvania produced during the year 1879 2,866,010 pounds of maple sugar.

The following extracts are made from Mr. Pringle's report upon the principal lumber-producing regions of the state:

"Originally the broad pine belt of northern Pennsylvania, occupying the region drained by the numerous streams constituting the headwaters of the Susquehanna, extended from Susquehanna county, in the northeastern corner of the state, westward through Bradford and Tioga counties to Potter county, although this county never had as much pine as the others, and thence southwestward over Cameron, Elk, and Clearfield counties. The heaviest growth of pine in all this region was on Pine creek, in the southwest part of Tioga county. Now there is but little pine left in Susquehanna and Bradford counties, these counties being thickly settled; and in Tioga county, from which one firm alone has cut four billion feet, there now remain standing but little over one billion feet. The greatest part of the pine now standing in the Pennsylvania forests is on the upper waters of the West Branch of the Susquehanna, in Cameron, Elk, and Clearfield counties. In some of the counties adjoining these, as McKean, there was once, and still may be, a little pine timber.

"Active lumbering operations on the West Branch of the Susquehanna were begun in 1850, when the boom of the Susquehanna Boom Company was constructed at Williamsport. At this place the greatest part of the lumber on the West Branch is sawed. At Lock Haven, 25 miles above, on the same river, advantage was taken of the feeder-dam of a canal to construct another boom, and a few companies operating in lumber are now located there, about one-tenth as much lumber being sawed as is handled at Williamsport. Some of the companies, however, are removing from Lock Haven to the larger center of Williamsport. Below Williamsport no logs are driven, but a little timber squared by the ax in the woods and left at full length is made into rafts and taken down the main Susquehanna. Some of this is sawed in the towns on the river, and the remainder is taken to the large markets to supply the demand for squared timber for ship-building, etc.

"Williamsport is situated on the north or left bank of the West Branch of the Susquehanna, and for 2 or 3 miles along the river side are ranged the mills and lumber-yards of the thirty-four lumber companies operating here. We visited a large number of mills and found much the same methods employed in all. The logs are first slit up by gang-saws; then each board or plank is put through an edger, where two circular saws cut a strip from each side to give the board a square and straight edge; the boards are then assorted into two or more grades, loaded on trucks, and moved over tramways which ramify through the lumber-yards adjacent to each mill. The fragments of boards and better portions of the edgings are made into fence pickets and other portions into laths, and the fragments and strips which will not even make laths are carried to one side and added to a burning pile. The fragments thus burned (rather than thrown into the river) constitute the only waste, for the sawdust supplies the engines with fuel. This being cut chiefly from heart-wood makes better and more easily handled fuel than the sap-wood strips. Even these are, however, often cut and put up into bundles of kindling-wood for city use.

"In the woods the trees are sawed into logs 12, 16, or 18 feet in length, as can be done to the best advantage and the least waste of timber.

"The West Branch of the Susquehanna must be an exceptionally fine river to drive, judging from the comparatively unbattered condition of the logs seen about the mills. The smaller streams in the woods are furnished with flood-dams, and from these extend throughout the timber belt numerous narrow-gauge railroads, tramways, and slides for bringing down the logs. Little hauling is done upon wagons or sleds, the ground in the woods being too rough, it is said, for hauling logs with teams. It is probable that snow does not fill up the depressions and smooth the surfaces to the same extent as in the northern woods.

"The lumbermen of this place at first were content to send their lumber to market in the simplest shape, but of late, as the supply diminishes more and more, mills and shops are being built for the manufacture of doors, sashes, blinds, packing-boxes, furniture, etc. Some companies have so exhausted their pine lands that they can in future only carry on business in this way, buying the rough timber from their neighbors. As the pine lands of one firm after another are exhausted the pine remaining comes to be held by a very few parties, who know its value. Not all of these are operators, but, living at a distance, sell stumpage to manufacturers.

"The following table, giving the amounts of lumber rafted out of the Susquehanna boom at Williamsport since the record has been kept, may be of interest as showing something of the rise and decline of the lumber business at this important center. The greatest prosperity or fullest development of the business was attained, as will be seen, in 1873. After that year, with the steady decrease of the supply of pine and the consequent increase of expense in securing logs, the annual stock steadily diminished until 1877. During the past three years the increasing demand for lumber has stimulated the operators to greater activity, but more than to this cause the recent gain in the yearly stocks is due to the substitution of hemlock for pine, the ratio of hemlock to pine

being at present as 1 to 4, although the average for the last seven years is but as 1 to 10. As the supply of pine timber is exhausted, hemlock will be more and more handled until it will become the most important timber of this region. The summary is made for the last eight years only:

Years.	Number logs.	Feet, board measure.	Years.	Number logs.	Feet, board measure.
1862.....	196,953	37,853,621	1875.....	1,096,897	210,746,956
1863.....	405,175	76,475,826	1876.....	715,087	134,396,293
1864.....	511,549	96,595,681	1877.....	589,827	106,944,257
1865.....	379,392	72,421,468	1878.....	617,552	112,069,602
1866.....	615,373	118,831,494	1879.....	1,040,278	190,549,111
1867.....	833,388	163,196,511	1880 (to November 21).....	763,768	128,558,959
1868.....	853,663	165,338,389			
1869.....	1,080,511	223,060,305	1873 to 1880 (eight years) ...	7,395,455	1,382,342,272
1870.....	1,099,777	225,180,973	Logs remaining in river November 21, 1880		25,000,000
1871.....	842,129	116,661,181			
1872.....	1,484,103	297,185,652			1,407,342,272
1873.....	1,582,460	318,342,712	Deduct hemlock		140,734,227
1874.....	989,586	180,734,362	Williamsport pine, 1873-1880		1,266,608,045

"It is proper to add that the variations in the yearly stock of logs shown above are in some measure due to a greater or less proportion of each annual cut being left behind in the woods or in the streams, from varying supplies of water or from other peculiarities of the season.

"The lumber manufactured at Lock Haven and Williamsport is shipped by railroad and canal to Baltimore and Philadelphia and to intermediate cities and stations.

"I found it more difficult to obtain information of the extent and limits of the hemlock woods of Pennsylvania, and of the amount of the standing timber and the annual crop of hemlock, than I did to get the same facts respecting the pine. Lumbermen agree that there was originally far more hemlock in this state than pine, and they speak of it now as inexhaustible, which is not strictly true, for it is doubtful if it holds out to supply the increasing drain made upon it by tanneries and saw-mills for more than twenty-five years to come. Large quantities of hemlock have been wasted. Much that grew intermingled with the pine has died after the pine has been removed, partly from exposure to fuller sunlight and summer drought, and partly to forest fires induced by and following lumber operations. In the early days of the tanning industry of this region, when hemlock lumber was esteemed of little value, and whenever of late years the lumber trade has been so dull as to offer no inducement to send to market the trunks of the trees felled for their bark, large quantities of these have been left in the woods to decay. Now, however, with a good market for hemlock lumber, tanning companies owning hemlock lands, or the contractors who furnish the tanneries with bark, buying for this purpose stumpage from the proprietors of the timber-lands, often own saw-mills in the timber region, and cut and ship this lumber to market by railroad.

"Inasmuch as hemlock, besides mingling more or less with pine throughout the pine belt, seems to have formed a border entirely around the pine, the extent of the hemlock woods, as well as the quantity of hemlock timber, has always been much greater than of pine. Beginning in Wayne county, in the extreme northeastern corner of the state, the original hemlock forest extended westward through the northern tier of counties as far as Warren county, in the vicinity of lake Erie. Thence its bounds may be traced southward through Forest, Clarion, and Jefferson, and thence eastward through Clearfield, Center, Clinton, Lycoming, and Sullivan counties. Now the northeastern counties are for the most part cleared, and not only have the outskirts of these woods been cut off on all sides, but their continuity has been completely broken up throughout its whole extent by countless clearings and settlements. Yet, however much the hemlock forest has suffered, it possesses to-day greater value than did all the pine standing in 1850. Quite neglected a few years ago, hemlock is appreciating rapidly in value and importance, and ere many years shall have passed it will be almost the only kind of lumber known in the Williamsport market. The best grades of hemlock bring as high a price as scrub pine, the product of the shorter and more knotty trees grown on high land. Although as a rule Pennsylvania hemlock is of superior quality, much of it being nearly as good as spruce, yet here, as well as elsewhere, considerable variation in quality is noticed. Lumbermen classify hemlock into two kinds, red and white, according to the character of the wood, but the more intelligent among them attribute the difference to soil and situation. White hemlock, being sounder, firmer, and straighter grained, constitutes the highest grade. Red hemlock is more brittle, more inclined to splinter, and liable to be found more or less decayed when the trees have gained full size. In this condition trees are said to be 'shaky'. Such timber is generally found on bottom lands, while the hemlock of high hillsides is apt to be short and scrubby. The quality of the hemlock seems to deteriorate west from the center of the state. The Pine Creek hemlock is considered better than that of the Sinnamahoning, and this better than that on the Alleghany. Seldom more than two good logs can be obtained from a trunk, the third and fourth logs being generally inferior and knotty; 8,000 feet per acre is here considered a good yield of hemlock, and 10,000 feet a large yield.

"From Williamsport to Lock Haven the valley of the West Branch of the Susquehanna is usually less than a mile in width, being bounded by abrupt and rocky ridges a few hundred feet in height. At Lock Haven we

ascended the ridge on the south side of the river, some 800 feet in altitude, in order to examine the moderate forest growth with which it was covered. In favorable places scattering specimens of white pine indicated the crop these hills have yielded the lumberman in former years. Hemlock, also, was scattered over the hillsides, but even as late as the present year most of the trees in this immediate neighborhood had been felled for their bark; their peeled trunks lay strewn over the hillsides, being left to decay within a mile or two of the saw-mills of Lock Haven. The summit of the ridge afforded a good view of the surrounding country. Parallel ridges of a similar altitude, and which appeared more heavily timbered, lay back of the one on which we stood; between them were seen narrow valleys occupied by farms. On the north or opposite side of the river successive ridges rose higher and higher as they receded from the river, and in the distance seemed to lose themselves in a plateau whose altitude was equal to that of the ground on which we were standing. The gentle slopes and rounded summits immediately above the river showed smooth, cultivated fields interspersed among woodlands of deciduous trees. The more distant heights displayed a darker forest growth where hemlock and pines predominated.

"From Lock Haven to Warren, the county-seat of Warren county, even on the hillsides overlooking the river, close to the banks of which the railroad crept, but especially where we were able to look into the deep runs coming down to the river by a gradual descent from the table-lands of the divides, seldom more than a few miles back above the river, we saw much original forest still standing and principally composed of hemlock. Some white pine appeared as scattering trees or in groves, and some hard wood. The proportion of hard wood increased as we ascended the divide between the waters of the Susquehanna and those of the Alleghany river.

"On the summit of this divide the forest had a truly northern aspect, except that we missed the spruce, not seen in Pennsylvania. The dark foliage of the hemlock mingled with sugar maples, beeches, and birches. For many miles above Lock Haven it was a second growth which occupied the hillsides, a thin growth of white oak, chestnut, locust, etc., which had followed the lumberman and forest fires. Considerable second-growth white pine was seen in a few places, but on this none of the present generation seem to set much value, and I have yet to meet any one in the state who gives a thought to encouraging and preserving such growth. To consume the forests as speedily as possible, satisfied with what can be realized from them in the operation, appears to be the spirit which rules this region. Alternating here and there with the original forest mentioned above were seen all along the railroad leading through this timber belt, but especially in the vicinity of the settlements and lumbered districts, tracts which have been ranged by fire. Sometimes the fires had spread from the clearings into unculled timber, killing everything, large and small. Sometimes 'hemlock slashes' had burned over after the trees had been cut and 'peeled'. Always the charred stumps thickly dotted the ground, and the blackened, half-consumed trunks strewn over the soil in confusion gave to the landscape an aspect of complete desolation. The bird cherries and poplars, which in the forests farther north soon cover and hide from view such wastes of ruin, are wanting here.

"I learned that the best hemlock grows on the steep sides of the deep runs, and that upon the summits of the divides were considerable barrens, the soil of which was sometimes too poor to support any arboreal growth. Farther to the west the summits of the dividing ridges are occupied by hard wood chiefly, although hemlocks mingle with the beeches and maples.

"Arrived at Warren, we find that we have passed through the woods and are in a long-settled and well-improved country, and, judging from the scattered patches of woodlands occupying the low hills within view, the region of hard-wood forest has been reached. The coniferous forest belt only extends into the southeastern quarter of Warren county; the northern and western portions, lying beyond the Alleghany river, yield oak, chestnut, hickory, etc. Originally there was a little pine scattered over the southeastern portion of Warren county, but this has been mostly cut, and hemlock remains, as it ever has been, the most important timber in this part of the county. In Forest county, next south of Warren, pine is local, being scattered in small quantities throughout the county. On the highlands there is much hard wood, beech, maple, and white wood existing in belts between the streams. This, however, may be called a hemlock county. In McKean county a central table-land is covered principally by a growth of maple, beech, etc. In the remaining portions of the county the timber is chiefly hemlock. The valley of the Alleghany river, in the eastern part of McKean county, is mostly cleared and improved. Elk county is one of the best counties for hemlock. Through Elk, the southwestern corner of McKean, and the southeastern corner of Warren runs the Philadelphia and Erie railroad. Along the line of this road, as it passes through this portion of the timber belt, are located the largest tanneries of the United States. These are consuming the hemlock of this region at an enormous rate, and, in addition to the vast amount of bark which they consume, large quantities are shipped out of the region by railroad. The first important tanneries of Warren county were established 12 or 15 years ago, and at the present rate of consumption the hemlock of this county can hardly hold out 20 years longer. The land, after the forest has been removed, is excellent for agricultural purposes throughout this region, and on all sides pioneers are making themselves farms. These men prefer to begin in the undisturbed forest rather than locate on the slashes, because they can pay for their land with the hemlock bark which it yields; and from a radius of 15 miles bark is drawn and sold at from \$4 50 to \$5 a cord to the tanneries. On an average, four trees yield a cord or ton of bark, the equivalent of 1,000 feet of lumber, board measure. In Warren county from 5,000 to 6,000 acres of hemlock were cut down in 1880, and there is no possibility of this growth being renewed, for every foot of slashed land is eventually burned over, and sometimes the burnings are repeated until the soil is nearly ruined for agricultural purposes. From the dry slashes the fires extend to a greater or less distance through the living

woods, ruining not only heavy bodies of hemlock, but also destroying the belts of hard wood intermixed with the hemlock. Notwithstanding stringent legislation in this state upon the subject of forest fires, they seem inevitable, and especially so in the slashes. They spread from the clearings constantly made throughout this timber belt by the settlers, and, as the forest abounds in deer and its streams are stocked with fish, hunters and fishermen are always in the woods, and from their camp fires spread many conflagrations. Many fires here also are set by a tribe of half-civilized Indians residing in this region, to burn over the huckleberry fields in order that the bushes may renew themselves and yield fuller crops; or, where it is so easy to start a fire and conceal its origin, many doubtless arise from malice.

"In this region the aspen springs up on land upon which the hemlock has been destroyed, but this tree manifestly does not thrive as it does in northern woods. Yellow and black birch, bird cherry, beech, maple, white oak, chestnut, black cherry, etc., are the trees which spring up slowly among the briers, and cover burned land with a rather meager second growth. If a few pines have been left on the hilltops they may scatter a few seeds and give rise to some saplings, but as regards hemlock, fires kill it out clean, seedlings and seed; and if the 'peelers' and the fires happen to leave any scattering trees standing, these, being more sensitive to changed conditions than pines, are seldom able long to survive as seed bearers. The bird cherry only thrives on cold, wet soils here. There is another phase of the slaughter of the hemlock forest: As the pine forest gives out, large numbers of laborers turn to the hemlock woods and find employment as bark peelers. In the pine woods work is mostly suspended when spring arrives; then larger numbers of men come into the hemlock woods than can find work at satisfactory wages, and these sometimes set fires in the slashes, which spread into the living woods and kill large quantities of hemlock. To save the bark it must be peeled at once, or before it adheres to the wood and becomes injured by worms, and thus employment is given to a larger force of men.

"The pine now remaining in Clearfield county is mostly found in the northern and the southwestern portions of the county. The eastern and southeastern portions are now principally cleared and improved, as the entire county is destined to be, the soil being principally a strong, clayey loam, excellent for farming purposes. Already four-fifths of the pine timber originally standing in the county has been removed; most of the hemlock, which originally about equaled in amount the pine, remains. There are no tanneries in this region, and after the pine is cut the hemlock is next harvested, the bark being saved and shipped to the tanneries below to the amount of from 5,000 to 6,000 cords annually. Fires are here sometimes started by hunters in order to clear away the young second growth, that they may be able better to see the deer. One important reason which lumbermen have for planting their saw-mills near the woods, in preference to driving all their logs to the sawing centers below, is that they can then work into shingles, etc., many trees which, being defective by reason of rotten spots or other blemishes, would not be worth driving down the river. Such trees are seen standing here and there all through the woods, having been left behind by the lumbermen. Sometimes persons buy this culled timber and erect shingle-mills, etc., to work it up.

"With respect to the maximum yield of pine per acre, it would seem that 10,000 feet was a good yield for tracts of 400 or 500 acres in extent, although smaller tracts of 50 acres and upward will often cut 25,000 feet to the acre, and even a yield of 100,000 feet to the acre has been reported. The rough nature of the surface in all this region often necessitates the use of slides to bring the logs from the forest to the streams. They are constructed by pinning to ties of hemlock some 3 feet in length hemlock logs about a foot in diameter placed side by side, their inner sides above the point of contact being hewn with care to form a broad V-shaped trough along which the logs may be slid. Except where there is considerable descent logs cannot be slid unless the weather is frosty, when the slide can be kept icy by means of water sprinkled over it from time to time. Slides sometimes are built for 6 or 8 miles back into the woods, usually following up some run so as to get an even and gentle grade. By this means the greatest part of the logs come down to the streams, for sleds are not used in this country. Most of the hazard of lumbering depends upon the lumberman's ability to slide his logs successfully. They can be cut at any time in the woods, and almost any year can be driven to the mills when once in the water, but mild weather interrupts sliding and deep snows impede the operation; so that in open winters lumbermen are sometimes compelled to do their sliding in the night time, when ice will form on the slide. The logs, stripped of their bark, are drawn singly, by horses with chains, from the places where they have fallen to the upper end of the slide. When a sufficient number—from 6 to 40, according to the grade and the size of the logs—have been placed end to end in the slide, the hook of a chain is driven into the rear log near its forward end, and horses are attached which walk a tow-path formed on one side of the slide, and push ahead of them the 'trail' of logs, thus bringing them down to the stream.

"Only in the late autumn and in the winter is it thought expedient in Pennsylvania to fell pine; if cut in summer, when the bark will part from the wood, the sap-wood soon assumes a blackish appearance and disfigures the lumber. As a rule hemlock is here cut and peeled in summer, at the time when operations in pine are suspended; thus by alternating operations in pine and hemlock the hands are kept employed throughout the whole year. In cutting trees the several parts of the work are allotted to different men; some merely fell the trees, others measure them off into suitable lengths and cut away the limbs as far as the upper end of the last log taken, where they sever the top of the tree from the trunk by means of the ax; others follow in pairs with cross-cut saws and cut the trunk into logs."

DENSITY OF FORESTS
 COMPILED UNDER THE DIRECTION OF
 C. S. SARGENT, SPECIAL AGENT
 1887.

Scale: 0 10 20 30 40 50 Miles

[illegible]

LEGEND.

I	Under 1 cord per acre
II	From 1 to 2
III	2 to 5
IV	5 to 10
V	10 to 20
VI	Over 20

Julius Fien & Co Ltd

SOUTH ATLANTIC DIVISION.

DELAWARE.

The northern portion of the state, comprising New Castle and Kent counties, was once covered with the deciduous forests of the Atlantic plain. Conifers, with the exception of the red cedar, were rare. In the sandy soil of the southern part of the state various pitch pines flourished, forming fully one-half of the forest growth. These pine forests were long ago consumed and are now replaced by a second growth, generally composed of the species which originally occupied the ground; and throughout the state the best hard-wood timber has been culled from the forest. Large quantities of wheel and cooperage stock were formerly manufactured in the northern counties; but of late years these and other industries using the products of the forest have, for want of material, generally decreased in importance. The manufacturers report a general scarcity of timber.

During the census year 3,305 acres of woodland were reported destroyed by fire, with a loss of \$15,675. Of such fires six were set by locomotives, six by the careless burners of brush upon farms, and two through malice.

KENT COUNTY.—About one-quarter of this county is reported covered with forest. A few small mills saw oak from the immediate neighborhood into shipstuff and car lumber, shipping to Wilmington, Philadelphia, and even to New York.

NEW CASTLE COUNTY.—About one-quarter of this county is reported covered with woodland, mostly of second growth and attached to farms. The large establishments for the manufacture of gunpowder, located in the neighborhood of Wilmington, consume large amounts of willow wood, generally grown for the purpose upon farms in their immediate vicinity.

SUSSEX COUNTY.—One-third to one-half of this county is reported covered with woodland. Numerous small mills, obtaining their supply of logs from the immediate neighborhood, saw oak for shipstuff.

MARYLAND.

The northwestern portion of the state, crossed by the ridges of the Appalachian system, was once covered with the forests of white pine, hemlock, birch, and maple peculiar to this mountain region. The central portion of the state, extending from the mountains to the shores of Chesapeake bay, was covered with oaks, hickories, gums, and other deciduous trees in great variety, the eastern peninsula largely with different species of pitch pine, occupying sandy plains, or mixed with deciduous trees.

In the mountain region considerable bodies of the original forest remain upon the highest and most inaccessible slopes; in the remainder of the state this, where the land has not been permanently cleared for agriculture, is now largely replaced by a second growth, or—the best timber at least—has been everywhere culled.

A large amount of cooperage stock was formerly manufactured in this state. This industry has, however, greatly suffered from the deterioration and exhaustion of the local supply of timber; manufacturers report the best stock nearly exhausted and the substitution for oak, formerly exclusively used, of elm and other inferior woods now brought from beyond the limits of the state.

During the census year 41,076 acres of woodland were reported destroyed by forest fires, with a loss of \$37,425. These fires were traced to the carelessness of hunters, to locomotives, and largely to the escape from farms to the forest of fires set in clearing land. The principal lumber manufacturing establishments using Maryland logs are situated in Garrett county; these saw white pine, hemlock, and oak to supply a limited local demand and ship to Baltimore, Philadelphia, Pittsburgh, and Wheeling; considerable oak timber is sent to Europe from this county. During the year 1879 the northern counties produced 176,076 pounds of maple sugar.

DISTRICT OF COLUMBIA.

The original forest has disappeared from the District of Columbia and has been replaced by a second and third growth of oaks, scrub pines, and other trees. The area occupied with woods is probably slowly increasing. A single saw-mill, situated in the city of Washington, saws logs grown beyond the limits of the District.

VIRGINIA.

The forests of Virginia, like those of the Carolinas and Georgia, fall naturally into three divisions, dependent upon the elevation and soil of the different parts of the state. The mountains and ridges of its western border are

covered with a heavy growth of pine, hemlock, white oak, cherry, yellow poplar, and other northern trees; over the region extending east of the mountains oaks, principally black oaks, once formed the prevailing forest growth; through these are now mingled long stretches of various pitch pines, occupying exhausted and barren soil once devoted to agriculture. The eastern counties are covered with the forests of the Maritime Pine Belt, generally confined to the Tertiary deposits of the coast and extending inland to the head of tide-water of the principal streams; along the western borders of this pine belt the forest growth is nearly equally divided between the pines and the broad-leaved species.

The inaccessible mountain region in the southwestern part of the state still contains immense quantities of the original oak, hickory, walnut, and cherry, the scanty population of these mountains having made but slight inroads upon the forests. Railroads have hardly penetrated them, while the streams which head here are unsuited to carry to market the hard woods of which this forest is largely composed. The most valuable hard-wood forest remaining on the continent exists in southwestern Virginia and the adjacent counties of West Virginia, Kentucky, Tennessee, and North Carolina. From the central and eastern portions of the state the original forest has almost entirely disappeared, and is now replaced by a second growth, in which the Jersey pine and the old-field pine are characteristic features, generally replacing more valuable species of the original growth.

During the census year 272,319 acres of woodland were reported ravaged by fire, with a loss of \$326,944. Of such fires the largest number was traced to the careless burning of brush upon farms and to locomotives.

The manufacture of cooperage stock is increasing rapidly in the western part of the state, and great quantities of staves are exported thence directly to Europe, as well as oak, yellow poplar, and walnut in the log. The manufacture of tobacco cases from sycamore lumber is an important industry in the neighborhood of Lynchburg and other tobacco-distributing centers. Considerable quantities of hand-made shingles are produced in the cypress swamps which occupy a large portion of Norfolk and other eastern counties. A large amount of second-growth pine (*Pinus Taeda*) is shipped from the different Virginia ports by schooner to New York for fuel, and this second-growth pine furnishes the principal building material used throughout the state. The grinding of oak and sumach bark and the manufacture of tanning extracts are important and profitable industries of the state.

WEST VIRGINIA.

The forests of West Virginia, with the exception of the belt of pine and spruce confined to the high ridges of the Alleghany mountains, are principally composed of broad-leaved trees, the most important of which are the white and chestnut oaks, the black walnut, the yellow poplar, and the cherry. The white pine and spruce forests reach within the state their southern limit as important sources of lumber supply.

The forests have been largely removed from the counties bordering the Ohio river, and the most valuable hard-wood timber adjacent to the principal streams, especially black walnut, cherry, and yellow poplar, has been culled in nearly every part of the state. But slight inroads, however, have yet been made into the magnificent body of hard-wood timber covering the extreme southern counties, which still contain vast quantities of oak, cherry, and poplar.

The black walnut found scattered everywhere in West Virginia is least plentiful in the northwestern and Ohio River counties, and most abundant along the upper waters of the rivers flowing into the Ohio through the southwestern part of the state. Yellow poplar is found throughout the state, and is still abundant about the headwaters of nearly all the principal streams. Large bodies of cherry are found in Greenbrier, Nicholas, Webster, and other counties immediately west of the mountains, and a large amount of hemlock is scattered through the valleys and ravines of the northeastern part of the state and along the western slopes of the Alleghanies. The area still occupied by white pine is estimated to extend over 310 square miles, and to contain about 990,000,000 feet of merchantable lumber. The principal centers of lumber manufacture are along the Kanawha river at Ronceverte, in Greenbrier county, at Parkersburg, and along the upper Potomac.


Partial returns of the hoop-pole industry gave a product during the census year of 3,549,000, valued at \$146,000.

During the census year 476,775 acres of woods were reported destroyed by fire, with a loss of \$155,280. Of these fires the largest number was traced to the careless clearing of land for agricultural purposes, although many had their origin in sparks from locomotives.

The manufacture of cooperage stock is fast increasing in importance, and seems destined, with the exhaustion of the more accessible hard-wood forests of the country, to assume a much greater development than at present. Large quantities of black walnut, yellow poplar, and oak in the log are shipped to northern markets and to Europe.

The following notes upon the forests of West Virginia are extracted from Mr. Pringle's report:

"Entering West Virginia at Keyser (New Creek) by way of Cumberland, Maryland, we find ourselves in one of the narrow valleys lying among the low abrupt ridges of the northern Alleghanies, among which we have been traveling since we reached the vicinity of Williamsport, Pennsylvania. Coming south from middle Pennsylvania, however, the forest growth covering the long mountain chains within view from the railroad becomes heavier and heavier, the evidences of fire and ax largely disappearing. On the hills above Keyser fewer evergreens appeared than I had previously seen. A few slopes were principally occupied by pine in variety, but the mountains of this



region were covered with a growth of deciduous trees, white, black, red, Spanish, and chestnut oaks, hickories, butternuts, black walnuts, yellow poplars, locusts, elms, sugar maples, etc. At Piedmont some \$200,000 have been expended in the construction of a boom on the North Branch of the Potomac. At this point, as well as at Swanton and Deer Park, on the Maryland side, there are mills sawing chiefly white oak, and also considerable white pine, spruce, hemlock, poplar, white ash, etc. Some spruce which had not been seen or heard of in the timber belt of Pennsylvania is found 20 miles above Piedmont. The market for lumber manufactured here is chiefly eastward. Much of the oak is sent to Europe, partly in the form of squared timber, partly cut 5 by 12 inches and from 15 to 20 feet long. The mills at Swanton and Deer Park are located on the railroad, and cut timber is hauled to them from the vicinity. The mills at Piedmont are fed by logs driven down the river from the western portions of Mineral and Grant counties, West Virginia. This lumber is chiefly oak, spruce, and hemlock. Great difficulty is experienced in driving this part of the Potomac, as it is a swift and rocky stream. Logs, especially oak, constantly lodge on the rocks or banks, and there has been great difficulty in maintaining the boom and dam at this point.

"Rowlesburg, in Preston county, owes its existence as a lumber depot to the fact that the Cheat river, upon which it is situated, as it passes through the Briery mountains, for a distance of 25 miles below this point has so narrow and rocky a channel and so swift a current that it is not possible to get the logs farther down the stream. Above Rowlesburg the Cheat river is a good stream to drive, and any one of its branches can be driven from a point 125 miles above that point. From the mouth of the Black Fork, 30 miles above, the timber is brought down in rafts rather than as separate logs; this is because there is no boom as yet at Rowlesburg to stop the logs. There are small booms on Black and Shaver's Forks, many miles above Rowlesburg. Scattered along the river at some distance above Rowlesburg there are a few small mills, the product of which is floated down the stream on rafts. The timber of Preston county between Rowlesburg and the vicinity of the mouth of the river is oak, poplar, chestnut, ash, beech, yellow beech, hemlock, basswood, and hickory.

"The timber of Canaan valley, in Tucker and Randolph counties, is largely hemlock on the lower lands, on the higher situations and slopes sugar maple and beech; and, as soon as a suitable elevation is reached, spruce is mingled with black cherry. In other portions of Tucker county and on the tributaries of the Cheat river, flowing out of Randolph county, the timber is chiefly oak, poplar, ash, spruce, cherry, black walnut, white pine, etc. This, however, is not a black-walnut region, and there are here nowhere more than scattered trees; a careful search has failed to find any great body of this timber here. It is estimated that 2,500,000,000 feet of yellow poplar are still standing in the valleys of the Cheat and its tributaries.

"Shaver's Fork is heavily timbered with spruce. A boom has been constructed at Grafton, on Tygart's Valley river, a main branch of the Monongahela. It is a rough stream, unfavorable for lumber operations, and for a distance only of 10 miles above Grafton is smooth enough to admit of the passage of rafts. All lumber has, therefore, to come down in separate logs, and only such kinds as are light enough to float well can be got down. For this reason there is very little except poplar sawed at Grafton. Oak is too heavy to be driven successfully, and as it cannot be tied up in rafts with poplar, as is done on the Cheat, the stores of oak timber growing in the valleys drained by this river must wait the building of a railroad to bring them to market. The yellow poplar still standing in this region is estimated at 300,000,000 feet, and on the higher grounds, especially about the headwaters of streams, there are fine bodies of black cherry mixed with other trees.

"At Parkersburg are located the mill and shops of the Parkersburg Mill Company, situated on the banks of the Little Kanawha, a short distance above its confluence with the Ohio. This is the only company operating in lumber within the city of Parkersburg. It manufactures about 6,000,000 feet of lumber annually, mostly poplar, some oak, and about a quarter of a million feet of beech. Little black walnut can now be obtained here, and that of inferior quality. Rough lumber and manufactured articles of wood find a market in nearly every direction, west, north, and east. I was astonished and delighted to see how closely the lumber was worked up and the great variety of articles manufactured from slabs, edgings, culls, etc., which in other mills are so generally thrown into the waste pile. Broom handles, corn-popper handles, brush handles, brush heads, tool handles of many descriptions, and fly-trap bottoms are but a few of the articles which are turned out by millions from odd bits of wood, few of which are too small to make something or other from. The company executes orders for articles used in manufactories widely distributed over the country from Cincinnati and Chicago to Boston and New York. Poplar is used for broom handles, and beech, maple, sycamore, black walnut, cherry, etc., for smaller articles. This company does not own and operate timber lands, but buys its logs from parties who deliver rafts to its mill. Formerly much lumber was wasted in this region in clearing lands for farms, but now proprietors of land find it to their advantage to cut and save their logs, which they bring down in rafts themselves or sell to parties who make a business of rafting. Once out of the small streams, the logs are easily rafted down the Little Kanawha during favorable seasons.

"There are no booms on the Little Kanawha, except temporary constructions for special purposes, which are broken up by every flood. Several years ago it was supposed that the timber on this river was nearly exhausted, but it continues to come down in undiminished quantities to the value of some hundred thousand dollars annually, in addition to railroad ties, staves, etc. It is only about 40 miles up the main river, and to no great distance back from the stream, that the supply of oak is exhausted. The river is a hundred miles long, and about its upper


waters and those of its tributaries the oak is comparatively untouched. Much of Wirt county and the greater part of Roane, Calhoun, and Gilmer, in the upper part of the valley of the Little Kanawha, are a vast virgin forest of oak and poplar, containing a good deal of black walnut and sugar maple and some black cherry. Baxter county is magnificently timbered, as is Webster, although the timber here is yet inaccessible.

"The Guyandotte is a good river for lumbering operations. Rafts can come down from a point 100 miles from its mouth. There are yet no booms on this river, except strings of logs occasionally stretched across it for temporary purposes. On its course above Guyandotte are four or five mills, doing for the most part a local business, their product for export being only about 1,000,000 feet of sawed lumber. The rafting of this sawed lumber is attended with some risk of loss, and therefore a much greater amount is brought down in unsawed logs bound together in rafts, which are taken down the Ohio and sold to various mills along its course. These rafts are usually made 11 logs wide, and three or four of these courses are placed end to end. White oak is made up into rafts with a poplar log in the center of each course, and thus the raft is made light enough to float easily. Along the Guyandotte, in the lower part of its course, the oak and poplar have been cut for a distance of from 1 mile to 2 miles from the stream, the black walnut for some 5 miles back; but nine-tenths of the area drained by this river is still in original forest, composed of white, chestnut, and other oaks, poplar, walnut, several hickories, beech, sugar maples, sycamore, ash, etc. In this region there is, in the aggregate, a good deal of black walnut, but it exists as scattered trees rather than in groves or tracts.

"Coal river is 160 miles long, and for 36 miles, or to Peytona, is navigable for barges. The valley of this river is covered with truly magnificent forests, in which the trees of the several species composing them attain remarkable dimensions. Poplar and white oak here exist in nearly equal proportions, and together constitute about a third of the timber. Besides these there is a good deal of black cherry, lin, and locust, as well as hemlock, the latter not being considered valuable in this country. Black walnut appears more abundant in this region than in any other of similar extent of which I have yet heard. But little timber has yet been removed from the valley of this river, and it is chiefly the lower portion and the immediate vicinity of the banks which have been lumbered.

"The Elk river empties into the Kanawha at Charleston. About 2 miles above its mouth are located a boom and several saw-mills, and here are also a dam and lock which secure slack-water for some 20 miles. The river is about 180 miles in length; logs have been driven from a point 150 miles above its mouth, but its valley has only been lumbered to any great extent in the immediate vicinity of the main river, and to a distance of some 110 miles from its mouth. Most of the original growth of the forest of the Elk basin still remains, and is composed largely of white oak, hickory, chestnut, and poplar. Black walnut here, as everywhere else in this state, is scattered, although it is estimated that 10,000,000 feet of this lumber still remain in this region. Above a certain altitude and about the upper waters of this river considerable black cherry, sugar maple, and birch is found. Here also beech and basswood abound, by the streams hemlock occurs, and on the mountains a little black spruce. About the upper settlements on this river miles of fence constructed with boards of black cherry and farms fenced with black-walnut rails may be seen. Formerly large numbers of coal-boats and salt-boats were built upon the Elk river. Once, also, the salt-works of the Kanawha required vast numbers of barrels; these were made of black as well as white oak; now but five of the sixty furnaces once boiling brine in this vicinity are in operation, and there is little demand for black oak for staves. The country along the Kanawha between the Elk and the Gauley rivers has been lumbered for 5 or 6 miles back from the streams, and about one-fourth of the timber has been cut from these valleys. The Gauley river with its several large tributaries drains a valley which covers nearly 5,000 square miles; its length is about 110 miles, much less than that of the Elk, which is a long, slender stream, but it occupies a much broader valley and has twice the volume of water of the Elk. Unlike the rivers just considered, which wear out for themselves smooth channels through the soft sandstone, the Gauley is a rough stream, tumbling rapidly over hard conglomerate rock, its bed being full of bowlders and ledges. For the first 10 miles from its mouth the fall averages 4 feet to the mile; above that 20 feet to the mile, while its upper waters are so swift and rough as to be unnavigable even for small boats. For these reasons the Gauley does not admit of the passing of rafts, and it is a difficult river upon which even to drive single logs. Its valley is but little settled, except on Meadow river and along its right bank below that stream. Above a point 15 miles from its mouth no timber has been touched except by the few settlers. In the lower part of the valley of the Gauley for 15 or more miles the timber is chiefly oak, poplar, walnut, etc.: The Gauley and its large affluents, the Cherry, Cranberry, and Williams rivers, all head back in the forests of black spruce, which sometimes take entire possession of the mountain tops; a little lower, yet often mingled with the spruce, hemlocks and black cherry abound. On Cherry river the cherry trees so predominate over all others as to have given their name to the stream. Here are trees often 4 feet in diameter. The region intermediate between the upper and the lower districts of the Gauley thus described contains much beech, sugar maple, and black cherry. The white oak which abounds in the lower basin of this river disappears above an altitude of 2,000 feet. I was informed that, although lumbering operations were but lately begun on the Gauley, nearly 1,000,000 feet of poplar were brought out of the river in 1879, and that it had yielded 50,000 feet of black walnut in 1880, while there were now in the river poplar logs enough to make 3,000,000 feet of lumber. About one-fourth of the cut of late years has been sawed at mills near the falls; the rest is rafted to Charleston.

DEPARTMENT OF THE INTERIOR





"The valley of the New river is only lumbered for from 3 to 5 miles from the stream, although the walnut has been gathered 10 miles farther back. This is a rough country in which to lumber, since the streams cut deep into the earth, and New river cannot be driven.

"Ronceverte is situated on the Greenbrier river at the point where the Chesapeake and Ohio railroad first meets this stream as it descends from the Alleghany mountains. Here is the boom of the Saint Lawrence Boom Company, and here are located three or four lumber firms operating steam-mills. One of these, the New York Hoop Company, uses two million hoop-poles per annum, chiefly hickory, manufacturing hoops for flour barrels, pork barrels, hogsheads, and tierces, besides strips for boxes, etc. The process of manufacturing hoops was explained to me as follows: The poles, of assorted lengths and sizes, are passed through machines which split each of them into two, three, or four pieces, and these are put through other machines which plane flat the inner side of each strip, leaving the bark intact. The hoops thus made are tied into bundles and shipped to New York.

"The Greenbrier river rises in the limestone sinks in Randolph county, whence it flows southwesterly through the fertile limestone valley between the Alleghany and the Greenbrier mountains for a distance of 120 miles, emptying into the New river at Hinton. Flowing through such a valley it is not a rapid stream, but from a point 12 miles below Travelers' Rest, on its headwaters, it is fine for rafting. Yet the stream needs some improvement, especially by the closing up of back channels into which the logs are borne by high water, to be left in swamps when the flood recedes.

"Only a small proportion of the timber of the Greenbrier river has been removed as yet, and it is estimated that in its valley white oak, white pine, poplar, cherry, hemlock, walnut, and ash enough remain to make 1,000,000,000 feet of boards, and that there are not less than 500,000,000 feet of white pine in this region, occupying a belt through the center of both Greenbrier and Pocahontas counties. The eastern limit of the black-spruce belt on the headwaters of the Elk and Gauley rivers, the most extensive and valuable in West Virginia, coincides with the western limits of the white-pine belt lying in Pocahontas county. Its southern line runs northwesterly from the south end of Pocahontas to near the center of Nicholas county. From this point its western line runs northeasterly through the center of Webster county to the vicinity of Huttonville, in Randolph county, the northern end of the belt covering the upper waters of Shaver's Fork of the Cheat river. Over this belt black spruce is scattered more or less densely, sometimes occupying almost exclusively the high slopes, particularly the northern slopes and the summits of the mountains.

"It is believed that 10,000,000 feet of black walnut, in paying quantities, could still be gathered in this part of the state, and that there would then be left an equal amount so scattered that it could not be profitably collected at present prices."

NORTH CAROLINA.

The forests of North Carolina were once hardly surpassed in variety and importance by those of any other part of the United States. The coast region was occupied by the coniferous forests of the southern Maritime Pine Belt; the middle districts of the state by a forest of oaks and other hard-wood trees, through which the old-field pine is now rapidly spreading over worn-out and abandoned farming lands. The high ridges and deep valleys of the Appalachian system which culminate in the western part of the state are still everywhere covered with dense forests of the most valuable hard-wood trees mingled with northern pines and hemlocks. The inaccessibility of this mountain region has protected these valuable forests up to the present time, and few inroads have yet been made into their stores of oak, cherry, yellow poplar, and walnut. The hard-wood forests of the middle districts, however, have been largely removed or culled of their finest timber, although the area of woodland in this part of the state is now increasing. These new forests, usually composed of inferior pine, are of little economic value, except as a source of abundant fuel and as a means of restoring fertility to the soil, preparing it to produce again more valuable crops. A larger proportion of the pine forest of the coast has been destroyed in North Carolina than in the other southern states. This part of the state has long been the seat of important lumbering operations, while the manufacture of naval stores, once almost exclusively confined to North Carolina, and always an important industry here, has seriously injured these forests. The original forests have been practically removed from the northeastern part of the state, the great region watered by the numerous streams flowing into Albemarle and Pamlico sounds; and although some lumber, largely second-growth pine trees of poor quality, is produced here, the importance of these forests is not great. The merchantable pine, too, has been removed from the banks of the Cape Fear and other rivers flowing through the southern part of the state, and although these streams still yield annually a large number of logs, they are only procured at a constantly increasing distance from their banks and with a consequent increasing cost for transport.

Forest fires inflict serious damage upon the pine forests of the south. During the census year 546,102 acres of woodland were reported destroyed by forest fires, with a loss of \$357,980. The largest number of these fires were traced to the carelessness of farmers in clearing land, to locomotives, hunters, and to malice.

Manufacturers of cooperage and wheel stock, industries which once flourished in the eastern and central portions of the state, already suffer from the exhaustion and deterioration of material. Such industries, however, are increasing in the extreme western counties, and promise to attain there an important development.

THE FORESTS OF THE UNITED STATES.

The following estimate, by counties, of the merchantable pine standing May 31, 1880, south of the Neuse river, the only part of the state where it is of commercial importance, was prepared by Mr. Edward Kidder, of Wilmington. It is based upon actual surveys and the reports of a large number of timber-land experts familiar with the different counties still occupied by the forests of long-leaved pine:

LONG-LEAVED PINE (*Pinus palustris*).

Counties.	Feet, board measure.
Bladen	288,000,000
Brunswick	141,000,000
Chatham	448,000,000
Columbus	288,000,000
Cumberland	806,000,000
Duplin	21,000,000
Harnett	486,000,000
Johnston	563,000,000
Moore	504,000,000
New Hanover	96,000,000
Onalaw	34,000,000
Robeson	864,000,000
Sampson	602,000,000
Wake	48,000,000
Wayne	40,000,000
Total	5,229,000,000
Cut for the census year ending May 31, 1880, exclusive of 50,190,000 feet cut in the counties adjacent to Albemarle and Pamlico sounds and along the Pamlico and Neuse rivers, which is largely loblolly pine (<i>Pinus Tæda</i>).	108,411,000

NAVAL STORES.

Small quantities of crude turpentine were produced upon the coast of North Carolina, between the Pamlico and Cape Fear rivers, soon after the earliest settlement of the country. It was sent to Great Britain or converted into spirits of turpentine and rosin for home consumption. The demand for ships' stores had greatly increased the North Carolina production as early as 1818, although the field of operations was not extended south of the Cape Fear river, nor more than 100 miles from the coast, until 1836. The large demand for spirits of turpentine created during that year induced manufacturers to test the yield of trees on the west side of the Cape Fear river, up to that time considered unproductive. The result was satisfactory, although overproduction and low prices deferred until 1840 the development of this region. Since 1840 this industry has been gradually carried southward. Naval stores were produced in South Carolina in 1840, and in Georgia two years later. Turpentine orchards were established in Florida and Alabama in 1855, and more recently in Mississippi and eastern Louisiana.

The naval stores manufactured in the United States are principally produced from the resinous exudations of the long-leaved pine (*Pinus palustris*), and in small quantities from the loblolly pine (*Pinus Tæda*), and the slash pine (*Pinus Cubensis*) of the Florida coast. The trees selected for "boxing" are usually from 12 to 18 inches in diameter, although trees with trunks only 8 inches through are now sometimes worked. A deep cut or "box" is made in the trunk of the tree, by a cut slanting downward, some 7 inches in depth, and generally 12 inches above the ground, and met by a second cut started 10 inches above the first and running down from the bark to meet it. In this manner a segment is removed from the trunk and a triangular trough formed 4 inches deep and 4 inches wide at the top.

Two such boxes, or upon a large trunk sometimes four, are made on each tree. A "crop", the unit of production among large operators, consists of 10,000 such boxes. The boxes are cut early in November with a narrow-bladed ax specially manufactured for the purpose, and the trees are worked on an average during thirty-two weeks. As soon as the upper surface of the box ceases to exude freely, it is "hacked" over and a fresh surface exposed, the dried resin adhering to the cut having been first carefully removed with a sharp, narrow, steel scraper. The boxes, especially after the first season, are often hacked as often as once a week, and are thus gradually extended upward until upon trees which have been worked during a number of seasons the upper surface of the box is often 10 or 12 feet above the ground. For these long boxes the scraper is attached to a wooden handle, generally loaded with iron at the lower end to facilitate the operation of drawing down the resin. Once in four weeks, or often less frequently, the resin caught in the bottom of the box is removed into a bucket with a small, sharp, oval steel spade attached to a short wooden handle. The product of these "dippings", as this operation is called, is placed in barrels and transported to the distillery. The first season a turpentine orchard is worked boxes are usually dipped eight times, yielding an average of 300 barrels of turpentine to the crop. The second year the

number of dippings is reduced to five, the product falling off to 150 barrels, while for the third season 100 barrels are considered a fair yield from three dippings. To this must be added the yield of the "scrapes", which for the first year is estimated, for one crop, at from 60 to 70 barrels of 280 pounds each, and for succeeding years at 100 barrels.

Trees can be profitably worked in North Carolina by experienced operators during four or five years, or, upon a small scale, in connection with farming operations and by actual residents, several years longer; farther south the trees seem to possess less recuperative power, and in South Carolina four years is given as the outside limit during which an orchard can be profitably worked, while in Georgia, Florida, and Alabama they are often abandoned at the end of the second and always at the end of the third year. Twenty-five men, including overseers, wagoners, distillers, coopers, and laborers can work ten crops. The average wages of such a force is \$1 a day per man, so that the cost of labor necessary to work a crop during the season of thirty-two weeks is \$480.

The following grades of turpentine are recognized in the trade: "Virgin dip", or "Soft white gum turpentine"—the product the first year the trees are worked; "Yellow dip"—the product of the second and succeeding years, and becoming darker colored and less liquid every year; "Scrape" or "Hard turpentine"—the product of the scrapings of the boxes.

Rosin is graded as follows: "W"—Window-glass; "N"—Extra pale; "M"—Pale; "K"—Low pale; "I"—Good No. 1; "H"—No. 1; "G"—Low No. 1; "F"—Good No. 2; "E"—No. 2; "D"—Good strain; "C"—Strain; "B"—Common strain; "A"—Black.

Window-glass is the lightest grade, and is only produced from the first dippings of "virgin" trees—that is, trees worked for the first time. The resinous exudation becomes darker colored and less volatile every year, as the box grows older, and the rosin produced is darker and less valuable. Trees worked during several years produce a very dark brown or black rosin. Spirits of turpentine made from virgin trees is light colored, light in weight, and free from any taste; the resinous matter yielded in succeeding years gains more and more body, and the additional heat required in distilling it throws off some resin combined with the spirits, producing in it a strong, biting taste and greater weight.

Tar, produced by burning the dead wood and most resinous parts of the long-leaved pine in covered kilns, is graded as follows: "Rope yellow", or Ropemakers' tar—the highest grade, produced with a minimum of heat from the most resinous parts of the wood; "Roany," or "Ship smearing"—the next running of the kiln; "Black" or "Thin"—the lowest grade, made from inferior wood, or the last running of the kiln, and therefore produced with the maximum of heat.

The following statistics of the production of naval stores during the census year were prepared by Mr. A. H. Van Bokkelen, of Wilmington, North Carolina, to whom I am indebted for much information in regard to the methods used in carrying on this industry:

States.	Turpentine.	Rosin.
	Gallons.	Barrels.
Alabama.....	2,005,000	158,482
Florida.....	1,036,350	68,281
Georgia.....	3,151,500	277,500
Louisiana.....	250,000	20,000
Mississippi.....	250,000	20,000
North Carolina.....	6,279,200	663,967
South Carolina.....	4,593,200	333,940
Total.....	17,565,250	1,542,170

Eighty thousand barrels of tar were manufactured during the census year in North Carolina, and 10,000 barrels in the other southern states.

The total value of this crop of naval stores at centers of distribution, and of course including freight from the forest and different brokerage charges, was not far from \$8,000,000. The net profits of the industry, even in the case of virgin trees, is very small, and at present prices is believed to be unprofitable except to the most skillful operators. The low price of southern timber-lands and the facility with which rights to operate tracts of forest for turpentine have been lately obtainable in several states have unnaturally stimulated production. The result of this has been that manufacturers, unable to make a profit except from virgin trees, abandon their orchards after one or two years' working and seek new fields of operation; the ratio of virgin forest to the total area worked over in the production of naval stores is therefore constantly increasing. It is estimated by Mr. Van Bokkelen that during the years between 1870 and 1880 an average of one-third of the total annual product of the country was obtained from virgin trees, and that in 1880 one-fourth of the crop was thus produced, necessitating the boxing in that year of the best trees upon 600,000 acres of forest. The production of naval stores is carried on in a wasteful, extravagant manner, and the net profits derived from the business are entirely out of proportion to the damage which it inflicts upon the forests of the country; the injury is enormous. Lumber made from trees

previously worked for turpentine is of inferior quality, although it is probably less injured than has been generally supposed. Comparatively few trees, however, once boxed are manufactured into lumber. It is estimated that 20 per cent. of them, weakened by the deep gashes inflicted upon their trunks, sooner or later are blown down and ruined; fires, too, every year destroy vast areas of the turpentine orchards, in spite of the care taken by operators to prevent their spread. It is customary in the winter, in order to prevent the fires which annually run through the forests of the Southern Pine Belt from spreading to the boxes, to "racket" the trees; that is, to remove all combustible material for a distance of 3 feet around the base of each boxed tree. Fire, carefully watched, has then been set to the dry grass between the trees, in order to prevent the spread of accidental conflagrations, and to give the box-choppers a firmer foothold than would be offered by the dry and slippery pine leaves. In spite of these precautions, however, turpentine orchards, especially when abandoned, are often destroyed by fire. The surface of the box, thickly covered with a most inflammable material, is easily ignited, and a fire once started in this way may rage over thousands of acres before its fury can be checked.

The manufacture of naval stores, then, decreases the value of the boxed tree for lumber, reduces the ability of the tree to withstand the force of gales, and enormously increases the danger to the forest of total destruction by fire.

Wilmington, the most important distributing point for this industry in the United States, handles 80 per cent. of all the naval stores manufactured in North Carolina. Previous to 1870 Swansboro', Washington, and New Berne were also large shipping points.

SOUTH CAROLINA.

The forest covering of South Carolina resembles in its general features that of the states immediately north and south of it. The pine forest of the coast, nearly coinciding in area with that of the Tertiary deposits, covers the eastern portion for a distance of 150 miles from the coast. The middle districts are occupied with hard-wood forests, or forests in which pines of various species are mixed with oaks, hickories, and other deciduous trees. The forests of the Alleghanies, rich in species and magnificent in the development of individual trees, spread over the mountains and valleys, which occupy the extreme western part of the state. The streams which flow through the Coast Pine Belt, often bordered by wide, deep swamps, are ill-suited to lumber operations, and less serious inroads have therefore been made into the pine forests of South Carolina than into those of North Carolina or Georgia. The merchantable pine, however, has been removed from the immediate neighborhood of the coast, from the banks of the Little Pedee river, and from along the lines of railroad.


The most accessible hard-wood timber has been cut from the forests of the middle districts, although vast quantities still remain remote from railroads or protected in deep river swamps, inaccessible except during a few months of summer. The western counties still contain great bodies of hard-wood timber, yet undisturbed except to supply the wants of the scattered population inhabiting this almost inaccessible mountain region.


The manufacture of rough red and white oak split staves and headings for the European and West Indian trade, already an important industry in this state, is capable of large development; rice tierces and rosin barrels are also largely made in the coast region from pine. At Plantersville, in Georgetown county, and at other points along the coast quantities of hand-made cypress shingles are manufactured in the swamps.

During the census year 431,730 acres of woodland were reported destroyed by forest fires, with a loss of \$291,225. These fires were set by careless hunters, by the careless burning of brush upon farms, and by sparks from locomotives.


BURNING OFF DEAD HERBAGE.

The pine belt of the coast, in South Carolina as well as through its entire extent from Virginia to Texas, suffers from fires set every spring by grazers for the purpose of improving the scanty herbage growing among the trees of this open forest. These fires run rapidly over the surface stripped by the fires of previous years of any accumulation of vegetable material, without inflicting any immediate injury upon the old trees of the forest unless a turpentine orchard is encountered, when, the resinous surfaces of the boxes being once fully ignited, nothing can save the trees from total destruction. If the mature trees of the forest are not under normal conditions greatly injured, however, by this annual burning of the dead herbage beneath them, the forest itself, as a whole, suffers enormously from this cause. Slight and short-lived as these fires are, they destroy the vegetable mold upon the surface of the ground, all seeds and seedling trees, and all shrubbery or undergrowth, which, in protecting the germination of seeds, insures the continuation of the forest. They deprive the soil of fertility and make it every year less able to support a crop of trees, and in thus robbing the soil they influence largely the composition of succeeding crops. Few young pines are springing up anywhere in the coast region to replace the trees destroyed, but where seedlings protected from fire appear upon land long subjected to annual burning, they are usually, although not universally, of less valuable species, and not the long-leaved pine which gives to this forest its principal economic importance. These annual fires are slowly but surely destroying the value of the Southern Pine Belt. They destroy all seeds and seedling trees, the fertility of the soil, and its power to produce again valuable species.









The following estimates of the amount of long-leaved pine standing in the state were made up from information obtained from Mr. Edward Kidder, of Wilmington, North Carolina, in regard to that part of the state north of the Edisto river, and from Mr. W. G. Norwood, of Blackshear, Georgia, for the southern part of the state. They are based on what is believed to be less accurate information respecting the northern part of the state than has been obtained in regard to the pine forests of the other states, and allowance should be made for possible large errors. The estimates are, however, probably largely below the actual productive capacity of the pine forests of the state which may be expected to exceed by 25 or 30 per cent. the following figures:

LONG-LEAVED PINE (*Pinus palustris*).

Counties.	Feet, board measure.	Counties.	Feet, board measure.
Aiken	209,000,000	Kershaw	171,000,000
Barnwell	340,000,000	Lancaster	5,000,000
Beaufort	49,000,000	Lexington	76,000,000
Charleston	458,000,000	Marion	328,000,000
Chesterfield	183,000,000	Marlborough	191,000,000
Clarendon	332,000,000	Orangeburgh	465,000,000
Colleton	453,000,000	Richland	88,000,000
Darlington	337,000,000	Sumter	380,000,000
Fairfield	7,000,000	Williamsburgh	536,000,000
Georgetown	128,000,000	Total	5,316,000,000
Hampton	202,000,000	Cut for the census year ending May 31, 1880...	124,492,000
Horry	380,000,000		

The principal centers of lumber manufacture are Georgetown, Charleston, and various points in Hampton and Barnwell counties, where small railroad mills are located. Charleston and Georgetown are the distributing centers for naval stores manufactured in the state.

GEORGIA.

The northern counties of Georgia are covered with the forests of the Alleghany Mountain region, here and in northern Alabama reaching the southern limits of their distribution and considerably reduced in the number of species composing them, the pines, firs, beeches, and other northern trees being generally replaced by the broad-leaved species of the Mississippi basin. From the base of the mountains forests of oak mixed with pines extend southward, occupying the central portion of the state and mingling with the trees of the Maritime Pine Belt along its northern limits. In the southern and coast counties great areas of swamps are still covered with forests of cypress, protected by their inaccessibility from the attacks of the lumberman.

The merchantable pine in the immediate vicinity of the principal streams and along the lines of railroad has been removed, and serious damage has been inflicted upon the pine forests of the state by the reckless manufacture of naval stores. Vast areas covered with pine, however, still remain, while the hard-wood forests of the central and northern portions of the state contain a large quantity of the most valuable hard woods.

The manufacture of cooperage stock is still in its infancy, and this and other industries requiring an abundant and cheap supply of hard wood seem destined soon to reach an enormous development in the upper districts of Georgia and the other states of the south Atlantic division.

During the census year 705,351 acres of woodland were reported devastated by fire, with a loss of \$167,620. The greatest number of these fires was traced to carelessness in clearing land, to sparks from locomotives, and to hunters.

The following estimates of the amount of long-leaved pine standing in the state of Georgia May 31, 1880, were prepared by Mr. W. G. Norwood, of Blackshear, in that state, a timber viewer and expert of high standing. He obtained his results by dividing the whole pine belt into irregular regions over which the average cut per acre could be obtained, allowance being made for clearings, farms, areas of culled forests, streams, swamps, etc. The area in each of these regions, by counties, was measured upon a large-scale map and the standing timber computed. These estimates include merchantable pine still standing on land partly cut over, or which has been worked in the manufacture of turpentine. The boxed areas include nearly all the regions from which any pine has been removed, and extend beyond them in all directions into the uncut forests and along rivers and railroads.

Similar methods, practically, were adopted in preparing the estimates of the amount of pine standing in Florida and the other Gulf states. The results thus obtained are not, of course, strictly accurate, and are not supposed to be so. The estimates are intended to show the average productive capacity of the pine forests over large areas, and to indicate generally in what part of the state the principal bodies of pine still occur. Liberal allowance has been made in computing areas of swamp and cleared land, and it will probably be safe to add 10 per cent. to these estimates of the pine standing in any of the southern states.

THE FORESTS OF THE UNITED STATES.

The following is an estimate of the amount of pine timber standing in the state May 31, 1880 :

LONG-LEAVED PINE (*Pinus palustris*).

Counties.	Feet, board measure.	Counties.	Feet, board measure.	Counties.	Feet, board measure.
Appling	543,000,000	Floyd	19,000,000	Polk	36,000,000
Baker	134,000,000	Glascock	17,000,000	Pulaski	408,000,000
Baldwin	35,000,000	Glynn	47,000,000	Randolph	128,000,000
Berrien	410,000,000	Hancock	76,000,000	Richmond	21,000,000
Bibb	38,000,000	Haralson	21,000,000	Schley	28,000,000
Brooks	281,000,000	Harris	22,000,000	Screven	188,000,000
Bryan	60,000,000	Houston	191,000,000	Sumter	191,000,000
Bulloch	733,000,000	Irwin	488,000,000	Talbot	44,000,000
Burke	298,000,000	Jefferson	206,000,000	Tattnall	768,000,000
Calhoun	117,000,000	Johnson	291,000,000	Taylor	58,000,000
Camden	82,000,000	Jones	40,000,000	Telfair	598,000,000
Charlton	246,000,000	Laurens	1,064,000,000	Terrell	104,000,000
Clay	96,000,000	Lee	128,000,000	Thomas	311,000,000
Clinch	350,000,000	Liberty	238,000,000	Twiggs	84,000,000
Coffee	578,000,000	Lowndes	236,000,000	Upson	32,000,000
Colquitt	339,000,000	McDuffie	10,000,000	Ware	161,000,000
Crawford	45,000,000	McIntosh	65,000,000	Warren	80,000,000
Decatur	653,000,000	Macon	52,000,000	Washington	240,000,000
Dodge	417,000,000	Miller	164,000,000	Wayne	160,000,000
Dooley	394,000,000	Mitchell	379,000,000	Webster	48,000,000
Dougherty	90,000,000	Monroe	18,000,000	Wilcox	292,000,000
Early	299,000,000	Montgomery	791,000,000	Wilkinson	152,000,000
Echols	183,000,000	Muscogee	35,000,000	Worth	512,000,000
Effingham	6,000,000	Paulding	2,000,000		
Emanuel	956,000,000	Pierce	220,000,000	Total	16,778,000,000
Cut for the census year ending May 31, 1880 (excluding 28,335,000 feet cut in the region of short-leaved pine and mixed growth).					272,743,000

The principal centers of lumber manufacture are situated along the coast at Brunswick, Darien, Savannah, and Saint Mary's. Logs sawed at these points are now driven down the various streams for a considerable distance from the coast. Large quantities of pine lumber are also manufactured in different mills located along the lines of railroad in Appling, Polk, Floyd, and other pine counties. Savannah and Brunswick are the principal points of distribution of the naval stores manufactured in the state.

FLORIDA.

The forests of the Southern Pine Belt cover the state as far south as cape Malabar and Charlotte harbor. The long-leaved pine is replaced along the sandy dunes and islands of the coast by oaks (of which the live oak is alone of commercial importance), scrub pines, and palmettos, while a deciduous forest, largely of northern composition, occupies the high, rolling lands in a large part of Gadsden, Leon, Jefferson, and Madison counties. The pine forests gradually decrease southward in density and value, and south of latitude 29° N. are of little present commercial value. Forests of pitch pine (*Pinus Cubensis*), however, extend far south of the region occupied by the more valuable long-leaved pine bordering the coast and covering the low ridges of the Everglades. Great areas of swamp occur everywhere through northern and central Florida, covered with forests of cypress, red cedar, gum, and bordered with bays, magnolias, and other broad-leaved evergreens; while the hummocks or low elevations, covered with rich soil and everywhere common, bear oaks and other deciduous trees, often of great size.

South of cape Malabar and Tampa bay the character of the vegetation changes, and the North American arborescent species are replaced by the semi-tropical trees of the West Indies. These occupy a narrow strip along the coast, cover the keys and reefs, and spread over some of the hummocks of the Everglades. This semi-tropical forest is confined to the saline shores of the innumerable bays and creeks of the region, or to the coral and sedimentary calcareous formation of the keys and hummocks. The species of which it is composed are here at the northern limits of their range; individual trees are comparatively small and the forests of the southern extremity of the Florida peninsula are commercially unimportant, although sufficiently extensive and varied to supply the scanty population of this region with lumber, fuel, and material for boat-building and the manufacture of fishing apparatus.

The forests of Florida have not suffered greatly from fire. Much of the state is uninhabited and unfit for agriculture or grazing. The danger, therefore, of fires set in clearing land for farms spreading to the forest is less than in other parts of the south, while the numerous streams and swamps everywhere intersecting the pine forests and the natural dryness of the sandy ridges, thinly covered with vegetable mold, check the spread of fires when started.

During the census year 105,320 acres of woodland were reported as burned over, with an estimated loss of \$69,900. The largest number of these fires was set by grazers to improve the pasturage for their stock.

MAP OF GEORGIA


SHOWING THE DISTRIBUTION OF PINE FORESTS,
WITH SPECIAL REFERENCE TO THE
LUMBER INDUSTRY.

COMPILED UNDER THE DIRECTION OF
C.S.SARGENT, SPECIAL AGENT.

1881.

LEGEND

-  (Short Leaved and Loblolly Pine (Pinus taeda) and Tardus; intermixed with Hardwood and scattered Long Leaved Pine.
-  Long Leaved Pine (Pinus palustris)
-  (Region from which Merchantable Pine has been cut.



Julius Bien & Co. Ltd.

The following estimates, by counties, of the long-leaved pine still standing in Florida east of the Apalachicola river were prepared by Mr. A. H. Curtiss, of Jacksonville; those for west Florida by Dr. Charles Mohr, of Mobile, Alabama:

LONG-LEAVED PINE (*Pinus palustris*).

Counties.	Feet, board measure.	Counties.	Feet, board measure.	Counties.	Feet, board measure.
Alachua	525,000,000	Holmes	150,000,000	Putnam	121,000,000
Baker	144,000,000	Jackson	233,000,000	Saint John's	66,000,000
Bradford	133,000,000	Jefferson	23,000,000	Santa Rosa	213,000,000
Brevard	68,000,000	Lafayette	425,000,000	Sumter	103,000,000
Calhoun	81,000,000	Levy	346,000,000	Suwannee	622,000,000
Clay	77,000,000	Liberty	75,000,000	Taylor	213,000,000
Columbia	455,000,000	Madison	122,000,000	Volusia	59,000,000
Duval	67,000,000	Manatee	200,000,000	Wakulla	72,000,000
Escambia	90,000,000	Marion	315,000,000	Walton	409,000,000
Hamilton	311,000,000	Nassau	104,000,000	Washington	187,000,000
Hernando	142,000,000	Orange	87,000,000	Total	6,615,000,000
Hillsborough	162,000,000	Polk	210,000,000		
Cut for the census year ending May 31, 1881 (excluding 77,500,000 feet, estimated, grown in Alabama and sawed in western Florida).					208,054,000

In this estimate no account is made of timber remaining on lands which have been cut over, or of that injured by the manufacture of turpentine.

The principal centers of lumber manufacture are Pensacola, Millview, and Blackwater, in Escambia and Santa Rosa counties. The logs sawed here and at other points upon Pensacola bay are driven down the streams from the forests of Alabama, the accessible pine in this part of Florida having been long exhausted. A large amount of pine lumber is also manufactured at Ellaville, in Madison county, upon the upper Suwannee river, and at Jacksonville, Saint Mary's, and at various points upon the lower Saint John river. Logs driven from the lower Suwannee river are sawed at Cedar Keys, where are situated the most important mills in the United States devoted to the manufacture of red cedar into pencil stuff.

Jacksonville, Saint Mary's, and Fernandina are the largest centers of distribution for the naval stores manufactured in the state.

The following extracts are taken from Mr. Curtiss' report upon the forests of Florida:

"In visiting western Florida I have had particularly in view the examination of the timber of a part of the state which is unlike all others in physical conformation, and consequently in vegetation. This region differs but little from the country bordering the southern Alleghanies, and may perhaps be regarded as the southern terminus of the Appalachian range. It commences about 40 miles north of the Gulf of Mexico, and extends northward between the Chipola and Okaloconee rivers into southwestern Georgia and southeastern Alabama. North of this there is little to connect it with the southern mountains except the rugged banks of the Chattahoochee river. The surface is undulating, hilly, often precipitous. The soil, like that of the Piedmont region of Virginia and Carolina, abounds in red clay, and is therefore adapted to crops which do not succeed in other portions of Florida. The vegetation is extremely varied and interesting, comprising most of the plants of northeastern Florida, a large portion of those found in the Piedmont country and in the rich river bottoms of the interior, and a considerable number found only on the limestone with which much of this country is underlaid. In the river bottoms, which are inundated at seasons, there is found a great variety of trees, some of which attain a size probably not equaled elsewhere. In this small portion of the state of Florida is to be found nearly every species of tree growing within the limits of the state, except those semi-tropical species found on the coast south of Cedar Keys and Mosquito inlet. Fully fifty American arborescent species here reach their southern limit. A few species show marked diminution in size, and all northern species which extend southward of this Chattahoochee region here attain in Florida their largest dimensions.

"There are two trees in this region of particular interest, as they are not known to grow anywhere else; these are the stinking cedar (*Torreya taxifolia*) and the yew (*Taxus Floridana*). There is reason to believe that the *Torreya* occurs also along the Wakulla river, and perhaps elsewhere in the state, but there is no positive knowledge of its occurrence except along the Apalachicola river, on the limestone hills which border it at intervals on the east.

"The forests of this region are still almost intact. Some poplar and tulip wood is cut from the river banks for northern markets, but the valuable timber on these rich shores is as yet almost untouched. The country southwest of this region, though of very little agricultural value, contains an immense quantity of the best cypress timber, hardly yet disturbed by the lumberman.

"Two mills have recently been established at Apalachicola, one of which saws nothing but cypress lumber. The product of this mill is sent to New Orleans. As white-pine lumber must soon become scarce, the attention of dealers ought to be directed to southern cypress, which will prove a good substitute for it. Although there is plenty of valuable pine in this country the swamps render it somewhat inaccessible, and the mills at Apalachicola

are more easily supplied with logs rafted down the river from Georgia. Many hewed logs of large dimensions are shipped from this point. The country near Apalachicola in surface and timber growth is much like that of northeastern Florida, all the good timber having been cut.

"PENCIL CEDAR.

"The favorite variety of red cedar, of tall and straight growth, is becoming scarce, but there remains a large quantity of quality sufficiently good for pencils in nearly all sections of the state north of a line drawn from cape Canaveral to the north end of Charlotte harbor. There is no red cedar in southern Florida, the Dixon mill at Tampa having exhausted the supply within reach of that place; but new mills have been established near Webster, in Sumter county, and at the head of Crystal river, at present the best source of supply.

"CYPRESS.

"The main body of cypress in southern Florida is located in the 'Big Cypress', a region of which I have heard much from persons who were in an expedition which went through it during the last Indian war. They entered it at the 'Little Palm hummock', 18 miles northeast of cape Romano. Traveling east about 12 miles they came to the 'Big Palm hummock', when they turned and traveled nearly due north for six days, averaging 12 miles a day. Their guide then informed them that the cypress extended 12 miles farther north; so it would seem that the main body of the 'Big Cypress' has a length of about 85 miles and a width, as they think, of about 20 miles. The cypress grows in belts running north and south, the main central belt being about 6 miles wide and consisting of large timber. There are narrow strips of cypress and pine alternating with prairie, although probably two-thirds of the whole region is covered with cypress. According to these estimates there must be at least 1,000 square miles covered with cypress timber in this region, which in times of high water could be floated out by the numerous creeks and inlets flowing toward the Gulf. There are also large quantities of heavy cypress on the swampy borders of Peace creek, the Hillsborough river, the Withlacoochee, etc., many trees squaring from 2 to 4 feet.

"The long-leaved pine extends south to Prairie creek, in about latitude 27° N. The pine between Prairie and Peace creeks, which is sawed at the mill near Ogden, belongs to this species. Timber in this region is quite shaky, and from all reports it is evident that the yellow pine in Manatee, Orange, and Hillsborough counties is quite inferior, being mostly of the rough-barked, sappy variety called in this region bastard pine. The long-leaved pine occupies nearly the whole of the interior of the peninsula north of a line drawn from Charlotte harbor to cape Malabar. At its southern limit I saw trees which measured over 2 feet in diameter and which would furnish logs 30 feet long.


"Pitch pine (*Pinus Oubensis*) appears on the west coast at Margo, 10 miles north of cape Romano, and extends northward to Prairie and Fishhead creeks, being the only pine of this region. From Charlotte harbor northward it is confined to a belt from 10 to 15 miles wide, bordering the Gulf, extending to Tampa and as far northward as Pensacola, being also scattered through the interior. This tree seldom exceeds 2 feet in diameter or 50 feet in height, and will afford a great quantity of framing timber, although it will be probably generally used in the production of naval stores, for which it is nearly or quite equal to the long-leaved pine.

"One of the most important facts in regard to the pine forests of Florida is their permanence. Owing to the sterility of soil and the liability to inundation of most of the state, it is certain that but a very small portion of Florida will ever be cleared of its forest covering. Taking into consideration the great area covered with valuable pine forests, and the fact that there will be a continuous new growth if the spread of forest fires can be checked, only trees of the largest size being cut, it is evident that Florida will furnish a perpetual supply of the most valuable pine lumber."

The following notes upon the pine forests of western Florida were furnished by Dr. Charles Mohr, of Mobile, Alabama:

"The pine forests occupying the region between the valley of the Apalachicola river and the banks of the Choctawhatchee, and from the headwaters of the Chipola to the bay of Saint Andrew's, are yet mostly in their primeval condition and contain a vast body of valuable timber. The district between the Choctawhatchee and the Perdido is the seat of the oldest and most active lumbering industry of the whole Gulf coast. The numerous streams flowing through the pine forests of eastern Alabama to the large bays upon the coast of western Florida make fully 4,000 square miles of southeastern Alabama comparatively accessible and tributary to the region from which the lumber finds an outlet by way of the bay of Pensacola.

"The better class of the somewhat elevated and undulating timber-lands which surround Escambia, Blackwater, and Saint Mary de Galves bay were long since stripped of their valuable timber. These forests having been culled time after time during the last quarter of a century, are now completely exhausted. The low, wet pine barrens, with their soil of almost pure sand, which trend eastward along the shores of Santa Rosa sound and Choctawhatchee bay, have never borne a growth of pine sufficiently large to furnish more than a small supply of timber of very inferior quality. The ridges between the Choctawhatchee river and the Yellow river are also, for the most part, arid, sandy wastes, never yielding more than a few hundred feet of lumber per acre.



"The well-timbered portion of west Florida commences with the southern border of Holmes county. This region is now, however, nearly exhausted along water-courses large enough for rafting, while of late years canals and ditches dug into the forest afford facilities for floating timber growing remote from streams to the mills. According to those best informed regarding the amount of timber still standing in this section, there is scarcely enough left between the Escambia and Choctawhatchee rivers, in western Florida, to keep the mills on the coast supplied for another half-dozen years, even if the whole of the pine standing could be made available.

"The lumber business of Perdido bay is entirely concentrated at Millview, where three large saw-mills are established. The production of lumber commenced here in 1865, increasing rapidly from 10,000,000 feet, board measure, in that year, to three and four times that amount. All the lumber manufactured upon Perdido bay is sent to Pensacola by a railroad constructed for the purpose. Only about 400 pieces of hewed timber are shipped from Millview, although the railroad has carried an average of 37,000,000 feet of lumber annually to Pensacola, the maximum annual yield of the Millview mills having been 45,000,000 feet.

"Pensacola is the most important port of lumber export on the Gulf coast. During the year ending August 30, 1879, 403 vessels, of a combined capacity of 217,487 tons, carried from the harbor of Pensacola 3,090,469 cubic feet of hewed square timber, 3,769,527 cubic feet of sawed square timber, and 60,000,000 feet of sawed lumber, board measure. Of the squared timber four-fifths is shipped to Great Britain.

"The peninsula between the junction of the Escambia and the bay of Saint Mary de Galves is low, and, along the shore-line, bordered with marshes. The timber needed to supply the mills located upon the shores of these waters has during the past forty years been drawn from this region, and when new forests have replaced the original growth they have been cut over and over again, and still furnish a small amount of timber, as the turpentine-distiller has not followed the log-getter in these regions. The supply of timber here, however, at present is too small to be taken into account in view of the enormously increased demands of the mills. There are three large mills on Blackwater bay producing 40,000,000 feet of lumber a year. Three-fourths of this lumber is produced in the establishment of Messrs. Simpson & Co., near the mouth of the Blackwater river, at Bagdad, about half a mile below Milton. Mills sawing square timber are situated 20 or 30 miles above the mouth of the Blackwater and use mostly water-power. The mill of Messrs. Milligan, Chaffin & Co., on this river, 20 miles above Milton, sends 28,000 pieces of square sawed timber to Pensacola, averaging 32 cubic feet each; 5,000 such pieces are furnished by a few very small water-mills higher up, swelling the whole amount of square timber to 33,000 pieces. The last-named firm has acquired by purchase large tracts of public land along Black and Coldwater rivers. To reach the timber growing on their land a canal 20 miles long, with sluices that intersect the small tributaries of these streams, has been dug. By means of this canal a sufficient supply of logs is secured to keep the mill running through the year. The large manufacturers of Bagdad have adopted a similar system, and by these means, and by the construction of tramways tapping the more remote and isolated regions tributary to the waters of Black and Yellowwater rivers toward the northern part of the state, the exhaustion of the timber-lands through the whole breadth of western Florida, as far as the banks of the Choctawhatchee river, will certainly be accomplished before the end of the next five years. A sash, door, and blind factory located at Bagdad consumes a large amount of cypress lumber. This is procured from the mills situated along the shores of the upper Choctawhatchee bay, and is grown along the banks of the Choctawhatchee river. The cypress lumber is exclusively used in the manufacture of sashes, blinds, doors, moldings, and particularly in the construction of houses, of which every year a considerable number is shipped by the way of New Orleans to the treeless regions of western Louisiana and Texas. This establishment manufactures a large amount of fencing, the rails of cypress, the posts of red and white cedar, rounded and capped. This is shipped to New Orleans and to the settlements in southern Florida. Of late years it has commenced sawing pencil-boards of red cedar. The logs, of very superior quality, are obtained from the hummocks and bottom lands bordering upon the Choctawhatchee. The lumber for this purpose must be entirely free from knots, of even, close grain, the woody fibers perfectly straight. These logs are cut in sections 6 inches in length, and the carefully-selected pieces sawed into slabs 2 inches broad and a quarter of an inch in thickness. Fifty gross of these slabs are packed in a case, and the establishment produces about six hundred cases annually. These are mostly shipped to a pencil factory in Jersey City, a small number going also to Germany.

"The saw-mills situated on the shores of Choctawhatchee bay extend from the mouth of Alaqua creek to Freeport, and westward to Point Washington; the logs sawed at these mills are for the most part brought down by raft from the upper waters of the Choctawhatchee and its tributaries. The lumber sawed here is mostly long-leaved pine, with a small amount of cypress. The product of these mills is mostly shipped to New Orleans in small schooners carrying from 15,000 to 20,000 feet each. The capacity of the mills upon this bay is in excess of their production, the difficulty of obtaining logs causing most of them to remain shut during half the year.

"The causes which up to the present time have prevented the destruction of the pine forests about Saint Andrew's bay, which is traversed by one fine river and bordered by another, must be traced to the difficulty of navigating these streams and to the want of a convenient outlet to the Gulf at Apalachicola. There are few saw-mills upon this bay, supplying only the local demand, and even these are furnished with logs floated down the Chattahoochee from beyond the confines of the state."

SOUTHERN CENTRAL DIVISION.

ALABAMA.

The northern and northeastern portions of Alabama, embracing the foot-hills of the southern Alleghany mountains and the valley of the Tennessee river, are covered with a rich and varied forest growth of broad-leaved trees, in which oaks, hickories, ashes, walnuts, and cherries abound. South of the Tennessee river the rolling country is covered with oaks, through which belts of short-leaved pine occur. In Cherokee and Saint Clair counties isolated bodies of long-leaved pine appear, while a narrow strip of the same species stretches nearly across the state between the thirty-third and thirty-second degrees of north latitude. South of this central belt the country is again covered with forests of hard woods, which farther south, in the rolling pine-hill region, are mixed with a heavy growth of the long-leaved pine; and this species occupies, or once occupied, almost exclusively, outside of the numerous river bottoms, the sandy plain extending along the coast and reaching nearly 100 miles inland from the shores of the Gulf. Great regions of swamp covered with heavy forests of cypress occur in the southern part of the state, especially in the region watered by the lower Tombigbee and Alabama rivers.

The forests of northern Alabama still contain great bodies of hard-wood timber, although the demands of the rapidly-increasing iron industry located here have already stripped of their tree covering many of the low hills of northeastern Alabama. The best pine has been gathered from Mobile and Baldwin counties, in the neighborhood of Mobile bay, from the lines of railroads and the banks of streams heading in the southern part of the state and flowing to the Gulf through western Florida.

The pine forests of southern Alabama have long suffered from the reckless manufacture of naval stores.

During the census year 569,160 acres of woodland were reported destroyed by fire, with an estimated loss of \$121,225. Of these fires the largest number were set to improve grazing, or by careless farmers and hunters.

The manufacture of cooperage and wheel stock, furniture, and other articles of wood is still in its infancy in Alabama and the other Gulf states. Such industries, in view of the magnificent forests of hard wood covering great areas in this region and the rapid exhaustion of the best material in the north and west, must in the near future be largely transferred to the southern states.

The cypress swamps adjacent to Mobile bay yield a large number of hand-split shingles and give employment to many persons, principally blacks.


The following estimate of the amount of pine standing in the state May 31, 1880, was prepared by Dr. Charles Mohr, of Mobile, who carefully examined the whole pine region of the Gulf states:

LONG-LEAVED PINE (*Pinus palustris*).

Regions.	Feet, board measure.
East of Perdido river.....	4,055,000,000
West of Perdido river.....	2,000,000,000
In the region of mixed growth.....	10,000,000,000
In the Central Pine Belt.....	1,750,000,000
In the Coosa River basin.....	900,000,000
In the Walker County district.....	180,000,000
Total.....	18,885,000,000
Cut for the census year ending May 31, 1880 (including 77,500,000 feet, estimated, grown in Alabama and sawed in western Florida).	245,398,000

SHORT-LEAVED PINE (*Pinus mitis*).

In the Central Pine Belt.....	1,875,000,000
In the Coosa River basin.....	482,000,000
Total.....	2,357,000,000
Cut for the census year ending May 31, 1880, none reported.	



2

In this estimate no account is made of small timber standing on some 1,282,000 acres which have been cut over, and from which the merchantable pine has been practically removed, or on 600,000 acres injured by the manufacture of turpentine.

There are fewer pine trees per acre in the region of mixed growth than in the pine belt proper, with which it mingles on the north; but the individual trees being larger, the average amount of standing pine per acre is greater, although generally of poorer quality.

Mobile is still the principal center in the state for the manufacture of pine and cypress lumber; a large amount of pine lumber is manufactured also along the line of the railroads penetrating the pine belt in Etowah county, and considerable hard wood is sawed in counties bordering the Tennessee river for local use and northern shipment. Mobile is also the distributing point for the naval stores manufactured in the state.

The following notes upon the forests of Alabama are extracted from Dr. Mohr's report:

"THE MARITIME PINE REGION.

"West of Mobile the road traversed for a distance of over 5 miles the plain, or so-called 'second bottom', composed of a more or less tenacious or sandy yellow clay. It has an elevation above the alluvial of the river of 15 to 25 feet, and is bordered on the west by the ridges of the stratified drift, which extend to within 6 to 18 miles of the shore-line. Near the coast this plain, flat and devoid of drainage, forms for many miles the low, wet savannas sparsely covered with a stunted growth of long-leaved pines; near the estuaries it is interspersed with tracts covered with a black, light soil, rich in humus and bearing a luxuriant growth of broad-leaved trees associated with a few *Coniferae*, and with the wooded swamps which extend over the depressions about the base of the higher land, and follow the low, inundated banks of the numerous streams. The prevailing forest tree of this plain, now much cultivated in the vicinity of Mobile, is the long-leaved pine. Situations offering a moister and somewhat richer soil along the hummocks and gentle acclivities bordering the swamps and the bottoms of the water-courses are occupied by the loblolly pine. With this is often associated the pitch pine (*Pinus Cubensis*), which prefers, however, the more or less inundated and always wet, swampy forest, where its spreading crown towers above the gum trees and white cedars. Wherever in the plain the long-leaved pine has been cut down, this pitch pine principally and the loblolly pine spring up to replace it.

"Many acres can be seen in this region covered with thrifty seedlings of this pitch pine, and trees have sprung up, to my own knowledge, since 1865, which are now from 20 to 25 feet in height with a diameter of trunk of from 4 to 6 inches; and trees from 50 to 60 feet in height with a circumference of from 3 to 4 feet, forming quite extensive forests, may be seen upon the shores of the bay from which the primeval forest was removed about fifty years ago.

"Ascending the highlands of drift, with its porous soil composed of irregular strata of white or ferruginous sands, gravels, and pebbles interspersed with layers of clay, the home of the long-leaved pine, which here arrives at perfection and forms the entire forest growth over immense areas, is entered. Upon this formation, after the removal of the original forest, either the long-leaved pine takes possession again of the soil or is replaced by a more or less stunted growth of various species of oak (*Quercus Catesbei*, *cinerea*, *nigra*, *obtusiloba*, and *falcata*), the mocker-nut, and a few other small trees and shrubs. What the conditions are by which such a rotation is regulated is not apparent. It is no doubt much influenced by the conflagrations which annually sweep through the woods and which are particularly destructive to the young pines, but it cannot be explained solely upon that ground. I have, however, observed that the more broken lands with the same sandy character of surface soil, but with a more argillaceous subsoil more or less impervious to water, are mostly covered with this second growth of deciduous trees, and that the flat table-lands with either a sandy or gravelly soil are invariably covered again with a second growth of the long-leaved pine. Among such young growths of this species I have never been able to discover a single seedling of the other pines.

"CYPRESS SWAMPS OF THE TENNESSEE RIVER.—The river was extraordinarily high, the lowlands being overflowed to a depth of more than 10 feet. The torrents which had fallen during the past three weeks caused a heavier freshet than any that had been experienced since the spring of 1875. Since that year no such opportunity has been offered for getting heavy cypress timber from the depths of these swampy forests. No idle man was to be found on shore; everybody who could swing an ax, paddle a boat, or pilot a log was in the swamp engaged in felling and floating cypress timber. All the mill-hands worked in the swamps; fields and gardens were left untouched, and even clerks from the stores were sent to the swamps as overseers.

"We soon entered the deep, dark forest stocked with some fine and large cypress trees, and came upon two negroes, each standing in his little skiff, engaged in felling a tree of the largest size. It was astonishing to witness the steadiness and celerity with which they performed their work, considering the instability of their footholds in the narrow boats. Every stroke of the ax told at the designated place, and it took them scarcely longer to cut a tree in this way than if they had been working upon solid ground. The top of the tree when felled is sawed off close to the first limbs by one man working under water a single-handled cross-cut saw. Another, provided with a long pole armed with a sharp iron spike, seizes the trunk and tows it, with the aid of

the slow current, to one of the lake-like sheets of still water which, interspersed with streams, are so common in these lowlands. Here the trunks are made into rafts and can be floated down the river to the mills along the banks below after the subsidence of the flood. The greatest part of this large timber is only accessible during the time of a high stage of water, so that the energies of the whole population are devoted during the times of freshets to getting out as much of it as possible. The large number of logs harvested shows clearly with what activity the destruction of these treasures of the forest is being pushed; and the reports, as of heavy thunder, caused by the fall of the mighty trees, resounding at short intervals from near and far, speak of its rapid progress.

"In 1831 Mr. Vaughn found these cypress swamps untouched by the ax. At present their resources are so diminished by the inroads made upon them during the last twelve years that, with a prospect of a rapidly-increasing demand for cypress lumber in the near future, he judges that they will be completely exhausted during the next ten years. This opinion is shared by all mill-owners here, who believe that in less than that time their business must come to an end. There is no hope that the supply will be continued by the natural increase of young trees. It is rare to find small trees among the large specimens. Seedlings and saplings are not found in these deep, swampy forests, and only occur in the openings and upon the banks of water-courses. The fact that the almost impenetrable shade, excluding the admission of light and air to a soil almost constantly drenched with water, is unfavorable to the growth of a new generation of the cypress, threatens to exclude it from localities where formerly this tree attained its greatest perfection. In swamps open to the influences of light and air, and not liable to prolonged periods of inundation, a growth of seedlings and small trees, especially along the banks of the smaller tributaries of the larger streams, springs up. The extremely slow growth of the cypress, however, during all stages of its existence, even if young trees spring up, destroys all hope of an adequate supply of this timber to meet the wants of coming generations. Trees of small size are as frequently cut as large ones. Saplings from 4 to 12 inches in diameter even are cut and supply the farmer, the builder, and the mechanic with material for many useful purposes. Logs not over 30 inches in diameter, however, are not worked up in the Tensas mills, which only use logs of larger size, the saplings being sent in rafts with pine logs to the saw-mills of Mobile. It is rare that a tree over 3 feet in diameter is found perfectly sound. Trees above 4 feet through are almost always invested with signs of decay. No timber seems to be open to so many defects as that of the cypress. Many of the trees are 'wind-shaken'; that is, portions of the body of the wood have separated in the direction of the concentric rings, causing annual splits which extend throughout a great length of the trunk, and if occurring repeatedly in the same stick render it unfit for use. A considerable number of the larger trees are rotten in sections. Logs cut from such trees may appear perfectly sound at both ends, but are found hollow and rotten in the interior. The inspection of cypress logs requires great experience and care to protect the buyer from loss. But there is one disease which particularly affects this timber, the cause of which is a perfect mystery to all interested in the matter. (a) From the center of the tree outward, although never extending into the sap-wood, occur great numbers of spindle-shaped, narrow excavations with perfectly smooth, rounded walls more or less tapering toward the ends, parallel with the bundles of woody fibers and nearly regularly disposed in the direction of the annual rings of growth. These cavities vary from one-half an inch or less to a foot in length, and are found from a few lines to an inch in width. They are filled with a yellowish-brown powder, the result of decayed, woody substance, although the walls of the cavities appear perfectly sound and unaffected by decay. These excavations are called 'pegs', and timber so affected 'peggy' timber. The cavities have no communication with the surface apparently, and remain always inclosed within the surrounding belt of sap-wood. It is only in the case of very old trees that the larger cavities produced by the junction of the pegs sometimes reach openings produced by external decay or accident. Undoubtedly these pegs cause the large hollows so often found in the center of large-sized and apparently perfectly healthy trees. Some of the timber of medium-sized specimens is honey-combed with these pegs. Such peggy stuff is useful for poles and pickets, which are found not less durable than if made from solid lumber.

"Two varieties of cypress timber are recognized according to the color, firmness, and heaviness of the wood, and are known as white cypress and black cypress; the latter has darker, closer grained, and more resinous wood than the former, and will sink in water. Its weight makes impossible the transportation of black-cypress logs by floating under ordinary circumstances, and the lumberman, unable always to recognize these peculiarities of the wood in the standing tree, cuts a chip before felling, which thrown into the water indicates, by its floating or sinking, whether it is black or white cypress. Trees of the heavy variety are deadened during the months of August and September by cutting a deep ring through the bark, and in the spring of the second season the timber is found sufficiently light to float.

"The cypress region of southern Alabama, which must be regarded as one of the great resources of its forest wealth, commences upon Mobile river, about 16 or 18 miles above its entrance into Mobile bay, extending through the lowlands upon both banks of this river, in Baldwin and Mobile counties, where it covers an extreme area of from 75 to 80 square miles. It extends northward to the junction of the Alabama and Tombigbee rivers, covering

^a This injury to the cypress is caused by a fungoid plant not yet determined, although widely distributed along the Gulf coast.—C. S. S.

large tracts in the delta between them, follows northward the course of these streams, and covers the extensive swamps which border their banks and the mouths of their numerous tributaries. Upon the Alabama the cypress swamps extend to the lower part of Clarke county. Next to the Mobile River region the largest supply of cypress can be drawn from the extensive bottoms of the Tombigbee, about the mouth of Bassett creek, near Jackson. During the freshet of the present year (1880) a large number of logs from this vicinity will be sent to the mills on the Tensas.

"BALDWIN COUNTY.—A quarter of a century ago a pine forest, unequaled in the magnificence of its tree growth, and supposed at that time to contain an inexhaustible supply of timber, covered Baldwin county through its whole extent. To-day this forest, from the line of the Mobile and Montgomery railroad, along the eastern shore of Mobile bay, and along all the water-courses as far as Bonsecours bay, upon the Gulf, is entirely destroyed, and presents a picture of ruin and utter desolation painful to behold.

"The production of naval stores has been carried on in this region without regard to any of its future interests, and, the forest being exhausted, manufacturers have been driven to seek new fields of operation. In the old turpentine orchards, long abandoned, no young trees have sprung up. Too far remote to make it possible to get their timber to the saw-mills, the large trees which have sufficient strength to withstand the effects of the barbarous process of boxing drag out their precarious existence for years after the smaller and weaker trees have been laid low, and shade the ground sufficiently to prevent the start of a young growth. The wood of these old boxes, as dead pines are called, is, after the loss of their vitality, charged throughout with an excess of resinous matter, and is in that condition sold as 'fat' or 'light' wood, being greatly esteemed as fuel for the generation of steam. For this purpose this final product of the pine forest is carried to the city of Mobile in broad flatboats, propelled by one huge square sail, and steered by a ponderous horizontal beam serving as a rudder. In a few years, however, this, the least valuable and the last product of the pine forest, will have forever disappeared, and with it the last remnant of the original forest growth of this part of the state. Occasionally, under the shade of the trees left standing, a young growth of pine is found, and on the high and undulating table-land between Mobile bay and Fish river, where the soil is light and very porous, a low and scanty oak scrub has taken possession of the ground. Toward the banks of the water-courses, however, where the largest trees were first cut to furnish timber to the mills once situated on Fish river, thus early leaving the ground open to atmospheric influences, fine and promising groves of long-leaved pine now often cover areas of wide extent. I measured many trees in these young second-growth pine forests, grown up within the last twelve to twenty-five years, standing from 15 to 30 feet in height with a diameter of trunk of from 4 to 6 inches, of thrifty growth, and rapidly overcoming the small oak growth with which it had to contend for the possession of the soil. It is the turkey and the upland willow oak alone which occur in these thin soils, too poor to support the Spanish and black oaks.

"The banks of the North Branch of the Fish river are composed of marsh or white drift sand. The arid, sandy ground is covered with a dwarf growth of live oak and myrtle live oak, observed here for the first time, and which farther east formed by far the largest part of the oak scrub covering the shore-lines of the large bays of western Florida. Two or three miles beyond the forks of Fish river a belt of pine forest is reached, not yet destroyed by the mutilations of the 'box-cutter' nor bereft of its best growth by the log-gatherer; it covers the highlands and declivities between Fish river and the waters which find their way into Perdido bay. This may be regarded as a virgin forest, only slightly invaded up to the present time along the Blackwater creek, Hollenger's creek, the Perdido river, and the bay shore. The mills situated on Perdido river and bay depend entirely for their present and future supply of logs upon this forest of southern Baldwin county, although I learn that it is expected to supply them during the next five years only, even if their production of lumber does not increase. This forest extends over six townships and covers an area estimated at from 125,000 to 150,000 acres.

"THE FORESTS OF THE CHATTAHOOCHEE IN EASTERN ALABAMA, MIXED FOREST GROWTH, ETC.

"The forests which once covered the wide bottom lands of the Chattahoochee in the neighborhood of Franklin, Alabama (opposite Fort Gaines), are now reduced to small patches of woodland confined to the base of ranges of low hills bordering the plain valley to the southeast. The tree growth was found here to differ in no way from that found lower down, except that the short-leaved pine (*Pinus mitis*) occurs more frequently. The crab apple and the cockspur thorn are frequent along the borders of the woods, but the pond pine (*Pinus serotina*), which might have been expected here, was not observed. In the sandy, wet, and deeply-shaded bottoms of a sluggish stream winding along the base of these hills I found the spruce pine (*Pinus glabra*) abundantly associated with the loblolly bay, red and sweet bays, and stately magnolias. The live oak is not found here, and it is doubtful if it extends in this part of the Gulf region more than a few miles north of the thirty-first degree of latitude. The low hills do not rise more than 150 feet above the plain; in entering them the second division of the sylvan vegetation characteristic of the eastern Gulf states is reached—a forest of mixed growth, which must be regarded, on account of its extent as well as the variety of its vegetation, as one of the important natural features of the region. I am of opinion that the deciduous-leaved trees have an equal representation in this forest with the

conifers. This certainly was the case before the settlement of the country, but as the broad-leaved trees occupy the best land, the areas of hard-wood forest have been more reduced by the demands of agriculture than have the forests of pine.

"The distribution of the different species of trees throughout this region depends upon the nature of the soil and the topographical features of the country. In general it can be stated that the marls and calcareous Tertiary strata which form the lower ridges and more or less undulating uplands and plains are chiefly occupied by trees with deciduous leaves, and by a few yellow pines. Here oaks predominate, and especially the post oak (*Quercus obtusiloba*), which prefers the level or gently-swelling ground with a generous, warm, and open soil; with it is frequently found the black oak (*Quercus tinctoria*), the Spanish oak and black-jack upon soils of poorer quality, the last, particularly, preferring one of closer, more argillaceous character mixed with fine sand. The black-jack finds here its best development, rivaling often in size the post oak; it enters largely also into the undergrowth of the post-oak woods, forming dense thickets on lands too poor to sustain a heavier tree growth.

"The hickories are unimportant features in the forests of this region. In the dry uplands they seldom attain more than medium size, although in the more shaded and richer situations the mocker-nut and pig-nut are not rare.

"The long-leaved pine, on account of the broad extent it covers, its gregarious habit, and the splendid growth it attains here, must be regarded as the most important timber tree of this region. Confined to a siliceous, dry, and porous soil, it occupies the high ridges invariably covered with a deposit of drift, often found widely spread over the more elevated highlands. For this reason the pine forests crown the hills and cover the more or less broken plateaus. They are found also toward the southern boundaries of this region, where the sands and gravels of the drift of the lower pine region encroach upon and mingle with the strata of older formations. Under these circumstances it is evident that the line of demarkation between this and the pine region of the coast is difficult to determine. The best distinction is found in the fact that in the pine forests of the lower pine region the growth of pines upon the uplands is never broken by patches of oak, and that the short-leaved pine never occurs there. Another point of distinction is found in the nature of the second growth, which springs up after the large pines have been removed. In the pine woods in the region of mixed tree growth the subsoil, of Tertiary origin, seems more favorable to the growth of oaks than to a second growth of the long-leaved pine. This is replaced generally by oaks mixed with the short-leaved pine and various deciduous trees. It is safe to assert that the southern limits of this region coincide with a line following the northern boundary of the coast drifts, along which the lower strata have completely disappeared beneath it.

"PIKE COUNTY.—On the broad ridges which form the divide between the waters of the Pea and Conecuh rivers, upon a purely sandy soil, are found, within the forest of long-leaved pine, tracts with strictly-defined outlines from a half mile to several miles in width, covered with a dense vegetation of small trees and shrubs peculiar to the perpetually moist and cool hummocks of the coast. The soil covered with this growth presents no unusual features; it is as poor and arid as that covering the rest of these heights. Surrounded on all sides by pine forests, not a single pine tree is seen within the limits of these glades, called by the inhabitants 'pogosines', an Indian name the meaning of which I was unable to learn.

"The trees are of small growth, the willow oak, the water oak, beech, red maple, and black gum rarely rising to a height of more than 30 feet among the sourwoods, junipers, hornbeams, hollies, papaws, fringe-trees, red bays, and other trees of the coast. These glades verge upon deep ravines from which issue large springs, and from this fact I conclude that, below their sandy, porous soil, strata must exist perpetually moistened by subterranean waters near enough to the surface to supply the moisture necessary to support such a luxuriant vegetation.

"FORESTS OF THE TENNESSEE VALLEY.

"The character of the forest vegetation changes upon the limestone formation of the valley of the Tennessee. This new region of tree growth extends from the northeastern confines of Alabama to a short distance beyond the Mississippi state line with a width of from 35 to 40 miles, and reaching beyond the northern boundary of the state. Its prominent feature is the total absence of pine and the scarcity of other evergreen trees. A few scattered saplings of the loblolly pine are found on its lower borders, waifs strayed from their natural habitats, the lower part of Morgan county, the true northern limit of this species, in Alabama at least. The red cedar is the only evergreen tree common among the forest growth of this limestone region, and the durability of its wood combined with its beauty places this tree among the most useful produced in this region. The red cedar forms here almost exclusively the second growth after the removal of the original forest, covering everywhere with extensive groves the dry, rocky hillsides and flats. The timber, however, of this second growth is only fit for the most ordinary purposes. The trees branch low, and the trunks are consequently full of knots and unfit for anything except fence posts. The fertile portions of this region have been largely denuded of their forest growth, although more than half is still covered with wood, a considerable portion with almost virgin forest. This is particularly true of Lauderdale and Colbert counties and the mountainous portions of the counties of Madison and Jackson. The vast quantities of oak, ash, walnut, and poplar timber contained in these counties can be sent to northern markets as soon as the Tennessee river has been made navigable by the removal of the obstacles at the Mussel shoals.

"The road from Decatur to Moulton, in Lawrence county, leads through broad and fertile valley lands, broken, as the mountains are approached, by limestone ridges jutting out into the plain. The beautiful Moulton valley, inclosed by the low foot-hills of the Sandy Mountain range which form its southern boundary, shows only along the base of the mountains a remnant of its original tree covering. Here the water oak, willow oak, red oak, mulberries, elms, and ashes were the trees found in the lower situations, and on rolling, higher land the white oak, the black oak, post oak, sassafras, and dogwood formed the prevailing forest growth. The lower flank of the steep escarpment of the highlands, a terrace of limestone cliffs mostly destitute of soil, bears a stunted tree growth. Here the red cedar and the upland hickory abound, and where the surface is less broken and a deeper soil covers the rock, chestnuts make their appearance with white oaks and the shell-bark and mocker-nut hickories. The ascent is less precipitous as the sandstone ledges are reached, and here the yellow pine (*Pinus mitis*) and the scrub pine (*Pinus inops*) are prominent among the oak forests of the mountains. When the crest of this abrupt decline is passed the oak forest is reached. It covers the extensive table-land between the Coosa and the eastern tributaries of the Tombigbee, and extends southward from the valley of the Tennessee to the lowlands commencing below Tuscaloosa, occupying an area of nearly 6,000 square miles.

"GENERAL REMARKS.

"The forests of long-leaved pine are principally confined to the following limited regions east of the Mississippi river: 1. The Great Maritime Pine region. 2. The Central Pine Belt of Alabama. 3. The Pine Region of the Coosa.

"Pine forests of more or less extent, too, mixed with woodlands composed of deciduous-leaved trees, occupy the ridges covered with a porous siliceous soil in the region of what I have called the mixed tree growth, and which upon its southern borders verges upon the Coast Pine Belt. Upon the heights of the low ranges of the metamorphic region of Alabama are also found more or less extensive tracts of this pine, generally, however, of inferior quality and size, while as far north as the thirty-fourth degree of latitude patches of thinly-scattered pine are met on the brows of the mountains, and, rarely, on the plateau of the carboniferous sand.

"The pine forests of Alabama, from the Escambia to the Mississippi state line, in the counties of Monroe, Baldwin, Washington, Mobile, and in portions of Clarke county, cover 3,500 square miles. Of these about 1,000 square miles have already been more or less destroyed in the manufacture of naval stores. Allowing 25 per cent. for land under cultivation, or covered by a forest of different trees, by water, etc., there are still 1,875 square miles left of this forest to supply the demands of the future.

"The whole amount of long-leaved pine lumber received at the port of Mobile averages about 60,000,000 feet, board measure, representing the product of mills at that place and along the various railroad lines leading to it. The amount of hewed square timber received is still small, but the business of exporting timber of this sort promises to assume large proportions in the near future.

"THE PINE BELT OF CENTRAL ALABAMA.—This forest occupies the deposits of drift which, in a strip varying from 10 to 30 miles in width, traverses the state from east to west. It is nearly in the center of the line connecting its eastern and western limits that its greatest width is found. This forest is estimated to cover 550 square miles, no allowance being made for lands cultivated or covered by other trees. The timber, both in quality and quantity, is unsurpassed by that growing on the best sections of the lower pine region. The manufacture of lumber and its export to northern markets has only been carried on in this region to any large extent during the last three or four years, and it is now rapidly assuming large proportions. The most important saw-mills in this region are situated on the line of the Louisville and Nashville railroad, between Clear creek and Elmore, Elmore county, and produced in the aggregate 67,000,000 feet of lumber, board measure, during the years 1879-'80. Considerable lumber is also produced along the line of the Selma, Rome and Dalton railroad, in Chilton county.

"Naval stores are not yet manufactured in this region.

"THE PINE REGION OF THE COOSA.—A detached belt of drift largely composed of coarse pebbles stretches from the eastern base of the Lookout Mountain range through the valley of the Coosa river, near Gadsden, covering nearly the whole of Cherokee county, to the Georgia state line. This forest is estimated to cover from 400 to 450 square miles, although much of the best timber nearest to the river has already been exhausted. Logs are driven down the Coosa and sawed at Gadsden. The manufacture of lumber at this place has been carried on for a number of years, and amounts to an average of 20,000,000 feet.

"NAVAL STORES.

"The manufacture of naval stores in the central Gulf states is almost entirely restricted for the present to the forest contiguous to Mobile and to the railroad lines leading to that port and to the southern confines of the pine belt in Mississippi. It is only during the past two seasons that turpentine orchards have been worked near Pascagoula, Mississippi, Pearl river, and in eastern Louisiana above Covington. The first turpentine distilleries were established on the Gulf coast a little more than a quarter of a century ago, along Fish river on the eastern and Dog river on the western shores of Mobile bay. The business soon assumed such proportions as to lead to the destruction of the

forests covering hundreds of square miles, particularly in Baldwin county. The production of naval stores in this county, as well as in the lower part of Mobile county, has at present nearly ceased, on account of the exhaustion of the forest. It is, however, now carried on with the greatest activity on the line of the Mobile and Ohio railroad. Between Mobile station, in Mobile county, and Quitman, Mississippi, there are at this date not less than thirty-three stills in operation, while along the Louisville and Nashville railroad there have been during the last five years fifty-three stills established in Alabama and Mississippi. These, with few exceptions, are controlled by Mobile capital, their whole product being handled from that market, so that the returns contained in the annual reports of the board of trade of Mobile fairly represent the whole production of naval stores in this pine region.

"According to the statements contained in the report for 1880, the crops amounted in the years 1879-'80 to 25,409 barrels of spirits of turpentine and 158,482 barrels of rosin. During a period of eight years, between 1873 and the close of the business year of 1880, 160,000 barrels of spirits of turpentine and 800,000 barrels of rosin have been produced in this same district. (a)

"The increase in prices during the last few years for all kinds of naval stores, and particularly the active demand for the best class of rosin, have given an increased impetus to this business, in consequence of which many of the older orchards have been abandoned and new ones started, while the number of new boxes cut during the present season is greater than ever before. There are no returns to be obtained of the production prior to 1875, but it can be safely assumed that up to that year 250 square miles of pine forest had been boxed. The production since 1875 must have involved a further destruction of 640,000 acres, or 1,000 square miles of forest. With the low price at which pine lands are held there is not the slightest regard paid to the utilization of their resources, and under the present system they are rapidly destroyed, regardless of the needs of the future and with the sole object of obtaining the quickest possible returns on the capital invested.

"It may be of interest to mention here the results obtained by a practical manufacturer by submitting the refuse of saw-mills, that is, slabs and sawdust, to a process of combined steam and dry distillation, with the view of utilizing the volatile products of such waste. He obtained from one cord of slabs 12 gallons of spirits of turpentine, 25 gallons of tar, 120 gallons of weak pyroligneous acid, and 12 barrels of charcoal. From one cord of lightwood he obtained 12 gallons of spirits of turpentine, 62½ gallons of tar, and 60 gallons of pyroligneous acid. The sawdust obtained from sawing 10,000 feet of pine lumber, subjected to distillation during one day, produced 22 gallons of spirits of turpentine."

MISSISSIPPI.

The forests of Mississippi originally extended over nearly the entire state. Prairies of no great area, situated in the northern central part of the state, presented the only break in its tree covering. The forest consisted of a belt of long-leaved pine, occupying the coast plain and reaching from the eastern confines of the state to the bottom lands of the Mississippi river, and from the coast nearly to the line of Vicksburg and Meridian. The northeastern portion of this long-leaved pine forest spread over a high rolling country, and here the pines were mixed with various hard-wood trees; north of the long-leaved pine forest a long belt gradually narrowing toward the north and occupied by a growth of short-leaved pine and of hard woods reached nearly to the northern boundary of the state, while south of the Tennessee river, in Tishomingo, Prentiss, and Itawamba counties, a considerable area was covered with forests of the short-leaved pine. The remainder of the state was clothed with a growth of hard woods, which in the swamps of the Yazoo delta and the bottom lands of the Mississippi river formed vast and almost impenetrable forests, where cypresses, gums, water oaks, ashes, and other trees which find their home in the deep, inundated swamps of the South Atlantic region attained noble dimensions and great value.

The pine forests have been removed from the immediate neighborhood of the Pascagoula and Pearl rivers and from their principal tributaries within the southern tier of counties; the most accessible timber has been cleared from the Biloxi, Blind, Jordan, Wolf, and Tchefuncta rivers, flowing into Mississippi sound, and from the line of the Chicago, Saint Louis, and New Orleans railroad. The long-leaved pine of Mississippi is, however, still practically intact, and these forests are capable of supplying an immense amount of timber as soon as the means of transportation can be furnished for it. A small amount of pine has been cut in the northeastern pine region from along the line of the Memphis and Charleston railroad.

The hard-wood forests outside of the bottom lands have been largely cleared from many counties in providing for the requirements of agriculture. Such land when abandoned is again covered in the central part of the state with a growth of old-field pine, and in the north, and especially in the northeastern counties, by a vigorous growth of short-leaved pine (*Pinus mitis*), which seems destined to become the most important timber tree of that region. The forests which cover the swamps of the state are still almost intact, although the most accessible cypress, which has long been cut in the Yazoo delta and the valley of the Pearl river to supply the New Orleans market, has become scarce.

During the census year 222,800 acres of woodland were reported destroyed by fire, with a loss of \$78,500. Of these fires the largest number was set by hunters, and by farmers carelessly starting fires in clearing land or to improve pasturage.

a These figures differ somewhat from those prepared by Mr. Van Bokkelen. See page 493.—C. S. S.

MAP OF MISSISSIPPI

SHOWING THE DISTRIBUTION OF THE PINE FORESTS.


WITH SPECIAL REFERENCE TO THE
LUMBER INDUSTRY.

COMPILED UNDER THE DIRECTION OF
C.S. SARGENT, SPECIAL AGENT.

1881.

LEGEND.

-  Short Leaved or Yellow Pine (*Pinus mitis*)
-  Short Leaved or Yellow Pine (*Pinus mitis*) mixed with Hardwoods
-  Long Leaved Pine (*Pinus palustris*)
-  Long Leaved Pine (*Pinus palustris*) with Hardwoods in about equal proportion
-  Region from which Merchantable Pine has been cut
-  Cypress Swamps
-  Turpentine Orchards



Scale

Julius Horn & Co. Lith.

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

Establishments for the manufacture of wagons, wheel stock, cooperage, etc., have been established at different times in the northern part of the state. The industries, however, which depend upon the hard-wood forests for material are still in their infancy in Mississippi, and are capable of enormous development.

The following estimates of the standing-pine supply of Mississippi, May 31, 1880, were prepared by Dr. Charles Mohr, who carefully explored the forests of the state:

LONG-LEAVED PINE (*Pinus palustris*).

Regions.	Feet, board measure.
In region west of Pearl river, tributary to the Chicago, Saint Louis, and New Orleans railroad.	6,800,000,000
East of Pearl river.....	7,600,000,000
Region of mixed growth, exclusive of 200,000 acres injured by the manufacture of turpentine.	3,800,000,000
Total.....	18,200,000,000
Cut for the census year ending May 31, 1880.....	108,000,000

SHORT-LEAVED PINE (*Pinus mitis*).

In the northeastern belt	1,600,000,000
In northern region of mixed growth.....	5,175,000,000
Total	6,775,000,000
Cut for the census year ending May 31, 1880.....	7,775,000

In this estimate no account is made of small timber standing on some 2,912,000 acres which have been cut over, and from which the merchantable pine has been practically removed.

The region of mixed growth, which adjoins the pine belt upon the north, contains a smaller number of pine trees per acre than the pine belt proper; but, the individual trees being larger, the average amount of standing pine per acre is here greater, although generally of poorer quality, than nearer the coast.

The principal centers of lumber manufacture are at the mouth of Pascagoula river, in Jackson county, at Mississippi City, in Harrison county, along the lower Pearl river, upon the line of the Chicago, Saint Louis, and New Orleans railroad in Lincoln county, and in the northeastern counties, where are located many small railroad mills, manufacturing in the aggregate a large amount of yellow-pine lumber (*Pinus mitis*).

The pine forests of the state have up to the present time suffered but little damage from the manufacture of naval stores. Turpentine orchards, however, have been recently established in the vicinity of the coast, near the mouth of the Pascagoula river, and at other points in the coast counties.

The following remarks are extracted from Dr. Charles Mohr's report upon the forests of Mississippi:

"THE PINE FORESTS OF SOUTHERN MISSISSIPPI.—In the vicinity of Scranton, near the mouth of the Pascagoula river, little is left of the original pine forest. The old clearings are covered with fine loblolly pine, from 40 to 60 feet high, upon rather close, dry soil. The pitch pine (*Pinus Cubensis*) forms dense groves, with seedling trees from 20 to 30 feet in height upon lands of lighter soil extending to the sea-shore. Oaks are not common. Fine groves of stately live oaks, however, line the banks of the river up to Moss Point, 4 miles distant.

"The annual export of lumber during the last four or five years has averaged 45,000,000 feet from the Pascagoula river. The largest percentage of this lumber is manufactured into boards and scantling for ordinary building purposes, and is shipped to Cuba, the Windward islands, to Mexico, Brazil, and a small part, in the form of deals 2 or 3 inches in thickness, intended for ship-building, to France, Spain, Holland, Belgium, and Germany. Large quantities of charcoal burned upon the banks of Black and Red creeks are sent to New Orleans in small coasting schooners, which run also from the bay of Biloxi and the bay of Saint Louis. At Moss Point eleven saw-mills, which furnish the lumber manufactured upon it, are situated on both banks of the East Pascagoula river. The combined capacity of these mills amounts to 220,000 feet a day, although the annual production during the past years has scarcely exceeded 40,000,000 feet. The timber manufactured in these mills comes from the Pascagoula and its tributaries, the Leaf and Chickasawha rivers and their sources, the Bogue Homo, Tallahala, Bay, and Okatuma creeks, as far up as the southern limits of Covington and Jones counties. A small number of logs also comes from the Escatawpa. The logs received at these mills average 20 inches in diameter and 40 feet in length. Sticks of such average dimensions are only furnished from first-class timber-lands, which, according to the best judges, produce six or seven trees of that size to the acre. Only lands lining the streams just mentioned, in a belt not exceeding 3 miles in width on each bank, have been up to this time invaded by the log-getter to supply these mills.

"The vastness of the timber resources yet contained in the region embraced in the northern half of Harrison and the whole of Greene and Perry, up to the southern confines of Marion and Jones counties, is astonishing. As is the case in Alabama, however, trees furnishing first-class spars for masts are difficult to find; they have been cut by spar-hunters in every part of the forest which could be reached by teams.

"Cypress lumber is not manufactured in this region, and the loblolly pine furnishes so small a part of the timber manufactured that it need not be considered. In Jones and Covington counties, about the headwaters of the upper tributaries of the Pascagoula, the country is rolling, intersected by numerous small, swift streams and rivulets. This region is magnificently timbered, and devoid of the barren ridges of almost pure sand so frequently found in the pine belt of Alabama.

"The low, flat, more or less wide pine lands bordering upon the marshes of the coast are sparsely covered with pine, while the trees growing in this wet, boggy soil, devoid of drainage and overlying a subsoil impervious to water, are stunted and of little value. The lower part of Harrison county is covered with these pine meadows, which fact accounts for the comparatively small importance of the bay of Saint Louis as a lumber-producing center.

"At Pearlinton, on the Pearl river, is established the large saw-mill of Poitevent & Favre, capable of producing 100,000 feet of lumber a day; at Logton, 2 miles farther up the river, are two mills, and 5 miles above these, at Gainesville, there is another. The largest part of the logs sawed at these mills is cut upon the banks of the Abolochitto creek, in Hancock county, and its tributaries extending into the lower part of Marion county, 50 or 60 miles distant. The remainder comes from the banks of the Pearl and the upper and lower Little rivers, which empty into it 10 miles above Columbia.

"The cypress is nearly exhausted from the lower Pearl river, and the 20,000 or 30,000 feet of this lumber which are sawed annually at Pearlinton are derived from the cypress swamps on the upper waters of the Pearl and Jackson rivers, where there is still a large amount of this timber of good size.

"The eastern bank of the Pearl river, within the Maritime Pine Belt, is sparsely settled, and forests, the especially in Hancock county and the upper part of Marion county, are unsurpassed in the quality and quantity of their pine timber. It is estimated by good judges that these forests will yield an average of 2,000 feet of lumber, board measure, to the acre. Up to the present time a strip of land scarcely 3 miles in width, embracing the banks of the water-courses, has been stripped of its timber growth, and fine spar timber is yet to be found here a few miles back from all the streams. Almost the whole of these rich timber-lands supplying the mills on Pearl river form a part of the public domain.

"The almost unbroken pine forests covering the upper tier of counties between the Pearl and Pascagoula rivers, toward the northern confines of the pine region, are still practically intact. The wealth of these forests has as yet found no outlet to the markets of the world. Thinly settled, they are still largely the property of the government, but in view of the speedily-increasing demand for lumber and the profits derived from the lumber business, such a condition of affairs must soon come to an end. It can be safely asserted that by far the largest part of the timber, felled in the Abolochitto region is taken from government land. There can be no question of this when it is considered how insignificantly small is the area of land which has been legally entered by private persons along that stream. The necessity of adopting proper measures to protect the timber wealth upon the public domain from depredations of such enormous extent forces itself upon the most casual observer, while to one who looks closer at the consequences of the continuance of the existing state of affairs the urgency becomes appallingly apparent. The ever-increasing consumption of timber at the mills upon Pearl river, of which one alone can cut 100,000 feet of lumber a day, will prove a powerful stimulus to a people who, since the development of the lumber business in these regions, have almost completely abandoned their former agricultural and pastoral pursuits and now depend entirely for their support upon cutting pine logs, to supply this enormous demand at the expense of the public property. Already plans have been made to invade this region by tramways and railroads, in order that its timber may be brought to market. This is true, too, of the region between the Pearl and the Amite rivers, down to the marshy lands of eastern Louisiana, a region in which the forests are also particularly good.

"In the state of Mississippi it is safe to estimate that, after deducting 25 per cent. for areas of swampy and cleared land, 9,000 square miles are still covered by forests of long-leaved pine. The production of this region during the census year amounts to 108,000,000 feet; of this, 60,000,000 finds its outlet at Pascagoula, 30,000,000 by Pearl river, 6,000,000 by bay of Saint Louis, and 12,000,000 by the Chicago, Saint Louis, and New Orleans railroad to northern markets.

"In the northern part of Harrison county we crossed a tract from which twelve years ago a hurricane swept a belt a quarter of a mile wide of all tree growth. It is interesting to note the growth which has since sprung up among the prostrate charred trunks of the pines still found lying about in large numbers. Black-jack oaks, the largest not over 12 feet in height, are mixed in almost equal numbers with stunted, thin saplings of the long-leaved pine. These plainly exhibit the helplessness of the struggle to which these offspring of the great timber tree are subjected under the influence of repeated conflagrations wherever the oak scrub has sprung up and added fuel, in the abundance of its leaves, to the fires which annually sweep through these woods.

"THE NORTHEASTERN COUNTIES.—After crossing the Sucarnoochee river below Scooba, in Kemper county, the pines which had covered the ridges near the borders of Lauderdale county disappear; scarcely a stray sapling

of the loblolly pine is seen as Scooba is reached. The cold, wet, calcareous soil of the flatwoods and prairies is unsuited to the growth of all coniferous trees, with the exception of the cypress. Along the railroad, as it traverses the flat prairie region, the country is sparsely wooded; large tracts of the prairie lands have always been destitute of trees, and the woodlands with which they were interspersed were cleared at the first settlement of the country. What remains of the original forest growth is now confined to localities too difficult of drainage to make agriculture profitable, and to the banks of streams subject to inundation. More or less extensive patches of woods are found also on the ledges where the limestone rock comes to the surface. In the swampy land the willow oak, the water oak, the black gum, sweet gum, white ash, and along the ponds willows and cottonwoods, prevail. The post oaks, white oaks, and cow oaks are mingled more or less freely with these trees in localities enjoying better drainage. Black-jack and black oaks, mixed with various haws, viburnums, and persimmons, occupy the rocky flats. No magnolias were seen in this region. The red, willow, and water oaks, the sycamore, and the sweet gum abound along the streams here, and are so common as to deserve special mention, while on the rolling uplands black oaks, post oaks, and white oaks, with poplars, shell-bark and pig-nut hickories, are common. From Tupelo toward Corinth the country is poorly wooded. The ascent is constant, reaching the point of highest elevation between the Gulf of Mexico and the Ohio river at Booneville. Corinth is situated on a wide pine plain, bounded on the west by the valley of the Tuscumbia river and east by the ridges which mark the water-shed of the Tennessee. The soil is here a deep calcareous clay, very stiff and heavy, hard as brick in warm, dry weather, and suddenly becoming a bottomless, stiff mire in seasons of rain. Below the valley of the Tuscumbia river the road passes over low and undulating ridges, of which the higher and steeper are yet covered with the remnants of the old oak forest. Here the Spanish and post oaks predominate in numbers; then follow the black oak and the scarlet oak, while the shell-bark hickory and the mockernut form but a small part of the tree growth of these uplands. The bottoms of the Tuscumbia, although subject to frequent overflows, are covered with a primeval forest not inferior in luxuriance and variety to that of the Mississippi river bottom lands. White-oak timber of the finest quality is found here in the greatest abundance and perfection. The most common species is the cow oak (*Quercus Michauxii*). I found that this river-bottom forest contained, by actual count, an average of from twelve to fourteen trees of this species, from 30 to 35 inches in diameter, to the acre. It is known to the inhabitants here by the name of cow oak or basket oak, being easily split into narrow, thin strips. The wood is extensively used in the manufacture of baskets used by the negroes in cotton-picking. These baskets are light, and of considerable strength and durability. Next in frequency follows the willow oak, and then the over-cup swamp oak (*Quercus lyrata*), and finally the red oak, found especially on the outskirts of the forest.

"The white ash is not so frequently seen here as elsewhere in similar localities, and does not seem to thrive on these stiff, cold soils. It is in part replaced by the green ash, which here attains the size of a large tree. The black gum is very common, and where the soil is least subjected to overflow the true white oak is found, with fine groups of beech, overtopped by large poplars. Among the smaller trees the mulberry, hornbeam, holly, and abundant papaws must be mentioned.

"The pine hills in the eastern part of Alcorn county are reached at a distance of 6 or 7 miles in a southerly direction from Corinth. Pine occurs on the dividing ridges between the waters of the Tuscumbia river and Yellow creek, or toward the south on those between the Tombigbee and the Tennessee rivers. A short distance west of Glendale station the Cretaceous strata disappear under the ferruginous sands, and mixed with a stunted growth of post oak and Spanish oak, pines appear, forming vast forests on the crests of the hills. This pine (*Pinus mitis*) takes possession of all the old clearings and fields thrown out of cultivation. The rapid growth of the seedlings, which spontaneously spring up thickly after the removal of the broad-leaved trees, leaves no chance for the seedling oaks. It is therefore a certainty that in the future the short-leaved pine will be almost the sole forest tree in this part of the state, outside of the bottom lands, and that it will probably extend its domain far beyond the original limits of its growth.

"The aspect of these pine woods resembles closely that of the lower pine region. The short-leaved pine replaces here the long-leaved pine of the coast, the scrubby post and Spanish oaks take the place of the turkey and the upland willow oaks, while the black-jack is common to both these regions of identical geological formation. The flora of the two regions also presents the same general features; the asters, goldenrods, sunflowers, and various leguminous plants are often the same or belong to closely-allied species. The pine-clad drift hills interspersed between the Carboniferous and Cretaceous regions are parts of the northern interior drift belt which extends throughout Alabama. The region of the short-leaved pine of northeastern Mississippi extends from the southern border of the valley of the Tennessee river to the southern extremity of Itawamba county, and is on an average 10 miles in width, embracing an area of nearly 600 square miles. Of this region, after the deduction of the fertile bottoms of the Tombigbee and Yellow Creek valleys, where no pines are found, two-thirds can be regarded as occupied by the pine forest. As the sole supply of pine lumber in the northern part of the state, this region is of great importance. Several saw-mills, none of which have an annual capacity of more than 3,000,000 feet, are established on the railroad line at Glendale, Burnsville, and near Iuka; portable saw-mills are worked also through this forest in its whole extent, their product being hauled in wagons for miles to the nearest station on the Mobile and Ohio and the Memphis and Charleston railroads. The largest shipments are made from Burnsville and Corinth.

"The second growth of the short-leaved pine, which is already growing with great rapidity in northern Mississippi upon exhausted fields thrown out of cultivation and wherever the forest has been cut from the ridges, should be protected and fostered by the owners of the soil. The care bestowed upon the natural seeding of this useful and valuable timber tree, and in assisting it to gain a permanent foothold on lands regarded as unfit or unprofitable for agriculture, of which tens of thousands of acres are now found in this state, would lead to results of great benefit to the community. The people have it in their power to replenish their timber resources, fast failing through the ever-progressing destruction of the original forest, without other outlay than simply assisting nature in her efforts to recover from injuries sustained in the wholesale destruction of the forest. The restoration of the forest over vast areas, now barren and unproductive wastes, would add vastly to the general welfare and prosperity through the influence such forests would exert upon the climate and salubrity of the country, by the shelter they would offer to insectivorous birds ever busy in the destruction of insects injurious to farm crops, and by the formation of protective screens against the cotton-worm, the most destructive of all insects in this part of the country; for it must be admitted as an undisputed fact that the destruction caused by the cotton-worm is far less upon the small farms where strips of woodland divide the fields than upon the plantations in the rich prairie lands where large areas are destitute of woods. Such forests would serve as windbreaks for crops growing in field and orchard, and as protection against the washing away of the light soil so peculiarly adapted to the cultivation of the great staple of the country, thus preventing the ruin of many productive fields, the *débris* from which, carried away by the rain and floods, fills the rivers and their estuaries, rendering navigation every year more dangerous.

"CENTRAL PINE HILLS.—A hilly region, the northern limit of which is near the center of Benton county, covered with upland oaks and short-leaved pines, extends eastward to the flatwoods in a belt from 8 to 12 miles in width. Farther south, in Calhoun and Sumter counties, this pine region is much wider, embracing the largest part of these and Choctaw and the western part of Oktibbeha counties; from Kosciusko, Attala county, it extends over the whole of Winston and the western part of Noxubee counties, being merged, south of Neshoba in the western part of Kemper county, with the region of mixed tree growth. This pine forest supplies a sufficient amount of lumber for the local demand, and portable saw-mills are found near the large settlements from Kosciusko to the southern limits of the region. It forms a prominent feature in the eastern Gulf states by its geographical position, and must be regarded as one of the distinct divisions which might be designated as the region of the central pine hills. Botanically this region differs from that of the mixed tree growth, upon which it borders toward the south, by the more equal distribution of the pines among the oaks, and particularly by the total absence of the long-leaved pine and other conifers, with the exception of the loblolly pine and of scattered cypress along the river banks, and by the absence of the great magnolia (*M. grandiflora*). The second forest growth in the northern part of this region consists almost exclusively of the short-leaved pine, which southward is associated with the loblolly pine. The short-leaved pine will in the future be the chief forest tree of this region.

"I have personally seen but little of the flatwoods proper, having only touched their southern limits in Kemper county. It is a region of close, cold soil, devoid of drainage, and covered with a stunted growth of post oak; and in its economic aspects as a timber region, or botanically, is of little interest or importance.

"WESTERN MISSISSIPPI.—In Copiah county, below the village of Terry, fifteen saw-mills are in operation along the railroad, obtaining their supply of logs from the heavily-timbered hills in the neighborhood. This lumber is shipped by rail to Saint Louis and Chicago. This business has already reached large proportions and is still increasing rapidly, the mills running without intermission at their full capacity throughout the year.

"Beyond Crystal Springs the country loses its rolling character; the pine hills disappear, and a short distance above the northern boundary of Copiah county, near Terry, a different geological formation is entered, and a strongly-marked change in the vegetation takes place. Horizontal strata of loam, inclosing layers of what appears a whitish sand, stretch northward over a vast extent of level country, and the long-leaved pine disappears with the gravels and sands of the drift.

"North of the pine region a large amount of rich land between the Pearl and Mississippi rivers has been brought under cultivation, especially along the bottoms of the Pearl river and along the principal railway lines. At Jackson, on the Pearl river, little is left of the original tree growth which covered its banks. Still enough is left, however, to show that it was chiefly composed of sweet gums, white oaks, elms, white ashes, etc. The railroad from Jackson to Vicksburg passes through a fertile agricultural country, where only small strips of forest remain between the large plantations and farms. Pines are not seen here, and the black walnut, originally so abundant among the oak and hickory forests which covered this region, must now be regarded as entirely exterminated. Beyond the Blackwater, in the hilly region of the bluff formation, the great magnolia covers the hillsides, although in the vicinity of Vicksburg the hills for miles around the city are entirely stripped of their forests.

"Vicksburg is the center of a considerable lumber industry, depending for its supply of timber upon the cypress rafted down from the mouth of the Yazoo river. The first mill devoted to the manufacture of cypress lumber was established in Vicksburg in 1865. Before that time all the timber from the Yazoo valley was rafted down the Mississippi river, mostly to New Orleans, as is still the case with the greatest number of the rafts. A second mill has lately been built at Vicksburg, and the combined annual capacity of the two is ten or twelve million feet. No

manufactured lumber is shipped from here farther south than Baton Rouge, nearly the whole production being consumed in the erection of small dwellings in the Mississippi and Yazoo bottoms. The logs received at these mills average 25 inches in diameter, with a length of from 30 to 70 feet.

"The hillsides in the neighborhood of Vicksburg, when thrown out of cultivation, are seen covered with a stunted growth of locust, Chickasaw plums, and other shrubs. The original forests of the bluff hills consist of extensive groves of stately magnolias, stretching down the slopes and mixing with large white oaks, Spanish oaks, beeches, and towering poplars, covering the mossy ground of the small valleys with delightful shade. Many of the magnolias are from 18 inches to 2 feet in diameter. The full-grown trees, however, show that they have already passed their prime; the upper limbs have begun to die, the base of their trunks being often rotten and hollow. Small specimens and sapling or seedling trees I could not find. The large trees are cut down to supply the neighboring city with fuel, and it is inevitable that in a comparatively short time these magnolia groves will have disappeared, and that these delightfully-shaded hills must share the desolation which surrounds the town.

"THE YAZOO DELTA.—Indian bayou, one of the small water-courses between Pearl river, Deer creek, and Sunflower river, has a sluggish current even in time of high water. As is the case with all the streams of the Yazoo delta, its banks are elevated often to a height of 10 or 15 feet above the surface of the water, thus affording excellent natural drainage for the adjacent country, which is covered with a yellow-brown loam of unsurpassed fertility. As the land, however, recedes from the banks it gradually sinks down again toward the level of the bed of the stream, and the water-courses, following the general direction of the Mississippi river, inclose corresponding lines of depression nearly level with the beds of the streams. These troughs between the bayous and rivers are one of the characteristic features in the topography of the Yazoo delta. They are of various extent, depth, and shape; flat and wide, they form tracts of dark, wet forest swamp, more or less dry in summer; or, narrower and deeper, they form swamps rarely ever entirely free from water; sometimes they are inundated wooded marshes and cane-brakes, or ponds and lagoons more or less shallow and studded with the mighty trunks of the cypress. When these depressions are of considerable depth, lakes, presenting open sheets of water sometimes miles in extent, are formed, their margins, only, overgrown with the cypress. Upon these features depend the great diversity of the forest growth which yet covers the largest part of the Yazoo valley. Along the elevated ridges fronting the streams the white oak, the willow oak, the shell-bark and mocker-nut hickories, the black walnut in great numbers, the yellow poplar and the sassafras large enough to furnish canoes of great size, the mulberry, the Spanish oak, the sweet and the black gums are the principal forest trees, with an undergrowth in the openings of dogwood, various haws, crab apples, wild grapes, buckthorns, etc. In the forests covering the lower lands, which slope back to the swamps and reservoirs, the cow oak takes the place of the white oak, while the over-cup white oak occurs everywhere in the more or less saturated soil. Here the sweet gum reaches its greatest size, and here grow also in great perfection the bitter-nut, the elms, hornbeams, white ash, box-elder, and red maples of enormous size. The honey locust, water oaks, and red and Spanish oaks are equally common. Here, among the smaller trees, the holly attains its greatest development, with hornbeans and wahoo elms, while papaws, haws, and privets form the mass of the dense undergrowth, which, interspersed with dense cane-brakes, covers the ground under the large trees.

"The region covered by these splendid forests of hard woods possesses a wealth of timber of the most valuable kinds and in surprising variety. They occupy by far the greatest part of Sunflower and the adjoining counties between the Mississippi river and the hills which border upon the Yazoo to the east. Most of the clearings made in this region before the outbreak of the war, by the planters settled lower down, have since been abandoned and are again densely covered with the young growth of the trees of which the forest was originally composed. During the last few years, however, the country has been entered again for cultivation by a class of small farmers, who from being farm hands have now risen to the position of independent landholders. It is astonishing to see the utter disregard of these settlers for the forest wealth of the country, which in a short time could not fail to be of great commercial value. On the shores of Indian bayou may be seen clearings with hundreds of the finest black walnuts among the deadened trees, while many of the noblest specimens of this valuable timber tree are felled for fence rails or trifling purposes. The amount of oak and hickory timber destroyed here annually is amazing. It is generally believed, however, that not one acre in fifty over this whole region of hard-wood forest has yet been stripped of its tree covering. Quite different is the condition of the cypress growth in the great Yazoo valley. This tree, confined to low and more or less inundated bottoms bordering on the Mississippi, the Lower Yazoo, Big Sunflower, and their numerous tributaries, was once found in the greatest abundance in this region, and immense quantities of cypress lumber have been furnished by the lower parts of Issaquena and Washington and the western parts of Warren and Yazoo counties. The most valuable timber has now, however, disappeared from the immediate neighborhood of the low river banks easily accessible at seasons of high water during every winter and spring. Only groves standing remote from the banks of the water-courses, and which are only accessible to the raftsmen during exceptionally high stages of water, now supply this lumber. In the upper portions of the valley, however, in the low depressions described as extending between the elevated banks of the streams, more or less limited areas of undisturbed cypress forest are found. The shallow lagoons, covered with water except during seasons of prolonged drought, and called cypress creeks, present in the spring of the year a strange sight. No object meets the eye between the immense trunks of the mighty trees, as in these cypress groves no other tree nor

shrub can live in the dark, shaded, water-covered soil. These reservoirs of drainage, generally without outlet, are called cypress lakes if the water in any part of them, too deep to allow the growth of trees, confines the cypress to their more shallow borders. Here the cypress arrives at its greatest dimensions and produces timber of the finest quality. These cypress lakes and cypress brakes, remote from streams, at no time of the year connected with them, and always surrounded with a mire of forest swamp impassable to wagons, still retain their best timber. Of late years, since swamp and overflowed lands have become the property of the state, planters have added many of these cypress tracts to their estates by purchase; many others have been acquired by companies formed to construct artificial channels by which the timber may be floated to the nearest streams. The richest and most extensive of these groves of cypress, already more or less in the hands of capitalists, are found along Steele's bayou, between Deer creek and the Sunflower river, in Washington county; between that stream and the lower course of Bogue Phalia, and between the Mississippi river and Black creek above Greenville. There is also a very large body of cypress inclosing the 'California brake', upon the Little Sunflower, in the counties of Bolivar and Coahoma, extending through Tallahatchie county to the Yazoo river.

"The traffic in cypress lumber in the Yazoo region dates from 1830. In 1838 it was commenced upon the Sunflower river and Deer creek, ten years after the first settlements were established upon the banks of these streams; since that time rafts have been sent regularly to New Orleans, and camps of lumbermen have been established in every direction, the forests, particularly those upon the public domains, being regarded as the undisputed property and lawful prey of the log-getter. In consequence the cypress groves have been, if not entirely destroyed, largely culled of their best timber wherever it could be obtained without investment of capital, that is by simply floating the logs to the streams at times of freshet and overflow.

"The cutting of these cypress forests is not wisely regulated under the ownership of the state. These lands have been thrown into the market at 50 cents an acre with the condition of settlement. Beneficial as such a law might prove in the disposal of lands fit for cultivation, it results, in the case of timber-land unfit for the plow, in the reckless destruction of one of the surest sources of public revenue. The state thus sells for 50 cents what on its face is worth to the purchaser hundreds of dollars, and which, when deprived of its value and rendered forever worthless, will be turned back to the state again.

"Much of the destruction of the timber can be traced to wasteful methods practiced by the negroes. Under present methods any one having rented a plantation will, for the most trifling wants, cut down a tree, regardless of size, and without any effort to preserve for future use the parts not immediately wanted, so that the next quarter of a century will probably see the entire destruction of the vast quantities of timber stored in the whole of this great territory."


LOUISIANA.

The coast of Louisiana is bordered by saline marshes and savannas extending inland from 10 to 40 miles, or is covered with a scattered growth of cypress occupying extensive fresh-water swamps peculiar to the region. In Vermillion, Calcasieu, Saint Martin's, and Saint Landry parishes considerable treeless areas, open grassy prairies in the borders of the forest, occur. With these exceptions Louisiana was originally covered with a dense and varied forest growth. The Maritime Pine Belt covered the eastern portion of the state nearly to the Amite river, or until checked from further western development by the alluvial deposits of the Mississippi. Forests of pine, too, occupied the western part of the state north and south of the Red river. The pine flats of Calcasieu were covered with forests formed almost exclusively of the long-leaved pine, which, farther north, mixed with oaks and various hard-wood trees, extends over the high rolling country which stretches from the Sabine northeasterly nearly to the Ouachita river. The northeastern part of the state was covered, outside of the broad bottom lands of the rivers, with a heavy forest of short-leaved pine (*Pinus mitis*) mixed with upland oaks, hickories, and other deciduous trees. The bottom lands and all that part of the state bordering the Mississippi were covered with a heavy growth of the trees peculiar to such low, rich soil throughout the Gulf region. The high bluffs which occur at different points along the Mississippi, the Atchafalaya, and other streams flowing through the western part of the state were covered with a noble forest of evergreen magnolias mingled with beeches, water oaks, and gums.

The most valuable forests of the state are still almost intact, although the pine has been cut from the banks of the Pearl river and some of its tributaries, and from along the line of the Chicago, Saint Louis, and New Orleans railroad, to furnish the New Orleans market with lumber. Pine has also been cut along the Sabine river, from both forks of the Calcasieu, along the Red river in the neighborhood of Alexandria and Shreveport, and more recently in Catahoula parish, along Little river. The river swamps and rolling hills in the eastern and northern parts of the state still contain vast bodies of valuable hard-wood forest yet untouched by the ax.

The forests of Louisiana, uninvaded as yet by the manufacturers of naval stores, have not greatly suffered from forest fires. During the census year only 64,410 acres of woodland were reported as burned over by fire, with a loss of only \$6,800. These fires were generally set to improve pasturage, or by careless hunters camping in the forest.

A small amount of cooperage stock is made in New Orleans almost entirely from cypress and pine, although that city has long been an important point of export for oak staves and headings brought there from Arkansas and



Tennessee by river. The magnificent hard woods common over much of the state can supply abundant material for many important industries which already at the north suffer from the exhaustion and deterioration of the local timber supply.

The following rough estimates of the amount of the long-leaved and short-leaved pine standing in the state have been prepared by measuring upon a large-scale map areas occupied by the pine forests, which coincide almost exactly with geological formations. From these areas the totals of clearings as returned by enumerators and all areas of swamp, bottom lands, and prairies are deducted to obtain the extent of territory covered with pine forests. By multiplying this area by the average stand of timber per acre, obtained by numerous observations in different parts of the state, the following estimate of the amount of merchantable pine standing May 31, 1880, is reached:

Parishes.	Long-leaved pine (<i>Pinus palustris</i>).	Short-leaved pine (<i>Pinus mitis</i>).
	<i>Feet, board measure.</i>	<i>Feet, board measure.</i>
Bienville.....	416, 000, 000	1, 837, 000, 000
Bossier		1, 574, 000, 000
Cadde		1, 696, 000, 000
Calcasieu	4, 219, 000, 000	
Caldwell	602, 000, 000	382, 000, 000
Catahoula	1, 558, 000, 000	304, 000, 000
Clalborne		1, 923, 000, 000
De Soto.....		1, 971, 000, 000
East Baton Rouge.....		157, 000, 000
East Feliciana	108, 000, 000	886, 000, 000
Grant.....	1, 574, 000, 000	
Jackson	483, 000, 000	1, 670, 000, 000
Livingston	300, 000, 000	
Morehouse		797, 000, 000
Natchitoches	1, 792, 000, 000	618, 000, 000
Onachita	16, 000, 000	1, 126, 000, 000
Rapides.....	2, 422, 000, 000	
Red River		643, 000, 000
Sabine	598, 000, 000	1, 974, 000, 000
Saint Helena	749, 000, 000	
Saint Landry	579, 000, 000	
Saint Tammany	1, 398, 000, 000	
Tangipahoa.....	1, 537, 000, 000	
Union.....		2, 522, 000, 000
Vernon	3, 741, 000, 000	
Washington	1, 734, 000, 000	
Webster.....		1, 443, 000, 000
West Feliciana.....		122, 000, 000
Winn.....	2, 062, 000, 000	
Total	26, 588, 000, 000	21, 625, 000, 000
Cut for the census year ending May 31, 1880 ...	61, 882, 000	22, 709, 000

The principal point of lumber manufacture is Saint Charles, in Calcasieu parish, on the southern border of the western pine forest. Lumber manufactured here is shipped east and west by rail, and in small schooners to Mexican and West Indian ports. A comparatively small amount of lumber is manufactured at New Orleans from logs cut in eastern Louisiana and towed through lake Pontchartrain and the canals to the city, and along the river front from logs rafted out of the Red, Little, Black, and other streams of northern Louisiana. New Orleans, however, is principally supplied with lumber sawed at Gulf ports, in spite of its position with reference to the most valuable hard-pine forests upon the continent, its large local demand for lumber and all saw-mill refuse, and its facilities for export, which would seem to indicate that it must become the most important center of lumber manufacture and distribution in the south. Small quantities of pine lumber have long been manufactured upon the Red river near Alexandria; short-leaved pine (*Pinus mitis*) is sawed at Shreveport, and in small quantities for local consumption at other points in the northern parishes.

MOSS GINNING.

New Orleans is the center of the "moss-ginning" industry of the United States. The "moss" (*Tillandsia usneoides*), a common epiphyte, growing in great quantities upon the cypress, live oak, and other southern trees, is gathered, by men known as "swampers", in the swamps of Louisiana, Mississippi, Alabama, and Florida. The moss when gathered is piled near the swamps and allowed to rot during ten or twelve months. It loses in this process about 90 per cent. of its weight, and is then shipped to New Orleans, where it is cleaned, dried, and ginned, losing in this latter operation 35 per cent. in weight. The prepared moss is used in upholstery, either alone or

mixed with hair. The product of the New Orleans factories is principally shipped to the western states, a comparatively small amount being sent to Europe. Six moss factories are located in New Orleans, and there are also small establishments at Plaquemine and at Morgan City, Louisiana, and at Pensacola, Florida. New Orleans received during the year ending August 31, 1881, 3,500 bales of rough moss, weighing 10,000,000 pounds, and valued at \$315,000. A considerable amount, however, is ginned in the country and shipped direct to consumers, or is prepared by the consumers themselves. Persons most familiar with the volume of this industry estimate that the value of the prepared moss gathered annually in Louisiana, the principal region of supply, is not far from \$550,000. The amount gathered, however, varies considerably from year to year. Moss can only be profitably collected at times of high floods, when the swamps are navigable to small boats, and the moss, hanging from the branches of the trees, can be easily gathered. The wages earned by the swamper, too, are not large, and the gathering of moss is only resorted to when more profitable employment upon farms cannot be obtained.

The following extracts are from notes of a hasty journey made through the forest region of western Louisiana by Dr. Charles Mohr:

"For the investigation of the important pine region of western Louisiana I selected Alexandria as my starting point. Situated almost centrally between the forests of long-leaved pine which skirt both sides of the Red River valley, Alexandria is the seat of the actual lumber trade and the point where the lumber interests of this great timber region must be developed in the future. Little is left of the vast cypress swamps which once covered the alluvial lands on the Mississippi river below the mouth of the Red river and the lower basin of that stream. It is only in the most inaccessible swamps, cut off from all communication with the rivers, that patches of this timber remain. The ever-increasing demand for this lumber has almost exhausted the available cypress of the Red River country, and cypress is now drawn from the forest farther north bordering the Black and Ouachita rivers. The lowlands along the river front, subject to inundation and devoid of drainage, present in their tree growth the same features as the low forests of the Mississippi and the Yazoo valleys. The bitter pecan flourishes here luxuriantly, and with it the white ash, the swamp over-cup oak, the persimmon, sycamore, sassafras, sweet gum, and cottonwood. The green ash is common, and in better-drained localities the willow, white, cow, and red oaks appear, with elms and occasional pecans. Twelve or 15 miles below Alexandria the first pines are seen looming up in the forest; upon a nearer approach they are recognized as the loblolly. A short distance farther up the river, upon sandy bluffs fronting the western shore, fine specimens of the short-leaved pine are observed, associated with black oaks, Spanish oak, the black-jack, and many of the shrubs peculiar to the drift of the coast pine region east of the Mississippi. The wide bottom lands of the river upon which Alexandria is situated extend west to bayou Boeuf. This district, unsurpassed in fertility and regarded as the garden of Louisiana, has but little left of the forest with which it was once covered. The pecan trees alone of the original forest growth have been spared from the general destruction. Of these, fine specimens line the roadsides and dot the fields. The unsightly honey locust occupies the waste low places, in company with a second growth of willows, hackberries, and catalpas. The shores of bayou Boeuf are covered with a variety of trees. Cypress line the brink of the water; beyond these, sycamores, bitter gums, sweet and white gums, pecans, water and willow oaks, red and white elms, red maple, and ash occupy the gentle acclivities, with a dense undergrowth of smaller trees—the dogwood, several haws, wahoos, catalpas, Carolina buckthorn, southern prickly ash, etc. Ascending the ridge to the uplands the deep alluvial soil is left behind, and the light sandy loams of the Tertiary strata make their appearance, and with this change of soil the vegetation changes as suddenly. Stately loblolly pines rise above the groves of post, black, and Spanish oaks, and where the ridge descends again to what might be called the second bottom of bayou Boeuf, a forest of white oak is entered, which contains a stand of timber seldom equaled. On the long, gentle swells these are associated with fine Spanish oaks, a few pig-nuts and mocker-nuts, and in the depressions with red oak, elms, ash, and other trees found on soil of good quality in the same latitude east of the Mississippi river.

"The hills formed by the sandstone drift gravels rise suddenly from the plain covered with the forest of the long-leaved pine, comparing favorably both in the size and number of the trees with the best timber districts in the Coast Pine Belt of the eastern Gulf states. Trees under 12 inches in diameter are rarely seen, as is the case everywhere in these undisturbed primeval pine forests. The soil of this region is closer, more retentive of moisture, and richer in plant-food than that in the Maritime Pine Region east of the Mississippi. The pines here are therefore of more rapid growth and below the standard of quality for which the pine produced on the poor, siliceous ridges of lower Mississippi and Alabama is so highly valued. The numerous streams which cut their way through these pine hills are fringed with many of the evergreens peculiar to the eastern Gulf coast; and magnolias, the red and white bay, wax myrtles, willows, and the devilwood are common.

"The pine region west of the Red River valley spreads westward to the Sabine, forming part of the great pine forest which extends far into eastern Texas. Southward it constantly increases in width; and its length from north to south, where it verges upon the lower maritime prairies of the Calcasieu, is not less than 100 miles. It includes the whole of the parish of Vernon, the largest part of Calcasieu, and portions of the parishes of Natchitoches and Rapides, covering an area of about 4,500 square miles. The northern portion of this belt is one vast primeval forest. The small inroads made by the scattered settlers and the few small saw-mills which supply a small local

demand are too insignificant to be taken into account. In the southern portion of this forest the saw-mills on the Sabine river and at Lake Charles have already removed some timber from the banks of the principal streams.

"The region of long-leaved pine which skirts the eastern confines of the Red River valley, and which at its southern extremity almost touches the river banks, may be called the central pine region of west Louisiana. The village of Pineville, opposite the city of Alexandria, is the center of the lumber trade of this region. The high, undulating uplands formed of the Pliocene-Tertiary strata which here front the river bear a growth of loblolly and short-leaved pine mixed with upland oaks. A few miles to the eastward, however, upon the hills covered with drift, the forest of long-leaved pine appears. The surface in this central pine region is more broken, the soil poorer, more porous and siliceous than west of the Red River valley, and the timber produced here is of unsurpassed quality. An average of not less than fifteen trees to the acre, with a diameter of over 15 inches 3 feet from the ground, grow here. The production of lumber is limited to saw-mills situated 7 or 8 miles from the river. They have been gradually removed from its banks as the timber was exhausted on a line 7 or 8 miles in length north and south from Pineville. The production of these mills amounts in the aggregate to 40,000 feet a day. The lumber manufactured here supplies the population of the Red River valley as far west as Shreveport.

"The rolling uplands which extend to the edge of the river at Shreveport are covered with a heavy, cold, clayey soil almost impervious to water; they bear an open growth of oaks, among which the post oak is the prevailing species, finding here the conditions most favorable to its growth. The Spanish oak, invariably called west of the Mississippi river red oak, with fine black-jack makes up the larger part of the tree growth. Hickories, represented by the pig-nut and mocker-nut, are not frequent, and are of small size. The black oak is found in localities with somewhat rocky surface and loose subsoil, while white oaks occur along the base of declivities where an accumulation of vegetable matter has been deposited. The undergrowth in these woods is scanty, and consists for the most part of seedling oaks. Where, however, the forest has been entirely removed, the loblolly pine takes exclusive possession of the soil. These oak forests reach to the northern confines of the state and extend west into Texas. In their southern extremity toward the pine region the soil is better, and the white oak becomes the prevailing forest tree. My attention was directed to the fact that since the removal of the raft of the Red river the drainage of the upper part of the valley has been greatly improved, and many of the lakes and swamps formerly continually inundated are now dry, while the swamp forest growth, including the cypress, is dying, or has already died.

"Opposite Shreveport the valley spreads out into an extensive plain from 8 to 10 miles in width, descending imperceptibly as it recedes from the bank of the river. These lowlands are mere swamps, often deeply overflowed by the backwater of the river, which finds its way through the numerous bayous and inlets which intersect this plain. The forest growth covering these swamps is of inferior size, and consists of but few species. The cypress occupies the overflowed swamps, but it is always below medium size, and I did not notice a single specimen 2 feet in diameter. The saline, gypsum soil does not seem suited to its full development. The water locust finds here its favorite home. It is very common in moist localities not subject to constant inundation. The wood of this tree is as hard and durable as that of the common honey locust, and is employed for the same purposes; that is, in the manufacture of stirrups, blocks, hubs, etc. The green ash is frequently seen here growing with the wahoo, hornbeam, holly, and privet, and forming broad clumps of great luxuriance beneath the larger trees. After passing Cross bayou the land gently rises, and, with better drainage, the trees of the swamps disappear and are replaced by a more varied and valuable timber growth. The white ash and white and red oaks are the more common trees in the woods which skirt the base of the ridges forming the eastern limits of the valley of the Red river. At this point they are separated from the low hills of the Pliocene sandy loams by a pretty, clear stream, the Red Chute, which runs swiftly over its bed along the base of the uplands; these form long, gentle, swelling slopes, or spread out into broad flats more or less deficient of drainage. The ridges are all wooded with upland oaks and short-leaved pines, while the loblolly pine, with water and willow oaks, sweet and black gums, cover the depressions and damp flats. The tree growth upon these ridges is vigorous. I have nowhere found the short-leaved pine of finer proportions, equaling in size and length of clear trunk the long-leaved species. This region of the short-leaved pine, with its low, heavily-timbered ridges, is similar in character of soil and vegetation to the pine hills of central and northern Mississippi, and might be designated as the region of the pine hills of northern Louisiana. Between lake Bodcan and lake Bisineau the surface of the country is very often imperfectly drained, and there the loblolly pine is the prevailing tree. A few miles back of Bellevue, in Bossier parish, the level forest is interrupted by a strip of prairie from 1 mile to 3 miles wide, covered with a cold, soapy, gray soil impervious to water. On these natural meadows no tree or shrub is growing, except a peculiar *Cratægus*, new to me. (a) It is a small tree or large shrub, forming strictly-defined, impenetrable, dense thickets a few rods or of several acres in extent. In its arborescent form it rises to a height of from 15 to 20 feet, with a more or less bent trunk 6 or 7 inches in diameter, spreading its crooked limbs at a height of from 4 to 6 feet above the ground. The fruit is said to be as large as that of the apple haw, sweet and edible; it is eagerly eaten by swine, which fatten upon it. This tree is here called by the people 'hogs' law'.

a *Cratægus brachyacantha*, Sargent and Engelmann.

"On the decline which leads to the valley of bayou Dauchitta, the flatwoods give way to a fine growth of Spanish and post oaks, elms, and gums.

"The western bank of the bayou is confronted by hills of the post-Tertiary sands and gravels which westward form a succession of steep ridges heavily wooded with the upland oaks and short-leaved pine. The narrow creek bottoms inclosed between these ridges are watered abundantly by springs and clear streams shaded by white and red bay, hollies, azaleas, and kalmias. The great magnolia is not seen here, and the American olive is missing. In these gravelly hills, extending westward to the valley of the Ouachita river, the short-leaved pine is very common and the characteristics of the pine-hill region are prominent. These hills cover a large area extending northward into Arkansas, and toward the south merging gradually into the oak woods which border upon the bottoms of the numerous tributaries of the Red river. This pine-hill region is sparsely settled, and, remote from water and rail communication, its original stores of pine and hard-wood timber have scarcely been touched.

"An intimate knowledge of the forest growth in this section was obtained by an excursion over the hills to bayou Dauchitta above its entrance to lake Bistineau. In the localities of the best drainage in this valley the cow oak is very common, mixed with the white and post oaks, while sweet gums, black gums, water and willow oaks, and hackberries occupy lower situations. On the immediate banks and in the sloughs small cypress trees are common, mixed with the bitter pecan, the hornbeam, the water locust, and the sycamore. The loblolly pine takes possession of every opening in the forest, descending the high hills, while numerous haws border the edges of the forest. In the bottoms and along the declivities, the Chickasaw and the American plum are found of larger size than farther east. Loblollies and hickories with the black and post oaks occupy the lower declivities, and upon the heights the yellow pine mixed with upland oaks forms fine forests."

TEXAS.

The most important forests of Texas are found in the extreme eastern part of the state, where the Maritime Pine Belt of the south Atlantic region extends to about midway between the Trinity and the Brazos rivers. A forest of long-leaved pine occupies most of the territory between the Sabine and the Brazos south of the thirty-first degree of north latitude, reaching south to within 20 miles of the coast. Beyond the long-leaved pine forests, forests of the loblolly pine, mixed with hard woods, stretch westward 50 or 60 miles, while north of these two regions a third division of the pine belt, composed of a heavy growth of short-leaved pine mingled with upland oaks, occupies the rolling ridges which extend northward to beyond the Red river. The swamps which line the larger streams flowing into the Gulf, especially within the limits of the pine belt, still contain large bodies of cypress. The quality of the Texas cypress, however, is inferior to that grown east of the Mississippi river, and probably one-third of the timber growing in the valleys of the Sabine and the Nueces rivers is "peggy" or affected by dry rot.

West of the pine belt open forests largely composed of post and black-jack oaks occur, gradually decreasing in density, and finally, west of the ninety-seventh degree of longitude, entirely disappearing. Farther west, however, the "lower" and "upper cross-timbers", two remarkable bodies of timber, composed of small and stunted specimens of these oaks, extend from the Indian territory far south into the prairie region, occupying long, narrow, irregular belts where sandy or gravelly alluvial deposits overlie the limestone of the prairie region. A belt of forest, largely composed of post and black-jack oaks, varying from 20 to 50 miles in width extends southwest of the Trinity nearly to the Nueces river, its eastern border following generally, at a distance of from 50 to 60 miles inland, the trend of the coast. The bottom lands east of the one hundredth meridian are lined with the deciduous trees which occupy similar situations in the eastern Gulf states. Near the coast the bottom lands of the large rivers, often several miles in width, are covered with dense forests composed of enormous trees. Farther west the bottoms gradually narrow, the number of arborescent species covering them decreases, and individual trees are small and stunted.

West of the Colorado river the forests of the Atlantic region are replaced outside of the bottom lands by Mexican forms of vegetation; the hills are covered with a stunted growth of mesquit, Mexican persimmon, various acacias, and other small trees of little value except for fuel and fencing.

An important tree in the forest of western Texas is the cedar covering the low limestone hills which occupy hundreds of square miles north and west of the Colorado river, in Travis, Bastrop, Hays, Comal, and adjacent counties. West of the one hundredth meridian all forest growth disappears, with the exception of a few scattered cottonwoods, elms, and hackberries, confined to the narrow bottoms, and a shrubby growth of mesquit, which covers the plains of western Texas, furnishing the only fuel of the region. The mountain ranges, outlying ridges of the Rocky mountains, which occupy the extreme western part of the state, are covered with an open, stunted forest of western pines and cedars, with which mingle the post oak, the yellow oak, and other species of the Atlantic region.

The pine belt covering the eastern counties of the state is alone important as a source of lumber supply. Areas of river-bottom land covered with trees are, as compared with the area of the state, insignificant in extent, and these river belts of forest are entirely insufficient to supply even the mere local wants of the nearest settlements. The oak forests, which stretch more or less continuously between the eastern pine belt and the treeless western prairies and plains, are, except along their extreme eastern borders, composed of small, stunted trees, often hollow, defective, and of little value except for fuel, fence rails, and railway ties. The forests of the western mountains are

DENSITY OF FORESTS
COMPILED UNDER THE DIRECTION OF
C. S. SARGENT, SPECIAL AGENT
1883.

Scale
0 10 20 30 40 50 60 Miles

LEGEND.

I	Under 1 cord per acre
II	From 1 to 2
III	" 2 to 3
IV	" 3 to 5
V	" 5 to 10
VI	" 10 to 20
VII	" 20 to 30
VIII	" 30 to 40
IX	" 40 to 50
X	" 50 to 60
XI	" 60 to 70
XII	" 70 to 80
XIII	" 80 to 90
XIV	" 90 to 100
XV	" 100 to 110
XVI	" 110 to 120
XVII	" 120 to 130
XVIII	" 130 to 140
XIX	" 140 to 150
XX	" 150 to 160
XXI	" 160 to 170
XXII	" 170 to 180
XXIII	" 180 to 190
XXIV	" 190 to 200
XXV	" 200 to 210
XXVI	" 210 to 220
XXVII	" 220 to 230
XXVIII	" 230 to 240
XXIX	" 240 to 250
XXX	" 250 to 260
XXXI	" 260 to 270
XXXII	" 270 to 280
XXXIII	" 280 to 290
XXXIV	" 290 to 300
XXXV	" 300 to 310
XXXVI	" 310 to 320
XXXVII	" 320 to 330
XXXVIII	" 330 to 340
XXXIX	" 340 to 350
XL	" 350 to 360
XLI	" 360 to 370
XLII	" 370 to 380
XLIII	" 380 to 390
XLIV	" 390 to 400
XLV	" 400 to 410
XLVI	" 410 to 420
XLVII	" 420 to 430
XLVIII	" 430 to 440
XLIX	" 440 to 450
L	" 450 to 460
LI	" 460 to 470
LII	" 470 to 480
LIII	" 480 to 490
LIV	" 490 to 500
LVI	" 500 to 510
LVII	" 510 to 520
LXIII	" 520 to 530
LXIV	" 530 to 540
LXV	" 540 to 550
LXVI	" 550 to 560
LXVII	" 560 to 570
LXVIII	" 570 to 580
LXIX	" 580 to 590
LXX	" 590 to 600
LXXI	" 600 to 610
LXXII	" 610 to 620
LXXIII	" 620 to 630
LXXIV	" 630 to 640
LXXV	" 640 to 650
LXXVI	" 650 to 660
LXXVII	" 660 to 670
LXXVIII	" 670 to 680
LXXIX	" 680 to 690
LXXX	" 690 to 700
LXXXI	" 700 to 710
LXXXII	" 710 to 720
LXXXIII	" 720 to 730
LXXXIV	" 730 to 740
LXXXV	" 740 to 750
LXXXVI	" 750 to 760
LXXXVII	" 760 to 770
LXXXVIII	" 770 to 780
LXXXIX	" 780 to 790
LXXXX	" 790 to 800
LXXXXI	" 800 to 810
LXXXXII	" 810 to 820
LXXXXIII	" 820 to 830
LXXXXIV	" 830 to 840
LXXXXV	" 840 to 850
LXXXXVI	" 850 to 860
LXXXXVII	" 860 to 870
LXXXXVIII	" 870 to 880
LXXXXIX	" 880 to 890
LXXXXX	" 890 to 900
LXXXXXI	" 900 to 910
LXXXXXII	" 910 to 920
LXXXXXIII	" 920 to 930
LXXXXXIV	" 930 to 940
LXXXXXV	" 940 to 950
LXXXXXVI	" 950 to 960
LXXXXXVII	" 960 to 970
LXXXXXVIII	" 970 to 980
LXXXXXIX	" 980 to 990
LXXXXXX	" 990 to 1000

DENSITY OF FORESTS
COMPILED UNDER THE DIRECTION OF
C. S. SARGENT, SPECIAL AGENT
1883.

Scale
0 10 20 30 Miles

LEGEND.

I	Under 1 cord per acre
II	From 1 to 2 "
III	" 2.5 "
IV	" 5.10 "
V	" 10.20 "
VI	" 20.30 "
Over	30 "

DENSITY OF FORESTS
COMPILED UNDER THE DIRECTION OF
C. S. SARGENT, SPECIAL AGENT
1883.

Scale
0 10 20 30 Miles

LEGEND.

I	Under 1 cord per acre
II	From 1 to 2 "
III	" 2.5 "
IV	" 5.10 "
V	" 10.20 "
VI	" 20.30 "
Over	30 "

DENSITY OF FORESTS
COMPILED UNDER THE DIRECTION OF
C. S. SARGENT, SPECIAL AGENT
1883.

Scale
0 10 20 30 40 50 60 Miles

LEGEND.

I	Under 1 cord per acre
II	From 1 to 2
III	" 2.5 "
IV	" 5, 10 "
V	" 20, 30 "
VI	" 40, 50 "
Over 60	"

DENSITY OF FORESTS
COMPILED UNDER THE DIRECTION OF
C. S. SARGENT, SPECIAL AGENT
1883.

Scale
0 10 20 30 40 50 60 Miles

LEGEND.

I	Under 1 cord per acre
II	From 1 to 2
III	" 2 to 3
IV	" 3 to 5
V	" 5 to 10
VI	" 10 to 20
VII	" 20 to 30
VIII	" 30 to 40
IX	" 40 to 50
X	" 50 to 60
XI	" 60 to 70
XII	" 70 to 80
XIII	" 80 to 90
XIV	" 90 to 100
XV	" 100 to 110
XVI	" 110 to 120
XVII	" 120 to 130
XVIII	" 130 to 140
XIX	" 140 to 150
XX	" 150 to 160
XXI	" 160 to 170
XXII	" 170 to 180
XXIII	" 180 to 190
XXIV	" 190 to 200
XXV	" 200 to 210
XXVI	" 210 to 220
XXVII	" 220 to 230
XXVIII	" 230 to 240
XXIX	" 240 to 250
XXX	" 250 to 260
XXXI	" 260 to 270
XXXII	" 270 to 280
XXXIII	" 280 to 290
XXXIV	" 290 to 300
XXXV	" 300 to 310
XXXVI	" 310 to 320
XXXVII	" 320 to 330
XXXVIII	" 330 to 340
XXXIX	" 340 to 350
XL	" 350 to 360
XLI	" 360 to 370
XLII	" 370 to 380
XLIII	" 380 to 390
XLIV	" 390 to 400
XLV	" 400 to 410
XLVI	" 410 to 420
XLVII	" 420 to 430
XLVIII	" 430 to 440
XLIX	" 440 to 450
L	" 450 to 460
LI	" 460 to 470
LII	" 470 to 480
LIII	" 480 to 490
LIV	" 490 to 500
LVI	" 500 to 510
LVII	" 510 to 520
LXIII	" 520 to 530
LXIV	" 530 to 540
LXV	" 540 to 550
LXVI	" 550 to 560
LXVII	" 560 to 570
LXVIII	" 570 to 580
LXIX	" 580 to 590
LXX	" 590 to 600
LXXI	" 600 to 610
LXXII	" 610 to 620
LXXIII	" 620 to 630
LXXIV	" 630 to 640
LXXV	" 640 to 650
LXXVI	" 650 to 660
LXXVII	" 660 to 670
LXXVIII	" 670 to 680
LXXIX	" 680 to 690
LXXX	" 690 to 700
LXXXI	" 700 to 710
LXXXII	" 710 to 720
LXXXIII	" 720 to 730
LXXXIV	" 730 to 740
LXXXV	" 740 to 750
LXXXVI	" 750 to 760
LXXXVII	" 760 to 770
LXXXVIII	" 770 to 780
LXXXIX	" 780 to 790
LXXXX	" 790 to 800
LXXXXI	" 800 to 810
LXXXXII	" 810 to 820
LXXXXIII	" 820 to 830
LXXXXIV	" 830 to 840
LXXXXV	" 840 to 850
LXXXXVI	" 850 to 860
LXXXXVII	" 860 to 870
LXXXXVIII	" 870 to 880
LXXXXIX	" 880 to 890
LXXXXX	" 890 to 900
LXXXXXI	" 900 to 910
LXXXXXII	" 910 to 920
LXXXXXIII	" 920 to 930
LXXXXXIV	" 930 to 940
LXXXXXV	" 940 to 950
LXXXXXVI	" 950 to 960
LXXXXXVII	" 960 to 970
LXXXXXVIII	" 970 to 980
LXXXXXIX	" 980 to 990
LXXXXXX	" 990 to 1000

not luxuriant, and at the best can only supply a limited local demand with inferior lumber. It is probably no exaggeration to say that west of the pine belt, and with the exception of the small amount of hard wood found on the bottom lands near the coast, the forests of Texas do not contain a single tree fit to manufacture into first-class lumber. The pine forests, therefore, of eastern Texas and western Louisiana are important factors in the future development of Texas, as well as of the treeless northeastern provinces of Mexico, which must draw their building material from these pineries. The position of these forests, therefore, with reference to an enormous territory destitute of timber, although adapted to agriculture and grazing, and which must soon be covered with a considerable population and a net-work of railroads, their richness of composition, and the facility with which they can be worked, give to them perhaps a greater prospective value than that possessed by any body of timber of similar extent in the United States.

During the census year 599,359 acres of woodland were reported damaged by fire, with an estimated loss of \$273,990. Of these fires the larger number was set to improve pasturage, in clearing land, or through malice. These returns do not include the large areas burned in western Texas by prairie fires, checking the growth of the mesquit over a great extent of territory.

Small amounts of cooperage stock and woodenware, principally for local consumption, are manufactured in the eastern counties from oak and cypress. Manufacturers report an abundant supply of material.

The following rough estimates of the amounts of the three kinds of pine standing in the state May 31, 1880, were made by multiplying the average stand of timber per acre by the county areas occupied by the pine forests, these being obtained by deducting, from total areas of the county, estimated areas covered by clearings, bottom lands, swamps, etc.:

Counties.	Long-leaved pine (<i>Pinus palustris</i>).	Short-leaved pine (<i>Pinus mitis</i>).	Loblolly pine (<i>Pinus Taeda</i>).
	<i>Feet, board measure.</i>	<i>Feet, board measure.</i>	<i>Feet, board measure.</i>
Anderson		336,000,000	1,763,600,000
Angelina	1,340,500,000		1,190,400,000
Bowie		2,380,800,000	
Camp		579,200,000	
Cass		2,470,400,000	
Cherokee		2,230,400,000	585,600,000
Franklin		448,000,000	
Gregg		598,400,000	
Grimes			211,200,000
Hardin	1,244,800,000		627,200,000
Harris			1,827,200,000
Harrison		2,328,400,000	
Henderson		521,600,000	
Hopkins		483,200,000	
Houston			3,216,000,000
Jasper	2,534,400,000		
Jefferson			288,000,000
Liberty	41,600,000		2,147,200,000
Madison			283,600,000
Marion		1,187,200,000	
Montgomery			2,326,400,000
Morris		729,600,000	
Nacogdoches	1,216,000,000	1,555,200,000	35,500,000
Newton	2,112,000,000		33,000,000
Orange	280,000,000		518,400,000
Panola	1,193,600,000	1,107,200,000	
Polk	2,720,000,000		473,600,000
Red River		272,000,000	
Rusk	115,200,000	2,492,800,000	
Sabine	1,648,000,000		
San Augustine	1,625,600,000		
San Jacinto			1,833,600,000
Shelby	1,884,800,000	425,600,000	
Smith		2,035,200,000	
Titus		806,000,000	
Trinity	51,000,000		1,987,200,000
Tyler	2,550,400,000		
Upshur		1,392,000,000	
Van Zandt		26,000,000	
Walker			1,590,400,000
Waller			19,000,000
Wood		1,600,000,000	
Total	20,508,200,000	28,093,200,000	20,907,100,000
Amount cut for the year ending May 31, 1880	66,450,000	a 146,420,000	61,570,000

a Including 30,290,000 shingles.

The principal centers of lumber manufacture in Texas are Orange and Beaumont, on the Sabine and Nueces rivers, above Sabine pass. Long-leaved pine and cypress are sawed here and shipped east and west by rail, and in small quantities by schooner to Texan and Mexican ports. Loblolly pine is sawed at a number of small mills upon the line of the International and Great Northern railroad in the counties south of the Trinity river, and a large amount of short-leaved pine is manufactured in the mills upon the line of the Texas Pacific railroad in the northeastern counties, Longview, in Gregg county, being the principal center of this industry. The product of these mills is shipped west by rail to supply settlers upon the prairies of northern Texas with building material.


The following extracts are derived from the notes upon the forests of Texas made by Dr. Charles Mohr, of Mobile:


"West of Marshall, upon the Texas Pacific railroad, the surface of the land becomes more broken; the soil is lighter, more porous, and favorable to the growth of the short-leaved pine, which soon becomes the prevailing forest tree in the woods extending toward the west. Longview, a small town at the junction of the International and Great Northern and Texas Pacific railroads, is situated almost in the center of the short-leaved pine region, and is the seat of an active lumber business. These forests of short-leaved pine, more or less interspersed with oaks, extend to the northern boundary of the state, and southward with an easterly trend to the confines of the region of the long-leaved pine. The short-leaved pine finds its western limits near Mineola.

At Palestine, in Anderson county, the uplands are covered with a loamy, somewhat sandy, soil underlaid with a heavy clay. Here a more or less open oak forest is common. The black oak abounds, with the Spanish, black-jack, blue-jack, and post oak, the last, however, always the prevailing species. Next to the post oak the black-jack is the species of widest distribution in Texas, the two species being always found associated together from the northern confines of the state to the prairies of the coast, and from the east to the treeless regions of western Texas. The bois d'arc (*Maclura aurantiaca*) is common along the banks of the water-courses in eastern Texas, attaining a size large enough to be economically valuable. It is here, however, most probably adventitious from the region in the northwest, where it forms an almost uninterrupted belt of woods from 4 to 10 miles wide, extending from a short distance south of the city of Dallas to the northern frontier of the state, entering the Indian territory between Sherman and Paris. This tree attains a height of from 45 to 50 feet, with a diameter of from 1 foot to 2 feet, and is of great value.

"The timber growth immediately west of the Brazos is stunted and scanty; large areas of grass land intervene between the scrubby woods until all at once ligneous growth disappears, and the seemingly boundless prairie, in gently undulating swells, expands before the view on all sides. Near the center of Milam county a belt of open post-oak woods from 20 to 25 miles in width is entered. It extends from Belton, in Bell county, southward to the upper confines of Gonzales county. Post oaks stand here from 20 to 30 feet apart, with black-jacks and blue-jacks between them, the trees being all of small size. The soil of these oak hills is of poor quality, sandy, gravelly, and more or less broken, arid, and devoid of vegetable mold. Toward the southern limit of this belt, near Bastrop, a tract of loblolly pine is found covering nearly four townships, or about 90,000 acres. During the last twelve years all the useful timber on this isolated tract has been cut down. A second growth of pine, however, has sprung up, and is now growing vigorously under the fostering care of the owners of the land, and promises in a short time to afford a new supply of timber. A belt of post oak is found intersecting the prairie from the upper part of McLennan county, near Waco, and extending to the northern frontier of the state, where it joins the cross-timbers of the Wichita. It is known as the 'lower cross-timbers'. This belt of oak wood is nearly 150 miles long, with its greatest width of about 20 miles between Dallas and Fort Worth. At a distance of from 20 to 40 miles west of the lower cross-timbers another belt of oak extends from Comanche county to the northern boundary of the state, with a long western spur following the valley of the Brazos as far as the ninety-ninth meridian. This oak forest is known as 'the cross-timbers'.

"Taken as a whole, the country west of the Brazos river, except the basin of the Colorado, is a poorly-timbered region. The mesquit was first met with on the declivities of the prairie, which verge here upon the valley of the Colorado. The wood of this tree is hard, fine-grained, tough, heavy, and of great durability. In the western portions of the state, almost entirely destitute of other timber growth, it serves, according to its size, a variety of purposes in the economy of the stock ranch, and is there invaluable for fencing. Burning with a clear, smokeless flame and possessing great heating powers, it is unsurpassed as fuel by any other Texas wood. It serves, moreover, another important purpose in furnishing an abundance of wholesome and nutritious food to large herds of cattle, at a season of the year when long-continued droughts have destroyed the grass upon the prairie. With the increasing settlement of the treeless-prairie region during the last 15 or 20 years, this tree has spread rapidly east and north. Near San Antonio I saw extensive districts, reported to have been, a few years ago, entirely destitute of even a trace of ligneous growth, and which are now covered with copses of mesquit. Similar growths have sprung up everywhere in the prairies of western Texas. The appearance of this new growth may be traced to the influence of the vast herds of stock which range over the prairies, and which, in voiding the seeds of this tree, assist its wider distribution, and, in keeping down the grass, diminish the quantity of combustible material which feeds the prairie fires, and thus check and finally prevent the spread of the frequent conflagrations which swept year after year over these grassy plains.





10

"West of the Colorado river the pecan-nut is an important product, forming one of the staple articles of export. Shipments of this nut from San Antonio average annually 1,250,000 pounds, obtained from the bottom lands of the Nueces, the Rio Frio, Medina, and Rio Concho. A million pounds, obtained from the Colorado, Guadalupe, Rio Blanco, Pierderelis, Sabinal, Llano, and San Saba rivers, are shipped from Austin, and about a quarter of a million more from Indianola, gathered on the lower Guadalupe, San Antonio, Colorado, and other streams flowing into the Gulf. The nuts are worth, on an average, 5 cents a pound to the gatherer.

"On the range of low hills extending from San Antonio to Austin, which rise at some points to a height of over 500 feet above the plain, forming the base of the terraces leading to the table-land of northern Mexico, the woods are confined to the barrens and the declivities bordering upon them. The open plains on these table-lands are either entirely destitute of ligneous growth, or, when covered with deeper and more fertile soil, support low copses of mesquit. The western juniper is observed here for the first time. It is a tree of low growth, seldom exceeding 35 feet in height, or more than a foot in diameter. It branches at a short distance from the base, forming a broad, round head. The wood is of a dingy, reddish color, fine-grained, hard, and heavy, and in density and durability is not inferior to that of the red cedar. It is knotty, however, from near the base, and furnishes no sticks sufficiently long to allow its use in cabinet-making, and can only be employed for rough construction, posts, palings, etc., for which purposes it is invaluable. The home of the western cedar is found on the rugged highlands which surround the channels of the headwaters of the numerous streams which flow from the eastern declivity of these hills. Here it forms open groves, with scarcely any other woody growth among the somewhat scattered trees. These cedar woods are particularly common upon the brows of the steep escarpments from the base of which issue the large springs which form such a striking feature in this part of the state. In the vicinity of the settlements few of the full-grown trees have been left. The improvidence of the first settlers in obtaining their timber supplies and the prairie fires which ran through these cedar woods in former years have caused the destruction of large areas once covered by this valuable tree. According to my observation, the western cedar prefers a calcareous, dry soil. Its range of distribution seems limited to the hilly region bordering upon the upper part of the Colorado valley, extending toward the south a short distance below New Braunfels, and westward to the sources of the Nueces and Guadalupe rivers. Well-timbered tracts of this tree are still found west of New Braunfels as far as Boerne, in Kendall county, and on the terraces of the higher ranges in Bandera and Kerr counties."

INDIAN TERRITORY.

The forests of the Indian territory are confined to its eastern portion. West of the ninety-ninth meridian trees are only found along the narrow river bottoms, the intervening ridges being bare of all forest growth. The extreme northeastern part of the territory contains numerous extensive open prairies, south of which a heavy body of forest composed of hard woods, mixed on the high ridges with the short-leaved pine, extends southward into Texas, with a maximum width in the Choctaw nation of 60 miles. In the Cherokee nation six considerable bodies of pine, varying from 10 to 30 miles in length and 2 to 4 miles in width, occur on Spavina creek, Illinois river, Salina river, Spring creek, and Bowman's Fork, tributaries of Grand river. A large body of pine occurs also 25 miles west of Reams, a station upon the Missouri, Kansas, and Texas railroad. Smaller bodies of pine are found, too, east of Reams, and at Stringtown, where lumber is manufactured and shipped southward by rail into northern Texas.

The bottom lands of all the streams flowing through the eastern portion of the territory are heavily timbered with hard woods, and especially those of the Neosho, Verdigris, Arkansas, and Canadian rivers contain great bodies of the finest black walnut now growing. A particularly fine growth of this timber extends along the Verdigris river for 50 miles above Coffeetown.

West of the region of heavy forest the country is covered with an open growth of upland oaks, among which the most prominent are the post oak and the black-jack. These forests are interspersed with prairies, often of considerable extent, which gradually occupy the whole country outside the bottom lands. Farther west, between the ninety-seventh and ninety-ninth degrees of west longitude, the "cross-timbers" enter the territory from the south. They are composed, as in Texas, of a stunted growth of post oak and black-jack, and extend northward across the territory in straggling patches into southern Kansas. The main belt of the "cross-timbers", about 70 miles wide at the Texas boundary, gradually becomes narrower toward the north and northwest, disappearing, at about longitude 99° west, upon the ridges south of the Cimarron river.

No returns of the amount of lumber manufactured in the territory have been received, nor other than the most general information in regard to its forest covering.

ARKANSAS.

Heavy forests cover the state of Arkansas, with the exception of a few isolated prairies principally confined to Prairie and Arkansas counties, north of the valley of the Arkansas river, and the western borders of the state. North of the Arkansas river the forests are mostly composed of the deciduous trees of the Mississippi basin, through which isolated belts occur, often of considerable extent, in which the short-leaved pine, the only species found in

northern Arkansas, is mixed with the hard woods. The southwestern part of the state south of the Arkansas river and west of the broad, level plain of the Mississippi is covered outside the river-bottom lands with an almost continuous forest of pine, in which the short-leaved species occupies the high, dry ridges and the loblolly the moist soil above the bottoms. Great bodies of cypress cover the extensive swamps that stretch along the eastern border of the state or line the bottoms of the White, Arkansas, Washita, and Red rivers. The hard-wood forests of the state are hardly surpassed in variety and richness, and contain inestimable bodies of the finest oak, walnut, hickory, and ash timber. Black walnut of large size is still widely scattered over the state, and is particularly abundant in the valley of the Red and other southern rivers. The pine forests are almost intact. Settlements made for agricultural purposes have been confined to bottom lands, and only during the last few years has pine lumber been manufactured in the state, except to supply a very limited local demand. Recently, however, comparatively small quantities of lumber manufactured at numerous railroad mills, principally established south of the Arkansas river, have been shipped north and south out of the state.

The forests of Arkansas have received comparatively little damage from fire. Pine generally succeeds pine even on burned land, although upon certain gravel and clay soils the second growth is largely composed of black and red oaks, or, in the southern part of the state, the sweet gum replaces other trees on bottom lands. During the census year 858,115 acres of woodland were reported devastated by fire, with an estimated loss of \$259,470. The largest number of these fires was due to the carelessness of farmers in clearing land, or to hunters camping in the forest.

Industries consuming hard woods are still in their infancy in Arkansas, although doubtless destined to attain an important development. Rough white-oak staves are largely manufactured in the White River country and in the northeastern part of the state for eastern and European markets.

A considerable traffic exists in the southwestern counties in the wood of the Osage orange, used for wheel stock, and more recently as pavement in Saint Louis and other northern cities.


The following estimates of the amount of short-leaved pine standing in Arkansas May 31, 1880, were prepared by Professor F. L. Harvey, of Fayetteville:

SHORT-LEAVED PINE (*Pinus mitis*).

Counties.	Feet, board measure.	Counties.	Feet, board measure.	Counties.	Feet, board measure.
Ashley	1,555,000,000	Hot Spring	1,348,000,000	Perry	1,023,000,000
Baxter	187,000,000	Howard	1,254,000,000	Phillips	21,000,000
Boone	124,000,000	Independence	93,000,000	Pike	1,695,000,000
Bradley	1,140,000,000	Izard	242,000,000	Poinsett	45,000,000
Calhoun	1,519,000,000	Jefferson	518,000,000	Polk	2,592,000,000
Carroll	159,000,000	Johnson	248,000,000	Pope	208,000,000
Clarke	1,280,000,000	La Fayette	588,000,000	Pulaski	668,000,000
Clay	8,000,000	Lee	14,000,000	Saint Francis	7,000,000
Columbia	1,866,000,000	Lincoln	105,000,000	Saline	933,000,000
Craighead	18,000,000	Little River	690,000,000	Scott	1,516,000,000
Cross	54,000,000	Logan	554,000,000	Searcy	166,000,000
Dallas	1,659,000,000	Lonoke	20,000,000	Sebastian	243,000,000
Dorsey	726,000,000	Madison	55,000,000	Sevier	969,000,000
Drew	482,000,000	Marion	207,000,000	Sharp	35,000,000
Faulkner	42,000,000	Miller	622,000,000	Stone	179,000,000
Fulton	146,000,000	Monroe	180,000,000	Union	2,364,000,000
Garland	1,865,000,000	Montgomery	2,281,000,000	Van Buren	435,000,000
Grant	207,000,000	Nevada	1,453,000,000	White	23,000,000
Greene	88,000,000	Newton	767,000,000	Yell	1,306,000,000
Hempstead	1,176,000,000	Osachita	1,384,000,000		
Total					41,315,000,000
Cut for the census year ending May 31, 1880 (including 57,943,000 shingles and 2,891,000 laths)					129,781,000

TENNESSEE.

The western counties of Tennessee are covered with heavy forests, similar in distribution and density to those which occupy the Yazoo region of western Mississippi. The river swamps in this part of the state still contain large bodies of cypress, while the hills are covered with oaks, hickories, and other hard-wood trees. The central portion of the state, now largely cleared for cultivation, was once covered with forests of hard wood, remnants of which are still found upon rocky ridges or land unfit for agriculture. Nearly through the center of this middle district, extending north and south, "the cedar glades" occupy an extensive region of Silurian limestone. Here the characteristic growth consists of red cedar (*Juniperus Virginiana*), often forming stunted forests of considerable extent, to the exclusion of other species, or is mixed with the honey locust, a characteristic species, also, of this well-marked region.



The eastern part of the state, occupied by the Cumberland plateau and the high ranges of the southern Alleghany mountains, is covered with a heavy forest of oak and other hard woods, mixed at high elevations with hemlock, pine, and spruce, and constituting one of the finest bodies of timber now standing in the United States. It contains, besides white and chestnut oak of fine quality, much yellow poplar, black walnut, and cherry. In the southeastern counties, especially in the valley of the Tennessee river, the hard-wood forests have been, however, already destroyed over large areas to furnish charcoal for the iron-manufacturing industry established here.

During the census year 985,430 acres of woodland were reported devastated by fire, with a loss of \$5,254,980. Of these fires the largest number was set in the careless clearing of land for agriculture or to improve grazing, and by hunters, locomotives, etc.

Mr. A. G. Willey, of Manchester, Tennessee, has supplied the following statement in regard to the effects produced upon the forest growth by the annual burning of dead herbage to improve pasturage:

"EFFECT OF FIRES UPON THE FOREST.

"The practice of burning timber-land, said to have been of Indian origin, has been continued by the white settlers. The native grasses do not die down when killed by frost; they simply die standing, and the young grass in the spring has to push through the old tuft, which is often 6 or 8 inches high. The fires are set in the timber and old fields to burn these tufts, that stock may graze four or six weeks earlier than if the old herbage had been left upon the ground. In the barrens and on the Cumberland plateau the timber is principally oak of various kinds, which do not shed their leaves at once when killed by frost, or rot when partially green, but remain dry upon the trees and fall gradually during winter and spring. The largest portion, therefore, are on the ground in February, the time when fires are set. The effect of these fires is to destroy all the natural sources of fertility, grass, leaves, and fallen timber. Had these been allowed to accumulate, what are now called barren lands would be the most fertile in the state. The practice kills, too, the young trees, so that some of the most valuable timber that the land is suitable to produce is unable to stand. The black-jack, post oak, black oak, etc., however, on account of the protection afforded by their thick bark, are able to gain some headway, and so crowd out more valuable trees. The state law makes it a misdemeanor with heavy penalty for any one to set fire to and burn a neighbor's land; but the difficulty of detection and conviction in such cases makes this law non-effective. These are the causes and effects of forest fires in this section; they never occur here in summer."

Considerable cooperage and wheel stock is manufactured in Tennessee, but, except in the eastern part of the state, manufacturers report a scarcity and deterioration of the best hard woods, especially white oak. In the eastern counties the manufacture of oak staves and other industries using hard woods are capable of large development.

The principal center of lumber manufacture in the state is Nashville, where several mills saw large quantities of black walnut, poplar, cherry, ash, oak, etc., received by raft from the upper Cumberland river in Tennessee and Kentucky. The local market takes about one-third of the lumber manufactured here, the remainder being sent north and east by rail. Memphis, on the Mississippi river, is also an important manufacturing center. The mills here are largely supplied by rafts from Missouri, Arkansas, and Tennessee, and saw large quantities of cypress, ash, poplar, hickory, gum, and black walnut. Considerable hard-wood lumber manufactured in Dyer, Lincoln, Obion, and Smith counties, and pine and hard-wood lumber in Knox and Jefferson, largely from logs obtained in the vicinity of the mills, is principally consumed locally.

KENTUCKY.

The forests of Kentucky resemble in general features those of Tennessee. Cypress, gum, and various water oaks occupy the river swamps of the western counties. The central region, now largely cleared and devoted to agriculture, was once covered with the oaks, walnuts, and hickories of the Atlantic region, while over the eastern and southeastern counties the dense forests of the Alleghany mountains extended. The eastern counties still contain great bodies of the best hard wood, especially black walnut, white oak, cherry, and yellow poplar, which are particularly fine and abundant in Bell, Harlan, and other southeastern counties. These forests, protected by the falls of the Cumberland river, which have prevented the driving of logs from its upper waters, and inaccessible to rail communication, are still practically uninjured, and probably unsurpassed in the amount, quality, and value of the timber which they contain. The destruction of forests to supply numerous iron furnaces with charcoal has been great in the northeastern counties, and no small part of this region has already been cut over.

During the census year 556,647 acres of woodland were reported devastated by fire, with an estimated loss of \$237,635. Of these fires by far the largest number was traced to farmers carelessly clearing land for agricultural purposes.

In Barren, Edmonson, and other central counties extensive tracts of prairie existed at the time of the earliest settlement of the state. The presence of these prairies in the midst of a heavily-timbered region is ascribed to the annual burning to which they were subjected by the aborigines. With the disappearance of the

Indians trees sprang up, and this region is now well covered with a vigorous growth of black oaks of different species. White oaks, however, are not abundant, and other species common to the region, such as the walnuts, the yellow poplar, and the beech, are wanting in these young forests, indicating perhaps the effect of fires in checking the subsequent growth or development of many useful timber trees.

PASTURAGE OF WOODLANDS.

The forests of Kentucky, as well as those of all the central and southern portion of the United States, suffer severely from the almost universal custom of using woodlands for pasturage. The evil resulting from this practice is only more apparent in Kentucky and Tennessee, because in these states the amount of live stock is proportionately larger than in other parts of the south, while in the thickly-settled agricultural sections of these states the ratio of woodland to total area is smaller. The pasturage of woodlands necessitates, or at least induces, the annual burning of the dead herbage, by which underbrush, young trees, seedlings, and seeds are destroyed and the succession and permanence of the forest endangered. What the fires spare, browsing animals devour; hogs root out seedlings, and by selecting the sweet acorns of the white oak in preference to the bitter fruit of the black oaks, are gradually changing the composition of the oak forests. Comparatively few white oaks spring up in the forests of the more thickly settled portions of the central Atlantic region, and this change of forest composition must be ascribed to the preference of domestic animals for the palatable fruit of what, as regards their timber, are the most valuable species. The injury, too, inflicted by the constant stamping of animals and consequent packing of the land about the stems of old trees is very great, and all reports speak of the gradual dying of old trees left standing in the grazing regions of Kentucky and Tennessee.

The spread of the mistletoe (*Phoradendron flavescens*), consequent upon the removal of the forest and the increase in the number of birds (the mistletoe seems to require a certain amount of light and air for its development; it does not flourish or increase rapidly in the dense forest, and cannot spread except by the agency of birds), is a cause of serious injury to the forest of this whole region. It slowly but surely destroys the trees upon which it obtains a foothold. The black walnut especially suffers from the growth of this parasite, which seems destined to destroy the finest walnut timber left standing in the settled portions of the southern central region.

Large quantities of cooperage and wheel stock are produced all over the state, and manufacturers generally report no scarcity or deterioration of timber, with the exception of white oak. The principal centers of lumber manufacture are at the mouth of the Tennessee river, in McCracken county, where a large amount of cypress, sycamore, gum, oak, walnut, and other hard wood is manufactured for the northern market from logs rafted down the Tennessee and other streams flowing into the Mississippi; at Frankfort, where poplar, oak, ash, walnut, pine, cherry, hickory, and maple logs, rafted from the upper waters of the Kentucky river, are sawed, the lumber being shipped north and east by rail; and at Louisville, where walnut, poplar, and oak lumber is manufactured for local consumption. The manufacture of pumps and water-pipes from logs of the Jersey pine (*Pinus inops*), at one time an important industry at Louisville, has, since the general introduction of city and town water-works, become unremunerative and unimportant.

NORTHERN CENTRAL DIVISION.

OHIO.

The forests of Ohio were originally composed of deciduous species, among which, in the eastern and especially in the northeastern counties, white pine and hemlock existed in isolated bodies of no great extent.

The original forest has now been generally removed, except from Ottawa, Miami, Montgomery, and a few other western counties, and from swamps and other lands unfit for agriculture; everywhere the walnut and other valuable timbers have been culled, and Ohio must soon depend almost exclusively for the lumber which it consumes upon the northern pineries and the hard-wood forests of the south.

During the census year 74,114 acres of woodland were reported destroyed by fire, with an estimated loss of \$797,170. Of these fires the largest number was traced to carelessness in clearing land, to hunters, sparks from locomotives, etc.

The production of cooperage stock has long been an important industry in the state; it has already suffered from a scarcity and deterioration of white oak, for which elm, beech, maple, and poplar are now often substituted. Manufacturers of wheel stock, furniture, woodenware, etc., report abundant material for present consumption.

Ohio is sixth among the states in the volume of its lumber-manufacturing interests. The business is widely distributed throughout the state, generally in the hands of small manufacturers operating portable mills, which threaten the rapid destruction of the remnants of her forests.

INDIANA.

Indiana was once almost entirely covered with noble forests of deciduous trees. Along its western borders these were interrupted, however, by numerous small prairies, the extreme eastern outposts of the great treeless region which, toward the north, extended over the counties of Benton, Newton, and Jasper, and over considerable portions of Lake, Porter, La Porte, Pulaski, White, Tippecanoe, and Warren counties. These prairies have gradually decreased in area with the settlement of the country, and those originally of small extent are now covered with a vigorous growth of the forest trees of the region.

The forests of Indiana are characterized by an almost entire absence of coniferous trees. Stunted white and gray pines occupy the sand-dunes which border the southern shores of lake Michigan, and "the knobs"—low, gravelly hills of small extent, in the southeastern river counties—are covered with a heavy growth of the Jersey pine. Swamps in the southwestern counties contain cypress, which finds here the northern limit of its distribution. The broad bottom lands and low ridges of this part of the state are covered with a forest growth probably unsurpassed in the development of individual trees, and rarely equaled in the richness of its composition.

The forests of the state have been largely removed in the development of its agriculture. No large bodies of the original timber remain. The black walnut with which the forests of Indiana once abounded has been everywhere culled and is now rare, while the best yellow poplar, oak, and other valuable timbers have been largely consumed.

During the census year 90,427 acres of woodland were reported injured by fire, with an estimated loss of \$130,335. These fires were set by farmers carelessly clearing land, by hunters, and by sparks from locomotives.

The forests of Indiana have long supplied material for a large manufacture of cooperage stock, furniture, wagon stock, woodenware, etc. The cooperage and furniture manufacturers already feel the scarcity and deterioration of the highest grades of oak and walnut, and very generally predict the entire exhaustion at no very distant day of the forests of the state.

Indiana is fifth among the states in the value of its lumber-manufacturing interests. Evansville, upon the Ohio river, in Vanderburgh county, is an important manufacturing center on account of the capital invested there in the lumber business and the amount of its product. The business, however, as in Ohio, is generally in the hands of small manufacturers operating portable mills and sawing logs hauled to them by farmers. At the present rate of destruction the forests of the state must soon lose all commercial importance.

ILLINOIS.

The forests of Illinois were originally confined to the southern portion of the state, the broad bottom lands of the Mississippi and the Illinois, and the southern third of the delta formed by these rivers. The remainder of the state was covered by broad, rolling prairies. The forest growth in this prairie region was confined to the narrow river bottoms and occasional open park-like groves of burr, scarlet, red, black-jack, or post oaks, known as "oak openings",

through which the prairie fires swept, destroying all undergrowth, without doing great injury to the full-grown trees. Prairie fires have gradually decreased in frequency and violence since the settlement of the state, and these open groves are now filled with a vigorous growth of young seedlings and shoots; their characteristic features have disappeared, and the area of the forest is gradually increasing.

The shores of lake Michigan are covered with a stunted growth of white pine; the dry, rocky hillsides in the western part of Union county, one of the southern counties of the state, bear a few yellow pines (*Pinus mitis*), and cypress is found in the southern river swamps. With these exceptions, of little importance commercially, the forests of Illinois are composed of deciduous species.

During the census year only 48,691 acres of woodland were reported destroyed by fire, with an estimated loss of \$45,775. These fires were generally traced to hunters, and to farmers permitting brush fires to escape to the forest.

The production of cooperage stock was once an important industry in southern Illinois. The business has greatly diminished, owing to the exhaustion of the local supply of the best hard woods. Bass, gum, hackberry, elm, sycamore, and other woods formerly considered of little value, are substituted for oak, and Illinois now receives most of its hard wood from Kentucky, Tennessee, and other southern states.

Illinois is eleventh among the states in the volume of its lumber-manufacturing interests. It owes this position to the fact that many large mills sawing pine logs rafted down the Mississippi river from the forests of Wisconsin are established within its borders, and not to the extent and value of the forests of the state. The manufacture of Illinois-grown lumber is small and totally inadequate to supply the wants of the present population of the state.

Chicago, owing to its general commercial importance and its position with reference to the great pine forests of the northwest, has become the greatest lumber-distributing center in the world.

According to the statistics gathered by the *Northwestern Lumberman* of Chicago, and published in that journal January 29, 1881, there were received in Chicago during the year 1880 1,419,974,000 feet of lumber by lake and 145,563,118 feet by rail, a total of 1,565,537,118 feet, an increase of 96,817,127 feet over the total receipts of 1879; 650,922,500 shingles were received during the same year.

Lumber was received from the lake ports during the year 1880, as follows:

Points of shipment.	Lumber.	Shingles.	Points of shipment.	Lumber.	Shingles.
	<i>Feet.</i>	<i>Number.</i>		<i>Feet.</i>	<i>Number.</i>
Ahnapee	150,000		Menominee	225,110,000	760,000
Alpena	4,517,000		Monastique	202,000	
Ashland	5,200,000	1,311,000	Muskegon	451,854,000	23,660,000
Bay de Noquet	3,670,000		North Bay	110,000	
Bayfield	980,000		Oconto	11,003,000	395,000
Benton	3,876,000		Ontonagon	2,503,000	2,886,000
Black Creek	4,825,000		Oscoda	739,000	
Black River	6,858,000		Packard's pier	2,681,000	
Canada ports	755,000		Paul's pier	500,000	
Caseville	200,000		Pensaukee	6,866,000	3,190,000
Cedar River	17,383,000	100,000	Pentwater	9,596,000	25,572,000
Charlevoix	1,541,000		Perry's pier	45,000	
Cheboygan	33,250,000		Peshtigo	51,600,000	7,920,000
Clay Bank		650,000	Pierport	3,355,000	
Copper Harbor	70,000		Point Saint Ignace	12,985,000	
Cross Village	233,000		Portage Lake	735,000	
Depere	250,000	1,611,000	Port Huron	344,000	
Duck lake	1,340,000		Port Sheldon	180,000	
Escanaba	5,182,000	3,457,000	Red River	200,000	3,857,000
Ford River	17,850,000	6,915,000	Rogers City	1,966,000	
Frankfort	9,565,000		Saginaw River	11,926,000	
Grand Haven	90,166,000	114,000,000	Saint Joseph	1,662,000	
Green Bay	1,577,000	22,562,000	Saugatuck	4,014,000	4,000,000
Hamlin	12,822,000	11,026,000	Sault Ste. Marie	522,000	
Hancock	300,000		Silver Lake	2,185,000	600,000
Holland	857,000		South Haven	3,650,000	300,000
Kewaunee	110,000	5,881,000	Sturgeon Bay	11,640,000	19,978,000
L'Anse	9,430,000	170,000	Suamico	3,065,000	2,480,000
Leland	970,000		Traverse	23,280,000	
Lincoln	1,235,000	300,000	Whitefish Bay	730,000	
Ludington	103,713,000	34,330,000	White Lake	66,603,000	24,756,000
Ludwig's pier	125,000				
Mackinaw City	275,000		Total	1,419,974,000	583,340,000
Manistee	165,217,000	259,911,000	Receipts by rail	145,563,118	67,582,500
Manitowoc	70,000	300,000			
Marquette	2,411,000	522,000			
Masonville	1,030,000		Grand total	1,565,537,118	650,922,500

Lumber was received by rail during the year 1880, as follows :

Names of lines.	Lumber.	Shingles.
	<i>Feet.</i>	<i>Number.</i>
Baltimore and Ohio railroad	9,696,000	
Chicago and Alton railroad	988,000	70,000
Chicago and Eastern Illinois railroad	26,799,000	
Chicago and Grand Trunk railway	1,506,418	80,000
Chicago and Northwestern railway	11,727,900	44,642,000
Chicago, Burlington, and Quincy railroad	3,716,800	
Chicago, Milwaukee, and Saint Paul railway	12,473,000	13,180,500
Chicago, Rock Island, and Pacific railway	2,224,000	
Illinois Central railroad	2,940,000	
Lake Shore and Michigan Southern railway	18,636,000	1,385,000
Michigan Central railroad	24,798,000	8,175,000
Pittsburgh, Cincinnati, and Saint Louis railway	12,481,000	
Pittsburgh, Fort Wayne, and Chicago railway	17,567,000	50,000
Wabash, Saint Louis, and Pacific railway	610,000	
Total	145,563,118	67,582,500

The following account of the early lumber trade of Chicago is condensed from a paper prepared by Mr. George W. Hotchkiss, secretary of the Chicago lumber exchange, and printed in the *Northwestern Lumberman* under date of March 19, 1881 :

"Colonel Mann, residing at Calumet, brought the first raft of lumber to Chicago. It was square building timber, poled from the mouth of the Calumet to the mouth of the Chicago river. The value of this raft was \$100, and its owner found considerable difficulty in disposing of it. In 1834 or 1835 Captain Carver opened a lumber-yard on the river bank, near the present site of the State-street bridge, and about the same time a man named Harrison owned a small schooner which went to some point across the lake and brought in white wood. This little vessel could not enter the river, on account of the bar across its mouth, and her cargo was unloaded upon scows and rafts, which were floated southward for half a mile or more, around the end of the bar, before they could be headed for the deep water of the river. In 1835 or 1836 a man named Rossiter had a small dock and yard on the river, between Clark and La Salle streets, and by this time other yards were started on the river. About the year 1836 a man named Cammack had a pit-mill on the north branch of the river. His son acted as pit-man, the old man being the top-sawyer. It is not unlikely that the first lumber used in Chicago was manufactured by this method, although about the same time a wind saw-mill was located not far from the present Kinzie-street bridge, which found abundant occupation in sawing white-wood timber, which then grew in the immediate neighborhood, mixed with elm, ash, basswood, and a few oak trees. History does not record, however, that the market was overstocked by the product of this mill, or that the lumber dealers of that day hurried to issue a new price-list low enough to crush the aspirations of their dangerous competitor. In fact, history is so perfectly silent upon the subject of this saw-mill that it is probable its work did not cut much of a figure in the lumber trade of that day, and that, in fact, it proved a veritable windmill, of less caliber than the muscle of the Cammacks, who no doubt found greater profit, if harder work, in driving their pit-saw. Captain Carver's lumber-yard was on the river bank, just west of the present State-street bridge, having a light, temporary dock, upon which the small vessels bringing lumber to the river unloaded. There was at this time (1836 or 1837) no other lumber-yard in the village upon the river. Captain Carver afterward (about 1839) sold out to George W. Snow, who occupied the same ground for a number of years.

"The earliest lumber of which Mr. Hilliard has any recollection came from Saint Joseph, Michigan; but shortly after his arrival at Chicago a man named Conroe built a mill at Manitowoc, Wisconsin, and Jones, King & Co., who were then doing a hardware and general business, received and handled his lumber as a side issue. A small pocket saw-mill, built by a man named Huntoon, in 1836, was located on the river bank not far from the present Chicago-avenue bridge. It was too small to do much work, but was esteemed a very useful and really wonderful mill at that time. The North Side was pretty well timbered with elm, oak, and white wood, and from this timber the mill obtained its stock. After the streets were cut out the wet nature of the ground compelled one who would visit this saw-mill to pick his way to it by jumping from log to log. It was so far from the village to the mill that it was seldom visited, except by those who enjoyed a Sunday walk and could find no objective point of greater interest for their stroll. The lumber-yard of Tuckerman & Higginson was located in 1843 on the north side of the river, near the present northwest end of Clark-street bridge. Clark street above Kinzie street had been cleared of timber, and a clear view was to be had as far as the eye could reach in a western direction, broken only by a few scattering trees which had been left as sentinels upon the plain. At this time George W. Snow had a yard on the river, near State street, and a Mr. Rossiter had also a yard between Newberry & Doles' warehouse, on the south branch of the river, west of what is now Clark street. Barber & Mason had a yard a little farther west, near Wells street. J. M. Underwood and Sylvester Lind each had a yard on the west side of the river, near Randolph

street. This was in 1844. Mr. Higginson obtained his supply of lumber in those days from Hall & Jerome, of Menominee, Michigan, Elisha Bailey, of Peshtigo, Wisconsin, and ——— Fisk, of Depere. In 1845 he had a contract for 1,000,000 feet with William F. Ferry, of Grand Haven. Lumber came also from Kalamazoo river, Saint Joseph, and Muskegon. In 1844 Mr. Higginson purchased a cargo from Mr. Rose, of Muskegon, and, as it was a beautiful lot of lumber, running 33½ per cent. upper grades, he was willing to pay a good price, obtaining it at \$5 75 per thousand feet. The first cargo of Saginaw lumber which reached Chicago was brought by James Fraser, one of the original proprietors of the plat of Bay City, who built two mills at Kawkawlin, in latter years known as the Ballon mills. This was in the year 1847 or 1848, and the cargo attracted a good deal of attention, because it was the first lot of circular-sawed lumber that had ever been seen by any of the dealers, and because of its general cleanness of appearance, the attractiveness of a lot of circular-sawed sidings among it, and its excellent quality. All these combined to make the cargo a novelty in its way, and it found a sale at \$8 per thousand feet, an extra good price for those days. Average cargoes at this time were quoted at \$6 50 to \$7 for mill-run lumber, culls out, and it did not need a very coarse piece to rank as a cull. Culls were rated at half price. The retail market held common lumber at about \$8 during the summer, and \$9 was asked for dry lumber through the winter. Common included everything below first and second clear; third clear, selects, picks, and finishing grades generally, being an invention of a later day. First clear sold at from \$12 to \$16, and second clear at \$10 to \$12; clear, undressed flooring brought \$12, and common flooring \$10. The lath trade was mostly in what was known as board lath, although narrow lath arrived in small quantities. The trade of the city in 1843 was about 12,000,000 feet, and this was considered as remarkable as to us were last year's sales of 1,500,000,000 feet, or about 140 times as much more."

MICHIGAN.

Michigan once possessed a tree covering of great density, richness, and variety. The hard-wood forests of the Ohio valley covered the southern portion of the state, extending to just north of the forty-third degree of latitude. North of this hard-wood belt the character of the forest changed; the white pine appeared, occupying the drier and more gravelly ridges, and, gradually increasing in size and frequency, became the most important element in the forests of the central and northern portions of the southern peninsula. In the northern peninsula, especially in the basin of the Menominee river, it covered the sandy plains almost to the exclusion of other species. The forests of hard wood, occupying low, rich soil between the pine-covered ridges, were valuable in their stores of sugar maple, birch, ash, beech, oak, and other northern trees, while the swamps common in the northern part of the state abounded in tamarack and yellow cedar of large size and excellent quality.


North of the central portion of the lower peninsula large tracts of barren plains exist. One of the most extensive of these tracts occupies a considerable portion of Crawford county, covering an area of several hundred square miles. A second barren region exists in Lake county, and there are others in Ogemaw and Iosco counties; similar barrens occur in the northern peninsula, the largest in Schoolcraft and Marquette counties. The soil covering these barrens is a light sandy loam, supporting a stunted growth of gray pine, birches, poplars, and scrub oak. These sandy plains owe their existence, perhaps, to the continual burning of the forest, prostrated possibly, in the first instance, by tornadoes, and thus affording abundant material for a fire hot enough to consume the vegetable mold of the surface and render the soil unfit to produce a second growth of heavy timber, or in many instances any tree growth whatever.


Serious inroads have already been made upon the forests of Michigan. The hard wood has been generally cleared from the southern counties, now largely occupied by farms, and the timber remaining in this part of the state, in small, scattered bodies, can hardly suffice for the wants of its agricultural population. The merchantable white pine has been cut from the banks of the principal streams and the shores of the lakes, and what now remains is remote from water transportation or scattered in isolated bodies of comparatively small extent. The hard-wood forests of the pine belt, however, although greatly injured by fire in parts of the state from which the pine has been cut, and invaded along their southern borders by agricultural settlements, contain, especially in the northern third of the lower peninsula and through the northern peninsula, vast quantities of valuable timber.

FOREST FIRES.

The forests of Michigan have long suffered from destructive fires. These have generally originated in the neighborhood of the loggers' camps or upon the farms of the agricultural pioneer, while the virgin forest has generally, although not always, escaped serious conflagrations. The timber-prospecter and the hunter are responsible for many fires in the primeval pine forest of the northwest; but, as a rule, fires follow and do not precede the lumberman. The reason is obvious: The logger in his operations leaves the resinous tops, branches, and chips of the pine trees scattered far and wide; these by the following midsummer become dry as tinder, and afford abundant material to feed a fire started by a careless hunter, log-cutter, or farmer clearing land near the forest. Such fires, which too often follow the cutting of pine forests of the northwest, have inflicted incalculable injury upon the country. They have destroyed vast quantities of hard-wood timber; they have consumed the young

DEPARTMENT OF THE INTERIOR






DENSITY OF FORESTS

COMPILED UNDER THE DIRECTION OF
C. S. SARGENT, SPECIAL AGENT
1883.

Scale






()

MAP OF THE
UPPER PENINSULA OF MICHIGAN
SHOWING THE DISTRIBUTION OF FORESTS,
WITH SPECIAL REFERENCE TO THE
LUMBER INDUSTRY
COMPILED UNDER THE DIRECTION OF
C.S. SARGENT, SPECIAL AGENT.
1881.

- LEGEND
- Hard Wood
 - Standing Pine
 - Pine and Hard Wood
 - Cut Pine
 - Cut Pine and Hard Wood
 - Barrens



Scale 0 25 50 75 100 Miles

pine trees left by the logger; they have robbed the soil of its fertility, and made it unfit to produce another crop of pine until the growth and decay of generations of other plants shall have restored its lost constituents. In the dense, uncultured forest, on the other hand, fires, although often destructive, are less dangerous in the absence of dead material to feed the flames than when the ground is strewn with dead branches, tops, and resinous chips.

During the census year only 238,271 acres of woodland were reported destroyed by fire, with an estimated loss of \$985,985. Of the 267 fires reported, 161 were traced to fires set in clearing land for agricultural purposes, and which escaped to the forests; 59 to hunters, 43 to sparks from locomotives, 3 to smokers, while only 1 was reported set by Indians.

The hard-wood forests of Michigan have long afforded abundant material for large and important industries engaged in the production of cooperage stock, handles, oars, agricultural implements, excelsior, wood pulp, etc. Manufacturers, especially in the southern part of the state, now report, however, a scarcity and general deterioration of stock. The best oak timber has been everywhere culled to supply the wants of railroads or the demands of the Canadian market. Elm, bass, and other soft woods, which a few years ago were considered of little value, are now in great demand and are fast disappearing, except from regions remote from railroads. Much hard wood, especially in the southern peninsula, has been destroyed by fire, or, if not destroyed, rendered almost worthless for manufacturing purposes by partial burning.

Next to Vermont and New York, Michigan produces a larger amount of maple sugar than any other state. During the year 1879 3,423,149 pounds were manufactured in the state.

STATISTICS OF GROWING TIMBER.

The following estimates of the merchantable timber standing in Michigan May 31, 1880, were prepared by Mr. H. C. Putnam, of Eau Claire, Wisconsin, with the assistance, in the lower peninsula especially, of Mr. G. W. Hotchkiss. These, as well as the estimates of the timber resources of Wisconsin and Minnesota, were obtained by compiling the results of actual surveys, and have been further verified by a large number of persons familiar with the forests in the different regions of these states. It must not, however, be forgotten that the figures given represent estimates, and not facts. Statistics of the volume of any growing crop are difficult to obtain and always liable to considerable error, and the forest, from its very nature and the extent over which it is spread, presents greater difficulties to the collector of statistics of productive capacity than the more compact and more easily studied crops of the field. The estimates of pine include all trees 12 inches in diameter 24 feet from the ground. Since they were prepared the scarcity of white pine has changed the methods of the lumberman, and trees are now generally estimated and cut as small as 8 inches in diameter 24 feet from the ground. If the amount of standing pine had been estimated upon the 8-inch basis it would have added (roughly) 10 per cent. to Mr. Putnam's figures. Small bodies of pine remote from streams no doubt exist in different parts of Michigan, Wisconsin, and Minnesota, in the aggregate of some commercial importance, which are not included in these estimates. The following figures, however, are believed to represent with as great accuracy as is attainable the productive capacity of the northwestern pineries. They cover the entire region, and these pine forests now contain no great body of unexplored timber, an unknown factor in the country's lumber supply:

WHITE PINE (*Pinus Strobus*).

Regions.	Feet, board measure.
LOWER PENINSULA.	
Basins of streams flowing into Saginaw bay, including Saginaw river and tributaries.	7,000,000,000
Basins of streams flowing into lake Huron.....	8,000,000,000
Basins of streams flowing into lake Michigan.....	14,000,000,000
Total	29,000,000,000
Cut for the census year ending May 31, 1880 (including 2,936,600,000 shingles and 428,445,000 laths, but exclusive of 36,000,000 staves and 8,380,000 sets headings).	4,068,773,000
UPPER PENINSULA.	
Basin of Menominee river and tributaries (Marquette and Menominee counties).	1,600,000,000
Ontonagon, Houghton, Keweenaw, Baraga, Marquette (west and north of Menominee basin), and Menominee (east of Menominee basin) counties.	2,400,000,000
Schoolcraft, Chippewa, Mackinac, and Delta counties.....	2,000,000,000
Total	6,000,000,000
Cut for the census year ending May 31, 1880 (including 106,482,000 shingles and 34,260,000 laths).	328,438,000

An estimated amount of 575,500,000 cords of hard wood is distributed over some 20,000,000 acres in the lower peninsula. Of this about 20 per cent. is suitable for lumber and cooperage stock. The cut of hard wood for the census year ending May 31, 1880 (exclusive of 163,821,000 staves and 18,567,000 sets headings, and including 6,038,000 feet of spool stock), was 440,944,000 feet. In scattered swamps there are standing some 5,000,000 cords of yellow cedar (*Thuya occidentalis*).

From Menominee and Delta counties the merchantable pine has been almost entirely removed. Baraga county contains little pine, and Keweenaw county a single considerable body some 30,000 acres in extent.

The northern portion of Ontonagon and Marquette counties is chiefly covered with hard wood.

An estimated amount of 124,500,000 cords of hard wood is distributed over some 10,000,000 acres in the upper peninsula. The cut of hard wood for the census year ending May 31, 1880 (exclusive of fuel and railroad ties), was 1,145,000 feet.

The southern counties of the upper peninsula contain large areas of swamp, covered with tamarack and yellow cedar (*Thuya occidentalis*), estimated, in the aggregate, at 62,500,000 cords.

Some 7,000,000,000 feet of hemlock lumber and 7,000,000 cords of bark still remain in the state.

Michigan is first among the states in the volume and value of its lumber product. Its principal centers of lumber manufacture are Muskegon, on the shores of lake Michigan, the shores of Saginaw bay, in Bay county, the Saginaw river, in Saginaw county, Manistee, and Menominee, in the upper peninsula. The valley of the Saginaw was long the seat of the most important lumber-manufacturing operations in the United States. Its supremacy, however, has departed with the destruction of the splendid pine forest which covered its water-shed, and the center of manufacture has moved westward from the shores of lake Huron across the peninsula to the waters flowing into lake Michigan.

Lumber was first manufactured in the Saginaw valley as early as 1832. Three years later a second mill, with an annual capacity of 3,000,000 feet, was built upon the Saginaw. In 1836 the first shipments of lumber were made from this mill, and from that time forward great attention was given to the manufacture of lumber for shipment. The commercial panic of 1837, however, seriously interfered with the development of this business, and it was not until 1849 that mills began to multiply. In 1844 there were 23 mills upon the Saginaw, with an aggregate capacity of 60,000,000 feet. Ten years later the number of mills had increased to 82, manufacturing 425,000,000 feet of lumber, while in 1873 there were 83 mills, which produced that year 567,000,000 feet. Since 1870 there has been an almost steady decrease in the number of mills operating in the Saginaw valley; the number finishing their "cut out" is fast increasing, and those destroyed by fire are not rebuilt. But, although the number of mills has decreased, their production has increased, their present capacity being estimated at 923,000,000 feet. A large part of the lumber manufactured upon the Saginaw is transported by lake to Ohio and New York ports, and thence to the principal eastern markets, although a considerable amount is shipped by vessel to Chicago and Milwaukee, and thence distributed by rail through the west. The wide market open to this lumber is due to its excellent quality. Twenty years ago logs which would run 25 per cent. "uppers" were considered common; 40 per cent. was the rule, and as high as 75 per cent. "uppers" was sometimes obtained. Logs were then cut from the lower trunk of the tree below the tops, and only the largest trees were selected. Now land which has been cut over three times is gone over again, and lumbermen are satisfied if logs yield 10 per cent. "uppers".

Of late years considerable changes have been introduced into Michigan lumbering operations by railroad logging; by this means mills are able to obtain a constant supply of logs by railroads built into the forest for the purpose, and this supply can be regulated almost entirely by the demand. There are several roads in different parts of the state doing this business, the principal being the Flint and Pere Marquette and the Lake George and Muskegon River railroads. The growth of this business in the Saginaw valley and at Muskegon, Manistee, and on the Flint and Pere Marquette road is shown by the following table extracted from *Bradstreet's* of February 5, 1881:

Years.	Saginaw valley.	Muskegon.	Manistee.	Flint and Pere Marquette railroad.
1865.....	200, 000, 000	108, 505, 700		
1866.....	209, 000, 000	157, 468, 300		
1867.....	429, 207, 808	288, 502, 200		
1868.....	446, 960, 583	213, 692, 000		
1869.....	321, 350, 663	267, 789, 900		
1870.....	623, 397, 353	198, 862, 600	121, 221, 395	
1871.....	521, 796, 927	250, 000, 000	142, 869, 617	
1872.....	645, 285, 278	815, 000, 000	155, 556, 729	
1873.....	690, 979, 461	376, 035, 037	179, 820, 243	
1874.....	589, 225, 404	224, 571, 527	182, 218, 383	
1875.....	584, 843, 701	309, 638, 418	168, 926, 197	
1876.....	572, 229, 472	299, 525, 919	147, 724, 241	
1877.....	651, 567, 948	312, 285, 951	152, 221, 548	
1878.....	558, 079, 674	340, 990, 055	178, 542, 889	
1879.....	780, 182, 286	432, 431, 679	211, 722, 030	14, 357, 670
1880.....	948, 174, 274	380, 000, 000	211, 971, 000	87, 485, 547

The following extracts are made from Mr. Putnam's report upon the forests of Michigan:

"The southern boundary of the pine forest in Michigan may be represented by a line drawn from Sarnia westward across the state nearly to the mouth of the Kalamazoo river. Originally the pine forest covered the northern two-thirds of the state, and estimates made in 1835 gave the amount of pine then standing as 150,000,000,000 feet. This estimate included the northern peninsula. The present estimate of the pine standing in the whole state, the northern peninsula also included, is 35,000,000,000 feet. There are now remaining no large bodies of standing pine in the state which have not been more or less cut into, and the timber adjacent to streams has all been cut. The pine now remaining is scattered generally through the northern half of the state, lying back at a distance of from 2 to 10 miles from streams large enough to float the logs. The best pine in the state has been cut. The belt of pine which ran through the center of the state, extending north from the southern boundary of the original pine forest for some 75 miles, contained the best pine in the northwest. This pine was what was called by lumbermen 'cork pine', a soft white pine, large and sound, with a thick bark. The quality of the pine of the Saginaw valley was particularly fine, too; that on the west shore was of smaller size.

"The standing pine on the lower peninsula of Michigan is estimated at 29,000,000,000 feet, of which there are in the Saginaw valley about 7,000,000,000 feet, including the pine upon the Saginaw, Au Sable, and Cheboygan rivers and their tributaries; on the streams flowing directly into lake Huron there are some 8,000,000,000 feet more; making 15,000,000,000 feet upon the streams of the east shore. On the western shore of the state there are 14,000,000,000 feet, including the pine upon the Kalamazoo, Black, Grand, Muskegon, White, Pentwater, Aux Bec Scies, Boardman, and Pine rivers. As before stated, the quality of the timber in the eastern portion of the state is better than that upon the west shore; this is smaller and partakes more of the sapling nature, while that on the east shore is largely cork pine. The pine of the east shore and Saginaw valley is largely used for finishing lumber, and should be transported to the east; indeed all the pine in the lower peninsula of Michigan is wanted at the east, and none should be sent west. The pine of the western shore is suited for fencing, flooring, and dimension stuff, being smaller and containing more knots and sap.

"The largest bodies of pine left in the lower peninsula are in the counties of Presque Isle, Montmorenci, Alpena, Alcona, Ogemaw, Roscommon, Crawford, Missaukee, Wexford, Manistee, Grand Traverse, Lake, Osceola, Clare, Gladwin, and Charlevoix. There are bodies of pine also in other counties from 15,000 to 20,000 acres in extent which have not yet been cut. The pine left in the lower peninsula is generally scattered through hard-wood timber, into which the settlers are now entering, clearing the hard-wood forests and exposing the pine to destruction by fire and windfall. This destruction has largely increased with the settlement of the country, and will increase still more unless stringent measures can be taken to protect the pine forests from waste.

"The southern part of the state outside the pine belt was originally covered with a dense forest of hard-wood timber; this region is now largely settled and is the farming region of Michigan. There is a large amount of hard-wood timber of commercial value still scattered through this farming country, particularly in its middle and northern parts. Along the west shore as far north as the straits of Mackinaw the pine has been cut in large quantities, but there is still a large amount of hard-wood timber left upon this area.

"The pine of the northern peninsula of Michigan is estimated at 6,000,000,000 feet. This includes the pine from the Saint Mary's river westward to the Wisconsin line and the mouth of the Montreal river, and upon the south shore of lake Superior. It is divided as follows:

"1. On the Menominee river and tributaries, 1,600,000,000 feet.

"2. In the western portion of the peninsula, not including the Menominee and tributaries, but including all west of the line of the Chicago and Northwestern railway between Escanaba and Marquette, 2,400,000,000 feet.

"3. East of the line of the Chicago and Northwestern railway, 2,000,000,000 feet.

"The largest bodies of pine in the northern peninsula are in the counties of Chippewa, Mackinac, Schoolcraft, Marquette, Houghton, and Ontonagon. There is also quite a large body in Keweenaw county, covering perhaps 30,000 acres. Ontonagon county, which extends along the south shore of lake Superior for nearly 100 miles, for 35 miles back from the lake is mostly covered with hard-wood timber, with a little pine along the streams, but not in sufficient quantities to estimate. This is also true of the northern part of Baraga and Marquette counties, extending along the southern shore of lake Superior, a distance of 125 miles from L'Anse to Onota, in Schoolcraft county. There are here a few small bodies of pine scattered through the hard wood, but it is needed by the settlers, and has no export value. The quality of the timber upon the Ontonagon and Presque Isle rivers and the upper Menominee, growing among the hard woods along the south slope of the Penoque iron range, is similar to that on the western shore of the lower peninsula. This timber is, however, somewhat difficult of access. The streams over which it must be driven (the Ontonagon and Presque Isle) are rough, broken, and require considerable improvement. The pine east of the line of the Chicago and Northwestern railway between Marquette and Escanaba, on the east half of the northern peninsula, is of poor quality, and may be classed as 'sapling pine', with occasional groves of what is called 'big sapling' scattered through the hard woods.

"In the upper peninsula of Michigan, according to the Lake Superior Canal Company's reports of examination and estimates of cord-wood, there is an average of about eighteen cords of wood per acre over the whole area of the peninsula, of which two-thirds is hard wood and one-third soft wood.

"In Menominee and Delta counties, the southern part of Schoolcraft county, and the extreme southern part of Marquette county are quite large quantities of tamarack and yellow cedar. From most of these lands the merchantable pine has been removed, and where the fires have not destroyed the cedar and tamarack the railroad companies are cutting the timber and shipping it to the prairies for telegraph poles, ties, and posts. It is stated by the owners of the lands, who long since cut the pine from them, that the cedar and tamarack trees left upon the land have netted them more than the original pine harvested. What makes this timber so valuable is its close proximity to the railroads and the ease with which it can be shipped by rail or over the waters of Green bay. This shows the necessity of preserving this kind of timber for future use, and of not abandoning it for taxes, as has heretofore been done, or allowing it to be destroyed by fires and windfalls.

"There are on the Menominee river some 9,000,000,000 feet of standing pine, one of the largest bodies left in the northwest. More than half of this, however, lies in the state of Wisconsin. About 200,000,000 feet of lumber are manufactured annually upon the Menominee. All the mills upon the river are located at its mouth, in the towns of Marinette and Menomonee, in Wisconsin, and it is considered next to impossible to build more mills at that point. The river is here narrow, and the facilities for holding logs, shipping lumber, dockage, etc., are quite limited in proportion to the amount of timber left in the region tributary to this stream; and this body of pine may therefore be considered to a certain extent in reserve, and likely to outlast many larger ones. There is little danger from fire on this river; the pine which is left grows upon the hard-wood ridges, interspersed with broad areas of swamp."

WISCONSIN.

The great prairies of the central Atlantic region once found their northeastern limits in southern Wisconsin. The forest covering of all the southern part of the state was confined to the bottom lands or open upland groves of stunted oaks of no great extent or of more than local importance. The central part of the state was covered with a dense forest of hard woods, oaks, ash, maple, cherry, birch, and the other trees of the northern forest, through which, upon gravelly or sandy ridges, great bodies of white pine were scattered. These pine forests gradually change in character and decrease in productiveness as they reach northward. Lakes are more common, and swamps of tamarack, cedar, and spruce occupy in the northern part of the state a considerable proportion of the forest area. The pine trees in these northern forests are smaller and more scattered than those farther south, although generally less intermixed with hard woods, and affording lumber of poorer quality.

The forest area has somewhat increased in the prairie region of the state since its first settlement and the consequent decrease of destructive prairie fires. The growth of trees has gradually spread from the bottom lands of the streams to the hills, and the oak forests upon the uplands have gradually encroached upon the prairie, losing their open, park-like character by the appearance of a young growth which has sprung up among the old trees.

The pine has been destroyed along the entire southern borders of the pine belt, along the banks of the principal streams, and from the lines of railroad, while the hard wood has been often greatly injured or destroyed by fire in those parts of the state where pine has been cut. The amount of pine still growing in Wisconsin is nevertheless large, although it should not be forgotten that the best and most easily accessible has already been harvested. What remains is generally remote from actual lines of transportation, and often, especially in the extreme northern part of the state, of comparatively poor quality.


During the census year 406,298 acres of woodland were reported destroyed by fire, with an estimated loss of \$725,610. The largest number of these fires was set by farmers in clearing land, or by sparks from locomotives.

The manufacturers of cooperage stock report a general deterioration and scarcity of the best varieties of hard woods, and the substitution of beech, elm, and other woods for oak.

The following estimates of the amount of timber standing in Wisconsin May 31, 1880, were prepared by Mr. H. C. Putnam:

WHITE PINE (*Pinus Strobus*).

Regions.	Feet, board measure.
Basin of Saint Croix river and tributaries	2,500,000,000
Basin of Chippewa river and tributaries	15,000,000,000
Basin of Black river and tributaries	900,000,000
Basin of Wisconsin river and tributaries	10,000,000,000
Basin of Wolf river and tributaries	600,000,000
Basin of Oconto river and tributaries	500,000,000
Basin of Peshtigo river and tributaries	1,500,000,000
Basin of Menomonie river and tributaries (in Wisconsin)	6,400,000,000
Shore of lake Superior	3,600,000,000
Total	41,000,000,000
Cut for census year ending May 31, 1880 (including 1,007,039,000 shingles and 348,301,000 laths).	2,097,299,000





Of this amount 485,552,000 feet were manufactured along the Mississippi river in Illinois, Iowa, and Missouri as far south as Saint Louis.

The wooded region in Crawford, Richland, Sauk, and Vernon counties is estimated to contain 12,000,000 cords of hard wood in addition to some timber of commercial value. The cut for the census year ending May 31, 1880 (exclusive of 86,545,000 staves and 7,498,000 sets of headings), was 117,041,000 feet.

Valuable oak timber exists in large quantities in Dunn, Pierce, and Saint Croix counties.

The cedar swamps scattered through the pine belt of the state cover an area of some 1,365,000 acres, and are estimated to contain 62,800,000 posts, telegraph poles, and railroad ties, in addition to large quantities of tamarack and spruce.

Wisconsin is the third state in the importance of its lumber-manufacturing interests. The great centers of manufacture are in the neighborhood of Eau Claire upon the Chippewa river, upon the Wisconsin river, and upon the shores of Green bay and lake Superior. Logs cut in the forests of Wisconsin supply also mills located on the Mississippi river in Illinois, Iowa, and Missouri with material equivalent to nearly 500,000,000 feet of lumber.

The following is extracted from Mr. Putnam's report upon the forests of Wisconsin:

"The southern boundary of the forest coincides with a line extending northwesterly from near the city of Milwaukee on lake Michigan, to the falls of Saint Croix on the Saint Croix river, and the western boundary of the state. This includes the heavy hard-wood as well as the pine forest. There is also, or there has been, a large amount of hard-wood timber in the southeastern part of the state, south of this line, but as no large bodies of forest of commercial value are now standing there, it will not be considered here. Large bodies of hard-wood timber exist in Vernon, Crawford, Richland, and Sauk counties, covering in the aggregate fully 400,000 acres and containing at least 12,000,000 cords of wood. This region, however, is already thickly settled, and the forests are being rapidly cleared for agricultural purposes. No estimate has ever been made of the amount of pine timber standing in Wisconsin at the time of its original settlement; at the present time it is estimated that 41,000,000,000 feet of merchantable pine remain in the state, situated as follows, river basins being taken as the natural divisions of these pineries:

"1. North of the Saint Croix river and upon the lands tributary to that stream there are 2,500,000,000 feet, covering 2,000,000 acres.

"2. On the southern shore of lake Superior, including all the waters tributary to the lake in the state of Wisconsin, extending from the Montreal river on the Michigan line westward to the western boundary of the state, and embracing the Wisconsin pine on the Montreal river and upon the Bad, White, Bois-Brulé, Black, and Left-Hand rivers, 3,600,000,000 feet, covering 1,800,000 acres.

"3. On the Chippewa river and its principal tributaries, the Red Cedar, West Branch, East Branch, Flambeau, Jump, Yellow, and Eau Claire, covering an area of some 6,253,000 acres, with an estimated stand of pine of 15,000,000,000 feet.

"4. In the Black River basin, with an area of 1,000,000 acres, containing an estimated stand of 900,000,000 feet.

"5. In the Wisconsin River basin, with an area of 4,500,000 acres, with an estimated stand of 10,000,000 feet.

"The remainder of the state, lying east of the east line of the Wisconsin River division and north of the southern boundary of the original forest, is divided by rivers as follows: (1) Wolf river, with 600,000,000 feet of pine; (2) the Oconto river, with 500,000,000 feet of pine; (3) the Peshtigo river, with 1,500,000,000 feet; (4) the Menomonee in Wisconsin, 6,400,000,000; making a total in the division east of the Wisconsin of some 7,000,000 acres, with an estimated stand of 9,000,000,000 feet of pine. This makes a grand total of pine forest area in the state of nearly 23,000,000 acres, still containing 41,000,000,000 feet of standing pine. This includes about 200,000,000 feet upon the Menomonee Indian reservation, in the county of Shawano, where very little pine has ever been cut; 100,000,000 feet on the Flambeau reservation, and 200,000,000 feet upon the Court Oreilles reservation. There is no merchantable pine standing on any of the other Indian reservations in the state.

"The quality of the pine in the state of Wisconsin varies largely with the differences in soil. The quality of the pine growing mixed with hard woods upon the southern boundary of the forest and bordering on the prairies was similar to that of the best Michigan pine. This is especially true of timber cut on the Wolf, Oconto, and Peshtigo rivers. The timber originally on the Wolf and Oconto rivers was especially fine. This has been largely cut, although there are still some very fine bodies of the best pine left on the Oconto and the western branch of the Peshtigo and northern branch of the Wolf rivers. The Black River district contained also a large amount of the best upper quality of pine, of which, however, more than half has been cut. The Eau de Galle River basin, in the counties of Pierce, Dunn, and Saint Croix, also contained at one time a large amount of the upper grade of pine, now, however, all removed. This grew among hard-wood timber, on good soil, which, when the timber is cut off, is valuable for farming purposes. The pine in this part of the state did not grow in extensive tracts. It was scattered through the hard-wood timber, from 1 to 10 large pine trees growing on an acre—trees which would scale from 1,000 to 3,500 feet of lumber each. There are still small bodies of this large pine left, but the great bulk of it is gone.

"As we go north from the southern boundary of the forest we enter a lighter soil, where cedar and tamarack swamps are interspersed between the hard-wood ridges. Many of these swamps are natural peat-bogs, covered with cedar, tamarack, and spruce. The tree growth upon them is heaviest near the outer edges, the centers

often being covered with grass or cranberry plants. These swamps, originally the beds of lakes, are now filling up and becoming gradually covered with timber. On the Wolf river the timber was very heavy. Instances are known of 10,000,000 or 12,000,000 feet of pine lumber having been cut from one section of 640 acres in the Lower Wolf River region.

"In the pine forest, away from the large bodies of mixed hard wood and pine previously described, the general character of the timber is about the same, varying somewhat in different localities, but still possessing the same general characteristics and qualities. Where the pine grows in large solid bodies there are many young trees mixed with the older, and the timber is generally of inferior or lower grade. This is true of pine growing about the head of the Flambeau and Wisconsin rivers, and the Menomonee river in Wisconsin. Large pine cannot grow and mature upon very poor soil, and where the soil is poor the trees, after reaching a certain size or age, decay and are thrown down by wind or destroyed by fire. The white pine in Wisconsin does not mature except upon the rich gravelly loam of the ridges.

"The principal points of lumber manufacture at present in Wisconsin are on lake Winnebago, at the cities of Oshkosh and Menasha, which take largely the product of the Wolf and Fox River pineries; at Green Bay and Oconto, which derive their logs principally from the Oconto river; at Peshtigo, on the Peshtigo river; at Marinette, on the Menomonee river; on the Wisconsin river, at Grand Rapids, Stevens Point, Mosinee, Wausan, and Jenny, the terminus of the Wisconsin Valley railroad, and at Necedah, on the Yellow river. Along the Wisconsin Central railroad, from Junction City to Ashland, are mills of more or less capacity at every station, the most important being at Ogemaw, Ashland, Medford, and Unity. Upon the Black river the principal manufacturing points are La Crosse and Black River Falls. On the Chicago, Saint Paul, Minneapolis, and Omaha railway, at Fairchild, are the large mills of Foster & Co., who are engaged in manufacturing the timber lying between the Black river and the waters of the Chippewa, included in the Chippewa estimate. On the Chippewa river the largest manufacturing establishment is the Mississippi River Logging Company, composed of fifteen of the heaviest concerns upon the Mississippi river. These firms obtain their stock mostly from the Chippewa river, the logs being driven down to its mouth into what is called the 'Beef Slough boom', where they are separated and formed into rafts and towed to the different mills below. This company cuts on the Chippewa about 400,000,000 feet a year. The principal manufacturing points on the Chippewa deriving their logs from its basin are situated at Waubeck, Dunnville, Menomonee, Meridian, and Eau Claire, where several large and important manufacturing establishments are located. Higher up the river the Badger State Lumber Company and the Grand Island Lumber Company are located, and at Chippewa Falls, the county-seat of Chippewa county, the Chippewa Lumber and Boom Company has a large water-mill, with a capacity of 65,000,000 feet a year, besides several smaller concerns. The railroad extending from Chippewa Falls eastward through Chippewa and Clark counties into Marathon county, and joining the Wisconsin Central railroad at Abbottsford, passes through a hard-wood country. Several firms are already established upon this line and have commenced the manufacture of staves and the production of hard-wood lumber for wagons, etc., and are developing a large business. This road runs through one of the finest bodies of hard wood in the state, containing large amounts of oak and maple growing on a fine soil suitable for farming. The Chippewa River country now contains the largest body of white pine of the best quality left in the states of Michigan, Wisconsin, and Minnesota. It is, however, being very rapidly cut.

"It is found in going north toward the heads of the streams that the timber stands more in large groves, and that there is less hard-wood timber mixed with the pine. When the loggers attack these forests they cut clean as they go, the timber being of more uniform size and age, and there being less undergrowth than farther down the streams. It is found, also, that the pineries on the heads of the streams do not hold out as well or yield as large an amount of timber as those farther south, where the forests border on the prairie lands and where the pines grow on better soil. This is true both of the Wisconsin and of the Michigan pineries. The poorer soils in the northern portion of the state do not grow and mature the large sapling forests of pine found in the southern portions of the pine belt. So that, while there is still a large area which has not been cut and which may appear inexhaustible, yet, owing to the fact that the timber lies more in groves, and that there are here wide extents of tamarack and cedar, swamps and open spaces, the ground will be cut over more rapidly than when the forest was first entered. This is true of the pine standing upon all the streams of northern Wisconsin in the Menomonee district—the Wisconsin, the Chippewa, Saint Croix—and on the southern shores of lake Superior. Commencing at Menomonee, on the Chicago, Saint Paul, Minneapolis, and Omaha railway, and running west through the 30 miles of 'big woods', large mills for the manufacture of hard-wood timber and of what little scattered pine there is left are established at Knapp, Wilson, Hersey, Woodville, and Baldwin stations. The principal manufactories in the Saint Croix district are at Hudson, on the Willow river, and at Stillwater, in Minnesota, which receives its logs from the Saint Croix, in Wisconsin, and which, therefore, should be treated as one of the Wisconsin pinery manufactories. At Somerset, on Apple river, there is one mill; there is one at Osceola, upon the Saint Croix, and upon the Northern Wisconsin railroad, which runs through the Saint Croix division; at Clayton, Granite Lake, and Shell Lake are large mills. There are also other mills along this road on the Lake Superior shore. There are mills of small capacity at Superior City, Bayfield, and Ashland; the latter receive their logs by the Wisconsin Central railroad from the Bad River pinery.

"On the Eau Claire river the timber is small and sound, growing very thick and long; there are frequent instances where 1,200,000 or even 1,500,000 feet of lumber have been cut upon a 40-acre lot. One tree was cut on Jump river some years ago which scaled 7,000 feet of lumber. The general character of this timber, especially upon the main Chippewa or West Branch and a portion of the Flambeau, is called 'big sapling pine'. Of the true cork pine very little is found in the northern part of Wisconsin, probably because the soil is not strong enough to permit its full development. The general character of the timber upon the Wisconsin river is very much the same as that upon the main Chippewa. There are instances of very fine pine having been cut in the hard-wood forest upon the lower part of the river, and some fine groves are found even as far north as the Tomahawk and East Branch. The Flambeau river, or East Branch of the Chippewa, has also, in ranges 2 and 3 east, extending from townships 35 to 41 north, inclusive, some excellent bodies of upper-grade pine.

"On the Jump river are some fine bodies of pine, nearly approaching in quality Michigan cork pine and running largely to 'uppers'. This is true also of the pine upon the Yellow river, where the timber grows largely scattered among hard woods and is of fine quality. One of the finest bodies of pine in Wisconsin is that which belongs to Cornell university, lying in townships 33 to 38, ranges 8 and 9, in the highest part of Chippewa county, on the divide between the Chippewa and Red Cedar rivers. On this body frequent estimates of 1,000,000 feet to 40 acres have been made. On the Saint Croix river are many barren areas timbered with scrub pine, patches of Norway pine, and small black and white oak. These barrens cover about 700,000 acres of the Saint Croix region. The soil is sandy, and fires run over the country every year. South of these barrens, in Polk, Barren, Saint Croix, Dunn, and Pierce counties, is a tract of very valuable hard-wood land, upon which the greatest portion of the timber is now standing, although settlements are already largely scattered through this region. This body of hard wood contains a large amount of valuable white-oak stave timber and much timber suitable for general manufacturing purposes. It is being, however, rapidly destroyed by settlers and by the fires incident to agricultural and logging operations.

"In Clark county, which lies partly in the Chippewa and partly in the Black River region, are large bodies of hard-wood timber as yet uncut and growing upon land valuable for farming purposes. This growth extends as far north as the northern line of the county. The same body of timber extends east through Marathon and Wood counties, and is particularly fine in the western portions. The same body of hard-wood timber continues east toward lake Michigan, including the counties of Portage, Waupaca, Shawano, Outagamie, Winnebago, Brown, Kewaunee, Manitowoc, Calumet, Fond du Lac, Sheboygan, and Ozaukee. Large tracts in these counties are, of course, cleared and settled; still they contain large bodies of unoccupied hard-wood timber, and the opportunities for cheap farms are plenty.

"Of the forest region proper of Wisconsin, fully 5 per cent. is not covered with timber; this includes swamps, lakes, rivers, bottoms, etc. In the extreme southern part of the forest area, over a region from 35 to 50 miles in width, the hard wood predominates, only about one-fifth of the forest growth being pine. North of this hard-wood region proper, perhaps one-half of the forest growth is pine and other soft woods and the rest hard woods. Hemlock is scattered through the pine forest outside of the heavy hard-wood areas. A careful estimate of the hemlock timber now standing gives the following results, the divisions agreeing with those used in estimating the standing pine: On the Chippewa river, upon 3,000,000 acres, 2,500,000,000 feet of hemlock; on the Saint Croix river, upon 1,000,000 acres, 500,000,000 feet of hemlock; on the Black river, upon 350,000 acres, 100,000,000 feet of hemlock; in the country east of the Wisconsin River division, and including the Wolf, Oconto, Peshtigo, and Menomonee rivers and their tributaries, upon 3,000,000 acres, 1,500,000,000 feet.

"The total area in the state on which hemlock timber grows is about 10,500,000 acres, containing, roughly, 5,500,000,000 feet. The quality of the hemlock timber in Wisconsin is not so good as that grown in New York and northern Pennsylvania, although it is valuable for its bark, and the timber when peeled can be driven down with the pine and sawed at the mills into dimension stuff for use where coarse lumber is required.

"Generally, therefore, the forests of Wisconsin may be divided into the hard-wood lands already described, along the southern borders, from which the pine has been mostly cut; north of this, and extending northward somewhat indefinitely, the mixed growth of hard wood and pine, growing upon soil adapted for agricultural purposes. The open meadows in this region are covered either with grass or cranberry marshes, alike valuable to the lumber and farming interests. About the head of the Flambeau river are large open spaces running into groves of heavy pine timber. These open spaces, once lakes or swamps, are drying up and the timber is gradually spreading over them. There are bodies of timber scattered through the southern portions of the state outside of the original forest area, but the amount of this timber is relatively so small that it cannot be considered of commercial importance, and hardly supplies the wants of the population occupying the thickly-settled southern counties.

"Five thousand men are employed in the pineries of the Chippewa river. They are expected to cut during the logging season about 600,000,000 feet of logs, or an average per man of over 100,000 feet. This rule is not applicable to the northwestern pineries generally, for in Michigan, as the timber is now farther from the streams, the average cut per man is not as great, and 80,000 feet per man would perhaps be a fair average, taking the pineries of the whole northwest.

"The annual increase or growth of timber is counterbalanced by the annual waste by windfalls and the natural decay of the old trees. The loss to the forest by fire is an unknown quantity, but it is quite a large amount, probably 5 per cent. of the whole. The lumbermen waste the log which runs into the top of the tree; this is knotty, but usually sound, and would make good merchantable lumber. It is left in the woods, however, because there is a good deal of work in trimming the knots and cutting off the limbs. From an ordinary-sized tree four 16-foot logs are usually taken, the rest being left. Often this top log is 22 inches in diameter at the butt and will scale from 100 to 120 feet. Loggers are paid so much per thousand feet by the lumberman, and the amount they receive is so small that they cannot afford to spend the time to finish up and take out the fifth or last log, which is therefore left in the woods and lost. Nearly one-tenth of the timber, therefore, is left in the woods and lost. The fires about the old choppings, or where lumber operations are going on, are principally caused by the carelessness of woodsmen in hunting up land-lines, or of driving-crews on the river in the spring who leave their fires, or by explorers in the forest during the month of May or June leaving their camp fires burning. In all the old cuttings the dried pine boughs and other timber left on the ground get very dry, and fire once started burns with great rapidity and violence.

"As a matter of fact, more than half the area from which pine forests have been cut in the northwest is sooner or later burned over. The fire destroys the young trees and changes the nature of the surface of the ground, so that the next crop which comes up consists of briars and poplars, and then hard woods. When pine is cut off or burned it does not come in again, and I have never seen any old choppings of pine come up with pine again, even when some trees were left and the ground had not been burned, although where a few large trees only are removed from a pine forest growing on good soil the small trees left standing, if protected from fire, will continue to grow."

MINNESOTA.

The Northern Pine Belt finds in Minnesota its extreme western limit in the United States in longitude $95^{\circ} 30'$, and its southwestern limit near the forty-sixth degree of latitude. Along its southern and western borders a narrow territory covered with an open growth of hard wood separates the forests of pine from the prairie, which occupies all the southern and western portions of the state.

The same general features which characterize the pine belt of Wisconsin extend into Minnesota. The pine in the southern portion, confined to gravelly ridges, is scattered through forests of hard wood. Farther north the forest changes in character, the pine being small and of inferior quality. Broad areas of barren land covered with stunted birch, gray pine, and scrub oak occur, while the whole country is thickly studded with lakes and with tamarack and cedar swamps. North of the Mississippi River divide the country is more open; the forest is stunted and of little value, and pine is only found in small, scattered clumps mixed with spruce, tamarack, and yellow cedar. The forest growth here occupies perhaps two-thirds of the rocky or swampy surface of the ground. Its productive capacity is not large, and the northern part of the state is not adapted to lumbering operations.

The pine has been removed from the principal streams of the state, and that which remains, except in the region tributary to lake Superior and in the vicinity of Red lake, is now inaccessible or of comparatively inferior quality. The best hard-wood forests of the state, as in Michigan and Wisconsin, have suffered seriously by fires started in abandoned pineries, or in clearing land for agriculture.

During the census year 250,805 acres of woodland were reported devastated by fire, with an estimated loss of \$1,395,110. The largest number of these fires was set in clearing land or by sparks from locomotives.

The manufacture of cooperage stock to supply the large flouring-mills of the state is an important industry. Manufacturers report a growing scarcity and general deterioration of material. Basswood, elm, and ash are largely used; oak is inferior in quality to that grown farther east and south.

The following estimates of the amount of pine timber standing in Minnesota May 31, 1880, were prepared by Mr. H. C. Putnam:

WHITE PINE (*Pinus Strobus*).

Regions.	Feet, board measure.
Mississippi river and tributaries.....	2,900,000,000
Rainy lake and Rainy Lake river.....	300,000,000
Red Lake river and other tributaries of the Red river.....	600,000,000
Saint Louis river and tributaries.....	3,500,000,000
Shore of lake Superior.....	870,000,000
Total.....	8,170,000,000
Cut for the census year, ending May 31, 1880 (including 187,836,000 shingles and 88,088,000 laths).	540,997,000


In the belt of hard wood extending west and south of the pine region, and consisting of white, red, and burr oak, sugar-maple, poplar, etc., it is estimated that 3,840,000 acres of forest remain, capable of yielding an average

MAP OF MINNESOTA.

SHOWING THE DISTRIBUTION OF FORESTS,
WITH SPECIAL REFERENCE TO THE
LUMBER INDUSTRY.

COMPILED UNDER THE DIRECTION OF
C. S. SARGENT, SPECIAL AGENT.

1881.



Scale

Julius L. Smith

of 15 cords of wood to the acre, or 57,600,000 cords. The cut for the census year ending May 31, 1880 (exclusive of 7,825,000 staves and 547,000 sets of headings), was 36,884,000 feet.

Minnesota is the eighth state in the importance of its lumber-manufacturing interests. The principal centers of manufacture are Minneapolis, upon the Mississippi river, the Saint Croix river in Washington county, the Mississippi river in Anoka county, and Duluth, near the mouth of the Saint Louis river.

The following is extracted from Mr. Putnam's report upon the forests of Minnesota:

"The great hard-wood forest of Minnesota lies to the south and west of the pine forest, extending north and northwest from Freeborn and Mower counties on the southeast into Marshall county, and to within 50 or 60 miles of the boundary-line between Canada and the United States. This body of hard wood, which is some 300 miles long by about 20 miles wide, borders upon the prairie, and is the extreme western body of timber of any commercial value east of the Rocky mountains. This forest covers about 3,840,000 acres of land generally valuable for agricultural purposes, besides its timber, which will average about 15 cords to the acre. The surface of the land is level or gently undulating, well watered, particularly the so-called 'park region' which lies in Becker, Otter Tail, Douglas, Stearns, and Todd counties, and in fact extends through Wright, Hennepin, Carver, Le Sueur, Rice, and Steele counties.

"North and east of this belt of hard wood the pine forests commence at a point where the southern line of the Wisconsin forest crosses the Saint Croix river, near Taylor's Falls. They extend northwesterly through the counties of Chisago, Isanti, Mille Lacs, Benton, Morrison, Todd, Otter Tail, Becker, Polk, and Beltrami, nearly parallel to the line of the hard-wood forest, and, crossing Red Lake river, extend round to the north of Red lake, and thence easterly, reaching the shore of lake Superior at the Grand Portage.

"The general character of the pine in Minnesota is similar to that of northern Wisconsin, although it contains more sapling pine and a smaller percentage of 'uppers.' It is generally somewhat scattering and in smaller groves. Large areas of barren land within the forest proper are covered with birch, through which are scattered patches of small pine, while large areas of swamp bear only tamarack and cedar. The pine of Minnesota is estimated as follows:

"1. On the portion tributary to the Rainy lake and Rainy Lake river, including the Big Fork, the Little Fork, and the Vermillion rivers, 300,000,000 feet. This stands upon streams which flow northward. This pine will naturally be sent to Manitoba.

"2. On the northern shore of lake Superior, east of Duluth, and covering the waters tributary to lake Superior, of which very little is surveyed and no area is given, 870,000,000 feet.

"3. On the waters of the Saint Louis, including the Cloquet, White Face, and other small streams, 3,500,000,000 feet.

"4. On Red Lake river and its tributaries. The great body of pine in this division is principally upon Red lake and Red Lake river. It is estimated to contain 600,000,000 feet, although it is nearly all unsurveyed.

"5. On the Mississippi river and tributaries above Minneapolis, 2,900,000,000 feet.

"About one-half of the pine has been cut in Carlton county; it has all been cut in Pine county with the exception of that growing in a few townships. It has nearly all been cut in Chisago, Kanabec, Morrison, and Crow Wing counties. A great deal of pine, too, has been cut in Cass county, while Todd, Otter Tail, and Wadena have all been cut over. The principal bodies of pine now remaining are located in Cook, Lake, Saint Louis, Cass, Itasca, and Beltrami counties. There were a few thousand acres growing on the Roseau river, where it runs into northwestern Minnesota, but this has all been cut by the Canadians. There is no hemlock or spruce in Minnesota. There are occasional ridges of hard wood within the pine forest, as in Wisconsin and Michigan.

"A large portion of the northern part of the state is as yet unsurveyed and but little known, except that, in the region extending from 30 to 100 miles south from the international line, there is little pine of commercial value. It is an open country, full of bogs, swamps, rocks, and wide areas of worthless land; this region extends from the Arrow river clear through to the international line, south and west of the Lake of the Woods, and to the Vermillion lake.

"Along the line of the Northern Pacific railroad and north and east of the Mille Lacs country are large swamps covered with tamarack timber of commercial value. Through this country are many marshy lakes containing floating islands, lands in process of formation by the accumulation of vegetation. The timber in this district is growing and increasing, and if fires can be kept out of the tamarack and cedar timber the small pine will grow rapidly.

"The timber which grew on the Saint Croix river in Minnesota was tributary to Stillwater, and has all been cut and manufactured there.

"The principal manufactories of pine on the Mississippi river are at Minneapolis.

"FORESTS ON INDIAN RESERVATIONS.

"Referring to the Red Lake Indian reservation in Minnesota, and other Indian reservations on which the pine remains uncut, amounting in the aggregate to 1,000,000,000 feet, it may be said that they are nearly all unsurveyed, and are generally covered with a heavy pine forest, and that the lands are unfit for agricultural purposes and only

valuable for the pine timber which grows upon them. These reservations should be held as long as possible by the government as a timber reserve. They should not be surveyed and subdivided except so far as may be necessary for their protection, and they should not be offered for sale until some necessity, now unforeseen, arises for their disposal. The 1,000,000,000 feet of pine should be held until the amount for which it can be sold is needed by the Indians, or until a price near its value can be obtained for it. By selling the land now the value of the timber cannot be realized, while the interest of the settlers who may hereafter enter upon the prairies would seem to demand that some reservation of pine should be made for them, if possible. The proposition to bring these lands into market, subject to pre-emption and homestead entry, is against the interest of every one except the few worthless tramps and irresponsible persons who may seek to enter and procure a title to these lands; and even if the land was so open to homestead and pre-emption entry, the aim and purpose of these laws could not be carried out, for no farms will be made nor homesteads improved in this Indian country.

"The White Earth Indian reservation is largely covered with hard wood, there being no pine upon more than a quarter of its area. The land is desirable for agricultural purposes, and may be utilized for the settlement of Indians, or under the homestead and pre-emption laws by whites, but the pine lands are unfit for cultivation, and the homesteading or pre-empting of them should not be allowed."

IOWA.

Iowa lies within the prairie region. The broad bottom lands along the river of the eastern part of the state once bore heavy forests of broad-leaved trees. Farther west the tree growth was less heavy in the narrower bottoms. All over the state, however, forests lined the streams and often spread, especially in the southwestern counties, over the uplands. Since the first settlement of the state the forest area has increased by the natural spread of trees over ground protected from fire, and by considerable plantations of cottonwood, maple, and other trees of rapid growth made by farmers to supply fuel and shelter.

The natural forests have been everywhere largely culled of their most valuable timber, and in spite of their increased acreage are, in their commercial aspect, in danger of speedy extermination. Manufacturers of cooperage stock and others using Iowa timber report great scarcity and general deterioration of stock.

During the census year 11,017 acres of woodland were reported destroyed by fire, with an estimated loss of \$45,470. These fires were largely the result of carelessness in clearing land.

Iowa is the ninth state in the importance of its lumber-manufacturing interests. It owes its position to numerous large mills situated along the Mississippi river entirely supplied with logs from the pineries of Wisconsin. The amount of Iowa-grown lumber manufactured is insignificant.

MISSOURI.

Southern and southwestern Missouri was originally covered with a dense forest of hard woods, through which in the southern counties extensive areas of the short-leaved pine (*Pinus mitis*), covering gravelly ridges and the low Ozark hills, were common. The northern and western limits of the true forest region may be defined by a line entering the state from the southwest, in the southern part of Jasper county, and passing northeasterly through Dade, Cedar, Saint Clair, Henry, Benton, Morgan, and Cooper counties, and then northward to the borders of the state. West of this line the timber is largely confined to the broad bottom lands, in belts often 2 or 3 miles in width. Farther west these become narrower and less heavily timbered. The extreme northwestern counties, Atchison and Nodaway, are almost destitute of timber.

The forests of southeastern Missouri still contain great stores of valuable timber, although the best trees have been cut in the neighborhood of all settlements, and for a distance varying from 5 to 20 miles back from all lines of railroad. This is especially true of the best white oak and of black walnut, once common, but now almost exterminated in all parts of the state.

Manufacturers of cooperage stock report a growing scarcity of material everywhere, and are now forced to obtain oak from Arkansas and elm and basswood from the rivers of southern Illinois and Indiana. The further development, however, of the railroad system of southern Missouri will make available for manufacturing purposes a large amount of valuable timber now remote from transportation.

During the census year 783,646 acres of woodland were reported destroyed by fire, with an estimated loss of \$294,865. These fires were traced to careless hunters, to fires set in clearing farming land, to sparks from locomotives, etc.

A gratifying improvement in the condition of the forest in the parts of the state first settled has followed the enactment of a fence law preventing the general ranging of stock through the timber-land. A young growth has sprung up among the older trees and along the borders of woodlands protected from browsing animals, and these young forests are valuable in their prospective yield and as an indication of the methods which must be adopted to preserve and perpetuate the forests of the whole Atlantic region.



LEGEND

- I Under 10 cord per acre
- II From 10 to 20
- III 25
- IV 50
- Over 100

DENSITY OF FORESTS

COMPILED UNDER THE DIRECTION OF
C.S. SARGENT, SPECIAL AGENT
1883.

Scale 0 20 40 60 80 100 120 STAT. M. E.

DAKOTA

Julius F. & Co. Ltd.

Missouri is the tenth state in the importance of its lumber-manufacturing interests. It owes its position in part to large mills located upon the Mississippi river manufacturing logs cut in the forests of Wisconsin. A much larger amount of lumber, however, in the aggregate, both pine and hard wood, is produced in numerous small railroad mills located along the line of the Iron Mountain and other railroads running through the southern part of the state.

Saint Louis is an important center of lumber distribution. It receives a large portion of the Wisconsin pine crop by raft, Michigan pine by rail, and southern pine and hard woods by rail and river.

DAKOTA.

Dakota, with the exception of its river lands and the small territory between the north and south forks of the Cheyenne river, is practically destitute of timber. The bottoms of the principal streams contain extensive groves of hard wood. As far west as the James river timber exists about the shores of the larger lakes, and upon the Low Turtle and Pembina mountains of the northern boundary, occasionally ascending the *côtes* or sides of low tables rising from the prairie. The Black hills, an extreme outpost of the Rocky Mountain system, were once heavily timbered. The yellow pine of the Pacific region is here mingled with the white spruce, the canoe birch, the burr oak, and the elm of the eastern forests, while poplars of the Atlantic and Pacific regions grow side by side.

Much timber has already been cut along the eastern rivers to supply the wants of a rapidly-increasing agricultural population, and the isolated pine forests of the Black hills, separated by hundreds of miles from any equally large or valuable body of building timber, have already suffered serious inroads. The best and most accessible pine has been cut and manufactured into lumber or consumed as fuel in the silver mines and stamping mills to which this region owes its population, and much timber has been allowed to perish in the fires which of late years have often swept through these forests.

The principal center of lumber manufacture is Deadwood, in the Black hills, where a comparatively large amount of pine is sawed. In the eastern counties a little oak and elm is manufactured, for the most part in small portable mills.

The following extracts are made from Mr. H. C. Putnam's report upon the eastern portions of the territory:

"Along the whole length of the Missouri river in Dakota there is a belt of hard-wood timber in the bottoms in bodies of from 100 to 500 acres in extent. This timber sometimes grows continuously, but more often there are open spaces between the groves. About three-fourths of the trees are burr oak, the remainder sycamore, cottonwood, green ash, box-elder, poplar, willow, etc. A similar forest growth lines the banks of the Red river north of Fort Abercrombie as far as Fort Pembina, near the international line. This strip of timber averages perhaps forty rods in width, and consists of the same varieties of trees that grow upon the Missouri river.

"In the Pembina mountains and west of Fort Pembina, on the Tongue and Pembina rivers, there are bodies of timber, generally of stunted growth, lying mostly along the streams or about the Pembina mountains in groves of from 160 to 3,000 acres in extent. This timber is situated principally in the two northern tiers of townships of Pembina county. It has no value except as fuel. The next body of timber in Dakota is in the neighborhood of Devil lake; it aggregates some 25,000 acres, distributed as follows: At Wood lake, some 20 miles north of Devil lake, there are 1,000 acres; on Graham's island, a promontory on the north shore of Devil lake, near the northwest end, are 2,500 acres of timber; east of this, on the north shore of the lake, are two groves of about 500 acres; at Rock island, which is really a promontory running into the lake, are 3,800 acres of timber; around the east and north shores, and around the whole southern shore of the lake, past Fort Totten to the extreme west end, are some 15,000 acres of forest adjacent to Devil lake; at Stump lake, a lake some 15 miles in diameter on the north side of Devil lake, there are 1,400 acres of timber; and commencing some 10 miles south of Fort Totten, and extending down along Cheyenne river into township 146, range 56, in Traill, Foster, and Grand Forks counties, are about 10,000 acres of timber. The valley here is only 1 or 2 miles in width, and the timber is generally distributed through it. Probably seven-eighths of all this Devil Lake timber is burr oak; the remainder is sycamore, green ash, etc. This timber in many places grows large, sometimes 30 or 40 feet to the first limb, and is valuable for fuel, for the construction of log houses, and for general use by settlers in the absence of other and better material.

"In the Turtle mountains, in Bottineau and Rolette counties, and extending into the British possessions, is quite a large tract of timber, principally oak of short, scrubby growth, and only valuable as fire-wood. A body of timber from 1 mile to 5 miles in width extends for 150 miles along the Mouse river, in the counties of Bottineau, McHenry, Stevens, and Renville. This timber is composed of burr oak, box-elder, sycamore, green ash, etc., and is suitable for fire-wood, house-building, and rough construction."

Mr. Robert Douglas, of Waukegan, Illinois, contributes the following remarks upon the forests of the Black Hills region, of which he made a critical examination:

"From Fort Meade the stage road runs about 2 miles along the base of the hills, and then follows up through heavy timber, gaining an altitude of over 1,500 feet above the fort when within 2 miles of Deadwood; thence down a

steep grade of about a mile until the valley is reached, and then up the valley by an easy grade to Deadwood. Five days' driving through the hills from the base of the foot-hills to one of the highest peaks shows little variation in the species of forest trees. The yellow pine (*Pinus ponderosa*) is the only tree of much value in the hills, and composes nineteen-twentieths of the forest, generally covering the hills from base to summit. The trees are larger and stand closer together than in Colorado, and grow here, too, more rapidly than farther south, as is shown by the width of the annual rings of growth and the shoots upon the standing trees. This is the only tree used for lumber at the saw-mills, and no other is used in the mines. The white spruce (*Picea alba*) grows principally near the water-courses, and here the largest trees of that species are to be found. It is scattered, however, through the pines even within 50 feet of the summit of Terry's peak. It is condemned by both saw-mill proprietors and miners as lacking strength and being very knotty, which cannot be doubted, as it retains its lower branches with wonderful tenacity, even when growing closely and in dense shade. These two species comprise all the *Coniferae* in the Black hills, with the exception of a prostrate juniper and rare specimens of the red cedar. The burr oak is found in the valleys extending into the foot-hills and along the creeks for 40 or 50 miles into the plains. It is short, gnarly, and apparently of little value, although exceptional trees in the valley are of fair size. In the narrow valleys and along water-courses are found the common cottonwood, black willow, narrow-leaved cottonwood, green ash, white elm, box-elder, ironwood, canoe birch, and quaking aspen; in the hills canoe birch, mountain ash, hazel, choke cherry, and juneberry are found growing side by side with the snowberry and mahonia of the Pacific region. The little aspen and the canoe birch perform the same service the aspen does in Colorado, and cover the ground after the timber is burned off, thus making a shade in which the pine seedlings find protection from sun and wind, and finally repopulate the hills. On the banks of the numerous creeks intersecting the 'bad lands' and plains from the Missouri river to the Black hills, box-elder, white elm, green ash, black willow, cottonwood, choke cherry, wild plum, and buffalo-berry occur, but the canoe birch is not found below the foot-hills, where it grows along the creeks 6 or 8 inches in diameter, or as a low shrub upon the hillsides.

"The region occupied by the Black Hills forests is 80 miles in length north and south, and about 30 miles wide from east to west. Forest fires are not so frequent nor so disastrous as in Colorado, although the 'big burning' of 1865, near Custer's peak, is estimated to have extended over 400 square miles. The yellow pine is largely reproducing itself over the whole of this area, the trees being now 3 or 4 feet high. As far as my observation went, this reproduction of the yellow pine over the old 'deadenings' is almost universal through the hills, although rarely or never seen in Colorado, and even in northern Wisconsin and the Michigan peninsula scarcely a single young pine has appeared in the whole burned district of 1871.

"The timber is disappearing rapidly in the vicinity of Deadwood, Lead City, Terryville, and Central."

NEBRASKA.


The forest growth of Nebraska was once confined to the eastern part of the state; the broad bottom lands of the Missouri and the lower Platte rivers contained groves of large oak, walnut, ash, and box-elder of considerable extent. These, under favorable conditions, spread to the bluffs and uplands. Westward the tree growth gradually became more scanty and stunted, until, west of the one hundredth meridian, only the large streams were lined with a few small cottonwoods and willows.

The best trees have already been culled from the scanty forest growth of the state, and if the area of natural woodland has somewhat increased along its eastern borders since the settlement of the country and the diminution of prairie fires, these forests are, in their commercial aspect, of little importance. Many small plantations of cottonwood and other trees of rapid growth have been made in connection with farms in the eastern counties, and these in some cases already furnish much-needed shelter to buildings and crops, and supply domestic fuel.

The lumber-manufacturing interests of Nebraska are not important. Mills at Omaha, the principal manufacturing center, saw cottonwood and a little walnut and oak, hauled to them from the neighborhood of the city, and small portable mills at other points along the Missouri saw a little cottonwood and such logs as the country tributary to them can furnish. The product of all the Nebraska mills is consumed in supplying the local demand.

KANSAS.

The heavy forest of the Mississippi basin just reaches the extreme southeastern corner of Kansas, covering nearly one-third of Cherokee county. North of this, and occupying the remaining eastern border of the state, a prairie region varying in width from 30 to 100 miles is still heavily wooded with valuable timber along the streams, the forest growth occasionally extending and covering areas of upland. West of this region of mixed prairie and woodland the timber is confined to the banks of streams. It is often, east of the ninety-seventh meridian, of considerable size and value, occurring in sufficient quantity to supply the most pressing wants of the agricultural population of this part of the state. West of the ninety-seventh meridian the tree growth gradually diminishes in vigor. Trees are here confined to the immediate banks of the large streams, and are small and of little value. West of the ninety-ninth meridian a few small stunted willows and cottonwoods, scattered at wide intervals along the large streams, represent the only forest-growth of this arid region.





A large amount of lumber is manufactured in the eastern counties in proportion to the extent of their forest area; but much of the best timber of the state has been cut, and Kansas must soon depend, even more than at present, upon the forests of Arkansas and Louisiana for its lumber supply.

Considerable plantations made in the eastern and southeastern counties by railroads and in connection with farms promise abundant success. All attempts, however, to raise trees in the arid central and western parts of the state have resulted in failure.

During the census year 7,080 acres of woodland were reported destroyed by fire, with an estimated loss of \$14,700. The largest number of these fires originated upon the prairie.

WESTERN DIVISION.

MONTANA.

The forests of Montana are confined to the high mountain ranges which occupy the western part of the territory. They are dense and important upon the slopes of the Cœur d'Alêne and other high ranges. Farther east, along the eastern slopes of the Rocky mountains and their outlying eastern ranges, the Big Belt, the Little Belt, the Crazy, the Snow, and the Bear mountains, and the ranges south of the Yellowstone river, the Yellowstone, Shoshonee, and Big Horn mountains, the forests are more open, stunted, and generally confined to the highest slopes, the borders of streams, or the sides of cañons. A narrow fringe of cottonwood, green ash, and willow lines the bottoms of the Missouri, Yellowstone, Tongue, Rosebud, Milk, and of the other large streams of the territory; and a few stunted pines and cedars are scattered along the river bluffs and the highest ridges of the Powder River, the Wolf, and other ranges in the southeastern part of the territory. The remainder of the territory, the eastern, northern, and southern portions, are destitute of timber.

The heavy forests of northwestern Montana, largely composed of red fir, yellow pine, and tamarack, and containing great bodies of white pine (*Pinus monticola*) and considerable valuable spruce (*Picea Engelmanni* and *P. alba*) constitute, with those covering the adjacent mountains of Idaho, one of the most important bodies of timber in the United States. East and west of this forest a treeless country, adapted to grazing and agriculture, and destined to support a large population which must obtain its building material and railroad supplies from it, extends over thousands of square miles. The development, too, of the important mining interests of southern Montana and Idaho is dependent upon these forests, their only valuable source of timber and fuel supply. These forests guard the headwaters of two of the great rivers of the continent, and in regulating their flow make possible through irrigation the devotion to profitable agriculture of a vast territory now an almost arid waste. The forests, largely composed of the lodge-pole pine (*Pinus Murrayana*), which cover the outlying eastern ranges of the Rocky mountains at an elevation of from 5,000 to 10,000 feet above the ocean level with a dense growth of slender trees or on poor soil and in exposed situations with an open, scattered forest, are, as sources of lumber supply, of comparatively little value. These forests, however, contain valuable supplies of fuel and abundant material for railway ties. They guard, too, the flow of numberless small streams, and their importance in this connection should not be overlooked.


The most important forests, commercially, of the territory are found along the valley of Clarke's Fork of the Columbia river, between the Horse Plains and the Idaho line; here the western white pine reaches its greatest development, becoming an important part of the forest growth. The valleys of the Saint Regis de Borgia and Missoula rivers contain great bodies of valuable fir and pine, which spread also in great luxuriance over the mountains east and south of Flathead lake.

Fires destroy every year large areas of the forest covering the mountains of the western division. The long, dry summers and the character of the forest, composed as it is almost entirely of coniferous resinous trees, favor the spread of forest fires. They increase rapidly in number with the increase of population, and threaten the entire extermination of the forests of the whole interior Pacific region. During the census year 88,020 acres of forest were reported destroyed by fire, with a loss of \$1,128,000. These fires, few in number, were traced for the most part to careless hunters, prospectors, and smokers.

Little lumber is manufactured in the territory. Red fir and spruce are sawed at Missoula and in the neighborhood of nearly all the mining centers in the western part of the territory. The product of the Montana mills is entirely used to supply the local demand.

The following report upon the forests of the northern Rocky Mountain region was prepared by Mr. Sereno Watson, of Cambridge, a special agent of the Census Office, in the division of Forestry:

"The territory whose forest resources I attempted under your instructions to examine includes an area of about 150,000 square miles, extending from the one hundred and thirteenth meridian to the summit of the Cascade mountains and from the parallel of $44\frac{1}{2}^{\circ}$ to the British boundary; or, according to political divisions, the western fifth of Montana, the northern two-fifths of Idaho, the eastern three-fifths of Washington territory, and the northeastern portion (or nearly one-half) of Oregon. It comprises a central treeless plateau of some 30,000 square miles in extent, the great 'plain of the Columbia', surrounded by more or less extensively timbered mountain systems. This tract is bounded on the east by the broad mountain range which separates Montana from Idaho, on





the north by the irregularly-broken country which lies north of the Spokane river and of that portion of the Columbia which has a westerly course in the same latitude, and on the west by the Cascade range, while on the south the circuit is less completely closed by the somewhat complicated system known as the Blue and Salmon River mountains.

"With the exception of a single county (Beaver Head) in Montana this entire region is drained by the Columbia river, since the Rocky Mountain divide, or the main central divide between the headwaters of the Columbia and of the Missouri, forms the boundary between Idaho and Montana only as far north as $45^{\circ} 40'$, when it turns abruptly eastward for 75 miles and then again northwestward to the British boundary in continuation of the line of the Wind River mountains of Wyoming.

"As will be seen from the detailed account which is to follow, the general character of the forest growth throughout this region is remarkably uniform, both in the kinds of trees found and in their manner of distribution. The trees of the most constant occurrence and that form the mass of the forests generally are, first, the red fir (*Pseudotsuga Douglasii*) and yellow pine (*Pinus ponderosa*), gradually giving place at higher altitudes to *Picea Engelmanni* and *Abies subalpina* or *Pinus Murrayana*; while of only somewhat less extended range, though sometimes more local in their distribution, are the larch (*Larix occidentalis*), cedar (*Thuja gigantea*), the white pine (*Pinus monticola*), the hemlocks (*Tsuga Mertensiana* and, less frequently, *Tsuga Pattoniana*), *Abies grandis*, and the balsam poplars. *Abies amabilis* only is confined to the Cascade mountains. No other species occur of sufficient size to be of importance as timber trees.

"In order to indicate more particularly the extent and distribution of the tree growth, the territories will be taken up in order by counties, and an estimate given of the area covered in each, though necessarily of the actual density of growth and amount of available timber (varying from square mile to square mile, and much of it never seen) no estimate can be given of such probably approximate accuracy as to be of any value whatever:

" MONTANA.

"BEAVER HEAD COUNTY (4,230 square miles).—This county, nearly equally divided by the one hundred and thirteenth meridian, is surrounded on three sides, north, south, and west, by the Rocky mountains, and is divided into two portions by a lofty spur which sets off in a northeasterly direction from the middle of the western side. The valleys of Beaver Head river and Red Rock creek to the east and south of this spur are treeless, except that the latter stream is bordered with a considerable growth of *Populus angustifolia*, often 60 feet high and a foot or two in diameter. The region to the east is scantily supplied with timber of any kind, while the northern flanks of the Rocky Mountain range as far as the head of Horse Plains creek are only sparingly timbered on the ridges and in some of the cañons, the trees small and mainly red fir, with some *Pinus albicaulis*, the summits and exposed ridges wholly bare.

"The broad dividing spur, which includes Bald mountain and several other peaks from 10,000 to 11,000 feet high, is about 30 miles in length by 20 in breadth. The peaks are bare above 9,000 feet, and the western slopes have some timber in the upper ravines alone. The trees at 7,000 feet are mainly red fir, giving place above to a small growth of *Pinus Murrayana* and *P. albicaulis*. On the eastern side of the spur there are deep, densely-wooded valleys, the timber said to be chiefly red fir, *Picea Engelmanni* and *Pinus Murrayana*. On Rattlesnake creek in this region there is a single saw-mill, 12 miles from Bannock City, which supplies the town and neighboring mining camps with lumber. When a better quality is needed it is brought across the mountains from the Lemhi River district in Idaho by a road crossing the range at the head of Horse Plains creek. The fuel used in Bannock City is hauled some 12 or 15 miles, chiefly from the Rocky mountains. Beyond the head of Horse Plains creek (where the Bald Mountain spur commences) the range for about 40 miles changes in character greatly, becoming higher, broader, and more rocky, with rugged, snow-clad peaks from 10,000 to 12,000 feet high, and with high, rocky spurs to the east, separating densely-wooded valleys difficult of access and rarely visited. The forests here come down to the western edge of Big Hole valley, and are continuous. The trees are said to be largely *Pinus Murrayana*, but there is probably a considerable proportion of red fir, *Picea Engelmanni* and *Abies subalpina*. The range now takes its turn to the east, forming the northern line of the county, and rises again into some high, snowy peaks, but is much less densely wooded. Where the pass crosses the range from the Big Hole valley to the Bitter Root, the prevalent tree is found to be *Pinus Murrayana*, mixed toward the summit of the divide (at 7,000 feet altitude) with some red fir and a small proportion of *Picea Engelmanni* and *Pinus albicaulis*. The trees are mostly young and small, evidently frequently overrun by fires, a dense new growth immediately in most cases replacing the old. The trunks very rarely reached a diameter of 15 or 20 inches. The timbered area of the county may be estimated at 1,000 square miles. No yellow pine was seen or heard of within its limits.

"DEER LODGE COUNTY (6,500 square miles).—This county, also nearly bisected by the one hundred and thirteenth meridian, is occupied by spurs of the Rocky mountains, which form its southern and eastern border, with the intervening open valleys of Deer Lodge river, Flint creek, and Big Blackfoot river. These spurs are to a large extent wholly bare of trees, only some of the ravines and ridges being covered by a more or less scattered growth of yellow pine and red fir of moderate size, and the higher northern slopes by a denser growth of *Pinus Murrayana*. North of the Big Blackfoot the timber is more dense, coming down into the valley, and consisting principally of

yellow pine, with some red fir and larch, and at the higher elevations (above 5,000 feet) of red fir, larch, and scrub pine. In the higher cañons of the main range to the south it is probable that *Abies subalpina* and *Picea Engelmanni* also occur, as I heard of a soft tamarack found at Gwendale, which appeared from the description to be the latter species. Total timbered area of the county is estimated at 2,250 square miles.

"MISSOULA COUNTY (21,000 square miles).—The Bitter Root mountains, which separate this county from Idaho, are a direct continuation of the Rocky mountains north from the point of divergence of that range in latitude 45° 40'. While broadening out until they cover a base of 100 miles or more, they rarely reach a height of 8,000 feet. There is nothing alpine in the character of their higher vegetation, nor do they anywhere rise above the limit of forest growth. The summits are not often very rugged, and though the slopes may be steep they are not generally greatly broken. For the most part they are well wooded upon both sides, with no meadows along the streams and little grass anywhere until the foot-hills are reached. Upon the Montana side it is from 20 to 40 miles from the base to the summit of the divides, and the Bitter Root valley, which skirts their feet for 60 miles, separates them from the low and comparatively bare spurs of the Rocky mountains on the east.

"Upon crossing the main divide upon the southern border, between the Big Hole and the Bitter Root valleys, at an altitude of 7,000 feet, the yellow pine immediately appears, of large size, and with its usual massive habit, and is henceforth the most conspicuous forest tree along the usual routes of travel, coming farther down into the valleys than any other tree, and more frequently attaining a large size, probably from its less liability to serious injury from fires. The descent from this southern divide to the Bitter Root valley is well wooded with large trees of the yellow pine and red fir (with at first some small *Pinus Murrayana*), which continue to be the only trees seen bordering the valley. These mountains were crossed by me on the Lolo trail up the Lolo creek, and by the Mullan road, which follows the Saint Regis de Borgia river. In the lower cañons only yellow pine and red fir are found, of the usual scattered growth—the trees rarely much over 2 feet in diameter—with larch and *Abies subalpina* coming down the side gulches, and white pine in the middle cañon. The yellow pine gradually gives place to *Picea Engelmanni*, *Abies subalpina*, and *A. grandis*, though none of these reach the size which they attain on the Idaho side of the range. The large timber is mostly confined to the neighborhood of the streams, where the larch and white pine sometimes reach a diameter of 3 or 4 feet, but it becomes much smaller upon the ridges, and in the upper cañons rarely is more than a foot or 18 inches through, while the mountain slopes are usually much burned over and covered with fallen timber, largely of *Pinus Murrayana*. The largest and most abundant tree in the upper cañons is probably the *Picea Engelmanni*. Small trees of the *Thuja gigantea* are also occasional, but nowhere in northern Montana does it become large enough to be of importance. The *Thuja* and *Abies grandis* extend as far south as the Nez Percé creek. I think that no hemlocks were seen on the Montana side of the range, but they may occur.


"North of the Mullan road to Clarke's Fork the eastern slopes of the range continue well wooded. On the eastern side of the county the low spur of the Rocky mountains lying to the east of Bitter Root valley is to a large extent bare, but has some young yellow pine (known as 'scrub pine' or 'black jack pine') and *Pinus Murrayana*. Approaching Hellgate river the timber becomes more prevalent in the ravines; and in the government timber reservation near Missoula, where there is a saw-mill run for government purposes, the timber was found to be yellow pine and red fir (not large) and considerable larch—fine trees 2 feet in diameter or more and 100 feet high. In Granite cañon, in the mountains north of Missoula, where there is also a saw-mill, the lower cañon was occupied by yellow pine and larch, with some red fir rarely over a foot through, and in the upper cañon *Abies subalpina* and *Picea Engelmanni* a foot in diameter. On the mountain sides above the cañon the timber is, as usual, small and worthless for lumber. In the eastern portion of the Flathead Indian reservation a very high and rugged range of mountains extends nearly as far north as the head of Flathead lake, and parallel with the main Rocky Mountain range, which here enters the county and continues across the northern boundary. Both of these ranges are throughout densely wooded, though on the eastern side of the Rocky mountains the timber wholly ceases a few miles (8 or 10) below the summit, giving place to the open grazing region of the upper Missouri. On crossing over the lower end of the western range, from the Big Blackfoot to the Jocko river, the timber was found to be at first almost entirely yellow pine, with red fir and larch in the gulches, the yellow pine ceasing toward the divide (at 5,000 feet altitude) and *Pinus Murrayana* taking its place, but reappearing on the northern side, with occasional *Picea Engelmanni* and even small *Thuja gigantea*. No white pine was seen. The same trees probably continue northward to the boundary (the larch is reported from about 30 miles south of the boundary).

"The remainder of the county, lying west of the Flathead River valley, is wholly occupied by mountains, of less altitude, but, so far as is known, generally densely wooded, with the exception of some of the spurs toward the Flathead river and Clarke's Fork and some small prairies bordering the streams.

"The total timbered area of the county is estimated at 17,000 square miles."

WYOMING.

The highest mountain ranges in Wyoming only are well timbered. The high rolling table-land which occupies the central part of the territory is destitute of all tree growth, while the low ranges which rise from this plateau



WYOMING

and border it on the south are either treeless or only support a few stunted yellow pines or firs widely scattered or forming small, isolated patches of open forest upon the highest slopes of the most sheltered ravines. The most important forests of the territory are those in the northeastern corner covering the western extension of the Black hills of Dakota, those upon the foot-hills and slopes of the cañons of the Big Horn mountains, and the dense forests of small lodge-pole pine (*Pinus Murrayana*) which occupy all the northwestern portion of the territory studded by the system of mountains surrounding the Yellowstone park.

The most valuable timber of the Big Horn mountains consists of yellow pine (*Pinus ponderosa*), attaining on the foot-hills sufficient size to furnish saw-logs. Probably one-third of this timber on the east side of the mountains has already been cut to supply mills located upon the streams from Crazy Woman creek to Tongue river. The table-land on both sides of the range between the crests of the foot-hills and the base of the Snow range is covered with a belt, from 8 to 10 miles in width, of small white fir (*Abies subalpina*). The trees are small, rarely exceeding 8 or 10 inches in diameter. They afford, however, useful material for fuel and fence and telegraph poles. This forest has suffered seriously from wind storms and fire. A heavy growth of cottonwood, with which is mingled a little green ash, occupies the banks of all the streams of the Big Horn region, with the exception of No-Wood creek, flowing from the western flank.

The forests of the Yellowstone region, composed for the most part of small lodge-pole pine, are confined to the mountain slopes and high valleys, at an elevation of between 5,000 and 10,000 feet. These forests are capable of supplying great quantities of fuel and fencing material. They contain, however, little timber suitable to manufacture into lumber.

The forests of all this arid central region suffer seriously from fire. These increase with the settlement of the country and inflict great damage upon the forest. In northwestern Wyoming, however, the forests of lodge-pole pine (*Pinus Murrayana*) destroyed by fire reproduce themselves, and the area occupied by this species in all the Rocky Mountain region is increasing. This is due no doubt to the fact that fire does not destroy the seeds of this species, protected in the cones, which remain closed upon the trees for years. The heat of the fire causes the cones to open and shed their seeds upon the burned surface of the soil, where they germinate quickly and freely.

During the census year 83,780 acres of woodland were reported destroyed by fire, with an estimated loss of \$3,255,000. These fires were set by Indians, trappers, and prospectors.

A little lumber, in addition to that manufactured in the Big Horn region, of which no returns have been received, is sawed in the Medicine Bow and other ranges in the southern part of the territory. A large amount of fire-wood and many railroad ties are cut in the southern mountains and delivered by chutes along the line of the Union Pacific railroad.

COLORADO.

The forests of Colorado are confined to the mountain ranges and high valleys which cover the western half of the state; the elevated, rolling plateau which extends from the eastern base of the mountains to the eastern boundary of the state is entirely destitute of tree growth, with the exception of an occasional stunted willow or cottonwood found in the bottom lands of the large streams. The important forests of the state cover the mountain slopes between 10,000 and 12,500 feet elevation, and are almost exclusively composed of spruce (*Picea Engelmanni*), with which are mingled different alpine pines of little economic value. Below the spruce belt a more open forest of red fir and yellow pine, occupying ravines or scattered over the ridges, extends down to the foot-hills. These are covered with an open growth in which the nut pine and the western juniper are the prevailing trees, while the borders of streams and bottoms of the cañons are occupied by cottonwoods, willows, cherries, oaks, and other deciduous trees and shrubs of little economic importance. Large areas upon the sides of the high Colorado mountains are exclusively covered with a dense growth of the quaking aspen. This tree very generally takes possession here of ground from which the coniferous forest has been removed by fire, and, as the number of forest fires is rapidly increasing in Colorado, it seems destined to become the only widely-distributed forest tree of this region. The high valleys, or "parks" as they are here locally called, when timbered at all, are covered with a dense forest growth in which the lodge pole pine (*Pinus Murrayana*), also common at high elevations in the spruce forests, is the prevailing and often the only species, disputing with the aspen the possession of the burned soil. The high plateau of southwestern Colorado is either treeless or is thinly covered with an open growth of small, stunted junipers.

The increase in the number of forest fires raging in the mountains of Colorado is alarming in a region where the forest once destroyed cannot easily reproduce itself, and upon mountains where forest covering is necessary to preserve the integrity of the channels and the constant flow of numerous important streams essential to the irrigation of wide areas of arid territory.

During the census year 113,820 acres of forest were reported destroyed by fire, with an estimated loss of \$935,500. These fires were set by careless hunters, miners, and prospectors, and by Indians or whites through malice.

The forests of the Colorado foot-hills afford abundant fuel and fencing material to supply the wants of the present population of this part of the state. Coarse lumber, suitable for the timbering of mines and railroad

construction, is manufactured from the fir and pines of the lower mountain slopes, which have also furnished immense quantities of fuel and railway ties. The timber, however, of this forest most accessible to mining centers and the lines of railroads has already been destroyed, while its productive capacity is everywhere impaired by wasteful methods of lumbering and destructive conflagrations. The elevated spruce forests, which contain the only great bodies of heavy timber found in the central Rocky Mountain region, have thus far, on account of the difficulties of operating in them, escaped all serious inroads from the ax of the lumberman. Small portable mills, however, have been established in these forests to supply the wants of some of the most elevated mining centers, and fires every year reduce their extent and value.

Colorado is principally supplied with lumber from Chicago; a small amount is manufactured, however, in the state, mostly upon the waters of the South Fork of the Platte river, in Jefferson county, and in the extensive pineries which cover the divide between the waters of the South Platte and the Arkansas rivers. A little lumber is also manufactured in small portable mills in nearly every county.

NEW MEXICO.

The forests of New Mexico are confined to the slopes and cañons of the high mountain ranges. The elevated plateau which occupies the whole of the eastern part of the territory is treeless, with the exception of occasional willows and cottonwoods bordering the large streams, while the high *mesas* of the southwest and west are sometimes dotted with an open growth of dwarf junipers and nut pines of considerable local importance as a source of fuel and fencing supply. The high mountain ranges extending southward into the northern part of the territory are covered with forests very similar in composition, density, and distribution to those covering the mountains of Colorado. Engelmann's spruce is here the important timber tree at high elevations; lower, open forests of red fir and yellow pine occupy the sides of cañons and the lower mountain slopes, and the nut pine and juniper cover the foot-hills with an open, scattered growth. The detached mountain ranges which spring from the central plateau of the territory are less heavily timbered than the higher mountains north and south. The yellow pine is here the most common and important tree, mingled in sheltered cañons and at highest elevations with occasional red firs.

The most important forests of the territory cover the high group of mountain ranges west of the Rio Grande and south of the thirty-fourth degree of latitude—the San Francisco, the Tulerosa, Sierra Blanca, Sierra Diablo, Mogollon, Pinos Altos, and Mimbres. The foot-hills and lower slopes of these mountains, between 5,000 and 7,000 feet elevation, are covered with a heavy growth of junipers, nut pines, and different evergreen oaks. The banks of streams are here lined with immense cottonwoods, sycamores, cherries, ashes, and hackberries, while the *arroyos* or depressions in the *mesas* contain fine groves of mesquit. Above an elevation of 7,000 feet the yellow pine appears, and mingled with it on north slopes the red fir and white pine (*Pinus reflexa*); the elevated valleys contain fine groves of cottonwood, box-elder, alder, and small oaks, while the most inaccessible slopes of some of the highest ranges are covered with forests of cypress (*Cupressus Guadalupensis*).

The coniferous forests of these mountains are dense and valuable, and, although not yet accessible for lumbering operations except at a few points, they seem destined to become an important factor in the future development of the whole region. They can, if properly protected, supply with lumber indefinitely a larger population than will probably occupy this part of the United States.

The deciduous trees of this entire southwestern region, often of considerable size, are generally hollow, especially the oaks; they are of little value for any mechanical purpose, although affording abundant and excellent fuel.


During the census year 64,034 acres of woodland only were reported destroyed by fire, with an estimated loss of \$142,075.


A small amount of coarse lumber, principally yellow pine, is manufactured in the territory, mostly in the counties of San Miguel and Santa Fé. New Mexico, however, like Colorado, obtains most of its lumber by rail from Chicago.

ARIZONA.

Northern, western, and southwestern Arizona are destitute of true forests. Ravines in the *mesas* of the high Colorado plateau of northern Arizona are occasionally covered, however, with stunted junipers. Cottonwoods and willows line the banks of the Colorado river, and the ironwood, the palo verde, the mesquit, the suwarrow, and other Mexican forms of arborescent vegetation are found in the valley of the Gila and the deserts of the southern part of the territory; individual trees are, however, widely scattered, nowhere forming forests in the true meaning of the word. The low lava ridges and arid lake beds with which the southwestern part of the territory is covered are entirely destitute of tree growth.

The mountain system culminating south of the Colorado plateau in the San Francisco mountains, and extending southeasterly through the middle of the territory into New Mexico, is well timbered. The high ranges which spring from this central elevated plateau bear heavy forests of yellow pine and red fir, the plateau itself





being covered, over thousands of square miles, with an open growth of yellow pine of considerable size. The streams and bottoms of the high mountains are lined with deciduous trees, of which the cottonwood, the cherry, the ash, the alder, and the walnut are the largest and most important. The group of short, detached mountain ranges which occupies with a general north and south trend the southeastern part of the territory is covered with a rich and varied forest growth. The highest slopes are covered with forests of pine, in which, in the Santa Catalina range at least, great bodies of splendid cypress (*Cupressus Guadalupensis*) are found; a little lower the red fir and white pine (*Pinus reflexa*), different oaks and junipers with a madroña, are scattered over the dry, gravelly slopes and ridges between 5,000 and 7,000 feet elevation. These in turn are replaced below 5,000 feet with an open growth of small evergreen oaks. The bottoms of the cañons and the borders of the streams between 4,000 and 8,000 feet elevation are lined in these mountains with hackberry, sycamore, cottonwood, willows, cherries, and ashes. The *arroyos* in the *mesas* are often covered, as in southern New Mexico, with noble groves of mesquit, or in drier situations support a stunted growth of acacias, yuccas, cacti, and other desert plants.

The yellow pine is the only tree of Arizona of great importance as a source of lumber supply. Oaks and other hard-wood trees are invariably defective and of little value except for fuel. The red fir, white pine, and cypress occur only at high elevations, and are generally too scattered and too difficult of access to make their manufacture into lumber practicable for the present at least.

The pine forests of central Arizona and southwestern New Mexico are of great importance to the development of the treeless regions which surround them. No other body of timber of any extent or value exists near the southern boundary of the United States between the pine belt of eastern Texas and the forests of the California mountains. These southern interior forests have nowhere yet greatly suffered. Their inaccessibility has protected them. Railroads, however, now either penetrate this forest region, or will soon do so, and these, with the rapid development of the mining industry now going on in the southwest, threaten these forests with the dangers which are fast exterminating those of Colorado and Utah.

During the census year 10,240 acres of woodland were destroyed by fire, with an estimated loss of \$56,000. These fires were set by careless hunters, prospectors, and Indians.

Pine lumber is sawed in Pima and Pinal counties, principally upon the Santa Catalina, Santa Rita, and Huachuca mountains, to supply important mining centers in this part of the territory. It is also manufactured in small quantities in portable mills near Indian reservations and other centers of population throughout the forest region. Returns from 13 mills only, situated in Pima, Pinal, Apache, and Yavapai counties, have been received. Southern Arizona is now, in spite of its fine forests of pine, almost entirely supplied by rail with lumber manufactured in California.

UTAH.

The Uintah range, occupying with an east and west trend the whole of the northeastern part of the territory, the Wahsatch mountains and their southern extension, the San Pitch and the Sanpete ranges, extending north and south nearly through the center of the territory, and the mountains which bound on the east the great Colorado plateau, bear at high elevations fir, spruce, and pine forests of considerable extent. The foot-hills of these mountains and their high valleys are dotted with an open growth of nut pine, juniper, and mountain mahogany (*Cercocarpus*). The high Colorado plateau and the arid deserts of western and southern Utah are treeless, with the exception of a few stunted junipers and nut pines which struggle for existence upon some of the low mountain ranges, and of willows and cottonwoods which line the banks of the infrequent and scanty streams.

The western flank of the Wahsatch mountains north of the fortieth degree of latitude has already been almost denuded of its best timber to supply the wants of the agricultural and mining settlements of the Salt Lake region, and the scanty forests of the territory have everywhere suffered serious loss from fire and wasteful methods of cutting timber and railway ties and of manufacturing charcoal.

During the census year 42,865 acres of woodland were reported destroyed by fire, with an estimated loss of \$1,042,800. These fires were set by Indians, wood-cutters, careless hunters, and prospectors.

Small quantities of lumber—pine, cottonwood, and a little spruce—are manufactured through the Wahsatch region, the principal centers of manufacture being Beaver City and Cedar City, in the south, the neighborhood of Salt Lake City, and Cache county in the extreme northern part of the territory. Utah is, however, almost entirely supplied with lumber from the eastern slopes of the California sierras and from Chicago. Small tanneries in Salt Lake City obtain a supply of red fir and spruce bark from the neighboring mountains.

The following notes upon Utah forests, made during the prosecution of a special investigation into the meat-producing capacity of the territory, have been supplied by Mr. E. C. Hall, a special agent of the Census, in the division of "Meat Production in the Grazing States and Territories":

"The timber of the Wahsatch mountains, in Cache, Rich, Morgan, and Weber counties of Utah, hardly suffices for the wants of the settlers. The trees from which lumber is obtained are cedar and a variety of white pine (*Pinus flexilis*). Some fir (*Pseudotsuga Douglasii*) is found, but it is not common north of the latitude of Salt Lake City. This tree likewise furnishes an inferior kind of lumber. In general, in Utah, north of latitude 40°, the west

base of the Wahsatch mountains has been stripped of the available timber, so that in the accessible cañons, especially in the neighborhood of settlements, it is laborious and expensive obtaining posts and poles for fencing, to say nothing of smooth planks, etc., for building. Cottonwood and occasionally box-elder are found fringing the river bottoms of the sections described.

"The Oquirrh mountains, on the east of Tooele county, and the Onaqui mountains, 30 miles west, contain cedar and considerable red fir, the latter a tree which I am told is not frequent in the Wahsatch range. The mining camps of Salt Lake and Tooele counties have largely depleted the timber areas of these mountains. More timber is standing on the Onaqui hills than on the Oquirrh range. From the vicinity of the latter to Cottonwood and Bingham Cañon mining districts the dearth of good fencing material is very noticeable throughout Box Elder, Cache, Rich, Weber, Morgan, and Salt Lake counties. Willow withes, stone walls, cottonwood poles, and sod walls flanked by ditches are among the devices for barriers against stock incursion, all pointing to the lack and costliness of lumber.

"LAKE RANGE, WEST OF UTAH LAKE.—This range of low mountains contains scattered black balsam and red fir. In winter this range is visited from the settlements of Utah valley, and the trees felled and sledged across the lake on the ice, to be used by the railroad and by farmers. No piñon pine was found in the Oquirrh or Onaqui mountains.

"The San Pitch mountains, in latitude $39^{\circ} 30'$, longitude $111^{\circ} 52'$, contain sparse timber—a so-called white pine (*Pinus flexilis*), scrubby cedar, and some other evergreen trees—at a high elevation and unavailable as lumber. No good clear planking suitable for building is obtained from these cuts.

"The low ranges west of Juab valley and flanking Dog valley, Dry valley, and Ferner valley, in latitude $39^{\circ} 30'$, longitude 112° , contain straggling cedar and some red fir difficult of access. The timber of the whole region north of latitude 39° and west of the main Wahsatch mountains is meager and inadequate for the purposes of the Mormon settlers.

"Fencing about Salt Lake City is of poor construction and costs \$200 per mile of pine poles and cedar posts. I saw some posts of white balsam (*Abies concolor*) 50 feet long, obtained from the cañons of the San Pitch range, used for fencing. At Springville, in Utah valley, posts of cedar were pointed out which were in good condition after fourteen years standing. Were it not for the existence of the 'no-fence' law, which enables a farmer to cultivate unfenced ground and claim damages from incursions of stock, the Utah farmer would be very badly off, not having means to purchase fencing material in a country so ill supplied with timber. The cedar which abounds here affords a lasting supply of fire-wood and posts, but for poles or plank the region depends largely upon imported lumber, especially for building plank, joists, etc.


"Upon ranges flanking East and West Tintic valleys, Juab county, in latitude $39^{\circ} 50'$, longitude $112^{\circ} 30'$, the timber is not abundant; it consists of red fir and black and white balsam, from which rough lumber for the mining camps of Mammoth, Tintic, and Silver City has been extensively taken. Cedar of the usual dwarfed kind grows abundantly along the upper slopes of the foot-hills, and is used for braces and posts in shafts of mines.

"SANPETE VALLEY RANGE (longitude $111^{\circ} 30'$; latitude $39^{\circ} 20'$).—The Wahsatch mountains, on the east of Sanpete valley, carry on their spurs and through the deep cañons facing the valley some of the best timber found in Utah. It is largely used by the Sanpete settlers. Yellow pine, black and white balsam, red fir, cedar, and poplar constitute the varieties of trees found. The yellow pine, less abundant now in accessible cañons, furnishes, it is claimed, a clear and firm lumber, fit for building, and not surpassed by any variety in Utah. The range west of the Sanpete mountains—i. e., the San Pitch mountains before spoken of—carries on its eastern slopes and cañons considerable balsam of both varieties and some red fir and poplar about the headwaters of creeks. Little yellow pine is found on the San Pitch range; at least, none is taken out at present, although I was told considerable had already been lumbered from such cañons as were penetrable. Fencing of cedar posts and poplar and balsam poles is largely used in the valley; cedar posts and pine plank are also used in fencing meadows and fields. In no other valley of Utah are the Mormons so well supplied, apparently, with fair lumber of native growth. Except for furniture and house trimmings, no imported wood is used here.

"SEVIER RIVER MOUNTAINS (latitude $38^{\circ} 30'$ to $39^{\circ} 10'$, longitude 112°).—The Tushar mountains and the Valley range, on the west of the Sevier valley, are supplied with meager timber, especially the Valley range. In no part of Utah have I noticed so few and so limited areas inclosed. Timber is said to exist in inaccessible places only on the Wahsatch range to the east of the valley. This is true in regard to the ranges west of the Sevier valley, where the character of the tree growth is inferior to even the average poor quality of Utah forests. Black balsam, white balsam and red fir grow in both ranges, but are approached with great difficulty. The indigenous scrub cedar prevails often in thick groves along the foot-hills, especially on the Valley and Tushar ranges to the west of the valley. Several saw-mills at the mouths of cañon streams on the East Wahsatch range have for several years worked up all the available lumber, but the prices asked for lumber—from \$35 to \$45 per 1,000 feet—place fencing material beyond the reach of the Sevier farmer.

"All the way up the Sevier valley, and along its south and east forks, fencing is limited and lumber high, a sure proof of the inadequate supply of trees on accessible mountains.

"Fish Lake plateau and mountains (latitude $38^{\circ} 33'$, longitude $111^{\circ} 50'$) contain a considerable quantity of the prevailing timber of Utah, as do also Thousand Springs mountains.







"The Aquarius plateau is said by Sevier Valley stockmen to be abundantly timbered with pine, balsam, and spruce, but Boulder valley (latitude $37^{\circ} 55'$, longitude $110^{\circ} 30'$) was destitute of standing timber, save cedar and poplar on its foot-hill fringe. The country was visited by fires, the Mormons told me, in 1872-'73, which destroyed large areas of the forests in the region southeast of the Grass Valley country. The whole section of Utah lying east of the Sevier valley to the Rio Colorado is better timbered, but from its rough and impenetrable location the timber is of no avail to most of the settlers, but only to such as penetrate the high valleys of Grass, Boulder, Potato, etc., lying adjacent to the timber. Fencing on Grass and Rabbit valleys, western Pinto county, is cheaper than in Sevier valley, but farmers and stockmen are so poor that they are forced to avail themselves of the no-fence law when breaking ground for crops.

"In the Paria River region fencing is very limited and lumber expensive, as timber is hard to get out of the mountain cañons.

"In the Kanab River region fencing at the settlements of Upper Kanab and Lower Kanab, Kane county, is said to be expensive, as material is difficult to obtain, the cañons leading to the valley affording a meager supply of cedar and black and white balsam, while some red fir and yellow pine is said to grow on the Sevier plateau (latitude $37^{\circ} 30'$); this, however, seldom reaches a market in the settlements, owing to the isolated situation of these forests.

"Considerable scrub oak is found on the slopes of the Oquirrh and Onaqui mountains, above referred to, in Tooele county, and many cedar thickets of considerable extent. In Tooele valley some fencing with cedar and panels of balsam occurs. Rush valley contains some bull fences of trunks of cedar, costing \$1 25 per rod, showing the cost of even poor material.

"It will be observed that outside of the Wahsatch mountains no building timber of value has been noted in Utah. The supply in this range has been largely consumed from the easily-approached cañons and slopes.

"In summing up my observations, which were made wholly with a view of investigating the fencing of pasture areas and cost of same, it may be stated that Utah seemed very generally lacking in serviceable material for fencing or building. The country settled for thirty years has drawn upon the near supply of standing timber, so that now lumber is obtained by great exertion and expense in most of the valley settlements. The labor and cost of fencing caused Brigham Young to enact the no-fence law, which enabled the destitute settlers to break ground, irrigate, and raise grain without the provision of any barrier against stock inroads, the cattleman being held responsible for the damages of his herd. This law in itself is a commentary on the scarcity of timber in Utah."

NEVADA.

The tree growth of Nevada, except in a portion of Douglas county, in the extreme western part of the state, which the forests of the California sierras just reach, is confined to the low ridges of the central and southern part of the state. The most important of these—the Humboldt, Toiyabe, Monitor Creek, Timpiute, Hot Creek, Kawich, and probably others—bear near their summits, in sheltered ravines, scattered patches of stunted white pine (*Pinus flexilis*) of sufficient size to furnish saw-logs. The lower slopes of the mountains of this region are often quite thickly covered with small nut-pines and groves of the mountain mahogany (*Cercocarpus*), here attaining its greatest development. Below the nut-pine low, stunted junipers cover the foot-hills, often extending, in the central part of the state, across the narrow elevated valleys which separate the low mountain ranges.

The great development of the mining interests of Nevada has already nearly exterminated its scanty and stunted forests. The white pine has been cut in the neighborhood of mines from all the mountain ranges, and the most accessible nut-pine, juniper, and mountain mahogany have been converted into cord-wood or made into charcoal. The forests of Nevada are nowhere reproducing themselves, and a scarcity of fuel, even for domestic purposes, must soon be felt.

A considerable amount of lumber is manufactured in the neighborhood of lake Tahoe, in Douglas county, and sent in flumes down the eastern slopes of the sierras to supply Carson City and Virginia City. The lumber-manufacturing interests of the remainder of the state are necessarily small and unimportant. Their entire extermination, with the forests which furnish them material, cannot be long delayed.

During the census year 8,710 acres of woodland were devastated by fire, with a loss of \$19,000. The fires were traced to hunters and Indians.

IDAHO.

The western slopes of the Bitter Root and Cœur d'Alène mountains, which form north of latitude 46° the eastern boundary of the territory of Idaho, are covered with dense, extensive, and valuable forests of fir, pine, and larch. The ridges of the Rocky mountains, which below latitude 46° occupy the eastern border of the territory, and the extreme eastern development of the Blue mountains of Oregon, just entering it from the west, are less heavily timbered with a scattered growth, in which yellow pine and red fir are still the prevailing trees. The great central region occupied by the Salmon River mountains is unexplored. These mountains are more or less timbered, but nothing is known of the composition or character of the forests which cover them. Judging, however, from the general elevation and climate of this region, its forests cannot be very important, nor capable of

supplying more than the local wants of its mining population. The great plains south and southeast of the Salmon River mountains, comprising fully one-third of the territory, are entirely destitute of tree covering, while the Snake River range and the ranges of the Bear River country contain in their more sheltered cañons only small areas of open, stunted forest.

During the census year 21,000 acres of woodland were reported destroyed by fire, with an estimated loss of \$202,000. These fires originated in the carelessness of hunters, prospectors, Indians, etc.

A small amount of pine and fir lumber is manufactured at Boise City and near other centers of population. The great forests of cedar, fir, and pine, however, in the Cœur d'Alène region are still almost intact. These forests, with proper care, are capable of furnishing indefinitely the treeless agricultural region of eastern Washington territory and Oregon with an abundant supply of excellent building material.

The following extracts are made from Mr. Sereno Watson's report upon the forests of the territory:

"This territory north of latitude 44½° is occupied by the Rocky and the Bitter Root mountains, forming its eastern boundary, with their broad, timbered, interlacing spurs, which terminate in the high, mostly treeless plateau which extends from near the Spokane river in a southeasterly direction to this parallel of latitude. The southern and southwestern portions were not visited by me, and the statements regarding them are to some extent conjectural.

"LEMHI COUNTY (5,530 square miles).—In the extreme eastern portion of this county, where the mountains are crossed by the Utah and Northern railroad, scattered trees of red fir are first met at an altitude of 6,000 feet. Beaver cañon, up which the railroad passes, is well timbered on both sides nearly to its head at 6,600 feet altitude with red fir only, but the broad plateau at the summit (6,869 feet) is treeless. In the lateral cañons (8 to 10 miles long), coming out near the mouth of Beaver cañon, there are two saw-mills, one of which was visited. The timber was here found to be confined to the south side of the cañon, and consisted almost wholly of red fir (here called 'red pine'), averaging from 20 to 22 inches in diameter. The largest log seen measured 32 inches at the butt. A 'white pine' proved to be *Picea Engelmanni*, and a 'bird's-eye pine' was *Pinus Murrayana*, both small, as was also the balsam (*Abies subalpina*), which was found some 3 or 4 miles up the cañon. The yellow pine did not occur here.

"It is probable that the cañons westward along the range are similarly timbered as far as the Lemhi agency. Here the character of the range changes (as stated under Beaver Head county, Montana), becoming higher and more rugged, and the *Pinus Murrayana* is probably more abundant, at least at the higher altitudes. The yellow pine also appears, but at what point is uncertain; it is certainly found at Gibbonsville, on the North Fork of the Salmon river, and it probably extends still farther southward. The Salmon River mountains, lying between the Lemhi river and Rock creek, are reported to be well timbered. The southwestern portion of the county I presume to be much more open.

"The total timbered area is estimated at from 1,500 to 2,000 square miles.

"IDAHO COUNTY (10,100 square miles).—The high and crowded spurs of the Bitter Root mountains fill the entire northeastern portion of this county, extending to the line of the South Fork of the Clearwater, mostly densely wooded from base to summit. The foot-hills and plateaus between the streams are more or less covered with scattered yellow pine and red fir. The valley of the Salmon river is probably comparatively treeless, and the low mountain range between that river and the Snake is scantily timbered.

"Estimated timber area, 4,000 square miles.

"WASHINGTON COUNTY (3,000 square miles).—I have but little information in regard to this county. The southern portion has been surveyed, and is probably nearly treeless. The rest appears to be more mountainous, and may be scantily timbered.

"Wooded area (say) 300 square miles.

"NEZ PERCÉ COUNTY (3,400 square miles).—Mainly high plateau, at about 3,000 feet altitude, in the southeast more or less covered with scattered yellow pine and red fir of good size, on the western side nearly without timber or with occasional yellow pine. Toward the head of Potluck creek some yellow pine and red fir are found in the valleys, and in the northeast the spurs from the Rocky mountains enter the county, covered in addition with the larch and *Thuya gigantea*. East of the Indian reservation the county extends up into the mountains in the form of a narrow gore, and is heavily timbered. The portion lying south of the reservation in the angle between the Snake and Salmon rivers is occupied by low mountains, mostly bare.

"Total timbered area estimated at 750 square miles.


"SHOSHONE COUNTY (5,950 square miles).—Wholly mountainous and covered with forests, with the exception of some prairies and open country near the Clearwater and lower portion of the Lolo Fork.

"Immediately after crossing the divide by the Lolo trail from Montana, at an altitude of 6,000 feet, the forest consisted of *Abies subalpina* and *Picea Engelmanni*, with young *Abies grandis* and *Tsuga Mertensiana*, and occasional larch and red fir, and upon the creeks some small *Thuya* and *Taxus*. The trail soon ascended the ridges and followed them for about 100 miles at an altitude of from 5,000 to over 7,000 feet, doubtless to avoid the fallen timber which made the cañons impassable, though enough of it was found on the route followed. The timber on these ridges was often small and scattered—*Abies subalpina* and *Picea Engelmanni*, with *Pinus Murrayana* and *P. albicaulis*—or on the damper northern slopes with larch and red fir, balsam, hemlock, and sometimes the mountain

DENSITY OF FORESTS


COMPILED UNDER THE DIRECTION OF
C.S. SARGENT, SPECIAL AGENT
1883.

Scale of Stat Miles



LEGEND

- I Under 1 cord per acre
- II From 1 to 2 " " "
- III 2 to 5 " " "
- IV 5 to 10 " " "
- V 10 to 20 " " "
- VI Over 20 " " "





hemlock (*Tsuga Pattoniana*), the trees larger (occasionally 2 feet through, the *Abies grandis* being the largest). The white pine (*Pinus monticola*) also frequently occurred. During the last day upon this ridge the trail was through heavy timber, chiefly of hemlock sometimes 3 feet in diameter, with some *Abies* and rarely *Pinus Murrayana* and *P. monticola*, the ridge even at 7,000 feet being covered with the same dense growth. Descending quite abruptly from the drier extremity of the spur, which was covered with *Abies*, *Tsuga*, and *Pinus* (*Murrayana*, *albicaulis*, and *monticola*), we passed through a forest of heavy balsam (*Abies grandis*), with a few larch and some red fir, and at about 4,000 feet came upon cedar (*Thuja gigantea*) to the exclusion of everything else—the trees from 2 to 4 feet in diameter. On the stream banks at the base were found the *Thuja*, *Pinus monticola*, *Abies subalpina* and *A. grandis*, *Picea Engelmanni*, and *Tsuga Mertensiana*, all growing together, with an undergrowth of maple, mountain ash, *Vaccinium*, *Ceanothus*, *Crataegus*, *Pachystima*, *Prunus*, etc. With timber of this character upon the high ridges it is evident that there must be much very heavy timber in the cañons.

"After crossing a low ridge covered with cedar, larch, and red fir, and following a narrow meadow frequently interrupted by clumps of timber, the trail at length came out upon an open *camass* prairie 25 miles northeast of Kamai. From this point the timber covering the plateau is an open growth of yellow pine and red fir, often quite large, with young trees intermixed, and some *Picea Engelmanni* and the two *Abies* in the wetter places. Considerable timber is cut upon the Lolo Fork and Clearwater and floated down to the mills at Lewiston. It is uncertain how far south along the main range the above large variety of trees continues. It is probable, in my opinion, that the *Thuja*, *Abies grandis*, *Tsuga*, *Pinus monticola*, and *Taxus* do not pass beyond the headwaters of the Clearwater, or, at the farthest, that some of them may reach the North Fork of the Salmon river, while the larch may possibly be found in the Salmon River mountains.

"At the northern extremity of the county, along the Mullan road, which from the Cœur d'Alène mission follows up the cañon of the Cœur d'Alène river, instead of following the spurs, a distance of 37 miles, the swampy bottoms were found heavily timbered with *Thuja*, red fir, *Abies grandis*, and *Tsuga Mertensiana*, with some larch and *Pinus monticola*. Some of the drier bottoms had been burned over, and were mostly covered with *Pinus Murrayana*. Some *Populus balsamifera* occurs, 3 feet through, or more (as also on the Montana side). The sides of the ridge were also nearly bare. The *Thuja*, which exclusively occupies some of the swamps, attaining a large size, ceases at the base of the dividing ridge, where also the *Picea Engelmanni* and *Abies subalpina* come in. The range above Cœur d'Alène cañon, and bounding the county on the north, is not heavily timbered, much of its upper slopes being bare.

"Total timbered area estimated at 5,000 square miles.

"KOOTENAI COUNTY (5,530 square miles).—The portion south of the Cœur d'Alène and Spokane rivers belongs mostly to the Cœur d'Alène Indian reservation, and is timbered, with the exception of open meadows upon the Cœur d'Alène and Saint Joseph rivers and upon Hangman creek. The timber is principally yellow pine and red fir, with some *Pinus Murrayana*, and fine bodies of cedar (*Thuja gigantea*) near the western borders of the lake. North of the Cœur d'Alène river the road from the mission to the fort passes through a cedar (*Thuja*) swamp, with many large trees, from 3 to 5 feet through, traversing cañons filled with a mixed growth of *Abies subalpina* and *A. grandis*, larch, hemlock, *Picea Engelmanni*, and red fir. This latter growth continues for some miles below the fort, where the valley opens out into the broad Spokane plain, which extends northeastward toward Pend d'Oreille lake without trees. The mountains south of the lake are low and not heavily timbered. The portion of the county north of Clarke's Fork and of Pend d'Oreille lake has, so far as I know, never been explored, but is probably mountainous and for the most part well timbered.

"Estimated timber area of the county, 4,500 square miles."

WASHINGTON.

Washington territory west of the summit of the Cascade range is covered with the heaviest continuous belt of forest growth in the United States. This forest extends over the slopes of the Cascade and Coast ranges, and occupies the entire drift plain surrounding the waters of Puget sound. The highest mountain peaks and the sand-dunes of the coast are treeless. The narrow valleys of the Cowlitz and Chehalis rivers are dotted with small oaks and other deciduous trees, and oaks and stunted yellow pines occupy with an open growth the barren Steilacoom plain south of Puget sound; with these exceptions western Washington territory is covered with a magnificent coniferous forest. The most valuable and generally distributed timber tree of this region is the red or yellow fir (*Pseudotsuga Douglasii*), forming about seven-eighths of the forest growth. The valuable red cedar (*Thuja gigantea*) and the hemlock (*Tsuga Mertensiana*), often covering extensive tracts, especially near the base of the Cascade mountains, are common; the noble tide-land spruce adds value and importance to the forests bordering the coast. The forests which cover the upper ridges of the Cascade mountains are principally composed of firs (*Abies amabilis* and *A. nobilis*), spruces (*Picea Engelmanni*), various small pines, hemlocks, etc. These elevated forests, often of great beauty, are of little economic importance.

East of the Cascade mountains the forests are less dense, and are confined to the mountain ranges. The great plains watered by the Columbia and Snake rivers are entirely destitute of tree covering.

Stevens county, which is broken and mountainous, with the exception of the narrow valleys and occasional small prairies, is covered with a heavy, open forest growth. The most valuable trees of the forests of this county are the red fir, the yellow pine (*Pinus ponderosa*), the white pine (*Pinus monticola*), the larch (*Larix occidentalis*), and the red cedar.

The forests of Spokane county are confined to the spurs and ridges of the extreme eastern part of the county, and consist of the yellow pine, red fir, and larch of small size and inferior quality.

The forests of Yakima county cover about one-half of its area, being confined to the eastern slope of the Cascade range. The forests covering the eastern slopes of these mountains are only surpassed in density and value by those extending over their western flanks. The yellow pine occupies the lowest slopes with an open growth of large trees. Above the pine the red fir is the prevailing tree. This at a greater elevation is succeeded by hemlock and larch, with which are mingled fine bodies of spruce (*Picea Engelmanni*) and hemlock, while the forest growth below the timber-line consists of firs, pines, and mountain hemlock.

The western portion of Kliekat county is covered with heavy forest growth, similar in composition and density to that of Yakima.

Walla Walla county is destitute of timber except in the extreme southeastern corner, where the spurs of the mountains are thinly covered with a sparse growth of yellow pine and larch.

Columbia county is without forest except along the ridges and summit of the Blue mountains, which are covered with yellow pine, larch, and, above 5,000 feet elevation, with a continuous growth of lodge-pole pine (*Pinus Murrayana*).

Whitman county is destitute of forest except in the extreme southeastern corner, where there is a scattered growth of small yellow pine.

An estimate of the actual amount of timber standing in the territory is not possible with the existing knowledge of the country, and none has been attempted. The quantity of merchantable timber, however, standing in western Washington territory is enormous; a yield of 200,000 feet of lumber to the acre is not at all exceptional, while over fully 20,000 square miles a yield of 25,000 feet to the acre might be expected; such estimates certainly would not exaggerate the productive capacity of these noble forests.

The forests of Washington territory, especially in the more thickly settled portions west of the Cascade mountains, have long suffered from destructive fires. The injury inflicted by such fires is proportionately less, however, in the humid coast region than east of the mountains, where the dryness of the climate prevents the reproduction of the forest once destroyed. West of the mountains young trees of the species of the original forest, and especially the red fir, soon densely cover the burned surface and grow with astonishing rapidity and vigor. It seems reasonably certain, therefore, that, whatever may be the fate of the forests which now cover western Washington territory and Oregon, they will be succeeded by forests of similar composition, and that this whole region, ill adapted in soil and topography to agriculture, will retain a permanent forest covering long after the other great forests of the continent have disappeared.

During the census year 37,910 acres of woodland were destroyed by fire, with an estimated loss of \$713,200. These fires were set by Indians, by whites in clearing land, by hunters, prospectors, etc.

The forests bordering the shores of Puget sound, the strait of Juan de Fuca, and the lower Columbia river have been culled of their best trees for a distance inland of 1 or 2 miles to supply the important lumber-manufacturing interests of this part of the territory. The product of western Washington territory during the census year was 153,986,000 feet of lumber, 6,550,000 laths, 910,000 shingles, and 23,666,000 staves—by far the largest part being manufactured in the mills located on the waters of Puget sound.

The first saw-mill built upon Puget sound was erected in 1851. It was a small water-power mill, with a daily capacity of about 1,000 feet. Two years later a similar mill was erected at Seattle, with a daily capacity of from 8,000 to 10,000 feet.

The centers of manufactures now are Port Gamble, Port Madison, Port Blakely, Port Discovery, Seabeck, Utsaladdy, Tacoma, and Seattle. At the last-named place there is a large establishment manufacturing sugar-barrel staves from cottonwood for the San Francisco market.


The lumber manufactured upon Puget sound is largely shipped to San Francisco and directly to China, Australia, New Zealand, and Mexican and South American Pacific ports.

The population of the southeastern part of the territory is principally supplied with lumber, largely coarse yellow pine of inferior quality, cut on the Blue mountains in small portable mills, and delivered at Dayton, in Walla Walla county, by a flume several miles in length. No statistics, however, have been received of the amount of lumber manufactured in this county.


The methods adopted by the lumbermen of western Washington territory are wasteful in the extreme. Loggers cut only timber growing within a mile or a mile and a half of shores accessible to good booming or shipping points, or which will yield not less than 30,000 feet of lumber to the acre. Only trees are cut which will produce at least three logs 24 feet long, with a minimum diameter of 30 inches. Trees are cut not less than 12 and often 20 feet from the ground, in order that the labor of cutting through the thick bark and enlarged base may be avoided, while 40 or 50 feet of the top of the tree are entirely wasted.

—

DEPARTMENT OF THE INTERIOR



WASHI





The following notes upon the forests of eastern Washington territory are extracted from Mr. Watson's report:

"WALLA WALLA COUNTY (1,260 square miles).—This county is wholly without timber, which is supplied from the Blue mountains of Oregon.

"COLUMBIA COUNTY (2,160 square miles).—A spur of the Blue mountains traverses the southern portion of this county, occupying about a fourth of its area, which is partially timbered, chiefly with red fir (*Pseudotsuga*), pine (*Pinus ponderosa*), and some *Picea Engelmanni*, none of it large. Elsewhere the county is nearly destitute of trees, though some of the streams, especially the Touchet, were at the first settling of the county bordered by scattered pines.

"WHITMAN COUNTY (5,000 square miles).—This county is destitute of timber. Some of the townships along the Idaho line were originally sparingly wooded with scattered pines upon the ridges, but these have nearly or wholly disappeared, and the supplies for fencing and fuel are brought from the neighboring mountains of Idaho. There is a saw-mill on the Palouse river, at Palouse, the logs for which are floated down from about 9 miles above.

"SPOKANE COUNTY (8,500 square miles).—The portion of this county to the west of the mouth of the Spokane river is wholly destitute of trees, with the exception of the high point or plateau opposite to the mouth of the Okinakane river. Here there is a small area thinly wooded, probably with yellow pine and red fir. On the eastern side of the county spurs from the mountains bordering Cœur d'Alène lake enter between Rock creek and Spokane river, and are covered more or less densely with a growth of yellow pine, often small, with some Douglas spruce and tamarack in the ravines. There is a saw-mill at Rock creek supplied from its immediate neighborhood. Crossing Hangman's creek a scattered growth of pine appears upon the ridges between Deep creek and the Spokane river, and as far west as the head of Crab creek. Trees also border the Spokane river below the falls and to within a few miles of its mouth. The region between the Spokane and Little Spokane rivers is mostly a broad, open valley, the hills bordering it upon the north being very thinly wooded. There are two saw-mills at Spokane Falls, but the logs for them are floated down from near Cœur d'Alène lake.

"The total area more or less covered with trees may be estimated at from 400 to 500 square miles.

"STEVENS COUNTY (14,760 square miles).—This county is broken and mountainous throughout, but with no high ranges east of the Cascade mountains. The portion lying east of the upper Columbia and north of the Spokane river has several small prairies upon Chamokane creek and Colville river, and there is a narrow, open valley along the Columbia for 20 miles below the mouth of the Colville. The mountains are all low, the ridges most frequently thinly wooded or nearly bare, with the timber becoming denser in the ravines, especially northward. The most common tree is the yellow pine, but in the ravines red fir is frequent, with tamarack and lodge-pole pine. Near the Colville river were seen *Picea Engelmanni*, *Abies grandis*, small *Thuyas*, and fine specimens of *Pinus monticola*, as well as *Populus balsamifera*, *Betula papyracea*, and *Alnus* of considerable size. The hills bordering the Columbia above Old Fort Colville are treeless. The drift-wood brought down by the river is said to be chiefly cedar (*Thuja gigantea*).

"The Colville Indian reservation, lying between the Okinakane and the Columbia eastward, is comparatively little known, being crossed by but two trails, one leading directly westward from Old Fort Colville, the other following the Kettle river, and for much of the way not far distant from the British boundary. As seen from the Columbia and from the heights bordering the Okinakane, this portion appears to be more open and grassy than that east of the Columbia, and, especially toward the south, more like the bare plateau of Spokane county. Okinakane valley itself is narrow, with mainly a desert vegetation of sage-brush, *Purshia*, and other like representatives of the Great Basin flora, which seems to find here its only passageway northward to the British boundary. The hills eastward have thinly-scattered pines, which occasionally descend into the valley. The northern trail from Old Fort Colville shows the lower valley of Kettle river to be well wooded, but above, opening out into grassy prairies and bordered by grass-covered hills or with scattered yellow pine, red fir, and larch. Upon the more densely wooded ridges and ravines were also found *Picea Engelmanni*, *Abies subalpina*, *Pinus Murrayana*, and *Thuja*.

"The main ridge separating Kettle river from the Okinakane (about 5,000 feet high and 12 miles from the latter stream) was well grassed upon both sides with large *Picea*, *Pseudotsuga*, *Pinus ponderosa*, and *Larix* along the creeks upon the eastern side, and on the west the *Pinus ponderosa* only. The ridges above the Okinakane to the north appeared treeless, while the northern slopes of the nearer hills to the south were pretty well covered with underbrush. West of the Okinakane, between that river and the Methow, the country is much like that to the east—high and broken, with scattered patches of timber, which becomes more general toward the northern boundary. Upon the Methow and Similkameen creeks there are open, grassy valleys of considerable extent, but for 12 miles from the mouth of the Methow the hills close in upon it and are considerably wooded. The rest of the county, from the Methow to the Wenatchee, is occupied by spurs from the Cascade mountains, which reach the banks of the Columbia; these are exceedingly rugged and almost impassable, being seldom traversed, even by Indians. A foot-trail leads from the headwaters of the Methow over to the Skagit, and a trail which has been passable for horses crosses the ridges between the upper Chelan lake and the Wenatchee, but it is described by the Indians as dangerous and long disused by them. The whole region is probably for the most part well timbered except along the Columbia river, where the mountains for from 10 to 15 miles back are but scantily wooded, the pine (*Pinus ponderosa*) and red fir occasionally reaching to the river. Heavy timber is reported about the head of Chelan lake, commencing at about 15 miles from

the foot, mostly yellow pine, but also red fir, some *Larix*, and small *Thuja*. The outlet to this lake is through a deep cañon, and is obstructed by falls and rapids. The Wenatchee flows through a more open valley, and, at least in high water, could be used for floating timber to the Columbia. For 7 miles from its mouth the ridges on each side are only scantily wooded, but from that point the trees (yellow pine and red fir, mostly young) occupy the valley, and at 20 miles the thick timber begins—pine, fir, red fir, larch, white pine (*Pinus monticola*), and cedar, the white pine sometimes 4 feet through, the cedar not large.

"YAKIMA COUNTY (8,900 square miles).—Immediately south of the Wenatchee the highest of the eastern spurs of the Cascade mountains extends in a southeasterly direction to the Columbia, forcing that river to make a bend eastward. This spur has an altitude of about 5,000 feet, and its higher northern slopes, overlooking the mouth of the Wenatchee and eastward, are somewhat densely covered with pine, red fir, and larch. The southern slope, as seen from Ellensburg, appeared nearly bare. I crossed the ridge about 17 miles above the mouth of the Wenatchee and a few miles east of the high, exceedingly rocky, and snow-covered peaks called by McClellan 'Mount Stuart'. It was found mostly well wooded, but the trees not exceeding 1 or 2 feet in diameter, and usually small red fir and yellow pine, with at length some *Abies grandis* and *Pinus monticola*, rarely a small *Thuja*, on the higher rocky ridges small larches, and at the summit some *Pinus Murrayana*. The same trees were found on the southern descent, excepting the *Pinus monticola*. Large cottonwoods (*Populus trichocarpa*) occurred on the creeks. South of this range the spurs recede, leaving a comparatively level sage-brush region, wholly treeless, from 50 to 70 miles broad, between the Columbia and Yakima, and crossing the lower portion of the latter river.

"Below the mouth of the Schwank, which is at the head of what is known as 'Killitas valley', on the Yakima, the foot-hills of the Cascade mountains extend to the Yakima river, a distance of about 50 miles from the summit of the range; but the lower portions of these spurs are bare, or with only scattered pines on their northern slopes, and the chief reliance of the settlers for fencing and fuel is upon the aspens and cottonwoods bordering the streams. Following up the Yakima from the mouth of the Schwank, the valley for 10 or 12 miles is thinly timbered with pine and red fir. For 17 miles more there is some larch on the ridges, and in the bottoms some *Abies grandis*, and rarely a small *Thuja*. Timber and ties had been extensively cut here for the railroad and floated down the river. At this point the yellow pine and tamarack ceased, and a dense, heavy growth began and continued for most of the way to the summit (20 or 25 miles), consisting of red fir, hemlock, *Abies grandis* and *A. amabilis* (all these from 3 to 5 feet through and 200 feet high or more), *Pinus monticola* (18 inches through), and *Thuja* (2 feet in diameter). One spruce, not over 2½ feet through, had a height of 225 feet.

"In like manner, upon the Nachess river, the open sage-brush country extended about 10 or 12 miles from its mouth, with only cottonwood along the stream. Scattered pines then commence, with at length red fir, but it is some 25 or 30 miles more before heavy timber is reached. A small grove of oak (*Quercus Garryana*) is found at the mouth of the Schwank, the only point upon the Yakima where it occurs. It is also frequent along the Nachess for 3 or 4 miles, commencing at about 12 miles from its mouth, but small and rarely over 6 inches in diameter or 15 feet in height. In Satas valley it is abundant. Along the southern border of the county there is again a long spur extending east from mount Adams to within about 40 miles of the mouth of the Yakima. This spur has an altitude of about 1,500 feet, and is mostly covered with a scattered growth of yellow pine, red fir, and *Abies grandis*.

"The entire wooded area of the county may be estimated at about 4,500 square miles.

"KLIKITAT COUNTY (2,300 square miles).—The spur eastward from mount Adams, just spoken of, covers much of the northern portion of this county and affords a good supply of excellent timber. The area may be estimated at 750 square miles. The high ridge overlooking the Columbia from The Dalles eastward is perfectly bare of trees."

OREGON.

The heavy forest of western Washington territory extends through western Oregon. The most valuable timber tree of the region is the red or yellow fir (*Pseudotsuga Douglasii*), which forms fully seven-eighths of the forest. The tide-land spruce (*Picea Sitchensis*) abounds along the coast, and the red cedar (*Thuja gigantea*) and the hemlock (*Tsuga Mertensiana*) are common and of large size. South of Coos bay an important forest of Port Orford cedar (*Chamaecyparis Lawsoniana*), mixed with the red fir and the tide-land spruce, occurs.

The valleys of the Willamette, Umpqua, and Rogue rivers contain an open, scattered growth of white oak (*Quercus Garryana*), now gradually increasing by the recent growth of young trees protected from the fires which formerly swept every season through these prairie-like valleys. South of the Rogue River valley the sugar pine (*Pinus Lambertiana*), the chestnut oak (*Quercus densiflora*), and other trees of the California forest occur in sufficient numbers to add economic value to the forests of the state.

The bottom lands of western Oregon are lined with a continuous growth of cottonwoods of immense size, willows, maples, ashes, and gigantic alders; those in the southwest, near the coast, contain great bodies of splendid hard maple (*Acer macrophyllum*) and laurel (*Umbellularia Californica*).

East of the Cascade mountains the forests are confined to the mountain ranges; they are open, scattered, and generally composed of comparatively small trees.




COMPILED UNDER THE DIRECTION OF
C. S. SARGENT, SPECIAL AGENT
1883.

LEGEND.

1	Under 1 cord per acre
	From 1 to 2 " " "
	" 2 " 5 " " "
	" 5 " 10 " " "
	" 10 " 20 " " "
	" 20 " 30 " " "
	" 30 " 100 " " "
	Over 100 " " "
	Range of Port Orford Cedar. (Chamaecyparis Lawsoniana)

TENTH CENSUS OF THE UNITED STATES



Johns, Horn & Co. Lith.

201
202
203

204

205

The forests of Wasco county, on the western slope of the Cascade range, when above 3,000 feet elevation are important. The most valuable trees are the red fir, the yellow pine, and the larch. The eastern part of the county is covered with a light growth of pine, principally yellow pine.

The slopes of the Blue mountains in Umatilla and Union counties are covered with an open, stunted forest, consisting of red fir, yellow pine, larch, and, above 4,000 feet elevation, a heavier continuous growth of lodge-pole pine (*Pinus Murrayana*).

Lake county is destitute of timber except on the eastern slope of the Cascade mountains and the southern part of the county, which contain a light forest growth confined to the high ridges of the mountains, and principally composed of yellow pine.

Grant and Baker counties are treeless except in the northern part, where the Blue mountains are covered with a light, open growth composed chiefly of yellow pine, with some larch and scrub pine.

The forests of Oregon have suffered serious losses from forest fires. Along the Coast Range, from the Columbia river to Port Orford and through the entire length of the Cascade mountains, fires have raged nearly every summer since the first settlement of the state, destroying thousands of acres of noble fir, spruce, and cedar. Forests similar in composition to those destroyed soon spring up again and cover the burned surface, but the loss in material which the state has suffered in this way is incalculable.

Forest fires are increasing in frequency, especially west of the summit of the Cascade mountains. During the census year, however, only 132,320 acres of woodland were reported destroyed by fire, with an estimated loss of \$593,850. These fires were set by hunters, Indians, and by farmers clearing land.

The abundant spruce, cedar, cottonwood, ash, maple, and alder of western Oregon have developed flourishing industries. At Portland large quantities of ash, maple, and alder are manufactured into furniture, and cottonwood, spruce, and cedar supply numerous establishments engaged in the production of cooperage stock and all kinds of woodenware. The supply of this material is large and of excellent quality.

The principal centers of the lumber-manufacturing interests are at Portland, where fir, spruce, cottonwood, and hard woods are sawed for the local market, and at Empire City and Marshfield upon Coos bay. Port Orford cedar and red-fir lumber are manufactured here, and shipped by schooner to Portland, San Francisco, and Mexican and South American Pacific ports. The first mill was established upon Coos bay, at North Bend, 4 miles above Empire City, in 1853; other mills were soon built, and in 1854 the first shipment of Port Orford cedar was made to San Francisco. Great quantities of this timber have been cut, while fires have destroyed even more than the ax. The fire which raged through the forests of Coos bay for three months in the summer of 1867 destroyed cedar estimated to amount to between 200,000,000 and 300,000,000 feet of lumber. This tree, however, reproduces itself very rapidly, and after the forest has been burned over it is the first arborescent species to reappear, springing up generally in the third year.

The heaviest continuous body of Port Orford cedar now standing is on cape Gregory, extending south to and beyond the mouth of the Coquille river. It is about 20 miles long by an average width of 12 miles, and lies along the western slope of the foot-hills of the Coast Range, extending to within 3 miles of the coast. In this forest two-thirds of the trees are Port Orford cedar, the others tide-land spruce and a few red firs. There is great danger, however, that the Port Orford cedar, one of the most valuable trees of the American forest, will soon be exterminated as a source of lumber supply, so far as this generation is concerned.

The following notes upon the forests of Wasco, Umatilla, Union, Grant, and Baker counties, the only portion of the state visited by Mr. Watson, are extracted from his report:

"WASCO COUNTY (17,760 square miles).—The timber of this county is confined almost wholly to the steep eastern slopes of the Cascade range; the low spurs of the Blue mountains, which enter the county on the east, bordering John Day's river and southward, being only partially supplied with pines, etc. I know nothing about Walker's range and the Paulina mountains in the southwest, but they are probably low, with little or no wood. The trees of the Cascades are doubtless nearly the same as those to the north of the Columbia, the larch reaching to the headwaters of the Deschutes river, the most southern locality for it that I have seen mentioned.

"The total more or less wooded area may be estimated at from 2,500 to 3,000 square miles.

"UMATILLA COUNTY (6,100 square miles).—The Blue mountains occupy the southern and eastern borders of this county, and are the only source of timber. They are for the most part well wooded, especially in the ravines, the trees growing to a fair size, and consisting of yellow and scrub pine, spruce and balsam (*Abies subalpina* and *A. grandis*).

"The wooded area is about 1,500 square miles.

"UNION COUNTY (4,300 square miles).—This county has the main range of the Blue mountains on the west and north and the Cedar mountains on the east, separated by the valleys of the Grande Ronde and Wallowa rivers. A large portion of these mountains is well timbered, the amount decreasing toward the east.

"The wooded area may be estimated at about 2,000 square miles.

"GRANT COUNTY NORTH OF LATITUDE 44° (5,800 square miles).—This portion of the county is traversed by the valley of John Day's river, to the north and east of which lie the main ranges of the Blue mountains, which are to a considerable extent well wooded. The mountains to the south are low and probably scantily timbered.

"Fifteen hundred square miles is probably a liberal estimate for the wooded area.

"BAKER COUNTY NORTH OF LATITUDE 44° (3,800 square miles).—This section is bordered on the west by a high range of the Blue mountains, which is well timbered. The remainder is almost wholly without timber.

"The estimated wooded area of this county is 900 square miles."

CALIFORNIA.

The heavy forests of California are confined to the Coast Range, the eastern and western slopes of the Sierra Nevada, and the group of mountains joining these ranges in the northern part of the state. They extend from the Oregon boundary south to latitude 34° 30' north. The most important trees of the Coast Range forest are the redwood and the red fir. The tide-land spruce and the hemlock of the Northern Coast Forest extend as far south as cape Mendocino, although less generally multiplied and less valuable than in Oregon and Washington territory. The chestnut oak (*Quercus densiflora*), of which the bark is largely used in tanning, is still common in the coast forests of the northern part of the state. The most valuable forest of the western slope of the Sierra Nevada is confined to a belt between 4,000 and 8,000 feet elevation, consisting of the sugar pine (*Pinus Lambertiana*), the yellow pine, and the red fir. Small scattered groves of the big trees (*Sequoia gigantea*) stretch along the southern portion of this belt. The western slopes of these mountains below 4,000 feet elevation are more or less densely covered with various species of pine of little economic importance, and the broad valleys of the Sacramento and the San Joaquin, lying between the Coast Range and the Sierra Nevada, are covered, except at the south, with an open growth of oaks, often of immense size, although of little value except as fuel. The eastern slopes of the Sierra Nevada are covered with a heavy forest, in which yellow pines (*Pinus ponderosa* and *P. Jeffreyi*) are the prevailing and most important trees.


South of latitude 36° 30' the forests, both of the Sierra Nevada and of the Coast Range, become gradually less heavy and less valuable than those covering the mountains farther north. Two degrees still farther south they are open and scattered, and have little economic value. The pine and fir forests, however, which cover the upper slopes of the San Bernardino and San Jacinto ranges are important on account of their isolated position in a region destitute of tree covering, and supply a considerable local market with lumber.


The northeastern and nearly all the southern and southeastern portions of the state are almost entirely destitute of forest covering. Oaks and occasional pines and junipers are, however, dotted over the low mountains of southwestern California, and willows and cottonwoods line the banks of streams. Forests of pine crown the highest ridges of the Inyo and other mountain ranges, rising from the desert east of the Sierra Nevada, and arborescent yuccas (*Yucca brevifolia*) form upon the high Mohave plateau an open forest, more remarkable in the strangeness of its growth than in economic value.

The narrow belt of redwood which extends along the western slopes of the Coast Range from the bay of Monterey to the northern boundary of the state is the most important forest of similar extent now standing. Few trees equal the redwood in economic value. No other forest can compare with this in productive capacity, and no other great body of timber in North America is so generally accessible or so easily worked. Single trees capable of producing 75,000 feet of lumber are not uncommon, while a yield of from 1,000,000 to 2,000,000 feet of lumber per acre is by no means rare. The redwood has already been practically destroyed in the neighborhood of San Francisco bay, both north and south, and through the entire extent of this forest the trees most accessible to streams and railroads have been culled. Heavy bodies of redwood are still standing, however, in the Santa Cruz region, and in Humboldt county in the valleys of Eel and Mud rivers and Redwood creek. The largest number of mills engaged in the manufacture of redwood lumber are located upon Humboldt bay, principally at Eureka and Arcata. The logs which supply these mills are generally cut within a distance of 1 or 2 miles from the shores of the bay, to which they are hauled by teams, made into rafts, and towed to the mills. Attempts made to raft logs down the mountain streams watering the redwood forests have not been successful. The rivers flowing west from the California Coast Range are short and rapid. Floods following the winter rains are sudden and severe, breaking up rafts and driving the logs out to sea, or lodging them far from the banks. At periods of low water numerous bars close these rivers to the navigation of the enormous redwood logs. The general destruction of these forests must therefore be accomplished by means of short logging railroads specially constructed to bring logs to the mills. Such a road has been built along Mad river, and there are others either built or projected near Trinidad and at other points along the coast.

Besides the mills upon Humboldt bay, there are others devoted entirely to the manufacture of redwood lumber at Crescent City, in Del Norte county; Trinidad, Rohnerville, and Bridgeville, in Humboldt county; Westport, Kibesillah, Albion, Little River, Caspar, Mendocino, Cuffey's Cove, Punta Arena, and Gualala, in Mendocino county; Duncan's mills, in Sonoma county; and at Santa Cruz.

Redwood lumber is principally shipped by schooner to San Francisco, the great point of lumber distribution upon the Pacific coast, and also direct by water to Wilmington, San Diego, and other ports of southern California, and to Mexico and South America.





DENSITY OF FORESTS

COMPILED UNDER THE DIRECTION OF
C. S. SARGENT, SPECIAL AGENT
1883.

Scale
0 10 20 30 40 50 60 70 80 90 100 110 120 130 140 150 160 170 180 190 200 210 220 230 240 250 260 270 280 290 300 310 320 330 340 350 360 370 380 390 400 410 420 430 440 450 460 470 480 490 500 510 520 530 540 550 560 570 580 590 600 610 620 630 640 650 660 670 680 690 700 710 720 730 740 750 760 770 780 790 800 810 820 830 840 850 860 870 880 890 900 910 920 930 940 950 960 970 980 990 1000


LEGEND.

I	Under record per acre
II	From 1 to 2 "
III	" 2 " 5 "
IV	" 5 " 10 "
V	" 10 " 20 "
VI	" 20 " 50 "
VII	" 50 " 100 "
VIII	" 100 " 200 "
IX	Over 200 "

The following estimates of the amount of accessible redwood standing May 31, 1880, were prepared by Mr. E. L. Allen, secretary of the Redwood Manufacturers' Association of San Francisco. They embrace only such portions of the forest as can be reached by water, or may in the future be penetrated by railroads, and do not include the small, isolated bodies of timber growing in inaccessible cañons:

REDWOOD (*Sequoia sempervirens*).

Regions.	Feet, board measure.
From the Oregon boundary to the mouth of Redwood creek	800,000,000
From the mouth of Redwood creek to the mouth of Mad river	9,000,000,000
From the mouth of Mad river to the mouth of Eel river	2,145,000,000
From the mouth of Eel river to the mouth of Mattole river	4,450,000,000
From the mouth of Mattole river to the mouth of Cottonwood creek	200,000,000
From the mouth of Cottonwood creek to the mouth of Russian river	7,680,000,000
In the Santa Cruz region	1,550,000,000
Total	25,825,000,000
Estimated cut for the census year ending May 31, 1880:	
Sawed lumber	125,390,000
Shingles and shakes	25,380,000
Split railroad ties	23,265,000
Posts, etc	12,600,000
Total	186,635,000



No estimate of the amount of pine and fir lumber standing in the state is now possible, and none has been attempted. An enormous amount of pine of excellent quality, both white and yellow, is contained in the sierra forests. These forests have been invaded by the lumberman at only a few points; their inaccessibility and the cost of getting to market the lumber manufactured in these mountains have thus far preserved them, and these sierra forests, if protected from fire, will serve as a reservoir from which the whole Pacific coast can draw its lumber supply long after its more accessible forests have disappeared.

The forests of California suffer seriously by fire; during the census year 356,815 acres of woodland were reported thus destroyed, with an estimated loss of \$440,750. These fires were set by careless hunters, prospectors, and by farmers in clearing land. Great injury, every year becoming greater, is inflicted on the mountain forests by stockmen starting fires to improve the herbage of the alpine pastures. These fires destroy undergrowth and young trees, and often consume great quantities of valuable timber, which does not grow again upon these exposed mountain slopes.

PASTURAGE OF MOUNTAIN FORESTS.

The permanence of the mountain forests of California is severely endangered, moreover, by the immense herds of sheep, cattle, and horses driven into the mountains every year, at the commencement of the dry season, to graze. From the foot-hills to the highest alpine meadows every blade of herbage and every seedling shrub and tree is devoured. Young trees are barked and ruined, and only the most rigid and thorny chaparral shrubs are able to resist the attacks of these ravenous herds. The sharp hoofs of sheep winding around the steep acclivities tread out the roots of grasses and other perennial plants and loosen the surface of the stony soil, which, deprived of the protection of its vegetable covering, is gradually washed into the valleys, choking the bottoms of streams and preparing the way for the disastrous torrents which must follow the destruction of the sierra forests; and the destruction of these forests is certain, if the practice of using them indiscriminately as sheep pastures is continued. The life of any forest in which all young trees are destroyed as soon as they appear above the surface of the soil is limited to the life of the fully grown individuals which compose it. A period of unusual climatic conditions, the demand of an increased population for lumber, or the now unforeseen attacks of some insect enemy may at any time sweep away the old trees of the sierra forests. There are no young trees growing to replace them, and it is doubtful if the forest could ever regain its foothold upon the steep and exposed slopes of these mountains once entirely stripped of the protection of their present covering of trees.

The sheep which threaten the destruction of the sierra forests threaten also the agricultural prosperity of the state; the streams heading in the sierras and watering the great interior valleys of California are protected in their flow by the forests growing about their upper sources. If these forests are destroyed, and the protection to the surface of the ground which they afford removed, the immense accumulation of the winter's snows must melt suddenly in the spring; brooks will become torrents, sweeping with irresistible force gravel and stones from the mountain sides down into the valleys below, and burying rich bottom lands in ruin. And this is not the only danger which must follow the destruction of these forests. If the snow which supplies the mountain streams melts slowly, a steady flow of water will be maintained late into the season; if, on the other hand, the snow melts suddenly and rapidly during the first warm days of spring, the unnatural flow of water in the stream must be followed by

its equally sudden disappearance, and the torrent will suddenly diminish to a slender brook or entirely disappear. Irrigation, without which agriculture in a large part of the Pacific region is impossible, is dependent upon the constant and steady flow of streams formed by melting snow, and as the forests which cover the mountain sides are essential to prevent the sudden melting of snow, their preservation is necessary for successful irrigation on any large or comprehensive scale.

The forests of California suffer from wasteful methods of cutting. Only the best and most accessible young trees are cut; often a noble pine capable of producing 25,000 or 30,000 feet of lumber is felled, a few split shingles made from the butt-cut, and the rest of the tree left to rot upon the ground. The preference of the railroad companies of the state for split rather than sawed redwood ties causes an immense and needless waste of this valuable timber. A great amount of material under the most favorable conditions is wasted in splitting out the ties, and when trees after being cut are found to split badly from any defect in the grain they are abandoned and left to waste.

The forests of California, unlike those of the Atlantic states, contain no great store of hard woods. The oaks of the Pacific forests, of little value for general mechanical purposes, are unfit for cooperage stock. No hickory, gum, elm, or ash of large size is found in these forests. California produces no tree from which a good wine cask or wagon wheel can be made. The cooperage business of the state, rapidly increasing with the development of grape culture, is entirely dependent upon the forests of the Atlantic region for its supply of oak. Woodenware and small cooperage stock are manufactured in large quantities, however, from cottonwood, spruce, alder, and red and white fir. Wine-butts and water-tanks are universally made from redwood, which is probably unsurpassed for such purposes.

The large tanning industry of the state consumes, in preference to all other material, large quantities of the bark of the chestnut oak (*Quercus densiflora*), once a common tree in the forests of the northern Coast ranges, but now becoming scarce and in danger of speedy extermination.

The principal centers of lumber manufacture outside of the redwood belt are situated along the line of the Central Pacific railroad, upon both flanks of the Sierra Nevada mountains, in Butte, Tehama, and Mono counties, and in the San Bernardino mountains. Lumber manufactured upon the eastern slope of the Sierra Nevadas is largely shipped eastward by rail to supply Nevada and Utah. The product of the mills situated west of the mountains is largely sent to San Francisco for distribution, or direct by rail to the mining centers of southern Arizona and New Mexico.

ALASKA.

Little is known to me of the present condition or productive capacity of the forests of Alaska. Their distribution, as shown on the forest map of North America, is based upon notes made by Mr. Ivan Petroff, a special agent of the Census Office, who has traced the timber limits of the territory, aided by Mr. C. W. Nelson, of the Smithsonian Institution, by whom the northern limits of the spruce forest are laid down.

The forests of the territory of any commercial value are confined to the islands and Coast ranges east and south of Prince William sound. The most valuable tree of this region is the Sitka cedar (*Chamaecyparis Nutkaensis*). The hemlock, the tide-land spruce, and the red cedar (*Thuja gigantea*) attain here also a considerable size. The importance, however, of these forests, both in extent and in the value of the timber they contain, has generally been greatly exaggerated. The Coast Forest north of the fiftieth degree of latitude rapidly diminishes in density and quality, and there is nothing in the climate or soil of Alaska to produce a forest growth more valuable than that covering the Coast ranges of British Columbia.


A few saw-mills of small capacity are located at different points in southeastern Alaska to supply the local demand for coarse lumber. Alaska is, however, largely supplied with lumber from Puget sound. The treeless Shumagin and Aleutian islands and the southern settlements of the peninsula are supplied with fire-wood brought from other portions of the territory.



Julius Bien & Co. lith.

Scale

0 10 20 30 40 50 60 70 80 90 100 MILES



GENERAL INDEX.

[NOTE.—In this index the names of families are indicated by "SMALL CAPS", of species by "Roman" type, and of synonyms by "Italics".]

A.		Page.		Page.
<i>Abies</i>	11, 12, 16, 573		<i>Abies magnifica</i> (<i>Abies nobilis</i>).....	214
<i>Abies alba</i> (<i>Picea alba</i>).....	204		<i>Abies Mariana</i>	202
<i>Abies alba</i> (<i>Picea Engelmanni</i>).....	205		<i>Abies Marylandica</i>	203
<i>Abies alba</i> (<i>Picea nigra</i>).....	203		<i>Abies Menziesii</i> (<i>Picea pungens</i>).....	205
<i>Abies alba</i> , var. <i>arctica</i>	204		<i>Abies Menziesii</i> (<i>Picea Sitchenensis</i>).....	206
<i>Abies alba</i> , var. <i>caerulea</i>	204		<i>Abies Menziesii</i> <i>l'arryana</i>	205
<i>Abies Albertiana</i>	208		<i>Abies Mertensiana</i>	207
<i>Abies amabilis</i>	213, 250, 255, 259, 346, 412, 478, 573, 576		<i>Abies microcarpa</i>	215
<i>Abies amabilis</i> (<i>Abies concolor</i>).....	212		<i>Abies mucronata</i>	209
<i>Abies amabilis</i> (<i>Abies grandis</i>).....	212		<i>Abies mucronata palustris</i>	209
<i>Abies amabilis</i> (<i>Abies magnifica</i>).....	214		<i>Abies nigra</i> (<i>Picea Engelmanni</i>).....	205
<i>Abies Americana</i>	203		<i>Abies nigra</i> (<i>Picea nigra</i>).....	208
<i>Abies arctica</i> (<i>Picea alba</i>).....	204		<i>Abies nigra</i> , var. <i>rubra</i>	203
<i>Abies arctica</i> (<i>Picea nigra</i>).....	203		<i>Abies nobilis</i>	214, 250, 255, 259, 346, 412, 478, 573
<i>Abies aromatica</i>	212		<i>Abies nobilis</i> (<i>Abies magnifica</i>).....	214
<i>Abies balsamea</i>	210, 251, 255, 259, 346, 412, 478		<i>Abies nobilis robusta</i>	214
<i>Abies balsamea</i> (<i>Abies concolor</i>).....	212		<i>Abies Parsonsii</i>	212
<i>Abies balsamea</i> , var. <i>Fraseri</i>	210		<i>Abies Pattoni</i>	208
<i>Abies balsamifera</i>	211		<i>Abies Pattoniana</i>	208
<i>Abies bifolia</i>	211		<i>Abies Pattonii</i> (<i>Tsuga Mertensiana</i>).....	208
<i>Abies bracteata</i>	8, 213, 249, 346		<i>Abies Pattonii</i> (<i>Tsuga Pattoniana</i>).....	208
<i>Abies Bridgesii</i>	207		<i>Abies pendula</i>	215
<i>Abies campylocarpa</i>	214		<i>Abies religiosa</i>	185
<i>Abies Canadensis</i> (<i>Picea alba</i>).....	204		<i>Abies rubra</i>	208
<i>Abies Canadensis</i> (<i>Tsuga Canadensis</i>).....	206		<i>Abies rubra</i> , var. <i>arctica</i>	203
<i>Abies Canadensis</i> (<i>Tsuga Mertensiana</i>).....	207		<i>Abies rubra</i> , var. <i>caerulea</i>	204
<i>Abies Caroliniana</i>	207		<i>Abies Sitchenensis</i>	206
<i>Abies caerulea</i>	204		<i>Abies speciosa</i> (<i>Tsuga Caroliniana</i>).....	207
<i>Abies concolor</i>	9, 212, 251, 255, 259, 264, 346, 412, 478, 570		<i>Abies subalpina</i>	211, 251, 255, 259, 264, 346, 412, 478, 564-567, 572, 573, 575, 577
<i>Abies denticulata</i>	203		<i>Abies subalpina</i> , var. <i>fallax</i>	211
<i>Abies Douglasii</i>	209		<i>Abies taxifolia</i> (<i>Pseudotsuga Douglasii</i>).....	209
<i>Abies Douglasii</i> , var. <i>macrocarpa</i>	210		<i>Abies taxifolia</i> (<i>Tsuga Mertensiana</i>).....	208
<i>Abies Douglasii</i> , var. <i>taxifolia</i>	209		<i>Abies trigona</i>	206
<i>Abies Engelmanni</i>	205		<i>Abies venusta</i>	213
<i>Abies Engelmanni glauca</i>	205		<i>Abies Williamsontii</i>	208
<i>Abies falcata</i>	206		<i>Acacia</i>	6, 10-12, 14
<i>Abies Fraseri</i>	210, 251, 255, 259, 346, 412, 478		<i>Acacia Bahamensis</i>	64
<i>Abies Gordoniana</i>	212		<i>Acacia Berlandieri</i>	63, 248
<i>Abies grandis</i>	7, 212, 251, 255, 259, 346, 412, 478, 565, 566, 572, 573, 575-577		<i>Acacia biceps</i>	62
<i>Abies grandis</i> (<i>Abies amabilis</i>).....	213		<i>Acacia ceculenta</i>	63
<i>Abies grandis</i> (<i>Abies concolor</i>).....	212		<i>Acacia frondosa</i>	62
<i>Abies grandis</i> (<i>Abies subalpina</i>).....	211		<i>Acacia glauca</i>	63
<i>Abies grandis</i> , var. <i>densiflora</i>	213		<i>Acacia, Green-bark</i>	60, 280, 362, 426
<i>Abies grandis</i> , var. <i>Lowiana</i>	212		<i>Acacia Greggii</i>	12, 63, 249, 282, 362, 426
<i>Abies heterophylla</i>	207		<i>Acacia latistylus</i>	64
<i>Abies Hookeriana</i>	208		<i>Acacia leucoccephala</i>	63
<i>Abies Hudsonica</i>	211		<i>Acacia pulverulenta</i>	63
<i>Abies lasiocarpa</i> (<i>Abies concolor</i>).....	212		<i>Acacia tephroleba</i>	63
<i>Abies lasiocarpa</i> (<i>Abies subalpina</i>).....	211		<i>Acacia, Three-thorned</i>	59, 280, 360, 426
<i>Abies laxa</i>	204		<i>Acacia Wrightii</i>	63, 249, 282
<i>Abies Lowiana</i>	212		<i>Acer</i>	10-12
<i>Abies macrocarpa</i>	210		<i>Acer barbatum</i> (<i>Acer glabrum</i>).....	48
<i>Abies magnifica</i>	214, 250, 255, 259, 346, 412, 478		<i>Acer barbatum</i> (<i>Acer saccharinum</i>).....	48
			<i>Acer Californicum</i>	51

	Page.		Page.
<i>Acer Canadense</i>	48	Alabama, pine region of the Coosa in.....	529
<i>Acer Caroliniana</i>	50	Alabama, production of naval stores in.....	517, 527, 529, 530
<i>Acer circinatum</i>	47, 249, 253, 256, 274, 358, 422	Alabama, rank of, according to value of lumber products.....	487
<i>Acer coccineum</i>	50	Alabama, remarks by Dr. Charles Mohr on the forests and the turpentine industry of.....	525-530
<i>Acer dasycarpum</i>	49, 250, 253, 256, 276, 358, 424	Alabama, statistics of forests, forest fires, standing timber, etc., in.....	491, 524-530
<i>Acer Douglasii</i>	48	Alabama, tabular statement of the amount of Long- and Short-leaved Pine standing in the forests of, May 31, 1880.....	524
<i>Acer Drummondii</i>	50	Alabama, the forests of Pike county in.....	528
<i>Acer eriocarpum</i>	49	Alabama, the Maritime Pine Region in.....	526
<i>Acer glabrum</i>	47, 250, 274	Alaska, statistics of forests, forest fires, standing timber, etc., in.....	580
<i>Acer glaucum</i>	50	Alaska, the Northern Coast Forest in.....	580
<i>Acer grandidentatum</i>	13, 48, 249, 276	Albany, New York, rank of, as a lumber market.....	503
<i>Acer macrophyllum</i>	47, 250, 253, 256, 259, 274, 358, 422, 576	Alder (<i>Alnus oblongifolia</i>).....	163, 326, 394, 460
<i>Acer montanum</i>	46	Alder (<i>Alnus rhombifolia</i>).....	163, 326, 394, 460
<i>Acer Negundo</i>	51	Alder (<i>Alnus rubra</i>).....	163, 326, 392, 416, 458
<i>Acer nigrum</i>	49	Alder, Black (<i>Alnus incana</i>).....	165, 326, 394, 460
<i>Acer palmatum</i>	47	Alder, Black (<i>Alnus serrulata</i>).....	161, 326
<i>Acer parviflorum</i>	46	Alder, Hoary.....	165, 326, 394, 460
<i>Acer Pennsylvanicum</i>	46, 250, 274	Alder, Seaside.....	162, 326, 458
<i>Acer Pennsylvanicum</i> (<i>Acer spicatum</i>).....	46	Alder, Smooth.....	164, 326
<i>Acer rubrum</i>	50, 250, 253, 256, 259, 276, 358, 424	Alder, Speckled.....	165, 326, 394, 460
<i>Acer rubrum</i> , var. <i>Drummondii</i>	50, 250, 276	Algaroba.....	62, 265, 280, 362, 476
<i>Acer rubrum</i> , var. <i>pallidum</i>	49	<i>Algarobia glandulosa</i>	61
<i>Acer saccharinum</i>	48, 249, 253, 256, 259, 276, 358, 422	<i>Alnus</i>	11, 12, 16, 575
<i>Acer saccharinum</i> (<i>Acer dasycarpum</i>).....	49	<i>Alnus Americana</i> (<i>Alnus incana</i>).....	164
<i>Acer saccharinum</i> (<i>Acer saccharinum</i> , var. <i>nigrum</i>).....	49	<i>Alnus Americana</i> (<i>Alnus serrulata</i>).....	164
<i>Acer saccharinum</i> , var. <i>nigrum</i>	49, 249, 253, 256, 259, 276, 350, 358, 424	<i>Alnus arguta</i>	162
<i>Acer saccharum</i>	48	<i>Alnus Canadensis</i>	164
<i>Acer sanguineum</i>	50	<i>Alnus genuina</i>	164
<i>Acer spicatum</i>	46, 250, 274	<i>Alnus glauca</i>	164
<i>Acer striatum</i>	46	<i>Alnus glutinosa</i> (<i>Alnus rubra</i>).....	163
<i>Acer tripartitum</i>	48	<i>Alnus glutinosa</i> , var. <i>rugosa</i>	164
<i>Acer virgatum</i>	47	<i>Alnus glutinosa</i> , var. <i>serrulata</i> (<i>Alnus rhombifolia</i>).....	163
<i>Achras cuneifolia</i>	103	<i>Alnus glutinosa</i> , var. <i>serrulata</i> (<i>Alnus serrulata</i>).....	164
<i>Achras mammosa</i>	103	<i>Alnus hybrida</i>	164
<i>Achras pallida</i>	101	<i>Alnus incana</i>	12, 164, 250, 326, 394, 460
<i>Achras salicifolia</i>	101	<i>Alnus incana</i> (Regel).....	164
<i>Achras serrata</i>	69	<i>Alnus incana</i> , var. <i>glauca</i> (<i>Alnus incana</i>).....	164
<i>Achras Zapotilla</i> , var. <i>parviflora</i>	103	<i>Alnus incana</i> , var. <i>glauca</i> (<i>Alnus incana</i> , var. <i>virescens</i>).....	165
Actual fuel value of some of the more important woods of the United States.....	350-353	<i>Alnus incana</i> , var. <i>rubra</i>	163
<i>Adelia acuminata</i>	112	<i>Alnus incana</i> , var. <i>virescens</i>	165
<i>Æsculus</i>	10-13	<i>Alnus Japonica</i>	162
<i>Æsculus Californica</i>	43, 250, 253, 256, 274, 422	<i>Alnus maritima</i>	162, 250, 326, 458
<i>Æsculus carnea</i>	42	<i>Alnus maritima typica</i>	162
<i>Æsculus discolor</i>	43	<i>Alnus oblongata</i> (<i>Alnus maritima</i>).....	162
<i>Æsculus echinata</i>	42	<i>Alnus oblongata</i> (<i>Alnus serrulata</i>).....	164
<i>Æsculus flava</i>	43, 250, 274	<i>Alnus oblongifolia</i>	163, 250, 255, 258, 326, 394, 460
<i>Æsculus flava</i> , var. <i>purpurascens</i>	43	<i>Alnus obtusifolia</i>	164
<i>Æsculus glabra</i>	42, 250, 253, 256, 274, 422	<i>Alnus Oregana</i>	163
<i>Æsculus Hippocastanum</i> , var. <i>glabra</i>	42	<i>Alnus rhombifolia</i>	163, 250, 255, 258, 326, 394, 460
<i>Æsculus Hippocastanum</i> , var. <i>Ohioensis</i>	42	<i>Alnus rubra</i>	163, 250, 255, 258, 262, 326, 392, 416, 458
<i>Æsculus Hippocastanum</i> , var. <i>pallida</i>	42	<i>Alnus rubra</i> (<i>Alnus serrulata</i>).....	164
<i>Æsculus hybrida</i>	43	<i>Alnus rugosa</i>	164
<i>Æsculus lutea</i>	43	<i>Alnus serrulata</i>	164, 250, 326
<i>Æsculus neglecta</i>	43	<i>Alnus serrulata</i> (Regel).....	164
<i>Æsculus octandra</i>	43	<i>Alnus serrulata genuina</i>	164
<i>Æsculus Ohioensis</i>	42	<i>Alnus serrulata</i> , var. <i>latifolia</i>	164
<i>Æsculus pallida</i>	42	<i>Alnus serrulata</i> , var. <i>macrophylla</i>	164
<i>Æsculus Pavia</i> , var. <i>discolor</i>	43	<i>Alnus serrulata</i> , var. <i>oblongata</i>	164
<i>Æsculus Watsoniana</i>	42	<i>Alnus serrulata</i> , var. <i>oblongifolia</i>	163
<i>Agastianis secundiflora</i>	57	<i>Alnus serrulata</i> , var. <i>rugosa</i> (<i>Alnus incana</i> , var. <i>virescens</i>).....	165
<i>Ageria Cassena</i>	36	<i>Alnus serrulata</i> , var. <i>rugosa</i> (<i>Alnus rhombifolia</i>).....	163
<i>Ageria geminata</i>	36	<i>Alnus serrulata</i> , var. <i>rugosa</i> (<i>Alnus serrulata</i>).....	164
<i>Ageria heterophylla</i>	35	<i>Alnus serrulata</i> , var. <i>vulgaris</i>	164
<i>Ageria obovata</i>	35	<i>Alnus undulata</i>	164
<i>Ageria opaca</i>	35	Amelanchier.....	10, 12, 14
<i>Ageria palustris</i>	35	Amelanchier <i>alnifolia</i>	85
Alabama, central, pine belt of.....	529	Amelanchier <i>Bartramiana</i>	84
Alabama, cypress swamps of the Tensas river in.....	525-527	Amelanchier <i>Botrysapium</i>	84
Alabama, eastern, forests of the Chattahoochee, mixed forest growth, etc., in.....	527, 528	Amelanchier <i>Canadensis</i>	84, 249, 253, 256, 364, 436
Alabama, forests of the Tennessee valley in.....	528, 529	Amelanchier <i>Canadensis</i> , var. <i>oblongifolia</i>	84
Alabama, lumber industry and manufactures from wood in.....	486, 487, 524, 525	Amelanchier <i>Canadensis</i> , var. <i>oligocarpa</i>	84
Alabama, Mobile the principal center of wood manufacture in.....	525	Amelanchier <i>Canadensis</i> , var. <i>rotundifolia</i>	84
Alabama, pine forests of Baldwin county in.....	527	Amelanchier <i>intermedia</i>	85

	Page.		Page.
<i>Amelanchier oblongifolia</i>	85	Arizona, lumber industry and manufactures from wood in	486, 487, 569
<i>Amelanchier ovalis</i> (<i>Amelanchier Canadensis</i>).....	84	Arizona, rank of, according to value of lumber products.....	487
<i>Amelanchier ovalis</i> (<i>Amelanchier Canadensis</i> , var. <i>oblongifolia</i>).....	85	Arizona, statistics of forests, forest fires, standing timber, etc., in	491, 508, 569
<i>Amelanchier spicata</i>	85	Arkansas, lumber industry and manufactures from wood in	486, 487, 544
<i>Amelanchier Wangenheimiana</i>	84	Arkansas, rank of, according to value of lumber products.....	487
American Crab.....	72, 284, 364, 428	Arkansas, statistics of forests, forest fires, standing timber, etc., in	491, 543, 544
American Crab Apple.....	72, 284	Arkansas, tabular statement of the amount of Short-leaved Pine standing in the forests of, May 31, 1880.....	544
American Elm.....	123, 304, 350, 374, 414, 440	<i>Aronia arborea</i>	84
American Holly.....	35, 270, 356, 420	<i>Aronia arbutifolia</i>	83
American Linden.....	27, 268, 350, 356, 418	<i>Aronia Botryopium</i>	84
Amount and value of charcoal used for domestic and manufacturing purposes during the census year.....	489	<i>Aronia cordata</i>	84
Amount and value of wood used as fuel in manufactures, etc., during the census year.....	489	<i>Aronia ovalis</i>	85
Amount and value of wood used for domestic purposes during the census year.....	489	Arrowwood.....	38, 272
<i>Amyris</i>	10, 13	<i>Asagraea spinosa</i>	55
<i>Amyris cymosa</i>	33	Ash.....	107, 296, 298, 370, 434
<i>Amyris dyatripa</i>	33	Ash and specific gravity of woods.....	248-251
<i>Amyris clemifera</i>	31	Ash, Black.....	111, 298, 372, 438
<i>Amyris Floridana</i>	33	Ash, Blue.....	111, 298, 370, 436
<i>Amyris maritima</i>	33	Ash, Green.....	109, 298, 370, 436
<i>Amyris Plumieri</i>	33	Ash, Ground.....	111, 298, 372, 438
<i>Amyris sylvatica</i>	33, 249, 253, 256, 270, 356, 420	Ash, Hoop.....	111, 298, 372, 438
ANACARDIACEÆ.....	52-54, 253, 256, 276, 360, 424	Ash, Mountain (<i>Pyrus Americana</i>).....	73, 284, 428
<i>Ananomia punctata</i>	88	Ash, Mountain (<i>Pyrus sambucifolia</i>).....	74, 284, 364, 428
Anaqua.....	114, 300, 372, 438	Ash, Oregon.....	111, 298, 372, 438
Andromeda.....	11, 15	Ash, Prickly (<i>Xanthoxylum Americanum</i>).....	29, 268
<i>Andromeda arborea</i>	98	Ash, Prickly (<i>Xanthoxylum Clava-Herculis</i>).....	30, 270, 356, 410
<i>Andromeda arborescens</i>	98	Ash, Red.....	109, 296, 370, 436
<i>Andromeda ferruginea</i>	96, 249, 254, 257, 292, 368, 432	Ash, Sea.....	30, 270, 356, 420
<i>Andromeda ferruginea</i> , var. <i>arborescens</i>	96	Ash, specific gravity, and weight per cubic foot of dry specimens of the woods of the United States.....	266-349
<i>Andromeda ferruginea</i> , var. <i>fruticosa</i>	96	Ash, Wafer.....	31, 270
<i>Andromeda plumata</i>	37	Ash, Water.....	110, 298, 370, 436
<i>Andromeda rhomboidalis</i>	96	Ash, White.....	107, 296, 350, 370, 436
<i>Andromeda rigida</i>	96	Ash, Yellow.....	57, 278, 300, 426
Anona.....	10, 13	Ash-leaved Maple.....	51, 276, 360, 424
<i>Anona glabra</i>	23	Asimina.....	10, 11, 13
<i>Anona laurifolia</i>	23, 250, 253, 256, 266, 354, 418	<i>Asimina campaniflora</i>	23
<i>Anona pendula</i>	23	<i>Asimina triloba</i>	23, 250, 253, 256, 266, 354, 418
<i>Anona speciosa</i> (<i>Anona laurifolia</i>).....	23	Asp. Quaking.....	171, 328, 350, 394, 460
<i>Anona triloba</i>	23	Aspects, economic, of the forests of the United States.....	483-560
ANONACEÆ.....	23, 253, 256, 260, 354, 418	Aspen.....	171, 328, 350, 394, 460
<i>Anonymos aquatica</i>	124	Atlantic Plain, Deciduous Forest of the Mississippi Basin and the.....	4
<i>Anthomeles festalis</i>	82	Atlantic region, the.....	3-6
<i>Anthomeles Douglasii</i>	75	Atlantic region, western third of the, remarks on.....	4
<i>Anthomeles flava</i>	82	Average number of hands employed in the lumbering industry of the United States.....	486
<i>Anthomeles glandulosa</i>	82	Average size and product of saw-mills in each state and territory.....	488
<i>Anthomeles rotundifolia</i>	77	Avicennia.....	11, 15
<i>Anthomeles turbinata</i>	82	<i>Avicennia nitida</i>	117, 249, 302
Ants' Wood.....	103, 294, 368, 434	<i>Avicennia oblongifolia</i>	117
Apple, American Crab.....	72, 284	<i>Avicennia tomentosa</i>	117
Apple, Custard.....	23, 266, 354, 418		
Apple Haw.....	82, 284, 364, 430		
Apple, Oregon Crab.....	73, 284		
Apple, Pond.....	23, 266, 354, 418		
Apple, Seven-year.....	95, 290		
Apple, Southern Crab.....	72, 284		
Arbol de Hierro.....	56, 278, 360, 426		
Arbor-vitæ.....	176, 330, 350, 396, 462		
Arbutus.....	11, 12, 15		
<i>Arbutus laurifolia</i>	97		
<i>Arbutus macrophylla</i>	97		
<i>Arbutus Menziesii</i>	97, 249, 254, 257, 292, 368, 432		
<i>Arbutus Menziesii</i> (<i>Arbutus Texana</i>).....	97		
<i>Arbutus Menziesii</i> (<i>Arbutus Xalapensis</i>).....	97		
<i>Arbutus procera</i>	97		
<i>Arbutus Texana</i>	97, 249, 292		
<i>Arbutus varians</i>	97		
<i>Arbutus Xalapensis</i>	97, 249, 254, 257, 292, 368, 434		
<i>Arbutus Xalapensis</i> (<i>Arbutus Texana</i>).....	97		
Ardisia.....	11, 15		
Ardisia Pickeringia.....	100, 249, 292		
Areas burned over and value of property destroyed by forest fires during the census year (see, also, under state headings).....	491, 492		

	Page.		Page.
Bay, Loblolly	25, 265, 268, 354, 418	BIGNONIACEÆ	115, 116, 254, 257, 300, 372, 438
Bay, Red	118, 302, 372, 438	Bilsted	86, 288, 350, 364, 414, 430
Bay, Rose	90, 292, 368, 434	Birch, Black (<i>Betula lenta</i>)	162, 324, 392, 458
Bay, Sweet	20, 266, 354, 414, 418	Birch, Black (<i>Betula occidentalis</i>)	160, 324, 392, 458
Bay, Tan	25, 265, 268, 354, 418	Birch, Canoe	160, 324, 350, 392, 458
Bay Tree, California	120, 302, 374, 440	Birch, Cherry	162, 324, 392, 458
Bay, White	20, 266, 354, 414, 418	Birch, Gray (<i>Betula alba</i> , var. <i>populifolia</i>)	159, 324, 350, 392, 458
Bayberry	136, 312, 380, 446	Birch, Gray (<i>Betula lutea</i>)	161, 324, 392, 458
Bean, Indian	115, 300, 372, 438	Birch, Mahogany	162, 324, 392, 458
Bean Tree	115, 300, 372, 438	Birch, Old-field	159, 324, 350, 392, 458
Bearberry	41, 272, 358, 422	Birch, Paper	160, 324, 350, 392, 458
Bearwood	41, 272, 358, 422	Birch, Red	161, 324, 392, 458
Beaver Tree	20, 266, 354, 414, 418	Birch, River	161, 324, 392, 458
Beech	158, 322, 350, 390, 450	Birch, Sweet	162, 324, 392, 458
Beech, Blue	150, 322, 392, 456	Birch, West-Indian	33, 270, 350, 356, 420
Beech, Water (<i>Carpinus Caroliniana</i>)	150, 322, 392, 456	Birch, White (<i>Betula alba</i> , var. <i>populifolia</i>)	159, 324, 350, 392, 458
Beech, Water (<i>Platanus occidentalis</i>)	129, 306, 350, 376, 442	Birch, White (<i>Betula papyrifera</i>)	160, 324, 350, 392, 458
Beefwood	117, 302, 372, 438	Birch, Yellow	161, 324, 392, 458
Bee Tree	27, 268, 356, 418	Bishop's Pine	200, 340, 404, 472
Behavior of selected woods of the United States under transverse strain: specimens eight centimeters square	414-417	Bitter Nut	135, 310, 350, 380, 446
Behavior of the principal woods of the United States under compression	418-481	Bitter Pecan	136, 310, 350, 380, 414, 446
Behavior of the principal woods of the United States under transverse strain	354-415	Black Alder (<i>Alnus incana</i>)	105, 328, 394, 460
<i>Benthamedia florida</i>	90	Black Alder (<i>Alnus serrulata</i>)	164, 326
<i>Betula</i>	11, 12, 15	Black and White Spruces the characteristic trees of the Northern Forest	8
<i>Betula acuminata</i>	159	Black Ash	111, 298, 372, 438
<i>Betula alba</i> , subspecies <i>commutata</i>	160	Black Birch (<i>Betula lenta</i>)	162, 324, 392, 458
<i>Betula alba</i> , subspecies <i>occidentalis typica</i>	160	Black Birch (<i>Betula occidentalis</i>)	160, 324, 392, 458
<i>Betula alba</i> , subspecies <i>papyrifera</i>	160	Black Calabash Tree	116, 300
<i>Betula alba</i> , subspecies <i>papyrifera</i> , var. <i>communis</i>	160	Black Cherry, Wild	68, 282, 362, 428
<i>Betula alba</i> , subspecies <i>papyrifera</i> , var. <i>cordifolia</i>	160	Black Cottonwood (<i>Populus angustifolia</i>)	174, 330, 394, 462
<i>Betula alba</i> , subspecies <i>papyrifera</i> , var. <i>cordifolia</i>	160	Black Cottonwood (<i>Populus trichocarpa</i>)	174, 330, 394, 462, 576
<i>Betula alba</i> , subspecies <i>populifolia</i>	159	Black Cypress	184, 334, 350, 398, 466
<i>Betula alba</i> , var. <i>papyrifera</i>	160	Black Gum	92, 290, 366, 432
<i>Betula alba</i> , var. <i>populifolia</i>	159, 250, 255, 258, 324, 350, 458	Black Haw	94, 290, 366, 432
<i>Betula alba</i> , var. <i>populifolia</i> (<i>Betula papyrifera</i>)	160	Black Hickory (<i>Carya porcina</i>)	134, 310, 350, 380, 446
<i>Betula angulata</i>	161	Black Hickory (<i>Carya tomentosa</i>)	134, 310, 350, 380, 444
<i>Betula Canadensis</i>	160	Black Hills region of Dakota, remarks by Mr. Robert Douglas on the forests of the	561, 562
<i>Betula carpinifolia</i>	162	Black Ironwood	39, 272, 358, 422
<i>Betula cordifolia</i>	160	Black Jack (<i>Quercus Catesbeii</i>)	151, 320, 388, 454
<i>Betula excelsa</i> (<i>Betula lenta</i>)	162	Black Jack (<i>Quercus nigra</i>)	150, 265, 320, 350, 388, 454
<i>Betula excelsa</i> (<i>Betula lutea</i>)	161	Black Jack, Forked-leaf	151, 320, 388, 454
<i>Betula grandis</i>	160	Black Larch	215, 346, 352, 412, 478
<i>Betula incana</i>	164	Black Locust (<i>Gleditsia triacanthos</i>)	59, 280, 360, 426
<i>Betula lanulosa</i>	161	Black Locust (<i>Robinia Pseudacacia</i>)	55, 278, 350, 360, 414, 424
<i>Betula lenta</i>	162, 249, 255, 258, 324, 392, 458	Black Mangrove	117, 302
<i>Betula lenta</i> (<i>Betula alba</i> , var. <i>populifolia</i>)	159	Black Oak (<i>Quercus Emoryi</i>)	146, 265, 318, 386, 452
<i>Betula lenta</i> (<i>Betula lutea</i>)	161	Black Oak (<i>Quercus Kelloggii</i>)	149, 265, 320, 388, 416, 454
<i>Betula lutea</i>	161, 250, 255, 258, 262, 324, 392, 458	Black Oak (<i>Quercus rubra</i>)	148, 265, 318, 350, 386, 452
<i>Betula nigra</i>	161, 250, 255, 258, 262, 324, 392, 458	Black Oak (<i>Quercus tinctoria</i>)	149, 265, 318, 350, 388, 454, 528
<i>Betula nigra</i> (<i>Betula lenta</i>)	162	Black Persimmon	105, 294
<i>Betula nigra</i> (<i>Betula papyrifera</i>)	160	Black Pine (<i>Pinus Jeffreyi</i>)	193, 338, 402, 470
<i>Betula occidentalis</i>	160, 250, 255, 258, 324, 392, 458	Black Pine (<i>Pinus Murrayana</i>)	195, 338, 350, 404, 470
<i>Betula occidentalis</i> (<i>Betula papyrifera</i>)	160	Black Sloe	67, 282, 428
<i>Betula papyracea</i>	159, 575	Black Spruce	208, 265, 342, 352, 408, 474, 496
<i>Betula papyrifera</i>	12, 159, 250, 255, 258, 262, 324, 350, 392, 458	Black Sugar Maple	48, 276, 350, 358, 424
<i>Betula populifolia</i>	159	Black Thorn	79, 286, 364, 430
<i>Betula rubra</i> (<i>Betula nigra</i>)	161	Black Tree	117, 302
<i>Betula rubra</i> (<i>Betula papyrifera</i>)	160	Black Walnut	131, 308, 350, 376, 414, 444
<i>Betula rugosa</i>	164	Black Willow (<i>Salix flaveocens</i> , var. <i>Scouleriana</i>)	170, 328, 394, 460
<i>Betula serrulata</i>	164	Black Willow (<i>Salix nigra</i>)	166, 328, 394, 460
<i>Betula-Alnus glauca</i>	164	Blackwood	117, 302
<i>Betula-Alnus maritima</i>	162	Blue Ash	111, 298, 370, 436
<i>Betula-Alnus rubra</i>	164	Blue Beech	159, 322, 392, 456
<i>Betula-Alnus</i> , var. <i>B. incana</i>	164	Blue Jack	153, 320, 390, 456
BETULACEÆ	159-165, 255, 258, 324, 392, 458	Blue Myrtle	41, 272
Big-bud Hickory	134, 310, 350, 380, 444	Blue Oak	143, 316, 386, 450
Big Cottonwood	175, 330, 350, 396, 462	Blue Spruce	205, 344, 406, 474
Big Cypress region of Florida	523	Blunewood	40, 272
Big Laurel	19, 266, 354, 414, 418	Bois d'Arc	128, 306, 376, 442, 542
Big Shell-bark	133, 310, 378, 444	BORRAGINACEÆ	113, 114, 254, 257, 300, 372, 438
Big Tree	184, 334, 398, 406	<i>Borya acuminata</i>	112
<i>Bigelovia acuminata</i>	112	<i>Borya ligustrina</i>	112
<i>Bignonia Catalpa</i>	115	<i>Borya nitida</i>	112
<i>Bignonia linearis</i>	110	Bottom Shell-bark	133, 310, 378, 444

	Page.		Page.
Bourreria.....	11, 15	Cabbage Palmetto.....	217, 348, 352
Bourreria Havanensis.....	114, 249, 254, 257, 300, 372, 438	Cabbage Tree.....	217, 348, 352
Bourreria Havanensis, var. radula.....	114	Caballeria Coriacea.....	99
Bourreria ovata.....	114	CACTACEÆ.....	89, 90, 288
Bourreria radula.....	114	Cactus, Giant.....	90, 288
Bourreria recurva.....	114	Cagiput.....	120, 302, 374, 440
Bourreria tomentosa.....	114	Calabash Tree, Black.....	116, 340
Egyrreria tomentosa, var. Havanensis.....	114	Calico Bush.....	98, 292, 368, 434
Box Elder (Negundo aceroides).....	51, 276, 380, 424	Calicowood.....	106, 296
Box Elder (Negundo Californicum).....	51, 276, 380, 424	California Bay Tree.....	120, 302, 374, 440
Boxwood (Cornus florida).....	91, 288, 366, 430	California Buckeye.....	44, 274, 358, 422
Boxwood (Schæfferia frutescens).....	39, 272	California, economic value of the Redwood of.....	579
Brahea dulcis.....	217	California Holly.....	84, 286
Brahea filamentosa.....	217	California, injury by grazing cattle to the mountain forests of.....	579
Brickley Thatch.....	218, 348	California Laurel.....	120, 302, 374, 440
Brittle Thatch.....	218, 348	California, lumber industry and manufactures from wood in.....	486, 487, 578, 580
Broad-leaved Maple.....	47, 274, 358, 422	California Nutmeg.....	180, 334, 400, 406
Droussonetia secundiflora.....	57	California Olive.....	120, 302, 374, 440
Droussonetia tinctoria.....	128	California, pasturage of mountain forests in.....	579, 580
Brown Hickory.....	134, 310, 350, 380, 446	California, rank of, according to value of lumber products.....	487
Bucida Buceras.....	87	California, statistics of forests, forest fires, standing timber, etc., in.....	491, 578-580
Buckeye, California.....	44, 274, 358, 422	California, tabular statement of the amount of Redwood standing in the forests of, May 31, 1880.....	579
Buckeye, Fetid.....	42, 274, 358, 422	Calothyrsus Californica.....	43
Buckeye, Ohio.....	42, 274, 358, 422	Calyptranthes.....	10, 14
Buckeye, Spanish.....	44, 274, 422	Calyptranthes Chytraculia.....	88, 249, 288
Buckeye, Sweet.....	43, 274	Canada Plum.....	65, 282, 362, 426
Buckthorn, Southern.....	103, 294, 368, 434	Canella.....	10, 13
Buckwheat Tree.....	38, 272, 356, 420	Canella alba.....	24, 249, 253, 256, 268, 354, 418
Bull Bay.....	19, 266, 354, 414, 418	Canella laurifolia.....	24
Bull Nut.....	134, 310, 350, 380, 444	Canella Winterana.....	24
Bull Pine (Pinus Jeffreyi).....	193, 338, 402, 470	CANELLACEÆ.....	24, 253, 256, 268, 354, 418
Bull Pine (Pinus mitis).....	200, 340, 350, 406, 472	Canoe Birch.....	160, 324, 350, 392, 458
Bull Pine (Pinus ponderosa).....	193, 338, 350, 402, 468	Canoe Cedar.....	177, 380, 396, 462
Bull Pine (Pinus Sabiniana).....	195, 338, 350, 404, 470	Canotia.....	10, 12, 13
Bumelia.....	11, 15	Canotia holocantha.....	32, 249, 270
Bumelia angustifolia.....	103	CAPPARIDACEÆ.....	24, 268
Bumelia arborea.....	102	Capparis.....	10, 13
Bumelia chrysophylloides.....	101	Capparis Breymia.....	24
Bumelia cuneata.....	103, 249, 254, 294, 368, 434	Capparis cynophyllophora.....	24
Bumelia ferruginea.....	102	Capparis emarginata.....	24
Bumelia fetidissima.....	101	Capparis Jamaicensis.....	24, 249, 268
Bumelia lanuginosa.....	102, 250, 254, 257, 294, 368, 434	Capparis siliquosa.....	24
Bumelia lanuginosa, var. macrocarpa.....	102	Capparis torulosa.....	24
Bumelia lycioides.....	102, 249, 254, 257, 294, 368, 434	Capparis uncinata.....	24
Bumelia lycioides, var. reclinatam.....	103	CAPRIFOLIACEÆ.....	98, 94, 253, 257, 260, 366, 432
Bumelia macrocarpa.....	102	Cardioplepis obtusa.....	41
Bumelia Mastichodendron.....	101	Carolina Poplar.....	175, 330, 350, 396, 462
Bumelia myrsinifolia.....	103	Carpinus.....	11, 16
Bumelia oblongifolia.....	102	Carpinus Americana.....	158
Bumelia pallida.....	101	Carpinus Betulus Virginiana.....	158
Bumelia parvifolia.....	103	Carpinus Caroliniana.....	158, 249, 255, 258, 322, 392, 456
Bumelia reclinata (Bumelia cuneata).....	103	Carpinus Ostrya.....	158
Bumelia reclinata (Bumelia lycioides).....	103	Carpinus Ostrya, var. Americana.....	158
Bumelia reclinata (Bumelia tenax).....	101	Carpinus triflora.....	158
Bumelia salicifolia.....	101	Carpinus Virginiana.....	158
Bumelia serrata.....	69	Carya.....	4, 11, 15
Bumelia spinosa.....	102, 249, 294	Carya alba.....	132, 249, 254, 257, 261, 308, 350, 378, 444
Bumelia tenax.....	101, 249, 254, 257, 294, 368, 434	Carya alba (Carya tomentosa).....	134
Bumelia tomentosa.....	102	Carya amara.....	135, 249, 254, 257, 310, 350, 380, 446
Bumwood.....	54, 278, 360, 424	Carya amara, var. myristicaeformis.....	135
Burlington, Vermont, as a lumber market, importance of.....	490	Carya amara, var. porcina.....	134
Burning Bush.....	38, 272	Carya angustifolia.....	132
Burr Oak.....	140, 265, 314, 384, 448	Carya aquatica.....	135, 249, 254, 257, 261, 310, 350, 380, 414, 446
Bursera.....	10, 13	Carya cathartica.....	130
Bursera acuminata.....	32	Carya cordiformis.....	133
Bursera gummiifera.....	32, 251, 253, 256, 270, 350, 356, 420	Carya glabra.....	134
BURSERACEÆ.....	32, 33, 253, 256, 270, 356, 420	Carya Illinoensis.....	132
Bustic.....	101, 294, 368, 434	Carya integrifolia.....	135
Butternut.....	130, 306, 376, 442	Carya microcarpa.....	133
Button-ball Tree.....	129, 306, 350, 376, 442	Carya myristicaeformis.....	135, 249, 254, 257, 310, 350, 380, 446
Buttonwood (Conocarpus erecta).....	87, 288, 364, 430	Carya oliviformis.....	132, 249, 254, 257, 308, 350, 378, 444
Buttonwood (Platanus occidentalis).....	129, 306, 350, 376, 442	Carya porcina.....	134, 249, 254, 257, 261, 310, 350, 380, 446
Buttonwood (Platanus racemosa).....	129, 306, 376, 442	Carya sulcata.....	133, 249, 254, 257, 261, 310, 378, 444
Buttonwood, White.....	87, 288, 364, 430	Carya tetraptera.....	132
Byrsionima.....	10, 13		
Byrsionima lucida.....	28, 250, 258, 266, 268, 356, 420		

	Page.		Page.
<i>Carya tomentosa</i>	183, 249, 254, 257, 310, 350, 380, 444	<i>Celtis cordata</i>	125
<i>Carya tomentosa</i> , var. <i>maxima</i>	134	<i>Celtis crassifolia</i>	125
<i>Caryotaxus Myristica</i>	186	<i>Celtis crassifolia</i> , var. <i>eucalyptifolia</i>	125
<i>Caryotaxus taxifolia</i>	186	<i>Celtis crassifolia</i> , var. <i>morifolia</i>	125
Cassida	101, 294, 368, 434	<i>Celtis crassifolia</i> , var. <i>thlasifolia</i>	125
Cassena	38, 272	<i>Celtis Douglasii</i>	126
<i>Cassine Caroliniana</i>	36	<i>Celtis Floridiana</i>	125
<i>Cassine Peragua</i>	36	<i>Celtis fuscata</i>	125
<i>Cassine ramulosa</i>	36	<i>Celtis heterophylla</i>	125
Castanea	11, 16	<i>Celtis integrifolia</i>	125
<i>Castanea alnifolia</i>	156	<i>Celtis laevigata</i>	125
<i>Castanea Americana</i>	157	<i>Celtis Lindheimeri</i>	126
<i>Castanea chrysophylla</i>	156	<i>Celtis longifolia</i>	125
<i>Castanea chrysophylla</i> , var. <i>minor</i>	156	<i>Celtis maritima</i> (in Am. Monthly Mag. and Crit. Rev.)	125
<i>Castanea nana</i>	156	<i>Celtis maritima</i> (in New Fl. and Bot.)	125
<i>Castanea pumila</i>	156, 250, 255, 258, 322, 390, 456	<i>Celtis Mississippiensis</i>	125
<i>Castanea sempervirens</i>	156	<i>Celtis morifolia</i>	125
<i>Castanea vesca</i> (<i>Castanea pumila</i>)	156	<i>Celtis obliqua</i>	125
<i>Castanea vesca</i> (<i>Castanea vulgaris</i> , var. <i>Americana</i>)	157	<i>Celtis occidentalis</i>	12, 125, 249, 254, 257, 260, 306, 376, 414, 442
<i>Castanea vesca</i> , var. <i>Americana</i>	157	<i>Celtis occidentalis</i> , var. <i>cordata</i>	125
<i>Castanea vulgaris</i> , var. <i>Americana</i>	157, 250, 255, 258, 262, 265, 322, 390, 456	<i>Celtis occidentalis</i> , var. <i>crassifolia</i>	126
Castanopsis	11, 12, 15	<i>Celtis occidentalis</i> , var. <i>grandidentata</i>	125
<i>Castanopsis chrysophylla</i>	156, 250, 255, 258, 322, 390, 456	<i>Celtis occidentalis</i> , var. <i>integrifolia</i>	125
<i>Castanopsis chrysophylla</i> , var. <i>minor</i>	156	<i>Celtis occidentalis</i> , var. <i>pumila</i> (<i>Celtis occidentalis</i>)	126
<i>Castanopsis chrysophylla</i> , var. <i>pumila</i>	156	<i>Celtis occidentalis</i> , var. <i>pumila</i> (<i>Celtis occidentalis</i> , var. <i>reticulata</i>)	126
Catalogue of Forest Trees	17-219	<i>Celtis occidentalis</i> , var. <i>reticulata</i>	126, 249, 254, 257, 306, 376, 442
Catalogue of Forest Trees, Index to	220-243	<i>Celtis occidentalis</i> , var. <i>scabruscula</i>	125
Catalpa	11, 15, 115, 300, 372, 438	<i>Celtis occidentalis</i> , var. <i>serrulata</i>	125
<i>Catalpa bignonioides</i>	115, 250, 254, 257, 300, 372, 438	<i>Celtis occidentalis</i> , var. <i>tenuifolia</i>	125
<i>Catalpa bignonioides</i> (<i>Catalpa speciosa</i>)	115	<i>Celtis patula</i>	125
<i>Catalpa communis</i>	115	<i>Celtis pumila</i>	125
<i>Catalpa cordifolia</i> (<i>Catalpa bignonioides</i>)	115	<i>Celtis reticulata</i>	126
<i>Catalpa cordifolia</i> (<i>Catalpa speciosa</i>)	115	<i>Celtis salicifolia</i>	125
<i>Catalpa speciosa</i>	115, 250, 254, 257, 300, 350, 372, 414, 438	<i>Celtis tenuifolia</i>	125
<i>Catalpa syriacaefolia</i>	115	<i>Celtis Texana</i>	126
Catalpa, Western	115, 300, 350, 372, 414, 438	Central pine hills of Mississippi, forests of the	534
Catawba	115, 300, 372, 438	<i>Cerasus Americana</i>	65
Cat's Claw (<i>Acacia Greggii</i>)	63, 282, 362, 426	<i>Cerasus borealis</i>	66
Cat's Claw (<i>Acacia Wrightii</i>)	63, 282	<i>Cerasus Capolin</i>	68
Cat's Claw (<i>Pithecolobium Unguis-cati</i>)	64, 282	<i>Cerasus Capuli</i>	68
Ceanothus	10, 12, 13, 573	<i>Cerasus Caroliniana</i>	69
<i>Ceanothus ferreus</i>	39	<i>Cerasus Ohioana</i>	66
<i>Ceanothus laevigatus</i>	39	<i>Cerasus demissa</i>	69
<i>Ceanothus reclinatus</i>	41	<i>Cerasus emarginata</i>	67
<i>Ceanothus thyrsiflorus</i>	41, 250, 272	<i>Cerasus erecta</i>	67
Cedar, Alaska, Tide-land Spruce, and Hemlock the characteristic and most valuable trees of the Northern Coast Forest	7	<i>Cerasus glandulosa</i>	67
Cedar an important tree in Texas	540	<i>Cerasus hiemalis</i>	65
Cedar, Bastard	176, 330, 396, 462	<i>Cerasus ilicifolia</i>	70
Cedar, Canoe	177, 330, 396, 462	<i>Cerasus mollis</i>	67
Cedar Elm	122, 304, 374, 440	<i>Cerasus nigra</i>	65
Cedar, Incense	176, 330, 396, 462	<i>Cerasus Pennsylvanica</i>	66
Cedar, Oregon	179, 332, 350, 396, 464	<i>Cerasus persicifolia</i>	66
Cedar, Pencil, of Florida	523	<i>Cerasus serotina</i> (<i>Prunus demissa</i>)	69
Cedar Pine	201, 340, 406, 472	<i>Cerasus serotina</i> (<i>Prunus serotina</i>)	68
Cedar, Port Orford	179, 332, 350, 396, 464, 576	<i>Cerasus sphaerocarpa</i>	70
Cedar, Post	176, 330, 396, 462	<i>Cerasus umbellata</i>	67
Cedar, Red (<i>Juniperus Virginiana</i>)	183, 332, 396, 464, 544	<i>Cerasus Virginiana</i>	68
Cedar, Red (<i>Thuya gigantea</i>)	177, 330, 396, 462, 573, 575, 576, 580	<i>Cercidium floridum</i>	60
Cedar, Stinking (<i>Torreya Californica</i>)	186, 334, 400, 466	<i>Cercis</i>	10, 12, 14
Cedar, Stinking (<i>Torreya taxifolia</i>)	186, 334, 400, 466, 521	<i>Cercis Canadensis</i>	61, 250, 253, 256, 280, 362, 426
Cedar, White (<i>Chamaecyparis Lawsoniana</i>)	179, 332, 350, 396, 464	<i>Cercis Canadensis</i> , var. <i>pubescens</i>	61
Cedar, White (<i>Chamaecyparis sphaeroloba</i>)	178, 330, 396, 416, 464	<i>Cercis occidentalis</i>	61
Cedar, White (<i>Libocedrus decurrens</i>)	176, 330, 396, 462	<i>Cercis occidentalis</i> , var.	61
Cedar, White (<i>Thuya occidentalis</i>)	176, 330, 350, 396, 462	<i>Cercis occidentalis</i> , var. <i>Texensis</i>	61
<i>Cedrus Mahagoni</i>	33	<i>Cercis reniformis</i>	61, 249, 280
CELASTRACE	38, 39, 272	<i>Cercocarpus</i>	9, 10, 12, 14, 569, 571
<i>Celtis</i>	11, 12, 15, 251	<i>Cercocarpus betulafolius</i>	71
<i>Celtis alba</i>	125	<i>Cercocarpus betuloides</i>	71
<i>Celtis Audubertiana</i>	126	<i>Cercocarpus brevifolius</i>	71
<i>Celtis Audubertiana</i> , var. <i>oblongata</i>	126	<i>Cercocarpus intricatus</i>	71
<i>Celtis Audubertiana</i> , var. <i>ovata</i>	126	<i>Cercocarpus ledifolius</i>	13, 71, 249, 284, 350, 428
<i>Celtis Berlandieri</i>	126	<i>Cercocarpus ledifolius</i> , var. <i>intricatus</i>	71
<i>Celtis brevipes</i>	126	<i>Cercocarpus parvifolius</i>	71, 249, 284
<i>Celtis canina</i>	125	<i>Cercocarpus parvifolius</i> , var. <i>glaber</i>	71
		<i>Cercocarpus parvifolius</i> , var. <i>paucidentatus</i>	71

	Page.		Page.
Cereus.....	10, 12, 14	<i>Cinchona Caribaea</i>	95
Cereus giganteus.....	89, 251, 288	<i>Cinchona Caroliniana</i>	95
Chapote.....	105, 294	<i>Cinchona Jamaicensis</i>	95
Chamaecyparis.....	11, 12, 16	Cinnamon Bark.....	24, 268, 354, 418
<i>Chamaecyparis Bourseri</i> (<i>Chamaecyparis Lawsoniana</i>).....	178	Cinnamon, Wild.....	24, 268, 354, 418
<i>Chamaecyparis Bourseri</i> (<i>Juniperus occidentalis</i>).....	181	Citharexylum.....	11, 15
<i>Chamaecyparis excelsa</i>	178	Citharexylum villosum.....	116, 249, 254, 257, 300, 372, 438
<i>Chamaecyparis Lawsoniana</i>	8, 178, 250, 255, 258, 350, 398, 464, 576	Cladrastis.....	10, 11, 14
<i>Chamaecyparis Nutkaensis</i>	178, 250, 255, 258, 263, 332, 398, 464, 580	<i>Cladrastis lutea</i>	57
<i>Chamaecyparis Nutkaensis</i> , var. <i>glauca</i>	178	Cladrastis tinctoria.....	57, 250, 253, 256, 278, 360, 426
<i>Chamaecyparis Nutkanus</i>	178	Clammy Locust.....	56, 278
Chamaecyparis, one of the characteristic and most valuable trees of the Coast Forest.....	7	Cliff Elm.....	123, 304, 374, 440
<i>Chamaecyparis sphaeroides</i>	177, 251, 255, 258, 262, 330, 396, 416, 464	<i>Cliftonia ligustrina</i>	38, 250, 253, 256, 272, 356, 420
<i>Chamaecyparis Palmetto</i>	217	Clusia.....	10, 13
Charcoal used for domestic and manufacturing purposes during the census year, amount and value of.....	489	<i>Clusia flava</i>	25
<i>Chelanthodendron Californicum</i>	26	<i>Clusia rosea</i>	25
Cherry.....	100, 292	Coast Forest in the Pacific region.....	7
Cherry Birch.....	162, 324, 392, 458	Coast Forest, Northern, in Alaska.....	580
Cherry, Indian.....	40, 272, 358, 422	Coast Live Oak.....	147, 318, 386, 452
Cherry, May.....	84, 286, 364, 430	Coast Pine Belt in South Carolina.....	518
Cherry, Pigeon.....	66, 282, 428	Coccoloba.....	11, 13
Cherry, Pin.....	66, 282, 428	<i>Coccoloba Floridana</i>	117, 249, 254, 257, 302, 372, 438
Cherry, Ram.....	66, 282, 362, 428	<i>Coccoloba Loganensis</i>	118
Cherry, Wild (<i>Prunus Capuli</i>).....	66, 282, 428	<i>Coccoloba parvifolia</i>	117
Cherry, Wild (<i>Prunus demissa</i>).....	66, 284, 362, 428	<i>Coccoloba uvifera</i>	118, 249, 302, 372, 448
Cherry, Wild Black.....	66, 282, 362, 428	<i>Coccoloba uvifera</i> , var. <i>Loganensis</i>	118
Cherry, Wild Red.....	66, 282, 428	<i>Coccoloba uvifera</i> , var. <i>ovalifolia</i>	118
Chestnut.....	157, 265, 322, 350, 390, 456	Cockspur Thorn.....	76, 286, 364, 430
Chestnut Oak (<i>Quercus densiflora</i>).....	155, 265, 322, 390, 456, 576, 578, 580	Cocoa Plum.....	65, 282, 362, 426
Chestnut Oak (<i>Quercus prinoides</i>).....	143, 265, 316, 384, 450	Coffee Nut.....	58, 280, 360, 426
Chestnut Oak (<i>Quercus Prinus</i>).....	142, 265, 316, 350, 384, 416, 450	Coffee Tree, Kentucky.....	58, 280, 360, 426
Chestnut Oak, Rock.....	142, 265, 316, 350, 384, 416, 450	Colorado, lumber industry and manufactures from wood in.....	486, 487, 567, 568
Chicago, amount of lumber and shingles received at, during the year 1880.....	548, 549	Colorado, rank of, according to value of lumber products.....	487
Chicago, commercial importance of, with reference to the lumber industry.....	548	Colorado, statistics of forests, forest fires, standing timber, etc., in.....	491, 567, 568
Chicago, early lumber trade of.....	549, 550	Colorado, Utah, and New Mexico, Chicago the principal source of supply of lumber for.....	568, 569
Chicago, the principal source of supply of lumber for Colorado, Utah, and New Mexico.....	568, 569	Colubrina.....	10, 13
Chickasaw Plum.....	66, 282, 362, 426	<i>Colubrina reclinata</i>	41, 249, 274, 358
Chilopsis.....	11, 12, 15	COMNETACEÆ.....	87, 253, 257, 288, 364, 430
<i>Chilopsis glutinosa</i>	116	Comparative value of woods.....	252-255
<i>Chilopsis linearis</i>	116	Compression, behavior of the principal woods of the United States under.....	418-481
<i>Chilopsis saligna</i>	13, 116, 250, 254, 257, 300, 372, 438	Condalia.....	10-12
<i>Chimanthus amygdalinus</i>	69	<i>Condalia ferrea</i>	39, 249, 253, 256, 272, 358, 422
China, Wild.....	44, 274, 358, 414, 422	<i>Condalia obovata</i>	12, 46, 249, 272
Chinquapin (<i>Castanea pumila</i>).....	156, 322, 390, 456	CONIFERÆ.....	4, 176-216, 255, 258, 330, 362, 396, 416, 525
Chinquapin (<i>Castanopsis chrysophylla</i>).....	156, 322, 390, 456	Connecticut, lumber industry and manufactures from wood in.....	486, 487, 500, 501
Chinquapin Oak.....	143, 265, 316, 384, 450	Connecticut, rank of, according to value of lumber products.....	487
Chionanthus.....	11, 15	Connecticut, statistics of forests, forest fires, standing timber, etc., in.....	500, 501
<i>Chionanthus angustifolia</i>	112	Connecticut, the Northern Pine belt in.....	500
<i>Chionanthus heterophylla</i>	112	Conocarpus.....	10, 14
<i>Chionanthus longifolia</i>	112	<i>Conocarpus acutifolia</i>	87
<i>Chionanthus maritima</i>	112	<i>Conocarpus erecta</i>	87, 249, 253, 257, 288, 364, 430
<i>Chionanthus montana</i>	112	<i>Conocarpus erecta</i> , var. <i>procumbens</i>	87
<i>Chionanthus trifida</i>	112	<i>Conocarpus procumbens</i>	87
<i>Chionanthus Virginica</i>	112, 250, 298	<i>Conocarpus racemosa</i>	87
<i>Chionanthus Virginica</i> , var. <i>angustifolia</i>	112	Consumption of wood for domestic purposes during the census year, estimated.....	489
<i>Chionanthus Virginica</i> , var. <i>latifolia</i>	112	Coral Sumach.....	54, 278, 360, 424
<i>Chionanthus Virginica</i> , var. <i>maritima</i>	112	Cordia.....	11, 15
<i>Chionanthus Virginica</i> , var. <i>montana</i>	112	<i>Cordia Bolanderi</i>	114, 249, 300
Chittamwood.....	52, 276	<i>Cordia Floridana</i>	114
<i>Chloromeles sempervirens</i>	72	<i>Cordia juglandifolia</i>	113
Chrysobalanus.....	10, 14	<i>Cordia Sebestena</i>	113, 249, 300
Chrysobalanus Icaco.....	64, 249, 282, 362, 426	<i>Cordia speciosa</i>	113
<i>Chrysobalanus Icaco</i> , var. <i>pellocarpa</i>	65	Cork Elm.....	123, 304, 374, 440
<i>Chrysobalanus pellocarpa</i>	65	Corkwood.....	117, 302, 372, 438
Chrysophyllum.....	11, 15	CORNACEÆ.....	90-93, 253, 257, 288, 366, 430
<i>Chrysophyllum Barbaco</i>	100	Cornus.....	10-13, 14
<i>Chrysophyllum Caneto</i> , β.....	100	<i>Cornus alternata</i>	90
<i>Chrysophyllum ferrugineum</i>	100	<i>Cornus alternifolia</i>	90, 249, 268
<i>Chrysophyllum microphyllum</i>	100	<i>Cornus florida</i>	90, 249, 253, 257, 280, 288, 366, 432
<i>Chrysophyllum monopyrenum</i>	100	<i>Cornus florida</i> (<i>Cornus Nuttallii</i>).....	91
<i>Chrysophyllum oliviforme</i>	100, 249, 254, 257, 292, 368, 434	<i>Cornus Nuttallii</i>	91, 249, 253, 257, 288, 366, 432
Cigar Tree.....	115, 300, 372, 438	<i>Corypha Palmetto</i>	217
		<i>Cotinus Americanus</i>	58

	Page.		Page.
<i>Ootinus cogggria</i>	52	<i>Oratægus lucida</i> (<i>Cratægus Crus-galli</i>)	76
Cotton Gum	93, 290, 350, 366, 432	<i>Oratægus macracantha</i>	77
Cottonwood (<i>Populus Fremontii</i>)	175, 300, 396, 462	<i>Oratægus Michauxii</i>	83
Cottonwood (<i>Populus Fremontii</i> , var. <i>Wislizeni</i>)	175, 300, 396, 462	<i>Oratægus microcarpa</i>	81
Cottonwood (<i>Populus monilifera</i>)	175, 330, 350, 396, 462	<i>Oratægus mollis</i>	78
Cottonwood, Balsam	174, 330, 394, 462	<i>Oratægus obovatifolia</i>	80
Cottonwood, Big	175, 330, 350, 396, 462	<i>Oratægus opaca</i>	82
Cottonwood, Black (<i>Populus angustifolia</i>)	174, 330, 394, 462	<i>Oratægus ovalifolia</i>	76
Cottonwood, Black (<i>Populus trichocarpa</i>)	174, 330, 394, 462, 576	<i>Oratægus oxyacantha</i>	81
Cottonwood, River	172, 328, 394, 460	<i>Oratægus oxyacantha</i> , var. <i>apiifolia</i>	81
Cottonwood, Swamp	172, 328, 394, 460	<i>Oratægus parvifolia</i>	83
Cottonwood, White	175, 330, 396, 462	<i>Oratægus populifolia</i> (<i>Cratægus coccinea</i>)	78
Cow Oak	141, 316, 384, 416, 450, 533	<i>Oratægus populifolia</i> (<i>Cratægus cordata</i>)	80
Crab, American	72, 284, 364, 428	<i>Oratægus prunifolia</i>	76
Crab, Sweet-scented	72, 284, 364, 428	<i>Oratægus prunifolia</i>	77
Crab Apple, American	72, 284	<i>Oratægus punctata</i>	80
Crab Apple, Oregon	73, 284	<i>Oratægus punctata</i> , var. <i>brevispina</i>	75
Crab Apple, Southern	72, 284	<i>Oratægus punctata</i> , var. <i>rubra</i> and <i>aurea</i>	80
Crabwood	121, 304	<i>Oratægus punctata</i> , var. <i>xanthocarpa</i>	80
<i>Cratægus</i>	10-12, 539, 573	<i>Oratægus pyrifolia</i>	79
<i>Cratægus æstivalis</i>	82, 250, 253, 256, 286, 304, 430	<i>Oratægus racemosa</i>	84
<i>Cratægus apiifolia</i>	81, 249, 286	<i>Cratægus rivularis</i>	13, 74, 249, 284
<i>Cratægus arborescens</i>	75, 250, 253, 256, 284, 364, 428	<i>Cratægus rivularis</i> (<i>Cratægus Douglasii</i>)	75
<i>Cratægus arbutifolia</i>	83	<i>Cratægus salicifolia</i>	78
<i>Cratægus berberifolia</i>	82, 248	<i>Cratægus sanguinea</i>	75
<i>Cratægus Bosciensis</i>	77	<i>Cratægus sanguinea</i> , var. <i>Douglasii</i> (<i>Cratægus Douglasii</i>)	75
<i>Cratægus brachyacantha</i>	75, 249, 284, 539	<i>Cratægus sanguinea</i> , var. <i>Douglasii</i> (<i>Cratægus rivularis</i>)	74
<i>Cratægus Caroliniana</i>	82	<i>Cratægus sanguinea</i> , var. <i>villosa</i>	78
<i>Cratægus coccinea</i>	77, 249, 286	<i>Cratægus spathulata</i>	81, 249, 253, 256, 286, 364, 430
<i>Oratægus coccinea</i> (<i>Cratægus tomentosa</i>)	79	<i>Cratægus spathulata</i> (<i>Cratægus flava</i> , var. <i>pubescens</i>)	83
<i>Oratægus coccinea</i> , var. <i>cordata</i>	79	<i>Cratægus spicata</i>	85
<i>Oratægus coccinea</i> , var. <i>mollis</i>	78	<i>Cratægus subvillosa</i>	78, 249, 253, 256, 286, 364, 430
<i>Cratægus coccinea</i> var. <i>oligandra</i>	78	<i>Cratægus Texana</i>	78
<i>Cratægus coccinea</i> , var. <i>populifolia</i>	78	<i>Cratægus tomentosa</i>	12, 79, 249, 253, 256, 286, 364, 430
<i>Cratægus coccinea</i> , var. <i>typica</i>	78	<i>Cratægus tomentosa</i> (<i>Amelanchier Canadensis</i>)	84
<i>Cratægus coccinea</i> , var. <i>viridis</i>	78	<i>Oratægus tomentosa</i> , var. <i>mollis</i>	78
<i>Oratægus coccinea</i> , var. <i>viridis</i> (<i>Cratægus tomentosa</i>)	79	<i>Oratægus tomentosa</i> , var. <i>plicata</i>	80
<i>Cratægus cordata</i>	80, 249, 286	<i>Cratægus tomentosa</i> , var. <i>punctata</i>	80
<i>Oratægus coronaria</i>	72	<i>Cratægus tomentosa</i> , var. <i>pyrifolia</i>	79
<i>Oratægus Coursetiana</i>	76	<i>Oratægus turbinata</i>	82
<i>Cratægus Crus-galli</i>	76, 249, 253, 256, 286, 364, 430	<i>Cratægus Virginica</i>	83
<i>Oratægus Crus-galli</i> (<i>Cratægus coccinea</i>)	77	<i>Cratægus viridis</i> (<i>Cratægus coccinea</i>)	78
<i>Oratægus Crus-galli</i> (<i>Cratægus tomentosa</i> , var. <i>punctata</i>)	80	<i>Oratægus viridis</i> (<i>Cratægus flava</i> , var. <i>pubescens</i>)	83
<i>Cratægus Crus-galli</i> , var. <i>linearis</i>	76	<i>Oratægus Watsoniana</i>	76
<i>Cratægus Crus-galli</i> , var. <i>ovalifolia</i>	76	<i>Crescentia</i>	11, 15
<i>Cratægus Crus-galli</i> , var. <i>prunifolia</i>	77	<i>Crescentia cucurbitina</i>	116, 250, 300
<i>Cratægus Crus-galli</i> , var. <i>pyracanthifolia</i>	78	<i>Crescentia latifolia</i>	116
<i>Oratægus Crus-galli</i> , var. <i>pyracanthifolia</i> (<i>Cratægus arborescens</i>)	75	<i>Crescentia lethifera</i>	116
<i>Oratægus Crus-galli</i> , var. <i>salicifolia</i>	76	<i>Crescentia obovata</i>	116
<i>Oratægus Crus-galli</i> , var. <i>splendens</i>	76	<i>Crescentia ovata</i>	116
<i>Cratægus cuneifolia</i>	80	<i>Crescentia toxicaria</i>	116
<i>Cratægus Douglasii</i>	75, 249, 284	Crop, forest, of the United States for the census year	485
<i>Oratægus elliptica</i> (<i>Cratægus æstivalis</i>)	82	Cucumber Tree (<i>Magnolia acuminata</i>)	20, 266, 354, 418
<i>Oratægus elliptica</i> (<i>Cratægus flava</i> , var. <i>pubescens</i>)	83	Cucumber Tree (<i>Magnolia cordata</i>)	21, 266, 354, 418
<i>Cratægus flava</i>	82, 249, 286	Cucumber Tree, Large-leaved	21, 266, 354, 418
<i>Oratægus flava</i> (<i>Cratægus flava</i> , var. <i>pubescens</i>)	83	Cucumber Tree, Long-leaved	22, 266, 354, 418
<i>Oratægus flava</i> (<i>Cratægus tomentosa</i>)	79	<i>Cupressinnata disticha</i>	183
<i>Oratægus flava</i> (<i>Cratægus tomentosa</i> , var. <i>punctata</i>)	80	<i>Cupressus</i>	11, 12, 16
<i>Oratægus flava</i> , var. <i>lobata</i>	82	<i>Cupressus Americana</i>	178
<i>Cratægus flava</i> , var. <i>pubescens</i>	83, 249, 253, 256, 286, 364, 430	<i>Cupressus Arbor-vita</i>	170
<i>Oratægus flexuosa</i>	80	<i>Cupressus Arizonica</i>	180
<i>Oratægus glandulosa</i> (<i>Cratægus coccinea</i>)	77	<i>Cupressus attenuata</i>	178
<i>Oratægus glandulosa</i> (<i>Cratægus Douglasii</i>)	75	<i>Cupressus Californica</i>	179
<i>Oratægus glandulosa</i> (<i>Cratægus flava</i>)	77	<i>Cupressus Californica gracilis</i> (<i>Cupressus Goveniana</i>)	179
<i>Oratægus glandulosa</i> (<i>Cratægus flava</i> , var. <i>pubescens</i>)	83	<i>Cupressus Californica gracilis</i> (<i>Cupressus Macnabiana</i>)	180
<i>Oratægus glandulosa</i> , var. <i>macracantha</i>	77	<i>Cupressus cornuta</i>	179
<i>Oratægus glandulosa</i> , var. <i>rotundifolia</i>	78	<i>Cupressus disticha</i>	183
<i>Oratægus latifolia</i> (<i>Cratægus tomentosa</i>)	79	<i>Cupressus disticha</i> , var. <i>imbricaria</i>	183
<i>Oratægus latifolia</i> (<i>Cratægus tomentosa</i> , var. <i>punctata</i>)	80	<i>Cupressus disticha</i> , var. <i>nulans</i>	183
<i>Oratægus leucocephalus</i>	79	<i>Cupressus disticha</i> , var. <i>patens</i>	183
<i>Oratægus leucophloeos</i>	79	<i>Cupressus fragrans</i>	178
<i>Oratægus linearis</i>	77	<i>Cupressus glandulosa</i>	180
<i>Oratægus lobata</i> (<i>Cratægus flava</i>)	82	<i>Cupressus Goveniana</i>	179, 250, 255, 258, 332, 398, 464
<i>Oratægus lobata</i> (<i>Cratægus tomentosa</i>)	79	<i>Cupressus Guadalupensis</i>	180, 250, 332, 568, 569
<i>Oratægus lucida</i> (<i>Cratægus æstivalis</i>)	82	<i>Cupressus Hartwegii</i>	179

	Page.		Page.
<i>Cupressus Hartwegii</i> , var. <i>fastigiata</i>	179	<i>Diospyros Virginiana</i> , var. <i>pubescens</i>	104
<i>Cupressus Lambertiana</i>	179	Diphollis.....	11, 15
<i>Cupressus Laosoniana</i>	178	Diphollis salicifolia.....	101, 249, 254, 257, 294, 368, 434
<i>Cupressus Macnabiana</i>	180, 248	Distribution of genera.....	10
<i>Cupressus macrocarpa</i>	8, 179, 250, 332, 398, 464	Distribution of species.....	12
<i>Cupressus macrocarpa</i> (<i>Cupressus Guadalupeensis</i>).....	180	District of Columbia, lumber industry and manufactures from wood in.....	486, 487
<i>Cupressus macrocarpa</i> , var. <i>fastigiata</i>	179	District of Columbia, original forest of, replaced by Oak, Scrub Pine, etc.....	511
<i>Cupressus Nootkalensis</i>	178	District of Columbia, rank of, according to value of lumber products.....	487
<i>Cupressus Nutkaensis</i>	178	District of Columbia, statistics of forests, forest fires, standing timber, etc., in.....	511
<i>Cupressus thyoides</i>	177	Division of the North American continent with reference to its forest geography.....	3
CUPULIFERÆ.....	137-159, 254, 257, 258, 312, 380, 414, 446	Doctor Gum.....	54, 278, 360, 424
Curtiss, A. H., remarks on the forests of Florida by.....	521, 522	Dogwood.....	90, 288
Custard Apple.....	23, 266, 354, 418	Dogwood, Flowering (<i>Cornus florida</i>).....	91, 288, 366, 430
Cypress, Bald.....	184, 334, 350, 398, 466	Dogwood, Flowering (<i>Cornus Nuttallii</i>).....	91, 288, 366, 432
Cypress, Black.....	184, 334, 350, 398, 466	Dogwood, Jammica.....	57, 278, 360, 426
Cypress, Deciduous.....	184, 334, 350, 398, 466	Dogwood, Striped.....	46, 274
Cypress, Lawson's.....	179, 332, 350, 398, 464	Douglas Fir.....	269, 265, 344, 352, 410, 476
Cypress, Monterey.....	179, 332, 398, 464	Douglas, Robert, remarks on the forests of the Black Hills region of Dakota by.....	561, 562
Cypress, Red.....	184, 334, 350, 398, 466	Downward Plum.....	103, 294, 368, 431
Cypress, Sitka.....	178, 332, 398, 464	<i>Drimophyllum pauciflorum</i>	120
Cypress, Southern, next to the Long-leaved Pine, the characteristic tree of the Southern Maritime Pine Belt.....	4	Drypetes.....	11, 15
Cypress swamps of the Tensas river in Alabama.....	525-527	<i>Drypetes alba</i> , var. <i>latifolia</i>	121
Cypress, White.....	184, 334, 350, 398, 466	<i>Drypetes crocea</i>	120, 249, 254, 257, 302, 374, 440
Cypress, Yellow.....	178, 332, 398, 464	<i>Drypetes crocea</i> , var. <i>latifolia</i>	121, 249, 254, 257, 302, 374, 440
CYBILLACEÆ.....	37, 38, 253, 256, 272, 358, 420	<i>Drypetes crocea</i> , var. <i>longipes</i>	120
Cyrilla.....	10, 11, 13	<i>Drypetes glauca</i> (<i>Drypetes crocea</i>).....	120
<i>Cyrilla Caroliniana</i>	37	<i>Drypetes glauca</i> (<i>Drypetes crocea</i> , var. <i>latifolia</i>).....	121
<i>Cyrilla fuscata</i>	37	<i>Drypetes sessiliflora</i>	120
<i>Cyrilla paniculata</i>	100	Dry wood of trees of the United States, weight of.....	249-251
<i>Cyrilla parvifolia</i>	37	Duck Oak.....	152, 320, 350, 388, 416, 454
<i>Cyrilla polystachia</i>	37	Dwarf Maple.....	48, 274
<i>Cyrilla racemiflora</i>	37, 249, 272, 356	Dwarf Sumach.....	53, 278, 360, 424
<i>Cyrilla racemosa</i>	37		
D.		E.	
Dahoon.....	35, 270, 356, 420	EBENACEÆ.....	104, 105, 254, 257, 294, 368, 454
Dahoon Holly.....	35, 270, 356, 420	Economic aspects of the forests of the United States.....	483-580
Dakota, lumber industry and manufacturers from wood in.....	486, 487, 561	Ehretia.....	11, 15
Dakota, rank of, according to value of lumber products.....	487	<i>Ehretia Bourreria</i>	114
Dakota, remarks by Mr. H. C. Putnam on the forests of the eastern portion of.....	561	<i>Ehretia elliptica</i>	114, 250, 254, 257, 300, 372, 438
Dakota, remarks by Mr. Robert Douglas on the forests of the Black Hills region of.....	561, 562	<i>Ehretia Havanensis</i>	114
Dakota, statistics of forests, forest fires, standing timber, etc., in.....	561, 562	<i>Ehretia radula</i>	114
Dalea.....	10, 12, 14	<i>Ehretia tomentosa</i>	114
Dalea spinosa.....	55, 250, 278	<i>Elaphrium integerrimum</i>	32
Darling Plum.....	39, 272, 358, 422	Elder (<i>Sambucus glauca</i>).....	93, 290, 366, 432
<i>Datisca hirta</i>	52	Elder (<i>Sambucus Mexicana</i>).....	94, 290
Deciduous Cypress.....	184, 334, 350, 398, 466	Elder, Box (<i>Negundo aceroides</i>).....	51, 276, 360, 424
Deciduous Forest of the Mississippi Basin and the Atlantic Plain.....	4	Elder, Box (<i>Negundo Californicum</i>).....	51, 276, 360, 424
Delaware, lumber industry and manufactures from wood in.....	486, 487, 511	Elder, Poison.....	54, 278
Delaware, rank of, according to value of lumber products.....	487	Elemi, Gum.....	33, 270, 350, 356, 420
Delaware, statistics of forests, forest fires, standing timber, etc., in.....	491, 511	Elkwood.....	21, 266, 354, 418
<i>Dermatophyllum spectosum</i>	57	Elm, American.....	123, 304, 350, 374, 414, 440
Desert Willow.....	116, 300, 372, 438	Elm, Cedar.....	122, 304, 374, 440
Devilwood.....	113, 300, 372, 438	Elm, Cliff.....	123, 304, 374, 440
Diamond Willow.....	170, 328	Elm, Cork.....	123, 304, 374, 440
Digger Pine.....	195, 338, 350, 404, 470	Elm, Hickory.....	123, 304, 374, 440
Dilly, Wild.....	108, 294, 368, 434	Elm, Moose.....	122, 304, 374, 440
<i>Diospyros</i>	11, 15	Elm, Red.....	122, 304, 374, 440
<i>Diospyros angustifolia</i>	104	Elm, Rock.....	123, 304, 374, 440
<i>Diospyros calycina</i>	104	Elm, Slippery (<i>Fremontia Californica</i>).....	26, 268
<i>Diospyros ciliata</i>	104	Elm, Slippery (<i>Ulmus fulva</i>).....	122, 304, 374, 440
<i>Diospyros concolor</i>	104	Elm, Water.....	123, 304, 350, 374, 414, 440
<i>Diospyros Guaiacana</i>	104	Elm, White (<i>Ulmus Americana</i>).....	123, 304, 350, 374, 414, 440
<i>Diospyros intermedia</i>	104	Elm, White (<i>Ulmus racemosa</i>).....	123, 304, 374, 440
<i>Diospyros lucida</i>	104	Elm, Winged.....	124, 304, 374, 440
<i>Diospyros Persimon</i>	104	<i>Emetilla ramulosa</i>	36
<i>Diospyros pubescens</i>	104	Enceno.....	147, 378, 388, 452
<i>Diospyros Texana</i>	105, 249, 294	<i>Endotropis oleifolia</i>	40
<i>Diospyros Virginiana</i>	104, 249, 254, 257, 260, 294, 350, 368, 434	ERICACEÆ.....	96-99, 254, 257, 292, 368, 432
<i>Diospyros Virginiana</i> , var. <i>concolor</i>	104	<i>Erythrina pisicupula</i>	57
<i>Diospyros Virginiana</i> , var. <i>macrocarpa</i>	104	Eugenia.....	10, 14
<i>Diospyros Virginiana</i> , var. <i>microcarpa</i>	104	<i>Eugenia axillaris</i>	89
		<i>Eugenia Baruensis</i>	89

	Page.
<i>Eugenia buxifolia</i>	88, 249, 253, 257, 288, 366, 430
<i>Eugenia dichotoma</i>	88, 249, 288
<i>Eugenia dichotoma</i> , var. <i>fragrans</i>	88
<i>Eugenia divaricata</i>	88
<i>Eugenia longipes</i>	89, 249, 288
<i>Eugenia montana</i>	88
<i>Eugenia monticola</i>	89, 249, 253, 257, 288, 366, 430
<i>Eugenia myrtilodes</i>	88
<i>Eugenia pallens</i>	88
<i>Eugenia procera</i>	89, 249, 253, 257, 288, 366, 430
<i>Eugenia pungens</i>	88
<i>Eugenia triplinervia</i> (<i>Eugenia buxifolia</i>)	88
<i>Eugenia triplinervia</i> (<i>Eugenia monticola</i>)	89
<i>Euonymus</i>	10, 12, 18
<i>Euonymus atropurpureus</i>	38, 249, 272
<i>Euonymus Carolinensis</i>	38
<i>Euonymus latifolius</i>	38
EUPHORBIAE	120, 121, 254, 257, 302, 374, 440
<i>Ezocarya lucida</i>	121
<i>Exostemma</i>	10, 14
<i>Exostemma Caribaeum</i>	95, 249, 253, 257, 265, 290, 366, 432
<i>Ezotheca oblongifolia</i>	45
Experiments by Mr. S. P. Sharples	247, 251, 264, 265
<i>Eysenhardtia</i>	10, 12, 14
<i>Eysenhardtia amorphoides</i>	55
<i>Eysenhardtia amorphoides</i> , var. <i>orthocarpa</i>	55
<i>Eysenhardtia orthocarpa</i>	13, 55, 249, 278
F.	
<i>Fagara fraxinifolia</i>	30
<i>Fagara lentiscifolia</i>	81
<i>Fagara Pterota</i>	81
<i>Fagus</i>	11, 16
<i>Fagus alba</i>	157
<i>Fagus Americana</i>	157
<i>Fagus Americana latifolia</i>	157
<i>Fagus Castanea</i>	157
<i>Fagus Castanea dentata</i>	157
<i>Fagus Castanea pumila</i>	156
<i>Fagus ferruginea</i>	157, 249, 255, 258, 262, 322, 350, 390, 456
<i>Fagus ferruginea</i> , var. <i>Caroliniana</i>	157
<i>Fagus pumila</i>	156
<i>Fagus pumila</i> , var. <i>præcox</i>	156
<i>Fagus sylvatica</i>	157
<i>Fagus sylvatica atropurpurea</i>	157
<i>Fagus sylvatica</i> , var. <i>Americana</i>	157
<i>Fagus sylvestris</i>	157
Fan-leaf Palm	217, 348, 414, 480
Farkleberry	96, 292, 432
Fetid Buckeye	42, 274, 358, 423
<i>Ficus</i>	11, 15
<i>Ficus aurea</i>	126, 251, 254, 257, 306, 376, 442
<i>Ficus aurea</i> , var. <i>latifolia</i>	126
<i>Ficus brevifolia</i>	127, 250, 306
<i>Ficus complicata</i>	127
<i>Ficus pedunculata</i>	127, 250, 254, 257, 306, 376, 442
Fiddlewood	116, 306, 372, 436
Fig, Wild	127, 306, 376, 442
Fir, Balm of Gilead	211, 346, 412, 478
Fir, Balsam (<i>Abies balsamea</i>)	211, 346, 412, 478
Fir, Balsam (<i>Abies concolor</i>)	213, 346, 412, 478
Fir, Douglas	209, 263, 344, 352, 410, 476
Fir, Red (<i>Abies magnifica</i>)	214, 346, 412, 478
Fir, Red (<i>Abies nobilis</i>)	214, 346, 412, 478, 573
Fir, Red (<i>Pseudotsuga Douglasii</i>)	209, 265, 344, 352, 410, 476, 565, 575
Fir, Red, the most important timber tree of the Pacific region	7
Fir, White (<i>Abies concolor</i>)	213, 346, 412, 478
Fir, White (<i>Abies grandis</i>)	212, 346, 412, 478
Fir, Yellow	209, 265, 344, 352, 410, 476, 576
Fires, effect of, upon the forests of Tennessee	546
Fires, forest, during the census year, causes of, areas burned over, and losses entailed by (<i>see, also</i> , under state headings)	491, 492
Fires, forest, in Michigan, destructiveness of	550
Fires, forest, forests, standing timber, etc., extent of, in the—	
North Atlantic division	494-510
Northern Central division	547-563

	Page.
Fires, forest, forests, standing timber, etc., extent of, in the—continued.	
South Atlantic division	511-522
Southern Central division	524-546
Western division	564-580
Fires, forest, forests, standing timber, etc., statistics of, in—	
Alabama	491, 524-530
Alaska	580
Arizona	491, 568, 569
Arkansas	491, 543, 544
California	491, 578-580
Colorado	491, 567, 568
Connecticut	504, 701
Dakota	491, 561, 562
Delaware	491, 511
District of Columbia	511
Florida	491, 520-523
Georgia	491, 519, 520
Idaho	491, 571-573
Illinois	491, 547-550
Indiana	491, 547
Indian territory	491, 542
Iowa	491, 569
Kansas	491, 562, 563
Kentucky	491, 545, 546
Louisiana	491, 526-40
Maine	491, 494-496
Maryland	491, 511
Massachusetts	491, 500, 501
Michigan	491, 550-554
Minnesota	491, 558-560
Mississippi	491, 530-536
Missouri	491, 500, 501
Montana	491, 564-568
Nebraska	502
Nevada	491, 571
New Hampshire	491, 496-498
New Jersey	491, 506
New Mexico	491, 506
New York	491, 501-506
North Carolina	491, 515-518
Ohio	491, 547
Oregon	491, 576-578
Pennsylvania	491, 506-510
Rhode Island	491, 500, 501
South Carolina	491, 518, 519
Tennessee	492, 544, 545
Texas	492, 540-543
Utah	492, 569-571
Vermont	492, 496-500
Virginia	492, 511, 512
Washington	492, 573-576
West Virginia	492, 512-515
Wisconsin	492, 564-568
Wyoming	492, 566, 567
Fires, forest, remarks respecting prevalence of and losses occasioned by	491-493
Florida, lumber industry and manufactures from wood in	486, 487, 521-523
Florida, Pencil Cedar of	522
Florida, production of naval stores in	517
Florida, rank of, according to value of lumber products	487
Florida, remarks by Dr. Charles Mohr on the forests of	522, 523
Florida, remarks by Mr. A. H. Curtiss on the forests of	521, 523
Florida, Semi-tropical Forest of	6
Florida, Southern Pine Belt in	520
Florida, statistics of forests, forest fires, standing timber, etc., in	491, 520-523
Florida, tabular statement of the amount of long-leaved pine standing in the forests of, May 31, 1880	521
Florida, the Big Cypress region of	522
Flowering Dogwood (<i>Cornus florida</i>)	91, 288, 306, 430
Flowering Dogwood (<i>Cornus Nuttallii</i>)	91, 288, 306, 432
<i>Fetidaurus montana</i>	186
<i>Fetidaurus Myristica</i>	186
Forestiera	11, 12, 15
<i>Forestiera acuminata</i>	112, 250, 254, 257, 298, 372, 436
Forest crop of the United States for the census year, value of the	485
Forest, Deciduous, of the Mississippi Basin and the Atlantic Plain	4
Forest fires during the census year, causes of, areas burned over, and losses entailed by (<i>see, also</i> , under state headings)	491, 492
Forest fires, remarks respecting prevalence of and losses occasioned by	491-493
Forest, Interior, in the Pacific region, extent of the	8-10
Forest, Mexican, of southern Texas, extent of the	6
Forest of Florida, Semi-tropical	6
Forest Trees, Catalogue of	17-219
Forest Trees, Index to Catalogue of	220-243
Forests, forest fires, standing timber, etc., extent of, in the—	
North Atlantic division	494-510
Northern Central division	547-563
South Atlantic division	511-523
Southern Central division	524-546
Western division	564-580
Forests, forest fires, standing timber, etc., statistics of, in—	
Alabama	491, 524-530
Alaska	580
Arizona	491, 568, 569
Arkansas	491, 543, 544
California	491, 578-580

GENERAL INDEX.

591

	Page.		Page.
Forests, forest fires, standing timber, etc., statistics of, in—continued.		<i>Fraxinus coriacea</i> (<i>Fraxinus Americana</i> , var. <i>Texensis</i>)	108
Colorado	491, 567, 568	<i>Fraxinus coriacea</i> (<i>Fraxinus pistaciifolia</i>)	108
Connecticut	491, 500, 501	<i>Fraxinus crispata</i>	111
Dakota	491, 561, 562	<i>Fraxinus Curtissii</i>	108
Delaware	491, 511	<i>Fraxinus curvidens</i>	110
District of Columbia	491, 511	<i>Fraxinus cuspidata</i>	112
Florida	491, 520-523	<i>Fraxinus dipetala</i>	112
Georgia	491, 519, 520	<i>Fraxinus discolor</i>	107
Idaho	491, 571-573	<i>Fraxinus elliptica</i>	112
Illinois	491, 547-550	<i>Fraxinus eptiptera</i>	107
Indiana	491, 547	<i>Fraxinus excelsior</i>	110
Indian territory	491, 543	<i>Fraxinus expansa</i>	109
Iowa	491, 560	<i>Fraxinus fusca</i>	112
Kansas	491, 562, 563	<i>Fraxinus grandifolia</i>	111
Kentucky	491, 545, 546	<i>Fraxinus Greggii</i>	106, 249, 251, 296
Louisiana	491, 530-540	<i>Fraxinus juglandifolia</i> (<i>Fraxinus Americana</i>)	107
Maine	491, 494-496	<i>Fraxinus juglandifolia</i> (<i>Fraxinus viridis</i>)	109
Maryland	491, 511	<i>Fraxinus juglandifolia</i> , var. <i>serrata</i>	107
Massachusetts	491, 500, 501	<i>Fraxinus juglandifolia</i> , var. <i>subintegerrima</i>	109
Michigan	491, 550-554	<i>Fraxinus juglandifolia</i> , var. <i>subserata</i>	107
Minnesota	491, 558-560	<i>Fraxinus lancea</i>	107
Mississippi	491, 530-536	<i>Fraxinus longifolia</i>	108
Missouri	491, 560, 561	<i>Fraxinus mixta</i>	112
Montana	491, 564-566	<i>Fraxinus nervosa</i>	110
Nebraska	491, 562	<i>Fraxinus nigra</i>	112
Nevada	491, 571	<i>Fraxinus nigra</i> (<i>Fraxinus pubescens</i>)	108
New Hampshire	491, 490-498	<i>Fraxinus nigra</i> (<i>Fraxinus sambucifolia</i>)	111
New Jersey	491, 506	<i>Fraxinus nigrescens</i>	110
New Mexico	491, 568	<i>Fraxinus Nova-Angliae</i> (<i>Fraxinus sambucifolia</i>)	111
New York	491, 501-506	<i>Fraxinus Nova-Angliae</i> (<i>Fraxinus viridis</i>)	109
North Carolina	491, 515-518	<i>Fraxinus Nuttallii</i>	110
Ohio	491, 547	<i>Fraxinus oblongocarpa</i>	108
Oregon	491, 570-578	<i>Fraxinus Oregana</i>	111, 250, 254, 257, 298, 372, 438
Pennsylvania	491, 506-510	<i>Fraxinus ovata</i>	112
Rhode Island	491, 500, 501	<i>Fraxinus pallida</i>	110
South Carolina	491, 518, 519	<i>Fraxinus pannonia</i>	112
Tennessee	492, 544, 545	<i>Fraxinus pauciflora</i>	110
Texas	492, 540-543	<i>Fraxinus Pennsylvanica</i>	108
Utah	492, 568-571	<i>Fraxinus pistaciifolia</i>	113, 106, 249, 254, 257, 296, 370, 434
Vermont	492, 498-500	<i>Fraxinus pistaciifolia</i> (<i>Fraxinus Americana</i> , var. <i>Texensis</i>)	108
Virginia	492, 511, 512	<i>Fraxinus pistaciifolia</i> , var. <i>coriacea</i>	108
Washington	492, 573-576	<i>Fraxinus platycarpa</i>	110, 251, 254, 257, 298, 370, 436
West Virginia	492, 512-515	<i>Fraxinus pubescens</i>	108, 250, 254, 257, 296, 370, 436
Wisconsin	492, 554, 558	<i>Fraxinus pubescens</i>	110
Wyoming	492, 566, 567	<i>Fraxinus pubescens</i> , var.	111
Forests, mountain, of California, injury by grazing cattle to the	570	<i>Fraxinus pubescens</i> , var. <i>latifolia</i>	108
Forests of North America, general remarks on the	3-16	<i>Fraxinus pubescens</i> , var. <i>longifolia</i>	108
Forests of the central pine hills of Mississippi	534	<i>Fraxinus pubescens</i> , var. <i>subpubescens</i>	108
Forests of the Chattahoochee, mixed forest growth, etc., in eastern Alabama	527, 528	<i>Fraxinus pulverulenta</i>	112
Forests of the Northern Pine Belt once extended over the state of Maine	494	<i>Fraxinus quadrangularis</i>	110
Forests of the Tennessee valley in Alabama	528, 529	<i>Fraxinus quadrangulata</i>	110, 249, 254, 257, 290, 298, 370, 436
Forests of the United States in their economic aspects	483-590	<i>Fraxinus quadrangulata</i> , var. <i>nervosa</i>	110
Forests of the Yazoo delta in Mississippi	535, 536	<i>Fraxinus Richardi</i>	112
Forests of western Mississippi	534, 535	<i>Fraxinus rubicunda</i>	112
Forests on Indian reservations in Minnesota	559, 560	<i>Fraxinus rufa</i>	112
Forked-leaf Black Jack	151, 320, 388, 454	<i>Fraxinus sambucifolia</i>	111, 250, 254, 257, 298, 372, 438
Foxtail Pine	191, 336, 402, 468	<i>Fraxinus sambucifolia</i> , var. <i>crispata</i>	111
<i>Frangula Californica</i>	40	<i>Fraxinus Schiedeana</i> , var. <i>parvifolia</i>	106
<i>Frangula Californica</i> , var. <i>tomentella</i>	41	<i>Fraxinus subvillosa</i>	108
<i>Frangula Caroliniana</i>	40	<i>Fraxinus tetragona</i>	110
<i>Frangula fragilis</i>	40	<i>Fraxinus tomentosa</i>	108
<i>Frangula Purshiana</i>	41	<i>Fraxinus trialata</i>	109
Franklinia	26	<i>Fraxinus triptera</i>	110
<i>Franklinia Alatomaha</i>	25	<i>Fraxinus velutina</i>	106
<i>Fraxinus</i>	11, 12, 15	<i>Fraxinus viridis</i>	12, 106, 249, 251, 254, 257, 290, 298, 370, 436
<i>Fraxinus acuminata</i>	107	<i>Fraxinus viridis</i> , var. <i>Berlandieriana</i>	109, 250, 298
<i>Fraxinus alba</i>	112	Fremontia	10, 12, 13
<i>Fraxinus alba</i> (<i>Fraxinus Americana</i>)	107	Fremontia Californica	26, 249, 268
<i>Fraxinus albicans</i> (<i>Fraxinus Americana</i> , var. <i>microcarpa</i>)	108	Frigolito	58, 278
<i>Fraxinus albicans</i> (<i>Fraxinus Americana</i> , var. <i>Texensis</i>)	108	Fringe Tree	113, 298
<i>Fraxinus Americana</i>	107, 250, 251, 254, 257, 290, 296, 350, 370, 436	Fuel in manufactures during the census year, amount and value of wood used as	489
<i>Fraxinus Americana</i> (<i>Fraxinus platycarpa</i>)	110	Fuel value, actual, of some of the more important woods of the United States	350-353
<i>Fraxinus Americana</i> , var. <i>Caroliniana</i>	110	Fuel value of woods, method of determining the	247, 251, 252
<i>Fraxinus Americana</i> , var. <i>juglandifolia</i>	109		
<i>Fraxinus Americana</i> , var. <i>latifolia</i>	107		
<i>Fraxinus Americana</i> , var. <i>microcarpa</i>	108		
<i>Fraxinus Americana</i> , var. <i>pubescens</i>	108		
<i>Fraxinus Americana</i> , var. <i>quadrangulata</i>	110		
<i>Fraxinus Americana</i> , var. <i>quadrangulata nervosa</i>	110		
<i>Fraxinus Americana</i> , var. <i>sambucifolia</i>	111		
<i>Fraxinus Americana</i> , var. <i>Texensis</i>	108, 249, 254, 257, 296, 370, 414, 436		
<i>Fraxinus Americana</i> , var. <i>triptera</i>	110		
<i>Fraxinus anomala</i>	13, 106, 249, 296		
<i>Fraxinus Berlandieriana</i>	109		
<i>Fraxinus Canadensis</i>	107		
<i>Fraxinus Caroliniana</i> (<i>Fraxinus platycarpa</i>)	110		
<i>Fraxinus Caroliniana</i> (<i>Fraxinus viridis</i>)	109		
<i>Fraxinus Carolinensis</i>	107		
<i>Fraxinus cinerea</i>	112		

G.	Page.		Page.
<i>Gardenia clusiae-folia</i>	95	Gum, Tupelo.....	93, 290, 350, 366, 432
Geiger Tree.....	113, 300	Gumbo Limbo.....	33, 270, 356, 429
Genera, distribution of.....	10	Gurgeon Stopper.....	83, 283, 366, 430
General remarks on the forests of North America.....	3-16	GUTTIFERÆ.....	23
Genipa.....	10, 14	<i>Gymnanthes lucida</i>	121
<i>Genipa clusiae-folia</i>	95, 249, 290	<i>Gymnabalanus Oatesbyana</i>	119
Georgia Bark.....	95, 290, 368, 432	<i>Gymnocladus</i>	10, 11, 14
Georgia, lumber industry and manufactures from wood in.....	486, 487, 510, 520	<i>Gymnocladus Canadensis</i>	53, 249, 253, 256, 280, 360, 426
Georgia Pine.....	202, 342, 352, 406, 416, 472	<i>Gymnocladus dioica</i>	58
Georgia, production of naval stores in.....	517		
Georgia, rank of, according to value of lumber products.....	487	III.	
Georgia, statistics of forests, forest fires, standing timber, etc., in.....	491, 519, 520	Hackberry (<i>Celtis occidentalis</i>).....	126, 306, 376, 414, 442
Georgia, tabular statement of the amount of Long-leaved Pine standing in the forests of, May 31, 1880.....	520	Hackberry (<i>Celtis occidentalis</i> , var. <i>reticulata</i>).....	126, 306, 376, 412
Georgia, the Maritime Pine Belt in.....	519	Hackmatack.....	215, 346, 352, 412, 478
Giant Cactus.....	90, 288	Halesia.....	11, 15
<i>Gigantabies taxifolia</i>	185	<i>Halesia diptera</i>	105, 250, 254, 257, 294, 563, 434
<i>Gigantabies Wellingtonia</i>	184	<i>Halesia parviflora</i>	106
Ginger Pine.....	179, 332, 350, 368, 464	<i>Halesia reticulata</i>	105
Glaberry.....	28, 268, 356, 420	<i>Halesia tetraptera</i>	106, 250, 296
Glaucous Willow.....	169, 328	<i>Halmia cornifolia</i>	80
<i>Gleditschia</i>	10, 11, 14	<i>Halmia flabellata</i>	77
<i>Gleditschia aquatica</i>	59	<i>Halmia lobata</i>	79
<i>Gleditschia brachycarpa</i>	59	<i>Halmia punctata</i>	80
<i>Gleditschia Carolinensis</i>	59	<i>Halmia tomentosa</i>	79
<i>Gleditschia elegans</i>	59	HAMAMELACEÆ.....	85, 86, 253, 256, 286, 364, 414, 430
<i>Gleditschia inermis</i> (<i>Gleditschia monosperma</i>).....	59	Hamamelis.....	10, 11, 14
<i>Gleditschia inermis</i> (<i>Gleditschia triacanthos</i> , var. <i>inermis</i>).....	59	<i>Hamamelis androgyna</i>	85
<i>Gleditschia macrantha</i>	59	<i>Hamamelis corylifolia</i>	85
<i>Gleditschia Meliloba</i>	59	<i>Hamamelis dioica</i>	85
<i>Gleditschia monosperma</i>	59, 249, 253, 256, 280, 362, 426	<i>Hamamelis macrophylla</i>	85
<i>Gleditschia spinosa</i>	59	<i>Hamamelis parvifolia</i>	85
<i>Gleditschia triacanthos</i>	59	<i>Hamamelis Virginiana</i> , var. <i>parvifolia</i>	85
<i>Gleditschia triacanthos</i> , var. <i>brachycarpus</i>	59	<i>Hamamelis Virginica</i>	85, 249, 286
<i>Gleditschia triacanthos</i> , var. <i>inermis</i>	59	Hard Maple.....	48, 276, 358, 422, 576
<i>Gleditschia triacanthos</i> , var. <i>monosperma</i>	59	Hard Pine.....	202, 342, 352, 406, 416, 472
Goose-foot Maple.....	46, 274	Haw, Apple.....	82, 286, 364, 430
Gopher Plum.....	91, 200, 366, 432	Haw, Black.....	94, 290, 366, 432
Gopherwood.....	87, 278, 360, 426	Haw, Hogs.....	75, 284
Gorlonia.....	10, 11, 13	Haw, May.....	82, 286, 364, 430
<i>Gordonia Franklini</i>	25	Haw, Paralely.....	81, 286
<i>Gordonia Lasianthus</i>	25, 250, 253, 256, 265, 268, 354, 418	Haw, Pear.....	79, 286, 364, 430
<i>Gordonia pubescens</i>	25, 248	Haw, Purple.....	40, 272
<i>Gordonia pyramidalis</i>	25	Haw, Red (<i>Crataegus coccinea</i>).....	78, 286
Grape, Sea.....	118, 302, 438	Haw, Red (<i>Crataegus flava</i> , var. <i>pubescens</i>).....	83, 286, 364, 430
Gravity, specific, ash, and weight per cubic foot of dry specimens of the woods of the United States.....	266-349	Haw, Scarlet (<i>Crataegus coccinea</i>).....	78, 286
Gray Birch (<i>Betula alba</i> , var. <i>populifolia</i>).....	159, 324, 350, 392, 458	Haw, Scarlet (<i>Crataegus subvillosa</i>).....	78, 286, 364, 430
Gray Birch (<i>Betula lutea</i>).....	161, 324, 392, 458	Haw, Small-fruited.....	81, 286, 364, 430
Gray Pine.....	201, 342, 352, 406, 472	Haw, Summer (<i>Crataegus flava</i>).....	83, 286
Great Laurel.....	99, 292, 368, 434	Haw, Summer (<i>Crataegus flava</i> , var. <i>pubescens</i>).....	83, 286, 364, 430
Great Plains, the.....	4-6	Haw, Yellow.....	83, 286
Green Ash.....	109, 298, 370, 436	Hazel, Witch.....	85, 286
Green-bark Acacia.....	60, 280, 362, 426	Hemlock and White Pine standing in the forests of Pennsylvania May 31, 1880, tabular statement of the amount of.....	506
Ground Ash.....	111, 298, 372, 438	Hemlock (<i>Pseudotsuga Douglasii</i> , var. <i>macrocarpa</i>).....	210, 346, 412, 478
Guaiacum.....	10, 13	Hemlock (<i>Tsuga Canadensis</i>).....	207, 265, 344, 352, 408, 474, 506
<i>Guaiacum angustifolium</i>	29	Hemlock (<i>Tsuga Caroliniana</i>).....	207, 265, 344, 410, 476
<i>Guaiacum sanctum</i>	28, 249, 253, 256, 268, 356, 420	Hemlock (<i>Tsuga Mertensiana</i>).....	208, 265, 344, 410, 476, 573, 576
<i>Guaiacum verticale</i>	28	Heteromeles.....	10, 12, 14
Guettarda.....	10, 15	<i>Heteromeles arbutifolia</i>	83, 249, 286
<i>Guettarda elliptica</i>	96, 249, 290	<i>Heteromeles Fremontiana</i>	83
<i>Guettarda Blodgettii</i>	96	<i>Heyderia decurrens</i>	176
Guiana Plum.....	121, 302, 374, 440	<i>Heymassoli spinosa</i>	34
<i>Guilandina dioica</i>	58	<i>Hickorea</i> , species.....	132
Gum, Black.....	92, 290, 368, 432	<i>Hickorius amara</i>	135
Gum, Cotton.....	93, 290, 350, 366, 432	Hickory, Big-bud.....	134, 310, 350, 380, 444
Gum, Doctor.....	54, 278, 360, 424	Hickory, Black (<i>Carya porcina</i>).....	134, 310, 350, 380, 446
Gum Elastic.....	102, 294, 368, 434	Hickory, Black (<i>Carya tomentosa</i>).....	134, 310, 350, 380, 444
Gum Elemi.....	33, 270, 350, 356, 420	Hickory, Brown.....	134, 310, 350, 380, 446
Gum, Red.....	86, 286, 350, 364, 414, 430	Hickory Elm.....	123, 304, 374, 440
Gum, Sour.....	92, 290, 368, 432	Hickory, Nutmeg.....	135, 310, 350, 380, 446
Gum, Star-leaved.....	86, 286, 350, 364, 414, 430	Hickory Pine (<i>Pinus Balfouriana</i> , var. <i>aristata</i>).....	191, 336, 402, 468
Gum, Sweet.....	86, 286, 350, 364, 414, 430	Hickory Pine (<i>Pinus pungens</i>).....	199, 340, 350, 404, 472
		Hickory, Shag-bark.....	133, 308, 350, 378, 444
		Hickory, Shell-bark.....	133, 308, 350, 378, 444
		Hickory, Swamp (<i>Carya amara</i>).....	135, 310, 350, 380, 446
		Hickory, Swamp (<i>Carya aquatica</i>).....	136, 310, 350, 380, 414, 446

	Page.		Page.
Hickory, Switch-bud	184, 810, 350, 380, 446	Incense Cedar	176, 380, 396, 462
Hickory, Water	186, 810, 350, 380, 414, 446	Index to Catalogue of Forest Trees	220-243
Hickory, White-heart	134, 810, 350, 380, 444	Indiana, lumber industry and manufactures from wood in	486, 487, 547
<i>Hicorius integrifolia</i>	185	Indiana, rank of, according to value of lumber products	487, 547
<i>Hierophyllum Cassine</i>	36	Indiana, statistics of forests, forest fires, standing timber, etc., in	491, 547
Hippomane	11, 15	Indian Bean	115, 300, 372, 438
Hippomane Mancinella	121, 250, 304, 400	Indian Cherry	40, 272, 358, 422
Hoary Alder	165, 326, 394, 400	Indian territory, statistics of forests, forest fires, standing timber, etc., in	491, 543
Hog Plum (<i>Prunus angustifolia</i>)	66, 282, 362, 426	India-rubber Tree	127, 306, 376, 442
Hog Plum (<i>Rhus Metopium</i>)	54, 278, 360, 424	Industry, lumber, commercial importance of Chicago with reference to the	548
Hog Plum (<i>Ximenia Americana</i>)	84, 270	Industry, lumbering, of the United States, average number of hands employed in the	486
Hogs' Haw	75, 284	Industry, lumbering, of the United States, general remarks on	485-493
Holly, American	35, 270, 356, 420	Industry, lumbering, of the United States, statistics of, for year ending May 31, 1890	486, 487
Holly, California	84, 286	<i>Inga forbesii</i>	64
Holly, Dahoon	35, 270, 356, 420	<i>Inga Guadalupeensis</i>	64
Honey Locust (<i>Gleditsia triacanthos</i>)	59, 280, 360, 426	<i>Inga microphylla</i>	64
Honey Locust (<i>Prosopis juliflora</i>)	62, 265, 280, 350, 362, 426	<i>Inga rosea</i>	64
Honey Pod	62, 265, 280, 350, 362, 426	<i>Inga Unguis-cati</i>	64
Honey Shucks	59, 280, 360, 426	Inkwood	45, 274, 358, 422
Hoop Ash	111, 238, 372, 438	Interior Forest in the Pacific region, extent of the	8-10
<i>Hopsea tinctoria</i>	105	Iowa, lumber industry and manufactures from wood in	486, 487, 500
Hop Hornbeam	158, 322, 390, 456	Iowa, rank of, according to value of lumber products	487, 500
Hop Tree	81, 270	Iowa, statistics of forests, forest fires, standing timber, etc., in	491, 500
Hornbeam	158, 322, 392, 456	<i>Isoxylon pomiferum</i>	128
Hornbeam, Hop	158, 322, 390, 456	Iron oak	139, 312, 362, 448
Horse Plum	65, 282, 362, 436	Ironwood (<i>Bumelia lycioides</i>)	103, 294, 368, 434
Horse Sugar	105, 284, 368, 434	Ironwood (<i>Carpinus Caroliniana</i>)	159, 322, 392, 456
Hypelate	10	Ironwood (<i>Cliftonia ligustrina</i>)	38, 272, 356, 420
<i>Hypelate oblongifolia</i>	45	Ironwood (<i>Cyrilla racemiflora</i>)	37, 272, 356
<i>Hypelate paniculata</i>	45, 249, 272, 358, 422	Ironwood (<i>Hypelate paniculata</i>)	45, 274, 358, 422
<i>Hypelate trifoliata</i>	45, 249, 272, 422	Ironwood (<i>Oleyna Tesota</i>)	56, 278, 360, 426
<i>Hyperanthus dioica</i>	58	Ironwood (<i>Ostrya Virginica</i>)	158, 322, 390, 456
<i>Hypericum Lasianthus</i>	25	Ironwood, Black	39, 272, 358, 422
I.		Ironwood, Red	39, 272, 358, 422
Idaho, lumber industry and manufactures from wood in	486, 487, 572	Ironwood, White	45, 274, 422
Idaho, rank of, according to value of lumber products	487	Islay	70, 284, 364, 428
Idaho, remarks by Mr. Sereno Watson on the forests of	572, 578	<i>Itea Cyrtilla</i>	87
Idaho, statistics of forests, forest fires, standing timber, etc., in	491, 571-578	Ivy	98, 292, 368, 434
Ilex	10, 11, 13	J.	
<i>Ilex aestivalis</i>	37	Jack, Black (<i>Quercus Catesbeii</i>)	151, 320, 388, 454
<i>Ilex ambiguus</i>	37	Jack, Black (<i>Quercus nigra</i>)	150, 265, 320, 350, 388, 454
<i>Ilex angustifolia</i>	35	Jack; Blue	153, 320, 390, 456
<i>Ilex aquifolium</i>	35	Jack, Forked-leaved Black	151, 320, 388, 454
<i>Ilex Canadensis</i>	35	Jack Oak	150, 265, 320, 350, 388, 454
<i>Ilex Cassena</i>	36	Jack, Sand	153, 320, 390, 456
<i>Ilex Cassine</i>	36, 249, 273	Jacquinia	11, 15
<i>Ilex Cassine</i> (<i>Ilex Dahoon</i>)	35	<i>Jacquinia armillaris</i>	100, 249, 292
<i>Ilex Cassine, B.</i>	36	Jamaica Dogwood	57, 278, 360, 426
<i>Ilex Cassine, var. angustifolia</i>	35	Jersey Pine	199, 340, 350, 404, 470, 546
<i>Ilex Cassine, var. latifolia</i>	35	Joe wood	100, 292
<i>Ilex cassinoides</i>	35	Joshua, The	219, 348
<i>Ilex Dahoon</i>	35, 250, 253, 256, 270, 356, 420	Joshua Tree	219, 348
<i>Ilex Dahoon, var. angustifolia</i>	35	Judas Tree	61, 280, 362, 426
<i>Ilex Dahoon, var. myrtifolia</i>	36, 250, 370	JUGLANDACEÆ	130-136, 254, 257, 306, 376, 414, 442
<i>Ilex decidua</i>	37, 249, 273	Juglans	11, 12, 15, 251
<i>Ilex Floridana</i>	36	<i>Juglans alba</i> (<i>Carya alba</i>)	133
<i>Ilex laurifolia</i>	35	<i>Juglans alba</i> (<i>Carya tomentosa</i>)	133
<i>Ilex lasiocarpa</i>	35	<i>Juglans alba acuminata</i>	134
<i>Ilex ligustrifolia</i>	36	<i>Juglans alba minima</i>	135
<i>Ilex ligustrina</i> (<i>Ilex Cassine</i>)	36	<i>Juglans alba ovata</i>	132
<i>Ilex ligustrina</i> (<i>Ilex Dahoon, var. angustifolia</i>)	36	<i>Juglans amara</i>	135
<i>Ilex myrtifolia</i>	36	<i>Juglans angustifolia</i> (<i>Carya amara</i>)	135
<i>Ilex opaca</i>	34, 250, 253, 256, 270, 356, 420	<i>Juglans angustifolia</i> (<i>Carya oliviformis</i>)	132
<i>Ilex prinoides</i>	37	<i>Juglans aquatica</i>	135
<i>Ilex quercifolia</i>	35	<i>Juglans Californica</i>	131
<i>Ilex religiosa</i>	36	<i>Juglans cathartica</i>	130
<i>Ilex rosmarifolia</i>	36	<i>Juglans cinerea</i>	130, 250, 254, 257, 260, 306, 376, 442
<i>Ilex vomitoria</i>	36	<i>Juglans compressa</i>	132
<i>Ilex Watsoniana</i>	36	<i>Juglans cordiformis</i>	135
ILICINÆ	34, 253, 256, 270, 356, 420	<i>Juglans cylindrica</i>	132
Illinois, lumber industry and manufactures from wood in	486, 487, 548-550	<i>Juglans exaltata</i>	132
Illinois Nut	132, 308, 350, 378, 444	<i>Juglans glabra</i>	134
Illinois, rank of, according to value of lumber products	487, 548		
Illinois, statistics of forests, forest fires, standing timber, etc., in	491, 547-550		

[illegible]

	Page.		Page.
<i>Liriodendron Tulipifera</i>	22, 250, 253, 256, 259, 266, 350, 354, 418	Lumber industry and manufactures from wood in—continued.	
Live Oak (<i>Quercus chrysolepis</i>)	146, 318, 386, 452	New Mexico	486, 487, 568
Live Oak (<i>Quercus virens</i>)	145, 265, 316, 386, 452	New York	486, 487, 502-505
Live Oak (<i>Quercus Wializeni</i>)	147, 318, 386, 452	North Carolina	486, 487, 515
Live Oak, Coast	147, 318, 386, 452	Ohio	486, 487, 547
Loblolly Bay	25, 265, 268, 354, 418	Oregon	486, 487, 577
Loblolly, Long- and Short-leaved Pine standing in the forests of Texas		Pennsylvania	486, 487, 506-510
May 31, 1880, tabular statement of the amount of	541	Rhode Island	486, 487, 500, 501
Loblolly Pine	197, 340, 350, 404, 416, 470, 516, 541	South Carolina	486, 487, 518, 519
Locust (<i>Robinia Neo-Mexicana</i>)	54, 278, 360, 426	Tennessee	486, 487, 545
Locust (<i>Robinia Pseudacacia</i>)	55, 278, 350, 360, 414, 424	Texas	486, 487, 541, 542
Locust, Black (<i>Gleditsia triacanthos</i>)	59, 280, 360, 426	Utah	486, 487, 569
Locust, Black (<i>Robinia Pseudacacia</i>)	55, 278, 350, 360, 414, 424	Vermont	486, 487, 498-500
Locust, Clammy	56, 278	Virginia	486, 487, 512
Locust, Honey (<i>Gleditsia triacanthos</i>)	59, 280, 360, 426	Washington	486, 487, 574
Locust, Honey (<i>Prosopis juliflora</i>)	62, 265, 280, 350, 362, 426	West Virginia	486, 487, 512-515
Locust, Sweet	59, 280, 360, 426	Wisconsin	486, 487, 554, 556
Locust, Water	60, 280, 362, 426	Wyoming	486, 487, 567
Locust, Yellow	55, 278, 350, 360, 414, 424	Lumber industry, commercial importance of Chicago with reference to the	548
Lodge-pole Pine	195, 338, 350, 404, 470, 504, 574, 577	Lumber industry of the Saginaw valley in Michigan	552
Logs rafted out of the Susquehanna boom at Williamsport, Pennsylv-		Lumber industry of the United States, general remarks on	485-493
vania, from 1862 to 1880, number of	508	Lumber market, importance of Burlington, Vermont, as a	499
Logwood	40, 272	Lumber market, rank of Albany, New York, as a	503
Long- and Short-leaved Pine standing in the forests of Alabama May 31,		Lumber trade of Chicago, early	549, 550
1880, tabular statement of the amount of	524	Lumbering industry of the United States, average number of hands	
Long- and Short-leaved Pine standing in the forests of Louisiana May 31,		employed in	486
1880, tabular statement of the amount of	537	Lumbering industry of the United States, statistics of, for year ending	
Long- and Short-leaved Pine standing in the forests of Mississippi May		May 31, 1880	486, 487
31, 1880, tabular statement of the amount of	581	Lumbermen of Washington territory, wasteful methods of	574
Long-leaved Cucumber Tree	22, 266, 354, 418	<i>Lyonia arborea</i>	98
Long-leaved, Loblolly, and Short-leaved Pine standing in the forests of		<i>Lyonia ferruginea</i>	96
Texas May 31, 1880, tabular statement of the amount of	541	<i>Lyonia rigida</i>	96
Long-leaved Pine	202, 342, 352, 406, 416, 472, 516, 519-521, 524, 531, 537, 541	Lysiloma	10, 14
Long-leaved Pine standing in the forests of Florida May 31, 1880, tabular		<i>Lysiloma Bahamensis</i>	64
statement of the amount of	521	<i>Lysiloma latifolia</i>	64, 250, 282, 362, 426
Long-leaved Pine standing in the forests of Georgia May 31, 1880, tabular			
statement of the amount of	520	Maclura	11, 15
Long-leaved Pine standing in the forests of North Carolina May 31, 1880,		<i>Maclura aurantiaca</i>	128, 249, 254, 257, 306, 376, 442, 542
tabular statement of the amount of	516	Madeira	34, 270, 350, 356, 420
Long-leaved Pine standing in the forests of South Carolina May 31, 1880,		Madrofia	97, 292, 368, 432
tabular statement of the amount of	519	Magnolia	10, 11, 13
Long-leaved Pine, the characteristic tree of the Southern Maritime Pine		<i>Magnolia acuminata</i>	20, 250, 253, 256, 259, 266, 354, 418
Belt	4	<i>Magnolia auricularis</i>	22
Louisiana, lumber industry and manufactures from wood in	486, 487, 536, 537	<i>Magnolia auriculata</i>	22
Louisiana, moses-ginning industry in	537, 538	<i>Magnolia cordata</i>	20, 250, 253, 256, 266, 354, 418
Louisiana, production of naval stores in	517, 536, 537	<i>Magnolia De Candollei</i>	20
Louisiana, rank of, according to value of lumber products	487	<i>Magnolia fragrans</i>	20
Louisiana, remarks by Dr. Charles Mohr on the forests of	538-540	<i>Magnolia Fraseri</i>	22, 250, 253, 256, 266, 354, 418
Louisiana, statistics of forests, forest fires, standing timber, etc., in	491, 536, 540	<i>Magnolia glauca</i>	19, 250, 253, 256, 266, 354, 414, 418
Louisiana, tabular statement of the amount of Long- and Short-leaved		<i>Magnolia glauca</i> , var. <i>latifolia</i>	20
Pine standing in the forests of, May 31, 1880	537	<i>Magnolia glauca</i> , var. <i>longifolia</i>	20
Lumber and shingles received at Chicago during the year 1880, amount of	548, 549	<i>Magnolia grandiflora</i>	19, 250, 253, 256, 266, 354, 414, 418, 534
Lumber for Colorado, Utah, and New Mexico, Chicago the principal		<i>Magnolia grandiflora</i> , var. <i>elliptica</i> and <i>obovata</i>	19
source of supply of	568, 569	<i>Magnolia grandiflora</i> , var. <i>lanceolata</i>	19
Lumber industry and manufactures from wood in—		<i>Magnolia longifolia</i>	20
Alabama	486, 487, 524, 525	<i>Magnolia macrophylla</i>	21, 250, 253, 256, 266, 354, 418
Arizona	486, 487, 569	Magnolia, Mountain	20, 266, 354, 418
Arkansas	486, 487, 544	<i>Magnolia pyramidata</i>	22
California	486, 487, 578, 580	<i>Magnolia tripetala</i>	21
Colorado	486, 487, 567, 568	Magnolia Umbrella	21, 250, 253, 256, 266, 354, 418
Connecticut	486, 487, 500, 501	<i>Magnolia Virginiana</i> , var. <i>a. glauca</i>	20
Dakota	486, 487, 561	<i>Magnolia Virginiana</i> , var. <i>β. fastida</i>	19
Delaware	486, 487, 511	<i>Magnolia Virginiana</i> , var. <i>c</i>	20
District of Columbia	486, 487	<i>Magnolia Virginiana</i> , var. <i>tripetala</i>	21
Florida	486, 487, 521, 523	MAGNOLIACEÆ	19-22, 253, 256, 266, 354, 414, 418
Georgia	486, 487, 519, 520	Mahogany	34, 270, 350, 356, 420
Idaho	486, 487, 573	Mahogany Birch	162, 324, 392, 458
Illinois	486, 487, 548-550	Mahogany, Mountain, and the Nut Pine the most important trees of the	
Indiana	486, 487, 547	Interior Forest	9
Iowa	486, 487, 500	Mahogany, Mountain (<i>Cercocarpus ledifolius</i>)	71, 284, 350, 428, 569, 571
Kansas	486, 487, 563	Mahogany, Mountain (<i>Cercocarpus parvifolius</i>)	71, 284, 569, 571
Kentucky	486, 487, 546	Maine, forests of the Northern Pine Belt once extended over the state of	494
Louisiana	486, 487, 536, 537	Maine, lumber industry and manufactures from wood in	486, 487, 494-496
Maine	486, 487, 494-496	Maine, rank of, according to value of lumber products	487
Maryland	486, 487, 511	Maine, statistics of forests, forest fires, standing timber, etc., in	491, 494, 496
Massachusetts	486, 487, 500, 501	Maine, tabular statement of the amount of Pine and Spruce standing in	
Michigan	486, 487, 551, 552	the forests of, May 31, 1880	494
Minnesota	486, 487, 558	MALPIGHIACEÆ	28, 253, 256, 268, 354, 420
Mississippi	486, 487, 531	<i>Malpighia lucida</i>	28
Missouri	486, 487, 500, 501	<i>Malus angustifolia</i>	72
Montana	486, 487, 504	<i>Malus coronaria</i>	72
Nebraska	486, 487, 502		
Nevada	486, 487, 571		
New Hampshire	486, 487, 497, 498		
New Jersey	486, 487, 506		

	Page.		Page.
<i>Malus diversifolia</i>	73	<i>Mespilus elliptica</i> (Cratægus flava, var. pubescens).....	83
<i>Malus microcarpa coronaria</i>	72	<i>Mespilus flabellata</i>	77
<i>Malus rivularis</i>	73	<i>Mespilus flava</i>	82
<i>Malus sempervirens</i>	72	<i>Mespilus flexispina</i>	82
<i>Malus subcordata</i>	73	<i>Mespilus glandulosa</i>	77
Manchineel.....	121, 304	<i>Mespilus hiemalis</i>	83
Manchineel, Mountain.....	54, 278, 300, 424	<i>Mespilus latifolia</i>	79
<i>Mancinella venenata</i>	121	<i>Mespilus linearis</i>	77
Mangrove.....	87, 285, 288, 364, 430	<i>Mespilus lobata</i>	9
Mangrove, Black.....	117, 302	<i>Mespilus lucida</i>	76
Mangrove, White.....	87, 288, 364, 430	<i>Mespilus lucida</i> , var. <i>angustifolia</i>	77
Maple, Ash-leaved.....	51, 276, 360, 424	<i>Mespilus Michauxii</i>	83
Maple, Black Sugar.....	49, 276, 350, 358, 424	<i>Mespilus monogyna</i> , var. <i>apifolia</i>	81
Maple, Broad-leaved.....	47, 274, 358, 422	<i>Mespilus nivea</i>	84
Maple, Dwarf.....	48, 274	<i>Mespilus odorata</i>	79
Maple, Goose-foot.....	48, 274	<i>Mespilus ovalifolia</i>	76
Maple, Hard.....	48, 276, 358, 422, 576	<i>Mespilus Phænopyrum</i>	80
Maple, Mountain.....	48, 274	<i>Mespilus populifolia</i>	78
Maple, Red.....	50, 276, 358, 424	<i>Mespilus pruinosa</i>	79
Maple, Rock.....	48, 276, 358, 422	<i>Mespilus prunellifolia</i>	76
Maple, Silver.....	49, 276, 358, 424	<i>Mespilus prunifolia</i>	77
Maple, Soft (Acer dasycarpum).....	49, 276, 358, 424	<i>Mespilus pubescens</i>	77
Maple, Soft (Acer rubrum).....	50, 276, 358, 424	<i>Mespilus punctata</i>	80
Maple, Striped.....	48, 274	<i>Mespilus pyrifolia</i> (Cratægus tomentosa).....	79
Maple, Sugar.....	48, 276, 358, 422	<i>Mespilus pyrifolia</i> (Cratægus tomentosa, var. <i>punctata</i>).....	80
Maple sugar and molasses produced in the United States in 1879, amount of.....	485	<i>Mespilus rotundifolia</i> (Cratægus coccinea).....	77
Maple, Swamp.....	50, 276, 358, 424	<i>Mespilus rotundifolia</i> (Cratægus Crus-galli, var. <i>prunifolia</i>).....	77
Maple, Vine.....	47, 274, 358, 422	<i>Mespilus salicifolia</i>	76
Maple, Water.....	50, 276, 358, 424	<i>Mespilus spathulata</i>	81
Maple, White.....	49, 276, 358, 424	<i>Mespilus tiliaefolia</i>	78
Maritime Pine Belt in Georgia.....	519	<i>Mespilus turbinata</i>	82
Maritime Pine Belt in Mississippi.....	582	<i>Mespilus viridis</i>	78
Maritime Pine Belt in North Carolina.....	515	<i>Mespilus Watsoniana</i>	76
Maritime Pine Region in Alabama.....	525	<i>Mespilus Wendlandii</i>	77
Marlberry.....	100, 292	Mesquit.....	62, 265, 280, 350, 362, 426
Maryland, lumber industry and manufactures from wood in.....	486, 487, 511	Mesquit, Sorew-pod.....	62, 280, 362, 426
Maryland, rank of, according to value of lumber products.....	487	Mesquit the most important species in the valleys of the Atlantic-Mexican region.....	9
Maryland, statistics of forests, forest fires, standing timber, etc., in.....	491, 511	Method of determining the fuel value of woods.....	247, 251, 252
Massachusetts, lumber industry and manufactures from wood in.....	486, 487, 500, 501	Method of determining the strength of woods.....	252
Massachusetts, rank of, according to value of lumber products.....	487	<i>Metopium Linnæi</i>	54
Massachusetts, statistics of forests, forest fires, standing timber, etc., in.....	491, 500, 501	Mexican Banana.....	219, 348
Massachusetts, the Northern Pine Belt in.....	500	Mexican Forest of southern Texas, extent of.....	6
Mastic.....	101, 292, 368, 434	Mexican Mulberry.....	123, 306
Maul Oak.....	148, 318, 386, 452	Mexican Persimmon.....	105, 294
May Cherry.....	84, 280, 364, 430	<i>Michauxia sessilis</i>	25
May Haw.....	82, 280, 364, 430	Michigan, destructiveness of forest fires in.....	550
Meadow Pine.....	202, 342, 352, 406, 410, 474	Michigan, lumber industry and manufactures from wood in.....	486, 487, 551, 552
MELIACEÆ.....	38, 34, 253, 256, 270, 356, 420	Michigan, lumber industry of the Saginaw valley in.....	552
<i>Melicocca paniculata</i>	45	Michigan, maple-sugar product of.....	551
<i>Melilotus heterophylla</i>	59	Michigan, rank of according to value of lumber products.....	487, 552
<i>Mespilus acerifolia</i>	80	Michigan, remarks by Mr. H. C. Putnam on the forests of.....	553, 554
<i>Mespilus acutivalis</i>	82	Michigan, statistics of forests, forest fires, standing timber, etc., in.....	491, 550-554
<i>Mespilus apifolia</i>	81	Michigan, tabular statement of the amount of White Pine standing in the forests of, May 31, 1880.....	551
<i>Mespilus arborea</i>	84	<i>Mimosa biceps</i>	62
<i>Mespilus arbutifolia</i>	83	<i>Mimosa frondosa</i>	62
<i>Mespilus Azarolus</i>	81	<i>Mimosa glauca</i>	62
<i>Mespilus berberifolia</i>	82	<i>Mimosa Guadalupeensis</i>	64
<i>Mespilus Bosciiana</i>	77	<i>Mimosa latissiliqua</i>	64
<i>Mespilus Calpodendron</i>	79	<i>Mimosa leucocephala</i>	62
<i>Mespilus Canadensis</i>	84	<i>Mimosa rosea</i>	64
<i>Mespilus Canadensis</i> , var. <i>cordata</i>	84	<i>Mimosa Unguis-cati</i>	64
<i>Mespilus Canadensis</i> , var. <i>obovatis</i>	85	Mimusops.....	11, 15
<i>Mespilus Caroliniana</i>	82	<i>Mimusops dissecta</i>	103
<i>Mespilus coccinea</i>	77	<i>Mimusops Sieberi</i>	103, 249, 254, 257, 294, 368, 434
<i>Mespilus cordata</i>	80	Minnesota, forests on Indian reservations in.....	550, 560
<i>Mespilus cornifolia</i>	80	Minnesota, lumber industry and manufactures from wood in.....	486, 487, 558
<i>Mespilus Crus-galli</i>	76	Minnesota, rank of, according to value of lumber products.....	487, 559
<i>Mespilus Crus-galli</i> , var. <i>pyracanthifolia</i>	76	Minnesota, remarks by Mr. H. C. Putnam on the forests of.....	550, 560
<i>Mespilus Crus-galli</i> , var. <i>salicifolia</i>	76	Minnesota, statistics of forests, forest fires, standing timber, etc., in.....	491, 558-560
<i>Mespilus cuneifolia</i> (Cratægus Crus-galli).....	76	Minnesota, tabular statement of the amount of White Pine standing in the forests of, May 31, 1880.....	558
<i>Mespilus cuneifolia</i> (Cratægus tomentosa, var. <i>punctata</i>).....	80	Minnesota, the Northern Pine Belt in.....	558
<i>Mespilus cuneiformis</i>	76	Mississippi Basin and the Atlantic Plain, Deciduous Forest of the.....	4
<i>Mespilus elliptica</i> (Cratægus Crus galli).....	76		

	Page.		Page.
Mississippi, forests of the central pine hills of	534	<i>Myrica cerifera</i> , var. <i>arborescens</i>	136
Mississippi, forests of the Yazoo delta in	535, 536	<i>Myrica cerifera</i> , var. <i>latifolia</i>	136
Mississippi, lumber industry and manufactures from wood in	486, 487, 531	<i>Myrica cerifera</i> , var. <i>media</i>	136
Mississippi, pine forests of the northeastern counties of	532-534	<i>Myrica cerifera</i> , var. <i>pumila</i>	136
Mississippi, production of naval stores in	517, 531, 532, 536	<i>Myrica Pennsylvanica</i>	136
Mississippi, rank of, according to value of lumber products	487	<i>Myrica Xalapensis</i>	137
Mississippi, remarks by Dr. Charles Mohr on the forests of	531-536	MYRICACEÆ	136, 137, 254, 257, 312, 380, 446
Mississippi, southern, pine forests of	531, 532	MYRSINACEÆ	99, 100, 292
Mississippi, statistics of forests, forest fires, standing timber, etc., in	491, 530-536	Myrsine	11, 15
Mississippi, tabular statement of the amount of Long- and Short-leaved Pine standing in the forests of, May 31, 1880	531	<i>Myrsine floribunda</i>	99
Mississippi, the Maritime Pine Belt in	532	<i>Myrsine Floridana</i>	99
Mississippi, western, forests of	534, 535	<i>Myrsine Rapanœ</i>	99, 243, 292
Missouri, lumber industry and manufactures from wood in	486, 487, 560, 561	MYRTACEÆ	88, 89, 253, 257, 288, 366, 430
Missouri, rank of, according to value of lumber products	487, 561	Myrtle, Blue	41, 272
Missouri, statistics of forests, forest fires, standing timber, etc., in	491, 560, 561	Myrtle, Wax	136, 312, 380, 446
Mobile the principal center of wood manufacture in Alabama	525	<i>Myrtus acillaris</i>	88
Mocker Nut	124, 310, 350, 380, 444	<i>Myrtus buxifolia</i>	88
Mock Orange	70, 284, 362, 428	<i>Myrtus Chytraeolia</i>	88
Mohr, Dr. Charles, remarks on the forests and the turpentine industry of Alabama by	525-530	<i>Myrtus dichotoma</i>	88
Mohr, Dr. Charles, remarks on the forests of Florida by	522, 523	<i>Myrtus monticola</i>	89
Mohr, Dr. Charles, remarks on the forests of Louisiana by	533-540	<i>Myrtus Poirœti</i>	88
Mohr, Dr. Charles, remarks on the forests of Mississippi by	531-536	<i>Myrtus procera</i>	89
Mohr, Dr. Charles, remarks on the forests of Texas by	542, 543		
Molasses and sugar, maple, produced in the United States in 1879, amount of	485	N.	
Montana, lumber industry and manufactures from wood in	486, 487, 564	Naked Wood (<i>Colubrina reclinata</i>)	42, 274, 358
Montana, rank of, according to value of lumber products	487	Naked Wood (<i>Eugenia dichotoma</i>)	88, 288
Montana, remarks by Mr. Sereno Watson on the forests of	565, 566	Nannyberry	94, 290, 423
Montana, statistics of forests, forest fires, standing timber, etc., in	491, 564-566	Naval stores, production of, in Alabama	517, 537, 539, 530
Monterey Cypress	179, 332, 398, 464	Naval stores, production of, in Florida	517
Monterey Pine	196, 340, 404, 470	Naval stores, production of, in Georgia	517
Moose Elm	122, 304, 374, 440	Naval stores, production of, in Louisiana	517, 536, 537
Moosewood	46, 274	Naval stores, production of, in Mississippi	517, 531, 532, 536
<i>Morus Canadensis</i> (Lamarck and Rafinesque)	127	Naval stores, production of, in North Carolina	516-518
<i>Morus microphylla</i>	12, 128, 249, 306	Naval stores, production of, in South Carolina	517
<i>Morus Missouriensis</i>	127	Nebraska, lumber industry and manufactures from wood in	486, 487, 562
<i>Morus parvifolia</i> (<i>Morus microphylla</i>)	128	Nebraska, rank of, according to value of lumber products	487
<i>Morus parvifolia</i> (<i>Morus rubra</i>)	127	Nebraska, statistics of forests, forest fires, standing timber, etc., in	562
<i>Morus reticulata</i>	127	Necklace Poplar	175, 330, 350, 396, 462
<i>Morus riparia</i>	127	Nectandra	11, 15
<i>Morus rubra</i>	127, 250, 254, 257, 280, 306, 374, 442	<i>Nectandra Bredemeieriana</i>	119
<i>Morus rubra</i> , var. <i>Canadensis</i>	127	<i>Nectandra Willdenoviana</i>	119, 249, 302
<i>Morus rubra</i> , var. <i>incisa</i>	127	<i>Negundium fraxinifolium</i>	51
<i>Morus rubra</i> , var. <i>tomentosa</i>	127	Negundo	10-18
<i>Morus scabra</i>	127	Negundo aceroides	12, 50, 250, 253, 256, 276, 380, 424
<i>Morus tomentosa</i>	127	Negundo aceroides (<i>Negundo Californicum</i>)	51
Moss-ginning industry in Louisiana	537, 538	Negundo Californicum	51, 250, 253, 256, 276, 380, 424
Mossy-cup Oak	140, 265, 314, 384, 448	Negundo Californicum (<i>Negundo aceroides</i>)	51
Mountain Ash (<i>Pyrus Americana</i>)	73, 284, 428	Negundo fraxinifolium	51
Mountain Ash (<i>Pyrus sambucifolia</i>)	74, 284, 364, 428	Negundo lobatum	51
Mountain forests in California, pasturage of	579, 580	Negundo Mexicanum	51
Mountain Laurel	120, 302, 374, 440	Negundo trifoliatum	51
Mountain Magnolia	20, 266, 354, 418	Nevada, lumber industry and manufactures from wood in	486, 487, 571
Mountain Mahogany and the Nut Pine the most important trees of the Interior Forest	9	Nevada, rank of, according to value of lumber products	487
Mountain Mahogany (<i>Cercocarpus ledifolius</i>)	71, 284, 350, 428, 569, 571	Nevada, statistics of forests, forest fires, standing timber, etc., in	491, 571
Mountain Mahogany (<i>Cercocarpus parvifolius</i>)	71, 284, 569, 571	Newcastle Thorn	76, 286, 364, 430
Mountain Manchineel	54, 278, 360, 424	New Hampshire, lumber industry and manufactures from wood in	486, 487,
Mountain Maple	46, 274		497, 498
Mountain Plum	84, 276	New Hampshire, rank of, according to value of lumber products	487
Mountain White Oak	143, 316, 386, 450	New Hampshire, remarks by Mr. C. G. Pringle on the forests of	497
Mulberry, Mexican	128, 306	New Hampshire, statistics of forests, forest fires, standing timber, etc., in	491, 496-498
Mulberry, Red	128, 306, 376, 442	New Hampshire, tabular statement of the amount of Spruce standing in the forests of, May 31, 1880	496
Myginda	10, 13	New Jersey, lumber industry and manufactures from wood in	486, 487, 506
Myginda pallens	38, 249, 272	New Jersey, rank of, according to value of lumber products	487
<i>Mylocaryum ligustrinum</i>	88	New Jersey, statistics of forests, forest fires, standing timber, etc., in	491, 506
Myrica	11, 12, 15	New Mexico, lumber industry and manufactures from wood in	486, 487, 568
Myrica Californica	137, 249, 254, 257, 312, 380, 446	New Mexico, rank of, according to value of lumber products	487
Myrica Carolinensis	136	New Mexico, statistics of forests, forest fires, standing timber, etc., in	491, 568
Myrica cerifera	136, 250, 254, 257, 312, 380, 446	New Mexico, Utah, and Colorado, Chicago the principal source of supply of lumber for	568, 569
Myrica cerifera humilis	136	New York, lumber industry and manufactures from wood in	486, 487, 502-505
Myrica cerifera sempervirens	136	New York, maple-sugar product of	501
Myrica cerifera, var. <i>angustifolia</i>	136	New York, rank of, according to value of lumber products	487
		New York, rank of Albany, as a lumber market	503

	Page.		Page.
New York, remarks by Mr. C. G. Pringle on the forests of	501-506	Oak, Black (Quercus Emoryi)	146, 265, 318, 388, 452
New York, statistics of forests, forest fires, standing timber, etc., in 1891, 501-506		Oak, Black (Quercus Kelloggii)	149, 265, 320, 388, 416, 454
New York, the Northern Pine Belt in	501	Oak, Black (Quercus rubra)	148, 265, 318, 350, 388, 452
North American continent, division of the, with reference to its forest geography	3	Oak, Black (Quercus tinctoria)	149, 265, 318, 350, 388, 454, 528
North American forests, general remarks on the	3-16	Oak, Blue	143, 316, 388, 450
North Atlantic division, extent of forests, standing timber, forest fires, etc., in the	494-510	Oak, Burr	140, 265, 314, 384, 448
North Carolina, grades of tar and rosin produced in	517	Oak, Chestnut (Quercus densiflora)	155, 265, 322, 390, 456, 576, 578, 580
North Carolina, lumber industry and manufactures from wood in	486, 487, 515	Oak, Chestnut (Quercus prinoides)	143, 265, 316, 384, 450
North Carolina, production of naval stores in	516-518	Oak, Chestnut (Quercus Prinus)	142, 265, 316, 350, 384, 416, 450
North Carolina, rank of, according to value of lumber products	487	Oak, Chinquapin	143, 265, 316, 384, 450
North Carolina, statistics of forests, forest fires, standing timber, etc., in	496, 515-518	Oak, Coast Live	147, 318, 388, 432
North Carolina, tabular statement of the amount of Long-leaved Pine standing in the forests of, May 31, 1880	516	Oak, Cow	141, 316, 384, 416, 450, 533
North Carolina, the Maritime Pine Belt in	515	Oak, Duck	152, 320, 350, 388, 416, 454
North Central division, extent of forests, forest fires, standing timber, etc., in the	547-563	Oak, Iron	139, 312, 382, 448
Northern Forest in the Atlantic region	3, 4	Oak, Jack	150, 265, 320, 350, 388, 454
Northern Forest in the Pacific region, extent of the	7	Oak, Laurel (Quercus imbricaria)	154, 322, 390, 456
Northern Pine Belt, forests of the, once extended over the State of Maine	494	Oak, Laurel (Quercus laurifolia)	153, 320, 390, 454
Northern Pine Belt in Connecticut	500	Oak, Live (Quercus chrysolepis)	146, 318, 388, 452
Northern Pine Belt in Massachusetts	500	Oak, Live (Quercus virens)	145, 265, 316, 388, 453
Northern Pine Belt in Minnesota	558	Oak, Live (Quercus Wislizeni)	147, 318, 388, 452
Northern Pine Belt in New York	501	Oak, Maul	146, 318, 388, 452
Northern Pine Belt in Rhode Island	500	Oak, Mossy-cup	140, 265, 314, 384, 448
Northern Pine Belt in the Atlantic region	4	Oak, Mountain White	143, 316, 388, 450
Norway Pine	192, 336, 350, 402, 468	Oak, Over-cup (Quercus lyrata)	140, 314, 350, 384, 450, 533
Nut, Bitter	135, 310, 350, 380, 446	Oak, Over-cup (Quercus macrocarpa)	140, 265, 314, 384, 448
Nut, Bull	134, 310, 350, 380, 444	Oak, Peach (Quercus densiflora)	155, 265, 322, 390, 456
Nut, Coffee	58, 280, 360, 426	Oak, Peach (Quercus Phellos)	154, 322, 390, 456
Nut, Illinois	132, 308, 350, 378, 444	Oak, Pin	152, 320, 388, 454
Nut, King	134, 310, 350, 380, 444	Oak, Possum	152, 320, 350, 388, 416, 454
Nut, Mocker	134, 310, 350, 380, 444	Oak, Post	139, 312, 382, 448, 528
Nut, Pig	134, 310, 350, 380, 416	Oak, Punk	152, 320, 350, 388, 416, 454
Nut Pine and the Mountain Mahogany the most important trees of the Interior Forest	9	Oak, Quercitron	149, 265, 318, 350, 388, 454
Nut Pine (Pinus cembroides)	190, 336	Oak, Red (Quercus falcata)	151, 265, 320, 350, 388, 454
Nut Pine (Pinus edulis)	190, 336, 402, 468	Oak, Red (Quercus rubra)	148, 265, 318, 350, 388, 452
Nut Pine (Pinus monophylla)	190, 336, 350, 402, 468	Oak, Red (Quercus rubra, var. Texana)	148, 318, 388, 452
Nut Pine (Pinus Parryana)	189, 336, 402, 468	Oak, Rock Chestnut	142, 265, 316, 350, 384, 416, 450
Nut, Tallow	34, 270	Oak, Scarlet	148, 318, 388, 452
Nutmeg, California	186, 334, 400, 466	Oak, Scrub (Quercus Catesbaei)	151, 320, 388, 454
Nutmeg, Hickory	135, 310, 350, 380, 446	Oak, Scrub (Quercus undulata, var. Gambellii)	139, 314, 384, 448
NYCTAGINACEÆ	117, 254, 257, 302, 372, 438	Oak, Shingle	154, 322, 390, 456
Nyssa	10, 11, 14	Oak, Spanish	151, 265, 320, 350, 388, 454, 528
Nyssa angulicarpa	93	Oak, Swamp Post	140, 314, 350, 384, 450
Nyssa angulosa	93	Oak, Swamp Spanish	152, 320, 388, 454
Nyssa aquatica (Nyssa sylvatica)	92	Oak, Swamp White	141, 314, 384, 450
Nyssa aquatica (Nyssa uniflora)	92	Oak, Tanbark	155, 265, 322, 390, 456
Nyssa biflora	92	Oak, Turkey	151, 320, 388, 454
Nyssa Canadensis	92	Oak, Upland Willow	153, 320, 390, 456
Nyssa candicans	91	Oak, Valparaiso	146, 318, 388, 452
Nyssa capitata	91, 250, 253, 257, 290, 366, 432	Oak, Water (Quercus aquatica)	152, 320, 350, 388, 416, 454
Nyssa capitata, var. grandidentata	93	Oak, Water (Quercus palustris)	152, 320, 388, 454
Nyssa Caroliniana	92	Oak, Water White	140, 314, 350, 384, 450
Nyssa coccinea	91	Oak, Weeping	138, 312, 382, 448
Nyssa denticulata	92	Oak, White (Quercus alba)	137, 265, 312, 350, 380, 414, 446
Nyssa grandidentata	93	Oak, White (Quercus Garryana)	138, 312, 350, 382, 448, 576
Nyssa integrifolia	92	Oak, White (Quercus grisea)	144, 316, 386, 452
Nyssa montana	91	Oak, White (Quercus lobata)	138, 312, 382, 448
Nyssa multiflora	92	Oak, White (Quercus oblongifolia)	144, 316, 386, 452
Nyssa multiflora, var. sylvatica	92	Oak, Willow	154, 322, 390, 456
Nyssa Ogeche	91	Oak, Yellow (Quercus prinoides)	143, 265, 316, 384, 450
Nyssa palustris	93	Oak, Yellow (Quercus tinctoria)	149, 265, 318, 350, 388, 454
Nyssa sylvatica	92, 250, 253, 257, 280, 290, 366, 432	Oak, Yellow-bark	149, 265, 318, 350, 388, 454
Nyssa tomentosa (Nyssa capitata)	91	Obispo Pine	200, 340, 404, 472
Nyssa tomentosa (Nyssa uniflora)	93	ENOCARPUS regia	218
Nyssa uniflora	92, 250, 253, 257, 280, 290, 350, 366, 432	Ogechee Lime	91, 290, 366, 432
Nyssa villosa	92	Ohio Buckeye	42, 274, 358, 422
		Ohio, lumber industry and manufactures from wood in	486, 487, 547
Oak, Bartram's	153, 320, 390, 456	Ohio, rank of, according to value of lumber products	487, 587
Oak, Basket	141, 316, 384, 416, 450	Ohio, statistics of forests, forest fires, standing timber, etc., in	491, 547
		OLACINÆ	34, 270
		Old-field Birch	159, 324, 350, 382, 458
		Old-field Pine	197, 340, 350, 404, 416, 470
		Old Man's Beard	113, 298
		Olea Americana	113
		OLEACEÆ	106-113, 254, 257, 296, 370, 434
		Olive, California	120, 302, 374, 440

	Page.		Page.
Olneya	10, 12, 14	Pecan	122, 308, 350, 378, 444
Olneya Tesota	56, 249, 253, 256, 272, 360, 426	Pecan, Bitter	122, 310, 350, 380, 414, 446
Orange, Mock	70, 284, 362, 428	Pecan-nut, the, an important product west of the Colorado river in Texas	543
Orange, Osage	123, 306, 376, 442	Pencil Cedar of Florida	522
Orange, Wild (<i>Prunus Caroliniana</i>)	70, 284, 362, 428	Pennsylvania, lumber industry and manufactures from wood in	488, 487, 506-510
Orange, Wild (<i>Xanthoxylum Clava-Herculis</i>)	30, 270, 356, 429	Pennsylvania, rank of, according to value of lumber products	487
<i>Orchidocarpum arietinum</i>	23	Pennsylvania, remarks by Mr. C. G. Pringle on the forests of	507-510
Oregon Ash	111, 298, 372, 438	Pennsylvania, statistics of forests, forest fires, standing timber, etc., in	491, 506-510
Oregon Cedar	179, 332, 350, 398, 404	Pennsylvania, tabular statement of the amount of White Pine and Hemlock standing in the forests of, May 31, 1880	506
Oregon Crab Apple	73, 284	Pepperidge	32, 290, 368, 432
Oregon, lumber industry and manufactures from wood in	488, 487, 577	Pepperwood	30, 270, 356, 429
Oregon Pine	209, 265, 344, 352, 410, 476	Per cent. of tannin in the bark of certain species	265
Oregon, rank of, according to value of lumber products	487	Persea	11, 15
Oregon, remarks by Mr. Sereno Watson on the forests of certain counties of	577, 578	<i>Persea Borbonia</i>	118
Oregon, statistics of forests, forest fires, standing timber, etc., in	491, 576-578	<i>Persea Carolinensis</i>	118, 250, 254, 257, 302, 372, 438
<i>Oreodaphne Californica</i>	120	<i>Persea Carolinensis</i> , var. <i>glabriuscula</i>	118
<i>Oreodora</i>	11, 16	<i>Persea Carolinensis</i> , var. <i>palustris</i>	119, 250, 254, 257, 302, 372, 438
<i>Oreodora oleracea</i>	218	<i>Persea Carolinensis</i> , var. <i>pubescens</i>	119
<i>Oreodora regia</i>	218, 250, 348, 352	<i>Persea Oatesbyana</i>	119
Original forest of the District of Columbia replaced by Oak, Scrub Pine, etc	511	<i>Persea Sassafras</i>	119
Osage Orange	123, 306, 376, 442	Persimmon	104, 294, 350, 368, 434
Osmanthus	11, 15	Persimmon, Black	105, 294
Osmanthus Americanus	113, 249, 254, 257, 300, 372, 438	Persimmon, Mexican	105, 294
Ostrya	11, 16	<i>Phanopyrum acerifolium</i>	80
<i>Ostrya Virginiana</i>	158	<i>Phanopyrum arborescens</i>	75
<i>Ostrya Virginica</i>	158, 249, 255, 258, 322, 390, 456	<i>Phanopyrum Carolinianum</i>	82
<i>Ostrya Virginica</i> , var. <i>eglandulosa</i>	158	<i>Phanopyrum coccineum</i>	77
<i>Ostrya Virginica</i> , var. <i>glandulosa</i>	158	<i>Phanopyrum cordatum</i>	80
Over-cup Oak (<i>Quercus lyrata</i>)	140, 314, 350, 384, 450, 533	<i>Phanopyrum ellipticum</i>	82
Over-cup Oak (<i>Quercus macrocarpa</i>)	140, 265, 314, 384, 448	<i>Phanopyrum populifolium</i>	78
Oxydendrum	11, 15	<i>Phanopyrum pruinatum</i>	79
Oxydendrum arboreum	98, 249, 254, 257, 292, 368, 434	<i>Phanopyrum spathulatum</i>	81
P.		<i>Phanopyrum subvillosum</i>	78
Pacific Coast Forest, the	7	<i>Phanopyrum Virginicum</i>	82
Pacific region, the	6-10	<i>Phanopyrum viride</i>	78
<i>Padus cartilaginea</i>	68	<i>Phanopyrum Wendlandii</i>	77
<i>Padus demissa</i>	69	<i>Photinia arbutifolia</i>	83
<i>Padus serotina</i>	68	<i>Photinia salicifolia</i>	83
<i>Padus Virginiana</i>	68	Picea	11, 12, 16, 575
Palm, Fan-leaf	217, 348, 414, 480	<i>Picea alba</i>	12, 204, 250, 255, 258, 263, 342, 408, 474, 562, 564
Palm, Royal	218, 368, 352	<i>Picea amabilis</i> (<i>Abies amabilis</i>)	213
<i>Palma argentea</i>	218	<i>Picea amabilis</i> (<i>Abies subalpina</i>)	211
PALMACEAE	217, 218, 255, 259, 348, 414, 480	<i>Picea balsamea</i>	211
Palmetto, Cabbage	217, 348, 352	<i>Picea balsamea</i> , var. <i>longifolia</i>	211
Palmetto Sabal	217, 250, 348, 352	<i>Picea bifolia</i>	211
Palmetto, Silk-top	217, 348	<i>Picea bracteata</i>	213
Palmetto, Silver-top	218, 348	<i>Picea Californica</i>	208
Palo Blanco	126, 306	<i>Picea Canadensis</i>	206
Palo Verde	60, 280, 362, 426	<i>Picea caerulea</i>	204
Papaw	23, 266, 354, 418	<i>Picea concolor</i>	212
Paper Birch	160, 324, 353, 392, 458	<i>Picea concolor</i> , var. <i>violacea</i>	212
Paradise Tree	32, 270, 356, 420	<i>Picea Douglasii</i>	209
Parkinsonia	10-12, 14	<i>Picea Engelmanni</i>	9, 205, 251, 255, 258, 263, 342, 408, 474, 564-567, 572-575
<i>Parkinsonia aculeata</i>	12, 60, 250, 280	<i>Picea Fraseri</i> (<i>Abies balsamea</i>)	211
<i>Parkinsonia microphylla</i>	60, 249, 280	<i>Picea Fraseri</i> (<i>Abies Fraseri</i>)	210
<i>Parkinsonia Torreyana</i>	60, 250, 253, 256, 280, 362, 426	<i>Picea glauca</i>	204
Paralely Haw	81, 286	<i>Picea grandis</i> (<i>Abies concolor</i>)	212
<i>Pasania densiflora</i>	155	<i>Picea grandis</i> (<i>Abies grandis</i>)	212
<i>Pavia Californica</i>	43	<i>Picea laza</i>	204
<i>Pavia carnea</i>	42	<i>Picea Louisiana</i>	212
<i>Pavia discolor</i>	43	<i>Picea magnifica</i>	214
<i>Pavia flava</i>	43	<i>Picea Menziesii</i> (<i>Picea pungens</i>)	205
<i>Pavia glabra</i>	42	<i>Picea Menziesii</i> (<i>Picea Sitichensis</i>)	206
<i>Pavia hybrida</i>	43	<i>Picea nigra</i>	202, 250, 255, 258, 263, 265, 342, 352, 408, 474, 484, 496, 498
<i>Pavia lutea</i>	43	<i>Picea nigra</i> , var. <i>glauca</i>	204
<i>Pavia neglecta</i>	43	<i>Picea nigra</i> , var. <i>rubra</i>	203
<i>Pavia pallida</i>	42	<i>Picea nobilis</i>	214
<i>Pavia Watsoniana</i>	42	<i>Picea pungens</i>	13, 205, 251, 255, 258, 344, 408, 474
Peach Oak (<i>Quercus densiflora</i>)	155, 265, 322, 390, 456	<i>Picea rubra</i>	303
Peach Oak (<i>Quercus Phellos</i>)	154, 322, 390, 456	<i>Picea Sitichensis</i>	206, 250, 255, 258, 263, 344, 408, 474, 576
Peach, Wild	70, 284, 362, 428	<i>Pickeringia paniculata</i>	100
Pear Haw	79, 286, 364, 430	Pigeon Cherry	66, 262, 428

	Page.		Page.
Pigeon Plum	117, 302, 372, 438	Pine, Scrub (<i>Pinus contorta</i>)	194, 338, 404, 470
Pigeonwood	117, 302, 372, 438	Pine, Scrub (<i>Pinus inopa</i>)	199, 340, 350, 404, 470, 529
Pig Nut	184, 310, 350, 380, 448	Pine, Short-leaved	200, 340, 350, 404, 472, 524, 527, 530, 531, 536, 537, 541, 544, 560
Pike county, Alabama, the forests of	528	Pine, Short-leaved, standing in the forests of Arkansas, May 31, 1880, tabular statement of the amount of	544
<i>Pilocereus Engelmanni</i>	80	Pine, Slash	202, 342, 352, 406, 416, 474, 516
Pin Cherry	66, 282, 428	Pine, Southern	202, 342, 352, 406, 416, 472
Pin Oak	152, 320, 388, 464	Pine, Spruce (<i>Pinus clausa</i>)	199, 340, 404, 472
Pinckneya	10, 11, 14	Pine, Spruce (<i>Pinus glabra</i>)	201, 340, 406, 472, 527
<i>Pinckneya pubens</i>	95, 250, 253, 257, 290, 306, 432	Pine, Spruce (<i>Pinus mitis</i>)	200, 340, 350, 406, 472
<i>Pinckneya pubescens</i>	95	Pine, Spruce (<i>Pinus Murrayana</i>)	195, 338, 350, 404, 470
Pine and Spruce standing in the forests of Maine May 31, 1880, tabular statement of the amount of	494	Pine, Sugar	188, 338, 350, 400, 466, 576, 578
Pine, Bastard	202, 342, 352, 406, 416, 474	Pine, Sugar, an important tree in the Coast Forest	8
Pine belt of central Alabama	529	Pine, Swamp	202, 342, 352, 406, 416, 474
Pine, Bishop's	200, 340, 404, 472	Pine, Table-mountain	199, 340, 350, 404, 472
Pine, Black (<i>Pinus Jeffreyi</i>)	193, 338, 402, 470	Pine, Weymouth	187, 334, 350, 400, 466
Pine, Black (<i>Pinus Murrayana</i>)	195, 338, 350, 404, 470	Pine, White, and Hemlock, standing in the forests of Pennsylvania, May 31, 1880, tabular statement of the amount of	506
Pine, Bull (<i>Pinus Jeffreyi</i>)	193, 338, 402, 470	Pine, White (<i>Pinus flexilis</i>)	188, 336, 400, 468, 569, 570, 571
Pine, Bull (<i>Pinus mitis</i>)	200, 340, 350, 406, 472	Pine, White (<i>Pinus glabra</i>)	201, 340, 406, 472
Pine, Bull (<i>Pinus ponderosa</i>)	193, 338, 350, 402, 468	Pine, White (<i>Pinus monticola</i>)	187, 336, 400, 466, 564, 574, 576
Pine, Bull (<i>Pinus Sabiniana</i>)	195, 338, 350, 404, 470	Pine, White (<i>Pinus reflexa</i>)	189, 336, 402, 468
Pine, Cedar	201, 340, 406, 472	Pine, White (<i>Pinus Strobus</i>)	187, 334, 350, 400, 466, 566, 551, 554, 558
Pine, Digger	195, 338, 350, 404, 470	Pine, White, standing in the forests of Michigan May 31, 1880, tabular statement of the amount of	551
Pine forests of Baldwin county, Alabama	527	Pine, White, standing in the forests of Minnesota May 31, 1880, tabular statement of the amount of	558
Pine forests of southern Mississippi	531, 532	Pine, White, standing in the forests of Wisconsin May 31, 1880, tabular statement of the amount of	554
Pine forests of the northeastern counties of Mississippi	532-534	Pine, White, the characteristic tree of the Northern Pine Belt	4
Pine, Foxtail	191, 336, 402, 468	Pine, Yellow, an important, characteristic tree of the Coast Forest	8
Pine, Georgia	202, 342, 352, 406, 416, 472	Pine, Yellow (<i>Pinus Arizonica</i>)	192, 338, 402, 468
Pine, Ginger	179, 332, 350, 398, 464	Pine, Yellow (<i>Pinus mitis</i>)	200, 340, 350, 406, 472, 529
Pine, Gray	201, 342, 352, 406, 472	Pine, Yellow (<i>Pinus palustris</i>)	202, 342, 352, 406, 416, 472
Pine, Hard	202, 342, 352, 406, 416, 472	Pine, Yellow (<i>Pinus ponderosa</i>)	193, 338, 350, 402, 468, 562, 565, 574, 578
Pine, Hickory (<i>Pinus Balfouriana</i> , var. <i>aristata</i>)	191, 336, 402, 468	Piñon (<i>Pinus edulis</i>)	190, 336, 402, 468
Pine, Hickory (<i>Pinus pungens</i>)	199, 340, 350, 404, 472	Piñon (<i>Pinus monophylla</i>)	190, 336, 350, 402, 468
Pine, Jersey	199, 340, 350, 404, 470, 546	Piñon (<i>Pinus Parryana</i>)	189, 336, 402, 468
Pine, Knob-cone	196, 340, 404, 470	Pinus	11, 12, 16
Pine, Loblolly	197, 340, 350, 404, 416, 470, 516, 541	<i>Pinus Abies Americana</i>	206
Pine, Lodge-pole	195, 338, 350, 404, 470, 564, 574, 577	<i>Pinus Abies Balsamea</i>	211
Pine, Long-leaved	202, 342, 352, 406, 416, 472, 516, 519-521, 524, 531, 537, 541	<i>Pinus Abies Canadensis</i>	202
Pine, Long- and Short-leaved, standing in the forests of Alabama May 31, 1880, tabular statement of the amount of	524	<i>Pinus adunca</i>	196
Pine, Long- and Short-leaved, standing in the forests of Louisiana May 31, 1880, tabular statement of the amount of	537	<i>Pinus alba</i>	204
Pine, Long-leaved, Loblolly, and Short-leaved, standing in the forests of Texas May 31, 1880, tabular statement of the amount of	541	<i>Pinus albicaulis</i>	189, 250, 255, 258, 336, 400, 468, 565, 572, 578
Pine, Long-leaved, standing in the forests of Florida May 31, 1880, tabular statement of the amount of	521	<i>Pinus amabilis</i> (<i>Abies amabilis</i>)	213
Pine, Long-leaved, standing in the forests of Georgia May 31, 1880, tabular statement of the amount of	520	<i>Pinus amabilis</i> (<i>Abies magnifica</i>)	214
Pine, Long-leaved, standing in the forests of North Carolina May 31, 1880, tabular statement of the amount of	516	<i>Pinus amabilis</i> (<i>Abies subalpina</i>)	211
Pine, Long-leaved, standing in the forests of South Carolina May 31, 1880, tabular statement of the amount of	519	<i>Pinus Americana</i> (<i>Picea nigra</i>)	203
Pine, Long-leaved, the characteristic tree of the Southern Maritime Pine Belt	4	<i>Pinus Americana</i> (<i>Tsuga Canadensis</i>)	206
Pine, Meadow	202, 342, 352, 406, 416, 474	<i>Pinus Americana rubra</i>	202
Pine, Monterey	196, 340, 404, 470	<i>Pinus aristata</i>	191
Pine, Norway	192, 336, 350, 402, 468	<i>Pinus arizonica</i>	192, 250, 255, 258, 338, 402, 468
Pine, Nut (<i>Pinus cembroides</i>)	190, 336	<i>Pinus australis</i>	201
Pine, Nut (<i>Pinus edulis</i>)	190, 336, 402, 468	<i>Pinus Balfouriana</i>	191, 250, 255, 258, 336, 402, 468
Pine, Nut (<i>Pinus monophylla</i>)	190, 336, 350, 402, 468	<i>Pinus Balfouriana</i> (<i>Pinus Balfouriana</i> , var. <i>aristata</i>)	191
Pine, Nut (<i>Pinus Parryana</i>)	189, 336, 402, 468	<i>Pinus Balfouriana</i> , var. <i>aristata</i>	191, 250, 255, 258, 336, 402, 468
Pine, Nut, and the Mountain Mahogany the most important trees of the Interior Forest	9	<i>Pinus balsamea</i>	210
Pine, Obispo	200, 340, 404, 472	<i>Pinus balsamea</i> , var. <i>Fraseri</i>	210
Pine, Old-field	197, 340, 350, 404, 416, 470	<i>Pinus Banksiana</i>	201, 250, 255, 258, 263, 342, 352, 406, 472
Pine, Oregon	209, 285, 344, 352, 410, 476	<i>Pinus Banksiana</i> (<i>Pinus contorta</i>)	194
Pine, Pitch	198, 340, 350, 404, 470, 520, 522, 525, 531	<i>Pinus Beardsleyi</i>	193
Pine, Pond	198, 340, 350, 404, 416, 470, 527	<i>Pinus Benthamiana</i>	193
Pine, Prince's	201, 342, 352, 406, 472	<i>Pinus Bolanderi</i>	194
Pine, Red	192, 336, 350, 402, 468	<i>Pinus Boursieri</i>	194
Pine region of the Coosa in Alabama	529	<i>Pinus brachyptera</i>	193
Pine, Rosemary	197, 340, 350, 404, 416, 470	<i>Pinus bracteata</i>	213
Pine, Sand	199, 340, 404, 472	<i>Pinus Californica</i> (<i>Pinus insignis</i>)	196
Pine, Scrub (<i>Pinus Banksiana</i>)	201, 342, 352, 406, 472	<i>Pinus Californica</i> (<i>Pinus tuberculata</i>)	196
Pine, Scrub (<i>Pinus clausa</i>)	199, 340, 404, 472	<i>Pinus Canadensis</i> (<i>Picea alba</i>)	204
		<i>Pinus Canadensis</i> (<i>Tsuga Canadensis</i>)	206
		<i>Pinus Canadensis</i> (<i>Tsuga Mertensiana</i>)	207
		<i>Pinus cembroides</i>	190, 250, 336
		<i>Pinus cembroides</i> (<i>Pinus albicaulis</i>)	189
		<i>Pinus cembroides</i> (<i>Pinus edulis</i>)	190

	Page.		Page.
<i>Pinus Chihuahuana</i>	194, 250, 255, 258, 338, 404, 470	<i>Pinus monophylla</i>	9, 18, 190, 250, 255, 258, 338, 350, 402, 468
<i>Pinus clausa</i>	190, 250, 255, 258, 340, 404, 472	<i>Pinus monticola</i>	7, 187, 250, 255, 258, 338, 400, 466, 564, 565, 573-576
<i>Pinus commutata</i>	205	<i>Pinus muricata</i>	190, 250, 255, 258, 340, 404, 472
<i>Pinus concolor</i>	212	<i>Pinus muricata</i> (Pinus contorta)	194
<i>Pinus contorta</i>	194, 250, 255, 258, 338, 404, 470	<i>Pinus Murrayana</i>	9, 194, 250, 255, 258, 338, 350, 404, 470, 564-567, 573-577
<i>Pinus contorta</i> (Pinus muricata)	199	<i>Pinus nigra</i>	202
<i>Pinus contorta</i> (Pinus Murrayana)	194	<i>Pinus nobilis</i>	214
<i>Pinus contorta</i> , var. <i>Bolanderi</i>	194	<i>Pinus Nuttallii</i>	216
<i>Pinus contorta</i> , var. <i>latifolia</i>	194	<i>Pinus ootocarpa</i>	190
<i>Pinus Coulteri</i>	195, 250, 255, 258, 340, 404, 470	<i>Pinus palustris</i>	4, 201, 249, 255, 258, 263, 342, 352, 406, 416, 472, 516, 519-521, 524, 531, 537, 541
<i>Pinus Oraigana</i>	193	<i>Pinus palustris</i> the characteristic tree of the Southern Maritime Pine Belt	4
<i>Pinus Cubensis</i>	202, 249, 255, 258, 263, 342, 352, 406, 416, 474, 516, 520, 522, 525	<i>Pinus Parryana</i>	189, 250, 255, 258, 338, 402, 468
<i>Pinus Oubensis</i> , var. <i>terthrocarpa</i>	202	<i>Pinus Parryana</i> (Pinus ponderosa)	193
<i>Pinus deflexa</i>	193	<i>Pinus Pattoniana</i> (Tsuga Mertensiana)	208
<i>Pinus Douglasii</i>	209	<i>Pinus Pattoniana</i> (Tsuga Pattoniana)	208
<i>Pinus Douglasii</i> , var. <i>brevibracteata</i>	209	<i>Pinus pendula</i>	215
<i>Pinus echinata</i>	200	<i>Pinus ponderosa</i>	8, 9, 192, 250, 255, 258, 263, 338, 350, 402, 468, 562, 565, 567, 574, 575, 578
<i>Pinus Edgariana</i>	199	<i>Pinus ponderosa</i> an important characteristic tree of the Coast Forest	8
<i>Pinus edulis</i>	9, 18, 190, 250, 255, 258, 338, 402, 468	<i>Pinus ponderosa</i> , var. <i>Benthianiana</i>	193
<i>Pinus Elliottii</i>	202	<i>Pinus ponderosa</i> , var. <i>Jeffreyi</i>	193
<i>Pinus Engelmannii</i> (Picea Engelmannii)	205	<i>Pinus ponderosa</i> , var. <i>scopulorum</i>	193
<i>Pinus Engelmannii</i> (Pinus ponderosa)	193	<i>Pinus porphyrocarpa</i>	187
<i>Pinus flexilis</i>	18, 188, 250, 255, 258, 338, 400, 468, 560-571	<i>Pinus pungens</i>	190, 250, 255, 258, 340, 350, 404, 472
<i>Pinus flexilis</i> (Pinus albicaulis)	189	<i>Pinus radiata</i>	196
<i>Pinus flexilis</i> , var. <i>albicaulis</i>	189	<i>Pinus reflexa</i>	189, 250, 255, 258, 338, 402, 468, 568, 569
<i>Pinus flexilis</i> , var. <i>macrocarpa</i>	188	<i>Pinus resinosa</i>	191, 250, 255, 258, 263, 338, 350, 402, 468
<i>Pinus flexilis</i> , var. <i>reflexa</i>	189	<i>Pinus resinosa</i> (Pinus ponderosa)	193
<i>Pinus flexilis</i> , var. <i>serrulata</i>	188	<i>Pinus rigida</i>	197, 250, 255, 258, 340, 350, 404, 470
<i>Pinus Fraseri</i> (Abies Fraseri)	210	<i>Pinus rigida</i> (Pinus insignis)	196
<i>Pinus Fraseri</i> (Pinus rigida)	197	<i>Pinus rigida</i> (Pinus mitis)	200
<i>Pinus Fremontiana</i>	190	<i>Pinus rigida</i> , var. <i>serotina</i>	198
<i>Pinus futilis</i>	190	<i>Pinus rubra</i> (Picea nigra)	203
<i>Pinus glabra</i>	200, 250, 255, 258, 263, 340, 406, 472, 527	<i>Pinus rubra</i> (Pinus resinosa)	191
<i>Pinus grandis</i> (Abies amabilis)	213	<i>Pinus rubra</i> , var. <i>violacea</i>	204
<i>Pinus grandis</i> (Abies concolor)	212	<i>Pinus rupestris</i>	201
<i>Pinus grandis</i> (Abies grandis)	212	<i>Pinus Sabiniana</i>	195, 250, 255, 258, 338, 404, 470
<i>Pinus Grozelleri</i>	187	<i>Pinus Sabiniana Coulteri</i>	195
<i>Pinus Hudsonica</i>	201	<i>Pinus Sabiniana macrocarpa</i>	195
<i>Pinus inops</i>	196, 250, 255, 258, 263, 340, 350, 404, 470, 529, 546	<i>Pinus serotina</i>	198, 349, 255, 258, 340, 350, 404, 416, 470, 527
<i>Pinus inops</i> (Pinus contorta)	194	<i>Pinus Shasta</i>	189
<i>Pinus inops</i> (Pinus Murrayana)	194	<i>Pinus Sinclairii</i>	196
<i>Pinus inops</i> , var. (Pinus muricata)	199	<i>Pinus Sitchensis</i>	206
<i>Pinus inops</i> , var. <i>clausa</i>	199	<i>Pinus species</i> (Abies subalpina)	211
<i>Pinus insignis</i>	8, 196, 250, 255, 258, 340, 404, 470	<i>Pinus Strobus</i>	4, 187, 251, 255, 258, 263, 334, 350, 400, 466, 494, 506, 551, 554, 558
<i>Pinus insignis macrocarpa</i>	196	<i>Pinus Strobus</i> the characteristic tree of the Northern Pine Belt	4
<i>Pinus insignis</i> , var. <i>binata</i>	196	<i>Pinus Strobus</i> , var. <i>alba</i>	187
<i>Pinus intermedia</i>	215	<i>Pinus Strobus</i> , var. <i>brevifolia</i>	187
<i>Pinus Jeffreyi</i>	9, 193, 250, 255, 258, 338, 402, 470, 578	<i>Pinus Strobus</i> , var. <i>compressa</i>	187
<i>Pinus Lambertiana</i>	8, 188, 251, 255, 258, 338, 350, 400, 466, 576, 578	<i>Pinus Strobus</i> , var. <i>menticola</i>	187
<i>Pinus Lambertiana</i> , var. (Pinus flexilis)	188	<i>Pinus Strobus</i> , var. <i>nivea</i>	187
<i>Pinus Lambertiana</i> , var. <i>brevifolia</i>	188	<i>Pinus sylvestris</i> , var. <i>divaricata</i>	201
<i>Pinus laricina</i>	215	<i>Pinus Teda</i>	197, 250, 255, 258, 263, 340, 350, 404, 416, 470, 512, 516, 541
<i>Pinus Laricio</i> , var. <i>resinosa</i>	191	<i>Pinus Teda</i> , var. <i>a</i>	197
<i>Pinus Larix</i>	216	<i>Pinus Teda</i> , var. <i>alopeuroidea</i>	198
<i>Pinus Larix alba</i>	215	<i>Pinus Teda</i> , var. <i>heterophylla</i>	202
<i>Pinus Larix nigra</i>	215	<i>Pinus Teda</i> , var. <i>rigida</i>	197
<i>Pinus Larix rubra</i>	215	<i>Pinus Teda</i> , var. <i>tenuifolia</i>	197
<i>Pinus lasiocarpa</i> (Abies concolor)	212	<i>Pinus Teda</i> , var. <i>variabilis</i>	200
<i>Pinus lasiocarpa</i> (Abies subalpina)	211	<i>Pinus Teda</i> , var. <i>Virginiana</i>	198
<i>Pinus Llaveana</i> (Pinus cembroides)	190	<i>Pinus taxifolia</i>	209
<i>Pinus Llaveana</i> (Pinus Parryana)	189	<i>Pinus tetragona</i>	204
<i>Pinus Loddigesii</i>	197	<i>Pinus Torreyana</i>	8, 192, 250, 255, 258, 338, 402, 468
<i>Pinus lophosperma</i>	192	<i>Pinus tuberculata</i>	196, 251, 255, 258, 340, 404, 470
<i>Pinus Louiana</i>	212	<i>Pinus tuberculata</i> (Pinus insignis)	196
<i>Pinus Lyallii</i>	216	<i>Pinus variabilis</i>	200
<i>Pinus macrocarpa</i>	195	<i>Pinus venusta</i>	218
<i>Pinus macrophylla</i>	193	<i>Pinus Virginiana</i>	198
<i>Pinus Mariana</i>	202	<i>Pinus Virginiana</i> , var. <i>echinata</i>	200
<i>Pinus Menziesii</i>	206	<i>Piscidia</i>	10, 14
<i>Pinus Menziesii</i> , var. <i>crilepa</i>	206	<i>Piscidia Carthagenensis</i>	57
<i>Pinus Mertensiana</i>	207	<i>Piscidia Erythrina</i>	57, 249, 253, 256, 278, 380, 426
<i>Pinus microcarpa</i>	215	<i>Pisonia</i>	11, 15
<i>Pinus mitis</i>	4, 200, 250, 255, 258, 340, 350, 406, 472, 524, 527, 529-531, 533, 536, 537, 541, 544, 560		
<i>Pinus mitis</i> , var. <i>paupera</i>	200		

	Page.		Page.
<i>Pisonia aculeata</i>	117	<i>Populus deltoides</i>	174
<i>Pisonia obtusata</i>	117, 250, 254, 257, 302, 372, 438	<i>Populus Fremontii</i>	175, 250, 255, 258, 330, 396, 462
<i>Pistacia</i>	10-12, 14	<i>Populus Fremontii</i> , var. <i>Wislizeni</i>	175, 250, 255, 258, 302, 330, 396, 462
<i>Pistacia Mexicana</i>	54	<i>Populus glandulosa</i>	175
Pitch Pine	198, 340, 350, 404, 470, 520, 522, 525, 531	<i>Populus grandidentata</i>	172, 250, 255, 258, 328, 394, 460
<i>Pithecolobium forbesi</i>	64	<i>Populus grandidentata</i> , var. <i>pendula</i>	172
<i>Pithecolobium Guadalupeensis</i>	64	<i>Populus heterophylla</i>	172, 250, 255, 258, 328, 394, 460
<i>Pithecolobium microphyllum</i>	64	<i>Populus heterophylla</i> (Populus balsamifera, var. <i>candicans</i>)	173
<i>Pithecolobium Ungula-cati</i>	64, 249, 282	<i>Populus heterophylla</i> , var. <i>argentea</i>	172
<i>Planera</i>	11, 15	<i>Populus laevigata</i> (Populus monilifera, <i>Atton</i> , etc.)	175
<i>Planera aquatica</i>	124, 250, 254, 257, 306, 376, 442	<i>Populus laevigata</i> (Populus monilifera, <i>Hort.</i>)	175
<i>Planera Gmelini</i>	124	<i>Populus latifolia</i>	173
<i>Planera Richardi</i>	124	<i>Populus Lindleyana</i>	175
<i>Planera ulmifolia</i>	124	<i>Populus macrophylla</i> (Populus balsamifera, var. <i>candicans</i>)	173
PLATANACEÆ	129, 130, 254, 257, 306, 376, 442	<i>Populus macrophylla</i> (Populus monilifera)	175
<i>Platanus</i>	11, 12, 15, 251	<i>Populus Marylandica</i>	175
<i>Platanus Californica</i>	129	<i>Populus monilifera</i>	174, 251, 255, 258, 302, 330, 350, 396, 462
<i>Platanus hybridus</i>	129	<i>Populus monilifera</i> (Populus Fremontii)	175
<i>Platanus lobata</i>	129	<i>Populus monilifera</i> (Populus Fremontii, var. <i>Wislizeni</i>)	175
<i>Platanus Mexicana</i> (<i>Platanus racemosa</i>)	129	<i>Populus neglecta</i>	175
<i>Platanus Mexicana</i> (<i>Platanus Wrightii</i>)	130	<i>Populus Ontariensis</i>	173
<i>Platanus occidentalis</i>	129, 250, 254, 257, 306, 350, 370, 442	<i>Populus Tacamahaca</i>	173
<i>Platanus occidentalis</i> (<i>Platanus racemosa</i>)	129	<i>Populus tremuliformis</i>	171
<i>Platanus racemosa</i>	129, 250, 254, 257, 306, 370, 442	<i>Populus tremuloides</i>	12, 171, 250, 255, 258, 328, 350, 394, 460
<i>Platanus racemosa</i> (<i>Platanus Wrightii</i>)	130	<i>Populus tremula</i>	171
<i>Platanus vulgaris</i> , var. <i>angulosa</i>	129	<i>Populus trichocarpa</i>	174, 251, 255, 258, 330, 394, 462, 576
<i>Platanus Wrightii</i>	130, 250, 254, 257, 306, 376, 442	<i>Populus trichocarpa</i> , var. <i>cupulata</i>	174
Plum, Canada	65, 282, 362, 426	<i>Populus viminea</i>	173
Plum, Chickasaw	66, 282, 362, 426	<i>Populus Virginiana</i>	175
Plum, Cocoa	65, 282, 362, 426	<i>Porcelia triloba</i>	23
Plum, Darling	39, 272, 358, 422	Porkwood	117, 302, 372, 438
Plum, Downward	103, 294, 368, 434	Porlira	10-13
Plum, Gopher	91, 290, 366, 432	Porlira angustifolia	29, 249, 268
Plum, Guiana	121, 302, 374, 440	Port Orford Cedar	179, 332, 350, 398, 464, 576
Plum, Hog (<i>Prunus angustifolia</i>)	66, 282, 362, 426	Possum Oak	152, 320, 350, 388, 410, 454
Plum, Hog (<i>Rhus Metopium</i>)	54, 278, 300, 424	Post Cedar	176, 330, 396, 462
Plum, Hog (<i>Ximenia Americana</i>)	34, 270	Post Oak	139, 312, 382, 448, 528
Plum, Horse	65, 282, 362, 426	Post Oak, Swamp	140, 314, 350, 384, 450
Plum, Mountain	34, 270	Prickly Ash (<i>Xanthoxylum Americanum</i>)	29, 268
Plum, Pigeon	117, 302, 372, 438	Prickly Ash (<i>Xanthoxylum Clava-Herculis</i>)	30, 270, 356, 420
Plum, Saffron	103, 294, 368, 434	Prince's Pine	201, 342, 352, 406, 472
Plum, Wild	65, 282, 362, 426	Principal woods of the United States under compression, behavior of the	418-481
Poison Elder	54, 278	Principal woods of the United States under transverse strain, behavior of the	354-414
Poison Sumach	54, 278	Pringle, Mr. C. G., remarks on the forests of New Hampshire by	497
Poisonwood (<i>Rhus Metopium</i>)	54, 278, 360, 424	Pringle, Mr. C. G., remarks on the forests of New York by	501-506
Poisonwood (<i>Sebastiania lucida</i>)	121, 304	Pringle, Mr. C. G., remarks on the forests of Pennsylvania by	507-510
POLYGONACEÆ	117, 118, 254, 257, 302, 372, 438	Pringle, Mr. C. G., remarks on the forests of Vermont by	498-500
<i>Polygonum uvifera</i>	118	Pringle, Mr. C. G., remarks on the forests of West Virginia by	512-515
Pond Apple	23, 266, 354, 418	<i>Prinos deciduus</i>	37
Pond Pine	198, 340, 350, 404, 416, 470, 527	<i>Pritchardia filamentosa</i>	217
Poplar	172, 328, 394, 460	Privet	112, 298, 372, 438
Poplar, Carolina	175, 330, 350, 396, 462	Product and average size of saw-mills in each state and territory	488
Poplar, Necklace	175, 330, 350, 396, 462	Production of naval stores in Alabama	517, 527, 520, 530
Poplar, Yellow	22, 266, 354, 418	Production of naval stores in Florida	517
<i>Populus</i>	11, 12, 16	Production of naval stores in Georgia	517
<i>Populus aladeeca</i>	173	Production of naval stores in Louisiana	517
<i>Populus angulata</i>	175	Production of naval stores in Mississippi	517
<i>Populus angulosa</i>	175	Production of naval stores in North Carolina	517
<i>Populus angustifolia</i>	13, 174, 250, 255, 258, 394, 462, 565	Production of naval stores in South Carolina	517
<i>Populus angustifolia</i> (Populus trichocarpa)	174	<i>Prosopis</i>	10-12, 14
<i>Populus argentea</i>	172	<i>Prosopis Emoryi</i>	62
<i>Populus Atheniensis</i> 1	171	<i>Prosopis glandulosa</i>	61
<i>Populus balsamifera</i>	12, 173, 251, 255, 258, 328, 394, 460, 573, 575	<i>Prosopis juliflora</i>	12, 61, 249, 253, 256, 265, 280, 350, 362, 426
<i>Populus balsamifera</i> (Populus trichocarpa)	174	<i>Prosopis odorata</i> (<i>Prosopis juliflora</i>)	61
<i>Populus balsamifera lanceolata</i>	173	<i>Prosopis odorata</i> (<i>Prosopis pubescens</i>)	62
<i>Populus balsamifera</i> , var.	174	<i>Prosopis pubescens</i>	18, 62, 249, 253, 256, 280, 362, 426
<i>Populus balsamifera</i> , var. <i>angustifolia</i>	174	<i>Prunus</i>	10-12, 14, 573
<i>Populus balsamifera</i> , var. <i>Californica</i>	174	<i>Prunus Americana</i>	12, 65, 249, 253, 256, 282, 362, 426
<i>Populus balsamifera</i> , var. <i>candicans</i>	173, 250, 255, 258, 330, 394, 460	<i>Prunus Americana</i> , var. <i>mollis</i>	65
<i>Populus balsamifera</i> , var. <i>genuina</i>	173	<i>Prunus angustifolia</i>	60, 249, 253, 256, 282, 362, 426
<i>Populus Canadensis</i> (Populus balsamifera, var. <i>candicans</i>)	173	<i>Prunus borealis</i>	66
<i>Populus Canadensis</i> (Populus monilifera)	175	<i>Prunus Canadensis</i>	68
<i>Populus Canadensis</i> , var. <i>angustifolia</i>	174	<i>Prunus Capulin</i>	68
<i>Populus candicans</i>	173	<i>Prunus Capuli</i>	68, 249, 282, 428
<i>Populus cordifolia</i>	172		

	Page.		Page.
<i>Prunus Carolina</i>	69	<i>Quassia Simaruba</i>	82
<i>Prunus Caroliniana</i>	69, 249, 253, 256, 284, 362, 428	<i>Quercitron Oak</i>	149, 265, 318, 350, 388, 454
<i>Prunus cartilaginea</i>	68	<i>Quercus</i>	11, 12, 15
<i>Prunus Chicasa</i>	66	<i>Quercus acutidens</i>	155
<i>Prunus cocinea</i>	65	<i>Quercus acutiglans</i>	146
<i>Prunus demissa</i>	69, 249, 253, 256, 284, 362, 428	<i>Quercus agrifolia</i>	146, 249, 254, 257, 318, 386, 452
<i>Prunus emarginata</i>	67, 250, 282	<i>Quercus agrifolia</i> , var. <i>frutescens</i>	147
<i>Prunus emarginata</i> , var. <i>mollis</i>	67, 253, 256, 282, 362, 428	<i>Quercus alba</i>	187, 249, 254, 257, 261, 265, 312, 350, 380, 414, 446
<i>Prunus hiemalis</i> (<i>Prunus Americana</i>).....	65	<i>Quercus alba minor</i>	139
<i>Prunus hiemalis</i> (<i>Prunus Americana</i> , var. <i>mollis</i>).....	65	<i>Quercus alba palustris</i>	141
<i>Prunus ilicifolia</i>	70, 249, 253, 256, 284, 364, 428	<i>Quercus alba</i> , var. <i>Gunnisonii</i>	139
<i>Prunus inistitia</i>	66	<i>Quercus alba</i> , var. <i>microcarpa</i>	137
<i>Prunus lanceolata</i>	66	<i>Quercus alba</i> , var. <i>pinnatifida</i>	137
<i>Prunus Lusitanica</i>	69	<i>Quercus alba</i> , var. <i>pinnatifido-sinuata</i>	137
<i>Prunus Mississippi</i>	65	<i>Quercus alba</i> , var. <i>repanda</i>	137
<i>Prunus mollis</i> (<i>Prunus Americana</i> , var. <i>mollis</i>).....	65	<i>Quercus alba</i> , var. <i>sinuata</i>	137
<i>Prunus mollis</i> (<i>Prunus emarginata</i> , var. <i>mollis</i>).....	67	<i>Quercus ambigua</i>	147
<i>Prunus nigra</i>	65	<i>Quercus annulata</i>	145
<i>Prunus Pennsylvanica</i>	12, 66, 250, 282, 428	<i>Quercus aquatica</i>	152, 249, 254, 258, 262, 320, 350, 388, 416, 454
<i>Prunus persicifolia</i>	60	<i>Quercus aquatica</i> (<i>Quercus nigra</i>).....	150
<i>Prunus pumila</i>	67	<i>Quercus aquatica</i> , var. <i>attenuata</i>	152
<i>Prunus sempervirens</i>	69	<i>Quercus aquatica</i> , var. <i>cuneata</i>	152
<i>Prunus serotina</i>	68, 250, 253, 256, 259, 282, 362, 428	<i>Quercus aquatica</i> , var. <i>elongata</i>	152
<i>Prunus serratifolia</i>	69	<i>Quercus aquatica</i> , var. <i>heterophylla</i>	15
<i>Prunus sphaerocarpa</i>	70, 249, 284	<i>Quercus aquatica</i> , var. <i>hybrida</i>	152
<i>Prunus spinosa</i>	65	<i>Quercus aquatica</i> , var. <i>indivisa</i>	152
<i>Prunus umbellata</i>	67, 249, 282, 428	<i>Quercus aquatica</i> , var. <i>laurifolia</i>	152
<i>Prunus Virginiana</i>	68	<i>Quercus aquatica</i> , var. <i>myrtifolia</i>	155
<i>Prunus Virginiana</i> (<i>Prunus serotina</i>).....	68	<i>Quercus Banisteri</i>	155
<i>Prunus Virginiana</i> , var. <i>demissa</i>	69	<i>Quercus berberidifolia</i>	155
<i>Pseudacacia odorata</i>	55	<i>Quercus bicolor</i>	141, 249, 254, 257, 261, 314, 384, 450
<i>Pseudopetalon glandulosum</i>	30	<i>Quercus bicolor</i> , var. <i>Michauxii</i>	141
<i>Pseudopetalon tricarpum</i>	30	<i>Quercus bicolor</i> , var. <i>mollis</i>	141
<i>Pseudotsuga</i>	7, 9, 11, 12, 16, 575	<i>Quercus bicolor</i> , var. <i>platanoides</i>	141
<i>Pseudotsuga Douglasii</i>	209, 250, 255, 259, 284, 265, 344, 352, 410, 476, 565, 569, 573, 576	<i>Quercus Breweri</i>	155
<i>Pseudotsuga Douglasii</i> , var. <i>macrocarpa</i>	210, 250, 255, 259, 346, 412, 478	<i>Quercus Californica</i>	149
<i>Pseudotsuga magnifica</i>	214	<i>Quercus Castanea</i> (<i>Quercus prinoides</i>).....	142
<i>Pseudotsuga nobilis</i>	214	<i>Quercus Castanea</i> (<i>Quercus Prinus</i>).....	142
<i>Pseudotsuga</i> the most important timber tree of the Pacific region.....	7	<i>Quercus Catesbaei</i>	151, 249, 254, 258, 320, 388, 454, 525
<i>Ptella</i>	10, 13	<i>Quercus Chinquapin</i>	143
<i>Ptella mollis</i>	31	<i>Quercus chrysolepis</i>	146, 249, 254, 258, 318, 386, 452
<i>Ptella trifoliata</i>	12, 31, 249, 270	<i>Quercus chrysolepis</i> , var. <i>vacciniifolia</i>	146
<i>Ptella trifoliata</i> , var. <i>mollis</i>	31	<i>Quercus cinerea</i>	153, 250, 255, 258, 320, 390, 456, 525
<i>Ptella viticifolia</i>	31	<i>Quercus cinerea</i> , var. <i>pumila</i>	155
<i>Punk Oak</i>	152, 320, 350, 388, 416, 454	<i>Quercus cinerea</i> , var. <i>sericea</i>	155
<i>Purple Haw</i>	40, 272	<i>Quercus coccinea</i>	148, 249, 254, 258, 318, 388, 452
<i>Purshia</i>	575	<i>Quercus coccinea</i> , var. <i>ambigua</i>	147
<i>Putnam</i> , Mr. H. C., remarks on the forests of Michigan by.....	553, 554	<i>Quercus coccinea</i> , var. <i>microcarpa</i>	148
<i>Putnam</i> , Mr. H. C., remarks on the forests of Minnesota by.....	559, 560	<i>Quercus coccinea</i> , var. <i>rubra</i>	147
<i>Putnam</i> , Mr. H. C., remarks on the forests of the eastern portion of Dakota by.....	561	<i>Quercus coccinea</i> , var. <i>tinctoria</i>	149
<i>Putnam</i> , Mr. H. C., remarks on the forests of Wisconsin by.....	555-558	<i>Quercus confertifolia</i>	154
<i>Pyrus</i>	10-12, 14	<i>Quercus crassipocula</i>	146
<i>Pyrus Americana</i>	73, 250, 284, 428	<i>Quercus cuneata</i>	150
<i>Pyrus Americana</i> (<i>Pyrus sambucifolia</i>).....	74	<i>Quercus decipiens</i>	144
<i>Pyrus Americana</i> , var. <i>microcarpa</i>	74	<i>Quercus densiflora</i>	154, 249, 254, 258, 265, 322, 390, 456, 570, 580
<i>Pyrus angustifolia</i>	72, 249, 284	<i>Quercus discolor</i>	149
<i>Pyrus aucuparia</i> (<i>Pyrus Americana</i>).....	73	<i>Quercus discolor</i> , var. <i>triloba</i>	151
<i>Pyrus aucuparia</i> (<i>Pyrus sambucifolia</i>).....	74	<i>Quercus Douglasii</i>	143, 249, 251, 254, 258, 316, 386, 450
<i>Pyrus Bartramiana</i>	84	<i>Quercus Douglasii</i> , var. <i>Gambelii</i>	139
<i>Pyrus Botryopium</i>	84	<i>Quercus Douglasii</i> , var. <i>Neesii</i>	138
<i>Pyrus coronaria</i>	72, 249, 253, 256, 259, 260, 284, 364, 428	<i>Quercus Drummondii</i>	139
<i>Pyrus coronaria</i> (<i>Pyrus angustifolia</i>).....	72	<i>Quercus dumosa</i>	155
<i>Pyrus coronaria</i> , var. <i>angustifolia</i>	72	<i>Quercus dumosa</i> , var. <i>bullata</i>	155
<i>Pyrus diversifolia</i>	73	<i>Quercus Durandii</i>	145, 249, 251, 254, 258, 316, 366, 452
<i>Pyrus fusca</i>	73	<i>Quercus echinacea</i>	153
<i>Pyrus glandulosa</i>	77	<i>Quercus echinoides</i>	155
<i>Pyrus microcarpa</i>	74	<i>Quercus elongata</i>	150
<i>Pyrus ovalis</i>	85	<i>Quercus Emoryi</i>	12, 146, 249, 254, 258, 265, 318, 386, 452
<i>Pyrus rivalaris</i>	73, 242, 284	<i>Quercus Emoryi</i> (<i>Quercus undulata</i> , var. <i>Gambelii</i>).....	139
<i>Pyrus sambucifolia</i>	12, 74, 250, 253, 256, 284, 364, 428	<i>Quercus falcata</i>	150, 249, 254, 258, 262, 265, 320, 350, 388, 454, 525
<i>Pyrus subcordata</i>	73	<i>Quercus falcata</i> , var. <i>Ludoviciana</i>	151
<i>Pyrus Wangenheimiana</i>	84	<i>Quercus falcata</i> , var. <i>pagodafolia</i>	151
		<i>Quercus falcata</i> , var. <i>triloba</i>	151
		<i>Quercus fulvescens</i>	146
		<i>Quercus Gambelii</i>	139
		<i>Quercus Garryana</i>	138, 249, 254, 257, 261, 312, 350, 382, 448, 576
Quaking Asp.....	171, 328, 350, 394, 460		
<i>Quassia dioica</i>	32		

	Page.		Page.
<i>Quercus Georgiana</i>	155	<i>Quercus Prinus</i>	142, 249, 254, 258, 261, 265, 316, 350, 384, 416, 450
<i>Quercus grisea</i>	144, 249, 251, 254, 258, 316, 386, 452	<i>Quercus Prinus</i> β. (<i>Quercus cinerea</i>)	153
<i>Quercus hastata</i>	146	<i>Quercus Prinus</i> (<i>Quercus Michauxii</i>)	141
<i>Quercus hemisphaerica</i>	152	<i>Quercus Prinus Chinquapin</i>	143
<i>Quercus hemisphaerica</i> , var. <i>nana</i>	152	<i>Quercus Prinus humilis</i>	142
<i>Quercus heterophylla</i>	153, 249, 255, 258, 320, 390, 456	<i>Quercus Prinus palustris</i>	141
<i>Quercus Hindsi</i>	138	<i>Quercus Prinus platanoides</i>	141
<i>Quercus humilis</i>	153	<i>Quercus Prinus pumila</i>	142
<i>Quercus hypoleuca</i>	154, 249, 255, 258, 322, 390, 456	<i>Quercus Prinus tomentosa</i>	141
<i>Quercus ilicifolia</i>	155	<i>Quercus Prinus</i> , var. <i>acuminata</i>	142
<i>Quercus imbricaria</i>	154, 249, 255, 258, 322, 390, 456	<i>Quercus Prinus</i> , var. <i>bicolor</i>	141
<i>Quercus Jacobi</i>	138	<i>Quercus Prinus</i> , var. <i>discolor</i>	141
<i>Quercus Kelloggii</i>	149, 250, 254, 258, 265, 320, 388, 416, 454	<i>Quercus Prinus</i> , var. <i>lata</i>	142
<i>Quercus laevis</i>	151	<i>Quercus Prinus</i> , var. <i>Michauxii</i>	141
<i>Quercus laurifolia</i>	152, 249, 254, 258, 320, 390, 454	<i>Quercus Prinus</i> , var. <i>monticola</i>	142
<i>Quercus laurifolia hybrida</i>	152	<i>Quercus Prinus</i> , var. <i>oblongata</i>	143
<i>Quercus laurifolia</i> , var. <i>acuta</i>	152	<i>Quercus Prinus</i> , var. <i>prinoides</i>	143
<i>Quercus laurifolia</i> , var. <i>obtus</i>	152	<i>Quercus pumila</i>	155
<i>Quercus lobata</i>	138, 249, 254, 257, 312, 382, 448	<i>Quercus pungens</i>	144
<i>Quercus lobata</i> , var. <i>fruticosa</i>	155	<i>Quercus Ransoni</i>	138
<i>Quercus lobulata</i>	139	<i>Quercus reticulata</i>	144, 249, 316
<i>Quercus longiglana</i>	138	<i>Quercus reticulata</i> , var. <i>Greggii</i>	144
<i>Quercus lyrata</i>	140, 249, 254, 257, 261, 314, 350, 384, 450, 533	<i>Quercus retusa</i>	145
<i>Quercus macrocarpa</i>	140, 249, 254, 257, 261, 265, 314, 384, 448	<i>Quercus rubra</i>	147, 250, 251, 254, 258, 262, 265, 318, 350, 386, 452
<i>Quercus macrocarpa</i> , var. <i>abbreviata</i>	140	<i>Quercus rubra</i> β. (<i>Quercus coccinea</i>)	148
<i>Quercus macrocarpa</i> , var. <i>minor</i>	140	<i>Quercus rubra</i> (<i>Quercus Kelloggii</i>)	149
<i>Quercus macrocarpa</i> , var. <i>oliveiformis</i>	140	<i>Quercus rubra</i> (<i>Quercus tinctoria</i>)	149
<i>Quercus Marylandica</i>	150	<i>Quercus rubra maxima</i>	147
<i>Quercus Michauxii</i>	141, 249, 254, 258, 261, 316, 384, 416, 450, 533	<i>Quercus rubra montana</i>	150
<i>Quercus montana</i>	142	<i>Quercus rubra ramosissima</i>	151
<i>Quercus Morehus</i>	147	<i>Quercus rubra</i> , var. <i>dissecta</i>	151
<i>Quercus Muhlenbergii</i>	143	<i>Quercus rubra</i> var. <i>latifolia</i>	147
<i>Quercus myrtifolia</i>	155	<i>Quercus rubra</i> , var. <i>montana</i>	147
<i>Quercus nana</i>	152	<i>Quercus rubra</i> , var. <i>runcinata</i>	147
<i>Quercus Nees</i>	138	<i>Quercus rubra</i> , var. <i>Texana</i>	148, 249, 254, 258, 318, 388, 452
<i>Quercus nigra</i>	150, 249, 254, 258, 265, 320, 350, 388, 454, 525	<i>Quercus San-Sabana</i>	145
<i>Quercus nigra</i> (<i>Quercus aquatica</i>)	152	<i>Quercus sempervirens</i>	145
<i>Quercus nigra</i> (<i>Quercus tinctoria</i>)	149	<i>Quercus sericea</i>	155
<i>Quercus nigra aquatica</i>	152	<i>Quercus Shumardii</i>	149
<i>Quercus nigra digitata</i>	150	<i>Quercus sinuata</i>	137
<i>Quercus nigra integrifolia</i>	150	<i>Quercus Sonomensis</i>	149
<i>Quercus nigra trifida</i>	152	<i>Quercus epicata</i>	144
<i>Quercus nigra</i> , var. (<i>Quercus heterophylla</i>)	153	<i>Quercus stellata</i>	139
<i>Quercus nigra</i> , var. <i>latifolia</i>	150	<i>Quercus stellata</i> , var. <i>depressa</i>	140
<i>Quercus oblongifolia</i>	148, 249, 251, 254, 258, 316, 386, 452	<i>Quercus stellata</i> , var. <i>Floridana</i>	139
<i>Quercus oblongifolia</i>	146	<i>Quercus stellata</i> , var. <i>Utahensis</i>	139
<i>Quercus oblongifolia</i> , var. <i>breviloba</i>	143	<i>Quercus Texana</i>	148
<i>Quercus obtusa</i>	152	<i>Quercus tinctoria</i>	149, 249, 254, 258, 262, 265, 318, 350, 388, 454, 528
<i>Quercus obtusifolia</i> , var. <i>breviloba</i>	145	<i>Quercus tinctoria</i> , var. <i>angulosa</i>	149
<i>Quercus obtusiloba</i>	138, 249, 254, 257, 261, 312, 382, 448, 525, 528	<i>Quercus tinctoria</i> , var. <i>Californica</i>	149
<i>Quercus obtusiloba</i> , var. <i>depressa</i>	140	<i>Quercus tinctoria</i> , var. <i>sinuosa</i>	149
<i>Quercus obtusiloba</i> , var. <i>parvifolia</i>	139	<i>Quercus triloba</i>	150
<i>Quercus Erstediana</i>	138	<i>Quercus uliginosa</i>	152
<i>Quercus oleoides</i>	145	<i>Quercus undulata</i>	13, 155
<i>Quercus oliveiformis</i>	140	<i>Quercus undulata</i> (<i>Quercus Durandii</i>)	145
<i>Quercus omyadenia</i>	146	<i>Quercus undulata</i> , var. <i>Gambelli</i>	139, 249, 254, 257, 314, 384, 448
<i>Quercus palustris</i>	151, 249, 254, 258, 320, 388, 454	<i>Quercus undulata</i> , var. <i>grisea</i>	144
<i>Quercus palustris</i> (<i>Quercus rubra</i> , var. <i>Texana</i>)	148	<i>Quercus undulata</i> , var. <i>oblongata</i>	143
<i>Quercus parvifolia</i>	139	<i>Quercus undulata</i> , var. <i>pungens</i>	144
<i>Quercus Phellos</i>	154, 249, 255, 258, 322, 390, 456	<i>Quercus undulata</i> , var. <i>Wrightii</i>	144
<i>Quercus Phellos angustifolia</i>	154	<i>Quercus vaccinifolia</i>	146
<i>Quercus Phellos</i> × <i>coccinea</i>	153	<i>Quercus velutina</i>	140
<i>Quercus Phellos latifolia</i>	154	<i>Quercus villosa</i>	139
<i>Quercus Phellos pumila</i>	155	<i>Quercus virens</i>	145, 249, 254, 258, 262, 265, 316, 388, 452
<i>Quercus Phellos</i> × <i>tinctoria</i>	158	<i>Quercus Virginiana</i>	145
<i>Quercus Phellos</i> , var. (<i>Quercus heterophylla</i>)	153	<i>Quercus Wislizeni</i>	147, 249, 254, 258, 318, 386, 452
<i>Quercus Phellos</i> , var. <i>arenaria</i>	155		
<i>Quercus Phellos</i> , var. <i>cinerea</i>	153		
<i>Quercus Phellos</i> , var. <i>humilis</i>	154		
<i>Quercus Phellos</i> , var. <i>imbricaria</i>	154		
<i>Quercus Phellos</i> , var. <i>laurifolia</i>	152		
<i>Quercus Phellos</i> , var. <i>sempervirens</i>	145		
<i>Quercus Phellos</i> , var. <i>sericea</i>	155		
<i>Quercus Phellos</i> , var. <i>viridis</i>	154		
<i>Quercus prinoides</i>	142, 249, 251, 254, 258, 261, 262, 265, 316, 384, 450		

B.

<i>Randia clusiaefolia</i>	95
<i>Rapanea Guyanensis</i>	99
Rattle-box	106, 296
Red Ash	109, 296, 370, 436
Red Bay	118, 302, 372, 438
Red Birch	161, 324, 392, 458
Red Cedar (<i>Juniperus Virginiana</i>)	183, 332, 398, 464, 544
Red Cedar (<i>Thuja gigantea</i>)	177, 330, 396, 462, 573, 575, 576, 580

	Page.		Page.
Red Cherry, Wild	66, 282, 428	<i>Rhus cotinoides</i>	52, 250, 276
Red Cypress	184, 334, 350, 398, 468	<i>Rhus cotinus</i>	52
Red Elm	122, 304, 374, 440	<i>Rhus glabra</i>	53
Red Fir (<i>Abies magnifica</i>)	214, 346, 412, 478	<i>Rhus hypselodendron</i>	52
Red Fir (<i>Abies nobilis</i>)	214, 346, 412, 478, 573	<i>Rhus leucantha</i>	53
Red Fir (<i>Pseudotsuga Douglasii</i>)	209, 265, 344, 352, 410, 478, 563, 575	<i>Rhus Metopium</i>	54, 249, 253, 256, 278, 360, 424
Red Fir the most important timber tree of the Pacific region	7	<i>Rhus Toxicodendron</i>	54
Red Gum	86, 286, 350, 364, 414, 430	<i>Rhus typhina</i>	52, 250, 278
Red Haw (<i>Crataegus coccinea</i>)	78, 286	<i>Rhus typhina</i> , var. <i>laciniata</i>	52
Red Haw (<i>Crataegus flava</i> , var. <i>pubescens</i>)	83, 286, 364, 430	<i>Rhus typhina</i> , var. <i>viridiflora</i>	52
Red Ironwood	39, 272, 358, 422	<i>Rhus venenata</i>	54, 250, 278
Red Maple	50, 276, 358, 424	<i>Rhus vernix</i>	54
Red Mulberry	128, 308, 376, 442	<i>Rhus viridiflora</i>	52
Red Oak (<i>Quercus falcata</i>)	151, 265, 320, 350, 388, 454	River Birch	161, 324, 392, 458
Red Oak (<i>Quercus rubra</i>)	148, 265, 318, 350, 388, 452	River Cottonwood	172, 328, 394, 400
Red Oak (<i>Quercus rubra</i> , var. <i>Texana</i>)	148, 318, 388, 452	Robinia	10, 13
Red Pine	192, 336, 350, 402, 468	<i>Robinia fragilis</i>	55
Red Stopper	89, 288, 368, 430	<i>Robinia glutinosa</i>	56
Redbud (<i>Cercis Canadensis</i>)	61, 280, 362, 426	<i>Robinia Neo-Mexicana</i>	13, 56, 249, 253, 256, 278, 360, 426
Redbud (<i>Cercis reniformis</i>)	61, 280	<i>Robinia Pseudacacia</i>	55, 249, 253, 256, 259, 278, 350, 360, 424
Redwood	185, 334, 350, 398, 460, 579	<i>Robinia viscosa</i>	56, 249, 278
Redwood Belt the most important feature in the Coast Range of the Coast Forest	8	Rock Chestnut Oak	142, 265, 316, 350, 384, 416, 450
Redwood of California, economic value of the	578	Rock Elm	122, 304, 374, 440
Redwood standing in the forests of California May 31, 1880, tabular statement of the amount of	579	Rock Maple	48, 276, 358, 422
Region, the Atlantic	2-6	Rocky Mountain region, northern, report by Mr. Sereno Watson on the forests of the	504, 565
Region, the Pacific	6-10	ROSEACEÆ	64-85, 253, 256, 282, 362, 426
Remarks, general, on the forests of North America	3-16	Rose Bay	89, 292, 368, 434
Reynolds	10, 13	Rosemary Pine	197, 340, 350, 404, 416, 470
<i>Reynoldsia latifolia</i>	39, 249, 358, 256, 272, 358, 422	Rosin and tar produced in North Carolina, grades of	517
<i>Rhamnidium revolutum</i>	39	Royal Palm	218, 348, 352
RHAMNACEÆ	39-42, 258, 256, 272, 358, 422	RUBIACEÆ	95, 96, 258, 257, 290, 366, 432
<i>Rhamnus</i>	10-13	Rum Cherry	68, 282, 362, 428
<i>Rhamnus alnifolia</i>	41	RUTACEÆ	29-32, 258, 256, 268, 356, 420
<i>Rhamnus Californica</i>	40, 250, 272		
<i>Rhamnus Californica</i> , var. <i>tomentella</i>	41	Sabal	11, 16
<i>Rhamnus Caroliniana</i>	40, 250, 253, 256, 272, 358, 422	Sabal Palmetto	217, 250, 348, 362
<i>Rhamnus ellipticus</i>	41	Saffron Plum	103, 294, 368, 434
<i>Rhamnus ferrea</i>	39	Sage-brush	575
<i>Rhamnus levigatus</i>	39	Saginaw valley in Michigan, lumber industry of	532
<i>Rhamnus laurifolia</i>	40	Sagunro	90, 288
<i>Rhamnus oleifolia</i>	40	SALICACEÆ	165-175, 255, 258, 328, 394, 400
<i>Rhamnus Purshiana</i>	41, 250, 253, 256, 272, 358, 422	Salix	11, 12, 16
<i>Rhamnus tomentellus</i>	41	<i>Salix</i> ———	170
Rhizophora	10, 11, 14	<i>Salix ambigua</i>	165
<i>Rhizophora Americana</i>	86	<i>Salix amygdaloides</i>	13, 166, 250, 255, 256, 328, 394, 460
<i>Rhizophora Mangle</i>	86, 249, 253, 257, 263, 288, 364, 430	<i>Salix argophylla</i>	168
<i>Rhizophora racemosa</i>	86	<i>Salix arguta</i>	167
RHIZOPHORACEÆ	86, 87, 253, 257, 288, 364, 430	<i>Salix arguta</i> , var. <i>lasiantha</i>	167
Rhode Island, lumber industry and manufactures from wood in	486, 487, 500, 501	<i>Salix Bigelovii</i>	170
Rhode Island, rank of, according to value of lumber products	487	<i>Salix Bigelovii</i> , var. <i>fuscor</i>	170
Rhode Island, statistics of forests, forest fires, standing timber, etc., in	491, 500, 501	<i>Salix brachycarpa</i>	168
Rhode Island, the Northern Pine Belt in	500	<i>Salix brachystachys</i>	170
Rhododendron	11, 12, 15	<i>Salix brachystachys</i> , var. <i>Scouleriana</i>	170
<i>Rhododendron maximum</i>	99, 250, 254, 257, 292, 368, 434	<i>Salix Caroliniana</i>	165
<i>Rhododendron maximum</i> , var. <i>album</i>	99	<i>Salix chlorophylla</i> , var. <i>pellita</i>	171
<i>Rhododendron maximum</i> , var. <i>purpureum</i>	99	<i>Salix cordata</i>	170
<i>Rhododendron maximum</i> , var. <i>roseum</i>	99	<i>Salix cordata</i> , var. <i>falcata</i>	165
<i>Rhododendron procerrum</i>	99	<i>Salix cordata</i> , var. <i>vestita</i>	170, 250, 328
<i>Rhododendron purpureum</i>	99	<i>Salix crassa</i>	169
<i>Rhododendron Purshii</i>	99	<i>Salix cuneata</i>	171
<i>Rhus</i>	10, 12, 14	<i>Salix discolor</i>	169, 250, 328
<i>Rhus arborescens</i>	53	<i>Salix discolor</i> , var. <i>eriocephala</i>	169
<i>Rhus Canadensis</i>	52	<i>Salix discolor</i> , var. <i>prinoides</i>	169
<i>Rhus copallina</i>	53, 250, 253, 256, 278, 360, 424	<i>Salix eriocephala</i>	169
<i>Rhus copallina</i> , var. <i>angustialata</i>	53	<i>Salix exigua</i>	168
<i>Rhus copallina</i> , var. <i>angustifolia</i>	53	<i>Salix falcata</i>	165
<i>Rhus copallina</i> , var. <i>integrifolia</i>	53	<i>Salix Fendleriana</i>	167
<i>Rhus copallina</i> , var. <i>lanceolata</i>	53, 250, 278, 424	<i>Salix flavescens</i>	169, 250, 255, 258, 328, 394, 460
<i>Rhus copallina</i> , var. <i>latifolia</i>	53	<i>Salix flavescens</i> (Salix <i>flavescens</i> , var. <i>Scouleriana</i>)	170
<i>Rhus copallina</i> , var. <i>latifolia</i>	53	<i>Salix flavescens</i> , var. <i>Scouleriana</i>	170, 250, 255, 258, 328, 394, 460
<i>Rhus copallina</i> , var. <i>leucantha</i>	53	<i>Salix flavo-virens</i>	165
<i>Rhus copallina</i> , var. <i>serrata</i>	53	<i>Salix fluviatilis</i>	168
		<i>Salix Hartwegi</i>	171
		<i>Salix Hindiana</i>	169

	Page.		Page.
<i>Salix Hindsiana</i> , var. <i>tenusifolia</i>	169	<i>Sapindus falcatus</i>	44
<i>Salix Hoffmanniana</i>	167	<i>Sapindus inaequalis</i>	44
<i>Salix Hookeriana</i>	170, 250, 328, 400	<i>Sapindus marginatus</i>	12, 44, 249, 253, 274, 358, 414, 422
<i>Salix Houstoniana</i>	165	<i>Sapindus Saponaria</i>	45, 249, 274
<i>Salix humilis</i> , var. <i>Hartwegi</i>	171	<i>Sapindus Saponaria</i> (<i>Sapindus marginatus</i>).....	44
<i>Salix laevigata</i>	167, 250, 255, 258, 368, 394, 400	SAPOTACEÆ.....	100-103, 254, 257, 292, 368, 434
<i>Salix laevigata</i> , var. <i>angustifolia</i>	167	<i>Sarcophagus Carolinianus</i>	40
<i>Salix laevigata</i> , var. <i>congesta</i>	167	<i>Sassafras</i>	120, 302, 374, 438
<i>Salix lanceifolia</i>	167	<i>Sassafras officinale</i>	119, 250, 254, 257, 260, 302, 374, 438
<i>Salix lasiandra</i>	167, 250, 328	Satinwood.....	11, 15, 31, 270, 356, 420
<i>Salix lasiandra</i> , var. <i>Fendleriana</i>	167, 250, 255, 258, 328, 394, 400	Savin (<i>Juniperus Virginiana</i>).....	183, 322, 398, 464
<i>Salix lasiandra</i> , var. <i>lancofolia</i>	167, 250, 255, 258, 328, 394, 400	Savin (<i>Torreya taxifolia</i>).....	186, 334, 400, 466
<i>Salix lasiolepis</i>	170, 250, 255, 258, 328, 394, 400	Saw-mills in each state and territory, average size and product of.....	488
<i>Salix lasiolepis</i> , var. <i>angustifolia</i>	171	Scarlet Haw (<i>Crataegus coccinea</i>).....	78, 286
<i>Salix lasiolepis</i> , var. <i>Bigelovii</i>	170	Scarlet Haw (<i>Crataegus subvillosa</i>).....	78, 286, 364, 430
<i>Salix lasiolepis</i> , var. <i>fallax</i>	170	Scarlet Oak.....	148, 318, 388, 452
<i>Salix lasiolepis</i> , var. <i>latifolia</i>	171	Schæfferia.....	10, 18
<i>Salix ligustrina</i>	165	<i>Schæfferia buxifolia</i>	39
<i>Salix longifolia</i>	12, 168, 250, 328	<i>Schæfferia completa</i>	30
<i>Salix longifolia</i> , var. <i>angustissima</i>	168	<i>Schæfferia frutescens</i>	39, 249, 272
<i>Salix longifolia</i> , var. <i>argyrophylla</i>	168	<i>Schæfferia lateriflora</i>	120
<i>Salix longifolia</i> , var. <i>argyrophylla angustissima</i>	168	<i>Schoubea commutata</i>	87
<i>Salix longifolia</i> , var. <i>argyrophylla opaca</i>	168	<i>Schubertia disticha</i>	183
<i>Salix longifolia</i> , var. <i>exigua</i>	168, 250, 328	<i>Schubertia sempervirens</i>	185
<i>Salix longifolia</i> , var. <i>opaca</i>	168	Screw Bean.....	62, 280, 362, 426
<i>Salix longifolia</i> , var. <i>pedicellata</i>	168	Screw-pod Mesquit.....	62, 280, 362, 426
<i>Salix longipes</i>	166	Scrub Oak (<i>Quercus Catesbeii</i>).....	151, 320, 388, 454
<i>Salix longipes</i> , var. <i>pubescens</i>	166	Scrub Oak (<i>Quercus undulata</i> , var. <i>Gambellii</i>).....	139, 314, 384, 448
<i>Salix lucida</i> , var. <i>angustifolia</i> , forma <i>lasiandra</i>	167	Scrub Pine (<i>Pinus Banksiana</i>).....	201, 342, 352, 406, 472
<i>Salix lucida</i> , var. <i>macrophylla</i>	167	Scrub Pine (<i>Pinus clausa</i>).....	199, 340, 404, 472
<i>Salix marginata</i>	166	Scrub Pine (<i>Pinus contorta</i>).....	194, 338, 404, 470
<i>Salix melanopis</i>	166	Scrub Pine (<i>Pinus inops</i>).....	199, 340, 350, 404, 470, 529
<i>Salix nigra</i>	12, 165, 250, 328, 394, 400	<i>Scutia ferrea</i> (<i>Condalia ferrea</i>).....	39
<i>Salix nigra</i> , var. <i>amygdaloides</i>	166	<i>Scutia ferrea</i> (<i>Reynosia latifolia</i>).....	39
<i>Salix nigra</i> , var. <i>angustifolia</i>	166	Sea Ash.....	30, 270, 356, 420
<i>Salix nigra</i> , var. <i>falcata</i>	165	Sea Grape.....	113, 302, 438
<i>Salix nigra</i> , var. <i>latifolia</i>	166	Seaside Alder.....	162, 328, 458
<i>Salix nigra</i> , var. <i>longifolia</i>	166	Sebastiania.....	11, 15
<i>Salix nigra</i> , var. <i>longipes</i>	166	<i>Sebastiania lucida</i>	121, 249, 304
<i>Salix nigra</i> , var. <i>longipes</i> , subvar. <i>gongylocarpa</i>	166	<i>Sebestena scabra</i>	113
<i>Salix nigra</i> , var. <i>longipes</i> , subvar. <i>venulosa</i>	166	Selected woods of the United States under transverse strain, behavior of: specimens eight centimeters square.....	414-417
<i>Salix nigra</i> , var. <i>marginata</i>	166	Semi-tropical Forest of Florida.....	6
<i>Salix nigra</i> , var. <i>Wardii</i>	166	Sequoia.....	8, 11, 12, 16
<i>Salix nigra</i> , var. <i>Wrightii</i>	166	<i>Sequoia gigantea</i>	184, 251, 255, 258, 334, 398, 466, 578
<i>Salix occidentalis</i>	166	<i>Sequoia gigantea</i> (<i>Sequoia sempervirens</i>).....	185
<i>Salix pentandra</i>	165	<i>Sequoia Rafinesquii</i>	206
<i>Salix pentandra</i> , var. <i>caudata</i>	167	<i>Sequoia religiosa</i>	185
<i>Salix prinoides</i>	169	<i>Sequoia sempervirens</i>	184, 250, 255, 258, 263, 334, 350, 398, 466, 579
<i>Salix Purshiana</i>	165	<i>Sequoia Wellingtonia</i>	184
<i>Salix rubra</i>	168	Service Tree.....	84, 286, 364, 430
<i>Salix Scouleriana</i>	170	Seven-year Apple.....	95, 290
<i>Salix sensitiva</i>	169	Shad Bush.....	84, 286, 364, 430
<i>Salix sessilifolia</i>	168, 250, 328	Shag-bark Hickory.....	183, 308, 350, 378, 444
<i>Salix sessilifolia</i> , var. <i>Hindsiana</i>	169	Sharples, Mr. S. P., experiments by.....	247, 251, 264, 265
<i>Salix sessilifolia</i> , var. <i>villosa</i>	168	She Balsam.....	210, 346, 412, 478
<i>Salix Sitchenensis</i>	171, 250, 328	Sheepberry.....	94, 290, 432
<i>Salix Sitchenensis</i> , var. <i>angustifolia</i>	171	Shell-bark, Big.....	183, 310, 378, 444
<i>Salix speciosa</i>	167	Shell-bark, Bottom.....	183, 310, 378, 444
<i>Salix subvillosa</i>	166	Shell-bark Hickory.....	183, 308, 350, 378, 444
<i>Salix Wrightii</i>	166	Shingle Oak.....	154, 322, 390, 456
<i>Samara floribunda</i>	99	Shingles and lumber received at Chicago during the year 1880.....	548, 549
<i>Samara pentandra</i>	99	Shittimwood (<i>Bumelia lanuginosa</i>).....	102, 294, 368, 434
<i>Sambucus</i>	10-12, 14	Shittimwood (<i>Rhamnus Purshiana</i>).....	41, 272, 358, 423
<i>Sambucus Californica</i>	93	Short- and Long-leaved Pine standing in the forests of Alabama May 31, 1880, tabular statement of the amount of.....	534
<i>Sambucus glauca</i>	93, 250, 253, 257, 290, 366, 432	Short- and Long-leaved Pine standing in the forests of Louisiana May 31, 1880, tabular statement of the amount of.....	537
<i>Sambucus glauca</i> (<i>Sambucus Mexicana</i>).....	93	Short- and Long-leaved Pine standing in the forests of Mississippi May 31, 1880, tabular statement of the amount of.....	531
<i>Sambucus Mexicana</i>	12, 93, 250, 290	Short-leaved, Loblolly, and Long-leaved Pine standing in the forests of Texas May 31, 1880, tabular statement of the amount of.....	541
<i>Sambucus Mexicana</i> (<i>Sambucus glauca</i>).....	93	Short-leaved Pine.....	200, 340, 350, 406, 472, 524, 527, 530, 531, 536, 537, 541, 544, 560
<i>Sambucus velutina</i>	93	Short-leaved Pine standing in the forests of Arkansas May 31, 1880, tabu- lar statement of the amount of.....	544
Sand Jack.....	153, 320, 390, 456	Shrubby Trefoll.....	31, 270
Sand Pine.....	199, 340, 464, 472		
Sand-bar Willow.....	168, 326		
SAPINDACEÆ.....	42-51, 253, 256, 274, 358, 414, 422		
<i>Sapindus</i>	10-13		
<i>Sapindus acuminata</i>	44		
<i>Sapindus Drummondii</i>	44		

GENERAL INDEX.

607

	Page.		Page.
Sideroxylon.....	11, 15	Southern Crab Apple.....	72, 284
Sideroxylon Carolinense.....	101	Southern Cypress, next to the Long-leaved Pine, the characteristic tree of the Southern Maritime Pine Belt.....	4
Sideroxylon chrysophyllodes.....	101	Southern Maritime Pine Belt in the Atlantic region.....	4
Sideroxylon euneatum.....	103	Southern Pine.....	202, 342, 352, 406, 416, 472
Sideroxylon decandrum.....	103	Southern Pine Belt in Florida.....	520
Sideroxylon lanuginosum.....	102	Spanish Bayonet (Yucca baccata).....	219, 348
Sideroxylon laeve.....	103	Spanish Bayonet (Yucca canaliculata).....	218, 348
Sideroxylon lycioides.....	103	Spanish Bayonet (Yucca elata).....	219, 348
Sideroxylon Mastichodendron.....	101, 249, 254, 257, 292, 368, 434	Spanish Buckeye.....	44, 274, 422
Sideroxylon pallidum.....	101	Spanish Oak.....	151, 265, 320, 350, 388, 454, 528
Sideroxylon reclinatum.....	103	Spanish Oak, Swamp.....	152, 320, 388, 454
Sideroxylon salicifolium.....	101	Spanish Stopper.....	88, 288, 366, 430
Sideroxylon sericeum.....	101	Species, distribution of.....	12
Sideroxylon tenax (Bumelia lanuginosa).....	102	Specific gravity and ash of woods.....	248-251
Sideroxylon tenax (Bumelia tenax).....	101	Specific gravity, ash, and weight per cubic foot of dry specimens of the woods of the United States.....	266-349
Siliquastrum cordatum.....	61	Specific gravity and transverse strength of certain woods, relation between.....	259-264
Silk-top Palmetto.....	217, 348	Specimens of woods, where deposited.....	247, 248
Silky Willow.....	171, 328	Speckled Alder.....	165, 328, 394, 460
Silver-bell Tree (Halesia diptera).....	105, 294, 368, 434	Spice Tree.....	120, 302, 374, 440
Silver-bell Tree (Halesia tetraptera).....	106, 296	Spindle Tree.....	38, 272
Silver Maple.....	49, 276, 358, 424	Spiraea Californica.....	70
Silver-top Palmetto.....	218, 348	Spoonwood.....	98, 292, 368, 434
Simaruba.....	10, 13	Spruce, a species of, the characteristic tree of the Colorado mountains.....	9
Simaruba amara.....	32	Spruce, Black.....	203, 265, 342, 352, 406, 474, 496
Simaruba glauca.....	32, 250, 253, 256, 270, 356, 420	Spruce, Blue.....	205, 344, 408, 474
Simaruba medicinale.....	32	Spruce Pine (Pinus clausa).....	199, 340, 404, 472
SIMARUBA.....	32, 253, 256, 270, 356, 420	Spruce Pine (Pinus glabra).....	201, 340, 406, 472, 527
Sitka Cypress.....	178, 332, 398, 464	Spruce Pine (Pinus mitis).....	200, 340, 350, 406, 472
Size, average, and product of saw-mills in each state and territory.....	488	Spruce Pine (Pinus Murrayana).....	195, 338, 350, 404, 470
Slash Pine.....	202, 342, 352, 406, 416, 474, 516	Spruce standing in the forests of New Hampshire May 31, 1880, tabular statement of the amount of.....	496
Slippery Elm (Fremontia Californica).....	26, 268	Spruce standing in the forests of Vermont May 31, 1880, tabular statement of the amount of.....	498
Slippery Elm (Ulmus fulva).....	122, 304, 374, 440	Spruce, Tide-land.....	206, 344, 408, 474, 580
Sloe.....	67, 282, 428	Spruce, White, the most important tree in the Pacific region.....	7
Sloe, Black.....	67, 282, 428	Spruce, White (Picea alba).....	204, 342, 408, 474, 562, 564
Small-fruited Haw.....	81, 286, 364, 430	Spruce, White (Picea Engelmanni).....	205, 265, 342, 408, 474, 564, 573, 574
Smooth Alder.....	164, 326	Spruce, White (Picea pungens).....	205, 344, 408, 474
Snow-drop Tree (Halesia diptera).....	105, 294, 368, 434	Spruces, Black and White, the characteristic trees of the Northern Forest.....	8
Snow-drop Tree (Halesia tetraptera).....	106, 296	Stag Bush.....	94, 290, 366, 432
Soapberry (Sapindus marginatus).....	44, 274, 358, 414, 422	Staghorn Sumach.....	53, 278
Soapberry (Sapindus Saponaria).....	45, 274	Standing timber, forests, forest fires, etc., extent of, in the—	
Soft Maple (Acer dasycarpum).....	49, 276, 358, 424	North Atlantic division.....	494-510
Soft Maple (Acer rubrum).....	50, 276, 358, 424	Northern Central division.....	547-563
Sophora.....	10, 11, 14	South Atlantic division.....	511-522
Sophora affinis.....	58, 249, 253, 256, 280, 360, 426	Southern Central division.....	524-546
Sophora secundiflora.....	57, 249, 278	Western division.....	564-586
Sophora speciosa.....	57	Standing timber, forests, forest fires, etc., statistics of, in—	
Sorbus Americana.....	73	Alabama.....	491, 524-530
Sorbus Americana, var. microcarpa.....	74	Alaska.....	530
Sorbus aucuparia (Pyrus Americana).....	73	Arizona.....	491, 568, 569
Sorbus aucuparia (Pyrus sambucifolia).....	74	Arkansas.....	491, 548, 544
Sorbus aucuparia, var. a. (Pyrus Americana, var. microcarpa).....	74	California.....	491, 578-580
Sorbus aucuparia, var. b. (Pyrus sambucifolia).....	74	Colorado.....	491, 567, 568
Sorbus aucuparia, var. Americana.....	73	Connecticut.....	500, 501
Sorbus humifusa.....	73	Dakota.....	491, 561, 562
Sorbus microcarpa.....	74	Delaware.....	491, 511
Sorbus riparia.....	74	District of Columbia.....	511
Sorbus sambucifolia.....	74	Florida.....	491, 520-522
Sorbus Sitchensis.....	74	Georgia.....	491, 519, 520
Sorrel Tree.....	98, 292, 368, 434	Idaho.....	491, 571-578
Sour Gum.....	92, 290, 366, 432	Illinois.....	491, 547-550
Sour Tupelo.....	91, 290, 366, 432	Indiana.....	491, 547
Sourwood.....	98, 292, 368, 434	Indian territory.....	491, 543
South Atlantic division, extent of forests, forest fires, standing timber, etc., in the.....	511-523	Iowa.....	491, 560
South Carolina, lumber industry and manufactures from wood in.....	486, 487, 518, 519	Kansas.....	491, 562, 563
South Carolina, production of naval stores in.....	517	Kentucky.....	491, 545, 546
South Carolina, rank of, according to value of lumber products.....	487	Louisiana.....	491, 536, 540
South Carolina, statistics of forests, forest fires, standing timber, etc., in.....	491, 518, 519	Maine.....	491, 494-496
South Carolina, tabular statement of the amount of Long-leaved Pine standing in the forests of, May 31, 1880.....	519	Maryland.....	491-511
South Carolina, the Coast Pine Belt in.....	518	Massachusetts.....	491, 500, 501
Southern Buckthorn.....	103, 294, 368, 434	Michigan.....	491, 550-554
Southern Central division, extent of forests, forest fires, standing timber, etc., in the.....	524-546	Minnesota.....	491, 558-560
		Mississippi.....	491, 530-536
		Missouri.....	491, 560-561
		Montana.....	491, 564, 566
		Nebraska.....	562
		Nevada.....	491, 571
		New Hampshire.....	491, 498-498
		New Jersey.....	491, 506
		New Mexico.....	491, 568
		New York.....	491, 501-506
		North Carolina.....	491, 515-518
		Ohio.....	491, 547
		Oregon.....	491, 576-578
		Pennsylvania.....	491, 506-510
		Rhode Island.....	491, 500, 501

	Page.		Page.
Standing timber, forests, forest fires, etc., statistics of, in—continued.		Sycamore (<i>Platanus occidentalis</i>)	129, 306, 350, 376, 442
South Carolina	491, 518, 519	Sycamore (<i>Platanus racemosa</i>)	129, 306, 376, 442
Tennessee	492, 544, 545	Sycamore (<i>Platanus Wrightii</i>)	130, 306, 376, 442
Texas	492, 540-543	Symplocos	11, 15
Utah	492, 569-571	Symplocos tinctoria	105, 250, 254, 257, 294, 368, 434
Vermont	492, 498-500		
Virginia	492, 511, 512	Table-mountain Pine	T. 199, 340, 350, 404, 472
Washington	492, 573-576	Tacamahac	173, 328, 394, 460
West Virginia	492, 512-515	Tallowberry	28, 268, 356, 420
Wisconsin	492, 554-558	Tallow Nut	34, 270
Wyoming	492, 568, 567	Tamarack (<i>Larix Americana</i>)	215, 346, 352, 412, 478
Star-leaved Gum	85, 288, 350, 364, 414, 430	Tamarack (<i>Larix occidentalis</i>)	216, 348, 414, 480
Statistics of the lumbering industry of the United States for year ending		Tamarack (<i>Pinus Murrayana</i>)	195, 338, 350, 404, 470
May 31, 1880	486, 487	Tamarind, Wild	64, 282, 362, 426
STERCULIACEÆ	26, 268	Tanbark Oak	155, 265, 322, 390, 456
Stinking Cedar (<i>Torreya Californica</i>)	186, 334, 400, 466	Tan Bay	25, 265, 268, 354, 418
Stinking Cedar (<i>Torreya taxifolia</i>)	186, 334, 400, 466, 521	Tannin in the bark of certain species, per cent. of	265
Stopper (<i>Eugenia longipes</i>)	89, 288	Tannin values of the bark of various trees	265
Stopper (<i>Eugenia monticola</i>)	89, 288, 366, 430	Tar and rosin produced in North Carolina, grades of	517
Stopper, Gurgeon	88, 288, 366, 430	Tar, turpentine, and rosin industry of North Carolina	516-518
Stopper, Red	89, 288, 366, 430	<i>Taxodium</i> species (<i>Sequoia sempervirens</i>)	185
Stopper, Spanish	88, 288, 366, 430	<i>Taxodium</i>	4, 11, 16
Stopper, White	89, 288, 366, 430	<i>Taxodium adscendens</i>	183
Strain, transverse, behavior of selected woods of the United States under:		<i>Taxodium distichum</i>	183, 250, 255, 258, 334, 350, 396, 466
specimens eight centimeters square	414-417	<i>Taxodium distichum fastigiatum</i>	183
Strain, transverse, behavior of the principal woods of the United States		<i>Taxodium distichum, var. microphyllum</i>	183
under	354-414	<i>Taxodium distichum, var. nutans</i>	183
Strength of woods, method of determining the	252	<i>Taxodium distichum, var. patens</i>	183
Strength, transverse, and specific gravity of certain woods, relation between	259-264	<i>Taxodium giganteum</i>	184
Striped Dogwood	46, 274	<i>Taxodium microphyllum</i>	183
Striped Maple	46, 274	<i>Taxodium</i> , next to the Long-leaved Pine, the characteristic tree of the	
<i>Strombocarpa odorata</i>	62	Southern Maritime Pine Belt	4
<i>Strombocarpa pubescens</i>	63	<i>Taxodium sempervirens</i>	185
Strong Bark	114, 300, 372, 438	<i>Taxodium Washingtonianum</i>	184
<i>Styphnolobium affine</i>	58	Taxus	11, 12, 16, 572, 573
SYTRACACEÆ	105, 106, 254, 257, 294, 368, 434	<i>Taxus baccata</i> (<i>Taxus brevifolia</i>)	185
Sugar and molasses, maple, produced in the United States in 1879,		<i>Taxus baccata, var. Canadensis</i>	185
amount of	485	<i>Taxus Bourserii</i>	185
Sugarberry	126, 306, 376, 414, 442	<i>Taxus brevifolia</i>	185, 250, 255, 258, 334, 400, 466
Sugar Maple	48, 276, 358, 422	<i>Taxus Canadensis</i>	185
Sugar Maple, Black	49, 276, 350, 358, 424	<i>Taxus Florida</i>	186, 250, 334, 521
Sugar, maple, product of Michigan	551	<i>Taxus Lindleyana</i>	185
Sugar, maple, product of New York	501	<i>Taxus montana</i>	186
Sugar, maple, product of Vermont	498	Tennessee, effect of fires upon the forests of	546
Sugar Pine	183, 336, 350, 400, 466, 576, 578	Tennessee, lumber industry and manufactures from wood in	486, 487, 545
Sugar Pine an important tree in the Coast Forest	8	Tennessee, rank of, according to value of lumber products	487
Sugar Tree	48, 276, 358, 422	Tennessee, statistics of forests, forest fires, standing timber, etc., in	492, 544, 545
Sumach, Coral	54, 278, 360, 424	Tennessee valley in Alabama, forests of the	528, 529
Sumach, Dwarf	53, 278, 360, 424	TERNSTROMIACEÆ	25, 26, 263, 266, 268, 354, 418
Sumach, Poison	54, 278	<i>Tetranthera Californica</i>	120
Sumach, Staghorn	53, 278	Texas, Cedar an important tree in	540
Summer Haw (<i>Crataegus flava</i>)	83, 286	Texas, lumber industry and manufactures from wood in	486, 487, 541, 542
Summer Haw (<i>Crataegus flava, var. pubescens</i>)	83, 286, 364, 430	Texas, rank of, according to value of lumber products	487
Susquehanna boom at Williamsport, Pennsylvania, number of logs rafted		Texas, remarks by Dr. Charles Mohr on the forests of	542, 543
out of, 1862 to 1880	508	Texas, statistics of forests, forest fires, standing timber, etc., in	492, 540-543
Suwarrow	90, 288	Texas, tabular statement of the amount of Long-leaved, Loblolly, and	
Swamp Cottenwood	172, 328, 394, 460	Short-leaved Pine standing in the forests of, May 31, 1880	541
Swamp Hickory (<i>Carya amara</i>)	135, 310, 350, 380, 446	Texas, the pecan-nut an important product west of the Colorado river in	543
Swamp Hickory (<i>Carya aquatica</i>)	136, 310, 350, 380, 414, 446	Thatch, Brickley	218, 348
Swamp Laurel	20, 266, 354, 414, 418	Thatch, Brittle	218, 348
Swamp Maple	50, 276, 358, 424	The Joshua	219, 348
Swamp Pine	202, 342, 352, 406, 416, 474	Thorn, Black	79, 286, 364, 430
Swamp Post Oak	140, 314, 350, 384, 450	Thorn, Cockspur	76, 286, 364, 430
Swamp Spanish Oak	152, 320, 388, 454	Thorn, Newcastle	76, 286, 364, 430
Swamp White Oak	141, 314, 384, 450	Thorn, Washington	81, 286
Swamps, Cypress, of the Tenase river in Alabama	525-527	Thorn, White	78, 286
Sweet Bay	20, 266, 354, 414, 418	Three-thorned Acacia	50, 280, 360, 426
Sweet Birch	162, 324, 392, 458	Thrinax	11, 16
Sweet Buckeye	43, 274	<i>Thrinax argentea</i>	218, 249, 348
Sweet Gum	86, 286, 350, 364, 414, 430	<i>Thrinax Garberi</i>	217
Sweet Leaf	105, 294, 368, 434	<i>Thrinax parviflora</i>	217, 250, 348
Sweet Locust	59, 280, 360, 426	Thuya	7, 11, 12, 16, 566, 572, 573, 575, 576
Sweet-scented Crab	72, 284, 364, 428	<i>Thuya Oraigana</i>	176
Swietenia	10, 13	<i>Thuya excelsa</i>	178
Swietenia Mahagoni	83, 249, 253, 256, 270, 350, 356, 420	<i>Thuya gigantea</i>	177, 251, 255, 258, 330, 396, 462, 565, 566, 572, 573, 575, 576, 580
<i>Swietenia Senegalensis</i>	83		
Switch-bud Hickory	134, 310, 350, 380, 446		

	Page.		Page.
<i>Thuja gigantea</i> (Libocedrus decurrens).....	176	Timber, standing, forests, forest fires, etc., statistics of, in—continued.	
<i>Thuja Lobbii</i>	177	Texas.....	492, 546, 548
<i>Thuja Menziesii</i>	177	Utah.....	492, 568-571
<i>Thuja obtusa</i>	176	Vermont.....	492, 498-509
<i>Thuja occidentalis</i>	176, 251, 255, 258, 262, 330, 350, 398, 462, 552	Virginia.....	492, 511, 512
<i>Thuja occidentalis</i> , var. <i>plicata</i>	177	Washington.....	492, 572-576
<i>Thuja odorata</i>	176	West Virginia.....	492, 512-515
<i>Thuja plicata</i>	177	Wisconsin.....	492, 554-558
<i>Thuja Sibirica</i>	176	Wyoming.....	492, 566, 567
<i>Thuja sphaeroidalis</i>	177	Tili.....	38, 272, 356, 439
<i>Thuja sphaeroides</i>	177	Tillon.....	84, 298
<i>Thuja Wareana</i>	178	Toothache Tree (<i>Xanthoxylum Americanum</i>).....	29, 298
<i>Thuyopsis borealis</i>	178	Toothache Tree (<i>Xanthoxylum Clava-Heronilla</i>).....	30, 270, 356, 429
<i>Thuyopsis cupressoides</i>	178	Torchwood.....	33, 270, 356, 429
<i>Thuyopsis Tehuacensis</i>	178	Tornilla.....	62, 280, 293
<i>Thuyopsis Tehuacensis</i>	178	Torreya.....	11, 12, 16, 521
<i>Thylax fraxineum</i>	29	Torreya Californica.....	198, 250, 255, 258, 334, 400, 468
Tide-land Spruce.....	206, 344, 408, 474, 580	Torreya Myristica.....	198
Tilia.....	10, 11, 13	Torreya taxifolia.....	198, 250, 255, 258, 334, 400, 468, 521
<i>Tilia alba</i>	27	<i>Toxicodendron arborescens</i>	33
<i>Tilia Americana</i>	26, 250, 253, 256, 268, 356, 418	<i>Toxylon MacLurea</i>	128
<i>Tilia Americana</i> (<i>Tilia Americana</i> , var. <i>pubescens</i>).....	27	Toyon.....	84, 298
<i>Tilia Americana</i> , var. <i>heterophylla</i>	27	Transmittal, letter of.....	ix
<i>Tilia Americana</i> , var. <i>pubescens</i>	27, 250, 253, 256, 268, 356, 420	Transverse strain, behavior of selected woods of the United States under: specimens eight centimeters square.....	414-417
<i>Tilia Americana</i> , var. <i>Walleri</i>	27	Transverse strain, behavior of the principal woods of the United States under.....	354-414
<i>Tilia Canadensis</i>	26	Transverse strength and specific gravity of certain woods, relation between.....	259-264
<i>Tilia Caroliniana</i>	27	Trees of the United States, weight of the dry wood of.....	249-251
<i>Tilia glabra</i>	26	Trees, tannin values of the bark of various.....	265
<i>Tilia grata</i>	27	Trefoil, Shrubby.....	31, 270
<i>Tilia heterophylla</i>	27, 250, 253, 256, 268, 356, 420	<i>Trilopus dentata</i>	85
<i>Tilia heterophylla</i> , var. <i>alba</i>	27	<i>Trilopus nigra</i>	85
<i>Tilia latifolia</i>	26	<i>Trilopus parvifolia</i>	85
<i>Tilia lasiflora</i> (<i>Tilia Americana</i> , var. <i>pubescens</i>).....	27	<i>Trilopus rotundifolia</i>	85
<i>Tilia lasiflora</i> (<i>Tilia heterophylla</i>).....	27	<i>Trilopus Virginiana</i>	85
<i>Tilia neglecta</i>	26	Tunga.....	11, 12, 16, 573
<i>Tilia nigra</i>	26	Tunga Canadensis.....	206, 250, 255, 258, 263, 264, 265, 344, 382, 406, 474, 506
<i>Tilia pubescens</i>	27	Tunga Caroliniana.....	207, 250, 255, 258, 344, 410, 476
<i>Tilia pubescens</i> , var. <i>leptophylla</i>	27	Tunga Douglasii.....	209
<i>Tilia stenopetala</i>	27	Tunga Lindleyana.....	209
<i>Tilia truncata</i>	27	Tunga Mortensiana.....	207, 250, 255, 258, 265, 344, 410, 476, 565, 572, 573, 576
TILIACEAE.....	26-28, 253, 256, 268, 356, 418	Tunga Pattoniana.....	7, 208, 250, 255, 258, 265, 344, 410, 476, 565, 573
Timber tree of the Pacific region, the Red Fir the most important.....	7	<i>Tulipastrum Americanum</i>	20
Timber, standing, forests, forest fires, etc., extent of, in the—		<i>Tulipastrum Americanum</i> , var. <i>subcordatum</i>	20
North Atlantic division.....	494-510	<i>Tulipifera Liriodendron</i>	23
Northern Central division.....	547-563	Tulip Tree.....	22, 296, 350, 354, 418
South Atlantic division.....	511-523	Tupelo.....	92, 290, 366, 432
Southern Central division.....	524-546	Tupelo Gum.....	93, 290, 350, 366, 432
Western division.....	504-580	Tupelo, Large.....	93, 290, 350, 366, 432
Timber, standing, forests, forest fires, etc., statistics of, in—		Tupelo, Sour.....	91, 290, 366, 432
Alabama.....	491, 524-530	Turkey Oak.....	151, 320, 388, 454
Alaska.....	580		
Arizona.....	491, 568, 569		
Arkansas.....	491, 543, 544		
California.....	491, 578-580		
Colorado.....	491, 567, 568		
Connecticut.....	491, 500, 501		
Dakota.....	491, 561, 562		
Delaware.....	491, 511		
District of Columbia.....	511		
Florida.....	491, 520, 523		
Georgia.....	491, 519, 520		
Idaho.....	491, 571-573		
Illinois.....	491, 547-550		
Indiana.....	491, 547		
Indian territory.....	543		
Iowa.....	491, 566		
Kansas.....	491, 562, 563		
Kentucky.....	491, 545, 546		
Louisiana.....	491, 536-540		
Maine.....	491, 494-496		
Maryland.....	491, 511		
Massachusetts.....	491, 500, 501		
Michigan.....	491, 550-554		
Minnesota.....	491, 558-560		
Mississippi.....	491, 530-536		
Missouri.....	491, 560, 561		
Montana.....	491, 564-566		
Nebraska.....	491, 562		
Nevada.....	491, 571		
New Hampshire.....	491, 496-498		
New Jersey.....	491, 506		
New Mexico.....	491, 568		
New York.....	491, 501-506		
North Carolina.....	491, 515-518		
Ohio.....	491, 547		
Oregon.....	491, 576-578		
Pennsylvania.....	491, 506-510		
Rhode Island.....	491, 500, 501		
South Carolina.....	491, 518, 519		
Tennessee.....	492, 544, 545		

	Page.
<i>Ulmus racemosa</i>	123, 249, 254, 257, 260, 304, 374, 440
<i>Ulmus rubra</i>	122
<i>Umbellularia</i>	8, 11, 12, 15
<i>Umbellularia Californica</i>	120, 250, 254, 257, 302, 374, 440, 576
Umbrella Tree	21, 266, 354, 418
<i>Ungnadia</i>	10-31
<i>Ungnadia heptaphylla</i>	44
<i>Ungnadia heterophylla</i>	44
<i>Ungnadia speciosa</i>	12, 44, 250, 274, 422
Upland Willow Oak	153, 320, 390, 456
<i>Urostigma pedunculatum</i>	127
URTICACEÆ	122-128, 254, 257, 304, 374, 414, 440
Utah, Colorado, and New Mexico, Chicago the principal source of supply of lumber for	568, 569
Utah, lumber industry and manufactures from wood in	486, 487, 569
Utah, rank of, according to value of lumber products	487
Utah, remarks by Mr. E. C. Hall, on the forests of	569-571
Utah, statistics of forests, forest fires, standing timber, etc., in	492, 569-571
<i>Uvaria triloba</i>	23
V.	
<i>Vaccinææ</i>	7
<i>Vaccinium</i>	11, 12, 15, 573
<i>Vaccinium arboreum</i>	96, 249, 292, 432
<i>Vaccinium difusum</i>	96
<i>Vaccinium mucronatum</i>	96
Valparaiso Oak	146, 318, 386, 452
Value and amount of charcoal used for domestic and manufacturing purposes during the census year	480
Value and amount of wood used as fuel in manufactures, etc., during the census year	489
Value and amount of wood used for domestic purposes during the census year	489
Value, economic, of the Redwood of California	578
Value of property destroyed and areas burned over by forest fires during the census year (<i>see, also</i> , under state headings)	491, 492
Value of the forest crop of the United States for the census year	485
Values of woods, comparative	252-255
<i>Vauquelinia</i>	10, 12, 14
<i>Vauquelinia corymbosa</i>	70
<i>Vauquelinia Torreyi</i>	70, 249, 284
VERBENACEÆ	116, 117, 254, 257, 300, 372, 438
Vermont, lumber industry and manufactures from wood in	486, 487, 498-500
Vermont, maple-sugar product of	498
Vermont, rank of, according to value of lumber products	487
Vermont, remarks by Mr. C. G. Pringle on the forests of	498-500
Vermont, statistics of forests, forest fires, standing timber, etc., in	492, 498-500
Vermont, tabular statement of the amount of Spruce standing in the state of, May 31, 1880	498
<i>Viburnum</i>	10-12, 14
<i>Viburnum Lentago</i>	94, 249, 290, 432
<i>Viburnum prunifolium</i>	94, 249, 253, 257, 366, 432
<i>Viburnum prunifolium</i> , var. <i>ferrugineum</i>	94
<i>Viburnum pyrifolium</i>	94
Vino Maple	47, 274, 358, 422
<i>Virgilia lutea</i>	57
<i>Virgilia secundiflora</i>	57
Virginia, lumber industry and manufactures from wood in	486, 487, 512
Virginia, rank of, according to value of lumber products	487
Virginia, statistics of forests, forest fires, standing timber, etc., in	492, 511, 512
W.	
Wafer Ash	81, 270
Wahoo (<i>Euonymus atropurpureus</i>)	88, 272
Wahoo (<i>Tilia heterophylla</i>)	28, 268, 356, 422
Wahoo (<i>Ulmus alata</i>)	124, 304, 374, 440
<i>Wallia cinerea</i>	130
<i>Wallia nigra</i>	131
Walnut	131, 308, 378, 444
Walnut, Black	131, 308, 350, 376, 414, 444
Walnut, White	130, 306, 370, 442
Washingtonia	11, 12, 16
Washingtonia filifera	217, 250, 255, 259, 348, 414, 480
Washington territory, lumber industry and manufactures from wood in	486, 487, 574
Washington territory, rank of, according to value of lumber products	487
Washington territory, remarks by Mr. Soreno Watson on the forests of	575, 578

	Page.
Washington territory, statistics of forests, forest fires, standing timber, etc., in.....	492, 573-576
Washington territory, wasteful methods of the lumbermen of.....	574
Washington Thorn.....	81, 286
Water Ash.....	110, 298, 370, 436
Water Beech (<i>Carpinus Caroliniana</i>).....	159, 322, 392, 456
Water Beech (<i>Platanus occidentalis</i>).....	129, 306, 350, 376, 442
Water Elm.....	123, 304, 350, 374, 414, 440
Water Hickory.....	136, 310, 350, 380, 414, 446
Water Locust.....	60, 280, 362, 426
Water Maple.....	50, 276, 358, 424
Water Oak (<i>Quercus aquatica</i>).....	152, 320, 350, 388, 416, 454
Water Oak (<i>Quercus palustris</i>).....	152, 320, 388, 455
Water White Oak.....	140, 314, 350, 384, 450
Watson, E. C., remarks on the forests of Utah by.....	569-571
Watson, Sereno, remarks on the forests of certain counties of Oregon by.....	577, 578
Watson, Sereno, remarks on the forests of Idaho by.....	572, 573
Watson, Sereno, remarks on the forests of Montana by.....	565, 566
Watson, Sereno, remarks on the forests of Washington territory by.....	575, 576
Watson, Sereno, report on the forests of the northern Rocky Mountain region by.....	564, 565
Wax Myrtle.....	136, 312, 380, 446
Weeping Oak.....	138, 312, 382, 448
Weight, ash, and specific gravity per cubic foot of dry specimens of the woods of the United States.....	266-349
Weight of the dry wood of trees of the United States.....	249-251
<i>Wellingtonia Californica</i>	184
<i>Wellingtonia gigantea</i>	184
Western Catalpa.....	115, 300, 350, 372, 414, 438
Western division, extent of forests, forest fires, standing timber, etc., in the.....	564-580
West-Indian Birch.....	33, 270, 350, 356, 420
West Virginia, lumber industry and manufactures from wood in.....	486, 487, 512-515
West Virginia, rank of, according to value of lumber products.....	487
West Virginia, remarks by Mr. C. G. Pringle on the forests of.....	512-515
West Virginia, statistics of forests, forest fires, standing timber, etc., in.....	492, 512-515
Weymouth Pine.....	187, 334, 350, 400, 466
Whistlewood.....	46, 274
White and Black Spruces the characteristic trees of the Northern Forest.....	3
White Ash.....	107, 296, 350, 370, 436
White Basswood.....	28, 266, 356, 420
White Bay.....	20, 266, 354, 414, 418
White Birch (<i>Betula alba</i> , var. <i>populifolia</i>).....	159, 324, 350, 392, 458
White Birch (<i>Betula papyrifera</i>).....	160, 324, 350, 392, 458
White Buttonwood.....	87, 288, 364, 430
White Cedar (<i>Chamaecyparis Lawsoniana</i>).....	179, 332, 350, 398, 404
White Cedar (<i>Chamaecyparis sphaeroides</i>).....	178, 330, 396, 416, 464
White Cedar (<i>Libocedrus decurrens</i>).....	176, 330, 396, 462
White Cedar (<i>Thuja occidentalis</i>).....	176, 330, 350, 396, 462
White Cottonwood.....	175, 330, 396, 462
White Cypress.....	184, 334, 350, 398, 466
White Elm (<i>Ulmus Americana</i>).....	123, 304, 350, 374, 414, 440
White Elm (<i>Ulmus racemosa</i>).....	123, 304, 374, 440
White Fir (<i>Abies concolor</i>).....	213, 346, 412, 478
White Fir (<i>Abies grandis</i>).....	212, 346, 412, 478
White Ironwood.....	45, 274, 422
White Laurel.....	20, 266, 354, 414, 418
White Mangrove.....	87, 288, 364, 430
White Maple.....	49, 276, 358, 424
White Oak (<i>Quercus alba</i>).....	137, 265, 312, 350, 380, 414, 446
White Oak (<i>Quercus Garryana</i>).....	138, 312, 350, 382, 448, 576
White Oak (<i>Quercus grisea</i>).....	144, 316, 396, 452
White Oak (<i>Quercus lobata</i>).....	138, 312, 382, 448
White Oak (<i>Quercus oblongifolia</i>).....	144, 316, 396, 452
White Oak, Mountain.....	143, 316, 386, 450
White Oak, Swamp.....	141, 314, 384, 450
White Oak, Water.....	140, 314, 350, 384, 450
White Pine (<i>Pinus flexilis</i>).....	188, 336, 400, 468, 569, 570, 571
White Pine (<i>Pinus glabra</i>).....	201, 340, 406, 472
White Pine (<i>Pinus monticola</i>).....	187, 336, 400, 468, 564, 574, 576
White Pine (<i>Pinus reflexa</i>).....	189, 336, 402, 468
White Pine (<i>Pinus Strobus</i>).....	187, 334, 350, 400, 460, 566, 551, 554, 558
White Pine standing in the forests of Michigan May 31, 1880, tabular statement of the amount of.....	561
White Pine standing in the forests of Minnesota May 31, 1880, tabular statement of the amount of.....	558

GENERAL INDEX.

611

	Page.
White Pine standing in the forests of Wisconsin May 31, 1880, tabular statement of the amount of	554
White Pine the characteristic tree of the Northern Pine Belt	4
White Spruce (<i>Picea alba</i>)	204, 342, 408, 474, 562, 564
White Spruce (<i>Picea Engelmanni</i>)	205, 265, 342, 408, 474, 564, 573, 574
White Spruce (<i>Picea pungens</i>)	205, 344, 408, 474
White Spruce the most important tree in the Pacific region	7
White Stopper	89, 288, 366, 430
White Thorn	78, 286
White Walnut	130, 306, 376, 442
White-heart Hickory	134, 310, 350, 380, 444
Whitewood (<i>Canella alba</i>)	24, 268, 354, 418
Whitewood (<i>Drypetes crocea</i>)	121, 302, 374, 440
Whitewood (<i>Liriodendron Tulipifera</i>)	22, 266, 350, 354, 418
Wild Black Cherry	68, 282, 362, 428
Wild Cherry (<i>Prunus Capuli</i>)	69, 282, 428
Wild Cherry (<i>Prunus demissa</i>)	69, 284, 362, 428
Wild China	44, 274, 358, 414, 422
Wild Cinnamon	24, 268, 354, 418
Wild Dilly	103, 294, 368, 434
Wild Fig	127, 306, 376, 442
Wild Lime (<i>Xanthoxylum Pterota</i>)	31, 270
Wild Lime (<i>Ximenia Americana</i>)	34, 270
Wild Orange (<i>Prunus Caroliniana</i>)	70, 284, 362, 428
Wild Orange (<i>Xanthoxylum Clava-Herculis</i>)	30, 270, 356, 420
Wild Peach	70, 284, 362, 428
Wild Plum	65, 282, 362, 426
Wild Red Cherry	66, 282, 428
Wild Tamarind	64, 282, 362, 426
Willow (<i>Salix amygdaloides</i>)	166, 326, 394, 460
Willow (<i>Salix laevigata</i>)	167, 326, 394, 460
Willow (<i>Salix lasiolepis</i>)	171, 328, 394, 460
Willow, Black (<i>Salix flavescens, var. Scouleriana</i>)	170, 328, 394, 460
Willow, Black (<i>Salix nigra</i>)	166, 326, 394, 460
Willow, Desert	116, 300, 372, 438
Willow, Diamond	170, 328
Willow, Glaucons	169, 328
Willow Oak	154, 322, 390, 456
Willow Oak, Upland	153, 320, 390, 456
Willow, Sand-bar	168, 326
Willow, Silky	171, 328
Winged Elm	124, 304, 374, 440
Wintera <i>Canella</i>	24
Wisconsin, lumber industry and manufactures from wood in	486, 487, 554, 556
Wisconsin, rank of, according to value of lumber products	487, 555
Wisconsin, remarks by Mr. H. C. Putnam on the forests of	555-558
Wisconsin, statistics of forests, forest fires, standing timber, etc., in	492, 554-558
Wisconsin, tabular statement of the amount of White Pine standing in the forests of, May 31, 1880	554
Witch Hazel	85, 286
Woodlands in Kentucky, pasturage of	546
Wood, manufactures from, and the lumber industry in—	
Alabama	486, 487, 524, 525
Arizona	486, 487, 569
Arkansas	486, 487, 544
California	486, 487, 578, 580
Colorado	486, 487, 567, 568
Connecticut	486, 487, 500, 501
Dakota	486, 487, 561
Delaware	486, 487, 511
District of Columbia	486, 487
Florida	486, 487, 521, 523
Georgia	486, 487, 519, 520
Idaho	486, 487, 572
Illinois	486, 487, 548-550
Indiana	486, 487, 547
Iowa	486, 487, 560
Kansas	486, 487, 563
Kentucky	486, 487, 546
Louisiana	486, 487, 536, 537
Maine	486, 487, 494-496
Maryland	486, 487, 511
Massachusetts	486, 487, 506, 501
Michigan	486, 487, 551, 552
Minnesota	486, 487, 558
Mississippi	486, 487, 531
Missouri	486, 487, 561
Montana	486, 487, 564
Nebraska	486, 487, 562
Nevada	486, 487, 571
New Hampshire	486, 487, 497, 498
New Jersey	486, 487, 506
New Mexico	486, 487, 568
New York	486, 487, 502-505
North Carolina	486, 487, 515

	Page.
Wood, manufactures from, and the lumber industry in—continued.	
Ohio	486, 487, 547
Oregon	486, 487, 577
Pennsylvania	486, 487, 506-510
Rhode Island	486, 487, 500, 501
South Carolina	486, 487, 518, 519
Tennessee	486, 487, 545
Texas	486, 487, 541, 542
Utah	486, 487, 539
Vermont	486, 487, 498-500
Virginia	486, 487, 512
Washington	486, 487, 574
West Virginia	486, 487, 512-515
Wisconsin	486, 487, 554, 556
Wyoming	486, 487, 567
Wood used as fuel in manufactures, etc., during the census year, amount and value of	489
Wood used for domestic purposes during the census year, amount and value of	489
Woods, comparative values of	252-255
Woods, method of determining the fuel value of	247, 251, 252
Woods, method of determining the strength of	252
Woods of the United States, actual fuel value of some of the more important	350-353
Woods of the United States under compression, behavior of the principal	418-481
Woods of the United States under transverse strain, behavior of the principal	354-414
Woods, relation between transverse strength and specific gravity of certain	259-264
Woods, specific gravity and ash of	248-251
Woods, where specimens of, were deposited	247, 248
Wyoming, lumber industry and manufactures from wood in	492, 567
Wyoming, rank of, according to value of lumber products	487
Wyoming, statistics of forests, forest fires, standing timber, etc., in	492, 566, 567
X.	
<i>Xanthoxylum</i>	10, 11, 13
<i>Xanthoxylum Americanum</i>	29, 250, 268
<i>Xanthoxylum aromaticum</i>	30
<i>Xanthoxylum Caribæum</i>	30, 249, 253, 256, 270, 356, 420
<i>Xanthoxylum Carolinianum</i>	30
<i>Xanthoxylum Catesbianum</i>	30
<i>Xanthoxylum Clava-Herculis</i>	30, 250, 253, 256, 270, 356, 420
<i>Xanthoxylum Clava-Herculis</i> (<i>Xanthoxylum Americanum</i>)	29
<i>Xanthoxylum Clava-Herculis</i> (<i>Xanthoxylum Caribæum</i>)	30
<i>Xanthoxylum Clava-Herculis, var.</i>	30
<i>Xanthoxylum Clava-Herculis, var. fruticosum</i>	30, 250, 270
<i>Xanthoxylum Floridanum</i>	30
<i>Xanthoxylum frazineum</i>	29
<i>Xanthoxylum frazinifolium</i> (<i>Xanthoxylum Americanum</i>)	29
<i>Xanthoxylum frazinifolium</i> (<i>Xanthoxylum Clava-Herculis</i>)	30
<i>Xanthoxylum hirsutum</i>	30
<i>Xanthoxylum lanceolatum</i>	30
<i>Xanthoxylum macrophyllum</i>	30
<i>Xanthoxylum mile</i>	29
<i>Xanthoxylum Pterota</i>	31, 249, 270
<i>Xanthoxylum ramiflorum</i>	29
<i>Xanthoxylum tricarpum</i> (<i>Xanthoxylum Americanum</i>)	29
<i>Xanthoxylum tricarpum</i> (<i>Xanthoxylum Clava-Herculis</i>)	30
<i>Ximenia</i>	10, 13
<i>Ximenia Americana</i>	34, 249, 270
<i>Ximenia montana</i>	34
<i>Ximenia multiflora</i>	34
Y.	
Yanpon	36, 272
Yazoo delta in Mississippi, forests of the	535, 536
Yellow Ash	57, 278, 360, 426
Yellow-bark Oak	149, 265, 318, 350, 388, 454
Yellow Birch	161, 324, 392, 458
Yellow Cypress	178, 332, 398, 464
Yellow Fir	209, 265, 344, 352, 410, 476, 576
Yellow Haw	82, 286
Yellow Locust	55, 278, 350, 360, 414, 424
Yellow Oak (<i>Quercus prinoides</i>)	143, 265, 318, 384, 450
Yellow Oak (<i>Quercus tinctoria</i>)	149, 265, 318, 350, 388, 454
Yellow Pine (<i>Pinus Arizona</i>)	192, 338, 402, 468
Yellow Pine (<i>Pinus mitis</i>)	200, 340, 350, 406, 472, 529
Yellow Pine (<i>Pinus palustris</i>)	202, 342, 352, 406, 416, 472
Yellow Pine (<i>Pinus ponderosa</i>)	193, 338, 350, 402, 468, 562, 565, 574, 578
Yellow Pine an important and characteristic tree of the Coast Forest	8

	Page.		Page.
Yellow Poplar.....	22, 266, 354, 418	<i>Yucca canaliculata</i>	212, 242, 345
Yellowwood (<i>Cladrastis tinctoria</i>).....	57, 278, 360, 426	<i>Yucca Dracontis</i> , var. <i>arborescens</i>	216
Yellowwood (<i>Schmæfferia frutescens</i>).....	39, 272	<i>Yucca elata</i>	219, 250, 348
Yew (<i>Taxus brevifolia</i>).....	183, 334, 400, 466	<i>Yucca filamentosa</i>	219
Yew (<i>Taxus Floridana</i>).....	186, 334, 521	<i>Yucca Treouliana</i>	226
Yopon.....	36, 272		
Yucca.....	11, 12, 16	H.	
<i>Yucca angustifolia</i> , var. <i>elata</i>	219	<i>Zizyphus Dominicensis</i>	41
<i>Yucca angustifolia</i> , var. <i>radicata</i>	219	<i>Zizyphus emarginatus</i>	30
<i>Yucca baccata</i>	6, 12, 219, 251, 348	ZYGOPHYLLACEÆ.....	26, 28, 353, 356, 263, 356, 490
<i>Yucca brevifolia</i>	8, 12, 218, 251, 348, 578		

2.

